1

DIRT/3D: 3D root phenotyping for field grown maize (*Zea mays*)

Suxing Liu[1,2,3], Carlos Sherard Barrow[4], Meredith Hanlon[5], Jonathan P. Lynch[5], Alexander Bucksch*[1,2,3]

[1] Department of Plant Biology, University of Georgia, Athens, GA, United States

[2] Warnell School of Forestry and Natural Resources, University of Georgia, Athens, GA, United States

[3] Institute of Bioinformatics, University of Georgia, Athens, GA, United States

[4] Instrumentation Shop, University of Georgia, Athens, GA, United States

[5] Department of Plant Science, Penn State University, State College, PA, United States

*corresponding author

Abstract

The development of crops with deeper roots holds substantial promise to mitigate the consequences of climate change. Deeper roots are an essential factor to improve water uptake as a way to enhance crop resilience to drought, to increase nitrogen capture to reduce fertilizer inputs and to increase carbon sequestration from the atmosphere to improve soil organic fertility. A major bottleneck to achieving these improvements is high-throughput phenotyping to quantify root phenotypes of field-grown roots. We address this bottleneck with DIRT/3D, a newly developed image-based 3D root phenotyping platform, which measures 18 architecture traits from mature field-grown maize root systems. DIRT/3D reliably computed all 18 traits, including distance between whorls and the number, angles, and diameters of crown and brace roots, on a test panel of 12 contrasting maize genotypes. The computed results were validated through comparison with manual measurements. Overall, we observed a coefficient of determination of $r^2 > 0.84$ and a high broad-sense heritability of $H^2_{mean} > 0.6$ for all important traits. The average values of the 18 traits and a newly developed descriptor to characterize a complete root architecture distinguished all genotypes. DIRT/3D is a step towards automated quantification of highly occluded maize root systems. Therefore, DIRT/3D supports breeders and root biologists in improving carbon sequestration and food security in the face of the adverse effects of climate change.

Introduction

Evaluating the information encoded in the shape of a plant as a response to environments is essential to understand the function of plant organs (Bucksch, 2011). In particular, roots exhibit shape diversity that is measurable as variation in rooting angles, numbers of roots per type, length, or diameter of roots within a root system (Lynch and Brown, 2012). An understanding of variation in root systems facilitates breeding for favorable root characteristics to improve yield in suboptimal conditions, including those resulting from climate change (Lynch, 2013). Improving root phenotypes through crop breeding and management holds promise for improved food security in developing nations, where drought, low soil fertility and biotic constraints to root function are primary causes of low yields, and also for reducing the environmental impacts of intensive agriculture by reducing the need for intensive fertilization and irrigation (Lynch, 2019). Root traits also offer opportunities to improve carbon sequestration (Paustian et al., 2016). These challenges demand efforts across a range of disciplines, from mathematics over computer science to plant biology and applied fields like plant breeding and agronomy (Bucksch et al., 2017; Bucksch et al., 2017).

A major interdisciplinary challenge in root biology is the development of deeper rooting crop varieties. Deeper roots promise a three-fold impact: they improve drought resilience, lower fertilizer input, and decrease atmospheric carbon. Deeper roots improve drought resilience to stronger and more frequently occurring droughts (Ault, 2020) by tapping into water in deep soil domains (Lynch and Wojciechowski, 2015). Nitrogen capture increases when roots grow deeper because nitrogen diffuses into and accumulates in deeper soil layers (Lynch, 2019). Deeper rooting crops increase carbon sequestration (Smith et al., 2007) mostly by depositing more organic residues in soils, thereby replenishing carbon after harvest (Paustian et al., 1997). An important tool in breeding deeper roots is the large-scale automated evaluation of root traits in the highly occluded root system architecture (Topp et al., 2016). Maize (*Zea mays*) in particular, with over 700 Mt of maize production world-wide (Ranum et al., 2014), is a prime target for improving rooting depth. However, measuring important root traits for deeper rooting in maize is hampered by the availability of advanced root phenotyping methods on the field-scale (Kuijken et al., 2015). Therefore, the research community has pushed for the development of a root phenotyping system that operates under field conditions (Paez-Garcia et al., 2015).

The field vs. lab complement

The majority of existing phenotyping methods to evaluate root architecture non-destructively emerged from laboratory settings. These methods range from fully automatic (Galkovskyi et al., 2012) to manually assisted (Lobet et al., 2011) and consider a variety of growth systems like gel cylinders (Iyer-Pascuzzi et al., 2010), rhizotrons (Nagel et al., 2012; Rellán-Álvarez et al., 2015), mesocosms (Nagel et al., 2012) and germination paper (Falk et al., 2020). Root phenotyping under lab conditions necessitates the use of constrained growth containers (Poorter et al., 2012; Bourgault et al., 2017), artificial growth media (Oliva and Dunand, 2007), and environments that potentially alter root system architecture (de Dorlodot, Bertin et al. 2005, Draye, Thaon et al. 2018). Therefore, it is essential to translate phenotyping experiments from the lab into repeatable field experiments (Zhu et al., 2011; Bucksch et al., 2014; Poorter et al., 2016).

In contrast, root phenotyping in the field is currently either invasive or destructive. Invasive procedures record small sections of the root with minirhizotron cameras placed in the soil (Gray et al., 2013; Yu et al., 2019). Still, these procedures are incapable of recording the full root system. Therefore, investigating root system architecture relies heavily on the destructive excavation of the root crown. Shovelomics is the standard field-ready protocol to excavate root systems in the field (Trachsel et al., 2011). It was initially developed for maize and has undergone a constant refinement by the root research community (Zheng et al., 2020). Other crops, including common bean (*Phaseolus vulgaris*) and cowpea (*Vigna unguiculata*) (Burridge et al., 2016), wheat (*T. aestivum*) (Slack et al., 2018; York et al., 2018), rapeseed (*Brassica napus*) (Arifuzzaman et al., 2019) and cassava (*Manihot esculenta*) (Kengkanna et al., 2019) have specialized shovelomics protocols in place. However, the manual excavation of the root system, followed by visual scoring and manual trait measurement, is difficult and subjective to the researcher.

Digital imaging of root traits in 2D

In response, software to measure root traits in simple digital images became available. Approaches to record root
traits in the field use different methods in terms of software platforms and imaging setups. According to the
software catalogue "The quantitative plant" (Lobet, 2020), DIRT is the only online platform (Das et al., 2015). The
DIRT platform provides image processing for over 750 root researchers following an easy to reproduce imaging
protocol. For imaging, DIRT needs a tripod, a consumer camera, and a black background with a white circle. Just
recently, DIRT enabled projects associated with root architecture and micronutrient content (Busener et al., 2020)
and translated traits from the lab to the field (Salungyu et al., 2020). More sophisticated and expensive imaging
setups use specialized tents (Colombi et al., 2015) and carefully designed imaging boxes (Grift et al., 2011;
Seethepalli et al., 2019), along with computationally cheap traits, for use on personal computers. The user can,
therefore, choose a system that suits the project needs. Systems generally vary in the number of instruments,
tools, and samples transported between the lab and field site, as well as the cost of the imaging setup and
hardware requirements. However, all these systems share a single obstacle: Resolving the highly occluded
branches of a dense 3D root system. The occlusion challenge arises when the 2D image projection of the 3D
branching structure 'hides' information of branching locations. Hence, the branching information is unrecoverable
and lost (Bray and Topp, 2018).

Digital imaging of root traits in 3D

3D approaches are capable of resolving even highly occluded branching structures (Bucksch, 2014). Gel systems
were among the first methods to measure fully resolved 3D root systems of younger roots (Clark et al., 2011; Topp
et al., 2013) and to capture some of their growth dynamics (Symonova et al., 2015). The bottleneck of imaging and
measuring older root systems in constrained growth containers filled with soil, however, remained. Therefore, X-
Ray computed tomography (CT) became a widely used tool to phenotype roots in pots filled with soil and soil-like
substrates (Pfeifer et al., 2015). These lab developments revealed for example new characteristics in maize root
development (Jiang et al., 2019). The X-ray CT approach is, in its applications, comparable to magnetic resonance
imaging (MRI) (Metzner et al., 2015). MRI also provides a 3D model of the root (van Dusschoten et al., 2016) and
can be used for time-lapse imaging of growth processes (Jahnke et al., 2009). The benefits of both X-Ray CT and
MRI are substantial and subject to scientific discussion (Fischer et al., 2016). However, X-Ray CT and MRI do not
meet the needs for large-scale field studies: Firstly, both technologies restrict plant growth to the size of a given
growth container. The restriction to smaller sizes is proportional to higher achievable resolution. Therefore, it is
common to observe an immature "pot phenotype" instead of a relevant phenotype grown in field soils (Poorter et
al., 2012; Bourgault et al., 2017). Secondly, both methods can take about 30 minutes or more to collect root
imaging data in soil. Extremes of several days are reported for X-Ray CT systems to achieve the resolution of root
hairs (Keyes et al., 2013; Sozzani et al., 2014). An additional constraint is the cost of constructing, operating, and
staffing such facilities, few of which are devoted to root studies.

In response to these phenotyping limitations, we developed DIRT/3D as an automatic 3D root phenotyping system
for excavated root systems grown in agricultural fields. Our approach consists of a newly developed 3D root
scanner and root analysis software. The 3D root scanner captures image data of one excavated maize root in about
five minutes. Our software uses the image data to produce a colored 3D point cloud model and to compute 16 root
traits. The computed traits measure individual roots and also characterize the complete root system. Individual
root traits include number, angle, and diameters of crown and brace roots. Traits like eccentricity or the distance
between brace and crown root whorls characterize the root system. The computed traits are known to be relevant
and reported frequently in literature as manual measurements (Saengwilai et al., 2014; Zhan et al., 2019). We also
introduce a new 3D whole root descriptor that encodes the arrangement of roots within the root system.

Results

DIRT/3D enables automatic measurement of 3D root traits for field-grown maize

We developed DIRT/3D (Digital Imaging of Root Traits in 3D) system to phenotype excavated root crowns of maize
(Figure 1). The system includes a 3D root scanner and a suite of parameter-free software that reconstructs field-
grown maize roots as a 3D point cloud model. The software also contains algorithms to compute 18 root traits.

Figure 1: Schematic overview of our DIRT/3D system. Field grown roots (a) excavated with the Shovelomics protocol (Trachsel et al., 2011) are
placed in the 3D root scanner (b). The scanner, with ten synchronized industrial cameras mounted on a curved frame, acquires about 2000
images of the root. The images are transferred to and stored in the CyVerse Data Store (Merchant et al., 2016) (c). The 3D reconstruction is
computed with DIRT/3D´s structure-from-motion software (d) and yields the resulting 3D root model (e). Overall, the analysis software
calculates 18 root traits from the 3D point cloud of the root system. The image in (f) shows examples for the trait classes angle, diameter, and
length. All developed hardware designs are open and software methods are open source. Executable are available as a Singularity or Docker
container (Kurtzer et al., 2017).

The 3D root scanner (Figure 2) utilizes ten industrial cameras mounted on a rotating curved frame (Figure 2a) to
capture images from all sides of the maize root (Supplementary Material 1). Scanning of one maize root completes
in five minutes. After obtaining the image data, we reconstruct a 3D point cloud of the root system. By analyzing
thin level sets of the 3D point cloud, DIRT/3D revealed traits behind multiple layers of occlusions (Detailed pipeline
in Supplementary Material 2). For example, DIRT/3D measures the distance between the brace and crown root
whorl and the number of brace and crown roots. DIRT/3D also tracks every individual root within the root system,
starting from the stem down to the emerging lateral in the excavated root crown. Each individually tracked root
enables the measurement of numbers, angles, and diameters at the individual root level (Figure 1f).

149 We used a panel of 12 maize genotypes with 5-10 replicates per genotype to validate the DIRT/3D pipeline. For our
150 validation trial, the 3D root scanner captured images at pan intervals of 1 degree and tilt intervals of 10 degrees.
151 Figure 3 shows a visual comparison of the captured root architectural variation between the genotypes.

Figure 3: The automatic DIRT/3D pipeline generates a detailed 3D point clouds of excavated roots. Examples show excavated maize root crowns their 3D point cloud models and structure graphs from the test panel of 12 genotypes. Visual comparison of the 2D views of real roots and their respective 3D root models shows that root architecture, along with the color, is reconstructed (Supplementary Material 4). All 3D models used in the paper are available as .ply file in Supplementary Material 17.

Level set scanning enables extraction of traits from 3D root models

157
We developed a new method to perform a top-down level set scan of the 3D root model to compute 3D root
traits. For a vertically aligned model, we slice the 3D root model from top to bottom at consecutive depth levels
(Figure 4). The number slices represent the constant imaging volume of the scanner and therefore, vary by root
system size. Two benefits result from the fixed scanning volume. First, the transformation to mm is constant, and
second, the optimal slice thickness can be determined experimentally for all roots. Therefore, all parameters are
constants in the algorithm. Here, a level set image is the commonly used vertical 2D projection of each slice onto a
plane (Bucksch, 2011; Mairhofer et al., 2012; Cochard et al., 2015; Dinas et al., 2015; Hyun et al., 2016)
representing the sequential distribution of roots into deeper soil levels (Figure 4 (b)-(e)).

Figure 4: The level set image sequence for the estimation of root traits. (a) A sliding plane scans the 3D root model from top to bottom to
acquire a level set image sequence. The information content per level set image varies with depth and generally encodes the points at a pre-
defined distance to the sliding plane. For example, at the top (b and d), only information about the stem appears in the level set image. At a
middle position (c and e), individual roots are visible as additional circles.

Ideally, each root is a closed circle in the level set plane. However, roots are under-sampled or affected by noise
such that the contours of some roots are disconnected. A video sequence of all level set images would result in a
flickering effect. Therefore, we use a phase-based frame interpolation technique (Meyer et al., 2015) to smooth
the level set image sequence. This method estimates transition frames between the level set images, which is
equal to an up-sampling process of the 3D point cloud (Waki, 2016). Insufficient sampled locations of the root
models are interpolated, which results in a smooth connection of formerly disconnected contours on level set
images (Zhao et al., 2018). A comparison of the original and smoothened sequence of level set images shows the
increased density of the 3D root model (Supplementary Material 5).

The active contour snake model identifies individual roots per level set image

179
The image sequence of the smoothened level sets is the key to compute the location and size parameters of
individual roots. Applying the active contour snake model (Kass et al., 1988) to each level set image results in a
curve that circumscribes each individual root in the level set image (Mugerwa et al., 2019). Each curve contracts
and moves towards the closed boundaries of an individual root by minimizing a partial differential equation, where

image boundaries represent a low energy state for the active contour. The partial differential equations formulate
a trade-off between an internal and external energy term. The internal energy describes the continuity and
smoothness of the contour to controls for curve deformations, and an external energy that describes how well the
contour fits the individual root (Zhao et al., 2018).

(a) Five sample images from smoothed level set image sequence

(b) Active snake results of (a)

Figure 5: Active contour snake method identifies individual roots by analyzing connected components. (a) shows five sample images from the
smoothened level set images. (b) shows the corresponding results from the active contour snake method. Each extracted individual root is
color-coded with connected component labeling. The active contour snake model detects individual roots in each level set image. A video
showing the active contour snake evolving process is available in Supplementary Material 6.

Our algorithm initializes a circle around each individual root in each level set as an initial input to the minimization
of the active contour snake (Supplementary Material 7). During the iterative minimization of the energy function,
we use a periodic boundary condition to enforce a closed curve. The resulting closed boundary curves represent
first estimates of individual roots and used as input to compute a binary mask for each level set image sequence
using Otsu's binarization method (Moghaddam and Cheriet, 2012). We adopt the connected components labeling
method to distinguish and label each closed-boundary object, representing individual roots (Playne and Hawick,
2018). The result of connected components labeling is a multiple segmentation of individual roots represented by
colored components (Figure 5).

However, due to the complexity of the maize root system, roots intertangle or adhere to each other. In an image
of the level set sequence, the entanglement will be visible as one connected component instead of two distinct
components. We use the watershed segmentation to segment the overlapping root (Supplementary Material 8).
The idea behind the watershed algorithm is to interpret grey values in the image as a local topography or
elevation. The algorithm uses pre-computed local minima to flood basins around them. The algorithm terminates
the flooding of a basin when the watershed lines of two basins meet. The Euclidean Distance Transform (EDT) of
the image allows for direct detection of the local minima (Fabbri et al., 2008). In that way, watersheds assign each
pixel to a unique component and allows the distinction entangled roots per level set image (Roshanian et al.,
2016).

A combination of Kalman filters and the Hungarian algorithm tracks individual roots

We developed an individual root tracking method by adopting a combination of Kalman filters and the Hungarian
algorithm (Sahbani and Adiprawita, 2016). The algorithm detects individual roots for consecutive level set images.
Once individual roots are detected, the Hungarian algorithm matches the corresponding individual roots across the

level set images. To improve the speed of the Hungarian algorithm, we use a Kalman filter to predict matching
individual roots in consecutive level set images (Figure 6). Behind the scenes, the tracking algorithm builds a
mathematical model of expected depth development of the root. In doing so, the algorithm uses the current
position, relative speed, and acceleration of individual roots to predict their location in the following level set
image. As a result, we obtain an initial root structure directly from the point cloud (see Figure 3 for examples of all
12 genotypes). An animation and video showing the individual root tracking process are available in
Supplementary Material 18.

(a) 3D root model

(b) Individual root tracking

(c) Computed root structure

(d) root structure details: detected whorl locations

222 **Figure 6: A combination of Kalman filters and the Hungarian algorithm tracks individual roots in the root system.** (a) 3D model of a field-
223 grown maize root generated by the DIRT/3D reconstruction. (b) Visualization of the tracking process of individual roots from level set images.
224 Two level set images are visualized with 50% transparency to show the tracking of individual roots in 3D space. (c) 3D visualization of the root
225 structure graph consisting of all tracked trajectories of brace and crown roots. (d) The structure graph data includes the detected whorls and
226 the corresponding brace and crown roots. Individual roots are colored by depth relative to the root system.

Trace connection and back tracking to improve the computed root structure

Under-sampled regions within the point cloud can result from left over soil that blocks the view into the root
system. Technically, this may lead to unconnected roots in the 3D model of the root system. As a solution to the
disconnection problem, we describe each root part by its curvature and Euclidean distance between every pair of
adjacent roots and root parts. If two close-by root parts have similar curvature value, we connect them by
interpolating a curved connection between both parts. We accept two connected parts as a valid solution if the
newly connected part does not deviate from the interpolated curve. Once we connected all root parts that fit the
same curve, we adopt the Iterative Back-Tracking (Liu et al., 2018) to connect remaining root parts to the root
structure either as continuous curve or as a new branching root.

During the sequential processing of all level set images, we calculate the diameters of the minimal bounding circle
that covers all points in all level set images in a 2D projection. Table 1 lists all 18 root architecture traits that
DIRT/3D computes.

Table 1: Description of DIRT/3D traits. Traits describe either a root system (RS) characteristic or measure an individual root (IR)
within the root system.

Trait name	Trait type	Trait description	Unit
Brace-crown whorl distance	RS	Distance between the last crown root whorl and first brace root whorl (Figure 1f).	mm
Crown-crown whorl distance	RS	Distance between the second last crown root whorl and the last crown root whorl (not always detectable)	mm
Brace/crown root angle	IR	Angle of the line fitted through 70% of the root length to the horizontal plane (Figure 1f)	degree
Brace/crown root diameter	IR	Average diameters of the fitted circles of the brace/crown roots at the detected whorl locations (Figure 1f)	mm
Lateral root diameter	IR	Average diameter of roots that are not identified as brace or crown roots.	mm
Brace root length	IR	Average length of the B-splines fit to all recognized individual brace roots (Figure 1f)	mm
Number of brace/crown root	RS	Count of the brace/crown roots in the root system (Figure 1f)	count
Occupancy index of brace/crown root	RS	Sum of all brace/crown root diameters at the respective whorl location divided by the stem perimeter	ratio
Root system diameter	RS	Average of maximum diameters measured at ten equidistant depth intervals (Figure 1f).	mm
Root system eccentricity	RS	Ratio of the averages of minimum and maximum diameter of the roots system at consecutive depth intervals (Figure 4)	count
Root system density	RS	Average root area divided by the convex hull area at consecutive depth intervals	ratio
Excavated root system depth	RS	The number of level set images multiplied by the level set thickness (Figure 4)	mm
Root system projection radius	RS	The radius of the smallest enclosing circle, which is the projection of all individual roots closest to the convex hull of the root system on the horizontal plane.	mm
Stem diameter	-	Diameter of the circle fit through a slice of the stem (Figure 1f)	mm
Root system volume	RS	Sum of all roots volume computed from diameter and length for each individual root	mm^3

3D root traits correlate at individual root and root system level

To test the accuracy and precision of DIRT/3D, we correlated the trait values measured automatically in the 3D point clouds with manually measured traits of the root crown. We validated manually measurable traits such as root system diameter, whorl distances, number of brace and crown roots, brace and crown root angle and weighed the root dry biomass with a precision scale to correlate it with root volume. The correlation analysis of the each of the chosen validation traits showed r^2>0.84 and P < 0.001 (Figure 7 a-d are selected examples among 10 traits validations in Supplementary Material 16). The results for the crown-crown whorl distance (Supplementary Material 16) indicate that at least 2 mm of whorl distance is needed to identify whorls with our methods.

Figure 7: Correlation of automatic and manual trait measurements. All measured traits resulted in correlations of r²>0.8 (See Supplementary
Material 16). The Figure above shows the examples of (a) the minimally occluded brace root angle; (b) crown root angle which is occluded by
the brace roots; (c) the number of crown roots nested inside the brace roots; (d) root system volume extracted with the root tracing algorithm
that generates the whole root descriptor correlated against manually weighed dry biomass.

Broad sense heritability suggests high repeatability of the observed root trait values

Broad-sense heritability,   for all traits (Figure 8) is computed as the ratio of total genetic variance to total
phenotypic variance (Falconer, 1989) to demonstrate the repeatability of the initial fields trial. For quantitative
plant traits, the broad-sense heritability across multiple varieties eliminates the time-consuming steps of
hybridization and population development for determining   . We observed a broad-sense heritability
              for all traits except brace root length, which indicates a moderately strong genetic basis of the
computed traits. Nine of the computed traits resulted in   (Figure 8), which indicates that the
calculated traits show minimal variation within genotypes sampled with the Shovelomics method. Note that the
crow-crown whorl distance could not be included into the heritability calculation because it is not always
detectable.

Figure 8: Broad sense heritability for 18 computed traits. Phenotypes vary between the individuals because of both environmental factors, the
genes that control traits, as well as various interactions between genes and environmental factors. We computed broad-sense heritability for all
3D traits in Table 1. All but one trait suggests a moderately to strong genetic basis to explain the observed inter-genotypic variation with
> 0.6.

3D root traits distinguish genotypes in the test panel

A principal component analysis of the 18 root traits shows clusters per genotype in the projection on the first and
second principal component. In our test data set, these two principal components explain 51.9% of the overall
observed variance (Figure 9a). The 3D root traits also distinguished genotypes by their means (Figure 9b). Overall,
we found that no single trait classifies all genotypes. However, an ANOVA test revealed that the means of each pair
of genotypes distinguishes in at least four traits (Supplementary Material 16). For example, genotype PA762 and
B101 show a significant difference with traits such as brace root diameter and lateral root diameter. However,
B101 and PHG50XHG47 do not show separable mean values in the brace and crown root angles. We excluded the
crown-crown whorl distance from the analysis because the distance was not detectable for some genotypes.

a.

 b.

Figure 9: Genotype differentiation of 12 maize genotypes. (A) We normalized all the mean trait values of computed 3D root traits from
DRIT/3D. Colored points denote the normalized mean values of the 16 root traits and error bars correspond to the standard error of the mean.
The lines guide the reader visually to explore the phenotypic variation between genotypes of the test panel. For example, the mean of
genotype PA762 and B101 distinguishes in brace root diameter and lateral root diameter; Genotype PHG50XHG47 and PA762 distinguish in the
root projection radius. However, B101, PA762, and PHG50XHG47 do not show distinguishing mean values in the brace root angle. (B) Principal
component analysis (PCA) of all extracted traits and individual root systems. Colors correspond to genotypes and points to measured root
systems.

Whole root descriptor distinguishes the unique spatial arrangement of individual roots for all genotypes

We introduce a 3D variation of the established D-curve for 2D images (Bucksch et al., 2014) as a whole root
descriptor. We compute the descriptor from the sequence of level set images derived from the reconstructed 3D
root models. For each level set image, we compute the number of pixels that represent roots as a measure for the
area. We found that the accumulation of root area across the level-set images is an intrinsic characteristic of each
genotype (Supplementary Material 9). The root area of each level set image corresponds to a certain thickness.
Therefore, the descriptor describes biologically the accumulation of root volume as a function of rooting depth.
The descriptor is robust to outliers and measurement errors because it relies on the cumulative distribution
function (Chun et al., 2000; Lee, 2001; Kyurkchiev, 2015). Figure 9 shows the results of the test panel of the 12
genotypes used the maize test panel. The whole root descriptor distinguished the unique arrangement of
individual roots for all 12 genotypes as a characteristic mean curve, as shown in Figure 10.

301
302 **Figure 10: The whole root descriptor of all 12 maize genotypes of the test panel.** The descriptor encodes the spatial arrangement of individual
303 roots within the root system as a function of the excavation depth. We define the curve of the cumulative root system area as the cumulative
304 distribution function (CDF) of the area per level set for each genotype. The error bar denotes the standard error of the normalized root area.
305 Each genotype associates with a characteristic CDF curve (colored coded). All genotypes distinguish visually from each other in their curve
306 characteristics.

Discussion

The presented system to measure 3D traits in highly occluded root systems advances root phenotyping because it
measures highly occluded traits such as whorl distances and number or brace and crown roots in the dense maize
root system without the need for more costly and cumbersome MRI and CT systems. Furthermore, we presented a
first 3D whole root descriptor for plant root architecture that measures the accumulation of root volume as a
function of rooting depth. For the test panel of twelve genotypes, a minimum of four traits distinguished all
genotypes. In contrast, the whole root descriptor is one characteristic that distinguished all genotypes with one
mathematical expression. Together, these make progress towards the unaccomplished goal of phenomics to
measure the comprehensive appearance of a continuously reshaping phenotype(Houle et al., 2010).

From a validation point of view, the top angle (r^2=0.87) and median root system width (r^2=0.88) published in the
2D DIRT system (Das et al., 2015) show equally good correlations with manual measurements as the comparable
brace root angle (r^2=0.88) and root system diameter (r^2=0.89) on a maize data set. Certainly, comparable results
could be expected for the crown root angles if occluding roots were mechanically removed from the root system.
The presented 3D system however extends the availability of traits compared to the 2D DIRT system in two ways:
Firstly, many occluded traits can be revealed without mechanical work and secondly, traits like brace and crown
root number or, root system eccentricity can be estimated without additional manual data collection.
Nevertheless, some algorithmic and technical challenges remain to exploit the utility of 3D root phenotyping for
breeders fully. To date, standard calibration procedures for structure from motion scanners with multiple cameras
are rare (Conte et al., 2018) and limit the achievable resolution to detect nodes within the maize root system.
Further research will focus on the details of the photogrammetric calibration of the 3D root scanner, which will
allow for thinner cross-section slices during level set extraction. We believe that root models of higher resolution
will enable the reconstruction of a more detailed root architecture to obtain measures of all nodes in the maize
root system.

We demonstrated the possibility to retrieve high geometric detail from field excavated roots. We argue that it will
be possible to obtain geometrically complete measurements in the sense of Euclid's definitions in Elements I-IV
and VI (Callahan and Casey, 2015). Local measurements of length, diameter, and angle are sufficient to reconstruct
every solid 3D object if sampled at sufficiently high rates. Again, research on the calibration technique used for
structure from motion scanners seems likely to be the limiting factor in achieving the needed resolution.
Assembling the complete geometrical system of the root system will allow us to describe the whole root system
and its spatial arrangements in one single mathematical construct. We presented a first 3D whole root descriptor
that is methodologically similar to the D- and DS-curve for 2D images (Bucksch et al., 2014). Here, we encoded root
architecture as an aggregate of the extracted 3D root volume and could reliably distinguish the roots of different
genotypes for a small diversity panel. However, our approach neglects the spatial arrangement of roots, which
limits the encoded detail. An extension of the presented whole root descriptor would enable the quantification of
morphological differences to understand the variation of architecture arrangements. Besides, the comparison
between plant species with similar topological organization but different geometric growth such as dense first
order laterals yet with varying patterns of curvature along the root, would be enabled.

The observed broad-sense heritability suggests a strong repeatability of our first study ($H^2_{mean} > 0.7$ for most traits).
The observed repeatability might overestimate the heritability values needed for genetic mapping studies due to
insufficiently captured root architecture variation with the Shovelomics method. However, the observed
repeatability, paired with near geometric completeness, indicates the presence of a local and global architecture
control by genes. Local control relates to phenes that assemble the architectural phenome of roots as a set of
mappable and locally measurable traits (Lynch and Brown, 2012). However, it is still an open question if a "global
blueprint" exists that controls the arrangement and state of root architecture phenes (Jiang et al., 2019) unless we
can map whole root architectures that are geometrically complete. A necessary step towards answering this
question from a mathematical point of view is to define a mathematical basis of locally controlled traits or phenes.
Since phenes are often mappable to genes (Yablokov, 1986), a mathematically independent basis formed by
phenes would open ways for the alternative hypothesis that roots have access to a spectrum of architectures that
acclimatize to their micro- and macro-environments via their species-specific phenes.

Conclusion

Our 3D phenotyping system is arguably the first optical system to handle highly occluded and mature root systems
collected in the field. It is worth noting that the time required to collect the imaging data is around five minutes,
which is similar than an X-Ray scan at a comparable resolution (Jiang et al., 2019). Unlike many root phenotyping
methods developed under lab conditions, our system measures maize roots grown under field conditions. We also
demonstrated that our system reliably computes previously inaccessible traits, such as whorl distances, the
number and angles of both brace and crown roots. We validated our system for the root trait classes of number,
angle, diameter, and length. Validation results demonstrate the reliability of our system with correlations of $r^2 >$

0.84 for all traits and P < 0.001. From our analysis, we concluded that DIRT/3D could extract 3D root traits
accurately at the individual and system levels.

Both, our software and hardware design are an open and inexpensive 3D root phenotyping solution. At time of
publication, the complete system was developed for about $6000, which includes labor costs to produce the frame
and rather high-end cameras. We currently explore options to build the complete 3D system for about $1500-2000
using cheaper cameras and other means to produce the rotation stand. Our open-source software is available to
the whole plant science community on GitHub, and can be deployed within a platform-agnostic Singularity/Docker
container to be executed independently of the operating system (Supplementary Material 10)
(https://github.com/Computational-Plant-Science). The use of Singularity/Docker containers will allow for
integration with cyber-infrastructures such as CyVerse. These containers can run on any high-performance
computing system that has the Singularity environment installed. The scanner design is part of the publication
(Supplementary Material 3) and can be reproduced, scaled and further developed by everyone.

The presented 3D system requires only one user interaction to place the root crown in the scanner. Placing the
root in the scanner could be replaced by a robot in future. Hence, we see our system as the first milestone towards
automated root trait measurements in the field. Our belief stems from ongoing developments in agricultural
robotics that will excavate field roots "on-the-go" (Shi et al., 2019) in the foreseeable future. In that way, our
system supports breeders and root biologists in the development of crops with increased water uptake, more
efficient nitrogen capture and improved sequestration of atmospheric carbon to mitigate the adverse effects of
climate change without compromising on yield gains.

Material and Methods

Plant material

Plants were grown at The Pennsylvania State University's Russell E. Larson Agricultural Research Center (40°
42'40.915" N, 77°, 57'11.120''W) which has a Hagerstown silt loam soil (fine, mixed, semi-active, mesic Typic
Hapludalf). Fields received fertilization with 190 kg nitrogen ha^{-1} applied as urea (46-0-0). The sites had drip
irrigation. The field management supplemented nutrients other than nitrogen, and applied pest management as
needed. We planted seeds using hand jab planters in rows with 76 cm row spacing, 91 cm alleys, 23 cm plant
spacing, 4.6 m plot length with 3.7 m planted, or ~56,800 plants ha^{-1}. We grew plants in three-row plots, and
sampled only the middle row. Planting occurred on June 5, 2018, and sampling on August 25, 2018, 81 days after
planting. Two fields provided 1ha of space for four replicates.

Twelve genotypes were selected based on previous knowledge of their architectural variation and sampling of a
larger set of genotypes. The twelve genotypes included six inbred lines (B101, B112, DKIB014, LH123HT, Pa762,
PHZ51) and six hybrid lines (DKPB80 x 3IIH6, H96 x 3IIH6, LH59 x PHG29, Pa762 x 3IIH6, PHG50 x PHG47, PHZ51 x
LH59). These genotypes represent the extremes of dense vs. sparse, large vs. small, and maximum and minimum
number of whorls selected from a full diversity panel. The lab of Shawn Kaeppler at the University of Wisconsin
provided the seeds. We selected ten representative plants for five of the genotypes (B112, Pa762, PHZ51, DKPB80
x 3IIH6, H96 x 3IIH6), and five representative plants of the remaining seven genotypes. Sampling followed the
shovelomics protocol (Trachsel et al., 2011), which minimizes variation by selecting similar representative
architectures. Shoots were removed above the first brace root whorl. We air-dried the roots on a greenhouse
bench and then transported the roots to the lab for imaging.

Obtaining the ground truth for root trait validation

Each root system was fixed on a board. We used a ruler to measure the length and diameter of the root system. A
second diameter was measured orthogonal to the board plane to determine the eccentricity of the root system.
We used a protractor to measure the brace and crown root angles to the horizontal from four sides. The average
angle of the four sides was taken to represents the rooting angle for brace or crown roots respectively. To measure

root diameters and whorl distances, we used a steel vernier caliper with a graduation of 0.02mm. The came caliper
was used to measure maximum root system diameters. The dry weight of the root systems was weighed with an
ADAM Core Portable Compact Balance CQT 202 (readability: 0.01g, linearity: 0.02g).

3D root scanner

We designed a 3D root scanner (Figure 2a) to capture images for 3D reconstruction of the root (Supplementary
Material 3). A stepper motor (Nema 34 CNC High Torque Stepper Motor 13Nm with Digital Stepper Driver
DM860I, Figure 2b) rotates a curved metal frame with ten low cost and highly versatile imaging cameras (Image
Source DFK 27BUJ003 USB 3.0, 6mm focal length) around the clamped root crown in a central fixture (Figure 2c).
From the stepper motor, we chose 12800 micro-step resolutions to rotate in 1-degree steps (Figure 2b). The
cameras ship with the 1/2.3" Aptina CMOS MT9J003 sensor and can achieve high image resolution at 3,856×2,764
(10.7 MP) up to 7 fps. We drilled 21 equidistant holes into the curved frame to provide a flexible arrangement of
each camera. A rail track along the curved frame allows for fine adjustment of the camera tilt and pan direction
without compromising stability (Figure 2d). Cameras are then arranged along the curved frame to achieve a
sampling of bigger and smaller root morphology that satisfy the Nyquist theorem to prevent aliasing (Liu et al.,
2009). In the case of maize roots, more cameras are concentrated to image the root crown with high amounts of
small occluded roots. Only 2 cameras cover the stem part of the because the surface area of the stem part usually
has minimal to no conclusion, which guarantees good 3D reconstruction results.

A computing cluster of ten Raspberry Pi 3 B+ synchronizes the image capture of the ten cameras using a master-
slave design (Supplementary Material 11). The synchronized cameras of our 3D root scanner capture
approximately 2000 images per maize root in about 5 min. The newly developed controller software on the
Raspberry Pi computing cluster synchronizes the camera's image capture and the stepper motor movement. Once
the stepper motor receives the "start move" signal via the master unit, it moves all the cameras into their
designated positions. Then, all ten cameras capture images simultaneously. Each Raspberry Pi stores the image
initially on a sim card. During the image capturing process, the stepper motor stands still and waits for the next
"start move" signal. The image data of all Raspberry Pi's automatically transfers to the CyVerse Data Store (Goff et
al., 2011; Merchant et al., 2016). Only the master unit stores information about the CyVerse user account. It uses
the iRods protocol (Ward et al., 2009) to transfer the images from each slave unit to the CyVerse Data Store. In the
following, the 3D reconstruction uses the image data in the online storage to generate the 3D point cloud of the
root system. Alternatively, the image data can be transferred manually to computers within the same WiFi
network.

Automatic reconstruction of the 3D root model with structure-from-motion

Fast Fourier Transform (FFT) detects blurry images

The structure from motion (SfM) method requires detected feature points to be visible in several camera views.
However, pose inaccuracy mechanically inferred by the scanning device or false feature matching may lead to
incomplete reconstructions (Zheng and Wu, 2015). As a result, not all feature points are triangulated to generate
3D points. In our case, a small number of images acquired with the 3D scanner appear dark or blurred as a result of
delayed image capture, frequency of surrounding light sources or vibrations of the scanner (see Supplementary
Material 19 for an example). We detect blurred images using Fast Fourier Transform (FFT) to transform the image
into the frequency domain. The absence or low number of high frequencies compared to the majority of images
indicates a blurred image. Removing blurred images results in higher confidence for feature matches and
therefore, improves model reconstruction quality and point density in SfM approaches.

Illumination adjustment and content-based segmentation to remove redundant information

We use standard deviation and a luminance-weighted gray world algorithm (Lam, 2005) to adjust and normalize
illumination across all captured images. The root is automatically separated from the background using a newly
developed content-based segmentation method (Supplementary Material 12). The method analyzes and compares
color-space differences across all normalized images and omits the redundant information of the background.
Overall, the size of the image data reduces to 30-50% of the original size. In later steps of the pipeline, the
segmentation decreases the number of false feature matches during the 3D reconstruction process as well as the

amount of data transmitted to online storage. The method is fully automatic and parameter-free and uses parallel
processing if available.

Improved feature matching to reduce computation time and improve 3D point cloud resolution

Given the images of segmented roots, we chose the Visual Structure From Motion method (Wu, 2011) as a basis to
develop 3D reconstruction software for roots. The computationally most expensive aspect of structure-from-
motion algorithms is the feature matching between image pairs. The amount and accuracy of the feature matching
determines the quality and resolution of the resulting 3D root model. In the original version, Visual Structure from
Motion performs a full pairwise image matching to build a feature space across all possible image pairs. For
example, the number of permutations P calculates for r images out of a set of n total images with the following
formula:

$$P_n^r = \frac{n!}{(n-r)!}$$

However, the computation of feature matches generates a large amount of false feature correspondences in the
dense root data. We found that image pairs that are not adjacent in the spherical scanner space are particularly
prone to incorrect matching (Supplementary Material 13). We observed that the false feature matches occur
predominantly between the dense and thin roots of the root system. Therefore, we optimized the feature
matching process to be suitable for dense root architectures.

The optimization in our algorithm generates a matching pair list inside a specified sliding window (Supplementary
Material 14). Sliding of the window allows for robust matching among all permutations of image pairs. For
example, given an image set captured around the individual root in the 1-degree interval (360 images in total), we
set the sliding window size as 10% of the image size. The widow size was found experimentally and is the optimum
for the 1-degree interval setting of the scanner. The total number of permutations of image pairs needed for
feature matching is $\frac{360!}{(360-2)!} = 129{,}240$ according to the formula above. For an image of size 1000 ×1000, we set
the sliding window size as 100×100, the number of permutations of image pairs needed will be reduced
to $\frac{100!}{(100-2)!} = 2450$. In that way, we need to compute only 1.89% of all permutations of image pairs.

As a next step, we utilize the RANSAC (random sample consensus) method to detect and remove the falsely
matched pairs. The RANSAC results usually contain only highly distinctive features to track between consecutive
images. Given the locations of multiple matched feature pairs in two or more images, we can produce an
estimation of the positions, orientations of cameras, and the coordinates of the features in a single step using
bundle adjustment (Wu et al., 2011).

Computing root traits from 3D models

We adopted a top-down level set scan of the 3D root model to compute 3D root traits (Supplementary Material
15). This scanning process generates a thin vertical 2D slice or level set image given a fixed plane. We use a phase-
based frame interpolation technique from video processing to smooth the image sequence. We developed a
method to extract individual roots in each level set image using the active contour snake model. Then we use the
watershed segmentation to segment the overlapping roots. Given a smoothed and segmented level set image
sequence, we used a combination of Kalman filters and the Hungarian algorithm (Sahbani and Adiprawita, 2016) to
track all individual roots, and build an embedded graph of the geometry of crown and brace roots. This embedded
graph forms the basis to compute all 18 root architecture traits. Crown and brace root traits are directly derived
from the embedded graph. In each level set image, we compute the lateral root diameter as an average of the
circular projections of point cloud points that are not identified as either crown or brace roots. For each level-set
image, we compute the area covered by physical roots. This area increases whenever roots emerge from a whorl
stays almost constant between whorls. If summarized as a cumulative function of area (see Supplemental Material

9) the starting location in the level set image stack corresponds to the starting point of a plateau in the cumulative
function.

Statistical Analyses

All statistics used python 3.7 and the modules NumPy 1.16 and SciPy 1.2.1 (Oliphant, 2007). Figures 7 and 10 used
matplotlib 3.2.1 (Hunter, 2007) for visualization of the statistics. Figures 8 and 9 used Microsoft Excel Version 16.34
to visualize trait and heritability data. Raw data are available in (Supplementary Material 16).

ACKNOWLEDGMENTS

The research was supported by the USDOE ARPA-E ROOTS Award Number DE-AR0000821 to A.B. and J.P.L.. The work was supported in part by the NSF CAREER Award No. *1845760* to A.B. Any Opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect those of the National Science Foundation. This work used the Extreme Science and Engineering Discovery Environment (XSEDE) resource Stampede2 at the Texas Advanced Computing Center through allocation TG-BIO160088. This material is partly based upon work supported by the National Science Foundation under Award Numbers DBI-0735191, DBI-1265383, and DBI-1743442. URL: www.cyverse.org.

CONTRIBUTIONS

S.L. wrote the manuscript, designed and implemented algorithms, designed and built hardware, performed data analysis and contributed to the experimental design. C.B.S. designed and built hardware. J.P.L. contributed to writing of the manuscript and the project idea, conceived experimental design. M.H. contributed to writing the manuscript, performed experiments and collected data. A.B. conceived the project idea, designed hardware, contributed to the data analysis, wrote manuscript and designed algorithms.

REFERENCES

Arifuzzaman M, Oladzadabbasabadi A, McClean P, Rahman M (2019) Shovelomics for phenotyping root architectural traits of rapeseed/canola (Brassica napus L.) and genome-wide association mapping. Molecular Genetics and Genomics: 1-16

Ault TR (2020) On the essentials of drought in a changing climate. Science **368**: 256-260

Bourgault M, James AT, Dreccer MF (2017) Pot size matters revisited: Does container size affect the response to elevated CO2 and our ability to detect genotypic variability in this response in wheat? Functional Plant Biology **44**: 52-61

Bray AL, Topp CN (2018) The quantitative genetic control of root architecture in maize. Plant and Cell Physiology **59**: 1919-1930

Bucksch A (2011) Revealing the skeleton from imperfect point clouds. Dr. Hut, Munchen

Bucksch A (2014) A practical introduction to skeletons for the plant sciences. Applications in plant sciences **2**: 1400005

Bucksch A, Atta-Boateng A, Azihou AF, Battogtokh D, Baumgartner A, Binder BM, Braybrook SA, Chang C, Coneva V, DeWitt TJ, Fletcher AG, Gehan MA, Diaz-Martinez DH, Hong L, Iyer-Pascuzzi AS, Klein LL, Leiboff S, Li M, Lynch JP, Maizel A, Maloof JN, Markelz RJC, Martinez CC, Miller LA, Mio W, Palubicki W, Poorter H, Pradal C, Price CA, Puttonen E, Reese JB, Rellan-Alvarez R, Spalding EP, Sparks EE, Topp CN, Williams JH, Chitwood DH (2017) Morphological Plant Modeling: Unleashing Geometric and Topological Potential within the Plant Sciences. Front Plant Sci **8**: 900

Bucksch A, Burridge J, York LM, Das A, Nord E, Weitz JS, Lynch JP (2014) Image-based high-throughput field phenotyping of crop roots. Plant Physiology **166**: 470-486

Bucksch A, Das A, Schneider H, Merchant N, Weitz JS (2017) Overcoming the Law of the Hidden in Cyberinfrastructures. Trends Plant Sci **22**: 117-123

Bucksch AK (2011) Revealing the skeleton from imperfect point clouds.

Burridge J, Jochua CN, Bucksch A, Lynch JP (2016) Legume shovelomics: high—throughput phenotyping of common bean (Phaseolus vulgaris L.) and cowpea (Vigna unguiculata subsp, unguiculata) root architecture in the field. Field Crops Research **192**: 21-32

Busener N, Kengkanna J, Saengwilai PJ, Bucksch A (2020) Image-based root phenotyping for plant breeding to improve the nutritional quality of cassava. Plants, People, Planet **(under revision)**

Callahan D, Casey J (2015) Euclid's Elements Redux. John Casey

Chun M-H, Han S-J, Tak N-I (2000) An uncertainty importance measure using a distance metric for the change in a cumulative distribution function. Reliability Engineering & System Safety **70**: 313-321

Clark RT, MacCurdy RB, Jung JK, Shaff JE, McCouch SR, Aneshansley DJ, Kochian LV (2011) Three-dimensional root phenotyping with a novel imaging and software platform. Plant physiology **156**: 455-465

Cochard H, Delzon S, Badel E (2015) X-ray microtomography (micro-CT): a reference technology for high-resolution quantification of xylem embolism in trees. Plant, Cell & Environment **38**: 201-206

Colombi T, Kirchgessner N, Le Marié CA, York LM, Lynch JP, Hund A (2015) Next generation shovelomics: set up a tent and REST. Plant and Soil **388**: 1-20

Conte P, Girelli VA, Mandanici E (2018) Structure from Motion for aerial thermal imagery at city scale: Pre-processing, camera calibration, accuracy assessment. ISPRS journal of photogrammetry and remote sensing **146**: 320-333

Das A, Schneider H, Burridge J, Ascanio AKM, Wojciechowski T, Topp CN, Lynch JP, Weitz JS, Bucksch A (2015) Digital imaging of root traits (DIRT): a high-throughput computing and collaboration platform for field-based root phenomics. Plant methods **11**: 51

Dinas PC, Nikaki A, Jamurtas AZ, Prassopoulos V, Efthymiadou R, Koutedakis Y, Georgoulias P, Flouris AD (2015) Association between habitual physical activity and brown adipose tissue activity in individuals undergoing PET-CT scan. Clinical endocrinology **82**: 147-154

Fabbri R, Costa LDF, Torelli JC, Bruno OM (2008) 2D Euclidean distance transform algorithms: A comparative survey. ACM Computing Surveys (CSUR) **40**: 2

Falconer DS (1989) Introduction to Quantiative Genetics 3rd Edition. Longman Scientific & Technical, Essex, UK

Falk KG, Jubery TZ, Mirnezami SV, Parmley KA, Sarkar S, Singh A, Ganapathysubramanian B, Singh AK (2020) Computer vision and machine learning enabled soybean root phenotyping pipeline. Plant Methods **16**: 5

Fischer A, Lasser T, Schrapp M, Stephan J, Noël PB (2016) Object specific trajectory optimization for industrial X-ray computed tomography. Scientific reports **6**: 19135

Galkovskyi T, Mileyko Y, Bucksch A, Moore B, Symonova O, Price CA, Topp CN, Iyer-Pascuzzi AS, Zurek PR, Fang S (2012) GiA Roots: software for the high throughput analysis of plant root system architecture. BMC plant biology **12**: 116

Goff S, Vaughn M, McKay S, Lyons E, Stapleton A, Gessler D, Matasci N, Wang L, Hanlon M, Lenards A, Muir A, Merchant N, Lowry S, Mock S, Helmke M, Kubach A, Narro M, Hopkins N, Micklos D, Hilgert U, Gonzales M, Jordan C, Skidmore E, Dooley R, Cazes J, McLay R, Lu Z, Pasternak S, Koesterke L, Piel W, Grene R, Noutsos C, Gendler K, Feng X, Tang C, Lent M, Kim S-j, Kvilekval K, Manjunath BS, Tannen V, Stamatakis A, Sanderson M, Welch S, Cranston K, Soltis P, Soltis D, O'Meara B, Ane C, Brutnell T, Kleibenstein D, White J, Leebens-Mack J, Donoghue M, Spalding E, Vision T, Myers C, Lowenthal D, Enquist B, Boyle B, Akoglu A, Andrews G, Ram S, Ware D, Stein L, Stanzione D (2011) The iPlant Collaborative: Cyberinfrastructure for Plant Biology. Frontiers in Plant Science **2**

Gray SB, Strellner RS, Puthuval KK, Ng C, Shulman RE, Siebers MH, Rogers A, Leakey AD (2013) Minirhizotron imaging reveals that nodulation of field-grown soybean is enhanced by free-air CO_2 enrichment only when combined with drought stress. Functional Plant Biology **40**: 137-147

Grift TE, Novais J, Bohn M (2011) High-throughput phenotyping technology for maize roots. Biosystems engineering **110**: 40-48

Houle D, Govindaraju DR, Omholt S (2010) Phenomics: the next challenge. Nat Rev Genet **11**: 855-866

Hunter JD (2007) Matplotlib: A 2D Graphics Environment. Computing in Science & Engineering **9**: 90-95

Hyun S-J, Bae C-W, Lee S-H, Rhim S-C (2016) Fatty degeneration of the paraspinal muscle in patients with degenerative lumbar kyphosis: a new evaluation method of quantitative digital analysis using MRI and CT scan. Clinical spine surgery **29**: 441-447

Iyer-Pascuzzi AS, Symonova O, Mileyko Y, Hao Y, Belcher H, Harer J, Weitz JS, Benfey PN (2010) Imaging and analysis platform for automatic phenotyping and trait ranking of plant root systems. Plant physiology **152**: 1148-1157

Jahnke S, Menzel MI, Van Dusschoten D, Roeb GW, Bühler J, Minwuyelet S, Blümler P, Temperton VM, Hombach T, Streun M (2009) Combined MRI–PET dissects dynamic changes in plant structures and functions. The Plant Journal **59**: 634-644

Jiang N, Floro E, Bray AL, Laws B, Duncan KE, Topp CN (2019) Three-dimensional time-lapse analysis reveals multiscale relationships in maize root systems with contrasting architectures. The Plant Cell **31**: 1708-1722

Kass M, Witkin A, Terzopoulos D (1988) Snakes: Active contour models. International journal of
computer vision **1**: 321-331
Kengkanna J, Jakaew P, Amawan S, Busener N, Bucksch A, Saengwilai P (2019) Phenotypic variation of
cassava root traits and their responses to drought. Applications in plant sciences **7**: e01238
Keyes SD, Daly KR, Gostling NJ, Jones DL, Talboys P, Pinzer BR, Boardman R, Sinclair I, Marchant A,
Roose T (2013) High resolution synchrotron imaging of wheat root hairs growing in soil and
image based modelling of phosphate uptake. New Phytologist **198**: 1023-1029
Kuijken RC, van Eeuwijk FA, Marcelis LF, Bouwmeester HJ (2015) Root phenotyping: from component
trait in the lab to breeding. Journal of experimental botany **66**: 5389-5401
Kurtzer GM, Sochat V, Bauer MW (2017) Singularity: Scientific containers for mobility of compute. PloS
one **12**: e0177459
Kyurkchiev N (2015) On the Approximation of the step function by some cumulative distribution
functions. Compt. rend. Acad. bulg. Sci **68**: 1475-1482
Lam EY (2005) Combining gray world and retinex theory for automatic white balance in digital
photography. *In* Proceedings of the Ninth International Symposium on Consumer Electronics,
2005.(ISCE 2005). IEEE, pp 134-139
Lee H-s (2001) Method and circuit for extracting histogram and cumulative distribution function for
image enhancement apparatus. *In*. Google Patents
Liu S, An P, Zhang Z, Zhang Q, Shen L, Jiang G (2009) On the relationship between multi-view data
capturing and quality of rendered virtual view. The Imaging Science Journal **57**: 250-259
Liu S, Zhang D, Song Y, Peng H, Cai W (2018) Automated 3-D neuron tracing with precise branch erasing
and confidence controlled back tracking. IEEE transactions on medical imaging **37**: 2441-2452
Lobet G (2020) The quantitative plant. *In*, Vol 2020
Lobet G, Pagès L, Draye X (2011) A novel image-analysis toolbox enabling quantitative analysis of root
system architecture. Plant physiology **157**: 29-39
Lynch JP (2013) Steep, cheap and deep: an ideotype to optimize water and N acquisition by maize root
systems. Ann Bot **112**: 347-357
Lynch JP (2019) Root phenotypes for improved nutrient capture: an underexploited opportunity for
global agriculture. New Phytologist **223**: 548-564
Lynch JP, Brown KM (2012) New roots for agriculture: exploiting the root phenome. Philos Trans R Soc
Lond B Biol Sci **367**: 1598-1604
Lynch JP, Brown KM (2012) New roots for agriculture: exploiting the root phenome. Philosophical
Transactions of the Royal Society B: Biological Sciences **367**: 1598-1604
Lynch JP, Wojciechowski T (2015) Opportunities and challenges in the subsoil: pathways to deeper
rooted crops. Journal of Experimental Botany **66**: 2199-2210
Mairhofer S, Zappala S, Tracy SR, Sturrock C, Bennett M, Mooney SJ, Pridmore T (2012) RooTrak:
automated recovery of three-dimensional plant root architecture in soil from X-ray
microcomputed tomography images using visual tracking. Plant physiology **158**: 561-569
Merchant N, Lyons E, Goff S, Vaughn M, Ware D, Micklos D, Antin P (2016) The iPlant collaborative:
cyberinfrastructure for enabling data to discovery for the life sciences. PLoS biology **14**:
e1002342
Metzner R, Eggert A, van Dusschoten D, Pflugfelder D, Gerth S, Schurr U, Uhlmann N, Jahnke S (2015)
Direct comparison of MRI and X-ray CT technologies for 3D imaging of root systems in soil:
potential and challenges for root trait quantification. Plant methods **11**: 17

Meyer S, Wang O, Zimmer H, Grosse M, Sorkine-Hornung A (2015) Phase-based frame interpolation for
video. *In* Proceedings of the IEEE conference on computer vision and pattern recognition, pp
1410-1418
Moghaddam RF, Cheriet M (2012) AdOtsu: An adaptive and parameterless generalization of Otsu's
method for document image binarization. Pattern Recognition **45**: 2419-2431
Mugerwa H, Rey MEC, Tairo F, Ndunguru J, Sseruwagi P (2019) Two sub-Saharan Africa 1 populations of
Bemisia tabaci exhibit distinct biological differences in fecundity and survivorship on cassava.
Crop Protection **117**: 7-14
Nagel KA, Putz A, Gilmer F, Heinz K, Fischbach A, Pfeifer J, Faget M, Blossfeld S, Ernst M, Dimaki C
(2012) GROWSCREEN-Rhizo is a novel phenotyping robot enabling simultaneous measurements
of root and shoot growth for plants grown in soil-filled rhizotrons. Functional Plant Biology **39**:
891-904
Oliphant TE (2007) Python for Scientific Computing. Computing in Science & Engineering **9**: 10-20
Oliva M, Dunand C (2007) Waving and skewing: how gravity and the surface of growth media affect root
development in Arabidopsis. New Phytologist **176**: 37-43
Paez-Garcia A, Motes CM, Scheible W-R, Chen R, Blancaflor EB, Monteros MJ (2015) Root traits and
phenotyping strategies for plant improvement. Plants **4**: 334-355
Paustian K, Agren G, Bosatta E (1997) Modelling litter quality effects on decomposition and soil organic
matter dynamics. Driven by nature: plant litter quality and decomposition
Paustian K, Lehmann J, Ogle S, Reay D, Robertson GP, Smith P (2016) Climate-smart soils. Nature **532**:
49-57
Pfeifer J, Kirchgessner N, Colombi T, Walter A (2015) Rapid phenotyping of crop root systems in
undisturbed field soils using X-ray computed tomography. Plant methods **11**: 41
Playne DP, Hawick K (2018) A new algorithm for parallel connected-component labelling on GPUs. IEEE
Transactions on Parallel and Distributed Systems **29**: 1217-1230
Poorter H, Bühler J, van Dusschoten D, Climent J, Postma JA (2012) Pot size matters: a meta-analysis of
the effects of rooting volume on plant growth. Functional Plant Biology **39**: 839-850
Poorter H, Fiorani F, Pieruschka R, Wojciechowski T, van der Putten WH, Kleyer M, Schurr U, Postma J
(2016) Pampered inside, pestered outside? Differences and similarities between plants growing
in controlled conditions and in the field. New Phytol **212**: 838-855
Ranum P, Peña-Rosas JP, Garcia-Casal MN (2014) Global maize production, utilization, and
consumption. Annals of the New York Academy of Sciences **1312**: 105-112
Rellán-Álvarez R, Lobet G, Lindner H, Pradier P-L, Sebastian J, Yee M-C, Geng Y, Trontin C, LaRue T,
Schrager-Lavelle A (2015) GLO-Roots: an imaging platform enabling multidimensional
characterization of soil-grown root systems. Elife **4**: e07597
Roshanian J, Yazdani S, Ebrahimi M (2016) Star identification based on euclidean distance transform,
voronoi tessellation, and k-nearest neighbor classification. IEEE Transactions on Aerospace and
Electronic Systems **52**: 2940-2949
Saengwilai P, Tian X, Lynch JP (2014) Low crown root number enhances nitrogen acquisition from low-
nitrogen soils in maize. Plant physiology **166**: 581-589
Sahbani B, Adiprawita W (2016) Kalman filter and iterative-hungarian algorithm implementation for low
complexity point tracking as part of fast multiple object tracking system. *In* 2016 6th
International Conference on System Engineering and Technology (ICSET). IEEE, pp 109-115
Salungyu J, Kengkanna J, Thaitad S, Bucksch A, Saengwilai P (2020) Translating maize root traits from
the lab to the field. Field Crops Research **(Under revision)**

Seethepalli A, Guo H, Liu X, Griffiths M, Almtarfi H, Li Z, Liu S, Zare A, Fritschi F, Blancaflor E (2019)
RhizoVision Crown: An Integrated Hardware and Software Platform for Root Crown
Phenotyping. bioRxiv: 569707
Shi X, Choi D, Heinemann PH, Hanlon M, Lynch J (2019) RootRobot: A Field-based Platform for Maize
Root System Architecture Phenotyping. *In* 2019 ASABE Annual International Meeting. American
Society of Agricultural and Biological Engineers, p 1
Slack S, York LM, Roghazai Y, Lynch J, Bennett M, Foulkes J (2018) Wheat shovelomics II: Revealing
relationships between root crown traits and crop growth. BioRxiv: 280917
Smith P, Martino D, Cai Z, Gwary D, Janzen H, Kumar P, McCarl B, Ogle S, O'Mara F, Rice C (2007)
Greenhouse gas mitigation in agriculture. Philosophical transactions of the royal Society B:
Biological Sciences **363**: 789-813
Sozzani R, Busch W, Spalding EP, Benfey PN (2014) Advanced imaging techniques for the study of plant
growth and development. Trends in plant science **19**: 304-310
Symonova O, Topp CN, Edelsbrunner H (2015) DynamicRoots: a software platform for the
reconstruction and analysis of growing plant roots. PLoS One **10**: e0127657
Topp CN, Bray AL, Ellis NA, Liu Z (2016) How can we harness quantitative genetic variation in crop root
systems for agricultural improvement? Journal of integrative plant biology **58**: 213-225
Topp CN, Iyer-Pascuzzi AS, Anderson JT, Lee C-R, Zurek PR, Symonova O, Zheng Y, Bucksch A, Mileyko
Y, Galkovskyi T (2013) 3D phenotyping and quantitative trait locus mapping identify core
regions of the rice genome controlling root architecture. Proceedings of the National Academy
of Sciences **110**: E1695-E1704
Trachsel S, Kaeppler SM, Brown KM, Lynch JP (2011) Shovelomics: high throughput phenotyping of
maize (Zea mays L.) root architecture in the field. Plant and Soil **341**: 75-87
van Dusschoten D, Metzner R, Kochs J, Postma JA, Pflugfelder D, Bühler J, Schurr U, Jahnke S (2016)
Quantitative 3D analysis of plant roots growing in soil using magnetic resonance imaging. Plant
physiology **170**: 1176-1188
Waki K (2016) 3-dimensional elastic image generation method and ultrasonic diagnostic apparatus. *In*.
Google Patents
Ward JH, de Torcy A, Chua M, Crabtree J (2009) Extracting and Ingesting DDI Metadata and Digital
Objects from a Data Archive into the iRODS extension of the NARA TPAP using the OAI-PMH. *In*
2009 Fifth IEEE International Conference on e-Science. IEEE, pp 185-192
Wu C (2011) VisualSFM: A visual structure from motion system.
Wu C, Agarwal S, Curless B, Seitz SM (2011) Multicore bundle adjustment. *In* CVPR 2011. IEEE, pp 3057-
3064
Yablokov AV (1986) Phenetics - Evolution, Population, Trait. Columbia University Press
York LM, Slack S, Bennett MJ, Foulkes MJ (2018) Wheat shovelomics I: A field phenotyping approach for
characterising the structure and function of root systems in tillering species. BioRxiv: 280875
Yu G, Zare A, Sheng H, Matamala R, Reyes-Cabrera J, Frischi FB, Juenger TE (2019) Root Identification in
Minirhizotron Imagery with Multiple Instance Learning. arXiv preprint arXiv:1903.03207
Zhan A, Liu J, Yue S, Chen X, Li S, Bucksch A (2019) Architectural and anatomical responses of maize
roots to agronomic practices in a semi-arid environment. Journal of Plant Nutrition and Soil
Science **182**: 751-762
Zhao F, He Q, Luo J (2018) Cardiac Deformation Imaging Based on Coherent Compounding of Diverging
Waves with Coded Excitation. *In* 2018 IEEE International Ultrasonics Symposium (IUS). IEEE, pp
1-4

Zheng E, Wu C (2015) Structure from motion using structure-less resection. *In* Proceedings of the IEEE
International Conference on Computer Vision, pp 2075-2083
Zheng Z, Hey S, Jubery T, Liu H, Yang Y, Coffey L, Miao C, Sigmon B, Schnable J, Hochholdinger F (2020)
Shared genetic control of root system architecture between Zea mays and Sorghum bicolor.
Plant physiology
Zhu J, Ingram PA, Benfey PN, Elich T (2011) From lab to field, new approaches to phenotyping root
system architecture. Current opinion in plant biology **14**: 310-317
750