Light into the darkness: Unifying the known and unknown coding sequence space in microbiome analyses

Chiara Vanni[1,2], Matthew S. Schechter[1,3], Silvia G. Acinas[4], Albert Barberán[5], Pier Luigi Buttigieg[6], Emilio O. Casamayor[7], Tom O. Delmont[8], Carlos M. Duarte[9], A. Murat Eren[3,10], Robert D. Finn[11], Renzo Kottmann[1], Alex Mitchell[11], Pablo Sanchez[4], Kimmo Siren[12], Martin Steinegger[13,14], Frank Oliver Glöckner[15,16,2], Antonio Fernandez-Guerra[1,17] *

Affiliations

Microbial Genomics and Bioinformatics Research Group, Max Planck Institute for Marine Microbiology, Celsiusstraße 1, 28359, Bremen, Germany

Jacobs University Bremen, Campus Ring 1, 28759 Bremen, Germany

Department of Medicine, University of Chicago, Chicago, IL 60637, USA

Department of Marine Biology and Oceanography, Institut de Ciènces del Mar, CSIC, Barcelona, Spain.

Department of Environmental Science, University of Arizona, Tucson, 85721 AZ, USA

Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Am Handelshafen 12, 27570 Bremerhaven, Germany

Center for Advanced Studies of Blanes CEAB-CSIC, Spanish Council for Research, Blanes, Spain

Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Univ Evry, Université Paris-Saclay, 91057 Evry, France

Red Sea Research Centre (RSRC) and Computational Bioscience Research Center (CBRC), King Abdullah University of Science and Technology, Thuwal 23955, Saudi Arabia

Josephine Bay Paul Center, Marine Biological Laboratory, Woods Hole, MA 02543, USA

European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI), Wellcome Genome Campus, Hinxton, Cambridge CB10 1SD, UK

Section for Evolutionary Genomics, The GLOBE Institute, University of Copenhagen, Copenhagen, Denmark

School of Biological Sciences, Seoul National University, Seoul, 08826, South Korea

Institute of Molecular Biology and Genetics, Seoul National University, Seoul, 08826, South Korea

University of Bremen, MARUM, Leobener Str. 8, 28359 Bremen, Germany
Life Sciences and Chemistry, Campus Ring 1, 28759 Bremen, Germany

16 Computing Center, Helmholtz Center for Polar and Marine Research, Am Handelshafen 12, 27570 Bremerhaven, Germany

38   17 Lundbeck GeoGenetics Centre, The Globe Institute, University of Copenhagen, 1350
39   Copenhagen, Denmark
40
41   *Corresponding author: Antonio Fernandez-Guerra, antonio.fernandez-guerra@sund.ku.dk
42

Abstract

Bridging the gap between the known and the unknown coding sequence space is one of the biggest challenges in molecular biology today. This challenge is especially extreme in microbiome analyses where between 40% and 60% of the coding sequences detected are of unknown function, and ignoring this fraction limits our understanding of microbial systems. Discarding the uncharacterized fraction is not an option anymore. Here, we present an in-depth exploration of the microbial unknown fraction through the lenses of a conceptual framework and a computational workflow we developed to unify the microbial known and unknown coding sequence space. Our approach partitions the coding sequence space in gene clusters and contextualizes them with genomic and environmental information. We analyzed 415,971,742 genes predicted from 1,749 metagenomes and 28,941 bacterial and archaeal genomes, putting into perspective the extent of the unknown fraction, its diversity, and its relevance in a genomic and environmental context. With the identification of a target gene of unknown function for antibiotic resistance, we demonstrate how a contextualized unknown coding sequence space provides a robust framework for the generation of hypotheses that can be used to augment experimental data.

Introduction

Thousands of isolate, single-cell, and metagenome-assembled genomes are guiding us towards a better understanding of how microbes shape life on Earth[1–7], thus bringing about a golden age of microbial genomics. An ever-increasing number of genomes and metagenomes are unlocking uncharted regions of microbial diversity[1,8,9], providing new perspectives on the evolution of life[10,11]. However, our rapidly growing inventories of new genes have a glaring issue: between 40% and 60% cannot be assigned to a known function[12–15]. Current analytical approaches for genomic and metagenomic data[16–20] generally do not include this uncharacterized fraction in downstream analyses, constraining their results to conserved pathways and housekeeping functions[17]. This inability to handle shades of the unknown is an immense impediment to realizing the potential for discovery of microbial genomics and microbiology at large[12,21].

Predicting function from traditional single sequence similarity appears to have yielded all it can[22–24], thus several groups have attempted to resolve gene function by other means. Such efforts include combining biochemistry and crystallography[25]; using environmental co-occurrence[26]; by grouping those genes into evolutionarily related families[27–30]; using remote homologies[31,32]; or more recently using deep learning approaches[33,34]. In 2018, Price et al.[13] developed a high-throughput experimental pipeline that provides mutant phenotypes for thousands of bacterial

genes of unknown function being one of the most promising methods to tackle the unknown.
Despite their promise, experimental methods are labor-intensive and require novel computational
methods that could bridge the existing gap between the known and unknown coding sequence
space (CDS-space).

Here we present a conceptual framework and a computational workflow that closes the gap
between the known and unknown CDS-space by connecting genomic and metagenomic gene
clusters. Our approach adds context to vast amounts of unknown biology, providing an invaluable
resource to get a better understanding of the unknown functional fraction and boost the current
methods for its experimental characterization. The application of our approach to 415,971,742
genes predicted from 1,749 metagenomes and 28,941 bacterial and archaeal genomes shows
that (1) the extent of the unknown fraction is smaller than expected, (2) that the diversity of gene
clusters in the unknown fraction is higher than in the known fraction, and (3) that the unknown
fraction is phylogenetically more conserved and is predominantly lineage-specific at the species
level. Finally, we show how we can connect all the outputs produced by our approach to augment
the results from experimental data and add context to genes of unknown function through
hypothesis-driven molecular investigations.

Results

A conceptual framework and a computational workflow to unify the known and the unknown microbial coding sequence space

We created the conceptual and technical foundations to unify the known and unknown CDS-space and provide a practical solution to one of the most significant ongoing challenges in microbiome analyses. First, we conceptually partitioned the known and unknown fractions into (1) Known with Pfam annotations (K), (2) Known without Pfam annotations (KWP), (3) Genomic unknown (GU), and (4) Environmental unknown (EU) (Fig. 1A). The framework introduces a subtle change of paradigm compared to traditional approaches where our objective is to provide the best representation of the unknown space. We gear all our efforts towards finding sequences without any evidence of known homologies by pushing the search space beyond the *twilight zone* of sequence similarity[35]. With this objective in mind, we use gene clusters (GCs) instead of genes as the fundamental unit to compartmentalize the CDS-space owing to their unique characteristics (Fig. 1B). GCs produce a structured CDS-space reducing its complexity (Fig. 1B), are independent of the known and unknown fraction, are conserved across environments and organisms, and can be used to aggregate information from different sources (Fig. 1A). Moreover, the GCs provide a good compromise in terms of resolution for analytical purposes, and owing to their unique properties, one can perform analyses at different scales. For fine-grained analyses, we can exploit the gene associations within each GC; and for coarse-grained analyses, we can create groups of GCs based on their shared homologies (Fig. 1B).

A

Known with Pfam
GC annotated to contain one or more Pfam entries but excluding DUFs

Known without Pfam
GC that have a known function but lack a Pfam annotation

Genomic unknown
GC of unknown function (DUFs are included here) and found in sequenced or draft genomes

Environmental unknown
GC of unknown function not detected in sequenced or draft genomes, but only in environmental metagenomes or MAGs

B

Figure 1: Conceptual framework to unify the known and unknown CDS-space and integration of the framework in the current analytical workflows (A) Link between the conceptual framework and the computational workflow to partition the CDS-space in the four conceptual categories. AGNOSTOS infers, validates and refines the GCs and combines them in gene cluster communities (GCCs). Then, it classifies them in one of the four conceptual categories based on their level of 'darkness'. Finally, we add context to each GC based on several sources of information, providing a robust framework for the generation of hypotheses that can be used to augment experimental data. (B) The computational workflow provides two mechanisms to structure the CDS-space using GCs, de novo creation of the GCs (*DB creation*), or integration of the dataset in an existing GC database (*DB update*). The structured CDS-space can then be plugged into traditional analytical workflows to annotate the genes within each GC of the known fraction. With AGNOSTOS, we provide the opportunity to easily integrate the unknown fraction into the current microbiome analyses. C) The versatility of the GCs enables analyses at different scales depending on the scope of our experiments. We can group GCs in gene cluster communities based on their shared homologies to perform coarse-grained analyses. On the other hand, we can design fine-grained analyses using the relationships between the genes in a GC, i.e., detecting network modules in the GC inner sequence similarity network. Additionally, the fact that GCs are conserved across environments, organisms and experimental conditions gives us access to an unprecedented amount of information to design and interpret experimental data.

Driven by the concepts defined in the conceptual framework, we developed AGNOSTOS, a computational workflow that infers, validates, refines, and classifies GCs in the four proposed categories (Fig. 1A; Fig. 1B; Supp. Fig 1). AGNOSTOS provides two operational modules (*DB creation* and *DB update*) to produce GCs with a highly conserved intra-homogeneous structure (Fig. 1B), both in terms of sequence similarity and domain architecture homogeneity; it exhausts any existing homology to known genes and provides a proper delimitation of the unknown CDS-space before classifying each GC in one of the four categories. In the last step, we decorate each GC with a rich collection of contextual data that we compile from different sources, or that we generate by analyzing the GC contents in different contexts (Fig. 1A). For each GC, we also offer several products that can be used for analytical purposes like improved representative sequences, consensus sequences, sequence profiles for MMseqs2[36] and HHblits[37], or the GC members as a sequence similarity network (see Online Methods). To complement the collection, we also provide a subset of what we define as *high-quality* GCs. The defining criteria are (1) the representative is a complete gene and (2) more than one-third of genes within a GC are complete genes.

Partitioning and contextualizing the coding sequence space of genomes and metagenomes

We used our approach to explore the unknown CDS-space of 1,749 microbial metagenomes derived from human and marine environments, and 28,941 genomes from GTDB_r86 (Supp Fig 2A).

The initial gene prediction of AGNOSTOS (Supp Fig 1) produced 322,248,552 genes from the environmental dataset and assigned Pfam annotations to 44% of them. Next, it clustered the predicted genes in 32,465,074 GCs. For the downstream processing, we kept 3,003,897 GCs (83% of the original genes) after filtering out any GC that contained less than ten genes[38] removing 9,549,853 clusters and 19,911,324 singletons (Supp Fig 2A; Supp. Note 1). The validation process selected 2,940,257 *good-quality* clusters (Fig. 1B; Supp. Table 1; Supp. Note 2), which resulted in 43% of them being members of the unknown CDS-space after the classification and remote homology refinement steps (Supp Fig 2A, Supp. Note 3).

We build the link between the environmental and genomic CDS-space by expanding the final collection of GCs with the genes predicted from GTDB_r86 (Supp Fig 2A). Our environmental GCs already included 72% of the genes from GTDB_r86; 22% of them created 2,400,037 new GCs, and the rest 6% resulted in singleton GCs (Supp Fig 2A; Supp. Note 4; Supp. Note 5). The final dataset includes 5,287,759 GCs (Supp Fig 2A), with both datasets sharing only 922,599 GCs (Supp Fig 2B). The addition of the GTDB_r86 genes increased the proportion of GCs in the unknown CDS-space to 54%. As the final step, the workflow generated a subset of 203,217 *high-quality* GCs (Supp Table 2; Supp Fig 3). In these *high-quality* clusters, we identified 12,313 clusters potentially encoding for small proteins (<= 50 amino acids). Most of these GCs are unknown (66% of them), which agrees with recent findings on novel small proteins from metagenomes[39].

The KWP category contains the largest proportion of incomplete ORFs (Supp. Table 3), impeding the detection and assignment of Pfam domains. But it also incorporates sequences with an unusual amino acid composition that has homology to proteins with high levels of disorder in the DPD database[40] and that have characteristic functions of the intrinsically disordered proteins[41] (IDP) like cellular processes and signaling as predicted by eggNOG annotations (Supp. Table 4). As part of the workflow, each GC is complemented with a rich set of information, as shown in Fig 1A (Supp. Table 5; Supp Note 6).

Beyond the twilight zone, communities of gene clusters

The method we developed to group GCs in gene cluster communities (GCCs) (Fig. 2A) reduced the final collection of GCs by 87%, producing 673,601 GCCs (Fig. 2B; Supp. Note 7). We validated the ability of our approach in capturing remote homologies between related GCs using two well-known gene families present in our environmental datasets, proteorhodopsins[42] and bacterial ribosomal proteins[43]. In our dataset, 64 GCs (12,184 genes) and 3 GCCs (Supp Note 8) contained sequences classified as proteorhodopsin (PR). One *Known* GCC contained 99% of the PR annotated genes (Fig. 2C), with the only exception of 85 genes taxonomically annotated as viral and assigned to the *PR Supercluster I*[44] enclosed in two GU communities (five GU gene clusters; Supp Note 8). For the ribosomal proteins, the results were not so satisfactory. We identified 1,843 GCs (781,579 genes) and 98 GCCs. The number of GCCs is larger compared to the expected number of ribosomal proteins families (16) used for validation. When we use *high-quality* GCs (Supp. Note 8), we get closer to the expected number of GCCs (Fig. 2D). With this subset, we identified 26 GCCs and 145 GCs (1,687 genes). The cross-validation of our method against the approach used in Méheust et al.[43] (Supp. Note 9) confirms the intrinsic complexity of analyzing metagenomic data. Both approaches showed a high agreement in the GCCs identified (Supp. Table 9-1). Still, our method inferred fewer GCCs for each of the ribosomal protein families (Supplementary Figure 9-3), coping better with the nuisances of a metagenomic setup, like incomplete genes (Supp. Table 6).

Figure 2: Overview and validation of the workflow to aggregate GCs in communities. (A) We inferred a gene cluster homology network using the results of an all-vs-all HMM gene cluster comparison with HHBLITS. The edges of the network are based on the HHblits-score/Aligned-columns. Communities are identified by an iterative screening of different MCL inflation parameters and evaluated using five different metrics that take into account the inter- and intra-community properties. (B) Comparison of the number of GCs and GCCs for each of the functional categories. (C) Validation of the GCCs inference based on the environmental genes annotated as proteorhodopsins. Ribbons in the alluvial plot are genes, and each

210   A smaller but highly diverse unknown coding sequence space

Combining clustering and remote homology searches reduce the extent of the unknown CDS-

space compared to the traditional genomic and metagenomic analysis approaches (Fig. 3A). Our

workflow recruited as much as 71% of genes in human-related metagenomic samples and 65%

of the genes in marine metagenomes into the known CDS-space. In both human and marine

microbiomes, the genomic unknown fraction showed a similar proportion of genes (21%, Fig. 3A).

The number of genes corresponding to EU gene clusters is higher in marine metagenomes; in

total, 12% of the genes are part of this GC category. We obtained a comparable result when we

evaluated the genes from the GTDB_r86, 75% of bacterial and 64% of archaeal genes were part

of the known CDS-space. Archaeal genomes contained more unknowns than those from Bacteria,

where 30% of the genes are classified as genomic unknowns in Archaea, and only 20% in

Bacteria (Fig. 3A; Supp. Table 7). We observed a similar trend when we evaluated the number of

amino acids belonging to the known and unknown CDS-space. From the 90,128,659,316 amino

acids analyzed, the majority of the amino acids in metagenomes (74%) and in GTDB_r86 (80%)

are in the known CDS-space (Fig. 3B; Supp. Table 7). In both cases, approximately 40% of the

amino acids in the known CDS-space were part of a Pfam domain (Fig. 3B; Supp. Table 7). The

proportion of amino acids in the unknown CDS-space ranged from the 22% in metagenomes and

15% in GTDB_r86. In both cases, only 2% of the amino acids in the unknown CDS-space were

covered by a Pfam domain.

To evaluate the coverage of our dataset, we calculated the accumulation rates of GCs and GCCs.

For the metagenomic dataset we used 1,264 metagenomes (18,566,675 GCs and 282,580

GCCs) and for the genomic dataset 28,941 genomes (9,586,109 GCs and 496,930 GCCs). The

rate of accumulation of unknown GCs was three times higher than the known (2 times for the

genomic), and both cases were far from reaching a plateau (Fig. 3C). This is not the case for the

GCC accumulation curves (Supp Fig 4B), where they reached a plateau. The rate of accumulation

is largely determined by the number of singletons, and especially singletons from EUs (Supp note

11 and Supp Fig 5). While the accumulation rate of known GCs between marine and human

metagenomes is almost identical, there are striking differences for the unknown GCs (Fig. 3D).

These differences are maintained even when we remove the virus-enriched samples from the

marine metagenomes (Supp Fig 4A). Although the marine metagenomes include a large variety

of environments, from coastal to the deep sea, the known space remains quite constrained.

Despite only including marine and human metagenomes in our database, our coverage to other

databases and environments is quite comprehensive, with an overall coverage of 76% (Supp.

Note 12). The lowest covered biomes are freshwater, soil and human non-digestive as revealed

by the screening of MGnify[16] (release 2018_09; 11 biomes; 843,535,6116 proteins) where we

assigned 74% of the MGnify proteins into one of our categories (Supplementary Fig. 6).

Figure 3: The extent of the known and unknown coding sequence space (A) Proportion of genes in the
known and unknown. (B) Amino acid distribution in the known and unknown CDS-space. (C) Accumulation
curves for the known and unknown CDS-space at the GC- level for the metagenomic and genomic data.
from TARA, MALASPINA, OSD2014 and HMP-I/II projects. (D) Collector curves comparing the human and
marine biomes. Colored lines represent the mean of 1000 permutations and shaded in grey the standard
deviation. Non-abundant singleton clusters were excluded from the accumulation curves calculation.

Revealing the importance of the unknown coding sequence space in marine and human environments

Although the role of the unknown fraction in the environment is still a mystery, the large number
of gene counts and abundance observed underlines its inherent ecological relevance (Fig. 4A).
In some samples, the genomic unknown fraction can account for more than 40% of the total gene
abundance observed (Fig. 4A). The environmental unknown fraction is also relevant in several
samples, where singleton GCs are the majority (Fig. 4A). We identified two metagenomes with
an unusual composition in terms of environmental unknown singletons. The marine metagenome
corresponds to a sample from Lake Faro (OSD42), a meromictic saline with a unique extreme
environment where Archaea plays an important role[45]. The HMP metagenome (SRS143565)
corresponds to a human sample from the right cubital fossa from a healthy female subject. To
understand the unusual composition of this metagenome, we should perform further analyses to
discard potential technical artifacts like sample contamination.

The ratio between the unknown and known GCs revealed that the metagenomes located at the
upper left quadrant in Fig. 4B-C are enriched in GCs of unknown function. In human
metagenomes, we can distinguish between body sites, with the gastrointestinal tract, where
microbial communities are expected to be more diverse and complex, especially enriched with
genomic unknowns. The HMP metagenomes with the largest ratio of unknowns are those
samples identified to contain crAssphages[46,47] and HPV viruses[48] (Supp. Table 8; Supp. Fig. 7).
Consistently, in marine metagenomes (Fig. 4D) we can separate between size fractions, where
the highest ratio in genomic and environmental unknowns corresponds to the ones enriched with
viruses and giant viruses.

To complement the previous findings, we performed a large-scale analysis to investigate the GC
occurrence patterns in the environment. The narrow distribution of the unknown fraction (Fig. 4D)
suggests that these GCs might provide a selective advantage and be necessary for the adaptation
to specific environmental conditions. But the pool of broadly distributed environmental unknowns
is the most interesting result. We identified traces of potential ubiquitous organisms left

uncharacterized by traditional approaches, as more than 80% of these GCs cannot be associated

with a metagenome-assembled genome (MAG) (Supp Table 9, Supp. Note 10).

Figure 4: Distribution of the unknown coding sequence space in the human and marine metagenomes
(A) Ratio between the proportion of the number of genes and their estimated abundances per cluster
category and biome. Columns represented in the facet depicts three cluster categories based on the size
of the clusters. (B) Relationship between the ratio of Genomic unknowns and Environmental unknowns in
the HMP-I/II metagenomes. Gastrointestinal tract metagenomes are enriched in Genomic unknown
coding sequences compared to the other body sites. (C) Relationship between the ratio of Genomic
unknowns and Environmental unknowns in the TARA Oceans metagenomes. Girus and virus enriched
metagenomes show a higher proportion of both unknown coding sequences (genomic and
environmental) compared to the Archaea|Bacteria enriched fractions. (D) Environmental distribution of
GCs and GCCs based on Levin's niche breadth index. We obtained the significance values after
generating 100 null gene cluster abundance matrices using the quasiswap algorithm.

The genomic unknown coding sequence space is lineage-specific

We already showed that the unknown CDS-space is habitat-specific and might be relevant for

organism adaptation. With the inclusion of the genomes from GTDB_r86, we have accessed a

phylogenomic framework to assess how conserved and exclusive is a GC within a lineage
(lineage-specifity[49]) and the clade depth where organisms share a GC (phylogenetic
conservation[50]). We identified 781,814 lineage-specific GCs and 464,923 phylogenetically
conserved (P < 0.05) GCs in Bacteria (Supp. Table 10; Supp. Note 13 for Archaea). The number
of lineage-specific GCs increases with the Relative Evolutionary Distance[11] (Fig. 5A) and
differences between the known and the unknown fraction start to be evident at the Family level.
The unknown GCs are more phylogenetically conserved than the known (Fig. 5B, p < 0.0001),
revealing the importance of the genome's uncharacterized fraction. However, this is not the case
for the lineage-specific and phylogenetically conserved GCs, where the unknown GCs are less
phylogenetically conserved (Fig. 5B), agreeing with the large number of lineage-specific GCs at
Genus and Species level. To discard the possibility that the lineage-specific GCs of unknown
function have a viral origin, we screened all GTDB_r86 genomes for prophages. We only found
37,163 lineage-specific GCs in prophage genomic regions, being 86% of them GCs of unknown
function. After unveiling the potential relevance of the GCs of unknown function in bacterial
genomes, we identified phyla in GTDB_r86 enriched with these types of clusters. A clear pattern
emerged when we partitioned the phyla based on the ratio of known to unknown GCs and vice
versa (Fig. 5D), the phyla with a larger number of MAGs are enriched in GCs of unknown function
Figure 5D. Phyla with a high proportion of non-classified GCs (those discarded during the
validation steps) contain a small number of genomes and are primarily composed of MAGs. These
groups of phyla highly enriched in unknowns and represented mainly by MAGs include newly
described phyla such as *Cand.* Riflebacteria and *Cand.* Patescibacteria[9,51,52], both with the largest
unknown to known ratio.
We demonstrate the possibility to bridge genomic and metagenomic data and simultaneously
unify the known and unknown CDS-space by integrating the new Ocean Microbial Reference
Gene Catalog[53] (OM-RGC v2) in our database. We assigned 26,170,875 genes to known GCs,
11,422,975 to genomic unknowns, 8,661,221 to environmental unknown and 520,083 were
discarded. From the 11,422,975 genes classified as genomic unknowns, we could associate
3,261,741 to a GTDB_r86 genome and we identified 113,175 as lineage-specific. The alluvial plot
in Fig. 5E depicts the new organization of the OM-RGC v2 after being integrated into our
framework, and how we can provide context to the two original types of unknowns in the OM-
RGC (those annotated as category S in eggNOG[54] and those without known homologs in the
eggNOG database[53]) that can lead to potential experimental targets at the organism level to
complement the metatranscriptomic approach proposed by Salazar et al[53].
333

Figure 5: Phylogenomic exploration of the unknown coding sequence space. (A) Distribution of the lineage-specific GCs by taxonomic level. Lineage-specific unknown GCs are more abundant in the lower taxonomic levels (genus, species). (B) Phylogenetic conservation of the known and unknown coding sequence space in 27,372 bacterial genomes from GTDB_r86. We observe differences in the conservation between the known and the unknown coding sequence space for lineage- and non-lineage specific GCs (paired Wilcoxon rank-sum test; all p-values < 0.0001). (C) The majority of the lineage-specific clusters are part of the unknown coding sequence space, being a small proportion found in prophages present in the GTDB_r86 genomes. (D) Known and unknown coding sequence space of the 27,732 GTDB_r86 bacterial genomes grouped by bacterial phyla. Phyla are partitioned based on the ratio of known to unknown GCs and vice versa. Phyla enriched in MAGs have higher proportions in GCs of unknown function. Phyla with a high proportion of non-classified clusters (NC; discarded during the validation steps) tend to contain a small number of genomes. (E) The left side of the alluvial plot shows the uncharacterized (OM-RGC v2 GC) and characterized (OM-RGC v2) fraction of the gene catalog. The functional annotation is based on the eggNOG annotations provided by Salazar et al.[53]. The right side of the alluvial plot shows the new organization of the OM-RGC v2 coding sequence space based on the approach described in this study. The treemap in the right links the metagenomic and genomic space adding context to the unknown fraction of the OM-RGC v2

Augmenting experimental data through a structured coding sequence space

We selected one of the experimental conditions tested in Price et al.[13] to demonstrate the potential
of our approach to augment experimental data. We compared the fitness values in plain rich
medium with added Spectinomycin dihydrochloride pentahydrate to the fitness in plain rich
medium (LB) in *Pseudomonas fluorescens FW300-N2C3* (Fig. 6A). This antibiotic inhibits protein
synthesis and elongation by binding to the bacterial 30S ribosomal subunit and interferes with the
peptidyl tRNA translocation. We identified the gene with locus id AO356_08590 that presents a
strong phenotype (fitness = -3.1; t = -9.1) and has no known function. This gene belongs to the
genomic unknown GC GU_19737823. We can track this GC into the environment and explore
the occurrence in the different samples we have in our database. As expected, the GC is mostly
found in non-human metagenomes (Fig. 6B) as *Pseudomonas* are common inhabitants of soil
and water environments[55].   However, finding this GC also in human-related samples is very
interesting, due to the potential association of *P. fluorescens* and human disease where Crohn's
disease patients develop serum antibodies to this microbe[56]. We can add another layer of
information to the selected GC by looking at the associated remote homologs in the GCC
GU_c_21103 (Fig. 6C). We identified all the genes in the GTDB_r86 genomes that belong to the
GCC GU_c_21103 (Supp. Table 11) and explored their genomic neighborhoods. All members
from GU_c_21103 are constrained to the class *Gammaproteobacteria*, and interestingly
GU_19737823 is mostly exclusive to the order *Pseudomonadales*. The gene order in the different
genomes analyzed is highly conserved, finding GU_19737823 after the *rpsF*::*rpsR* operon and
before *rplI*. *rpsF* and *rpsR* encode for 30S ribosomal proteins, the prime target of spectinomycin.
The combination of the experimental evidence and the associated data inferred by our approach
provides strong support to generate the hypothesis that the gene AO356_08590 might be involved
in the resistance to spectinomycin.
378

Figure 6: Augmenting experimental data with GCs of unknown function. (A) We used the fitness values from the experiments from Price et al.[13] to identify genes of unknown function that are important for fitness under certain experimental conditions. The selected gene belongs to the genomic unknown GC GU_19737823 and presents a strong phenotype (fitness = -3.1; t = -9.1) (B) Occurrence of GU_19737823 in the metagenomes used in this study. Darker bars depict the number of metagenomes where the GC is found. (C) GU_19737823 is a member of the GCC GU_c_21103. The network shows the relationships between the different GCs members of the gene cluster community GU_c_21103. The size of the node corresponds to the node degree of each GC. Edge thickness corresponds to the bitscore/column metric. Highlighted in red is GU_19737823. (D) We identified all the genes in the GTDB_r86 genomes that belong to the GCC GU_c_21103 and explored their genomic neighborhoods. GU_c_21103 members were constrained to the class Gammaproteobacteria, and GU_19737823 is mostly exclusive to the order Pseudomonadales. The gene order in the different genomes analyzed is highly conserved, finding GU_19737823 after the *rpsF::rpsR* operon and before *rpll. rpsF* and *rpsR* encode for the *30S ribosomal protein S6* and *30S ribosomal protein S18* respectively. The GTDB_r86 subtree only shows RefSeq genomes. Branch colors correspond to the different GCs found in GU_c_21103. Bubble plot depicts the number of genomes with a gene that belongs to GU_c_21103.

Discussion

We present a new conceptual framework and computational workflow to unify the known and
unknown CDS-space in microbial analyses. Using this framework, we performed an in-depth
exploration of the microbial unknown CDS-space. We demonstrated that we could link the
unknown fraction of metagenomic studies to specific genomes and provide a powerful tool for
hypothesis generation. During the last years, the microbiome community has established a
standard operating procedure[17] for analyzing metagenomes that we can briefly summarize into
(1) assembly, (2) gene prediction, (3) gene catalog inference, (4) binning, and (5) characterization.
Thanks to recent computational developments[36,57], we envisioned an alternative to this workflow
where we can maximize the information used when analyzing genomic and metagenomic data.
In addition, we provide a mechanism to reconcile top-down and bottom-up approaches, thanks to
the well-structured CDS-space proposed by our framework. AGNOSTOS can create
environmental- and organism-specific variations of a seed GC database. Then, it integrates the
predicted genes from new genomes and metagenomes and dynamically creates and classifies
new GCs with those genes not integrated during the initial step (Fig. 1B). Afterward, the potential
functions of the known GCs can be carefully characterized by incorporating them into the
traditional workflows.

One of the most appealing characteristics of our approach is that the GCs provide unified groups
of homologous genes across environments and organisms indifferently if they belong to the
known or unknown CDS-space, and we can contextualize the unknown fraction using this
genomic and environmental information. Our combination of partitioning and contextualization
features a smaller unknown CDS-space than we expected. On average, for our genomic and
metagenomic data, only 30% of the genes fall in the unknown fraction. One hypothesis to
reconcile this surprising finding is that until recently, the methodologies to identify remotely
homologous sequences in large datasets were computationally prohibitive. New methods[36,37], like
the ones used in AGNOSTOS, are enabling large scale distant homology searches. Still, one has
to apply conservative measures to control the trade-off between specificity and sensitivity to avoid
overclassification.

We found that the majority of the coding sequence space at gene and amino acid is known, both
in genomes and metagenomes. However, it presents a high diversity as shown in the GC
accumulation curves highlighting the vast remaining untapped microbial fraction and its potential
importance for niche adaptation owing to its narrow ecological distribution. In a genomic context,
the unknown fraction is predominantly species' lineage-specific and phylogenetically more

conserved than the known fraction, supporting the signal observed in the environmental data and
emphasizing that the unknown fraction should not be ignored. We also ruled out the effect of
prophages, strengthening the hypothesis that the lineage-specific GCs of unknown function might
be associated with the mechanisms of microbial diversification and niche adaptation as a result
of the constant diversification of gene families and the survival of new gene lineages[58,59]. It is
worth noting that we need to explore further the unknown fraction to identify new potential protein
domains. Only 10% of the unknown CDS-space amino acids are part of a Pfam domain (DUF and
others); this contrasts with the numbers observed in the known CDS-space, where Pfam domains
include 50% of the amino acids.

Metagenome-assembled genomes are not only unveiling new regions of the microbial universe
(42% of the genomes in GTDB_r86), but they are also enriching genes of unknown function in
the tree of life. We investigated the unknown CDS-space of *Cand*. Patescibacteria, more
commonly known as Candidate Phyla Radiation (CPR), a phylum that has raised considerable
interest due to their unusual biology[9]. We provide a collection of 54,343 lineage-specific GCs of
unknown function at different taxonomic level resolutions (Supp. Table 12; Supp. Note 14), which
will be a valuable resource for the advancement of knowledge in the CPR research efforts.

Our effort to tackle the unknown provides a pathway to unlock a large pool of likely relevant data
that remains untapped to analysis and discovery. With the identification of a potential target gene
of unknown function for antibiotic resistance, we demonstrate the value of our approach and how
it can boost insights from model organism experiments. But severe challenges remain, such as
the dependence on the quality of the assemblies and their gene predictions, as shown by the
analysis of the ribosomal protein GCCs where many of the recovered genes are incomplete. While
sequence assembly has been an active area of research[60], this has not been the case for gene
prediction methods[60], which are becoming outdated[61] and cannot cope with the current amount
of data. Alternatives like protein-level assembly[62] combined with the exploration of the assembly
graphs' neighborhoods[63] become very attractive for our purposes. In any case, we still face the
challenge of discriminating between real and artifactual singletons[64]. At the moment, there are no
methods available to provide a plausible solution and, at the same time, being scalable. We devise
a potential solution in the recent developments in unsupervised deep learning methods where
they use large corpora of proteins to define a language model *embedding* for protein sequences[65].
These models could be applied to predict *embeddings* in singletons, which could be clustered or
used to determine their coding potential. Another issue is that we might be creating more GCs
than expected. We follow a conservative approach to avoid mixing multidomain proteins in GCs
owing to the fragmented nature of the metagenome assemblies that could result in the split of a

GC. However, not only splitting can be a problem, but also lumping unrelated genes or GCs owing

to the use of remote homologies. Although the inference of GCCs is using very sensitive methods

to compare profile HMMs, low sequence diversity in GCs can limit its effectiveness. Our approach

is affected by the presence and propagation of contamination in reference databases, a significant

problem in 'omics[66,67]. In our case, we only use Pfam as a source for annotation owing to its high-

quality and manual curation process. The categorization process of our GCs depends on the

information from other databases, and to minimize the potential impact of contamination, we apply

methods that weight the annotations of the identified homologs to discriminate if a GC belongs to

the known or unknown CDS-space. We foresee the integration of our approach to assist in the

manual curation process and increase the quality of the recovered MAGs[68].

The work presented here should incentivize the scientific community to build a collective effort to

define the different levels of unknown[69] where clear guidelines and protocols should be

established. Our work proves that the integration of the unknown fraction is possible and aims to

provide a new brighter future for microbiome analyses.

Material and methods

Genomic and metagenomic dataset

We used a set of 583 marine metagenomes from four of the major metagenomic surveys of the

ocean microbiome: Tara Oceans expedition (TARA)[2], Malaspina expedition[70], Ocean Sampling

Day (OSD)[3], and Global Ocean Sampling Expedition (GOS)[71]. We complemented this set with

1,246 metagenomes obtained from the Human Microbiome Project (HMP) phase I and II[72]. We

used the assemblies provided by TARA, Malaspina, OSD and HMP projects and the long Sanger

reads from GOS[73]. A total of 156M (156,422,969) contigs and 12.8M long-reads were collected

(Supp. Table 6).

For the genomic dataset, we used the 28,941 prokaryotic genomes (27,372 bacterial and 1,569

archaeal) from the Genome Taxonomy Database[11] (GTDB) Release 03-RS86 (19th August

2018).

Computational workflow development

We implemented a computation workflow based on Snakemake[74] for the easy processing of large

datasets in a reproducible manner. The workflow provides three different strategies to analyze

the data. The module *DB-creation* creates the gene cluster database, validates and partitions the
gene clusters (GCs) in the main functional categories. The module *DB-update* allows the
integration of new sequences (either at the contig or predicted gene level) in the existing gene
cluster database. In addition, the workflow has a *profile-search* function to quickly screen samples
using the gene cluster PSSM profiles in the database.

Metagenomic and genomic gene prediction

499   We used Prodigal (v2.6.3)[75] in metagenomic mode to predict the genes from the metagenomic
500   dataset. For the genomic dataset, we used the gene predictions provided by Annotree[76], since
501   they were obtained, consistently, with Prodigal v2.6.3. We identified potential spurious genes
502   using the *AntiFam* database[77]. Furthermore, we screened for '*shadow' genes* using the procedure
503   described in Yooseph et al.[78].

PFAM annotation

505   We annotated the predicted genes using the *hmmsearch* program from the *HMMER* package
506   (version: 3.1b2)[79] in combination with the Pfam database v31[80]. We kept the matches exceeding
507   the internal gathering threshold and presenting an independent e-value < 1e-5 and coverage >
508   0.4. In addition, we took into account multi-domain annotations, and we removed overlapping
509   annotations when the overlap is larger than 50%, keeping the ones with the smaller e-value.

Determination of the gene clusters

We clustered the metagenomic predicted genes using the cascaded-clustering workflow of the
MMseqs2 software[57] ("*--cov-mode 0 -c 0.8 --min-seq-id 0.3*"). We discarded from downstream
analyses the singletons and clusters with a size below a threshold identified after applying a
broken-stick model[81]. We integrated the genomic data into the metagenomic cluster database
using the "DB-update" module of the workflow. This module uses the *clusterupdate* module of
MMseqs2[36], with the same parameters used for the metagenomic clustering.

Quality-screening of gene clusters

We examined the GCs to ensure their high intra-cluster homogeneity. We applied two
methodologies to validate their cluster sequence composition and functional annotation
homogeneity. We identified non-homologous sequences inside each cluster combining the

identification of a new cluster representative sequence via a sequence similarity network (SSN)

analysis, and the investigation of intra-cluster multiple sequence alignments (MSAs), given the

new representative. Initially, we generated an SSN for each cluster, using the semi-global

alignment methods implemented in *PARASAIL*[82] (version 2.1.5). We trimmed the SSN using a

custom algorithm[83,84] that removes edges while maintaining the network structural integrity and

obtaining the smallest connected graph formed by a single component. Finally, the new cluster

representative was identified as the most central node of the trimmed SSN by the eigenvector

centrality algorithm, as implemented in igraph[85]. After this step, we built a multiple sequence

alignment for each cluster using *FAMSA*[86] (version 1.1). Then, we screened each cluster-MSA

for non-homologous sequences to the new cluster representative. Owing to computational

limitations, we used two different approaches to evaluate the cluster-MSAs. We used *LEON-BIS*[87]

for the clusters with a size ranging from 10 to 1,000 genes and OD-SEQ[88] for the clusters with

more than 1,000 genes. In the end, we applied a broken-stick model[81] to determine the threshold

to discard a cluster.

The predicted genes can have multi-domain annotations in different orders, therefore to validate

the consistency of intra-cluster Pfam annotations, we applied a combination of w-shingling[89] and

Jaccard similarity. We used w-shingling (k-shingle = 2) to group consecutive domain annotations

as a single object. We measured the homogeneity of the *shingle sets* (sets of domains) between

genes using the Jaccard similarity and reported the median similarity value for each cluster.

Moreover, we took into consideration the Clan membership of the Pfam domains and that a gene

might contain N-, C- and M-terminal domains for the functional homogeneity validation. We

discarded clusters with a median similarity < 1.

After the validation, we refined the gene cluster database removing the clusters identified to be

discarded and the clusters containing ≥ 30% *shadow genes*. Lastly, we removed the single

shadow, spurious and non-homologous genes from the remaining clusters (Supplementary Note

2).

547   Remote homology classification of gene clusters

To partition the validated GCs into the four main categories, we processed the set of GCs

containing Pfam annotated genes and the set of not annotated GCs separately. For the annotated

GCs, we inferred a consensus protein domain architecture (DA) (an ordered combination of

protein domains) for each annotated gene cluster. To identify each gene cluster consensus DA,

we created directed acyclic graphs connecting the Pfam domains based on their topological order

on the genes using *igraph*[85]. We collapsed the repetitions of the same domain. Then we used the

gene completeness as a positive-weighting value for the selection of the cluster consensus DA.
Within this step, we divided the GCs into "Knowns" (Known) if annotated to at least one Pfam
domains of known function (DKFs) and "Genomic unknowns" (GU) if annotated entirely to Pfam
domains of unknown function (DUFs).

We aligned the sequences of the non-annotated GCs with FAMSA[86] and obtained cluster
consensus sequences with the *hhconsensus* program from *HH-SUITE*[37]. We used the cluster
consensus sequences to perform a nested search against the UniRef90 database (release
2017_11)[90] and NCBI *nr* database (release 2017_12)[91] to retrieve non-Pfam annotations with
MMSeqs2[36] ("*-e 1e-05 --cov-mode 2 -c 0.6*"). We kept the hits within 60% of the Log(best-e-value)
and searched the annotations for any of the terms commonly used to define proteins of unknown
function (Supp. Table 12). We used a quorum majority voting approach to decide if a gene cluster
would be classified as *Genomic Unknown* or *Known without Pfams* based on the annotations
retrieved. We searched the consensus sequences without any homologs in the UniRef90
database against NCBI *nr*. We applied the same approach and criteria described for the first
search. Ultimately, we classified as *Environmental Unknown* those GCs whose consensus
sequences did not align with any of the NCBI *nr* entries.

In addition, we developed some conservative measures to control the trade-off between specificity
and sensitivity for the remote homology searches such as (1) a modification of the algorithm
described in Hingamp et al.[92] to get a confident group of homologs to determine if a query protein
is known or unknown by a quorum majority voting approach (Supp Note 3); (2) strict parameters
in terms of iterations, bidirectional coverage and probability thresholds for the HHblits alignments
to minimize the inclusion of non-homologous sequences; and (3) avoid providing annotations for
our gene clusters, as we believe that annotation should be a careful process done on a smaller
scale and with experimental context.

Gene cluster remote homology refinement

We refined the *Environmental Unknown* GCs to ensure the lack of any characterization by
searching for remote homologies in the Uniclust database (release 30_2017_10) using the
HMM/HMM alignment method *HHblits*[93]. We created the HMM profiles with the *hhmake* program
from the *HH-SUITE*[37]. We only accepted those hits with an *HHblits-probability* ≥ 90% and we re-
classified them following the same majority vote approach as previously described. The clusters
with no hits remained as the refined set of EUs. We applied a similar refinement approach to the
KWP clusters to identify GCs with remote homologies to Pfam protein domains. The KWP HMM
profiles were searched against the Pfam *HH-SUITE* database (version 31), using *HHblits*. We

accepted hits with a probability ≥ 90% and a target coverage > 60% and removed overlapping
domains as described earlier. We moved the KWP with remote homologies to known Pfams to
the Known set, and those showing remote homologies to Pfam DUFs to the GUs. The clusters
with no hits remained as the refined set of KWP.

Gene cluster characterization

To retrieve the taxonomic composition of our clusters we applied the *MMseqs2 taxonomy* program
(version: b43de8b7559a3b45c8e5e9e02cb3023dd339231a), which allows computing the lowest
common ancestor through the implementation of the 2bLCA protocol [92]. We searched all cluster
genes against UniProtKB (release of January 2018) [94] using the following parameters "*-e 1e-05 -*
-cov-mode 0 -c 0.6". We parsed the results to keep only the hits within 60% of the log10(best-e-
value). To retrieve the taxonomic lineages, we used the R package *CHNOSZ*[95]. We measured
the intra-cluster taxonomic admixture by applying the *entropy.empirical()* function from the *entropy*
R package[96]. This function estimates the Shannon entropy based on the different taxonomic
annotation frequencies. For each cluster, we also retrieved the cluster consensus taxonomic
annotation, which we defined as the taxonomic annotation of the majority of the genes in the
cluster.
In addition to the taxonomy, we evaluated the clusters' level of darkness and disorder using the
Dark Proteome Database (DPD)[40] as reference. We searched the cluster genes against the DPD,
applying the MMseqs2 search program[36] with "*-e 1e-20 --cov-mode 0 -c 0.6*". For each cluster,
we then retrieved the mean and the median level of darkness, based on the gene DPD
annotations.

High-quality clusters

609 We defined a subset of high-quality clusters based on the completeness of the cluster genes and
610 their representatives*.* We identified the minimum required percentage of complete genes per
611 cluster by a broken-stick model[81] applied to the percentage distribution. Then, we selected the
612 GCs found above the threshold and with a complete representative.

A set of non-redundant domain architectures

We estimated the number of potential domain architectures present in the *Known* GCs taking into
account the large proportion of fragmented genes in the metagenomic dataset and that could
inflate the number of potential domain architectures. To identify fragments of larger domain

architecture, we took into account their topological order in the genes. To reduce the number of

comparisons, we calculated the pairwise string cosine distance (q-gram = 3) between domain

architectures and discarded the pairs that were too divergent (cosine distance ≥ 0.9). We

collapsed a fragmented domain architecture to the larger one when it contained less than 75% of

complete genes.

622   Inference of gene cluster communities

We aggregated distant homologous GCs into GCCs. The community inference approach

combined an all-vs-all HMM gene cluster comparison with Markov Cluster Algorithm (MCL)[97]

community identification. We started performing the inference on the Known GCs to use the Pfam

DAs as constraints. We aligned the gene cluster HMMs using HHblits[93] (-n 2 -Z 10000000 -B

10000000 -e 1) and we built a homology graph using the cluster pairs with probability ≥ 50% and

bidirectional coverage > 60%. We used the ratio between HHblits-bitscore and aligned-columns

as the edge weights (Supp. Note 9). We used MCL[97] (v. 12-068) to identify the communities

present in the graph. We developed an iterative method to determine the optimal MCL inflation

parameter that tries to maximize the relationship of five intra-/inter-community properties: (1) the

proportion of MCL communities with one single DA, based on the consensus DAs of the cluster

members; (2) the ratio of MCL communities with more than one cluster; (3) the proportion of MCL

communities with a PFAM clan entropy equal to 0; (4) the intra-community HHblits-score/Aligned-

columns score (normalized by the maximum value); and (5) the number of MCL communities,

which should, in the end, reflect the number of non-redundant DAs. We iterated through values

ranging from 1.2 to 3.0, with incremental steps of 0.1. During the inference process, some of the

GCs became orphans in the graph. We applied a three-step approach to assigning a community

membership to these GCs. First, we used less stringent conditions (probability ≥ 50% and

coverage >= 40%) to find homologs in the already existing GCCs. Then, we ran a second iteration

to find secondary relationships between the newly assigned GCs and the missing ones. Lastly,

we created new communities with the remaining GCs. We repeated the whole process with the

other categories (KWP, GU and EU), applying the optimal inflation value found for the Known (2.2

for metagenomic and 2.5 for genomic data).

645   Gene cluster communities validation

We tested the biological significance of the GCCs using the phylogeny of proteorhodopsin[44] (PR).

We used the proteorhodopsin HMM profiles[42] to screen the marine metagenomic datasets using

hmmsearch (version 3.1b2)[79]. We kept the hits with a coverage > 0.4 and e-value <= 1e-5. We

removed identical duplicates from the sequences assigned to PR with CD-HIT[98] (v4.6) and

cleaned from sequences with less than 100 amino acids. To place the identified PR sequences

into the MicRhode[44] PR tree first, we optimized the initial tree parameters and branch lengths with

RAxML (v8.2.12)[99]. We used PaPaRA (v2.5)[100] to incrementally align the query PR sequences

against the MicRhode PR reference alignment and *pplacer*[101] (v1.1.alpha19-0-g807f6f3) to place

the sequences into the tree. Finally, we assigned the query PR sequences to the MicRhode PR

Superclusters based on the phylogenetic placement. We further investigated the GCs annotated

as viral (196 genes, 14 GC) comparing them to the six newly discovered viral PRs[102] using

Parasail[82] (-a sg_stats_scan_sse2_128_16 -t 8 -c 1 -x). As an additional evaluation, we

investigated the distributions of standard GCCs and HQ GCCs within ribosomal protein families.

We obtained the ribosomal proteins used for the analysis combining the set of 16 ribosomal

proteins from Méheust et al.[43] and those contained in the collection of bacterial single-copy genes

of Anvi'o[103]. Also, for the ribosomal proteins, we compared the outcome of our method to the one

proposed by Méheust et al.[43] (Supp. Note 9).

Metagenomic sample selection for downstream analyses

664   For the subsequent ecological analyses we selected those metagenomes with a number of genes

665   larger or equal to the first quartile of the distribution of all the metagenomic gene counts. (Supp.

666   Table 13).

Gene cluster abundance profiles in genomes and metagenomes

We estimated abundance profiles for the metagenomic cluster categories using the read coverage

to each predicted gene as a proxy for abundance. We calculated the coverage by mapping the

reads against the assembly contigs using the *bwa-mem* algorithm from *BWA mapper*[104]. Then,

we used *BEDTOOLS*[105], to find the intersection of the gene coordinates to the assemblies, and

normalize the per-base coverage by the length of the gene. We calculated the cluster abundance

in a sample as the sum of the cluster gene abundances in that sample, and the cluster category

abundance in a sample as the sum of the cluster abundances. We obtained the proportions of

the different gene cluster categories applying a total-sum-scaling normalization. For the genomic

abundance profiles, we used the number of genes in the genomes and normalized by the total

gene counts per genome.

Rate of genomic and metagenomic gene clusters accumulation

We calculated the cumulative number of known and unknown GCs as a function of the number of metagenomes and genomes. For each metagenome count, we generated 1000 random sets, and we calculated the number of GCs and GCCs recovered. For this analysis, we used 1,246 HMP metagenomes and 358 marine metagenomes (242 from TARA and 116 from Malaspina). We repeated the same procedure for the genomic dataset. We removed the singletons from the metagenomic dataset with an abundance smaller than the mode abundance of the singletons that got reclassified as good-quality clusters after integrating the GTDB data to minimize the impact of potential spurious singletons. To complement those analyses, we evaluated the coverage of our dataset by searching seven different state-of-the-art databases against our set of metagenomic GC HMM profiles (Supp. Note 12).

Occurrence of gene clusters in the environment

We used 1,264 metagenomes from the TARA Oceans, MALASPINA Expedition, OSD2014 and HMP-I/II to explore the properties of the unknown CDS-space in the environment. We applied the Levins Niche Breadth (NB) index[106] to investigate the GCs and GCCs environmental distributions. We removed the GCs and cluster communities with a mean relative abundance < 1e-5. We followed a divide-and-conquer strategy to avoid the computational burden of generating the null-models to test the significance of the distributions owing to the large number of metagenomes and GCs. First, we grouped similar samples based on the gene cluster content using the Bray-Curtis dissimilarity[107] in combination with the *Dynamic Tree Cut*[108] R package. We created 100 random datasets picking up one random sample from each group. For each of the 100 random datasets, we created 100 random abundance matrices using the *nullmodel* function of the *quasiswap* count method[109]. Then we calculated the *observed* NB and obtained the 2.5% and 97.5% quantiles based on the randomized sets. We compared the observed and quantile values for each gene cluster and defined it to have a *Narrow distribution* when the *observed* was smaller than the 2.5% quantile and to have a *Broad distribution* when it was larger than the 97.5% quantile. Otherwise, we classified the cluster as *Non-significant*[110]. We used a majority voting approach to get a consensus distribution classification based on the ten random datasets.

Identification of prophages in genomic sequences

We used PhageBoost (https://github.com/ku-cbd/PhageBoost/) to find gene regions in the microbial genomes that result in high viral signals against the overall genome signal. We set the following thresholds to consider a region prophage: minimum of 10 genes, maximum 5 gaps, single-gene probability threshold 0.9. We further smoothed the predictions using Parzen rolling windows of 20 periods and looked at the smoothed probability distribution across the genome. We disregarded regions that had a summed smoothed probability less than 0.5, and those regions that did differ from the overall population of the genes in a genome by using Kruskal–Wallis rank test (p-value 0.001).

Lineage-specific gene clusters

We used the F1-score developed for AnnoTree[76] to identify the lineage-specific GCs and to which rank they are specific. Following similar criteria to the ones used in Mendler et al.[76], we considered a gene cluster to be lineage-specific if it is present in less than half of all genomes and at least 2 with F1-score > 0.95.

Phylogenetic conservation of gene clusters

We calculated the phylogenetic conservation (τD) of each gene cluster using the *consenTRAIT*[50] function implemented in the R package *castor*[50]. We used a paired Wilcoxon rank-sum test to compare the average τD values for lineage-specific and non-specific GCs.

Evaluation of the OM-RGC v2 uncharacterized fraction

We integrated the 46,775,154 genes from the second version of the TARA Ocean Microbial Reference Gene Catalog (OM-RGC v2)[53] into our cluster database using the same procedure as for the genomic data. We evaluated the uncharacterized fraction and the genes classified into the eggNOG[54] category S within the context of our database.

Augmenting experimental data

We searched the 37,684 genes of unknown function associated with mutant phenotypes from Price et al.[13] against our gene cluster profiles. We kept the hits with e-value ≤ 1e-20 and a query coverage > 60%. Then we filtered the results to keep the hits within 90% of the Log(best-e-value), and we used a majority vote function to retrieve the consensus category for each hit. Lastly, we

735 selected the best-hits based on the smallest e-value and the largest query and target coverage
736 values. We used the fitness values from the RB-TnSeq experiments from Price et al. to identify
737 genes of unknown function that are important for fitness under certain experimental conditions.

Code and data availability

The code used for the analyses in the manuscript is available at https://github.com/functional-
dark-side/functional-dark-side.github.io/tree/master/scripts. The code to recreate the figures is
available at https://github.com/functional-dark-side/vanni_et_al-figures. Detailed descriptions of
the different methods and results of this manuscript are available at
https://dark.metagenomics.eu. The workflow AGNOSTOS is available at
https://github.com/functional-dark-side/agnostos-wf, and its database can be downloaded from
https://doi.org/10.6084/m9.figshare.12459056.

746
747

Acknowledgments

The authors thankfully acknowledge the computer resources at MareNostrum and the technical
support provided by Barcelona Supercomputing Center (RES-AECT-2014-2-0085), the BMBF-
funded de.NBI Cloud within the German Network for Bioinformatics Infrastructure (de.NBI)
(031A537B, 031A533A, 031A538A, 031A533B, 031A535A, 031A537C, 031A534A, 031A532B),
the University of Oxford Advanced Research Computing
(http://dx.doi.org/10.5281/zenodo.22558) and the MARBITS bioinformatics core at ICM-CSIC. CV
was supported by the Max Planck Society. AFG received funding from the European Union's
Horizon 2020 research and innovation program Blue Growth: Unlocking the potential of Seas and
Oceans under grant agreement no. 634486 (project acronym INMARE). AM was supported by
the Biotechnology and Biological Sciences Research Council [BB/M011755/1, BB/R015228/1]
and RDF by the European Molecular Biology Laboratory core funds. EOC was supported by
project INTERACTOMA RTI2018-101205-B-I00 from the Spanish Agency of Science MICIU/AEI.
SGA and PS received additional funding by the project MAGGY (CTM2017-87736-R) from the
Spanish Ministry of Economy and Competitiveness. The Malaspina 2010 Expedition was
supported by the Spanish Ministry of Economy and Competitiveness (MINECO) through the

Consolider-Ingenio program (ref. CSD2008-00077). The authors thank Johannes Söding and
Alex Bateman for helpful discussions.
766

Author Contributions

CV, MSS and AF-G performed the analyses and wrote the computational workflow. MS assisted
with the clustering and remote homology searches. KS helped with the identification of prophages
in genomic sequences. PLB and AB provided feedback and assisted with the ecological analyses.
RDF and AM provided feedback and information on the MGnify and Pfam databases. CMD, PS
and SGA provided the Malaspina metagenomes. TOD and AME analyzed data in the context of
metagenome-assembled genomes. AF-G conceived the study and supervised the work. CV and
AF-G wrote the manuscript. All authors read, edited and approved the final manuscript.
775
776

777 # Competing Interests

778 The authors declare no competing interests.
779
780

References

1. Hug, L. A. *et al.* A new view of the tree of life. *Nat Microbiol* **1**, 16048 (2016).

2. Sunagawa, S. *et al.* Ocean plankton. Structure and function of the global ocean microbiome. *Science* **348**, 1261359 (2015).

3. Kopf, A. *et al.* The ocean sampling day consortium. *Gigascience* **4**, 27 (2015).

4. Almeida, A. *et al.* A new genomic blueprint of the human gut microbiota. *Nature* **568**, 499–504 (2019).

5. Pasolli, E. *et al.* Extensive Unexplored Human Microbiome Diversity Revealed by Over 150,000 Genomes from Metagenomes Spanning Age, Geography, and Lifestyle. *Cell* **176**, 649-662.e20 (2019).

6. Pachiadaki, M. G. *et al.* Charting the Complexity of the Marine Microbiome through Single-Cell Genomics. *Cell* **179**, 1623-1635.e11 (2019).

7. Cross, K. L. *et al.* Targeted isolation and cultivation of uncultivated bacteria by reverse genomics. *Nat. Biotechnol.* **37**, 1314–1321 (2019).

8. Eloe-Fadrosh, E. A. *et al.* Global metagenomic survey reveals a new bacterial candidate phylum in geothermal springs. *Nat. Commun.* **7**, 10476 (2016).

9. Brown, C. T. *et al.* Unusual biology across a group comprising more than 15% of domain Bacteria. *Nature* **523**, 208–211 (2015).

10. Spang, A. *et al.* Complex archaea that bridge the gap between prokaryotes and eukaryotes. *Nature* **521**, 173–179 (2015).

11. Parks, D. H. *et al.* A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. *Nat. Biotechnol.* **36**, 996–1004 (2018).

12. Bernard, G., Pathmanathan, J. S., Lannes, R., Lopez, P. & Bapteste, E. Microbial Dark Matter Investigations: How Microbial Studies Transform Biological Knowledge and

808  Empirically Sketch a Logic of Scientific Discovery. *Genome Biol. Evol.* **10**, 707–715 (2018).

809 13. Price, M. N. *et al.* Mutant phenotypes for thousands of bacterial genes of unknown function.

810  *Nature* **557**, 503–509 (2018).

811 14. Carradec, Q. *et al.* A global ocean atlas of eukaryotic genes. *Nat. Commun.* **9**, 373 (2018).

812 15. Almeida, A. *et al.* A unified catalog of 204,938 reference genomes from the human gut

813  microbiome. *Nat. Biotechnol.* (2020) doi:10.1038/s41587-020-0603-3.

814 16. Mitchell, A. L. *et al.* MGnify: the microbiome analysis resource in 2020. *Nucleic Acids Res.*

815  **48**, D570–D578 (2020).

816 17. Quince, C., Walker, A. W., Simpson, J. T., Loman, N. J. & Segata, N. Shotgun

817  metagenomics, from sampling to analysis. *Nat. Biotechnol.* **35**, 833–844 (2017).

818 18. Franzosa, E. A. *et al.* Species-level functional profiling of metagenomes and

819  metatranscriptomes. *Nat. Methods* **15**, 962–968 (2018).

820 19. Huerta-Cepas, J. *et al.* Fast Genome-Wide Functional Annotation through Orthology

821  Assignment by eggNOG-Mapper. *Mol. Biol. Evol.* **34**, 2115–2122 (2017).

822 20. Chen, I.-M. A. *et al.* IMG/M v.5.0: an integrated data management and comparative

823  analysis system for microbial genomes and microbiomes. *Nucleic Acids Res.* **47**, D666–

824  D677 (2019).

825 21. Hanson, A. D., Pribat, A., Waller, J. C. & Crécy-Lagard, V. de. 'Unknown'proteins and

826  'orphan'enzymes: the missing half of the engineering parts list--and how to find it. *Biochem.*

827  *J* **425**, 1–11 (2010).

828 22. Arnold, F. H. Design by Directed Evolution. *Acc. Chem. Res.* **31**, 125–131 (1998).

829 23. Brandenberg, O. F., Fasan, R. & Arnold, F. H. Exploiting and engineering hemoproteins for

830  abiological carbene and nitrene transfer reactions. *Curr. Opin. Biotechnol.* **47**, 102–111

831  (2017).

832 24. Arnold, F. H. Directed Evolution: Bringing New Chemistry to Life. *Angew. Chem. Int. Ed*

833  *Engl.* **57**, 4143–4148 (2018).

25. Jaroszewski, L. *et al.* Exploration of uncharted regions of the protein universe. *PLoS Biol.* **7**,

(2009).

26. Buttigieg, L. P. *et al.* Ecogenomic Perspectives on Domains of Unknown Function:

Correlation-Based Exploration of Marine Metagenomes. *PLoS One* **8**, (2013).

27. Yooseph, S. *et al.* The Sorcerer II global ocean sampling expedition: Expanding the

universe of protein families. *PLoS Biol.* **5**, 0432–0466 (2007).

28. Wyman, S. K., Avila-Herrera, A., Nayfach, S. & Pollard, K. S. A most wanted list of

conserved microbial protein families with no known domains. *PLoS One* **13**, e0205749

(2018).

29. Brum, J. R. *et al.* Illuminating structural proteins in viral "dark matter" with metaproteomics.

Proc. Natl. Acad. Sci. U. S. A. **113**, 2436–2441 (2016).

30. Bateman, A., Coggill, P. & Finn, D. R. DUFs: Families in search of function. *Acta*

Crystallogr. Sect. F Struct. Biol. Cryst. Commun. **66**, 1148–1152 (2010).

31. Lobb, B., Kurtz, D. A., Moreno-Hagelsieb, G. & Doxey, A. C. Remote homology and the

functions of metagenomic dark matter. *Front. Genet.* **6**, 1–12 (2015).

32. Bitard-Feildel, T. & Callebaut, I. Exploring the dark foldable proteome by considering

hydrophobic amino acids topology. *Sci. Rep.* **7**, 41425 (2017).

33. Bileschi, M. L. *et al.* Using Deep Learning to Annotate the Protein Universe. *bioRxiv* 626507

(2019) doi:10.1101/626507.

34. Liu, X. L. Deep Recurrent Neural Network for Protein Function Prediction from Sequence.

bioRxiv 103994 (2017) doi:10.1101/103994.

35. Rost, B. Twilight zone of protein sequence alignments. *Protein Eng. Des. Sel.* **12**, 85–94

(1999).

36. Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for the

analysis of massive data sets. *Nat. Biotechnol.* **advance on**, (2017).

37. Steinegger, M. *et al.* HH-suite3 for fast remote homology detection and deep protein

annotation. *BMC Bioinformatics* **20**, 473 (2019).

38. Skewes-Cox, P., Sharpton, T. J., Pollard, K. S. & DeRisi, J. L. Profile hidden Markov

models for the detection of viruses within metagenomic sequence data. *PLoS One* **9**,

e105067 (2014).

39. Sberro, H. *et al.* Large-Scale Analyses of Human Microbiomes Reveal Thousands of Small,

Novel Genes. *Cell* **178**, 1245-1259.e14 (2019).

40. Perdigão, N., Rosa, A. C. & O'Donoghue, S. I. The Dark Proteome Database. *BioData Min.*

10, 1–11 (2017).

41. Habchi, J., Tompa, P., Longhi, S. & Uversky, V. N. Introducing protein intrinsic disorder.

Chem. Rev. **114**, 6561–6588 (2014).

42. Olson, D. K., Yoshizawa, S., Boeuf, D., Iwasaki, W. & DeLong, E. F. Proteorhodopsin

variability and distribution in the North Pacific Subtropical Gyre. *ISME J.* **12**, 1047–1060

(2018).

43. Méheust, R., Burstein, D., Castelle, C. J. & Banfield, J. F. The distinction of CPR bacteria

from other bacteria based on protein family content. *Nat. Commun.* **10**, 4173 (2019).

44. Boeuf, D., Audic, S., Brillet-Guéguen, L., Caron, C. & Jeanthon, C. MicRhoDE: a curated

database for the analysis of microbial rhodopsin diversity and evolution. *Database* **2015**,

(2015).

45. La Cono, V. *et al.* Partaking of Archaea to biogeochemical cycling in oxygen-deficient

zones of meromictic saline Lake Faro (Messina, Italy). *Environ. Microbiol.* **15**, 1717–1733

(2013).

46. Edwards, R. A. *et al.* Global phylogeography and ancient evolution of the widespread

human gut virus crAssphage. *Nat Microbiol* **4**, 1727–1736 (2019).

47. Dubinkina, V. B., Ischenko, D. S., Ulyantsev, V. I., Tyakht, A. V. & Alexeev, D. G.

Assessment of k-mer spectrum applicability for metagenomic dissimilarity analysis. *BMC*

Bioinformatics vol. 17 (2016).

48. Ma, Y. *et al.* Human papillomavirus community in healthy persons, defined by

metagenomics analysis of human microbiome project shotgun sequencing data sets. *J.*

Virol. **88**, 4786–4797 (2014).

49. Mendler, K. *et al.* AnnoTree: visualization and exploration of a functionally annotated

microbial tree of life. *Nucleic Acids Res.* **47**, 4442–4448 (2019).

50. Martiny, A. C., Treseder, K. & Pusch, G. Phylogenetic conservatism of functional traits in

microorganisms. *ISME J.* **7**, 830–838 (2013).

51. Rinke, C. *et al.* Insights into the phylogeny and coding potential of microbial dark matter.

Nature **499**, 431–437 (2013).

52. Anantharaman, K. *et al.* Expanded diversity of microbial groups that shape the dissimilatory

sulfur cycle. *ISME J.* **12**, 1715–1728 (2018).

53. Salazar, G. *et al.* Gene Expression Changes and Community Turnover Differentially Shape

the Global Ocean Metatranscriptome. *Cell* **179**, 1068-1083.e21 (2019).

54. Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and phylogenetically

annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids*

Res. **47**, D309–D314 (2019).

55. Heffernan, B., Murphy, C. D. & Casey, E. Comparison of planktonic and biofilm cultures of

Pseudomonas fluorescens DSM 8341 cells grown on fluoroacetate. *Appl. Environ.*

Microbiol. **75**, 2899–2907 (2009).

56. Scales, B. S., Dickson, R. P., LiPuma, J. J. & Huffnagle, G. B. Microbiology, genomics, and

clinical significance of the Pseudomonas fluorescens species complex, an unappreciated

colonizer of humans. *Clin. Microbiol. Rev.* **27**, 927–948 (2014).

57. Steinegger, M. & Söding, J. Clustering huge protein sequence sets in linear time. *Nat.*

Commun. **9**, 2542 (2018).

58. Francino, M. P. The ecology of bacterial genes and the survival of the new. *Int. J. Evol.*

Biol. **2012**, 394026 (2012).

59. Muller, E. E. L. Determining Microbial Niche Breadth in the Environment for Better

Ecosystem Fate Predictions. *mSystems* **4**, (2019).

60. Roumpeka, D. D., Wallace, R. J., Escalettes, F., Fotheringham, I. & Watson, M. A Review

of Bioinformatics Tools for Bio-Prospecting from Metagenomic Sequence Data. *Front.*

Genet. **8**, 23 (2017).

61. Ivanova, N. N. *et al.* Stop codon reassignments in the wild. *Science* **344**, 909–913 (2014).

62. Steinegger, M., Mirdita, M. & Söding, J. Protein-level assembly increases protein sequence

recovery from metagenomic samples manyfold. *Nat. Methods* **16**, 603–606 (2019).

63. Titus Brown, C. *et al.* Exploring neighborhoods in large metagenome assembly graphs

reveals hidden sequence diversity. *bioRxiv* 462788 (2018) doi:10.1101/462788.

64. Höps, W., Jeffryes, M. & Bateman, A. Gene Unprediction with Spurio: A tool to identify

spurious protein sequences. *F1000Res.* **7**, 261 (2018).

65. Heinzinger, M. *et al.* Modeling aspects of the language of life through transfer-learning

protein sequences. *BMC Bioinformatics* **20**, 723 (2019).

66. Breitwieser, F. P., Pertea, M., Zimin, A. & Salzberg, S. L. Human contamination in bacterial

genomes has created thousands of spurious proteins. *Genome Res.* (2019)

doi:10.1101/gr.245373.118.

67. Steinegger, M. & Salzberg, S. L. Terminating contamination: large-scale search identifies

more than 2,000,000 contaminated entries in GenBank. *Genome Biol.* **21**, 115 (2020).

68. Chen, L.-X., Anantharaman, K., Shaiber, A., Eren, A. M. & Banfield, J. F. Accurate and

complete genomes from metagenomes. *Genome Res.* **30**, 315–333 (2020).

69. Thomas, A. M. & Segata, N. Multiple levels of the unknown in microbiome research. *BMC*

Biol. **17**, 48 (2019).

70. Duarte, C. M. Seafaring in the 21St Century: The Malaspina 2010 Circumnavigation

Expedition. *Limnol. Oceanog. Bull.* **24**, 11–14 (2015).

71. Rusch, D. B. *et al.* The Sorcerer II Global Ocean Sampling Expedition: Northwest Atlantic

through Eastern Tropical Pacific. *PLoS Biol.* **5**, 1–34 (2007).

72. Lloyd-Price, J. *et al.* Strains, functions and dynamics in the expanded Human Microbiome

Project. *Nature* **550**, 61–66 (2017).

73. Sanger, F., Nicklen, S. & Coulson, A. R. DNA sequencing with chain-terminating inhibitors.

Proc. Natl. Acad. Sci. U. S. A. **74**, 5463–5467 (1977).

74. Köster, J. Reproducible data analysis with Snakemake. *F1000Res.* **7**, (2018).

75. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* **11**, 119–119 (2010).

76. Mendler, K. *et al.* AnnoTree: visualization and exploration of a functionally annotated

microbial tree of life. *Nucleic Acids Res.* **47**, 4442–4448 (2019).

77. Eberhardt, R. Y. *et al.* AntiFam: a tool to help identify spurious ORFs in protein annotation.

Database **2012**, bas003–bas003 (2012).

78. Yooseph, S., Li, W. & Sutton, G. Gene identification and protein classification in microbial

metagenomic sequence data via incremental clustering. *BMC Bioinformatics* **9**, 1–13

(2008).

79. Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence similarity

searching. *Nucleic Acids Res.* **39**, W29–W37 (2011).

80. Finn, R. D. *et al.* The Pfam protein families database: towards a more sustainable future.

Nucleic Acids Res. **44**, D279–D285 (2016).

81. Bennett, K. D. Determination of the number of zones in a biostratigraphical sequence. *New*

Phytol. **132**, 155–170 (1996).

82. Daily, J. Parasail: SIMD C library for global, semi-global, and local pairwise sequence

alignments. *BMC Bioinformatics* **17**, 81–81 (2016).

83. Žure, M., Fernandez-Guerra, A., Munn, C. B. & Harder, J. Geographic distribution at

subspecies resolution level: closely related Rhodopirellula species in European coastal

sediments. *ISME J.* **11**, 478–489 (2017).

84. Chafee, M. *et al.* Recurrent patterns of microdiversity in a temperate coastal marine

environment. *ISME J.* **12**, 237–252 (2018).

85. Csardi, G. & Nepusz, T. The igraph software package for complex network research.

InterJournal vol. Complex Systems 1695 (2006).

86. Deorowicz, S., Debudaj-Grabysz, A. & Gudyś, A. FAMSA: Fast and accurate multiple

sequence alignment of huge protein families. *Sci. Rep.* **6**, 33964–33964 (2016).

87. Vanhoutreve, R. *et al.* LEON-BIS: multiple alignment evaluation of sequence neighbours

using a Bayesian inference system. *BMC Bioinformatics* **17**, 271–271 (2016).

88. Jehl, P., Sievers, F. & Higgins, D. G. OD-seq: outlier detection in multiple sequence

alignments. *BMC Bioinformatics* **16**, 269–269 (2015).

89. Broder, A. Z. On the resemblance and containment of documents. in *Proceedings.*

Compression and Complexity of SEQUENCES 1997 (Cat. No. 97TB100171) 21–29 (IEEE,

1997).

90. The UniProt Consortium. UniProt: the universal protein knowledgebase. *Nucleic Acids Res.*

45, D158–D169 (2017).

91. NCBI Resource Coordinators. Database resources of the National Center for Biotechnology

Information. *Nucleic Acids Res.* **46**, D8–D13 (2018).

92. Hingamp, P. *et al.* Exploring nucleo-cytoplasmic large DNA viruses in Tara Oceans

microbial metagenomes. *ISME J.* **7**, 1678–1695 (2013).

93. Remmert, M., Biegert, A., Hauser, A. & Söding, J. HHblits: Lightning-fast iterative protein

sequence searching by HMM-HMM alignment. *Nat. Methods* **9**, 173–175 (2012).

94. UniProt Consortium, T. UniProt: the universal protein knowledgebase. *Nucleic Acids Res.*

46, 2699 (2018).

95. Dick, J. M. Calculation of the relative metastabilities of proteins using the CHNOSZ

software package. *Geochem. Trans.* **9**, 10 (2008).

96. Hausser, J. & Strimmer, K. Entropy inference and the James-Stein estimator, with

application to nonlinear gene association networks. *arXiv [stat.ML]* (2008).

97. van Dongen, S. & Abreu-Goodger, C. Using MCL to Extract Clusters from Networks. in

Bacterial Molecular Networks: Methods and Protocols (eds. van Helden, J., Toussaint, A. &

Thieffry, D.) 281–295 (Springer New York, 2012).

98. Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of protein

or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

99. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

100.Berger, S. A. & Stamatakis, A. PaPaRa 2.0: a vectorized algorithm for probabilistic

phylogeny-aware alignment extension. *Heidelberg Institute for Theoretical Studies,*

http://sco. h-its. org/exelixis/publica tions. html. Exelixis-RRDR-2012-2015 (2012).

101.Matsen, F. A., Kodner, R. B. & Armbrust, E. V. pplacer: linear time maximum-likelihood and

Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*

Bioinformatics **11**, 538 (2010).

102.Needham, D. M. *et al.* A distinct lineage of giant viruses brings a rhodopsin photosystem to

unicellular marine predators. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 20574–20583 (2019).

103.Murat Eren, A. *et al.* Anvi'o: an advanced analysis and visualization platform for 'omics

data. *PeerJ* **3**, e1319 (2015).

104.Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows–Wheeler transform.

Bioinformatics **26**, 589–595 (2010).

105.Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841–842 (2010).

106.Levins, R. THE STRATEGY OF MODEL BUILDING IN POPULATION BIOLOGY. *Am. Sci.*

54, 421–431 (1966).

107.Bray, J. R., Roger Bray, J. & Curtis, J. T. An Ordination of the Upland Forest Communities

of Southern Wisconsin. *Ecological Monographs* vol. 27 325–349 (1957).

108.Langfelder, P., Zhang, B. & Horvath, S. Defining clusters from a hierarchical cluster tree:

the Dynamic Tree Cut package for R. *Bioinformatics* **24**, 719–720 (2008).

109.Miklós, I. & Podani, J. RANDOMIZATION OF PRESENCE–ABSENCE MATRICES:

COMMENTS AND NEW ALGORITHMS. *Ecology* vol. 85 86–92 (2004).

110.Salazar, G. *et al.* Particle-association lifestyle is a phylogenetically conserved trait in

bathypelagic prokaryotes. *Mol. Ecol.* **24**, 5692–5706 (2015).