Unifying the global coding sequence space enables the study of genes with unknown function across biomes

Chiara Vanni[1,2], Matthew S. Schechter[1,3], Silvia G. Acinas[4], Albert Barberán[5], Pier Luigi Buttigieg[6], Emilio O. Casamayor[7], Tom O. Delmont[8], Carlos M. Duarte[9], A. Murat Eren[3,10], Robert D. Finn[11], Renzo Kottmann[1], Alex Mitchell[11], Pablo Sanchez[4], Kimmo Siren[12], Martin Steinegger[13,14], Frank Oliver Glöckner[15,16,2], Antonio Fernandez-Guerra[1,17] *

Affiliations

Microbial Genomics and Bioinformatics Research Group, Max Planck Institute for Marine Microbiology, Celsiusstraße 1, 28359, Bremen, Germany

Jacobs University Bremen, Campus Ring 1, 28759 Bremen, Germany

Department of Medicine, University of Chicago, Chicago, IL 60637, USA

Department of Marine Biology and Oceanography, Institut de Ciènces del Mar, CSIC, Barcelona, Spain.

Department of Environmental Science, University of Arizona, Tucson, 85721 AZ, USA

Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Am Handelshafen 12, 27570 Bremerhaven, Germany

Center for Advanced Studies of Blanes CEAB-CSIC, Spanish Council for Research, Blanes, Spain

8 Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Univ Evry,

Université Paris-Saclay, 91057 Evry, France

9 Red Sea Research Centre (RSRC) and Computational Bioscience Research Center (CBRC),

King Abdullah University of Science and Technology, Thuwal 23955, Saudi Arabia

10 Josephine Bay Paul Center, Marine Biological Laboratory, Woods Hole, MA 02543, USA

11 European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI),

Wellcome Genome Campus, Hinxton, Cambridge CB10 1SD, UK

12 Section for Evolutionary Genomics, The GLOBE Institute, University of Copenhagen,

Copenhagen, Denmark

13 School of Biological Sciences, Seoul National University, Seoul, 08826, South Korea

14 Institute of Molecular Biology and Genetics, Seoul National University, Seoul, 08826, South

Korea

15 University of Bremen, MARUM, Leobener Str. 8, 28359 Bremen, Germany

Life Sciences and Chemistry, Campus Ring 1, 28759 Bremen, Germany

16 Computing Center, Helmholtz Center for Polar and Marine Research, Am Handelshafen 12,

27570 Bremerhaven, Germany

17 Lundbeck GeoGenetics Centre, The Globe Institute, University of Copenhagen, 1350

Copenhagen, Denmark

*Corresponding author: Antonio Fernandez-Guerra, antonio.fernandez-guerra@sund.ku.dk

Abstract

One of the biggest challenges in molecular biology is bridging the gap between the known and the unknown coding sequence space. This challenge is especially extreme in microbial systems, where between 40% and 60% of the predicted genes are of unknown function. Discarding this uncharacterized fraction should not be an option anymore. Here, we present a conceptual framework and a computational workflow that bridges this gap and provides a powerful strategy to contextualize the investigations of genes of unknown function. Our approach partitions the coding sequence space removing the known-unknown dichotomy, unifies genomic and metagenomic data and provides a framework to expand those investigations across environments and organisms. By analyzing 415,971,742 genes predicted from 1,749 metagenomes and 28,941 bacterial and archaeal genomes we showcase our approach and its application in ecological, evolutionary and biotechnological investigations. As a result, we put into perspective the extent of the unknown fraction, its diversity, and its relevance in genomic and environmental contexts. By identifying a target gene of unknown function for antibiotic resistance, we demonstrate how a contextualized unknown coding sequence space enables the generation of hypotheses that can be used to augment experimental data.

Main

Thousands of isolate, single-cell, and metagenome-assembled genomes are guiding us towards a better understanding of life on Earth[1–8]. At the same time, the ever-increasing number of genomes and metagenomes unlocking uncharted regions of life's diversity[1,9,10] are providing new perspectives on the evolution of life[11,12]. However, our rapidly growing inventories of new genes have a glaring issue: between 40% and 60% cannot be assigned to a known function[13–16]. Current analytical approaches for genomic and metagenomic data[17–21] generally do not include this uncharacterized fraction in downstream analyses, constraining their results to

conserved pathways and housekeeping functions[18]. This inability to handle shades of the

unknown is an immense impediment to realizing the potential for discovery of microbiology and

molecular biology at large[13,22]. Predicting function from traditional single sequence similarity

appears to have yielded all it can[23–25], thus several groups have attempted to resolve gene

function by other means. Such efforts include combining biochemistry and crystallography[26];

using environmental co-occurrence[27]; by grouping those genes into evolutionarily related

families[28–31]; using remote homologies[32,33]; or more recently using deep learning

approaches[34,35]. In 2018, Price et al.[14] developed a high-throughput experimental pipeline that

provides mutant phenotypes for thousands of bacterial genes of unknown function being one of

the most promising methods to tackle the unknown. Despite their promise, experimental

methods are labor-intensive and require novel computational methods that could bridge the

existing gap between the known and unknown coding sequence space (CDS-space).

Here we present a conceptual framework and a computational workflow that closes the gap

between the known and unknown CDS-space by connecting genomic and metagenomic gene

clusters. Our approach adds context to vast amounts of unknown biology, providing an

invaluable resource to get a better understanding of the unknown functional fraction and boost

the current methods for its experimental characterization. The application of our approach to

415,971,742 genes predicted from 1,749 metagenomes and 28,941 bacterial and archaeal

genomes revealed that the unknown fraction (1) is smaller than expected, (2) is exceptionally

diverse, and (3) is phylogenetically more conserved and predominantly lineage-specific at the

Species level. Finally, we show how we can connect all the outputs produced by our approach

to augment the results from experimental data and add context to genes of unknown function

through hypothesis-driven molecular investigations.

Results

A conceptual framework and a computational workflow to unify the known and the unknown coding sequence space

We created the conceptual and technical foundations to unify the known and unknown CDS-space facilitating the integration of the genes of unknown function into ecological, evolutionary and biotechnological investigations. First, we conceptually partitioned the known and unknown fractions into (1) Known with Pfam annotations (K), (2) Known without Pfam annotations (KWP), (3) Genomic unknown (GU), and (4) Environmental unknown (EU) (Fig. 1A). The framework introduces a subtle change of paradigm compared to traditional approaches where our objective is to provide the best representation of the unknown space. We gear all our efforts towards finding sequences without any evidence of known homologies by pushing the search space beyond the *twilight zone* of sequence similarity[36]. With this objective in mind, we use gene clusters (GCs) instead of genes as the fundamental unit to compartmentalize the CDS-space owing to their unique characteristics (Fig. 1B). (1) GCs produce a structured CDS-space reducing its complexity (Fig. 1B), (2) are independent of the known and unknown fraction, (3) are conserved across environments and organisms, and (4) can be used to aggregate information from different sources (Fig. 1A). Moreover, the GCs provide a good compromise in terms of resolution for analytical purposes, and owing to their unique properties, one can perform analyses at different scales. For fine-grained analyses, we can exploit the gene associations within each GC; and for coarse-grained analyses, we can create groups of GCs based on their shared homologies (Fig. 1B).

Figure 1: Conceptual framework to unify the known and unknown CDS-space and integration of the
framework in the current analytical workflows (A) Link between the conceptual framework and the
computational workflow to partition the CDS-space in the four conceptual categories. AGNOSTOS infers,
validates and refines the GCs and combines them in gene cluster communities (GCCs). Then, it classifies
them in one of the four conceptual categories based on their level of 'darkness'. Finally, we add context to
each GC based on several sources of information, providing a robust framework for the generation of
hypotheses that can be used to augment experimental data. (B) The computational workflow provides two
mechanisms to structure the CDS-space using GCs, de novo creation of the GCs (*DB creation*), or
integration of the dataset in an existing GC database (*DB update*). The structured CDS-space can then
be plugged into traditional analytical workflows to annotate the genes within each GC of the known
fraction. With AGNOSTOS, we provide the opportunity to easily integrate the unknown fraction into the
current microbiome analyses. C) The versatility of the GCs enables analyses at different scales
depending on the scope of our experiments. We can group GCs in gene cluster communities based on
their shared homologies to perform coarse-grained analyses. On the other hand, we can design fine-
grained analyses using the relationships between the genes in a GC, i.e., detecting network modules in
the GC inner sequence similarity network. Additionally, the fact that GCs are conserved across
environments, organisms and experimental conditions gives us access to an unprecedented amount of
information to design and interpret experimental data.

Driven by the concepts defined in the conceptual framework, we developed AGNOSTOS, a

computational workflow that infers, validates, refines, and classifies GCs in the four proposed

categories (Fig. 1A; Fig. 1B; Supp. Fig 1). AGNOSTOS provides two operational modules (*DB*

creation and *DB update*) to produce GCs with a highly conserved intra-homogeneous structure

(Fig. 1B), both in terms of sequence similarity and domain architecture homogeneity; it exhausts

any existing homology to known genes and provides a proper delimitation of the unknown CDS-

space before classifying each GC in one of the four categories (Methods). In the last step, we

decorate each GC with a rich collection of contextual data that we compile from different

sources, or that we generate by analyzing the GC contents in different contexts (Fig. 1A). For

each GC, we also offer several products that can be used for analytical purposes like improved

representative sequences, consensus sequences, sequence profiles for MMseqs2[37] and

HHblits[38], or the GC members as a sequence similarity network (Methods). To complement the

collection, we also provide a subset of what we define as *high-quality* GCs. The defining criteria

are (1) the representative is a complete gene and (2) more than one-third of genes within a GC

are complete genes.

Partitioning and contextualizing the coding sequence space of genomes and metagenomes

We used our approach to explore the unknown CDS-space of 1,749 microbial metagenomes derived from human and marine environments, and 28,941 genomes from GTDB_r86 (Supp Fig 2A). The initial gene prediction of AGNOSTOS (Supp Fig 1) produced 322,248,552 genes from the environmental dataset and assigned Pfam annotations to 44% of them. Next, it clustered the predicted genes in 32,465,074 GCs. For the downstream processing, we kept 3,003,897 GCs (83% of the original genes) after filtering out any GC that contained less than ten genes[39] removing 9,549,853 clusters and 19,911,324 singletons (Supp Fig 2A; Supp. Note 1). The validation process selected 2,940,257 *good-quality* clusters (Fig. 1B; Supp. Table 1; Supp. Note 2), which resulted in 43% of them being members of the unknown CDS-space after the classification and remote homology refinement steps (Supp Fig 2A, Supp. Note 3). We build the link between the environmental and genomic CDS-space by expanding the final collection of GCs with the genes predicted from GTDB_r86 (Supp Fig 2A). Our environmental GCs already included 72% of the genes from GTDB_r86; 22% of them created 2,400,037 new GCs, and the rest 6% resulted in singleton GCs (Supp Fig 2A; Supp. Note 4; Supp. Note 5). The final dataset includes 5,287,759 GCs (Supp Fig 2A), with both datasets sharing only 922,599 GCs (Supp Fig 2B). The addition of the GTDB_r86 genes increased the proportion of GCs in the unknown CDS-space to 54%. As the final step, the workflow generated a subset of 203,217 *high-quality* GCs (Supp Table 2; Supp Fig 3). In these *high-quality* clusters, we identified 12,313 clusters potentially encoding for small proteins (<= 50 amino acids). Most of these GCs are unknown (66% of them), which agrees with recent findings on novel small proteins from metagenomes[40]. The KWP category contains the largest proportion of incomplete ORFs (Supp. Table 3), disrupting the detection and assignment of Pfam domains. But it also incorporates sequences with an unusual amino acid composition that has homology to proteins with high levels of

disorder in the DPD database[41] and that have characteristic functions of the intrinsically

disordered proteins[42] (IDP) like cellular processes and signaling as predicted by eggNOG

annotations (Supp. Table 4).

As part of the workflow, each GC is complemented with a rich set of information, as shown in

Fig 1A (Supp. Table 5; Supp Note 6).

Beyond the twilight zone, communities of gene clusters

The method we developed to group GCs in gene cluster communities (GCCs) (Fig. 2A) reduced

the final collection of GCs by 87%, producing 673,601 GCCs (Methods; Fig. 2B; Supp. Note 7).

We validated the ability of our approach in capturing remote homologies between related GCs

using two well-known gene families present in our environmental datasets, proteorhodopsins[43]

and bacterial ribosomal proteins[44]. Our dataset contained 64 GCs (12,184 genes) and 3 GCCs

(Supp Note 8) classified as proteorhodopsin (PR). One *Known* GCC contained 99% of the PR

annotated genes (Fig. 2C), with the only exception of 85 genes taxonomically annotated as viral

and assigned to the *PR Supercluster I*[45] within two GU communities (five GU gene clusters;

Supp Note 8). For the ribosomal proteins, the results were not so satisfactory. We identified

1,843 GCs (781,579 genes) and 98 GCCs. The number of GCCs is larger compared to the

expected number of ribosomal proteins families (16) used for validation. When we use *high-*

quality GCs (Supp. Note 8), we get closer to the expected number of GCCs (Fig. 2D). With this

subset, we identified 26 GCCs and 145 GCs (1,687 genes). The cross-validation of our method

against the approach used in Méheust et al.[44] (Supp. Note 9) confirms the intrinsic complexity of

analyzing metagenomic data. Both approaches showed a high agreement in the GCCs

identified (Supp. Table 9-1). Still, our method inferred fewer GCCs for each of the ribosomal

protein families (Supplementary Figure 9-3), coping better with the nuisances of a metagenomic

setup, like incomplete genes (Supp. Table 6).

Figure 2: Overview and validation of the workflow to aggregate GCs in communities. (A) We inferred a
gene cluster homology network using the results of an all-vs-all HMM gene cluster comparison with
HHBLITS. The edges of the network are based on the HHblits-score/Aligned-columns. Communities are
identified by an iterative screening of different MCL inflation parameters and evaluated using five different
metrics that take into account the inter- and intra-community properties. (B) Comparison of the number of
GCs and GCCs for each of the functional categories. (C) Validation of the GCCs inference based on the
environmental genes annotated as proteorhodopsins. Ribbons in the alluvial plot are genes, and each
stacked bar corresponds (from left to right) to the (1) gene taxonomic classification at the domain level,

A smaller but highly diverse unknown coding sequence space

Combining clustering and remote homology searches reduce the extent of the unknown CDS-

space compared to what is reported by the traditional genomic and metagenomic analysis

approaches (Fig. 3A). Our workflow recruited as much as 71% of genes in human-related

metagenomic samples and 65% of the genes in marine metagenomes into the known CDS-

space. In both human and marine microbiomes, the genomic unknown fraction showed a similar

proportion of genes (21%, Fig. 3A). The number of genes corresponding to EU gene clusters is

higher in marine metagenomes; in total, 12% of the genes are part of this GC category. We

obtained a comparable result when we evaluated the genes from the GTDB_r86, 75% of

bacterial and 64% of archaeal genes were part of the known CDS-space. Archaeal genomes

contained more unknowns than those from Bacteria, where 30% of the genes are classified as

genomic unknowns in Archaea, and only 20% in Bacteria (Fig. 3A; Supp. Table 7). Our

approach allows us to go beyond genes, and for the first time we can provide a detailed

characterization of the CDS-space at the amino acid level. From the 90,128,659,316 amino

acids analyzed, the majority of the amino acids in metagenomes (74%) and in GTDB_r86 (80%)

are in the known CDS-space (Fig. 3B; Supp. Table 7). In both cases, approximately 40% of the

amino acids in the known CDS-space were part of a Pfam domain (Fig. 3B; Supp. Table 7). The

proportion of amino acids in the unknown CDS-space ranged from the 22% in metagenomes

and 15% in GTDB_r86. In both cases, only 2% of the amino acids in the unknown CDS-space

were covered by a Pfam domain. To evaluate the differences between both fractions of the

CDS-space, we calculated the accumulation rates of GCs and GCCs. For the metagenomic

dataset we used 1,264 metagenomes (18,566,675 GCs and 282,580 GCCs) and for the

genomic dataset 28,941 genomes (9,586,109 GCs and 496,930 GCCs). The rate of

accumulation of unknown GCs was three times higher than the known (2 times for the

genomic), and both cases were far from reaching a plateau (Fig. 3C). This is not the case for

the GCC accumulation curves (Supp Fig 4B), where they reached a plateau. The rate of

accumulation is largely determined by the number of singletons, and especially singletons from

EUs (Supp note 11 and Supp Fig 5). While the accumulation rate of known GCs between

marine and human metagenomes is almost identical, there are striking differences for the

unknown GCs (Fig. 3D). These differences are maintained even when we remove the virus-

enriched samples from the marine metagenomes (Supp Fig 4A). Although the marine

metagenomes include a large variety of environments, from coastal to the deep sea, the known

space remains quite constrained.

Despite only including marine and human metagenomes in our database, our coverage to other

databases and environments is quite comprehensive, with an overall coverage of 76% (Supp.

Note 12). The lowest covered biomes are freshwater, soil and human non-digestive as revealed

by the screening of MGnify[17] (release 2018_09; 11 biomes; 843,535,6116 proteins) where we

assigned 74% of the MGnify proteins into one of our categories (Supplementary Fig. 6).

Furthermore, as a result of this evaluation, we classified 20% of the FunkFams[29] and 44% of the

unknowns used by Price et al.[14] to the known fraction (Supp. Table 12-1).

The unknown coding sequence space has a limited ecological distribution in human and marine environments

Although the role of the unknown fraction in the environment is still a mystery, the large number

of gene counts and abundance observed underlines its inherent ecological relevance (Fig. 4A).

In some metagenomes, the genomic unknown fraction can account for more than 40% of the

total gene abundance observed (Fig. 4A). The environmental unknown fraction is also relevant

in several samples, where singleton GCs are the majority (Fig. 4A). We identified two

metagenomes with an unusual composition in terms of environmental unknown singletons. The

marine metagenome corresponds to a sample from Lake Faro (OSD42), a meromictic saline

with a unique extreme environment where Archaea plays an important role[46]. The HMP

metagenome (SRS143565) corresponds to a human sample from the right cubital fossa from a

healthy female subject. To understand the unusual composition of this metagenome, we should

perform further analyses to discard potential technical artifacts like sample contamination. The

ratio between the unknown and known GCs revealed that the metagenomes located at the

upper left quadrant in Fig. 4B-C are enriched in GCs of unknown function. In human

metagenomes, we can distinguish between body sites, with the gastrointestinal tract, where

microbial communities are expected to be more diverse and complex, especially enriched with

genomic unknowns. The HMP metagenomes with the largest ratio of unknowns are those

samples identified to contain crAssphages[47,48] and HPV viruses[49] (Supp. Table 8; Supp. Fig. 7).

Consistently, in marine metagenomes (Fig. 4D) we can separate between size fractions, where

the highest ratio in genomic and environmental unknowns corresponds to the ones enriched

with viruses and giant viruses.

Figure 4: Distribution of the unknown coding sequence space in the human and marine metagenomes
(A) Ratio between the proportion of the number of genes and their estimated abundances per cluster
category and biome. Columns represented in the facet depicts three cluster categories based on the size
of the clusters. (B) Relationship between the ratio of Genomic unknowns and Environmental unknowns in
the HMP-I/II metagenomes. Gastrointestinal tract metagenomes are enriched in Genomic unknown
coding sequences compared to the other body sites. (C) Relationship between the ratio of Genomic
unknowns and Environmental unknowns in the TARA Oceans metagenomes. Girus and virus enriched
metagenomes show a higher proportion of both unknown coding sequences (genomic and
environmental) compared to the Archaea|Bacteria enriched fractions. (D) Environmental distribution of
GCs and GCCs based on Levin's niche breadth index. We obtained the significance values after
generating 100 null gene cluster abundance matrices using the *quasiswap* algorithm.

To complement the previous findings, we performed a large-scale analysis to investigate the GC

occurrence patterns in the environment. The narrow distribution of the unknown fraction (Fig.

4D) suggests that these GCs might provide a selective advantage and be necessary for the

adaptation to specific environmental conditions. But the pool of broadly distributed

environmental unknowns is the most interesting result. We identified traces of potential

ubiquitous organisms left uncharacterized by traditional approaches, as more than 80% of these

GCs cannot be associated with a metagenome-assembled genome (MAG) (Supp Table 9,

Supp. Note 10).

The genomic unknown coding sequence space is lineage-specific

With the inclusion of the genomes from GTDB_r86, we have access to a phylogenomic

framework that can be used to assess how exclusive is a GC within a lineage (lineage-

specifity[50]) and how conserved is a GC across clades (phylogenetic conservation[51]). We

identified 781,814 lineage-specific GCs and 464,923 phylogenetically conserved (P < 0.05) GCs

in Bacteria (Supp. Table 10; Supp. Note 13 for Archaea). The number of lineage-specific GCs

increases with the Relative Evolutionary Distance[12] (Fig. 5A) and differences between the

known and the unknown fraction start to be evident at the Family level resulting in 4X more

unknown lineage-specific GCs at the Species level. The unknown GCs are more

phylogenetically conserved than the known (Fig. 5B, p < 0.0001), revealing the importance of

the genome's uncharacterized fraction. However, this is not the case for the lineage-specific and

phylogenetically conserved GCs, where the unknown GCs are less phylogenetically conserved

(Fig. 5B), agreeing with the large number of lineage-specific GCs at Genus and Species level.

To discard the possibility that the lineage-specific GCs of unknown function have a viral origin,

we screened all GTDB_r86 genomes for prophages. We only found 37,163 lineage-specific

GCs in prophage genomic regions, being 86% of them GCs of unknown function. After unveiling

the potential relevance of the GCs of unknown function in bacterial genomes, we identified

phyla in GTDB_r86 enriched with these types of clusters. A clear pattern emerged when we

partitioned the phyla based on the ratio of known to unknown GCs and vice versa (Fig. 5D), the

phyla with a larger number of MAGs are enriched in GCs of unknown function Figure 5D. Phyla

with a high proportion of non-classified GCs (those discarded during the validation steps)

contain a small number of genomes and are primarily composed of MAGs. These groups of

phyla highly enriched in unknowns and represented mainly by MAGs include newly described

phyla such as *Cand.* Riflebacteria and *Cand.* Patescibacteria[10,52,53], both with the largest

unknown to known ratio.

325
326
Figure 5: Phylogenomic exploration of the unknown coding sequence space. (A) Distribution of the
lineage-specific GCs by taxonomic level. Lineage-specific unknown GCs are more abundant in the lower
taxonomic levels (Genus, Species). (B) Phylogenetic conservation of the known and unknown coding
sequence space in 27,372 bacterial genomes from GTDB_r86. We observe differences in the
conservation between the known and the unknown coding sequence space for lineage- and non-lineage
specific GCs (paired Wilcoxon rank-sum test; all p-values < 0.0001). (C) The majority of the lineage-
specific clusters are part of the unknown coding sequence space, being a small proportion found in
prophages present in the GTDB_r86 genomes. (D) Known and unknown coding sequence space of the
27,732 GTDB_r86 bacterial genomes grouped by bacterial phyla. Phyla are partitioned based on the ratio
of known to unknown GCs and vice versa. Phyla enriched in MAGs have higher proportions in GCs of
unknown function. Phyla with a high proportion of non-classified clusters (NC; discarded during the
validation steps) tend to contain a small number of genomes. (E) The left side of the alluvial plot shows
the uncharacterized (OM-RGC v2 GC) and characterized (OM-RGC v2) fraction of the gene catalog. The
functional annotation is based on the eggNOG annotations provided by Salazar et al.[54]. The right side of

We demonstrate the possibility to bridge genomic and metagenomic data and simultaneously

unify the known and unknown CDS-space by integrating the new Ocean Microbial Reference

Gene Catalog[54] (OM-RGC v2) in our database. We assigned 26,170,875 genes to known GCs,

11,422,975 to genomic unknowns, 8,661,221 to environmental unknown and 520,083 were

discarded. From the 11,422,975 genes classified as genomic unknowns, we could associate

3,261,741 to a GTDB_r86 genome and we identified 113,175 as lineage-specific. The alluvial

plot in Fig. 5E depicts the new organization of the OM-RGC v2 after being integrated into our

framework, and how we can provide context to the two original types of unknowns in the OM-

RGC (those annotated as category S in eggNOG[55] and those without known homologs in the

eggNOG database[54]) that can lead to potential experimental targets at the organism level to

complement the metatranscriptomic approach proposed by Salazar et al[54]..

A structured coding sequence space augments the interpretation of experimental data

We selected one of the experimental conditions tested in Price et al.[14] to demonstrate the

potential of our approach to augment experimental data. We compared the fitness values in

plain rich medium with added Spectinomycin dihydrochloride pentahydrate to the fitness in plain

rich medium (LB) in *Pseudomonas fluorescens FW300-N2C3* (Fig. 6A). This antibiotic inhibits

protein synthesis and elongation by binding to the bacterial 30S ribosomal subunit and

interferes with the peptidyl tRNA translocation. We identified the gene with locus id

AO356_08590 that presents a strong phenotype (fitness = -3.1; t = -9.1) and has no known

function. This gene belongs to the genomic unknown GC GU_19737823. We can track this GC

into the environment and explore the occurrence in the different samples we have in our

database. As expected, the GC is mostly found in non-human metagenomes (Fig. 6B) as

Pseudomonas are common inhabitants of soil and water environments[56]. However, finding this

GC also in human-related samples is very interesting, due to the potential association of *P.*

fluorescens and human disease where Crohn's disease patients develop serum antibodies to

this microbe[57]. We can add another layer of information to the selected GC by looking at the

associated remote homologs in the GCC GU_c_21103 (Fig. 6C). We identified all the genes in

the GTDB_r86 genomes that belong to the GCC GU_c_21103 (Supp. Table 11) and explored

their genomic neighborhoods. All members from GU_c_21103 are constrained to the class

Gammaproteobacteria, and interestingly GU_19737823 is mostly exclusive to the order

Pseudomonadales. The gene order in the different genomes analyzed is highly conserved,

finding GU_19737823 after the *rpsF*::*rpsR* operon and before *rplI*. *rpsF* and *rpsR* encode for

30S ribosomal proteins, the prime target of spectinomycin. The combination of the experimental

evidence and the associated data inferred by our approach provides strong support to generate

the hypothesis that the gene AO356_08590 might be involved in the resistance to

spectinomycin.

Figure 6: Augmenting experimental data with GCs of unknown function. (A) We used the fitness values from the experiments from Price et al.[14] to identify genes of unknown function that are important for fitness under certain experimental conditions. The selected gene belongs to the genomic unknown GC GU_19737823 and presents a strong phenotype (fitness = -3.1; t = -9.1) (B) Occurrence of GU_19737823 in the metagenomes used in this study. Darker bars depict the number of metagenomes where the GC is found. (C) GU_19737823 is a member of the GCC GU_c_21103. The network shows the relationships between the different GCs members of the gene cluster community GU_c_21103. The size of the node corresponds to the node degree of each GC. Edge thickness corresponds to the bitscore/column metric. Highlighted in red is GU_19737823. (D) We identified all the genes in the GTDB_r86 genomes that belong to the GCC GU_c_21103 and explored their genomic neighborhoods. GU_c_21103 members were constrained to the class Gammaproteobacteria, and GU_19737823 is mostly exclusive to the order Pseudomonadales. The gene order in the different genomes analyzed is highly conserved, finding GU_19737823 after the *rpsF::rpsR* operon and before *rplI*. *rpsF* and *rpsR* encode for the *30S ribosomal protein S6* and *30S ribosomal protein S18,* respectively. The GTDB_r86 subtree only shows RefSeq

Discussion

We present a new conceptual framework and computational workflow to unify the known and

unknown CDS-space. Using this framework, we performed an in-depth exploration of the

microbial unknown CDS-space, demonstrating that we can link the unknown fraction of

metagenomic studies to specific genomes and provide a powerful tool for hypothesis

generation. During the last years, the microbiome community has established a standard

operating procedure[18] for analyzing metagenomes that we can briefly summarize into (1)

assembly, (2) gene prediction, (3) gene catalog inference, (4) binning, and (5) characterization.

Thanks to recent computational developments[37,58], we envisioned an alternative to this workflow

to maximize the information used when analyzing genomic and metagenomic data. In addition,

we provide a mechanism to reconcile top-down and bottom-up approaches, thanks to the well-

structured CDS-space proposed by our framework. AGNOSTOS can create environmental- and

organism-specific variations of a seed GC database. Then, it integrates the predicted genes

from new genomes and metagenomes and dynamically creates and classifies new GCs with

those genes not integrated during the initial step (Fig. 1B). Afterward, the potential functions of

the known GCs can be carefully characterized by incorporating them into the traditional

workflows.

One of the most appealing characteristics of our approach is that the GCs provide unified

groups of homologous genes across environments and organisms indifferently if they belong to

the known or unknown CDS-space, and we can contextualize the unknown fraction using this

genomic and environmental information. Our combination of partitioning and contextualization

features a smaller unknown CDS-space than we expected. On average, for our genomic and

metagenomic data, only 30% of the genes fall in the unknown fraction. One hypothesis to

reconcile this surprising finding is that the methodologies to identify remotely homologous

sequences in large datasets were computationally prohibitive until recently. New methods[37,38],

like the ones used in AGNOSTOS, are enabling large scale distant homology searches. Still,

one has to apply conservative measures to control the trade-off between specificity and

sensitivity to avoid overclassification.

We found that most of the coding sequence space at the gene and amino acid level is known,

both in genomes and metagenomes. However, it presents a high diversity, as shown in the GC

accumulation curves highlighting the vast remaining untapped microbial fraction and its potential

importance for niche adaptation owing to its narrow ecological distribution. In a genomic context

and after ruling out the effect of prophages, the unknown fraction is predominantly Species'

lineage-specific and phylogenetically more conserved than the known fraction, supporting the

signal observed in the environmental data emphasizing that we should not ignore the unknown

fraction. It is worth noting that the high diversity observed in the unknowns only represents the

20% of the amino acids in the CDS-space we analyzed, and only 10% of this unknown amino

acid space is part of a Pfam domain (DUF and others). This contrasts with the numbers

observed in the known CDS-space, where Pfam domains include 50% of the amino acids. All

this evidence combined strengthens the hypothesis that the genes of unknown function,

especially the lineage-specific ones, might be associated with the mechanisms of microbial

diversification and niche adaptation due to the constant diversification of gene families and the

survival of new gene lineages[59,60].

Metagenome-assembled genomes are not only unveiling new regions of the microbial universe

(42% of the genomes in GTDB_r86), but they are also enriching genes of unknown function in

the tree of life. We investigated the unknown CDS-space of *Cand*. Patescibacteria, more

commonly known as Candidate Phyla Radiation (CPR), a phylum that has raised considerable

interest due to its unusual biology[10]. We provide a collection of 54,343 lineage-specific GCs of

unknown function at different taxonomic level resolutions (Supp. Table 12; Supp. Note 14),

which will be a valuable resource for the CPR advancement research efforts.

Our effort to tackle the unknown provides a pathway to unlock a large pool of likely relevant data

that remains untapped to analysis and discovery. By identifying a potential target gene of

unknown function for antibiotic resistance, we demonstrate the value of our approach and how it

can boost insights from model organism experiments. But severe challenges remain, such as

the dependence on the quality of the assemblies and their gene predictions, as shown by the

analysis of the ribosomal protein GCCs where many of the recovered genes are incomplete.

While sequence assembly has been an active area of research[61], this has not been the case for

gene prediction methods[61], which are becoming outdated[62] and cannot cope with the current

amount of data. Alternatives like protein-level assembly[63] combined with exploring the assembly

graphs' neighborhoods[64] become very attractive for our purposes. In any case, we still face the

challenge of discriminating between real and artifactual singletons[65]. Currently, there are no

methods available to provide a plausible solution and, at the same time, being scalable. We

devise a potential solution in the recent developments in unsupervised deep learning methods

where they use large corpora of proteins to define a language model *embedding* for protein

sequences[66]. These models could be applied to predict *embeddings* in singletons, which could

be clustered or used to determine their coding potential. Another issue is that we might be

creating more GCs than expected. We follow a conservative approach to avoid mixing multi-

domain proteins in GCs owing to the fragmented nature of the metagenome assemblies that

could result in the split of a GC. However, not only splitting can be a problem, but also lumping

unrelated genes or GCs owing to the use of remote homologies. Although the inference of

GCCs is using very sensitive methods to compare profile HMMs, low sequence diversity in GCs

can limit its effectiveness. Moreover, our approach is affected by the presence and propagation

of contamination in reference databases, a significant problem in 'omics[67,68]. In our case, we

only use Pfam as a source for annotation owing to its high-quality and manual curation process.

The categorization process of our GCs depends on the information from other databases, and

to minimize the potential impact of contamination, we apply methods that weight the annotations

of the identified homologs to discriminate if a GC belongs to the known or unknown CDS-space.

The results presented here prove that the integration and the analysis of the unknown fraction

are possible. We are unveiling a brighter future, not only for microbiome analyses but also for

boosting eukaryotic-related studies, thanks to the increasing number of projects including

metatranscriptomic data[8,69]. Furthermore, our work lays the foundations for further

developments of clear guidelines and protocols to define the different levels of unknown[70] and

should encourage the scientific community for a collaborative effort to tackle this challenge.

Material and methods

Genomic and metagenomic dataset

We used a set of 583 marine metagenomes from four of the major metagenomic surveys of the

ocean microbiome: Tara Oceans expedition (TARA)[2], Malaspina expedition[71], Ocean Sampling

Day (OSD)[3], and Global Ocean Sampling Expedition (GOS)[72]. We complemented this set with

1,246 metagenomes obtained from the Human Microbiome Project (HMP) phase I and II[73]. We

used the assemblies provided by TARA, Malaspina, OSD and HMP projects and the long

Sanger reads from GOS[74]. A total of 156M (156,422,969) contigs and 12.8M long-reads were

collected (Supp. Table 6).

For the genomic dataset, we used the 28,941 prokaryotic genomes (27,372 bacterial and 1,569

archaeal) from the Genome Taxonomy Database[12] (GTDB) Release 03-RS86 (19th August

2018).

Computational workflow development

We implemented a computation workflow based on Snakemake[75] for the easy processing of

large datasets in a reproducible manner. The workflow provides three different strategies to

analyze the data. The module *DB-creation* creates the gene cluster database, validates and

partitions the gene clusters (GCs) in the main functional categories. The module *DB-update*

allows the integration of new sequences (either at the contig or predicted gene level) in the

existing gene cluster database. In addition, the workflow has a *profile-search* function to quickly

screen samples using the gene cluster PSSM profiles in the database.

Metagenomic and genomic gene prediction

506   We used Prodigal (v2.6.3)[76] in metagenomic mode to predict the genes from the metagenomic

507   dataset. For the genomic dataset, we used the gene predictions provided by Annotree[50], since

508   they were obtained, consistently, with Prodigal v2.6.3. We identified potential spurious genes

509   using the *AntiFam* database[77]. Furthermore, we screened for '*shadow*' genes using the

510   procedure described in Yooseph et al.[78].

PFAM annotation

512   We annotated the predicted genes using the *hmmsearch* program from the *HMMER* package

513   (version: 3.1b2)[79] in combination with the Pfam database v31[80]. We kept the matches exceeding

514   the internal gathering threshold and presenting an independent e-value < 1e-5 and coverage >

515   0.4. In addition, we took into account multi-domain annotations, and we removed overlapping

516   annotations when the overlap is larger than 50%, keeping the ones with the smaller e-value.

Determination of the gene clusters

We clustered the metagenomic predicted genes using the cascaded-clustering workflow of the MMseqs2 software[58] ("*--cov-mode 0 -c 0.8 --min-seq-id 0.3*"). We discarded from downstream analyses the singletons and clusters with a size below a threshold identified after applying a broken-stick model[81]. We integrated the genomic data into the metagenomic cluster database using the "DB-update" module of the workflow. This module uses the *clusterupdate* module of MMseqs2[37], with the same parameters used for the metagenomic clustering.

Quality-screening of gene clusters

We examined the GCs to ensure their high intra-cluster homogeneity. We applied two methodologies to validate their cluster sequence composition and functional annotation homogeneity. We identified non-homologous sequences inside each cluster combining the identification of a new cluster representative sequence via a sequence similarity network (SSN) analysis, and the investigation of intra-cluster multiple sequence alignments (MSAs), given the new representative. Initially, we generated an SSN for each cluster, using the semi-global alignment methods implemented in *PARASAIL*[82] (version 2.1.5). We trimmed the SSN using a custom algorithm[83,84] that removes edges while maintaining the network structural integrity and obtaining the smallest connected graph formed by a single component. Finally, the new cluster representative was identified as the most central node of the trimmed SSN by the eigenvector centrality algorithm, as implemented in igraph[85]. After this step, we built a multiple sequence alignment for each cluster using *FAMSA*[86] (version 1.1). Then, we screened each cluster-MSA for non-homologous sequences to the new cluster representative. Owing to computational limitations, we used two different approaches to evaluate the cluster-MSAs. We used *LEON-BIS*[87] for the clusters with a size ranging from 10 to 1,000 genes and OD-SEQ[88] for the clusters

with more than 1,000 genes. In the end, we applied a broken-stick model[81] to determine the

threshold to discard a cluster.

The predicted genes can have multi-domain annotations in different orders, therefore to validate

the consistency of intra-cluster Pfam annotations, we applied a combination of w-shingling[89] and

Jaccard similarity. We used w-shingling (k-shingle = 2) to group consecutive domain

annotations as a single object. We measured the homogeneity of the *shingle sets* (sets of

domains) between genes using the Jaccard similarity and reported the median similarity value

for each cluster. Moreover, we took into consideration the Clan membership of the Pfam

domains and that a gene might contain N-, C- and M-terminal domains for the functional

homogeneity validation. We discarded clusters with a median similarity < 1.

After the validation, we refined the gene cluster database removing the clusters identified to be

discarded and the clusters containing ≥ 30% *shadow genes*. Lastly, we removed the single

shadow, spurious and non-homologous genes from the remaining clusters (Supplementary Note

2).

Remote homology classification of gene clusters

To partition the validated GCs into the four main categories, we processed the set of GCs

containing Pfam annotated genes and the set of not annotated GCs separately. For the

annotated GCs, we inferred a consensus protein domain architecture (DA) (an ordered

combination of protein domains) for each annotated gene cluster. To identify each gene cluster

consensus DA, we created directed acyclic graphs connecting the Pfam domains based on their

topological order on the genes using *igraph*[85]. We collapsed the repetitions of the same domain.

Then we used the gene completeness as a positive-weighting value for the selection of the

cluster consensus DA. Within this step, we divided the GCs into "Knowns" (Known) if annotated

to at least one Pfam domains of known function (DKFs) and "Genomic unknowns" (GU) if

annotated entirely to Pfam domains of unknown function (DUFs).

565 We aligned the sequences of the non-annotated GCs with FAMSA[86] and obtained cluster

566 consensus sequences with the *hhconsensus* program from *HH-SUITE*[38]. We used the cluster

567 consensus sequences to perform a nested search against the UniRef90 database (release

568 2017_11)[90] and NCBI *nr* database (release 2017_12)[91] to retrieve non-Pfam annotations with

569 *MMSeqs2*[37] ("*-e 1e-05 --cov-mode 2 -c 0.6*"). We kept the hits within 60% of the Log(best-e-

570 value) and searched the annotations for any of the terms commonly used to define proteins of

571 unknown function (Supp. Table 12). We used a quorum majority voting approach to decide if a

572 gene cluster would be classified as *Genomic Unknown* or *Known without Pfams* based on the

573 annotations retrieved. We searched the consensus sequences without any homologs in the

574 UniRef90 database against NCBI *nr*. We applied the same approach and criteria described for

575 the first search. Ultimately, we classified as *Environmental Unknown* those GCs whose

576 consensus sequences did not align with any of the NCBI *nr* entries.

577 In addition, we developed some conservative measures to control the trade-off between

578 specificity and sensitivity for the remote homology searches such as (1) a modification of the

579 algorithm described in Hingamp et al.[92] to get a confident group of homologs to determine if a

580 query protein is known or unknown by a quorum majority voting approach (Supp Note 3); (2)

581 strict parameters in terms of iterations, bidirectional coverage and probability thresholds for the

582 HHblits alignments to minimize the inclusion of non-homologous sequences; and (3) avoid

583 providing annotations for our gene clusters, as we believe that annotation should be a careful

584 process done on a smaller scale and with experimental context.

585 Gene cluster remote homology refinement

586 We refined the *Environmental Unknown* GCs to ensure the lack of any characterization by

587 searching for remote homologies in the Uniclust database (release 30_2017_10) using the

588 HMM/HMM alignment method *HHblits*[93]. We created the HMM profiles with the *hhmake*

589 program from the *HH-SUITE*[38]. We only accepted those hits with an *HHblits-probability* ≥ 90%

and we re-classified them following the same majority vote approach as previously described.

The clusters with no hits remained as the refined set of EUs. We applied a similar refinement

approach to the KWP clusters to identify GCs with remote homologies to Pfam protein domains.

The KWP HMM profiles were searched against the Pfam *HH-SUITE* database (version 31),

using *HHblits*. We accepted hits with a probability ≥ 90% and a target coverage > 60% and

removed overlapping domains as described earlier. We moved the KWP with remote

homologies to known Pfams to the Known set, and those showing remote homologies to Pfam

DUFs to the GUs. The clusters with no hits remained as the refined set of KWP.

Gene cluster characterization

To retrieve the taxonomic composition of our clusters we applied the *MMseqs2 taxonomy*

program (version: b43de8b7559a3b45c8e5e9e02cb3023dd339231a), which allows computing

the lowest common ancestor through the implementation of the 2bLCA protocol [92]. We searched

all cluster genes against UniProtKB (release of January 2018) [94] using the following parameters

"*-e 1e-05 --cov-mode 0 -c 0.6*". We parsed the results to keep only the hits within 60% of the

log10(best-e-value). To retrieve the taxonomic lineages, we used the R package *CHNOSZ*[95].

We measured the intra-cluster taxonomic admixture by applying the *entropy.empirical()* function

from the *entropy* R package[96]. This function estimates the Shannon entropy based on the

different taxonomic annotation frequencies. For each cluster, we also retrieved the cluster

consensus taxonomic annotation, which we defined as the taxonomic annotation of the majority

of the genes in the cluster.

In addition to the taxonomy, we evaluated the clusters' level of darkness and disorder using the

Dark Proteome Database (DPD)[41] as reference. We searched the cluster genes against the

DPD, applying the MMseqs2 search program[37] with "*-e 1e-20 --cov-mode 0 -c 0.6*". For each

cluster, we then retrieved the mean and the median level of darkness, based on the gene DPD

annotations.

High-quality clusters

We defined a subset of high-quality clusters based on the completeness of the cluster genes and their representatives. We identified the minimum required percentage of complete genes per cluster by a broken-stick model[81] applied to the percentage distribution. Then, we selected the GCs found above the threshold and with a complete representative.

A set of non-redundant domain architectures

We estimated the number of potential domain architectures present in the *Known* GCs taking into account the large proportion of fragmented genes in the metagenomic dataset and that could inflate the number of potential domain architectures. To identify fragments of larger domain architecture, we took into account their topological order in the genes. To reduce the number of comparisons, we calculated the pairwise string cosine distance (q-gram = 3) between domain architectures and discarded the pairs that were too divergent (cosine distance ≥ 0.9). We collapsed a fragmented domain architecture to the larger one when it contained less than 75% of complete genes.

Inference of gene cluster communities

We aggregated distant homologous GCs into GCCs. The community inference approach combined an all-vs-all HMM gene cluster comparison with Markov Cluster Algorithm (MCL)[97] community identification. We started performing the inference on the Known GCs to use the Pfam DAs as constraints. We aligned the gene cluster HMMs using HHblits[93] (-n 2 -Z 10000000 -B 10000000 -e 1) and we built a homology graph using the cluster pairs with probability ≥ 50% and bidirectional coverage > 60%. We used the ratio between HHblits-bitscore and aligned-columns as the edge weights (Supp. Note 9). We used MCL[97] (v. 12-068) to identify the communities present in the graph. We developed an iterative method to determine the optimal

MCL inflation parameter that tries to maximize the relationship of five intra-/inter-community

properties: (1) the proportion of MCL communities with one single DA, based on the consensus

DAs of the cluster members; (2) the ratio of MCL communities with more than one cluster; (3)

the proportion of MCL communities with a PFAM clan entropy equal to 0; (4) the intra-

community HHblits-score/Aligned-columns score (normalized by the maximum value); and (5)

the number of MCL communities, which should, in the end, reflect the number of non-redundant

DAs. We iterated through values ranging from 1.2 to 3.0, with incremental steps of 0.1. During

the inference process, some of the GCs became orphans in the graph. We applied a three-step

approach to assigning a community membership to these GCs. First, we used less stringent

conditions (probability ≥ 50% and coverage >= 40%) to find homologs in the already existing

GCCs. Then, we ran a second iteration to find secondary relationships between the newly

assigned GCs and the missing ones. Lastly, we created new communities with the remaining

GCs. We repeated the whole process with the other categories (KWP, GU and EU), applying

the optimal inflation value found for the Known (2.2 for metagenomic and 2.5 for genomic data).

Gene cluster communities validation

We tested the biological significance of the GCCs using the phylogeny of proteorhodopsin[45]

(PR). We used the proteorhodopsin HMM profiles[43] to screen the marine metagenomic datasets

using *hmmsearch* (version 3.1b2)[79]. We kept the hits with a coverage > 0.4 and e-value <= 1e-

5. We removed identical duplicates from the sequences assigned to PR with CD-HIT[98] (v4.6)

and cleaned from sequences with less than 100 amino acids. To place the identified PR

sequences into the MicRhode[45] PR tree first, we optimized the initial tree parameters and

branch lengths with RAxML (v8.2.12)[99]. We used PaPaRA (v2.5)[100] to incrementally align the

query PR sequences against the MicRhode PR reference alignment and *pplacer*[101]

(v1.1.alpha19-0-g807f6f3) to place the sequences into the tree. Finally, we assigned the query

PR sequences to the MicRhode PR Superclusters based on the phylogenetic placement. We

further investigated the GCs annotated as viral (196 genes, 14 GC) comparing them to the six

newly discovered viral PRs[102] using Parasail[82] (-a sg_stats_scan_sse2_128_16 -t 8 -c 1 -x). As

an additional evaluation, we investigated the distributions of standard GCCs and HQ GCCs

within ribosomal protein families. We obtained the ribosomal proteins used for the analysis

combining the set of 16 ribosomal proteins from Méheust et al.[44] and those contained in the

collection of bacterial single-copy genes of Anvi'o[103]. Also, for the ribosomal proteins, we

compared the outcome of our method to the one proposed by Méheust et al.[44] (Supp. Note 9).

670   Metagenomic sample selection for downstream analyses

671   For the subsequent ecological analyses, we selected those metagenomes with a number of

672   genes larger or equal to the first quartile of the distribution of all the metagenomic gene counts.

673   (Supp. Table 13).

674   Gene cluster abundance profiles in genomes and metagenomes

We estimated abundance profiles for the metagenomic cluster categories using the read

coverage to each predicted gene as a proxy for abundance. We calculated the coverage by

mapping the reads against the assembly contigs using the *bwa-mem* algorithm from *BWA*

mapper[104]. Then, we used *BEDTOOLS*[105], to find the intersection of the gene coordinates to the

assemblies, and normalize the per-base coverage by the length of the gene. We calculated the

cluster abundance in a sample as the sum of the cluster gene abundances in that sample, and

the cluster category abundance in a sample as the sum of the cluster abundances. We obtained

the proportions of the different gene cluster categories applying a total-sum-scaling

normalization. For the genomic abundance profiles, we used the number of genes in the

genomes and normalized by the total gene counts per genome.

Rate of genomic and metagenomic gene clusters accumulation

We calculated the cumulative number of known and unknown GCs as a function of the number of metagenomes and genomes. For each metagenome count, we generated 1000 random sets, and we calculated the number of GCs and GCCs recovered. For this analysis, we used 1,246 HMP metagenomes and 358 marine metagenomes (242 from TARA and 116 from Malaspina). We repeated the same procedure for the genomic dataset. We removed the singletons from the metagenomic dataset with an abundance smaller than the mode abundance of the singletons that got reclassified as good-quality clusters after integrating the GTDB data to minimize the impact of potential spurious singletons. To complement those analyses, we evaluated the coverage of our dataset by searching seven different state-of-the-art databases against our set of metagenomic GC HMM profiles (Supp. Note 12).

Occurrence of gene clusters in the environment

We used 1,264 metagenomes from the TARA Oceans, MALASPINA Expedition, OSD2014 and HMP-I/II to explore the properties of the unknown CDS-space in the environment. We applied the Levins Niche Breadth (NB) index[106] to investigate the GCs and GCCs environmental distributions. We removed the GCs and cluster communities with a mean relative abundance < 1e-5. We followed a divide-and-conquer strategy to avoid the computational burden of generating the null-models to test the significance of the distributions owing to the large number of metagenomes and GCs. First, we grouped similar samples based on the gene cluster content using the Bray-Curtis dissimilarity[107] in combination with the *Dynamic Tree Cut*[108] R package. We created 100 random datasets picking up one random sample from each group. For each of the 100 random datasets, we created 100 random abundance matrices using the *nullmodel* function of the *quasiswap* count method[109]. Then we calculated the *observed* NB and obtained

the 2.5% and 97.5% quantiles based on the randomized sets. We compared the observed and

quantile values for each gene cluster and defined it to have a *Narrow distribution* when the

observed was smaller than the 2.5% quantile and to have a *Broad distribution* when it was

larger than the 97.5% quantile. Otherwise, we classified the cluster as *Non-significant*[110]. We

used a majority voting approach to get a consensus distribution classification based on the ten

random datasets.

715 Identification of prophages in genomic sequences

We used PhageBoost (https://github.com/ku-cbd/PhageBoost/) to find gene regions in the

microbial genomes that result in high viral signals against the overall genome signal. We set the

following thresholds to consider a region prophage: minimum of 10 genes, maximum 5 gaps,

single-gene probability threshold 0.9. We further smoothed the predictions using Parzen rolling

windows of 20 periods and looked at the smoothed probability distribution across the genome.

We disregarded regions that had a summed smoothed probability less than 0.5, and those

regions that did differ from the overall population of the genes in a genome by using Kruskal–

Wallis rank test (p-value 0.001).

724 Lineage-specific gene clusters

725   We used the F1-score developed for AnnoTree[50] to identify the lineage-specific GCs and to

726   which rank they are specific. Following similar criteria to the ones used in Mendler et al.[50], we

727   considered a gene cluster to be lineage-specific if it is present in less than half of all genomes

728   and at least 2 with F1-score > 0.95.

Phylogenetic conservation of gene clusters

730 We calculated the phylogenetic conservation (τD) of each gene cluster using the

731 *consenTRAIT*[51] function implemented in the R package *castor*[51]. We used a paired Wilcoxon

732 rank-sum test to compare the average τD values for lineage-specific and non-specific GCs.

Evaluation of the OM-RGC v2 uncharacterized fraction

734 We integrated the 46,775,154 genes from the second version of the TARA Ocean Microbial

735 Reference Gene Catalog (OM-RGC v2)[54] into our cluster database using the same procedure

736 as for the genomic data. We evaluated the uncharacterized fraction and the genes classified

737 into the eggNOG[55] category S within the context of our database.

Augmenting RB-TnSeq experimental data

We searched the 37,684 genes of unknown function associated with mutant phenotypes from

Price et al.[14] against our gene cluster profiles. We kept the hits with e-value ≤ 1e-20 and a query

coverage > 60%. Then we filtered the results to keep the hits within 90% of the Log(best-e-

value), and we used a majority vote function to retrieve the consensus category for each hit.

Lastly, we selected the best-hits based on the smallest e-value and the largest query and target

coverage values. We used the fitness values from the RB-TnSeq experiments from Price et al.

to identify genes of unknown function that are important for fitness under certain experimental

conditions.

Code and data availability

The code used for the analyses in the manuscript is available at https://github.com/functional-

dark-side/functional-dark-side.github.io/tree/master/scripts. The code to recreate the figures is

available at https://github.com/functional-dark-side/vanni_et_al-figures. Detailed descriptions of

the different methods and results of this manuscript are available at

https://dark.metagenomics.eu. The workflow AGNOSTOS is available at

https://github.com/functional-dark-side/agnostos-wf, and its database can be downloaded from

https://doi.org/10.6084/m9.figshare.12459056.

Acknowledgements

The authors thankfully acknowledge the computer resources at MareNostrum and the technical

support provided by Barcelona Supercomputing Center (RES-AECT-2014-2-0085), the BMBF-

funded de.NBI Cloud within the German Network for Bioinformatics Infrastructure (de.NBI)

(031A537B, 031A533A, 031A538A, 031A533B, 031A535A, 031A537C, 031A534A, 031A532B),

the University of Oxford Advanced Research Computing

(http://dx.doi.org/10.5281/zenodo.22558) and the MARBITS bioinformatics core at ICM-CSIC.

CV was supported by the Max Planck Society. AFG received funding from the European

Union's Horizon 2020 research and innovation program Blue Growth: Unlocking the potential of

Seas and Oceans under grant agreement no. 634486 (project acronym INMARE). AM was

supported by the Biotechnology and Biological Sciences Research Council [BB/M011755/1,

BB/R015228/1] and RDF by the European Molecular Biology Laboratory core funds. EOC was

supported by project INTERACTOMA RTI2018-101205-B-I00 from the Spanish Agency of

Science MICIU/AEI. SGA and PS received additional funding by the project MAGGY

(CTM2017-87736-R) from the Spanish Ministry of Economy and Competitiveness. The

Malaspina 2010 Expedition was supported by the Spanish Ministry of Economy and

Competitiveness (MINECO) through the Consolider-Ingenio program (ref. CSD2008-00077).

The authors thank Johannes Söding and Alex Bateman for helpful discussions.

Author Contributions

CV, MSS and AF-G performed the analyses and wrote the computational workflow. MS assisted

with the clustering and remote homology searches. KS helped with the identification of

prophages in genomic sequences. PLB and AB provided feedback and assisted with the

ecological analyses. RDF and AM provided feedback and information on the MGnify and Pfam

databases. CMD, PS and SGA provided the Malaspina metagenomes. TOD and AME analyzed

data in the context of metagenome-assembled genomes. AF-G conceived the study and

supervised the work. CV and AF-G wrote the manuscript. All authors read, edited and approved

the final manuscript.

Competing Interests

788 The authors declare no competing interests.

References

1. Hug, L. A. *et al.* A new view of the tree of life. *Nat Microbiol* **1**, 16048 (2016).

2. Sunagawa, S. *et al.* Ocean plankton. Structure and function of the global ocean

microbiome. *Science* **348**, 1261359 (2015).

3. Kopf, A. *et al.* The ocean sampling day consortium. *Gigascience* **4**, 27 (2015).

4. Almeida, A. *et al.* A new genomic blueprint of the human gut microbiota. *Nature* **568**, 499–

504 (2019).

5. Pasolli, E. *et al.* Extensive Unexplored Human Microbiome Diversity Revealed by Over

150,000 Genomes from Metagenomes Spanning Age, Geography, and Lifestyle. *Cell* **176**,

649-662.e20 (2019).

6. Pachiadaki, M. G. *et al.* Charting the Complexity of the Marine Microbiome through Single-

Cell Genomics. *Cell* **179**, 1623-1635.e11 (2019).

7. Cross, K. L. *et al.* Targeted isolation and cultivation of uncultivated bacteria by reverse

genomics. *Nat. Biotechnol.* **37**, 1314–1321 (2019).

8. Delmont, T. O. *et al.* Functional repertoire convergence of distantly related eukaryotic

plankton lineages revealed by genome-resolved metagenomics. *Cold Spring Harbor*

Laboratory 2020.10.15.341214 (2020) doi:10.1101/2020.10.15.341214.

9. Eloe-Fadrosh, E. A. *et al.* Global metagenomic survey reveals a new bacterial candidate

phylum in geothermal springs. *Nat. Commun.* **7**, 10476 (2016).

10. Brown, C. T. *et al.* Unusual biology across a group comprising more than 15% of domain

Bacteria. *Nature* **523**, 208–211 (2015).

11. Spang, A. *et al.* Complex archaea that bridge the gap between prokaryotes and eukaryotes.

Nature **521**, 173–179 (2015).

12. Parks, D. H. *et al.* A standardized bacterial taxonomy based on genome phylogeny

substantially revises the tree of life. *Nat. Biotechnol.* **36**, 996–1004 (2018).

13. Bernard, G., Pathmanathan, J. S., Lannes, R., Lopez, P. & Bapteste, E. Microbial Dark

Matter Investigations: How Microbial Studies Transform Biological Knowledge and

Empirically Sketch a Logic of Scientific Discovery. *Genome Biol. Evol.* **10**, 707–715 (2018).

14. Price, M. N. *et al.* Mutant phenotypes for thousands of bacterial genes of unknown function.

Nature **557**, 503–509 (2018).

15. Carradec, Q. *et al.* A global ocean atlas of eukaryotic genes. *Nat. Commun.* **9**, 373 (2018).

16. Almeida, A. *et al.* A unified catalog of 204,938 reference genomes from the human gut

microbiome. *Nat. Biotechnol.* (2020) doi:10.1038/s41587-020-0603-3.

17. Mitchell, A. L. *et al.* MGnify: the microbiome analysis resource in 2020. *Nucleic Acids Res.*

48, D570–D578 (2020).

18. Quince, C., Walker, A. W., Simpson, J. T., Loman, N. J. & Segata, N. Shotgun

metagenomics, from sampling to analysis. *Nat. Biotechnol.* **35**, 833–844 (2017).

19. Franzosa, E. A. *et al.* Species-level functional profiling of metagenomes and

metatranscriptomes. *Nat. Methods* **15**, 962–968 (2018).

20. Huerta-Cepas, J. *et al.* Fast Genome-Wide Functional Annotation through Orthology

Assignment by eggNOG-Mapper. *Mol. Biol. Evol.* **34**, 2115–2122 (2017).

21. Chen, I.-M. A. *et al.* IMG/M v.5.0: an integrated data management and comparative

analysis system for microbial genomes and microbiomes. *Nucleic Acids Res.* **47**, D666–

D677 (2019).

22. Hanson, A. D., Pribat, A., Waller, J. C. & Crécy-Lagard, V. de. 'Unknown'proteins and

'orphan'enzymes: the missing half of the engineering parts list--and how to find it. *Biochem.*

J **425**, 1–11 (2010).

23. Arnold, F. H. Design by Directed Evolution. *Acc. Chem. Res.* **31**, 125–131 (1998).

24. Brandenberg, O. F., Fasan, R. & Arnold, F. H. Exploiting and engineering hemoproteins for

abiological carbene and nitrene transfer reactions. *Curr. Opin. Biotechnol.* **47**, 102–111

(2017).

25. Arnold, F. H. Directed Evolution: Bringing New Chemistry to Life. *Angew. Chem. Int. Ed*

Engl. **57**, 4143–4148 (2018).

26. Jaroszewski, L. *et al.* Exploration of uncharted regions of the protein universe. *PLoS Biol.* **7**,

(2009).

27. Buttigieg, L. P. *et al.* Ecogenomic Perspectives on Domains of Unknown Function:

Correlation-Based Exploration of Marine Metagenomes. *PLoS One* **8**, (2013).

28. Yooseph, S. *et al.* The Sorcerer II global ocean sampling expedition: Expanding the

universe of protein families. *PLoS Biol.* **5**, 0432–0466 (2007).

29. Wyman, S. K., Avila-Herrera, A., Nayfach, S. & Pollard, K. S. A most wanted list of

conserved microbial protein families with no known domains. *PLoS One* **13**, e0205749

(2018).

30. Brum, J. R. *et al.* Illuminating structural proteins in viral "dark matter" with metaproteomics.

Proc. Natl. Acad. Sci. U. S. A. **113**, 2436–2441 (2016).

31. Bateman, A., Coggill, P. & Finn, D. R. DUFs: Families in search of function. *Acta*

Crystallogr. Sect. F Struct. Biol. Cryst. Commun. **66**, 1148–1152 (2010).

32. Lobb, B., Kurtz, D. A., Moreno-Hagelsieb, G. & Doxey, A. C. Remote homology and the

functions of metagenomic dark matter. *Front. Genet.* **6**, 1–12 (2015).

33. Bitard-Feildel, T. & Callebaut, I. Exploring the dark foldable proteome by considering

hydrophobic amino acids topology. *Sci. Rep.* **7**, 41425 (2017).

34. Bileschi, M. L. *et al.* Using Deep Learning to Annotate the Protein Universe. *bioRxiv* 626507

(2019) doi:10.1101/626507.

35. Liu, X. L. Deep Recurrent Neural Network for Protein Function Prediction from Sequence.

bioRxiv 103994 (2017) doi:10.1101/103994.

36. Rost, B. Twilight zone of protein sequence alignments. *Protein Eng. Des. Sel.* **12**, 85–94

(1999).

37. Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for the

analysis of massive data sets. *Nat. Biotechnol.* **advance on**, (2017).

38. Steinegger, M. *et al.* HH-suite3 for fast remote homology detection and deep protein

annotation. *BMC Bioinformatics* **20**, 473 (2019).

39. Skewes-Cox, P., Sharpton, T. J., Pollard, K. S. & DeRisi, J. L. Profile hidden Markov

models for the detection of viruses within metagenomic sequence data. *PLoS One* **9**,

e105067 (2014).

40. Sberro, H. *et al.* Large-Scale Analyses of Human Microbiomes Reveal Thousands of Small,

Novel Genes. *Cell* **178**, 1245-1259.e14 (2019).

41. Perdigão, N., Rosa, A. C. & O'Donoghue, S. I. The Dark Proteome Database. *BioData Min.*

10, 1–11 (2017).

42. Habchi, J., Tompa, P., Longhi, S. & Uversky, V. N. Introducing protein intrinsic disorder.

Chem. Rev. **114**, 6561–6588 (2014).

43. Olson, D. K., Yoshizawa, S., Boeuf, D., Iwasaki, W. & DeLong, E. F. Proteorhodopsin

variability and distribution in the North Pacific Subtropical Gyre. *ISME J.* **12**, 1047–1060

(2018).

44. Méheust, R., Burstein, D., Castelle, C. J. & Banfield, J. F. The distinction of CPR bacteria

from other bacteria based on protein family content. *Nat. Commun.* **10**, 4173 (2019).

45. Boeuf, D., Audic, S., Brillet-Guéguen, L., Caron, C. & Jeanthon, C. MicRhoDE: a curated

database for the analysis of microbial rhodopsin diversity and evolution. *Database* **2015**,

(2015).

46. La Cono, V. *et al.* Partaking of Archaea to biogeochemical cycling in oxygen-deficient

zones of meromictic saline Lake Faro (Messina, Italy). *Environ. Microbiol.* **15**, 1717–1733

(2013).

47. Edwards, R. A. *et al.* Global phylogeography and ancient evolution of the widespread

human gut virus crAssphage. *Nat Microbiol* **4**, 1727–1736 (2019).

48. Dubinkina, V. B., Ischenko, D. S., Ulyantsev, V. I., Tyakht, A. V. & Alexeev, D. G.

894 Assessment of k-mer spectrum applicability for metagenomic dissimilarity analysis. *BMC*

895 *Bioinformatics* vol. 17 (2016).

896 49. Ma, Y. *et al.* Human papillomavirus community in healthy persons, defined by

897 metagenomics analysis of human microbiome project shotgun sequencing data sets. *J.*

898 *Virol.* **88**, 4786–4797 (2014).

899 50. Mendler, K. *et al.* AnnoTree: visualization and exploration of a functionally annotated

900 microbial tree of life. *Nucleic Acids Res.* **47**, 4442–4448 (2019).

901 51. Martiny, A. C., Treseder, K. & Pusch, G. Phylogenetic conservatism of functional traits in

902 microorganisms. *ISME J.* **7**, 830–838 (2013).

903 52. Rinke, C. *et al.* Insights into the phylogeny and coding potential of microbial dark matter.

904 *Nature* **499**, 431–437 (2013).

905 53. Anantharaman, K. *et al.* Expanded diversity of microbial groups that shape the dissimilatory

906 sulfur cycle. *ISME J.* **12**, 1715–1728 (2018).

907 54. Salazar, G. *et al.* Gene Expression Changes and Community Turnover Differentially Shape

908 the Global Ocean Metatranscriptome. *Cell* **179**, 1068-1083.e21 (2019).

909 55. Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and phylogenetically

910 annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids*

911 *Res.* **47**, D309–D314 (2019).

912 56. Heffernan, B., Murphy, C. D. & Casey, E. Comparison of planktonic and biofilm cultures of

913 Pseudomonas fluorescens DSM 8341 cells grown on fluoroacetate. *Appl. Environ.*

914 *Microbiol.* **75**, 2899–2907 (2009).

915 57. Scales, B. S., Dickson, R. P., LiPuma, J. J. & Huffnagle, G. B. Microbiology, genomics, and

916 clinical significance of the Pseudomonas fluorescens species complex, an unappreciated

917 colonizer of humans. *Clin. Microbiol. Rev.* **27**, 927–948 (2014).

918 58. Steinegger, M. & Söding, J. Clustering huge protein sequence sets in linear time. *Nat.*

919 *Commun.* **9**, 2542 (2018).

59. Francino, M. P. The ecology of bacterial genes and the survival of the new. *Int. J. Evol.*

Biol. **2012**, 394026 (2012).

60. Muller, E. E. L. Determining Microbial Niche Breadth in the Environment for Better

Ecosystem Fate Predictions. *mSystems* **4**, (2019).

61. Roumpeka, D. D., Wallace, R. J., Escalettes, F., Fotheringham, I. & Watson, M. A Review

of Bioinformatics Tools for Bio-Prospecting from Metagenomic Sequence Data. *Front.*

Genet. **8**, 23 (2017).

62. Ivanova, N. N. *et al.* Stop codon reassignments in the wild. *Science* **344**, 909–913 (2014).

63. Steinegger, M., Mirdita, M. & Söding, J. Protein-level assembly increases protein sequence

recovery from metagenomic samples manyfold. *Nat. Methods* **16**, 603–606 (2019).

64. Titus Brown, C. *et al.* Exploring neighborhoods in large metagenome assembly graphs

reveals hidden sequence diversity. *bioRxiv* 462788 (2018) doi:10.1101/462788.

65. Höps, W., Jeffryes, M. & Bateman, A. Gene Unprediction with Spurio: A tool to identify

spurious protein sequences. *F1000Res.* **7**, 261 (2018).

66. Heinzinger, M. *et al.* Modeling aspects of the language of life through transfer-learning

protein sequences. *BMC Bioinformatics* **20**, 723 (2019).

67. Breitwieser, F. P., Pertea, M., Zimin, A. & Salzberg, S. L. Human contamination in bacterial

genomes has created thousands of spurious proteins. *Genome Res.* (2019)

doi:10.1101/gr.245373.118.

68. Steinegger, M. & Salzberg, S. L. Terminating contamination: large-scale search identifies

more than 2,000,000 contaminated entries in GenBank. *Genome Biol.* **21**, 115 (2020).

69. Vorobev, A. *et al.* Transcriptome reconstruction and functional analysis of eukaryotic

marine plankton communities via high-throughput metagenomics and metatranscriptomics.

Genome Res. (2020) doi:10.1101/gr.253070.119.

70. Thomas, A. M. & Segata, N. Multiple levels of the unknown in microbiome research. *BMC*

Biol. **17**, 48 (2019).

71. Duarte, C. M. Seafaring in the 21St Century: The Malaspina 2010 Circumnavigation

Expedition. *Limnol. Oceanog. Bull.* **24**, 11–14 (2015).

72. Rusch, D. B. *et al.* The Sorcerer II Global Ocean Sampling Expedition: Northwest Atlantic

through Eastern Tropical Pacific. *PLoS Biol.* **5**, 1–34 (2007).

73. Lloyd-Price, J. *et al.* Strains, functions and dynamics in the expanded Human Microbiome

Project. *Nature* **550**, 61–66 (2017).

74. Sanger, F., Nicklen, S. & Coulson, A. R. DNA sequencing with chain-terminating inhibitors.

Proc. Natl. Acad. Sci. U. S. A. **74**, 5463–5467 (1977).

75. Köster, J. Reproducible data analysis with Snakemake. *F1000Res.* **7**, (2018).

76. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* **11**, 119–119 (2010).

77. Eberhardt, R. Y. *et al.* AntiFam: a tool to help identify spurious ORFs in protein annotation.

Database **2012**, bas003–bas003 (2012).

78. Yooseph, S., Li, W. & Sutton, G. Gene identification and protein classification in microbial

metagenomic sequence data via incremental clustering. *BMC Bioinformatics* **9**, 1–13

(2008).

79. Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence similarity

searching. *Nucleic Acids Res.* **39**, W29–W37 (2011).

80. Finn, R. D. *et al.* The Pfam protein families database: towards a more sustainable future.

Nucleic Acids Res. **44**, D279–D285 (2016).

81. Bennett, K. D. Determination of the number of zones in a biostratigraphical sequence. *New*

Phytol. **132**, 155–170 (1996).

82. Daily, J. Parasail: SIMD C library for global, semi-global, and local pairwise sequence

alignments. *BMC Bioinformatics* **17**, 81–81 (2016).

83. Žure, M., Fernandez-Guerra, A., Munn, C. B. & Harder, J. Geographic distribution at

subspecies resolution level: closely related Rhodopirellula species in European coastal

sediments. *ISME J.* **11**, 478–489 (2017).

84. Chafee, M. *et al.* Recurrent patterns of microdiversity in a temperate coastal marine

environment. *ISME J.* **12**, 237–252 (2018).

85. Csardi, G. & Nepusz, T. The igraph software package for complex network research.

InterJournal vol. Complex Systems 1695 (2006).

86. Deorowicz, S., Debudaj-Grabysz, A. & Gudyś, A. FAMSA: Fast and accurate multiple

sequence alignment of huge protein families. *Sci. Rep.* **6**, 33964–33964 (2016).

87. Vanhoutreve, R. *et al.* LEON-BIS: multiple alignment evaluation of sequence neighbours

using a Bayesian inference system. *BMC Bioinformatics* **17**, 271–271 (2016).

88. Jehl, P., Sievers, F. & Higgins, D. G. OD-seq: outlier detection in multiple sequence

alignments. *BMC Bioinformatics* **16**, 269–269 (2015).

89. Broder, A. Z. On the resemblance and containment of documents. in *Proceedings.*

Compression and Complexity of SEQUENCES 1997 (Cat. No. 97TB100171) 21–29 (IEEE,

1997).

90. The UniProt Consortium. UniProt: the universal protein knowledgebase. *Nucleic Acids Res.*

45, D158–D169 (2017).

91. NCBI Resource Coordinators. Database resources of the National Center for Biotechnology

Information. *Nucleic Acids Res.* **46**, D8–D13 (2018).

92. Hingamp, P. *et al.* Exploring nucleo-cytoplasmic large DNA viruses in Tara Oceans

microbial metagenomes. *ISME J.* **7**, 1678–1695 (2013).

93. Remmert, M., Biegert, A., Hauser, A. & Söding, J. HHblits: Lightning-fast iterative protein

sequence searching by HMM-HMM alignment. *Nat. Methods* **9**, 173–175 (2012).

94. UniProt Consortium, T. UniProt: the universal protein knowledgebase. *Nucleic Acids Res.*

46, 2699 (2018).

95. Dick, J. M. Calculation of the relative metastabilities of proteins using the CHNOSZ

software package. *Geochem. Trans.* **9**, 10 (2008).

96. Hausser, J. & Strimmer, K. Entropy inference and the James-Stein estimator, with

application to nonlinear gene association networks. *arXiv [stat.ML]* (2008).

97. van Dongen, S. & Abreu-Goodger, C. Using MCL to Extract Clusters from Networks. in

Bacterial Molecular Networks: Methods and Protocols (eds. van Helden, J., Toussaint, A. &

Thieffry, D.) 281–295 (Springer New York, 2012).

98. Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of protein

or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

99. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

100. Berger, S. A. & Stamatakis, A. PaPaRa 2.0: a vectorized algorithm for probabilistic

phylogeny-aware alignment extension. *Heidelberg Institute for Theoretical Studies,*

http://sco. h-its. org/exelixis/publica tions. html. Exelixis-RRDR-2012-2015 (2012).

101. Matsen, F. A., Kodner, R. B. & Armbrust, E. V. pplacer: linear time maximum-likelihood and

Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*

Bioinformatics **11**, 538 (2010).

102. Needham, D. M. *et al.* A distinct lineage of giant viruses brings a rhodopsin photosystem to

unicellular marine predators. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 20574–20583 (2019).

103. Murat Eren, A. *et al.* Anvi'o: an advanced analysis and visualization platform for 'omics

data. *PeerJ* **3**, e1319 (2015).

104. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows–Wheeler transform.

Bioinformatics **26**, 589–595 (2010).

105. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841–842 (2010).

106. Levins, R. THE STRATEGY OF MODEL BUILDING IN POPULATION BIOLOGY. *Am. Sci.*

54, 421–431 (1966).

107. Bray, J. R., Roger Bray, J. & Curtis, J. T. An Ordination of the Upland Forest Communities

of Southern Wisconsin. *Ecological Monographs* vol. 27 325–349 (1957).

108. Langfelder, P., Zhang, B. & Horvath, S. Defining clusters from a hierarchical cluster tree:

the Dynamic Tree Cut package for R. *Bioinformatics* **24**, 719–720 (2008).

109. Miklós, I. & Podani, J. RANDOMIZATION OF PRESENCE–ABSENCE MATRICES:

COMMENTS AND NEW ALGORITHMS. *Ecology* vol. 85 86–92 (2004).

110. Salazar, G. *et al.* Particle-association lifestyle is a phylogenetically conserved trait in

bathypelagic prokaryotes. *Mol. Ecol.* **24**, 5692–5706 (2015).