Unifying the known and unknown microbial coding sequence space

Chiara Vanni[1,2], Matthew S. Schechter[1,3], Silvia G. Acinas[4], Albert Barberán[5], Pier Luigi Buttigieg[6], Emilio O. Casamayor[7], Tom O. Delmont[8], Carlos M. Duarte[9], A. Murat Eren[3,10], Robert D. Finn[11], Renzo Kottmann[1], Alex Mitchell[11], Pablo Sanchez[4], Kimmo Siren[12], Martin Steinegger[13,14], Frank Oliver Glöckner[15,16,2], Antonio Fernandez-Guerra[1,17 *]

Affiliations

Microbial Genomics and Bioinformatics Research Group, Max Planck Institute for Marine Microbiology, Celsiusstraße 1, 28359, Bremen, Germany

Jacobs University Bremen, Campus Ring 1, 28759 Bremen, Germany

Department of Medicine, University of Chicago, Chicago, IL 60637, USA

Department of Marine Biology and Oceanography, Institut de Ciènces del Mar, CSIC, Barcelona, Spain.

Department of Environmental Science, University of Arizona, Tucson, 85721 AZ, USA

Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Am Handelshafen 12, 27570 Bremerhaven, Germany

Center for Advanced Studies of Blanes CEAB-CSIC, Spanish Council for Research, Blanes, Spain

8 Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Univ Evry,

Université Paris-Saclay, 91057 Evry, France

9 Red Sea Research Centre (RSRC) and Computational Bioscience Research Center (CBRC),

King Abdullah University of Science and Technology, Thuwal 23955, Saudi Arabia

10 Josephine Bay Paul Center, Marine Biological Laboratory, Woods Hole, MA 02543, USA

11 European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI),

Wellcome Genome Campus, Hinxton, Cambridge CB10 1SD, UK

12 Section for Evolutionary Genomics, The GLOBE Institute, University of Copenhagen,

Copenhagen, Denmark

13 School of Biological Sciences, Seoul National University, Seoul, 08826, South Korea

14 Institute of Molecular Biology and Genetics, Seoul National University, Seoul, 08826, South

Korea

15 University of Bremen, MARUM, Leobener Str. 8, 28359 Bremen, Germany

Life Sciences and Chemistry, Campus Ring 1, 28759 Bremen, Germany

16 Computing Center, Helmholtz Center for Polar and Marine Research, Am Handelshafen 12,

27570 Bremerhaven, Germany

17 Lundbeck GeoGenetics Centre, The Globe Institute, University of Copenhagen, 1350

Copenhagen, Denmark

*Corresponding author: Antonio Fernandez-Guerra, antonio.fernandez-guerra@sund.ku.dk

Abstract

Genes of unknown function are among the biggest challenges in molecular biology, especially in microbial systems, where 40%-60% of the predicted genes are unknown. Despite previous attempts, systematic approaches to include the unknown fraction into analytical workflows are still lacking. Here, we propose a conceptual framework and a computational workflow that bridge the known-unknown gap in genomes and metagenomes. We showcase our approach by exploring 415,971,742 genes predicted from 1,749 metagenomes and 28,941 bacterial and archaeal genomes. We quantify the extent of the unknown fraction, its diversity, and its relevance across multiple biomes. Furthermore, we provide a collection of 283,874 lineage-specific genes of unknown function for *Cand*. Patescibacteria, being a significant resource to expand our understanding of their unusual biology. Finally, by identifying a target gene of unknown function for antibiotic resistance, we demonstrate how we can enable the generation of hypotheses that can be used to augment experimental data.

Introduction

Thousands of isolate, single-cell, and metagenome-assembled genomes are guiding us towards a better understanding of life on Earth (Almeida et al., 2019; Cross et al., 2019; Delmont et al., 2020; Hug et al., 2016; Kopf et al., 2015; Pachiadaki et al., 2019; Pasolli et al., 2019; Sunagawa et al., 2015). At the same time, the ever-increasing number of genomes and metagenomes, unlocking uncharted regions of life's diversity, (Brown et al., 2015; Eloe-Fadrosh et al., 2016; Hug et al., 2016) are providing new perspectives on the evolution of life (Parks et al., 2018; Spang et al., 2015). However, our rapidly growing inventories of new genes have a glaring issue: between 40% and 60% cannot be assigned to a known function (Almeida et al., 2020; Bernard, Pathmanathan, Lannes, Lopez, & Bapteste, 2018; Carradec et al., 2018; Price et al., 2018). Current analytical approaches for genomic and metagenomic data (Chen et al., 2019; Franzosa et al., 2018; Huerta-Cepas et al., 2017; Mitchell et al., 2020; Quince, Walker, Simpson, Loman, & Segata, 2017) generally do not include this uncharacterized fraction in downstream analyses, constraining their results to conserved pathways and housekeeping functions (Quince et al., 2017). This inability to handle the unknown is an immense impediment to realizing the potential for discovery of microbiology and molecular biology at large (Bernard et al., 2018; Hanson, Pribat, Waller, & Crécy-Lagard, 2010). Predicting function from traditional single sequence similarity appears to have yielded all it can (Arnold, 1998, 2018; Brandenberg, Fasan, & Arnold, 2017), thus several groups have attempted to resolve gene function by other means. Such efforts include combining biochemistry and crystallography (Jaroszewski et al., 2009); using environmental co-occurrence (Buttigieg et al., 2013); by grouping those genes into evolutionarily related families (Bateman, Coggill, & Finn, 2010; Brum et al., 2016; Wyman, Avila-Herrera, Nayfach, & Pollard, 2018; Yooseph et al., 2007); using remote homologies (Bitard-Feildel & Callebaut, 2017; Lobb, Kurtz, Moreno-Hagelsieb, & Doxey, 2015); or more recently using deep learning approaches (Bileschi et al., 2019; Liu, 2017). In 2018, Price et al. (Price et

al., 2018) developed a high-throughput experimental pipeline that provides mutant phenotypes

for thousands of bacterial genes of unknown function being one of the most promising methods

to tackle the unknown. Despite their promise, experimental methods are labor-intensive and

require novel computational methods that could bridge the existing gap between the known and

unknown coding sequence space (CDS-space).

Here we present a conceptual framework and a computational workflow that closes the gap

between the known and unknown CDS-space by connecting genomic and metagenomic gene

clusters. Our approach adds context to vast amounts of unknown biology, providing an

invaluable resource to understand the unknown functional fraction better and boost the current

methods for its experimental characterization. The application of our approach to 415,971,742

genes predicted from 1,749 metagenomes and 28,941 bacterial and archaeal genomes

revealed that the unknown fraction (1) is smaller than expected, (2) is exceptionally diverse, and

(3) is phylogenetically more conserved and predominantly lineage-specific at the Species level.

Finally, we show how we can connect all the outputs produced by our approach to augment the

results from experimental data and add context to genes of unknown function through

hypothesis-driven molecular investigations.

Results

A conceptual framework and a computational workflow to unify the known and the unknown coding sequence space

We created the conceptual and technical foundations to unify the known and unknown CDS-space facilitating the integration of the genes of unknown function into ecological, evolutionary and biotechnological investigations. First, we conceptually partitioned the known and unknown fractions into (1) Known with Pfam annotations (K), (2) Known without Pfam annotations (KWP), (3) Genomic unknown (GU), and (4) Environmental unknown (EU) (Fig. 1A). The framework introduces a subtle change of paradigm compared to traditional approaches where our objective is to provide the best representation of the unknown space. We gear all our efforts towards finding sequences without any evidence of known homologies by pushing the search space beyond the *twilight zone* of sequence similarity (Rost, 1999). With this objective in mind, we use gene clusters (GCs) instead of genes as the fundamental unit to compartmentalize the CDS-space owing to their unique characteristics (Fig. 1B). (1) GCs produce a structured CDS-space reducing its complexity (Fig. 1B), (2) are independent of the known and unknown fraction, (3) are conserved across environments and organisms, and (4) can be used to aggregate information from different sources (Fig. 1A). Moreover, the GCs provide a good compromise in terms of resolution for analytical purposes, and owing to their unique properties, one can perform analyses at different scales. For fine-grained analyses, we can exploit the gene associations within each GC; and for coarse-grained analyses, we can create groups of GCs based on their shared homologies (Fig. 1B).

A

CONCEPTUAL FRAMEWORK   COMPUTATIONAL WORKFLOW

CDS-space

KNOWN WITH PFAM   KNOWN WITHOUT PFAM   CDS-space DARKNESS

GENOMIC UNKNOWN   ENVIRONMENTAL UNKNOWN

Known with Pfam
GC annotated to contain one or more Pfam entries but excluding DUFs

Known without Pfam
GC that have a known function but lack a Pfam annotation

Genomic unknown
GC of unknown function (DUFs are included here) and found in sequenced or draft genomes

Environmental unknown
GC of unknown function not detected in sequenced or draft genomes, but only in environmental metagenomes or MAGs

Groups of GCs

GENE CLUSTER COMMUNITIES

AGNOSTOS → GENE CLUSTERS (GC)

Modules
- Create GC DB
- Update GC DB
- GC profile search

KNOWLEDGE

Rich contextual data

'OMICS DATA   PHYLOGENETIC DATA

ECOLOGICAL DATA   EXPERIMENTAL DATA

Conserved sequence composition   Domain architecture homogeneity

Remote homology characterization

GC properties

'OMICS DATA
- expression data
- genomic context/distribution

PHYLOGENETIC DATA
- lineage specificity
- phylogenetic conservation

ECOLOGICAL DATA
- environmental range
- distribution patterns

EXPERIMENTAL DATA
- experimental conditions
- mutant phenotypes

B

AGNOSTOS (*DB creation*)   TRADITIONAL DOWNSTREAM PROCESSING

GENE → (META)GENOME CDS-SPACE

CDS-SPACE → CDS-SPACE (KNOWN / UNKNOWN) → CDS-SPACE (KNOWN / UNKNOWN)

GENE CLUSTER

GENE → (META)GENOME CDS-SPACE

AGNOSTOS (*DB update*) → GENE CLUSTER DB

EXISTING GENE CLUSTERS

NEW GENE CLUSTERS

CDS-SPACE (KNOWN / UNKNOWN)

NEW GENE CLUSTER

GENE CLUSTER with annotations

C

GROUPS OF GENE CLUSTERS   GENE CLUSTER

GENE CLUSTER   SEQUENCE SIMILARITY NETWORK   GENE

GENE CLUSTER   GENE CLUSTER

MARINE
SOIL
EXPERIMENTAL CONDITIONS
HUMAN GUT

COARSE-GRAINED ◄——— ANALYSIS SCOPE ———► FINE-GRAINED

Figure 1: Conceptual framework to unify the known and unknown CDS-space and integration of the
framework in the current analytical workflows (A) Link between the conceptual framework and the
computational workflow to partition the CDS-space in the four conceptual categories. AGNOSTOS infers,
validates and refines the GCs and combines them in gene cluster communities (GCCs). Then, it classifies
them in one of the four conceptual categories based on their level of 'darkness'. Finally, we add context to
each GC based on several sources of information, providing a robust framework for generating
hypotheses that can be used to augment experimental data. (B) The computational workflow provides two
mechanisms to structure the CDS-space using GCs, de novo creation of the GCs (*DB creation*), or
integrating the dataset in an existing GC database (*DB update*). The structured CDS-space can then be
plugged into traditional analytical workflows to annotate the genes within each GC of the known fraction.
With AGNOSTOS, we provide the opportunity to integrate the unknown fraction into the current
microbiome analyses easily. C) The versatility of the GCs enables analyses at different scales depending
on the scope of our experiments. We can group GCs in gene cluster communities based on their shared
homologies to perform coarse-grained analyses. On the other hand, we can design fine-grained analyses
using the relationships between the genes in a GC, i.e., detecting network modules in the GC inner
sequence similarity network. Additionally, given that GCs are conserved across environments, organisms
and experimental conditions give us access to an unprecedented amount of information to design and
interpret experimental data.

Driven by the concepts defined in the conceptual framework, we developed AGNOSTOS, a

computational workflow that infers, validates, refines, and classifies GCs in the four proposed

categories (Fig. 1A; Fig. 1B; Supp. Fig 1). AGNOSTOS provides two operational modules (*DB*

creation and *DB update*) to produce GCs with a highly conserved intra-homogeneous structure

(Fig. 1B), both in terms of sequence similarity and domain architecture homogeneity; it exhausts

any existing homology to known genes and provides a proper delimitation of the unknown CDS-

space before classifying each GC in one of the four categories (Methods). In the last step, we

decorate each GC with a rich collection of contextual data compiled from different sources or

generated by analyzing the GC contents in different contexts (Fig. 1A). For each GC, we also

offer several products that can be used for analytical purposes like improved representative

sequences, consensus sequences, sequence profiles for MMseqs2 (Steinegger & Soding,

2017) and HHblits (Steinegger, Meier, et al., 2019), or the GC members as a sequence

similarity network (Methods). To complement the collection, we also provide a subset of what

we define as *high-quality* GCs. The defining criteria are (1) the representative is a complete

gene and (2) more than one-third of genes within a GC are complete genes.

Partitioning and contextualizing the coding sequence space of genomes and metagenomes

We used our approach to explore the unknown CDS-space of 1,749 microbial metagenomes

derived from human and marine environments, and 28,941 genomes from the GTDB_r86 (Supp

Fig 2A). The initial gene prediction of AGNOSTOS (Supp Fig 1) produced 322,248,552 genes

from the environmental dataset and assigned Pfam annotations to 44% of them. Next, it

clustered the predicted genes in 32,465,074 GCs. For the downstream processing, we kept

3,003,897 GCs (83% of the original genes) after filtering out any GC that contained less than

ten genes (Skewes-Cox, Sharpton, Pollard, & DeRisi, 2014) removing 9,549,853 clusters and

19,911,324 singletons (Supp Fig 2A; Supp. Note 1). The validation process selected 2,940,257

good-quality clusters (Fig. 1B; Supp. Table 1; Supp. Note 2), which resulted in 43% of them

being members of the unknown CDS-space after the classification and remote homology

refinement steps (Supp Fig 2A, Supp. Note 3). We build the link between the environmental and

genomic CDS-space by expanding the final collection of GCs with the genes predicted from

GTDB_r86 (Supp Fig 2A). Our environmental GCs already included 72% of the genes from

GTDB_r86; 22% of them created 2,400,037 new GCs, and the rest 6% resulted in singleton

GCs (Supp Fig 2A; Supp. Note 4; Supp. Note 5). The final dataset includes 5,287,759 GCs

(Supp Fig 2A), with both datasets sharing only 922,599 GCs (Supp Fig 2B). The addition of the

GTDB_r86 genes increased the proportion of GCs in the unknown CDS-space to 54%. As the

final step, the workflow generated a subset of 203,217 *high-quality* GCs (Supp. Table 2; Supp.

Fig 3). In these *high-quality* clusters, we identified 12,313 clusters potentially encoding for small

proteins (<= 50 amino acids). Most of these GCs are unknown (66%), which agrees with recent

findings on novel small proteins from metagenomes (Sberro et al., 2019). The KWP category

contains the largest proportion of incomplete genes (Supp. Table 3), disrupting the detection

and assignment of Pfam domains. But it also incorporates sequences with an unusual amino

acid composition that has homology to proteins with high levels of disorder in the DPD database

(Perdigão, Rosa, & O'Donoghue, 2017) and has characteristic functions of intrinsically

disordered proteins (Habchi, Tompa, Longhi, & Uversky, 2014) (IDP) like cellular processes and

signaling as predicted by eggNOG annotations (Supp. Table 4).

As part of the workflow, each GC is complemented with a rich set of information, as shown in

Fig 1A (Supp. Table 5; Supp. Note 6).

184   Beyond the twilight zone, communities of gene clusters

The method we developed to group GCs in gene cluster communities (GCCs) (Fig. 2A) reduced

the final collection of GCs by 87%, producing 673,601 GCCs (Methods; Fig. 2B; Supp. Note 7).

We validated our approach to capture remote homologies between related GCs using two well-

known gene families present in our environmental datasets, proteorhodopsins (Olson,

Yoshizawa, Boeuf, Iwasaki, & DeLong, 2018) and bacterial ribosomal proteins (Méheust,

Burstein, Castelle, & Banfield, 2019). Our dataset contained 64 GCs (12,184 genes) and 3

GCCs (Supp Note 8) classified as proteorhodopsin (PR). One *Known* GCC contained 99% of

the PR annotated genes (Fig. 2C), except 85 genes taxonomically annotated as viral and

assigned to the *PR Supercluster I* (Boeuf, Audic, Brillet-Guéguen, Caron, & Jeanthon, 2015)

within two GU communities (five GU gene clusters; Supp. Note 8). For the ribosomal proteins,

the results were not so satisfactory. We identified 1,843 GCs (781,579 genes) and 98 GCCs.

The number of GCCs is larger than the expected number of ribosomal protein families (16) used

for validation. When we use *high-quality* GCs (Supp. Note 8), we get closer to the expected

number of GCCs (Fig. 2D). With this subset, we identified 26 GCCs and 145 GCs (1,687

genes). The cross-validation of our method against the approach used in Méheust et al.

(Méheust et al., 2019) (Supp. Note 9) confirms the intrinsic complexity of analyzing

metagenomic data. Both approaches showed a high agreement in the GCCs identified (Supp.

Table 9-1). Still, our method inferred fewer GCCs for each of the ribosomal protein families

(Supplementary Figure 9-3), coping better with the nuisances of a metagenomic setup, such as

incomplete genes (Supp. Table 6).

Figure 2: Overview and validation of the workflow to aggregate GCs in communities. (A) We inferred a
gene cluster homology network using the results of an all-vs-all HMM gene cluster comparison with
HHBLITS. The edges of the network are based on the HHblits-score/Aligned-columns. Communities are
identified by an iterative screening of different MCL inflation parameters and evaluated using five different
metrics that consider the inter- and intra-community properties. (B) Comparison of the number of GCs
and GCCs for each of the functional categories. (C) Validation of the GCCs inference based on the
environmental genes annotated as proteorhodopsins. Ribbons in the alluvial plot are genes, and each
stacked bar corresponds (from left to right) to the (1) gene taxonomic classification at the domain level,
(2) GC membership, (3) GCC membership and (4) MicRhoDE operational classification. (D) Validation of
the GCCs inference based on ribosomal proteins based on standard and high-quality GCs.

A smaller but highly diverse unknown coding sequence space

Combining clustering and remote homology searches reduces the extent of the unknown CDS-space compared to what is reported by the traditional genomic and metagenomic analysis approaches (Fig. 3A). Our workflow recruited as much as 71% of genes in human-related metagenomic samples and 65% of the genes in marine metagenomes into the known CDS-space. In both human and marine microbiomes, the genomic unknown fraction showed a similar proportion of genes (21%, Fig. 3A). The number of genes corresponding to EU gene clusters is higher in marine metagenomes; 12% of the genes are part of this GC category. We obtained a comparable result when we evaluated the genes from the GTDB_r86, 75% of bacterial and 64% of archaeal genes were part of the known CDS-space. Archaeal genomes contained more unknowns than those from Bacteria, where 30% of the genes are classified as genomic unknowns in Archaea, and only 20% in Bacteria (Fig. 3A; Supp. Table 7). Our approach allows us to go beyond genes, and for the first time, we can provide a detailed characterization of the CDS-space at the amino acid level. From the 90,128,659,316 amino acids analyzed, the majority of the amino acids in metagenomes (74%) and GTDB_r86 (80%) are in the known CDS-space (Fig. 3B; Supp. Table 7). In both cases, approximately 40% of the amino acids in the known CDS-space were part of a Pfam domain (Fig. 3B; Supp. Table 7). The proportion of amino acids in the unknown CDS-space ranged from 22% in metagenomes and 15% in GTDB_r86. Pfam domains covered only 2% of the amino acids in the unknown CDS-space in both cases. To evaluate the differences between the two CDS-space fractions, we calculated the accumulation rates of GCs and GCCs. For the metagenomic dataset we used 1,264 metagenomes (18,566,675 GCs and 282,580 GCCs) and for the genomic dataset 28,941 genomes (9,586,109 GCs and 496,930 GCCs). The rate of accumulation of unknown GCs was three times higher than the known (2 times for the genomic), and both cases were far from reaching a plateau (Fig. 3C). This is not the case for the GCC accumulation curves (Supp Fig

4B), where they reached a plateau. The accumulation rate is largely determined by the number

of singletons, especially singletons from EUs (Supp note 11 and Supp. Fig 5). While the

accumulation rate of known GCs between marine and human metagenomes is almost identical,

there are striking differences for the unknown GCs (Fig. 3D). These differences are maintained

even when we remove the virus-enriched samples from the marine metagenomes (Supp Fig

4A). Although the marine metagenomes include a large variety of environments, from coastal to

the deep sea, the known space remains quite constrained.

Despite only including marine and human metagenomes in our database, our coverage of other

databases and environments is quite comprehensive, with an overall coverage of 76% (Supp.

Note 12). The lowest covered biomes are freshwater, soil and human non-digestive as revealed

by the screening of MGnify (Mitchell et al., 2020) (release 2018_09; 11 biomes; 843,535,6116

proteins) where we assigned 74% of the MGnify proteins into one of our categories

(Supplementary Fig. 6). Furthermore, as a result of this evaluation, we classified 20% of the

FunkFams (Wyman et al., 2018) and 44% of the unknowns used by Price et al. (Price et al.,

2018) to the known fraction (Supp. Table 12-1).

Figure 3: The extent of the known and unknown coding sequence space (A) Proportion of genes in the known and unknown. (B) Amino acid distribution in the known and unknown CDS-space. (C) Accumulation curves for the known and unknown CDS-space at the GC- level for the metagenomic and genomic data. from TARA, MALASPINA, OSD2014 and HMP-I/II projects. (D) Collector curves comparing the human and marine biomes. Colored lines represented the mean of 1000 permutations and shaded in grey the standard deviation. Non-abundant singleton clusters were excluded from the accumulation curves calculation.

The unknown coding sequence space has a limited ecological distribution in human and marine environments

Although the role of the unknown fraction in the environment is still a mystery, the large number

of gene counts and abundance observed underlines its inherent ecological relevance (Fig. 4A).

In some metagenomes, the genomic unknown fraction can account for more than 40% of the

total gene abundance observed (Fig. 4A). The environmental unknown fraction is also relevant

in several samples, where singleton GCs are the majority (Fig. 4A). We identified two

metagenomes with an unusual composition in terms of environmental unknown singletons. The

marine metagenome corresponds to a sample from Lake Faro (OSD42), a meromictic saline

with a unique extreme environment where Archaea plays an important role (La Cono et al.,

2013). The HMP metagenome (SRS143565) corresponds to a human sample from the right

cubital fossa from a healthy female subject. To understand this unusual composition, we should

perform further analyses to discard potential technical artifacts like sample contamination. The

ratio between the unknown and known GCs revealed that the metagenomes located at the

upper left quadrant in Fig. 4B-C are enriched in GCs of unknown function. In human

metagenomes, we can distinguish between body sites, with the gastrointestinal tract, where

microbial communities are expected to be more diverse and complex, significantly enriched with

genomic unknowns. The HMP metagenomes with the largest ratio of unknowns are those

samples identified to contain crAssphages (Dubinkina, Ischenko, Ulyantsev, Tyakht, & Alexeev,

2016; Edwards et al., 2019) and HPV viruses (Ma et al., 2014) (Supp. Table 8; Supp. Fig. 7).

Consistently, in marine metagenomes (Fig. 4D) we can separate between size fractions, where

the highest ratio in genomic and environmental unknowns corresponds to the ones enriched

with viruses and giant viruses.

Figure 4: Distribution of the unknown coding sequence space in the human and marine metagenomes (A) Ratio between the proportion of the number of genes and their estimated abundances per cluster category and biome. Columns represented in the facet depicts three cluster categories based on the size of the clusters. (B) Relationship between the ratio of Genomic unknowns and Environmental unknowns in the HMP-I/II metagenomes. Gastrointestinal tract metagenomes are enriched in Genomic unknown coding sequences compared to the other body sites. (C) Relationship between the ratio of Genomic unknowns and Environmental unknowns in the TARA Oceans metagenomes. Girus and virus enriched metagenomes show a higher proportion of both unknown coding sequences (genomic and environmental) than the Archaea|Bacteria enriched fractions. (D) Environmental distribution of GCs and GCCs based on Levin's niche breadth index. We obtained the significance values after generating 100 null gene cluster abundance matrices using the *quasiswap* algorithm.

To complement the previous findings, we performed a large-scale analysis to investigate the GC occurrence patterns in the environment. The narrow distribution of the unknown fraction (Fig. 4D) suggests that these GCs might provide a selective advantage and be necessary to adapt to specific environmental conditions. But the pool of broadly distributed environmental unknowns is the most exciting result. We identified traces of potential ubiquitous organisms left

306   uncharacterized by traditional approaches, as more than 80% of these GCs cannot be

307   associated with a metagenome-assembled genome (MAG) (Supp Table 9, Supp. Note 10).

The genomic unknown coding sequence space is lineage-specific

With the inclusion of the genomes from GTDB_r86, we have access to a phylogenomic

framework that can be used to assess how exclusive is a GC within a lineage (lineage-specifity

(Mendler et al., 2019)) and how conserved is a GC across clades (phylogenetic conservation

(Martiny, Treseder, & Pusch, 2013)). We identified 781,814 lineage-specific GCs and 464,923

phylogenetically conserved (P < 0.05) GCs in Bacteria (Supp. Table 10; Supp. Note 13 for

Archaea). The number of lineage-specific GCs increases with the Relative Evolutionary

Distance (Parks et al., 2018) (Fig. 5A) and differences between the known and the unknown

fraction start to be evident at the Family level resulting in 4X more unknown lineage-specific

GCs at the Species level. The unknown GCs are more phylogenetically conserved than the

known (Fig. 5B, p < 0.0001), revealing the importance of the genome's uncharacterized fraction.

However, this is not the case for the lineage-specific and phylogenetically conserved GCs,

where the unknown GCs are less phylogenetically conserved (Fig. 5B), agreeing with the large

number of lineage-specific GCs at Genus and Species level. To discard the possibility that the

lineage-specific GCs of unknown function have a viral origin, we screened all GTDB_r86

genomes for prophages. We only found 37,163 lineage-specific GCs in prophage genomic

regions, being 86% GCs of unknown function. After unveiling the potential relevance of the GCs

of unknown function in bacterial genomes, we identified phyla in GTDB_r86 enriched with these

types of clusters. A clear pattern emerged when we partitioned the phyla based on the ratio of

known to unknown GCs and vice versa (Fig. 5D), the phyla with a larger number of MAGs are

enriched in GCs of unknown function Figure 5D. Phyla with a high proportion of non-classified

GCs (those discarded during the validation steps) contain a small number of genomes and are

primarily composed of MAGs. These groups of phyla highly enriched in unknowns and

represented mainly by MAGs include newly described phyla such as *Cand.* Riflebacteria and

Cand. Patescibacteria (Anantharaman et al., 2018; Brown et al., 2015; Rinke et al., 2013), both

with the largest unknown to known ratio.

335
336
Figure 5: Phylogenomic exploration of the unknown coding sequence space. (A) Distribution of the
lineage-specific GCs by taxonomic level. Lineage-specific unknown GCs are more abundant in the lower
taxonomic levels (Genus, Species). (B) Phylogenetic conservation of the known and unknown coding
sequence space in 27,372 bacterial genomes from GTDB_r86. We observe differences in the
conservation between the known and the unknown coding sequence space for lineage- and non-lineage
specific GCs (paired Wilcoxon rank-sum test; all p-values < 0.0001). (C) The majority of the lineage-
specific clusters are part of the unknown coding sequence space, being a small proportion found in
prophages present in the GTDB_r86 genomes. (D) Known and unknown coding sequence space of the
27,732 GTDB_r86 bacterial genomes grouped by bacterial phyla. Phyla are partitioned based on the ratio
of known to unknown GCs and vice versa. Phyla enriched in MAGs have higher proportions in GCs of
unknown function. Phyla with a high proportion of non-classified clusters (NC; discarded during the
validation steps) tend to contain a small number of genomes. (E) The alluvial plot's left side shows the
uncharacterized (OM-RGC v2 GC) and characterized (OM-RGC v2) fraction of the gene catalog. The
functional annotation is based on the eggNOG annotations provided by Salazar et al.(Salazar et al.,
2019). The right side of the alluvial plot shows the new organization of the OM-RGC v2 coding sequence
space based on the approach described in this study. The treemap in the right links the metagenomic and
genomic space adding context to the unknown fraction of the OM-RGC v2

We demonstrate the possibility of bridging genomic and metagenomic data and simultaneously

unifying the known and unknown CDS-space by integrating the new Ocean Microbial Reference

Gene Catalog (Salazar et al., 2019) (OM-RGC v2) into our database. We assigned 26,170,875

genes to known GCs, 11,422,975 to genomic unknowns, 8,661,221 to environmental unknown

and 520,083 were discarded. From the 11,422,975 genes classified as genomic unknowns, we

could associate 3,261,741 to a GTDB_r86 genome and we identified 113,175 as lineage-

specific. The alluvial plot in Fig. 5E depicts the new organization of the OM-RGC v2 after being

integrated into our framework and how we can provide context to the two original types of

unknowns in the OM-RGC (those annotated as category S in eggNOG (Huerta-Cepas et al.,

2019) and those without known homologs in the eggNOG database (Salazar et al., 2019)) that

can lead to potential experimental targets at the organism level to complement the

metatranscriptomic approach proposed by Salazar et al. (Salazar et al., 2019).

A structured coding sequence space augments the interpretation
of experimental data

We selected one of the experimental conditions tested in Price et al. (Price et al., 2018) to

demonstrate the potential of our approach to augment experimental data. We compared the

fitness values in plain rich medium with added Spectinomycin dihydrochloride pentahydrate to

the fitness in plain rich medium (LB) in *Pseudomonas fluorescens FW300-N2C3* (Fig. 6A). This

antibiotic inhibits protein synthesis and elongation by binding to the bacterial 30S ribosomal

subunit and interferes with the peptidyl tRNA translocation. We identified the gene with locus id

AO356_08590 that presents a strong phenotype (fitness = -3.1; t = -9.1) and has no known

function. This gene belongs to the genomic unknown GC GU_19737823. We can track this GC

into the environment and explore the occurrence in the different samples we have in our

database. As expected, the GC is mostly found in non-human metagenomes (Fig. 6B) as

Pseudomonas are common inhabitants of soil and water environments(Heffernan, Murphy, &

Casey, 2009). However, finding this GC also in human-related samples is very interesting due

to the potential association of *P. fluorescens* and human disease where Crohn's disease

patients develop serum antibodies to this microbe (Scales, Dickson, LiPuma, & Huffnagle,

2014). We can add another layer of information to the selected GC by looking at the associated

remote homologs in the GCC GU_c_21103 (Fig. 6C). We identified all the genes in the

GTDB_r86 genomes that belong to the GCC GU_c_21103 (Supp. Table 11) and explored their

genomic neighborhoods. All members from GU_c_21103 are constrained to the class

Gammaproteobacteria, and interestingly GU_19737823 is mostly exclusive to the order

Pseudomonadales. The gene order in the different genomes analyzed is highly conserved,

finding GU_19737823 after the *rpsF*::*rpsR* operon and before *rplI*. *rpsF* and *rpsR* encode for

30S ribosomal proteins, the prime target of spectinomycin. The combination of the experimental

evidence and the associated data inferred by our approach provides strong support to generate

the hypothesis that the gene AO356_08590 might be involved in the resistance to

spectinomycin.

Figure 6: Augmenting experimental data with GCs of unknown function. (A) We used the fitness values from the experiments from Price et al.(Price et al., 2018) to identify genes of unknown function that are important for fitness under certain experimental conditions. The selected gene belongs to the genomic unknown GC GU_19737823 and presents a strong phenotype (fitness = -3.1; t = -9.1) (B) Occurrence of GU_19737823 in the metagenomes used in this study. Darker bars depict the number of metagenomes where the GC is found. (C) GU_19737823 is a member of the GCC GU_c_21103. The network shows the relationships between the different GCs members of the gene cluster community GU_c_21103. The size of the node corresponds to the node degree of each GC. Edge thickness corresponds to the bitscore/column metric. Highlighted in red is GU_19737823. (D) We identified all the genes in the GTDB_r86 genomes that belong to the GCC GU_c_21103 and explored their genomic neighborhoods. GU_c_21103 members were constrained to the class Gammaproteobacteria, and GU_19737823 is mostly exclusive to the order Pseudomonadales. The gene order in the different genomes analyzed is highly conserved, finding GU_19737823 after the *rpsF::rpsR* operon and before *rplI*. *rpsF* and *rpsR* encode for the *30S ribosomal protein S6* and *30S ribosomal protein S18,* respectively. The GTDB_r86 subtree only shows RefSeq genomes. Branch colors correspond to the different GCs found in GU_c_21103. The bubble plot depicts the number of genomes with a gene that belongs to GU_c_21103.

Discussion

We present a new conceptual framework and computational workflow to unify the known and unknown CDS-space. Using this framework, we performed an in-depth exploration of the microbial unknown CDS-space, demonstrating that we can link the unknown fraction of metagenomic studies to specific genomes and provide a powerful tool for hypothesis generation. During the last years, the microbiome community has established a standard operating procedure(Quince et al., 2017) for analyzing metagenomes that we can briefly summarize into (1) assembly, (2) gene prediction, (3) gene catalog inference, (4) binning, and (5) characterization. Thanks to recent computational developments (Steinegger & Soding, 2017; Steinegger & Söding, 2018), we envisioned an alternative to this workflow to maximize the information used when analyzing genomic and metagenomic data. In addition, we provide a mechanism to reconcile top-down and bottom-up approaches, thanks to the well-structured CDS-space proposed by our framework. AGNOSTOS can create environmental- and organism-specific variations of a seed GC database. Then, it integrates the predicted genes from new genomes and metagenomes and dynamically creates and classifies new GCs with those genes not integrated during the initial step (Fig. 1B). Afterward, the potential functions of the known GCs can be carefully characterized by incorporating them into the traditional workflows.

One of the most appealing characteristics of our approach is that the GCs provide unified groups of homologous genes across environments and organisms indifferently if they belong to the known or unknown CDS-space, and we can contextualize the unknown fraction using this genomic and environmental information. Our combination of partitioning and contextualization features a smaller unknown CDS-space than we expected. On average, for our genomic and metagenomic data, only 30% of the genes fall in the unknown fraction. One hypothesis to reconcile this surprising finding is that the methodologies to identify remotely homologous sequences in large datasets were computationally prohibitive until recently. New methods

(Steinegger, Meier, et al., 2019; Steinegger & Soding, 2017), like the ones used in AGNOSTOS,

are enabling large scale distant homology searches. Still, one has to apply conservative

measures to control the trade-off between specificity and sensitivity to avoid overclassification.

We found that most of the coding sequence space at the gene and amino acid level is known,

both in genomes and metagenomes. However, it presents a high diversity, as shown in the GC

accumulation curves highlighting the vast remaining untapped microbial fraction and its potential

importance for niche adaptation owing to its narrow ecological distribution. In a genomic context

and after ruling out the effect of prophages, the unknown fraction is predominantly Species'

lineage-specific and phylogenetically more conserved than the known fraction, supporting the

signal observed in the environmental data emphasizing that we should not ignore the unknown

fraction. It is worth noting that the high diversity observed in the unknowns only represents the

20% of the amino acids in the CDS-space we analyzed, and only 10% of this unknown amino

acid space is part of a Pfam domain (DUF and others). This contrasts with the numbers

observed in the known CDS-space, where Pfam domains include 50% of the amino acids. All

this evidence combined strengthens the hypothesis that the genes of unknown function,

especially the lineage-specific ones, might be associated with the mechanisms of microbial

diversification and niche adaptation due to the constant diversification of gene families and the

survival of new gene lineages (Francino, 2012; Muller, 2019).

Metagenome-assembled genomes are not only unveiling new regions of the microbial universe

(42% of the genomes in GTDB_r86), but they are also enriching genes of unknown function in

the tree of life. We investigated the unknown CDS-space of *Cand*. Patescibacteria, more

commonly known as Candidate Phyla Radiation (CPR), a phylum that has raised considerable

interest due to its unusual biology (Brown et al., 2015). We provide a collection of 54,343

lineage-specific GCs of unknown function at different taxonomic level resolutions (Supp. Table

12; Supp. Note 14), which will be a valuable resource for the CPR advancement research

efforts.

Our effort to tackle the unknown provides a pathway to unlock a large pool of likely relevant data

that remains untapped to analysis and discovery. By identifying a potential target gene of

unknown function for antibiotic resistance, we demonstrate the value of our approach and how it

can boost insights from model organism experiments. But severe challenges remain, such as

the dependence on the quality of the assemblies and their gene predictions, as shown by the

analysis of the ribosomal protein GCCs where many of the recovered genes are incomplete.

While sequence assembly has been an active area of research (Roumpeka, Wallace,

Escalettes, Fotheringham, & Watson, 2017), this has not been the case for gene prediction

methods (Roumpeka et al., 2017), which are becoming outdated(Ivanova et al., 2014) and

cannot cope with the current amount of data. Alternatives like protein-level assembly

(Steinegger, Mirdita, & Söding, 2019) combined with exploring the assembly graphs'

neighborhoods (Titus Brown et al., 2018) become very attractive for our purposes. In any case,

we still face the challenge of discriminating between real and artifactual singletons (Höps,

Jeffryes, & Bateman, 2018). There are currently no methods available to provide a plausible

solution and, at the same time, being scalable. We devise a potential solution in the recent

developments in unsupervised deep learning methods where they use large corpora of proteins

to define a language model *embedding* for protein sequences (Heinzinger et al., 2019). These

models could be applied to predict *embeddings* in singletons, which could be clustered or used

to determine their coding potential. Another issue is that we might be creating more GCs than

expected. We follow a conservative approach to avoid mixing multi-domain proteins in GCs

owing to the fragmented nature of the metagenome assemblies that could result in the split of a

GC. However, not only splitting can be a problem, but also lumping unrelated genes or GCs

owing to the use of remote homologies. Although the inference of GCCs is using very sensitive

methods to compare profile HMMs, low sequence diversity in GCs can limit its effectiveness.

Moreover, our approach is affected by the presence and propagation of contamination in

reference databases, a significant problem in 'omics (Breitwieser, Pertea, Zimin, & Salzberg,

2019; Steinegger & Salzberg, 2020). In our case, we only use Pfam as a source for annotation

owing to its high-quality and manual curation process. The categorization process of our GCs

depends on the information from other databases, and to minimize the potential impact of

contamination, we apply methods that weight the annotations of the identified homologs to

discriminate if a GC belongs to the known or unknown CDS-space.

The results presented here prove that the integration and the analysis of the unknown fraction

are possible. We are unveiling a brighter future, not only for microbiome analyses but also for

boosting eukaryotic-related studies, thanks to the increasing number of projects, including

metatranscriptomic data (Delmont et al., 2020; Vorobev et al., 2020). Furthermore, our work

lays the foundations for further developments of clear guidelines and protocols to define the

different levels of unknown (Thomas & Segata, 2019) and should encourage the scientific

community for a collaborative effort to tackle this challenge.

502 Material and methods

503 Genomic and metagenomic dataset

We used a set of 583 marine metagenomes from four of the major metagenomic surveys of the

ocean microbiome: Tara Oceans expedition (TARA) (Sunagawa et al., 2015), Malaspina

expedition (Duarte, 2015), Ocean Sampling Day (OSD) (Kopf et al., 2015), and Global Ocean

Sampling Expedition (GOS) (Rusch et al., 2007). We complemented this set with 1,246

metagenomes obtained from the Human Microbiome Project (HMP) phase I and II (Lloyd-Price

et al., 2017). We used the assemblies provided by TARA, Malaspina, OSD and HMP projects

and the long Sanger reads from GOS (Sanger, Nicklen, & Coulson, 1977). A total of 156M

(156,422,969) contigs and 12.8M long-reads were collected (Supp. Table 6).

For the genomic dataset, we used the 28,941 prokaryotic genomes (27,372 bacterial and 1,569

archaeal) from the Genome Taxonomy Database (Parks et al., 2018) (GTDB) Release 03-RS86

(19th August 2018).

Computational workflow development

We implemented a computation workflow based on Snakemake (Köster, 2018) for the easy

processing of large datasets in a reproducible manner. The workflow provides three different

strategies to analyze the data. The module *DB-creation* creates the gene cluster database,

validates and partitions the gene clusters (GCs) in the main functional categories. The module

DB-update allows the integration of new sequences (either at the contig or predicted gene level)

in the existing gene cluster database. In addition, the workflow has a *profile-search* function to

quickly screen samples using the gene cluster PSSM profiles in the database.

Metagenomic and genomic gene prediction

We used Prodigal (v2.6.3) (Hyatt et al., 2010) in metagenomic mode to predict the genes from

the metagenomic dataset. For the genomic dataset, we used the gene predictions provided by

Annotree (Mendler et al., 2019), since they were obtained, consistently, with Prodigal v2.6.3.

We identified potential spurious genes using the *AntiFam* database (Eberhardt et al., 2012).

Furthermore, we screened for '*shadow*' genes using the procedure described in Yooseph et al.

(Yooseph, Li, & Sutton, 2008).

PFAM annotation

We annotated the predicted genes using the *hmmsearch* program from the *HMMER* package

(version: 3.1b2) (R. D. Finn, Clements, & Eddy, 2011) in combination with the Pfam database

v31 (Robert D. Finn et al., 2016). We kept the matches exceeding the internal gathering

threshold and presenting an independent e-value < 1e-5 and coverage > 0.4. In addition, we

took into account multi-domain annotations, and we removed overlapping annotations when the

overlap is larger than 50%, keeping the ones with the smaller e-value.

537   Determination of the gene clusters

We clustered the metagenomic predicted genes using the cascaded-clustering workflow of the

MMseqs2 software (Steinegger & Söding, 2018) ("*--cov-mode 0 -c 0.8 --min-seq-id 0.3"*). We

discarded from downstream analyses the singletons and clusters with a size below a threshold

identified after applying a broken-stick model (Bennett, 1996). We integrated the genomic data

into the metagenomic cluster database using the "DB-update" module of the workflow. This

module uses the *clusterupdate* module of MMseqs2 (Steinegger & Soding, 2017), with the same

parameters used for the metagenomic clustering.

545   Quality-screening of gene clusters

We examined the GCs to ensure their high intra-cluster homogeneity. We applied two

methodologies to validate their cluster sequence composition and functional annotation

homogeneity. We identified non-homologous sequences inside each cluster combining the

identification of a new cluster representative sequence via a sequence similarity network (SSN)

analysis, and the investigation of intra-cluster multiple sequence alignments (MSAs), given the

new representative. Initially, we generated an SSN for each cluster, using the semi-global

alignment methods implemented in *PARASAIL* (Daily, 2016) (version 2.1.5). We trimmed the

SSN using a custom algorithm (Chafee et al., 2018; Žure, Fernandez-Guerra, Munn, & Harder,

2017) that removes edges while maintaining the network structural integrity and obtaining the

smallest connected graph formed by a single component. Finally, the new cluster representative

was identified as the most central node of the trimmed SSN by the eigenvector centrality

algorithm, as implemented in igraph (Csardi & Nepusz, 2006). After this step, we built a multiple

sequence alignment for each cluster using *FAMSA* (Deorowicz, Debudaj-Grabysz, & Gudyś,

2016) (version 1.1). Then, we screened each cluster-MSA for non-homologous sequences to

the new cluster representative. Owing to computational limitations, we used two different

approaches to evaluate the cluster-MSAs. We used *LEON-BIS* (Vanhoutreve et al., 2016) for

the clusters with a size ranging from 10 to 1,000 genes and OD-SEQ (Jehl, Sievers, & Higgins,

2015) for the clusters with more than 1,000 genes. In the end, we applied a broken-stick model

(Bennett, 1996) to determine the threshold to discard a cluster.

The predicted genes can have multi-domain annotations in different orders, therefore to validate

the consistency of intra-cluster Pfam annotations, we applied a combination of w-shingling

(Broder, 1997) and Jaccard similarity. We used w-shingling (k-shingle = 2) to group consecutive

domain annotations as a single object. We measured the homogeneity of the *shingle sets* (sets

of domains) between genes using the Jaccard similarity and reported the median similarity

value for each cluster. Moreover, we took into consideration the Clan membership of the Pfam

domains and that a gene might contain N-, C- and M-terminal domains for the functional

homogeneity validation. We discarded clusters with a median similarity < 1.

After the validation, we refined the gene cluster database removing the clusters identified to be

discarded and the clusters containing ≥ 30% *shadow genes*. Lastly, we removed the single

shadow, spurious and non-homologous genes from the remaining clusters (Supplementary Note

2).

Remote homology classification of gene clusters

To partition the validated GCs into the four main categories, we processed the set of GCs

containing Pfam annotated genes and the set of not annotated GCs separately. For the

annotated GCs, we inferred a consensus protein domain architecture (DA) (an ordered

combination of protein domains) for each annotated gene cluster. To identify each gene cluster

consensus DA, we created directed acyclic graphs connecting the Pfam domains based on their

topological order on the genes using *igraph* (Csardi & Nepusz, 2006). We collapsed the

repetitions of the same domain. Then we used the gene completeness as a positive-weighting

value for the selection of the cluster consensus DA. Within this step, we divided the GCs into

"Knowns" (Known) if annotated to at least one Pfam domains of known function (DKFs) and

"Genomic unknowns" (GU) if annotated entirely to Pfam domains of unknown function (DUFs).

We aligned the sequences of the non-annotated GCs with FAMSA (Deorowicz et al., 2016) and

obtained cluster consensus sequences with the *hhconsensus* program from *HH-SUITE*

(Steinegger, Meier, et al., 2019). We used the cluster consensus sequences to perform a

nested search against the UniRef90 database (release 2017_11) (The UniProt Consortium,

2017) and NCBI *nr* database (release 2017_12) (NCBI Resource Coordinators, 2018) to retrieve

non-Pfam annotations with *MMSeqs2* (Steinegger & Soding, 2017) ("*-e 1e-05 --cov-mode 2 -c*

0.6"). We kept the hits within 60% of the Log(best-e-value) and searched the annotations for

any of the terms commonly used to define proteins of unknown function (Supp. Table 12). We

used a quorum majority voting approach to decide if a gene cluster would be classified as

Genomic Unknown or *Known without Pfams* based on the annotations retrieved. We searched

the consensus sequences without any homologs in the UniRef90 database against NCBI *nr*. We

applied the same approach and criteria described for the first search. Ultimately, we classified

as *Environmental Unknown* those GCs whose consensus sequences did not align with any of

the NCBI *nr* entries.

In addition, we developed some conservative measures to control the trade-off between

specificity and sensitivity for the remote homology searches such as (1) a modification of the

algorithm described in Hingamp et al. (Hingamp et al., 2013) to get a confident group of

homologs to determine if a query protein is known or unknown by a quorum majority voting

approach (Supp Note 3); (2) strict parameters in terms of iterations, bidirectional coverage and

probability thresholds for the HHblits alignments to minimize the inclusion of non-homologous

sequences; and (3) avoid providing annotations for our gene clusters, as we believe that

annotation should be a careful process done on a smaller scale and with experimental context.

Gene cluster remote homology refinement

We refined the *Environmental Unknown* GCs to ensure the lack of any characterization by

searching for remote homologies in the Uniclust database (release 30_2017_10) using the

HMM/HMM alignment method *HHblits* (Remmert, Biegert, Hauser, & Söding, 2012). We created

the HMM profiles with the *hhmake* program from the *HH-SUITE* (Steinegger, Meier, et al.,

2019). We only accepted those hits with an *HHblits-probability* ≥ 90% and we re-classified them

following the same majority vote approach as previously described. The clusters with no hits

remained as the refined set of EUs. We applied a similar refinement approach to the KWP

clusters to identify GCs with remote homologies to Pfam protein domains. The KWP HMM

profiles were searched against the Pfam *HH-SUITE* database (version 31), using *HHblits*. We

accepted hits with a probability ≥ 90% and a target coverage > 60% and removed overlapping

domains as described earlier. We moved the KWP with remote homologies to known Pfams to

the Known set, and those showing remote homologies to Pfam DUFs to the GUs. The clusters

with no hits remained as the refined set of KWP.

Gene cluster characterization

To retrieve the taxonomic composition of our clusters we applied the *MMseqs2 taxonomy*

program (version: b43de8b7559a3b45c8e5e9e02cb3023dd339231a), which allows computing

the lowest common ancestor through the implementation of the 2bLCA protocol (Hingamp et al.,

2013). We searched all cluster genes against UniProtKB (release of January 2018) (UniProt

Consortium, 2018) using the following parameters "*-e 1e-05 --cov-mode 0 -c 0.6*". We parsed

the results to keep only the hits within 60% of the log10(best-e-value). To retrieve the taxonomic

lineages, we used the R package *CHNOSZ* (Dick, 2008). We measured the intra-cluster

taxonomic admixture by applying the *entropy.empirical()* function from the *entropy* R package

(Hausser & Strimmer, 2008). This function estimates the Shannon entropy based on the

different taxonomic annotation frequencies. For each cluster, we also retrieved the cluster

consensus taxonomic annotation, which we defined as the taxonomic annotation of the majority

of the genes in the cluster.

In addition to the taxonomy, we evaluated the clusters' level of darkness and disorder using the

Dark Proteome Database (DPD) (Perdigão et al., 2017) as reference. We searched the cluster

genes against the DPD, applying the MMseqs2 search program (Steinegger & Soding, 2017)

with "*-e 1e-20 --cov-mode 0 -c 0.6*". For each cluster, we then retrieved the mean and the

median level of darkness, based on the gene DPD annotations.

High-quality clusters

643 We defined a subset of high-quality clusters based on the completeness of the cluster genes

644 and their representatives. We identified the minimum required percentage of complete genes

645 per cluster by a broken-stick model (Bennett, 1996) applied to the percentage distribution. Then,

646 we selected the GCs found above the threshold and with a complete representative.

A set of non-redundant domain architectures

We estimated the number of potential domain architectures present in the *Known* GCs taking

into account the large proportion of fragmented genes in the metagenomic dataset and that

could inflate the number of potential domain architectures. To identify fragments of larger

domain architecture, we took into account their topological order in the genes. To reduce the

number of comparisons, we calculated the pairwise string cosine distance (q-gram = 3) between

domain architectures and discarded the pairs that were too divergent (cosine distance ≥ 0.9).

We collapsed a fragmented domain architecture to the larger one when it contained less than

75% of complete genes.

Inference of gene cluster communities

We aggregated distant homologous GCs into GCCs. The community inference approach combined an all-vs-all HMM gene cluster comparison with Markov Cluster Algorithm (MCL) (van Dongen & Abreu-Goodger, 2012) community identification. We started performing the inference on the Known GCs to use the Pfam DAs as constraints. We aligned the gene cluster HMMs using HHblits (Remmert et al., 2012) (-n 2 -Z 10000000 -B 10000000 -e 1) and we built a homology graph using the cluster pairs with probability ≥ 50% and bidirectional coverage > 60%. We used the ratio between HHblits-bitscore and aligned-columns as the edge weights (Supp. Note 9). We used MCL (van Dongen & Abreu-Goodger, 2012) (v. 12-068) to identify the communities present in the graph. We developed an iterative method to determine the optimal MCL inflation parameter that tries to maximize the relationship of five intra-/inter-community properties: (1) the proportion of MCL communities with one single DA, based on the consensus DAs of the cluster members; (2) the ratio of MCL communities with more than one cluster; (3) the proportion of MCL communities with a PFAM clan entropy equal to 0; (4) the intra-community HHblits-score/Aligned-columns score (normalized by the maximum value); and (5) the number of MCL communities, which should, in the end, reflect the number of non-redundant DAs. We iterated through values ranging from 1.2 to 3.0, with incremental steps of 0.1. During the inference process, some of the GCs became orphans in the graph. We applied a three-step approach to assigning a community membership to these GCs. First, we used less stringent conditions (probability ≥ 50% and coverage >= 40%) to find homologs in the already existing GCCs. Then, we ran a second iteration to find secondary relationships between the newly assigned GCs and the missing ones. Lastly, we created new communities with the remaining GCs. We repeated the whole process with the other categories (KWP, GU and EU), applying the optimal inflation value found for the Known (2.2 for metagenomic and 2.5 for genomic data).

Gene cluster communities validation

We tested the biological significance of the GCCs using the phylogeny of proteorhodopsin

(Boeuf et al., 2015) (PR). We used the proteorhodopsin HMM profiles (Olson et al., 2018) to

screen the marine metagenomic datasets using *hmmsearch* (version 3.1b2) (R. D. Finn et al.,

2011). We kept the hits with a coverage > 0.4 and e-value <= 1e-5. We removed identical

duplicates from the sequences assigned to PR with CD-HIT (W. Li & Godzik, 2006) (v4.6) and

cleaned from sequences with less than 100 amino acids. To place the identified PR sequences

into the MicRhode (Boeuf et al., 2015) PR tree first, we optimized the initial tree parameters and

branch lengths with RAxML (v8.2.12) (Stamatakis, 2014). We used PaPaRA (v2.5) (Berger &

Stamatakis, 2012) to incrementally align the query PR sequences against the MicRhode PR

reference alignment and *pplacer* (Matsen, Kodner, & Armbrust, 2010) (v1.1.alpha19-0-g807f6f3)

to place the sequences into the tree. Finally, we assigned the query PR sequences to the

MicRhode PR Superclusters based on the phylogenetic placement. We further investigated the

GCs annotated as viral (196 genes, 14 GC) comparing them to the six newly discovered viral

PRs (Needham et al., 2019) using Parasail (Daily, 2016) (-a sg_stats_scan_sse2_128_16 -t 8 -c

1 -x). As an additional evaluation, we investigated the distributions of standard GCCs and HQ

GCCs within ribosomal protein families. We obtained the ribosomal proteins used for the

analysis combining the set of 16 ribosomal proteins from Méheust et al. (Méheust et al., 2019)

and those contained in the collection of bacterial single-copy genes of Anvi'o (Murat Eren et al.,

2015). Also, for the ribosomal proteins, we compared the outcome of our method to the one

proposed by Méheust et al. (Méheust et al., 2019) (Supp. Note 9).

Metagenomic sample selection for downstream analyses

For the subsequent ecological analyses, we selected those metagenomes with a number of genes larger or equal to the first quartile of the distribution of all the metagenomic gene counts. (Supp. Table 13).

Gene cluster abundance profiles in genomes and metagenomes

We estimated abundance profiles for the metagenomic cluster categories using the read coverage to each predicted gene as a proxy for abundance. We calculated the coverage by mapping the reads against the assembly contigs using the *bwa-mem* algorithm from *BWA mapper* (H. Li & Durbin, 2010). Then, we used *BEDTOOLS* (Quinlan & Hall, 2010), to find the intersection of the gene coordinates to the assemblies, and normalize the per-base coverage by the length of the gene. We calculated the cluster abundance in a sample as the sum of the cluster gene abundances in that sample, and the cluster category abundance in a sample as the sum of the cluster abundances. We obtained the proportions of the different gene cluster categories applying a total-sum-scaling normalization. For the genomic abundance profiles, we used the number of genes in the genomes and normalized by the total gene counts per genome.

Rate of genomic and metagenomic gene clusters accumulation

We calculated the cumulative number of known and unknown GCs as a function of the number of metagenomes and genomes. For each metagenome count, we generated 1000 random sets, and we calculated the number of GCs and GCCs recovered. For this analysis, we used 1,246 HMP metagenomes and 358 marine metagenomes (242 from TARA and 116 from Malaspina). We repeated the same procedure for the genomic dataset. We removed the singletons from the metagenomic dataset with an abundance smaller than the mode abundance of the singletons

724   that got reclassified as good-quality clusters after integrating the GTDB data to minimize the

725   impact of potential spurious singletons. To complement those analyses, we evaluated the

726   coverage of our dataset by searching seven different state-of-the-art databases against our set

727   of metagenomic GC HMM profiles (Supp. Note 12).

729   Occurrence of gene clusters in the environment

We used 1,264 metagenomes from the TARA Oceans, MALASPINA Expedition, OSD2014 and

HMP-I/II to explore the properties of the unknown CDS-space in the environment. We applied

the Levins Niche Breadth (NB) index (Levins, 1966) to investigate the GCs and GCCs

environmental distributions. We removed the GCs and cluster communities with a mean relative

abundance < 1e-5. We followed a divide-and-conquer strategy to avoid the computational

burden of generating the null-models to test the significance of the distributions owing to the

large number of metagenomes and GCs. First, we grouped similar samples based on the gene

cluster content using the Bray-Curtis dissimilarity(Bray, Roger Bray, & Curtis, 1957) in

combination with the *Dynamic Tree Cut* (Langfelder, Zhang, & Horvath, 2008) R package. We

created 100 random datasets picking up one random sample from each group. For each of the

100 random datasets, we created 100 random abundance matrices using the *nullmodel* function

of the *quasiswap* count method (Miklós & Podani, 2004). Then we calculated the *observed* NB

and obtained the 2.5% and 97.5% quantiles based on the randomized sets. We compared the

observed and quantile values for each gene cluster and defined it to have a *Narrow distribution*

when the *observed* was smaller than the 2.5% quantile and to have a *Broad distribution* when it

was larger than the 97.5% quantile. Otherwise, we classified the cluster as *Non-significant*

(Salazar et al., 2015). We used a majority voting approach to get a consensus distribution

classification based on the ten random datasets.

Identification of prophages in genomic sequences

We used PhageBoost (https://github.com/ku-cbd/PhageBoost/) to find gene regions in the microbial genomes that result in high viral signals against the overall genome signal. We set the following thresholds to consider a region prophage: minimum of 10 genes, maximum 5 gaps, single-gene probability threshold 0.9. We further smoothed the predictions using Parzen rolling windows of 20 periods and looked at the smoothed probability distribution across the genome. We disregarded regions that had a summed smoothed probability less than 0.5, and those regions that did differ from the overall population of the genes in a genome by using Kruskal–Wallis rank test (p-value 0.001).

Lineage-specific gene clusters

We used the F1-score developed for AnnoTree (Mendler et al., 2019) to identify the lineage-specific GCs and to which rank they are specific. Following similar criteria to the ones used in Mendler et al. (Mendler et al., 2019), we considered a gene cluster to be lineage-specific if it is present in less than half of all genomes and at least 2 with F1-score > 0.95.

Phylogenetic conservation of gene clusters

We calculated the phylogenetic conservation (τD) of each gene cluster using the *consenTRAIT* (Martiny et al., 2013) function implemented in the R package *castor* (Martiny et al., 2013). We used a paired Wilcoxon rank-sum test to compare the average τD values for lineage-specific and non-specific GCs.

Evaluation of the OM-RGC v2 uncharacterized fraction

We integrated the 46,775,154 genes from the second version of the TARA Ocean Microbial Reference Gene Catalog (OM-RGC v2) (Salazar et al., 2019) into our cluster database using

770   the same procedure as for the genomic data. We evaluated the uncharacterized fraction and the

771   genes classified into the eggNOG (Huerta-Cepas et al., 2019) category S within the context of

772   our database.

Augmenting RB-TnSeq experimental data

We searched the 37,684 genes of unknown function associated with mutant phenotypes from

Price et al. (Price et al., 2018) against our gene cluster profiles. We kept the hits with e-value ≤

1e-20 and a query coverage > 60%. Then we filtered the results to keep the hits within 90% of

the Log(best-e-value), and we used a majority vote function to retrieve the consensus category

for each hit. Lastly, we selected the best-hits based on the smallest e-value and the largest

query and target coverage values. We used the fitness values from the RB-TnSeq experiments

from Price et al. to identify genes of unknown function that are important for fitness under

certain experimental conditions.

Availability of data and materials

The code used for the analyses in the manuscript is available at https://github.com/functional-

dark-side/functional-dark-side.github.io/tree/master/scripts. The code to recreate the figures is

available at https://github.com/functional-dark-side/vanni_et_al-figures. Detailed descriptions of

the different methods and results of this manuscript are available at

https://dark.metagenomics.eu. The workflow AGNOSTOS is available at

https://github.com/functional-dark-side/agnostos-wf, and its database can be downloaded from

https://doi.org/10.6084/m9.figshare.12459056.

Acknowledgements

The authors thankfully acknowledge the computer resources at MareNostrum and the technical

support provided by Barcelona Supercomputing Center (RES-AECT-2014-2-0085), the BMBF-

funded de.NBI Cloud within the German Network for Bioinformatics Infrastructure (de.NBI)

(031A537B, 031A533A, 031A538A, 031A533B, 031A535A, 031A537C, 031A534A, 031A532B),

the University of Oxford Advanced Research Computing

(http://dx.doi.org/10.5281/zenodo.22558) and the MARBITS bioinformatics core at ICM-CSIC.

CV was supported by the Max Planck Society. AFG received funding from the European

Union's Horizon 2020 research and innovation program Blue Growth: Unlocking the potential of

Seas and Oceans under grant agreement no. 634486 (project acronym INMARE). AM was

supported by the Biotechnology and Biological Sciences Research Council [BB/M011755/1,

BB/R015228/1] and RDF by the European Molecular Biology Laboratory core funds. EOC was

supported by project INTERACTOMA RTI2018-101205-B-I00 from the Spanish Agency of

Science MICIU/AEI. SGA and PS received additional funding by the project MAGGY

(CTM2017-87736-R) from the Spanish Ministry of Economy and Competitiveness. The

Malaspina 2010 Expedition was supported by the Spanish Ministry of Economy and

Competitiveness (MINECO) through the Consolider-Ingenio program (ref. CSD2008-00077).

The authors thank Johannes Söding and Alex Bateman for helpful discussions.

Contributions

CV, MSS and AF-G performed the analyses and wrote the computational workflow. MS assisted

with the clustering and remote homology searches. KS helped with the identification of

prophages in genomic sequences. PLB and AB provided feedback and assisted with the

ecological analyses. RDF and AM provided feedback and information on the MGnify and Pfam

816   databases. CMD, PS and SGA provided the Malaspina metagenomes. TOD and AME analyzed

817   data in the context of metagenome-assembled genomes. AF-G conceived the study and

818   supervised the work. CV and AF-G wrote the manuscript. All authors read, edited and approved

819   the final manuscript.

820   Competing Interests

821   The authors declare no competing interests.

References

Almeida, A., Mitchell, A. L., Boland, M., Forster, S. C., Gloor, G. B., Tarkowska, A., … Finn, R.

D. (2019). A new genomic blueprint of the human gut microbiota. *Nature*, *568*(7753),

499–504.

Almeida, A., Nayfach, S., Boland, M., Strozzi, F., Beracochea, M., Shi, Z. J., … Finn, R. D.

(2020). A unified catalog of 204,938 reference genomes from the human gut

microbiome. *Nature Biotechnology*. doi:10.1038/s41587-020-0603-3

Anantharaman, K., Hausmann, B., Jungbluth, S. P., Kantor, R. S., Lavy, A., Warren, L. A., …

Banfield, J. F. (2018). Expanded diversity of microbial groups that shape the

dissimilatory sulfur cycle. *The ISME Journal*, *12*(7), 1715–1728.

Arnold, F. H. (1998). Design by Directed Evolution. *Accounts of Chemical Research*, *31*(3),

125–131.

Arnold, F. H. (2018). Directed Evolution: Bringing New Chemistry to Life. *Angewandte Chemie* ,

57(16), 4143–4148.

Bateman, A., Coggill, P., & Finn, D. R. (2010). DUFs: Families in search of function. *Acta*

Crystallographica. Section F, Structural Biology and Crystallization Communications,

66(10), 1148–1152.

Bennett, K. D. (1996). Determination of the number of zones in a biostratigraphical sequence.

The New Phytologist, *132*(1), 155–170.

Berger, S. A., & Stamatakis, A. (2012). PaPaRa 2.0: a vectorized algorithm for probabilistic

phylogeny-aware alignment extension. *Heidelberg Institute for Theoretical Studies,*

Http://Sco. h-Its. Org/Exelixis/Publica Tions. Html. Exelixis-RRDR-2012-2015. Retrieved

from https://cme.h-its.org/exelixis/pubs/Exelixis-RRDR-2012-5.pdf

Bernard, G., Pathmanathan, J. S., Lannes, R., Lopez, P., & Bapteste, E. (2018). Microbial Dark

Matter Investigations: How Microbial Studies Transform Biological Knowledge and

Empirically Sketch a Logic of Scientific Discovery. *Genome Biology and Evolution*, *10*(3),

707–715.

Bileschi, M. L., Belanger, D., Bryant, D., Sanderson, T., Carter, B., Sculley, D., … Colwell, L. J.

(2019). Using Deep Learning to Annotate the Protein Universe (p. 626507).

doi:10.1101/626507

Bitard-Feildel, T., & Callebaut, I. (2017). Exploring the dark foldable proteome by considering

hydrophobic amino acids topology. *Scientific Reports*, *7*, 41425.

Boeuf, D., Audic, S., Brillet-Guéguen, L., Caron, C., & Jeanthon, C. (2015). MicRhoDE: a

curated database for the analysis of microbial rhodopsin diversity and evolution.

Database: The Journal of Biological Databases and Curation, *2015*.

doi:10.1093/database/bav080

Brandenberg, O. F., Fasan, R., & Arnold, F. H. (2017). Exploiting and engineering hemoproteins

for abiological carbene and nitrene transfer reactions. *Current Opinion in Biotechnology*,

47, 102–111.

Bray, J. R., Roger Bray, J., & Curtis, J. T. (1957). An Ordination of the Upland Forest

Communities of Southern Wisconsin. *Ecological Monographs*, Vol. 27, pp. 325–349.

doi:10.2307/1942268

Breitwieser, F. P., Pertea, M., Zimin, A., & Salzberg, S. L. (2019). Human contamination in

bacterial genomes has created thousands of spurious proteins. *Genome Research*.

doi:10.1101/gr.245373.118

Broder, A. Z. (1997). On the resemblance and containment of documents. *Proceedings.*

Compression and Complexity of SEQUENCES 1997 (Cat. No. 97TB100171), 21–29.

IEEE.

Brown, C. T., Hug, L. A., Thomas, B. C., Sharon, I., Castelle, C. J., Singh, A., … Banfield, J. F.

(2015). Unusual biology across a group comprising more than 15% of domain Bacteria.

Nature, *523*(7559), 208–211.

Brum, J. R., Cesar Ignacio-Espinoza, J., Kim, E.-H., Trubl, G., Jones, R. M., Roux, S., …

Sullivan, M. B. (2016). Illuminating structural proteins in viral "dark matter" with

metaproteomics. *Proceedings of the National Academy of Sciences of the United States*

of America, *113*(9), 2436–2441.

Buttigieg, L. P., Hankeln, W., Kostadinov, I., Kottmann, R., Yilmaz, P., Duhaime, B. M., &

Gl??ckner, O. F. (2013). Ecogenomic Perspectives on Domains of Unknown Function:

Correlation-Based Exploration of Marine Metagenomes. *PloS One*, *8*(3).

Carradec, Q., Pelletier, E., Da Silva, C., Alberti, A., Seeleuthner, Y., Blanc-Mathieu, R., …

Wincker, P. (2018). A global ocean atlas of eukaryotic genes. *Nature Communications*,

9(1), 373.

Chafee, M., Fernàndez-Guerra, A., Buttigieg, P. L., Gerdts, G., Eren, A. M., Teeling, H., &

Amann, R. I. (2018). Recurrent patterns of microdiversity in a temperate coastal marine

environment. *The ISME Journal*, *12*(1), 237–252.

Chen, I.-M. A., Chu, K., Palaniappan, K., Pillay, M., Ratner, A., Huang, J., … Kyrpides, N. C.

(2019). IMG/M v.5.0: an integrated data management and comparative analysis system

for microbial genomes and microbiomes. *Nucleic Acids Research*, *47*(D1), D666–D677.

Cross, K. L., Campbell, J. H., Balachandran, M., Campbell, A. G., Cooper, S. J., Griffen, A., …

Podar, M. (2019). Targeted isolation and cultivation of uncultivated bacteria by reverse

genomics. *Nature Biotechnology*, *37*(11), 1314–1321.

Csardi, G., & Nepusz, T. (2006). The igraph software package for complex network research.

InterJournal, p. 1695. Retrieved from http://igraph.org

Daily, J. (2016). Parasail: SIMD C library for global, semi-global, and local pairwise sequence

alignments. *BMC Bioinformatics*, *17*(1), 81–81.

Delmont, T. O., Gaia, M., Hinsinger, D. D., Fremont, P., Guerra, A. F., Murat Eren, A., … Jaillon,

O. (2020). Functional repertoire convergence of distantly related eukaryotic plankton

lineages revealed by genome-resolved metagenomics (p. 2020.10.15.341214).

doi:10.1101/2020.10.15.341214

Deorowicz, S., Debudaj-Grabysz, A., & Gudyś, A. (2016). FAMSA: Fast and accurate multiple

sequence alignment of huge protein families. *Scientific Reports*, *6*(1), 33964–33964.

Dick, J. M. (2008). Calculation of the relative metastabilities of proteins using the CHNOSZ

software package. *Geochemical Transactions*, *9*, 10.

Duarte, C. M. (2015). Seafaring in the 21St Century: The Malaspina 2010 Circumnavigation

Expedition. *Limnology and Oceanography Bulletin*, *24*(1), 11–14.

Dubinkina, V. B., Ischenko, D. S., Ulyantsev, V. I., Tyakht, A. V., & Alexeev, D. G. (2016).

Assessment of k-mer spectrum applicability for metagenomic dissimilarity analysis. *BMC*

Bioinformatics, Vol. 17. doi:10.1186/s12859-015-0875-7

Eberhardt, R. Y., Haft, D. H., Punta, M., Martin, M., O'Donovan, C., & Bateman, A. (2012).

AntiFam: a tool to help identify spurious ORFs in protein annotation. *Database*, *2012*(0),

bas003–bas003.

Edwards, R. A., Vega, A. A., Norman, H. M., Ohaeri, M., Levi, K., Dinsdale, E. A., … Dutilh, B.

E. (2019). Global phylogeography and ancient evolution of the widespread human gut

virus crAssphage. *Nature Microbiology*, *4*(10), 1727–1736.

Eloe-Fadrosh, E. A., Paez-Espino, D., Jarett, J., Dunfield, P. F., Hedlund, B. P., Dekas, A. E., …

Ivanova, N. N. (2016). Global metagenomic survey reveals a new bacterial candidate

phylum in geothermal springs. *Nature Communications*, *7*, 10476.

Finn, R. D., Clements, J., & Eddy, S. R. (2011). HMMER web server: interactive sequence

similarity searching. *Nucleic Acids Research*, *39*(suppl), W29–W37.

Finn, Robert D., Coggill, P., Eberhardt, R. Y., Eddy, S. R., Mistry, J., Mitchell, A. L., … Bateman,

A. (2016). The Pfam protein families database: towards a more sustainable future.

Nucleic Acids Research, *44*(D1), D279–D285.

Francino, M. P. (2012). The ecology of bacterial genes and the survival of the new. *International*

Journal of Evolutionary Biology, *2012*, 394026.

Franzosa, E. A., McIver, L. J., Rahnavard, G., Thompson, L. R., Schirmer, M., Weingart, G., …

Huttenhower, C. (2018). Species-level functional profiling of metagenomes and

metatranscriptomes. *Nature Methods*, *15*(11), 962–968.

Habchi, J., Tompa, P., Longhi, S., & Uversky, V. N. (2014). Introducing protein intrinsic disorder.

Chemical Reviews, *114*(13), 6561–6588.

Hanson, A. D., Pribat, A., Waller, J. C., & Crécy-Lagard, V. de. (2010). 'Unknown'proteins and

'orphan'enzymes: the missing half of the engineering parts list--and how to find it.

Biochemical Journal, *425*(1), 1–11.

Hausser, J., & Strimmer, K. (2008). Entropy inference and the James-Stein estimator, with

application to nonlinear gene association networks. Retrieved from

http://arxiv.org/abs/0811.3579

Heffernan, B., Murphy, C. D., & Casey, E. (2009). Comparison of planktonic and biofilm cultures

of Pseudomonas fluorescens DSM 8341 cells grown on fluoroacetate. *Applied and*

Environmental Microbiology, *75*(9), 2899–2907.

Heinzinger, M., Elnaggar, A., Wang, Y., Dallago, C., Nechaev, D., Matthes, F., & Rost, B.

(2019). Modeling aspects of the language of life through transfer-learning protein

sequences. *BMC Bioinformatics*, *20*(1), 723.

Hingamp, P., Grimsley, N., Acinas, S. G., Clerissi, C., Subirana, L., Poulain, J., … Ogata, H.

(2013). Exploring nucleo-cytoplasmic large DNA viruses in Tara Oceans microbial

metagenomes. *The ISME Journal*, *7*(9), 1678–1695.

Höps, W., Jeffryes, M., & Bateman, A. (2018). Gene Unprediction with Spurio: A tool to identify

spurious protein sequences. *F1000Research*, *7*, 261.

Huerta-Cepas, J., Forslund, K., Coelho, L. P., Szklarczyk, D., Jensen, L. J., von Mering, C., &

Bork, P. (2017). Fast Genome-Wide Functional Annotation through Orthology

Assignment by eggNOG-Mapper. *Molecular Biology and Evolution*, *34*(8), 2115–2122.

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K., Cook, H., …

Bork, P. (2019). eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated

orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids*

Research, *47*(D1), D309–D314.

Hug, L. A., Baker, B. J., Anantharaman, K., Brown, C. T., Probst, A. J., Castelle, C. J., …

Banfield, J. F. (2016). A new view of the tree of life. *Nature Microbiology*, *1*, 16048.

Hyatt, D., Chen, G.-L., LoCascio, P. F., Land, M. L., Larimer, F. W., & Hauser, L. J. (2010).

Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC*

Bioinformatics, *11*(1), 119–119.

Ivanova, N. N., Schwientek, P., Tripp, H. J., Rinke, C., Pati, A., Huntemann, M., … Rubin, E. M.

(2014). Stop codon reassignments in the wild. *Science*, *344*(6186), 909–913.

Jaroszewski, L., Li, Z., Krishna, S. S., Bakolitsa, C., Wooley, J., Deacon, M. A., … Godzik, A.

(2009). Exploration of uncharted regions of the protein universe. *PLoS Biology*, *7*(9).

Jehl, P., Sievers, F., & Higgins, D. G. (2015). OD-seq: outlier detection in multiple sequence

alignments. *BMC Bioinformatics*, *16*(1), 269–269.

Kopf, A., Bicak, M., Kottmann, R., Schnetzer, J., Kostadinov, I., Lehmann, K., … Glöckner, F. O.

(2015). The ocean sampling day consortium. *GigaScience*, *4*, 27.

Köster, J. (2018). Reproducible data analysis with Snakemake. *F1000Research*, *7*. Retrieved

from http://www.dodsc.tu-

dortmund.de/cms/Medienpool/files/002_Kolloquium/09_Koester.pdf

La Cono, V., La Spada, G., Arcadi, E., Placenti, F., Smedile, F., Ruggeri, G., … Yakimov, M. M.

(2013). Partaking of Archaea to biogeochemical cycling in oxygen-deficient zones of

meromictic saline Lake Faro (Messina, Italy). *Environmental Microbiology*, *15*(6), 1717–

1733.

Langfelder, P., Zhang, B., & Horvath, S. (2008). Defining clusters from a hierarchical cluster

tree: the Dynamic Tree Cut package for R. *Bioinformatics* , *24*(5), 719–720.

Levins, R. (1966). THE STRATEGY OF MODEL BUILDING IN POPULATION BIOLOGY.

American Scientist, *54*(4), 421–431.

Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows–Wheeler

transform. *Bioinformatics* , *26*(5), 589–595.

Li, W., & Godzik, A. (2006). Cd-hit: a fast program for clustering and comparing large sets of

protein or nucleotide sequences. *Bioinformatics* , *22*(13), 1658–1659.

Liu, X. L. (2017). Deep Recurrent Neural Network for Protein Function Prediction from

Sequence (p. 103994). doi:10.1101/103994

Lloyd-Price, J., Mahurkar, A., Rahnavard, G., Crabtree, J., Orvis, J., Hall, A. B., … Huttenhower,

C. (2017). Strains, functions and dynamics in the expanded Human Microbiome Project.

Nature, *550*(7674), 61–66.

Lobb, B., Kurtz, D. A., Moreno-Hagelsieb, G., & Doxey, A. C. (2015). Remote homology and the

functions of metagenomic dark matter. *Frontiers in Genetics*, *6*(JUL), 1–12.

Ma, Y., Madupu, R., Karaoz, U., Nossa, C. W., Yang, L., Yooseph, S., … Pei, Z. (2014). Human

papillomavirus community in healthy persons, defined by metagenomics analysis of

human microbiome project shotgun sequencing data sets. *Journal of Virology*, *88*(9),

4786–4797.

Martiny, A. C., Treseder, K., & Pusch, G. (2013). Phylogenetic conservatism of functional traits

in microorganisms. *The ISME Journal*, *7*(4), 830–838.

Matsen, F. A., Kodner, R. B., & Armbrust, E. V. (2010). pplacer: linear time maximum-likelihood

and Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*

Bioinformatics, *11*, 538.

Méheust, R., Burstein, D., Castelle, C. J., & Banfield, J. F. (2019). The distinction of CPR

bacteria from other bacteria based on protein family content. *Nature Communications*,

10(1), 4173.

Mendler, K., Chen, H., Parks, D. H., Lobb, B., Hug, L. A., & Doxey, A. C. (2019). AnnoTree:

visualization and exploration of a functionally annotated microbial tree of life. *Nucleic*

Acids Research, *47*(9), 4442–4448.

Miklós, I., & Podani, J. (2004). RANDOMIZATION OF PRESENCE–ABSENCE MATRICES: COMMENTS AND NEW ALGORITHMS. *Ecology*, Vol. 85, pp. 86–92. doi:10.1890/03-0101

Mitchell, A. L., Almeida, A., Beracochea, M., Boland, M., Burgin, J., Cochrane, G., … Finn, R. D. (2020). MGnify: the microbiome analysis resource in 2020. *Nucleic Acids Research*, *48*(D1), D570–D578.

Muller, E. E. L. (2019). Determining Microbial Niche Breadth in the Environment for Better Ecosystem Fate Predictions. *MSystems*, *4*(3). doi:10.1128/mSystems.00080-19

Murat Eren, A., Esen, Ö. C., Quince, C., Vineis, J. H., Morrison, H. G., Sogin, M. L., & Delmont, T. O. (2015). Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ*, *3*, e1319.

NCBI Resource Coordinators. (2018). Database resources of the National Center for Biotechnology Information. *Nucleic Acids Research*, *46*(D1), D8–D13.

Needham, D. M., Yoshizawa, S., Hosaka, T., Poirier, C., Choi, C. J., Hehenberger, E., … Worden, A. Z. (2019). A distinct lineage of giant viruses brings a rhodopsin photosystem to unicellular marine predators. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(41), 20574–20583.

Olson, D. K., Yoshizawa, S., Boeuf, D., Iwasaki, W., & DeLong, E. F. (2018). Proteorhodopsin variability and distribution in the North Pacific Subtropical Gyre. *The ISME Journal*, *12*(4), 1047–1060.

Pachiadaki, M. G., Brown, J. M., Brown, J., Bezuidt, O., Berube, P. M., Biller, S. J., … Stepanauskas, R. (2019). Charting the Complexity of the Marine Microbiome through Single-Cell Genomics. *Cell*, *179*(7), 1623-1635.e11.

Parks, D. H., Chuvochina, M., Waite, D. W., Rinke, C., Skarshewski, A., Chaumeil, P.-A., & Hugenholtz, P. (2018). A standardized bacterial taxonomy based on genome phylogeny

substantially revises the tree of life. *Nature Biotechnology*, *36*(10), 996–1004.

Pasolli, E., Asnicar, F., Manara, S., Zolfo, M., Karcher, N., Armanini, F., … Segata, N. (2019).

Extensive Unexplored Human Microbiome Diversity Revealed by Over 150,000

Genomes from Metagenomes Spanning Age, Geography, and Lifestyle. *Cell*, *176*(3),

649-662.e20.

Perdigão, N., Rosa, A. C., & O'Donoghue, S. I. (2017). The Dark Proteome Database. *BioData*

Mining, *10*(1), 1–11.

Price, M. N., Wetmore, K. M., Waters, R. J., Callaghan, M., Ray, J., Liu, H., … Deutschbauer, A.

M. (2018). Mutant phenotypes for thousands of bacterial genes of unknown function.

Nature, *557*(7706), 503–509.

Quince, C., Walker, A. W., Simpson, J. T., Loman, N. J., & Segata, N. (2017). Shotgun

metagenomics, from sampling to analysis. *Nature Biotechnology*, *35*(9), 833–844.

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* , *26*(6), 841–842.

Remmert, M., Biegert, A., Hauser, A., & Söding, J. (2012). HHblits: Lightning-fast iterative

protein sequence searching by HMM-HMM alignment. *Nature Methods*, *9*(2), 173–175.

Rinke, C., Schwientek, P., Sczyrba, A., Ivanova, N. N., Anderson, I. J., Cheng, J.-F., … Woyke,

T. (2013). Insights into the phylogeny and coding potential of microbial dark matter.

Nature, *499*(7459), 431–437.

Rost, B. (1999). Twilight zone of protein sequence alignments. *Protein Engineering, Design &*

Selection: PEDS, *12*(2), 85–94.

Roumpeka, D. D., Wallace, R. J., Escalettes, F., Fotheringham, I., & Watson, M. (2017). A

Review of Bioinformatics Tools for Bio-Prospecting from Metagenomic Sequence Data.

Frontiers in Genetics, *8*, 23.

Rusch, D. B., Halpern, A. L., Sutton, G., Heidelberg, K. B., Williamson, S., Yooseph, S., …

Venter, J. C. (2007). The Sorcerer II Global Ocean Sampling Expedition: Northwest

Atlantic through Eastern Tropical Pacific. *PLoS Biology*, *5*(3), 1–34.

Salazar, G., Cornejo-Castillo, F. M., Borrull, E., Díez-Vives, C., Lara, E., Vaqué, D., … Acinas,

S. G. (2015). Particle-association lifestyle is a phylogenetically conserved trait in

bathypelagic prokaryotes. *Molecular Ecology*, *24*(22), 5692–5706.

Salazar, G., Paoli, L., Alberti, A., Huerta-Cepas, J., Ruscheweyh, H.-J., Cuenca, M., …

Sunagawa, S. (2019). Gene Expression Changes and Community Turnover Differentially

Shape the Global Ocean Metatranscriptome. *Cell*, *179*(5), 1068-1083.e21.

Sanger, F., Nicklen, S., & Coulson, A. R. (1977). DNA sequencing with chain-terminating

inhibitors. *Proceedings of the National Academy of Sciences of the United States of*

America, *74*(12), 5463–5467.

Sberro, H., Fremin, B. J., Zlitni, S., Edfors, F., Greenfield, N., Snyder, M. P., … Bhatt, A. S.

(2019). Large-Scale Analyses of Human Microbiomes Reveal Thousands of Small,

Novel Genes. *Cell*, *178*(5), 1245-1259.e14.

Scales, B. S., Dickson, R. P., LiPuma, J. J., & Huffnagle, G. B. (2014). Microbiology, genomics,

and clinical significance of the Pseudomonas fluorescens species complex, an

unappreciated colonizer of humans. *Clinical Microbiology Reviews*, *27*(4), 927–948.

Skewes-Cox, P., Sharpton, T. J., Pollard, K. S., & DeRisi, J. L. (2014). Profile hidden Markov

models for the detection of viruses within metagenomic sequence data. *PloS One*, *9*(8),

e105067.

Spang, A., Saw, J. H., Jørgensen, S. L., Zaremba-Niedzwiedzka, K., Martijn, J., Lind, A. E., …

Ettema, T. J. G. (2015). Complex archaea that bridge the gap between prokaryotes and

eukaryotes. *Nature*, *521*(7551), 173–179.

Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. *Bioinformatics* , *30*(9), 1312–1313.

Steinegger, M., Meier, M., Mirdita, M., Vöhringer, H., Haunsberger, S. J., & Söding, J. (2019).

HH-suite3 for fast remote homology detection and deep protein annotation. *BMC*

Bioinformatics, *20*(1), 473.

Steinegger, M., Mirdita, M., & Söding, J. (2019). Protein-level assembly increases protein

sequence recovery from metagenomic samples manyfold. *Nature Methods*, *16*(7), 603–

606.

Steinegger, M., & Salzberg, S. L. (2020). Terminating contamination: large-scale search

identifies more than 2,000,000 contaminated entries in GenBank. *Genome Biology*,

21(1), 115.

Steinegger, M., & Soding, J. (2017). MMseqs2 enables sensitive protein sequence searching for

the analysis of massive data sets. *Nature Biotechnology*, *advance on*.

doi:10.1038/nbt.3988

Steinegger, M., & Söding, J. (2018). Clustering huge protein sequence sets in linear time.

Nature Communications, *9*(1), 2542.

Sunagawa, S., Coelho, L. P., Chaffron, S., Kultima, J. R., Labadie, K., Salazar, G., … Bork, P.

(2015). Ocean plankton. Structure and function of the global ocean microbiome.

Science, *348*(6237), 1261359.

The UniProt Consortium. (2017). UniProt: the universal protein knowledgebase. *Nucleic Acids*

Research, *45*(D1), D158–D169.

Thomas, A. M., & Segata, N. (2019). Multiple levels of the unknown in microbiome research.

BMC Biology, *17*(1), 48.

Titus Brown, C., Moritz, D., O'Brien, M. P., Reidl, F., Reiter, T., & Sullivan, B. D. (2018).

Exploring neighborhoods in large metagenome assembly graphs reveals hidden

sequence diversity (p. 462788). doi:10.1101/462788

UniProt Consortium, T. (2018). UniProt: the universal protein knowledgebase. *Nucleic Acids*

Research, *46*(5), 2699.

van Dongen, S., & Abreu-Goodger, C. (2012). Using MCL to Extract Clusters from Networks. In

J. van Helden, A. Toussaint, & D. Thieffry (Eds.), *Bacterial Molecular Networks: Methods*

and Protocols (pp. 281–295). New York, NY: Springer New York.

Vanhoutreve, R., Kress, A., Legrand, B., Gass, H., Poch, O., & Thompson, J. D. (2016). LEON-

BIS: multiple alignment evaluation of sequence neighbours using a Bayesian inference

system. *BMC Bioinformatics*, *17*(1), 271–271.

Vorobev, A., Dupouy, M., Carradec, Q., Delmont, T., Annamale, A., Wincker, P., & Pelletier, E.

(2020). Transcriptome reconstruction and functional analysis of eukaryotic marine

plankton communities via high-throughput metagenomics and metatranscriptomics.

Genome Research. doi:10.1101/gr.253070.119

Wyman, S. K., Avila-Herrera, A., Nayfach, S., & Pollard, K. S. (2018). A most wanted list of

conserved microbial protein families with no known domains. *PloS One*, *13*(10),

e0205749.

Yooseph, S., Li, W., & Sutton, G. (2008). Gene identification and protein classification in

microbial metagenomic sequence data via incremental clustering. *BMC Bioinformatics*,

9, 1–13.

Yooseph, S., Sutton, G., Rusch, B. D., Halpern, L. A., Williamson, J. S., Remington, K., …

Venter, C. J. (2007). The Sorcerer II global ocean sampling expedition: Expanding the

universe of protein families. *PLoS Biology*, *5*(3), 0432–0466.

Žure, M., Fernandez-Guerra, A., Munn, C. B., & Harder, J. (2017). Geographic distribution at

subspecies resolution level: closely related Rhodopirellula species in European coastal

sediments. *The ISME Journal*, *11*(2), 478–489.