Unifying the known and unknown microbial coding sequence space

Chiara Vanni[1,2], Matthew S. Schechter[1,3], Silvia G. Acinas[4], Albert Barberán[5], Pier Luigi Buttigieg[6], Emilio O. Casamayor[7], Tom O. Delmont[8], Carlos M. Duarte[9], A. Murat Eren[3,10], Robert D. Finn[11], Renzo Kottmann[1], Alex Mitchell[11], Pablo Sanchez[4], Kimmo Siren[12], Martin Steinegger[13,14], Frank Oliver Glöckner[15,16,2], Antonio Fernandez-Guerra[1,17] *

Affiliations

Microbial Genomics and Bioinformatics Research Group, Max Planck Institute for Marine Microbiology, Celsiusstraße 1, 28359, Bremen, Germany

Jacobs University Bremen, Campus Ring 1, 28759 Bremen, Germany

Department of Medicine, University of Chicago, Chicago, IL 60637, USA

Department of Marine Biology and Oceanography, Institut de Ciènces del Mar, CSIC, Barcelona, Spain.

Department of Environmental Science, University of Arizona, Tucson, 85721 AZ, USA

Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Am Handelshafen 12, 27570 Bremerhaven, Germany

Center for Advanced Studies of Blanes CEAB-CSIC, Spanish Council for Research, Blanes, Spain

8 Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Univ Evry,

Université Paris-Saclay, 91057 Evry, France

9 Red Sea Research Centre (RSRC) and Computational Bioscience Research Center (CBRC),

King Abdullah University of Science and Technology, Thuwal 23955, Saudi Arabia

10 Josephine Bay Paul Center, Marine Biological Laboratory, Woods Hole, MA 02543, USA

11 European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI),

Wellcome Genome Campus, Hinxton, Cambridge CB10 1SD, UK

12 Section for Evolutionary Genomics, The GLOBE Institute, University of Copenhagen,

Copenhagen, Denmark

13 School of Biological Sciences, Seoul National University, Seoul, 08826, South Korea

14 Institute of Molecular Biology and Genetics, Seoul National University, Seoul, 08826, South

Korea

15 University of Bremen, MARUM, Leobener Str. 8, 28359 Bremen, Germany

Life Sciences and Chemistry, Campus Ring 1, 28759 Bremen, Germany

16 Computing Center, Helmholtz Center for Polar and Marine Research, Am Handelshafen 12,

27570 Bremerhaven, Germany

17 Lundbeck GeoGenetics Centre, The Globe Institute, University of Copenhagen, 1350

Copenhagen, Denmark

*Corresponding author: Antonio Fernandez-Guerra, antonio.fernandez-guerra@sund.ku.dk

Abstract

Genes of unknown function are among the biggest challenges in molecular biology, especially in microbial systems, where 40%-60% of the predicted genes are unknown. Despite previous attempts, systematic approaches to include the unknown fraction into analytical workflows are still lacking. Here, we propose a conceptual framework and a computational workflow that bridge the known-unknown gap in genomes and metagenomes. We showcase our approach by exploring 415,971,742 genes predicted from 1,749 metagenomes and 28,941 bacterial and archaeal genomes. We quantify the extent of the unknown fraction, its diversity, and its relevance across multiple biomes. Furthermore, we provide a collection of 283,874 lineage-specific genes of unknown function for *Cand*. Patescibacteria, being a significant resource to expand our understanding of their unusual biology. Finally, by identifying a target gene of unknown function for antibiotic resistance, we demonstrate how we can enable the generation of hypotheses that can be used to augment experimental data.

Introduction

Thousands of isolate, single-cell, and metagenome-assembled genomes are guiding us towards a better understanding of life on Earth (Almeida et al., 2019; Cross et al., 2019; Delmont et al., 2020; Hug et al., 2016; Kopf et al., 2015; Pachiadaki et al., 2019; Pasolli et al., 2019; Sunagawa et al., 2015). At the same time, the ever-increasing number of genomes and metagenomes, unlocking uncharted regions of life's diversity, (Brown et al., 2015; Eloe-Fadrosh et al., 2016; Hug et al., 2016) are providing new perspectives on the evolution of life (Parks et al., 2018; Spang et al., 2015). However, our rapidly growing inventories of new genes have a glaring issue: between 40% and 60% cannot be assigned to a known function (Almeida et al., 2020; Bernard et al., 2018; Carradec et al., 2018; Price et al., 2018). Current analytical approaches for genomic and metagenomic data (Chen et al., 2019; Franzosa et al., 2018; Huerta-Cepas et al., 2017; Mitchell et al., 2020; Quince et al., 2017) generally do not include this uncharacterized fraction in downstream analyses, constraining their results to conserved pathways and housekeeping functions (Quince et al., 2017). This inability to handle the unknown is an immense impediment to realizing the potential for discovery of microbiology and molecular biology at large (Bernard et al., 2018; Hanson et al., 2010). Predicting function from traditional single sequence similarity appears to have yielded all it can (Arnold, 2018, 1998; Brandenberg et al., 2017), thus several groups have attempted to resolve gene function by other means. Such efforts include combining biochemistry and crystallography (Jaroszewski et al., 2009); using environmental co-occurrence (Buttigieg et al., 2013); by grouping those genes into evolutionarily related families (Bateman et al., 2010; Brum et al., 2016; Wyman et al., 2018; Yooseph et al., 2007); using remote homologies (Bitard-Feildel and Callebaut, 2017; Lobb et al., 2015); or more recently using deep learning approaches (Bileschi et al., 2019; Liu, 2017). In 2018, Price et al. (Price et al., 2018) developed a high-throughput experimental pipeline that provides mutant phenotypes for thousands of bacterial genes of unknown function being one of

the most promising methods to tackle the unknown. Despite their promise, experimental

methods are labor-intensive and require novel computational methods that could bridge the

existing gap between the genes with known and unknown function.

Here we present a conceptual framework and a computational workflow that closes the gap by

connecting genomic and metagenomic data by the exploitation of groups of homologous genes,

and facilitates the integration of genes of unknown function into ecological, evolutionary, and

biotechnological investigations. The conceptual framework is based on the partitioning of the

known and unknown fractions into four different categories that reflects the level of darkness

(Figure 1A). The category "Known" (K) contains these sequences predicted to harbor domains

of known function described by Pfam (Finn et al., 2016). Some sequences without Pfam

domains of unknown function, like the ones encoding for small and intrinsically disordered

proteins, can have homology to characterized proteins, hence we classify them as "Known

without Pfam" (KWP). Finally, sequences that cannot be classified by either of these criteria, are

classified as "Genomic Unknown" (GU) when they may be found in the genomes in reference

databases; and "Environmental Unknown" (EU) when only have been observed in

environmental samples. By contextualizing the different categories with information from several

sources (Figure 1C), we provide an invaluable resource for the investigation of the genes with

unknown function and boost the current methods for its experimental characterization.

The application of our approach to 415,971,742 genes predicted from 1,749 metagenomes and

28,941 bacterial and archaeal genomes revealed that the unknown fraction (1) is smaller than

has been previously reported, (2) is exceptionally diverse, and (3) is phylogenetically more

conserved than the known fraction and predominantly lineage-specific at the Species level.

Finally, we show how we can connect all the outputs produced by our approach to augment the

results from experimental data and add context to genes of unknown function through

hypothesis-driven molecular investigations.

Results

A computational workflow to unify the known and the unknown coding sequence space

Driven by the concepts defined in the conceptual framework, we developed AGNOSTOS, a computational workflow that infers, validates, refines, and classifies groups of homologous genes or gene clusters (GCs) in the four proposed categories (Fig. 1A; Fig. 1B; Supp. Fig 1). AGNOSTOS produces GCs with a highly conserved intra-homogeneous structure (Fig. 1B), both in terms of sequence similarity and domain architecture homogeneity; it exhausts any existing homology to known genes and provides a proper delimitation of the unknown genes before classifying each GC in one of the four categories (Methods). In the last step, we decorate each GC with a rich collection of contextual data compiled from different sources or generated by analyzing the GC contents in different contexts (Fig. 1A). For each GC, we also offer several products that can be used for analytical purposes like improved representative sequences, consensus sequences, sequence profiles for MMseqs2 (Steinegger and Soding, 2017) and HHblits (Steinegger et al., 2019a), or the GC members as a sequence similarity network (Methods). To complement the collection, we also provide a subset of what we define as *high-quality* GCs. The defining criteria are (1) the representative is a complete gene and (2) more than one-third of genes within a GC are complete genes.

Partitioning and contextualizing the coding sequence space of genomes and metagenomes

First, we used AGNOSTOS in a metagenomic context to show its ability to process complex and noisy datasets. We explored the unknown sequence space of 1,749 human and marine metagenomes. In total, we predicted 322,248,552 genes from the environmental dataset and assigned Pfam annotations to 44% of them (Supp. Fig. 2A). Next, AGNOSTOS clustered the predicted genes in 32,465,074 GCs and flagged those gene clusters that contain less than ten genes (Skewes-Cox et al., 2014). Flagged gene clusters are not processed through the validation workflow because they don't have enough sequences to provide reliable results. We flagged 29,461,177 gene clusters, 19,911,324 of them being singletons (Supp. Fig. 2A; Supp. Note 1). A total of 3,003,897 GCs (83% of the original genes) will go through the validation steps. The validation process selected 2,940,257 *good-quality* clusters (Fig. 1B; Supp. Table 1; Supp. Note 2), which resulted in 43% of them being members of the unknowns after the classification and remote homology refinement steps (Supp. Fig. 2A, Supp. Note 3).

Lastly, we demonstrate how AGNOSTOS can integrate a new dataset into the already existing

metagenomic database by integrating 28,941 genomes from the GTDB_r86 (Supp. Fig. 2A).

With this integration we can build links between the environmental and genomic sequence

space by expanding the final collection of GCs with the genes predicted from GTDB_r86.

Surprisingly, the integration showed that the environmental GCs (human and marine) not only

included a large proportion of the genes predicted from GTDB_r86 (72%) but also that most of

these genes (90%) are part of the known sequence space (Supp. Fig. 2A; Supp. Note 4; Supp.

Note 5). Only 22% of the GTDB_r86 genes created new GCs (2,400,037), being most of these

genes (84%) classified as unknowns by AGNOSTOS (Supp. Fig. 2A; Supp. Note 4; Supp. Note

5). Finally, only a small proportion of the genes from GTDB_r86 (6%) resulted in singletons

(Supp. Fig. 2A; Supp. Note 4; Supp. Note 5).

The final dataset after being validated and categorized resulted in 5,287,759 GCs (Supp. Fig.

2A), with both datasets sharing only 922,599 GCs (Supp. Fig. 2B). The integration of GTDB_r86

into the metagenomic database increased the proportion of GCs in the unknown sequence

space from 43% to the 54%.

Additionally, AGNOSTOS identified a subset of 203,217 *high-quality* GCs (Supp. Table 2). In

these *high-quality* GCs, we identified 12,313 clusters potentially encoding for small proteins (<=

50 amino acids). Most of these GCs are unknown (66%), which agrees with recent findings on

novel small proteins from metagenomes (Sberro et al., 2019). We also observed that the KWP

category contains the largest proportion of incomplete genes (Supp. Table 3), disrupting the

detection and assignment of Pfam domains. But it also incorporates sequences with an unusual

amino acid composition that have homology to proteins with high levels of disorder in the DPD

database (Perdigão et al., 2017) and has characteristic functions of intrinsically disordered

proteins (Habchi et al., 2014) (IDP) like cellular processes and signaling as predicted by

eggNOG annotations (Supp. Table 4).

As part of the workflow, each GC is complemented with a rich set of information, as shown in

Fig 1A (Supp. Table 5; Supp. Note 6).

Beyond the twilight zone, communities of gene clusters

To find relationships between gene clusters, we grouped them in gene cluster communities

(GCCs) (Fig. 2A) using remote homologies. As we are dealing with the unknown, we identified

GCCs independently in each category, using the gene clusters from category K as a reference

to identify the best parameters for the clustering of the gene cluster homology network

(Methods; Fig. 2B; Supp. Note 7) and applying the learnt parameters to the other categories.

The grouping reduced the final collection of GCs by 87%, producing 673,601 GCCs (Methods;

Fig. 2B; Supp. Note 7).

We validated our approach to capture remote homologies between related GCs using two well-

known gene families present in our environmental datasets, proteorhodopsins (Olson et al.,

2018) and bacterial ribosomal proteins (Méheust et al., 2019). Theoretically, we would expect to

have one or a very low number of GCCs for each of these gene families.

We identified 64 GCs (12,184 genes) and 3 GCCs (Supp. Note 8) containing genes classified

as proteorhodopsin (PR). One community from the category K contained 99% of the PR

annotated genes (Fig. 2C), except 85 genes taxonomically annotated as viral and assigned to

the *PR Supercluster I* (Boeuf et al., 2015) within two GU communities (five GU gene clusters;

Supp. Note 8).

For the ribosomal proteins, the results were not so satisfactory. We identified 1,843 GCs

(781,579 genes) and 98 GCCs. The number of GCCs is larger than the expected number of

ribosomal protein families (16) used for validation. When we used *high-quality* GCs (Supp. Note

8), we got closer to the expected number of GCCs (Fig. 2D). With this subset, we identified 26

GCCs and 145 GCs (1,687 genes). The cross-validation of our method against the approach

used in Méheust et al. (Méheust et al., 2019) (Supp. Note 9) confirms the intrinsic complexity of

analyzing metagenomic data. Both approaches showed a high agreement in the GCCs

identified (Supp. Table 9-1). Still, our method inferred fewer GCCs for each of the ribosomal

protein families (Supp. Fig. 9-3), coping better with the complexities of a metagenomic setup,

such as incomplete genes (Supp. Table 6).

Figure 2: Overview and validation of the workflow to aggregate GCs in communities. (A) We inferred a
gene cluster homology network using the results of an all-vs-all HMM gene cluster comparison with
HHBLITS. The edges of the network are based on the HHblits-score/Aligned-columns. Communities are
identified by an iterative screening of different MCL inflation parameters and evaluated using five different
metrics that consider the inter- and intra-community properties. (B) Comparison of the number of GCs
and GCCs for each of the functional categories. (C) Validation of the GCCs inference based on the
environmental genes annotated as proteorhodopsins. Ribbons in the alluvial plot are genes, and each
stacked bar corresponds (from left to right) to the (1) gene taxonomic classification at the domain level,

A smaller but highly diverse unknown coding sequence space

By combining clustering and remote homology searches we reduced the extent of the unknown

sequence space compared to what is reported by the traditional genomic and metagenomic

analysis approaches where the unknown fraction can add up to 60% in marine metagenomes

(Salazar et al., 2019) or up to 40% in human metagenomes (Thomas and Segata, 2019) (Fig.

3A). Our workflow recruited as much as 71% of genes in human-related metagenomic samples

and 65% of the genes in marine metagenomes into the known sequence space. In both human

and marine microbiomes, the genomic unknown fraction showed a similar proportion of genes

(21%, Fig. 3A). The number of genes corresponding to EU gene clusters is higher in marine

metagenomes; 12% of the genes are part of this GC category. We obtained a comparable result

when we evaluated the genes from the GTDB_r86, 75% of bacterial and 64% of archaeal genes

were part of the known sequence space. Archaeal genomes contained more unknowns than

those from Bacteria, where 30% of the genes are classified as genomic unknowns in Archaea,

and only 20% in Bacteria (Fig. 3A; Supp. Table 7).

To further evaluate the differences between the known and unknown sequence space, we

calculated the accumulation rates of GCs and GCCs, using them as operational protein families

(Schloss and Handelsman, 2008). For the metagenomic dataset we used 1,264 metagenomes

(18,566,675 GCs and 282,580 GCCs) and for the genomic dataset 28,941 genomes (9,586,109

GCs and 496,930 GCCs). The rate of accumulation of unknown GCs was three times higher

than the known (2X for the genomic), and in both cases the curves were far from reaching a

plateau (Fig. 3B). This is not the case for the GCC accumulation curves (Supp. Fig. 4B), which

reached a plateau.

Figure 3: The extent of the known and unknown sequence space (A) Proportion of genes in the known
and unknown. (B) Accumulation curves for the known and unknown sequence space at the GC- level for
the metagenomic and genomic data. from TARA, MALASPINA, OSD2014 and HMP-I/II projects. (C)
Collector curves comparing the human and marine biomes. Colored lines represented the mean of 1000
permutations and shaded in grey the standard deviation. Non-abundant singleton clusters were excluded
from the accumulation curves calculation. (D) Amino acid distribution in the known and unknown
sequence space.
253

The accumulation rate is largely determined by the number of singletons, especially singletons

from EUs (Supp. note 10 and Supp. Fig. 5). While the accumulation rate of known GCs between

marine and human metagenomes is almost identical, there are striking differences for the

unknown GCs (Fig. 3C). These differences are maintained even when we remove the virus-

enriched samples from the marine metagenomes (Supp. Fig. 4A). Although the marine

metagenomes include a large variety of environments, from coastal to the deep sea, the known

space remains quite constrained.

Next, we wanted to know how much of the sequence space we integrated with AGNOSTOS

was found in other databases (Supp. Note 11). Despite only including marine and human

metagenomes in our database, we already cover, in average, 76% of the sequence space of

seven datasets spanning different databases and environments (Supp. Fig. 11-1). By screening

MGnify (Mitchell et al., 2020) (release 2018_09; 11 biomes; 843,535,6116 proteins) we

identified freshwater, soil and human non-digestive as the biomes less covered by our data

(Supp. Fig. 6). Two of the seven analyzed datasets are designed to study genes of unknown

function (Supp. Table 11-1). On (Wyman et al., 2018), where they defined Function Unknown

Families of homologous proteins (FunkFams), we identified 20% of their FunkFams to be

members of the known sequence space. On (Price et al., 2018), we classified as known, 44% of

the genes of unknown function used in their experimental conditions.

One indirect consequence of our approach is that we can provide a detailed view of the

sequence space at the amino acid level. We estimated the number of amino acids belonging to

the known or unknown sequence space, and how many of these amino acids are contained

within Pfam domain boundaries. From the 90,128,659,316 amino acids analyzed, most of the

amino acids in metagenomes (74%) and genomes (80%) are in the known sequence space

(Fig. 3D; Supp. Table 7) while only 22% in metagenomes and 15% in genomes are part of the

unknowns. In both cases, approximately 40% of the amino acids in the known sequence space

were part of a Pfam domain (Fig. 3D; Supp. Table 7). While this result is expected based on the

large number of genes present in the known space (Fig. 3A), what is surprising is the low

proportion of amino acids (2%) corresponding to DUF Pfam domains in genomes and

metagenomes (Fig. 3D). If we use as reference the proportions of amino acids observed in the

known sequence space, we can hypothesize that there are still many DUFs to be unearthed.

With AGNOSTOS and its throughout validation and characterization of the genomic and

environmental unknowns, we provide the basic building blocks (gene clusters' multiple

sequence alignments) to identify conserved regions that might become new potential DUFs.

287 The unknown sequence space has a limited ecological
288 distribution in human and marine environments

Although the role of the unknown fraction in the environment is still a mystery, the large number

of gene counts and abundance observed underlines its inherent ecological relevance (Fig. 4A).

In some metagenomes, the genomic unknown fraction can account for more than 40% of the

total gene abundance observed (Fig. 4A). The environmental unknown fraction is also relevant

in several samples, where singleton GCs are the majority (Fig. 4A). We identified two

metagenomes with an unusual composition in terms of environmental unknown singletons. The

marine metagenome corresponds to a sample from Lake Faro (OSD42), a meromictic saline

with a unique extreme environment where Archaea plays an important role (La Cono et al.,

2013). And the HMP metagenome (SRS143565) that corresponds to a human sample from the

right cubital fossa from a healthy female subject. To understand this unusual composition, we

should perform further analyses to discard potential technical artifacts like sample

contamination.

The ratio between the unknown and known GCs is useful to reveal which metagenomes are

enriched in GCs of unknown function (upper left quadrant in Fig. 4B-C) and it can be used as a

proxy to assess the sequence contained in a metagenome. In human metagenomes, this ratio

can distinguish between body sites, with the gastrointestinal tract, an ecologically complex

environment (Qin et al., 2010), significantly enriched with genomic unknowns. Furthermore, it is

not surprising that the human and marine metagenomes with the largest ratio of unknowns are

those samples enriched with viral sequences. Specifically, in the HMP metagenomes are those

samples identified to contain crAssphages (Dubinkina et al., 2016; Edwards et al., 2019) and

HPV viruses (Ma et al., 2014) (Supp. Table 8; Supp. Fig. 7). While, in marine metagenomes

(Fig. 4C) the highest ratio in genomic and environmental unknowns correspond to the ones

enriched with viruses and giant viruses.

Figure 4: Distribution of the unknown sequence space in the human and marine metagenomes (A) Ratio
between the proportion of the number of genes and their estimated abundances per cluster category and
biome. Columns represented in the facet depicts three cluster categories based on the size of the
clusters. (B) Relationship between the ratio of Genomic unknowns and Environmental unknowns in the
HMP-I/II metagenomes. Gastrointestinal tract metagenomes are enriched in Genomic unknown
sequences compared to the other body sites. (C) Relationship between the ratio of Genomic unknowns
and Environmental unknowns in the TARA Oceans metagenomes. Girus and virus enriched
metagenomes show a higher proportion of both unknown sequences (genomic and environmental) than
the Archaea|Bacteria enriched fractions. (D) Environmental distribution of GCs and GCCs based on
Levin's niche breadth index. We obtained the significance values after generating 100 null gene cluster
abundance matrices using the *quasiswap* algorithm.

We performed a large-scale analysis to investigate the occurrence patterns of the GCs in the

environment by analyzing their abundance and distribution breadth. The narrow distribution of

the unknown fraction (Fig. 4D) suggests that these GCs might provide a selective advantage

and be necessary to adapt to specific environmental conditions. But the pool of broadly

distributed environmental unknowns is the most exciting result. We identified traces of potential

ubiquitous organisms left uncharacterized by traditional approaches, as more than 80% of these

GCs cannot be associated with a metagenome-assembled genome (MAG) (Supp. Table 9,

Supp. Note 12).

The genomic unknown coding sequence space is lineage-specific

With the inclusion of the genomes from GTDB_r86, we have access to a phylogenomic

framework that can be used to assess how exclusive is a GC within a lineage (lineage-specifity

(Mendler et al., 2019)) and how conserved (phylogenetic conservation) is a GC across the

different clades in the GTDB_r86 phylogenomic tree (Martiny et al., 2013). We identified

781,814 lineage-specific GCs and 464,923 phylogenetically conserved (P < 0.05) GCs in

Bacteria (Supp. Table 10; Supp. Note 13 for Archaea). The number of lineage-specific GCs

increases with the Relative Evolutionary Distance (Parks et al., 2018) (Fig. 5A) and differences

between the known and the unknown fraction start to be evident at the Family level resulting in

4X more unknown lineage-specific GCs at the Species level. In general terms, the unknown

GCs are more phylogenetically conserved (GCs shared among members of deep clades) than

the known (Fig. 5B, p < 0.0001), revealing the importance of the genome's uncharacterized

fraction. However, the lineage-specific unknown GCs are less phylogenetically conserved (Fig.

5B) than the known, agreeing with the large number of lineage-specific GCs observed at Genus

and Species level (Fig. 5A).

One potential confounding factor that might contribute to inflate the number of lineage-specific

GCs in the unknown fraction, is the presence of prophages owing to their potential host

specificity (Ross et al., 2016). To discard the possibility that the lineage-specific GCs of

unknown function have a viral origin, we screened all GTDB_r86 genomes for prophages. We

only found 37,163 lineage-specific GCs in prophage genomic regions, being 86% of these GCs

of unknown function.

After unveiling the potential relevance of the GCs of unknown function in bacterial genomes, we

identified phyla in GTDB_r86 enriched with these types of clusters. A clear pattern emerged

when we partitioned the phyla based on the ratio of known to unknown GCs and vice versa (Fig.

5D), the phyla with a larger number of MAGs are enriched in GCs of unknown function (Fig.

5D). Phyla with a high proportion of non-classified GCs (those discarded during the validation

steps) contain a small number of genomes and are primarily composed of MAGs. These groups

of phyla highly enriched in unknowns and represented mainly by MAGs include newly described

phyla such as *Cand.* Riflebacteria and *Cand.* Patescibacteria (Anantharaman et al., 2018;

Brown et al., 2015; Rinke et al., 2013), both with the largest unknown to known ratio.

Figure 5: Phylogenomic exploration of the unknown sequence space. (A) Distribution of the lineage-specific GCs by taxonomic level. Lineage-specific unknown GCs are more abundant in the lower taxonomic levels (Genus, Species). (B) Phylogenetic conservation of the known and unknown sequence space in 27,372 bacterial genomes from GTDB_r86. We observe differences in the conservation between the known and the unknown sequence space for lineage- and non-lineage specific GCs (paired Wilcoxon rank-sum test; all p-values < 0.0001). (C) The majority of the lineage-specific clusters are part of the unknown sequence space, being a small proportion found in prophages present in the GTDB_r86 genomes. (D) Known and unknown sequence space of the 27,732 GTDB_r86 bacterial genomes grouped by bacterial phyla. Phyla are partitioned based on the ratio of known to unknown GCs and vice versa. Phyla enriched in MAGs have higher proportions in GCs of unknown function. Phyla with a high proportion of non-classified clusters (NC; discarded during the validation steps) tend to contain a small number of genomes. (E) The alluvial plot's left side shows the uncharacterized (OM-RGC v2 GC) and characterized (OM-RGC v2) fraction of the gene catalog. The functional annotation is based on the eggNOG annotations provided by Salazar et al.(Salazar et al., 2019). The right side of the alluvial plot shows the new organization of the OM-RGC v2 sequence space based on the approach described in this study. The treemap in the right links the metagenomic and genomic space adding context to the unknown fraction of the OM-RGC v2

One of the strengths of AGNOSTOS is the possibility of bridging genomic and metagenomic

data and simultaneously unifying the known and unknown sequence space by integrating the

new Ocean Microbial Reference Gene Catalog (Salazar et al., 2019) (OM-RGC v2) into our

database. We assigned 26,170,875 genes to known GCs, 11,422,975 to genomic unknowns,

8,661,221 to environmental unknown and 520,083 were discarded. From the 11,422,975 genes

classified as genomic unknowns, we could associate 3,261,741 to a GTDB_r86 genome and we

identified 113,175 as lineage-specific. The alluvial plot in Fig. 5E depicts the new organization of

the OM-RGC v2 after being integrated into our framework and how we can provide context to

the two original types of unknowns in the OM-RGC (those annotated as category S in eggNOG

(Huerta-Cepas et al., 2019) and those without known homologs in the eggNOG database

(Salazar et al., 2019)) that can lead to potential experimental targets at the organism level to

complement the metatranscriptomic approach proposed by Salazar et al. (Salazar et al., 2019).

A structured sequence space augments the interpretation of experimental data

We selected one of the experimental conditions tested in Price et al. (Price et al., 2018) to

demonstrate the potential of our approach to augment experimental data. We compared the

fitness values in plain rich medium with added Spectinomycin dihydrochloride pentahydrate to

the fitness in plain rich medium (LB) in *Pseudomonas fluorescens FW300-N2C3* (Fig. 6A). This

antibiotic inhibits protein synthesis and elongation by binding to the bacterial 30S ribosomal

subunit and interferes with the peptidyl tRNA translocation. We identified the gene with locus id

AO356_08590 that presents a strong phenotype (fitness = -3.1; t = -9.1) and has no known

function. This gene belongs to the genomic unknown GC GU_19737823. We can track this GC

into the environment and explore the occurrence in the different samples we have in our

database. As expected, the GC is mostly found in non-human metagenomes (Fig. 6B) as

Pseudomonas are common inhabitants of soil and water environments (Heffernan et al., 2009).

However, finding this GC also in human-related samples is very interesting due to the potential

association of *P. fluorescens* and human disease where Crohn's disease patients develop

serum antibodies to this microbe (Scales et al., 2014). We can add another layer of information

to the selected GC by looking at the associated remote homologs in the GCC GU_c_21103

(Fig. 6C). We identified all the genes in the GTDB_r86 genomes that belong to the GCC

GU_c_21103 (Supp. Table 11) and explored their genomic neighborhoods. All members from

GU_c_21103 are constrained to the class *Gammaproteobacteria*, and interestingly

GU_19737823 is mostly exclusive to the order *Pseudomonadales*. The gene order in the

different genomes analyzed is highly conserved, finding GU_19737823 after the *rpsF*::*rpsR*

operon and before *rplI*. *rpsF* and *rpsR* encode for 30S ribosomal proteins, the prime target of

spectinomycin. The combination of the experimental evidence and the associated data inferred

by our approach provides strong support to generate the hypothesis that the gene

AO356_08590 might be involved in the resistance to Spectinomycin.

Figure 6: Augmenting experimental data with GCs of unknown function. (A) We used the fitness values from the experiments from Price et al. (Price et al., 2018) to identify genes of unknown function that are important for fitness under certain experimental conditions. The selected gene belongs to the genomic unknown GC GU_19737823 and presents a strong phenotype (fitness = -3.1; t = -9.1) (B) Occurrence of GU_19737823 in the metagenomes used in this study. Darker bars depict the number of metagenomes where the GC is found. (C) GU_19737823 is a member of the GCC GU_c_21103. The network shows the relationships between the different GCs members of the gene cluster community GU_c_21103. The size of the node corresponds to the node degree of each GC. Edge thickness corresponds to the bitscore/column metric. Highlighted in red is GU_19737823. (D) We identified all the genes in the GTDB_r86 genomes that belong to the GCC GU_c_21103 and explored their genomic neighborhoods. GU_c_21103 members were constrained to the class *Gammaproteobacteria*, and GU_19737823 is mostly exclusive to the order *Pseudomonadales*. The gene order in the different genomes analyzed is highly conserved, finding GU_19737823 after the *rpsF::rpsR* operon and before *rplI. rpsF* and *rpsR* encode for the *30S ribosomal protein S6* and *30S ribosomal protein S18,* respectively. The GTDB_r86 subtree only shows RefSeq genomes. Branch colors correspond to the different GCs found in GU_c_21103. The bubble plot depicts the number of genomes with a gene that belongs to GU_c_21103.

Discussion

We present a new conceptual framework and computational workflow to unify the known and unknown sequence space. Using this framework, we performed an in-depth exploration of the microbial unknown sequence space, demonstrating that we can link the unknown fraction of metagenomic studies to specific genomes and provide a powerful tool for hypothesis generation. The framework introduces a subtle change of paradigm compared to traditional approaches where our objective is to provide the best representation of the unknown space. We gear all our efforts towards finding sequences without any evidence of known homologies by pushing the search space beyond the *twilight zone* of sequence similarity (Rost, 1999). With this objective in mind, we use gene clusters instead of genes as the fundamental unit to compartmentalize the sequence space owing to their unique properties (Fig. 1B). Gene clusters (1) provide a structured sequence space that helps to reduce its complexity, (2) are independent of the known and unknown fraction, (3) are conserved across environments and organisms, and (4) can be used to aggregate information from different sources (Fig. 1A). Moreover, GCs provide a good compromise in terms of resolution for analytical purposes, and owing to their unique properties, one can perform analyses at different scales. For fine-grained analyses, we can exploit the gene associations within each GC; and for coarse-grained analyses, we can create groups of GCs based on their shared homologies (Fig. 1B).

AGNOSTOS integrates transparently into the standard operating procedure for analyzing metagenomes (Quince et al., 2017) adopted by the microbiome community. It can briefly be summarized into (1) assembly, (2) gene prediction, (3) gene catalog inference, (4) binning, and (5) characterization. AGNOSTOS exploits recent computational developments (Steinegger and Söding, 2018; Steinegger and Soding, 2017) to maximize the information used when analyzing genomic and metagenomic data. In addition, we provide a mechanism to reconcile top-down

and bottom-up approaches, thanks to the well-structured sequence space proposed by our

framework. AGNOSTOS can create environmental- and organism-specific variations of a seed

database based on gene clusters. Then, it integrates the predicted genes from new genomes

and metagenomes and dynamically creates and classifies new GCs with those genes not

integrated during the initial step (Fig. 1B). Afterward, the potential functions of the known GCs

can be carefully characterized by incorporating them into the traditional standard operating

procedure described previously.

One of the most appealing characteristics of our approach is that the GCs provide unified

groups of homologous genes across environments and organisms indifferently if they belong to

the known or unknown sequence space, and we can contextualize the unknown fraction using

this genomic and environmental information. Our combination of partitioning and

contextualization features a smaller unknown sequence space than previously reported

(Steinegger et al., 2019b). On average, only 30% of the genes fall in the unknown fraction for

our genomic and metagenomic data. One hypothesis to reconcile this surprising finding is that

the methodologies to identify remotely homologous sequences in large datasets were

computationally prohibitive until recently. New methods (Steinegger et al., 2019a; Steinegger

and Soding, 2017), like the ones used in AGNOSTOS, enable large-scale remote homology

searches. Still, one must apply conservative measures to control the trade-off between

specificity and sensitivity to avoid overclassification.

We found that most of the sequence space at the gene and amino acid level is known, both in

genomes and metagenomes. However, the GC accumulation curves showed that the unknown

fraction is far more diverse than the known. When we combine the high diversity and its narrow

ecological distribution, we can unveil the magnitude of the untapped unknown functional fraction

and its potential importance for niche adaptation. In a genomic context and after ruling out the

effect of prophages, the unknown fraction is predominantly Species' lineage-specific and

phylogenetically more conserved than the known fraction, supporting the signal observed in the

environmental data emphasizing that we should not ignore the unknown fraction. It is worth

noting that the high diversity observed in the unknowns only represents the 20% of the amino

acids in the sequence space we analyzed, and only 10% of this unknown amino acid space is

part of a Pfam domain (DUF and others). This contrasts with the numbers observed in the

known sequence space, where Pfam domains include 50% of the amino acids. All this evidence

combined strengthens the hypothesis that the genes of unknown function, especially the

lineage-specific ones, might be associated with the mechanisms of microbial diversification and

niche adaptation due to the constant diversification of gene families and the survival of new

gene lineages (Francino, 2012; Muller, 2019).

Metagenome-assembled genomes are not only unveiling new regions of the microbial universe

(42% of the genomes in GTDB_r86), but they are also enriching the tree of life with genes of

unknown function. One excellent example is *Cand*. Patescibacteria, more commonly known as

Candidate Phyla Radiation (CPR), a phylum that has raised considerable interest due to its

unusual biology (Brown et al., 2015). We performed an in-depth exploration of this phylum, and

we provide a collection of 54,343 lineage-specific GCs of unknown function at different

taxonomic level resolutions (Supp. Table 12; Supp. Note 14), which will be a valuable resource

for the CPR advancement research efforts.

One of the ultimate goals of our approach is to provide a mechanism to unlock the large pool of

likely relevant data that remains untapped to analysis and discovery and boost insights from

model organism experiments. We demonstrated the value of our approach by identifying a

potential target gene of unknown function for antibiotic resistance. But severe challenges

remain, such as the dependence on the quality of the assemblies and their gene predictions, as

shown by the analysis of the ribosomal protein GCCs where many of the recovered genes are

incomplete. While sequence assembly has been an active area of research (Roumpeka et al.,

2017), this has not been the case for gene prediction methods (Roumpeka et al., 2017), which

are becoming outdated (Ivanova et al., 2014) and cannot cope with the current amount of data.

Alternatives like protein-level assembly (Steinegger et al., 2019b) combined with exploring the

assembly graphs' neighborhoods (Brown et al., 2020) become very attractive for our purposes.

In any case, we still face the challenge of discriminating between genuine and artifactual

singletons (Höps et al., 2018). There are currently no methods available to provide a plausible

solution and, at the same time, being scalable. We devise a potential solution in the recent

developments in unsupervised deep learning methods where they use large corpora of proteins

to define a language model *embedding* for protein sequences (Heinzinger et al., 2019). These

models could be applied to predict *embeddings* in singletons, which could be clustered or used

to determine their coding potential. Another issue is that we might be creating more GCs than

expected. We follow a conservative approach to avoid mixing multi-domain proteins in GCs

owing to the fragmented nature of the metagenome assemblies that could result in the split of a

GC. However, not only splitting can be a problem, but also lumping unrelated genes or GCs

owing to the use of remote homologies. Although the inference of GCCs is using very sensitive

methods to compare profile HMMs, low sequence diversity in GCs can limit its effectiveness.

Moreover, our approach is affected by the presence and propagation of contamination in

reference databases, a significant problem in 'omics (Breitwieser et al., 2019; Steinegger and

Salzberg, 2020). In our case, we only use Pfam (Finn et al., 2016) as a source for annotation

owing to its high-quality and manual curation process. The categorization process of our GCs

depends on the information from other databases, and to minimize the potential impact of

contamination, we apply methods that weight the annotations of the identified homologs to

discriminate if a GC belongs to the known or unknown sequence space.

The results presented here prove that the integration and the analysis of the unknown fraction

are possible. We are unveiling a brighter future, not only for microbiome analyses but also for

boosting eukaryotic-related studies, thanks to the increasing number of projects, including

metatranscriptomic data (Delmont et al., 2020; Vorobev et al., 2020). Furthermore, our work

lays the foundations for further developments of clear guidelines and protocols to define the

different levels of unknown (Thomas and Segata, 2019) and should encourage the scientific

community for a collaborative effort to tackle this challenge.

551 Material and methods

552 Genomic and metagenomic dataset

We used a set of 583 marine metagenomes from four of the major metagenomic surveys of the

ocean microbiome: Tara Oceans expedition (TARA) (Sunagawa et al., 2015), Malaspina

expedition (Duarte, 2015), Ocean Sampling Day (OSD) (Kopf et al., 2015), and Global Ocean

Sampling Expedition (GOS) (Rusch et al., 2007). We complemented this set with 1,246

metagenomes obtained from the Human Microbiome Project (HMP) phase I and II (Lloyd-Price

et al., 2017). We used the assemblies provided by TARA, Malaspina, OSD and HMP projects

and the long Sanger reads from GOS (Sanger et al., 1977). A total of 156M (156,422,969)

contigs and 12.8M long-reads were collected (Supp. Table 6).

For the genomic dataset, we used the 28,941 prokaryotic genomes (27,372 bacterial and 1,569

archaeal) from the Genome Taxonomy Database (Parks et al., 2018) (GTDB) Release 03-RS86

(19th August 2018).

Computational workflow development

We implemented a computation workflow based on Snakemake (Köster, 2018) for the easy

processing of large datasets in a reproducible manner. The workflow provides three different

strategies to analyze the data. The module *DB-creation* creates the gene cluster database,

validates and partitions the gene clusters (GCs) in the main functional categories. The module

DB-update allows the integration of new sequences (either at the contig or predicted gene level)

in the existing gene cluster database. In addition, the workflow has a *profile-search* function to

quickly screen samples using the gene cluster PSSM profiles in the database.

Metagenomic and genomic gene prediction

We used Prodigal (v2.6.3) (Hyatt et al., 2010) in metagenomic mode to predict the genes from

the metagenomic dataset. For the genomic dataset, we used the gene predictions provided by

Annotree (Mendler et al., 2019), since they were obtained, consistently, with Prodigal v2.6.3.

We identified potential spurious genes using the *AntiFam* database (Eberhardt et al., 2012).

Furthermore, we screened for '*shadow*' genes using the procedure described in Yooseph et al.

(Yooseph et al., 2008).

PFAM annotation

We annotated the predicted genes using the *hmmsearch* program from the *HMMER* package

(version: 3.1b2) (Finn et al., 2011) in combination with the Pfam database v31 (Finn et al.,

2016). We kept the matches exceeding the internal gathering threshold and presenting an

independent e-value < 1e-5 and coverage > 0.4. In addition, we considered multi-domain

annotations, and we removed overlapping annotations when the overlap is larger than 50%,

keeping the ones with the smaller e-value.

Determination of the gene clusters

We clustered the metagenomic predicted genes using the cascaded-clustering workflow of the

MMseqs2 software (Steinegger and Söding, 2018) ("*--cov-mode 0 -c 0.8 --min-seq-id 0.3"*). We

discarded from downstream analyses the singletons and clusters with a size below a threshold

identified after applying a modification of the broken-stick model (Macarthur, 1957). We

randomly split the number of gene clusters into p subsets, where p is defined by the proportion

of outlier genes per gene cluster. The subsets are then sorted by decreasing size. We iterated

over all subsets averaging the results over all iterations. The broken stick model generates the

outlier gene proportions, which would occur by chance alone, that is, the distribution of outlier

gene proportions if there were no structure in the data.

We integrated the genomic data into the metagenomic cluster database using the "DB-update"

module of the workflow. This module uses the *clusterupdate* module of MMseqs2 (Steinegger

and Soding, 2017), with the same parameters used for the metagenomic clustering.

Quality-screening of gene clusters

We examined the GCs to ensure their high intra-cluster homogeneity. We applied two

methodologies to validate their cluster sequence composition and functional annotation

homogeneity. We identified non-homologous sequences inside each cluster combining the

identification of a new cluster representative sequence via a sequence similarity network (SSN)

analysis, and the investigation of intra-cluster multiple sequence alignments (MSAs), given the

new representative. Initially, we generated an SSN for each cluster, using the semi-global

alignment methods implemented in *PARASAIL* (Daily, 2016) (version 2.1.5). We trimmed the

SSN using a filtering algorithm (Chafee et al., 2018; Žure et al., 2017) that removes edges while

maintaining the network structural integrity and obtaining the smallest connected graph formed

by a single component. Finally, the new cluster representative was identified as the most central

node of the trimmed SSN by the eigenvector centrality algorithm, as implemented in igraph

(Csardi and Nepusz, 2006). After this step, we built a multiple sequence alignment for each

cluster using *FAMSA* (Deorowicz et al., 2016) (version 1.1). Then, we screened each cluster-

MSA for non-homologous sequences to the new cluster representative. Owing to computational

limitations, we used two different approaches to evaluate the cluster-MSAs. We used *LEON-BIS*

(Vanhoutreve et al., 2016) for the clusters with a size ranging from 10 to 1,000 genes and OD-

SEQ (Jehl et al., 2015) for the clusters with more than 1,000 genes. In the end, we applied a

broken-stick model (Macarthur, 1957) to determine the threshold to discard a cluster.

The predicted genes can have multi-domain annotations in different orders, therefore to validate

the consistency of intra-cluster Pfam annotations, we applied a combination of w-shingling

(Broder, 1997) and Jaccard similarity. We used w-shingling (k-shingle = 2) to group consecutive

domain annotations as a single object. We measured the homogeneity of the *shingle sets* (sets

of domains) between genes using the Jaccard similarity and reported the median similarity

value for each cluster. Moreover, we took into consideration the Clan membership of the Pfam

domains and that a gene might contain N-, C- and M-terminal domains for the functional

homogeneity validation. We discarded clusters with a median similarity < 1.

After the validation, we refined the gene cluster database removing the clusters identified to be

discarded and the clusters containing ≥ 30% *shadow genes*. Lastly, we removed the single

shadow, spurious and non-homologous genes from the remaining clusters (Supp. Note 2).

Remote homology classification of gene clusters

To partition the validated GCs into the four main categories, we processed the set of GCs

containing Pfam annotated genes and the set of not annotated GCs separately. For the

annotated GCs, we inferred a consensus protein domain architecture (DA) (an ordered

combination of protein domains) for each annotated gene cluster. To identify each gene cluster

consensus DA, we created directed acyclic graphs connecting the Pfam domains based on their

topological order on the genes using *igraph* (Csardi and Nepusz, 2006). We collapsed the

repetitions of the same domain. Then we used the gene completeness as a positive-weighting

value for the selection of the cluster consensus DA. Within this step, we divided the GCs into

"Knowns" (Known) if annotated to at least one Pfam domains of known function (DKFs) and

"Genomic unknowns" (GU) if annotated entirely to Pfam domains of unknown function (DUFs).

We aligned the sequences of the non-annotated GCs with FAMSA (Deorowicz et al., 2016) and

obtained cluster consensus sequences with the *hhconsensus* program from *HH-SUITE*

(Steinegger et al., 2019a). We used the cluster consensus sequences to perform a nested

search against the UniRef90 database (release 2017_11) (The UniProt Consortium, 2017) and

NCBI *nr* database (release 2017_12) (NCBI Resource Coordinators, 2018) to retrieve non-Pfam

annotations with *MMSeqs2* (Steinegger and Soding, 2017) ("*-e 1e-05 --cov-mode 2 -c 0.6*"). We

kept the hits within 60% of the Log(best-e-value) and searched the annotations for any of the

terms commonly used to define proteins of unknown function (Supp. Table 12). We used a

quorum majority voting approach to decide if a gene cluster would be classified as *Genomic*

Unknown or *Known without Pfams* based on the annotations retrieved. We searched the

consensus sequences without any homologs in the UniRef90 database against NCBI *nr*. We

applied the same approach and criteria described for the first search. Ultimately, we classified

as *Environmental Unknown* those GCs whose consensus sequences did not align with any of

the NCBI *nr* entries.

In addition, we developed some conservative measures to control the trade-off between

specificity and sensitivity for the remote homology searches such as (1) a modification of the

algorithm described in Hingamp et al. (Hingamp et al., 2013) to get a confident group of

homologs to determine if a query protein is known or unknown by a quorum majority voting

approach (Supp. Note 3); (2) strict parameters in terms of iterations, bidirectional coverage and

probability thresholds for the HHblits alignments to minimize the inclusion of non-homologous

sequences; and (3) avoid providing annotations for our gene clusters, as we believe that

annotation should be a careful process done on a smaller scale and with experimental context.

Gene cluster remote homology refinement

We refined the *Environmental Unknown* GCs to ensure the lack of any characterization by

searching for remote homologies in the Uniclust database (release 30_2017_10) using the

HMM/HMM alignment method *HHblits* (Remmert et al., 2012). We created the HMM profiles

with the *hhmake* program from the *HH-SUITE* (Steinegger et al., 2019a). We only accepted

those hits with an *HHblits-probability* ≥ 90% and we re-classified them following the same

majority vote approach as previously described. The clusters with no hits remained as the

refined set of EUs. We applied a similar refinement approach to the KWP clusters to identify

GCs with remote homologies to Pfam protein domains. The KWP HMM profiles were searched

against the Pfam *HH-SUITE* database (version 31), using *HHblits*. We accepted hits with a

probability ≥ 90% and a target coverage > 60% and removed overlapping domains as described

earlier. We moved the KWP with remote homologies to known Pfams to the Known set, and

those showing remote homologies to Pfam DUFs to the GUs. The clusters with no hits remained

as the refined set of KWP.

Gene cluster characterization

We used the *MMseqs2 taxonomy* module (commit:

b43de8b7559a3b45c8e5e9e02cb3023dd339231a) in combination with the UniProtKB (release

of January 2018) (UniProt Consortium, 2018) to retrieve the taxonomic ids of all genes in a gene

cluster. The *taxonomy* module implements the 2bLCA (Hingamp et al., 2013) to compute the

lowest common ancestor of query sequence. We used the following parameters "*-e 1e-05 –cov-*

mode 0 -c 0.6" for the search. To retrieve the taxonomic lineages, we used the R package

CHNOSZ (Dick, 2008).

We used eggNOG-mapper (Huerta-Cepas et al., 2017) and the EggNog5 database (Huerta-

Cepas et al., 2019) to provide functional annotations for each gene in a gene cluster. We refined

the functional annotations by selecting the orthologous group within the lowest taxonomic level

predicted by EggNog-mapper.

We measured the intra-cluster taxonomic and functional admixture by applying the

entropy.empirical() function from the *entropy* R package (Hausser and Strimmer, 2008). This

function estimates the Shannon entropy based on the different taxonomic and functional

annotation frequencies. For each cluster, we also retrieved the cluster consensus taxonomic

and functional annotation using a quorum majority voting approach..

In addition to the taxonomic and functional annotations, we evaluated the clusters' level of

darkness and disorder using the Dark Proteome Database (DPD) (Perdigão et al., 2017) as

reference. We searched the cluster genes against the DPD, applying the MMseqs2 search

program (Steinegger and Soding, 2017) with "*-e 1e-20 --cov-mode 0 -c 0.6*". For each cluster,

we then retrieved the mean and the median level of darkness, based on the gene DPD

annotations.

699   High-quality clusters

700   We defined a subset of high-quality clusters based on the completeness of the cluster genes

701   and their representatives. We identified the minimum required percentage of complete genes

702   per cluster by a broken-stick model (Macarthur, 1957) applied to the percentage distribution.

703   Then, we selected the GCs found above the threshold and with a complete representative.

704   A set of non-redundant domain architectures

We estimated the number of potential domain architectures present in the *Known* GCs

considering the large proportion of fragmented genes in the metagenomic dataset and that

could inflate the number of potential domain architectures. To identify fragments of larger

domain architecture, we considered their topological order in the genes. To reduce the number

of comparisons, we calculated the pairwise string cosine distance (q-gram = 3) between domain

architectures and discarded the pairs that were too divergent (cosine distance ≥ 0.9). We

collapsed a fragmented domain architecture to the larger one when it contained less than 75%

of complete genes.

Inference of gene cluster communities

We aggregated distant homologous GCs into GCCs. The community inference approach

combined an all-vs-all HMM gene cluster comparison with Markov Cluster Algorithm (MCL) (van

Dongen and Abreu-Goodger, 2012) community identification. We started performing the

inference on the Known GCs to use the Pfam DAs as constraints. We aligned the gene cluster

HMMs using HHblits (Remmert et al., 2012) (-n 2 -Z 10000000 -B 10000000 -e 1) and we built a

homology graph using the cluster pairs with probability ≥ 50% and bidirectional coverage > 60%.

We used the ratio between HHblits-bitscore and aligned-columns as the edge weights (Supp.

Note 9). We used MCL (van Dongen and Abreu-Goodger, 2012) (v. 12-068) to identify the

communities present in the graph. We developed an iterative method to determine the optimal

MCL inflation parameter that tries to maximize the relationship of five intra-/inter-community

properties: (1) the proportion of MCL communities with one single DA, based on the consensus

DAs of the cluster members; (2) the ratio of MCL communities with more than one cluster; (3)

the proportion of MCL communities with a PFAM clan entropy equal to 0; (4) the intra-

community HHblits-score/Aligned-columns score (normalized by the maximum value); and (5)

the number of MCL communities, which should, in the end, reflect the number of non-redundant

DAs. We iterated through values ranging from 1.2 to 3.0, with incremental steps of 0.1. During

the inference process, some of the GCs became orphans in the graph. We applied a three-step

approach to assigning a community membership to these GCs. First, we used less stringent

conditions (probability ≥ 50% and coverage >= 40%) to find homologs in the already existing

GCCs. Then, we ran a second iteration to find secondary relationships between the newly

assigned GCs and the missing ones. Lastly, we created new communities with the remaining

GCs. We repeated the whole process with the other categories (KWP, GU and EU), applying

the optimal inflation value found for the Known (2.2 for metagenomic and 2.5 for genomic data).

Validation of gene cluster communities

We tested the biological significance of the GCCs using the phylogeny of proteorhodopsin

(Boeuf et al., 2015) (PR). We used the proteorhodopsin HMM profiles (Olson et al., 2018) to

screen the marine metagenomic datasets using *hmmsearch* (version 3.1b2) (Finn et al., 2011).

We kept the hits with a coverage > 0.4 and e-value <= 1e-5. We removed identical duplicates

from the sequences assigned to PR with CD-HIT (Li and Godzik, 2006) (v4.6) and cleaned from

sequences with less than 100 amino acids. To place the identified PR sequences into the

MicRhode (Boeuf et al., 2015) PR tree first, we optimized the initial tree parameters and branch

lengths with RAxML (v8.2.12) (Stamatakis, 2014). We used PaPaRA (v2.5) (Berger and

Stamatakis, 2012) to incrementally align the query PR sequences against the MicRhode PR

reference alignment and *pplacer* (Matsen et al., 2010) (v1.1.alpha19-0-g807f6f3) to place the

sequences into the tree. Finally, we assigned the query PR sequences to the MicRhode PR

Superclusters based on the phylogenetic placement. We further investigated the GCs annotated

as viral (196 genes, 14 GC) comparing them to the six newly discovered viral PRs (Needham et

al., 2019) using Parasail (Daily, 2016) (-a sg_stats_scan_sse2_128_16 -t 8 -c 1 -x). As an

additional evaluation, we investigated the distributions of standard GCCs and HQ GCCs within

ribosomal protein families. We obtained the ribosomal proteins used for the analysis combining

the set of 16 ribosomal proteins from Méheust et al. (Méheust et al., 2019) and those contained

in the collection of bacterial single-copy genes of anvi'o (Eren et al., 2021). Also, for the

ribosomal proteins, we compared the outcome of our method to the one proposed by Méheust

et al. (Méheust et al., 2019) (Supp. Note 9).

Metagenomic sample selection for downstream analyses

For the subsequent ecological analyses, we selected those metagenomes with a number of genes larger or equal to the first quartile of the distribution of all the metagenomic gene counts. (Supp. Table 13).

Gene cluster abundance profiles in genomes and metagenomes

We estimated abundance profiles for the metagenomic cluster categories using the read coverage to each predicted gene as a proxy for abundance. We calculated the coverage by mapping the reads against the assembly contigs using the *bwa-mem* algorithm from *BWA mapper* (Li and Durbin, 2010). Then, we used *BEDTOOLS* (Quinlan and Hall, 2010), to find the intersection of the gene coordinates to the assemblies, and normalize the per-base coverage by the length of the gene. We calculated the cluster abundance in a sample as the sum of the cluster gene abundances in that sample, and the cluster category abundance in a sample as the sum of the cluster abundances. We obtained the proportions of the different gene cluster categories applying a total-sum-scaling normalization. For the genomic abundance profiles, we used the number of genes in the genomes and normalized by the total gene counts per genome.

Rate of genomic and metagenomic gene clusters accumulation

We calculated the cumulative number of known and unknown GCs as a function of the number of metagenomes and genomes. For each metagenome count, we generated 1000 random sets, and we calculated the number of GCs and GCCs recovered. For this analysis, we used 1,246 HMP metagenomes and 358 marine metagenomes (242 from TARA and 116 from Malaspina). We repeated the same procedure for the genomic dataset. We removed the singletons from the metagenomic dataset with an abundance smaller than the mode abundance of the singletons

781 that got reclassified as good-quality clusters after integrating the GTDB data to minimize the

782 impact of potential spurious singletons. To complement those analyses, we evaluated the

783 coverage of our dataset by searching seven different state-of-the-art databases against our set

784 of metagenomic GC HMM profiles (Supp. Note 11).

786 Occurrence of gene clusters in the environment

We used 1,264 metagenomes from the TARA Oceans, MALASPINA Expedition, OSD2014 and

HMP-I/II to explore the properties of the unknown sequence space in the environment. We

applied the Levins Niche Breadth (NB) index (Levins, 1966) to investigate the GCs and GCCs

environmental distributions. We removed the GCs and cluster communities with a mean relative

abundance < 1e-5. We followed a divide-and-conquer strategy to avoid the computational

burden of generating the null-models to test the significance of the distributions owing to the

large number of metagenomes and GCs. First, we grouped similar samples based on the gene

cluster content using the Bray-Curtis dissimilarity (Bray et al., 1957) in combination with the

Dynamic Tree Cut (Langfelder et al., 2008) R package. We created 100 random datasets

picking up one random sample from each group. For each of the 100 random datasets, we

created 100 random abundance matrices using the *nullmodel* function of the *quasiswap* count

method (Miklós and Podani, 2004). Then we calculated the *observed* NB and obtained the 2.5%

and 97.5% quantiles based on the randomized sets. We compared the observed and quantile

values for each gene cluster and defined it to have a *Narrow distribution* when the *observed*

was smaller than the 2.5% quantile and to have a *Broad distribution* when it was larger than the

97.5% quantile. Otherwise, we classified the cluster as *Non-significant* (Salazar et al., 2015).

We used a majority voting approach to get a consensus distribution classification based on the

ten random datasets.

Identification of prophages in genomic sequences

We used PhageBoost (Sirén et al., 2021) to find gene regions in the microbial genomes that result in high viral signals against the overall genome signal. We set the following thresholds to consider a region prophage: minimum of 10 genes, maximum 5 gaps, single-gene probability threshold 0.9. We further smoothed the predictions using Parzen rolling windows of 20 periods and looked at the smoothed probability distribution across the genome. We disregarded regions that had a summed smoothed probability less than 0.5, and those regions that did differ from the overall population of the genes in a genome by using Kruskal–Wallis rank test (p-value 0.001).

Lineage-specific gene clusters

We used the F1-score developed for AnnoTree (Mendler et al., 2019) to identify the lineage-specific GCs and to which rank they are specific. Following similar criteria to the ones used in Mendler et al. (Mendler et al., 2019), we considered a gene cluster to be lineage-specific if it is present in less than half of all genomes and at least 2 with F1-score > 0.95.

Phylogenetic conservation of gene clusters

We calculated the phylogenetic conservation (τD) of each gene cluster using the *consenTRAIT* (Martiny et al., 2013) function implemented in the R package *castor* (Martiny et al., 2013). We used a paired Wilcoxon rank-sum test to compare the average τD values for lineage-specific and non-specific GCs.

Evaluation of the OM-RGC v2 uncharacterized fraction

We integrated the 46,775,154 genes from the second version of the TARA Ocean Microbial Reference Gene Catalog (OM-RGC v2) (Salazar et al., 2019) into our cluster database using the same procedure as for the genomic data. We evaluated the uncharacterized fraction and the

827   genes classified into the eggNOG (Huerta-Cepas et al., 2019) category S within the context of

828   our database.

Augmenting RB-TnSeq experimental data

We searched the 37,684 genes of unknown function associated with mutant phenotypes from

Price et al. (Price et al., 2018) against our gene cluster profiles. We kept the hits with e-value ≤

1e-20 and a query coverage > 60%. Then we filtered the results to keep the hits within 90% of

the Log(best-e-value), and we used a majority vote function to retrieve the consensus category

for each hit. Lastly, we selected the best-hits based on the smallest e-value and the largest

query and target coverage values. We used the fitness values from the RB-TnSeq experiments

from Price et al. to identify genes of unknown function that are important for fitness under

certain experimental conditions.

Availability of data and materials

The code used for the analyses in the manuscript is available at https://github.com/functional-

dark-side/functional-dark-side.github.io/tree/master/scripts. The code to recreate the figures is

available at https://github.com/functional-dark-side/vanni_et_al-figures. Detailed descriptions of

the different methods and results of this manuscript are available at

https://dark.metagenomics.eu. The workflow AGNOSTOS is available at

https://github.com/functional-dark-side/agnostos-wf (the version used for the manuscript can be

found here https://doi.org/10.5281/zenodo.4557847) and its database can be downloaded from

https://doi.org/10.6084/m9.figshare.12459056.

Acknowledgements

The authors thankfully acknowledge the computer resources at MareNostrum and the technical support provided by Barcelona Supercomputing Center (RES-AECT-2014-2-0085), the BMBF-funded de.NBI Cloud within the German Network for Bioinformatics Infrastructure (de.NBI) (031A537B, 031A533A, 031A538A, 031A533B, 031A535A, 031A537C, 031A534A, 031A532B), the University of Oxford Advanced Research Computing (http://dx.doi.org/10.5281/zenodo.22558) and the MARBITS bioinformatics core at ICM-CSIC. CV was supported by the Max Planck Society. AFG received funding from the European Union's Horizon 2020 research and innovation program Blue Growth: Unlocking the potential of Seas and Oceans under grant agreement no. 634486 (project acronym INMARE). AM was supported by the Biotechnology and Biological Sciences Research Council [BB/M011755/1, BB/R015228/1] and RDF by the European Molecular Biology Laboratory core funds. EOC was supported by project INTERACTOMA RTI2018-101205-B-I00 from the Spanish Agency of Science MICIU/AEI. SGA and PS received additional funding by the project MAGGY (CTM2017-87736-R) from the Spanish Ministry of Economy and Competitiveness. The Malaspina 2010 Expedition was supported by the Spanish Ministry of Economy and Competitiveness (MINECO) through the Consolider-Ingenio program (ref. CSD2008-00077). The authors thank Johannes Söding and Alex Bateman for helpful discussions.

Contributions

CV, MSS and AF-G performed the analyses and wrote the computational workflow. MS assisted with the clustering and remote homology searches. KS helped with the identification of prophages in genomic sequences. PLB and AB provided feedback and assisted with the ecological analyses. RDF and AM provided feedback and information on the MGnify and Pfam

873   databases. CMD, PS and SGA provided the Malaspina metagenomes. TOD and AME analyzed

874   data in the context of metagenome-assembled genomes. AF-G conceived the study and

875   supervised the work. CV and AF-G wrote the manuscript. All authors read, edited and approved

876   the final manuscript.

877 Competing Interests

878   The authors declare no competing interests.

References

Almeida A, Mitchell AL, Boland M, Forster SC, Gloor GB, Tarkowska A, Lawley TD, Finn RD. 2019. A new genomic blueprint of the human gut microbiota. *Nature* **568**:499–504.

Almeida A, Nayfach S, Boland M, Strozzi F, Beracochea M, Shi ZJ, Pollard KS, Sakharova E, Parks DH, Hugenholtz P, Segata N, Kyrpides NC, Finn RD. 2020. A unified catalog of 204,938 reference genomes from the human gut microbiome. *Nat Biotechnol*. doi:10.1038/s41587-020-0603-3

Anantharaman K, Hausmann B, Jungbluth SP, Kantor RS, Lavy A, Warren LA, Rappé MS, Pester M, Loy A, Thomas BC, Banfield JF. 2018. Expanded diversity of microbial groups that shape the dissimilatory sulfur cycle. *ISME J* **12**:1715–1728.

Arnold FH. 2018. Directed Evolution: Bringing New Chemistry to Life. *Angew Chem Int Ed Engl* **57**:4143–4148.

Arnold FH. 1998. Design by Directed Evolution. *Acc Chem Res* **31**:125–131.

Bateman A, Coggill P, Finn DR. 2010. DUFs: Families in search of function. *Acta Crystallogr Sect F Struct Biol Cryst Commun* **66**:1148–1152.

Berger SA, Stamatakis A. 2012. PaPaRa 2.0: a vectorized algorithm for probabilistic phylogeny-aware alignment extension. *Heidelberg Institute for Theoretical Studies, http://sco h-its org/exelixis/publica tions html Exelixis-RRDR-2012-2015*.

Bernard G, Pathmanathan JS, Lannes R, Lopez P, Bapteste E. 2018. Microbial Dark Matter Investigations: How Microbial Studies Transform Biological Knowledge and Empirically Sketch a Logic of Scientific Discovery. *Genome Biol Evol* **10**:707–715.

Bileschi ML, Belanger D, Bryant D, Sanderson T, Carter B, Sculley D, DePristo MA, Colwell LJ. 2019. Using Deep Learning to Annotate the Protein Universe. *bioRxiv*. doi:10.1101/626507

Bitard-Feildel T, Callebaut I. 2017. Exploring the dark foldable proteome by considering hydrophobic amino acids topology. *Sci Rep* **7**:41425.

Boeuf D, Audic S, Brillet-Guéguen L, Caron C, Jeanthon C. 2015. MicRhoDE: a curated database for the analysis of microbial rhodopsin diversity and evolution. *Database* **2015**. doi:10.1093/database/bav080

Brandenberg OF, Fasan R, Arnold FH. 2017. Exploiting and engineering hemoproteins for abiological carbene and nitrene transfer reactions. *Curr Opin Biotechnol* **47**:102–111.

Bray JR, Roger Bray J, Curtis JT. 1957. An Ordination of the Upland Forest Communities of Southern Wisconsin. *Ecological Monographs*. doi:10.2307/1942268

Breitwieser FP, Pertea M, Zimin A, Salzberg SL. 2019. Human contamination in bacterial genomes has created thousands of spurious proteins. *Genome Res*. doi:10.1101/gr.245373.118

Broder AZ. 1997. On the resemblance and containment of documentsProceedings. Compression and Complexity of SEQUENCES 1997 (Cat. No. 97TB100171). IEEE. pp. 21–29.

Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A, Wilkins MJ, Wrighton KC, Williams KH, Banfield JF. 2015. Unusual biology across a group comprising more than 15% of domain Bacteria. *Nature* **523**:208–211.

Brown CT, Moritz D, O'Brien MP, Reidl F, Reiter T, Sullivan BD. 2020. Exploring neighborhoods in large metagenome assembly graphs using spacegraphcats reveals hidden sequence diversity. *Genome Biol* **21**:164.

Brum JR, Cesar Ignacio-Espinoza J, Kim E-H, Trubl G, Jones RM, Roux S, VerBerkmoes NC, Rich VI, Sullivan MB. 2016. Illuminating structural proteins in viral "dark matter" with metaproteomics. *Proc Natl Acad Sci U S A* **113**:2436–2441.

Buttigieg LP, Hankeln W, Kostadinov I, Kottmann R, Yilmaz P, Duhaime BM, Gl??ckner OF.

2013. Ecogenomic Perspectives on Domains of Unknown Function: Correlation-Based
Exploration of Marine Metagenomes. *PLoS One* **8**.

Carradec Q, Pelletier E, Da Silva C, Alberti A, Seeleuthner Y, Blanc-Mathieu R, Lima-Mendez
G, Rocha F, Tirichine L, Labadie K, Kirilovsky A, Bertrand A, Engelen S, Madoui M-A,
Méheust R, Poulain J, Romac S, Richter DJ, Yoshikawa G, Dimier C, Kandels-Lewis S,
Picheral M, Searson S, Tara Oceans Coordinators, Jaillon O, Aury J-M, Karsenti E,
Sullivan MB, Sunagawa S, Bork P, Not F, Hingamp P, Raes J, Guidi L, Ogata H, de
Vargas C, Iudicone D, Bowler C, Wincker P. 2018. A global ocean atlas of eukaryotic
genes. *Nat Commun* **9**:373.

Chafee M, Fernàndez-Guerra A, Buttigieg PL, Gerdts G, Eren AM, Teeling H, Amann RI. 2018.
Recurrent patterns of microdiversity in a temperate coastal marine environment. *ISME J*
12:237–252.

Chen I-MA, Chu K, Palaniappan K, Pillay M, Ratner A, Huang J, Huntemann M, Varghese N,
White JR, Seshadri R, Smirnova T, Kirton E, Jungbluth SP, Woyke T, Eloe-Fadrosh EA,
Ivanova NN, Kyrpides NC. 2019. IMG/M v.5.0: an integrated data management and
comparative analysis system for microbial genomes and microbiomes. *Nucleic Acids*
Res **47**:D666–D677.

Cross KL, Campbell JH, Balachandran M, Campbell AG, Cooper SJ, Griffen A, Heaton M, Joshi
S, Klingeman D, Leys E, Yang Z, Parks JM, Podar M. 2019. Targeted isolation and
cultivation of uncultivated bacteria by reverse genomics. *Nat Biotechnol* **37**:1314–1321.

Csardi G, Nepusz T. 2006. The igraph software package for complex network research.
InterJournal.

Daily J. 2016. Parasail: SIMD C library for global, semi-global, and local pairwise sequence
alignments. *BMC Bioinformatics* **17**:81–81.

Delmont TO, Gaia M, Hinsinger DD, Fremont P, Guerra AF, Murat Eren A, Vanni C, Kourlaiev
A, d'Agata L, Clayssen Q, Villar E, Labadie K, Cruaud C, Poulain J, Da Silva C, Wessner
M, Noel B, Aury J-M, Coordinators TO, de Vargas C, Bowler C, Karsenti E, Pelletier E,
Wincker P, Jaillon O. 2020. Functional repertoire convergence of distantly related
eukaryotic plankton lineages revealed by genome-resolved metagenomics. *Cold Spring*
Harbor Laboratory. doi:10.1101/2020.10.15.341214

Deorowicz S, Debudaj-Grabysz A, Gudyś A. 2016. FAMSA: Fast and accurate multiple
sequence alignment of huge protein families. *Sci Rep* **6**:33964–33964.

Dick JM. 2008. Calculation of the relative metastabilities of proteins using the CHNOSZ
software package. *Geochem Trans* **9**:10.

Duarte CM. 2015. Seafaring in the 21St Century: The Malaspina 2010 Circumnavigation
Expedition. *Limnol Oceanog Bull* **24**:11–14.

Dubinkina VB, Ischenko DS, Ulyantsev VI, Tyakht AV, Alexeev DG. 2016. Assessment of k-mer
spectrum applicability for metagenomic dissimilarity analysis. *BMC Bioinformatics*.
doi:10.1186/s12859-015-0875-7

Eberhardt RY, Haft DH, Punta M, Martin M, O'Donovan C, Bateman A. 2012. AntiFam: a tool to
help identify spurious ORFs in protein annotation. *Database* **2012**:bas003–bas003.

Edwards RA, Vega AA, Norman HM, Ohaeri M, Levi K, Dinsdale EA, Cinek O, Aziz RK, McNair
K, Barr JJ, Bibby K, Brouns SJJ, Cazares A, de Jonge PA, Desnues C, Díaz Muñoz SL,
Fineran PC, Kurilshikov A, Lavigne R, Mazankova K, McCarthy DT, Nobrega FL, Reyes
Muñoz A, Tapia G, Trefault N, Tyakht AV, Vinuesa P, Wagemans J, Zhernakova A,
Aarestrup FM, Ahmadov G, Alassaf A, Anton J, Asangba A, Billings EK, Cantu VA,
Carlton JM, Cazares D, Cho G-S, Condeff T, Cortés P, Cranfield M, Cuevas DA, De la
Iglesia R, Decewicz P, Doane MP, Dominy NJ, Dziewit L, Elwasila BM, Eren AM, Franz
C, Fu J, Garcia-Aljaro C, Ghedin E, Gulino KM, Haggerty JM, Head SR, Hendriksen RS,
Hill C, Hyöty H, Ilina EN, Irwin MT, Jeffries TC, Jofre J, Junge RE, Kelley ST, Khan
Mirzaei M, Kowalewski M, Kumaresan D, Leigh SR, Lipson D, Lisitsyna ES, Llagostera

M, Maritz JM, Marr LC, McCann A, Molshanski-Mor S, Monteiro S, Moreira-Grez B,
Morris M, Mugisha L, Muniesa M, Neve H, Nguyen N-P, Nigro OD, Nilsson AS,
O'Connell T, Odeh R, Oliver A, Piuri M, Prussin AJ Ii, Qimron U, Quan Z-X, Rainetova P,
Ramírez-Rojas A, Raya R, Reasor K, Rice GAO, Rossi A, Santos R, Shimashita J,
Stachler EN, Stene LC, Strain R, Stumpf R, Torres PJ, Twaddle A, Ugochi Ibekwe M,
Villagra N, Wandro S, White B, Whiteley A, Whiteson KL, Wijmenga C, Zambrano MM,
Zschach H, Dutilh BE. 2019. Global phylogeography and ancient evolution of the
widespread human gut virus crAssphage. *Nat Microbiol* **4**:1727–1736.
Eloe-Fadrosh EA, Paez-Espino D, Jarett J, Dunfield PF, Hedlund BP, Dekas AE, Grasby SE,
Brady AL, Dong H, Briggs BR, Li W-J, Goudeau D, Malmstrom R, Pati A, Pett-Ridge J,
Rubin EM, Woyke T, Kyrpides NC, Ivanova NN. 2016. Global metagenomic survey
reveals a new bacterial candidate phylum in geothermal springs. *Nat Commun* **7**:10476.
Eren AM, Kiefl E, Shaiber A, Veseli I, Miller SE, Schechter MS, Fink I, Pan JN, Yousef M,
Fogarty EC, Trigodet F, Watson AR, Esen ÖC, Moore RM, Clayssen Q, Lee MD,
Kivenson V, Graham ED, Merrill BD, Karkman A, Blankenberg D, Eppley JM, Sjödin A,
Scott JJ, Vázquez-Campos X, McKay LJ, McDaniel EA, Stevens SLR, Anderson RE,
Fuessel J, Fernandez-Guerra A, Maignien L, Delmont TO, Willis AD. 2021. Community-
led, integrated, reproducible multi-omics with anvi'o. *Nat Microbiol* **6**:3–6.
Finn RD, Clements J, Eddy SR. 2011. HMMER web server: interactive sequence similarity
searching. *Nucleic Acids Res* **39**:W29–W37.
Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M, Qureshi
M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A. 2016. The Pfam protein
families database: towards a more sustainable future. *Nucleic Acids Res* **44**:D279–
D285.
Francino MP. 2012. The ecology of bacterial genes and the survival of the new. *Int J Evol Biol*
2012:394026.
Franzosa EA, McIver LJ, Rahnavard G, Thompson LR, Schirmer M, Weingart G, Lipson KS,
Knight R, Caporaso JG, Segata N, Huttenhower C. 2018. Species-level functional
profiling of metagenomes and metatranscriptomes. *Nat Methods* **15**:962–968.
Habchi J, Tompa P, Longhi S, Uversky VN. 2014. Introducing protein intrinsic disorder. *Chem*
Rev **114**:6561–6588.
Hanson AD, Pribat A, Waller JC, Crécy-Lagard V de. 2010. 'Unknown' proteins and
'orphan' enzymes: the missing half of the engineering parts list--and how to find it.
Biochem J **425**:1–11.
Hausser J, Strimmer K. 2008. Entropy inference and the James-Stein estimator, with application
to nonlinear gene association networks. *arXiv [statML]*.
Heffernan B, Murphy CD, Casey E. 2009. Comparison of planktonic and biofilm cultures of
Pseudomonas fluorescens DSM 8341 cells grown on fluoroacetate. *Appl Environ*
Microbiol **75**:2899–2907.
Heinzinger M, Elnaggar A, Wang Y, Dallago C, Nechaev D, Matthes F, Rost B. 2019. Modeling
aspects of the language of life through transfer-learning protein sequences. *BMC*
Bioinformatics **20**:723.
Hingamp P, Grimsley N, Acinas SG, Clerissi C, Subirana L, Poulain J, Ferrera I, Sarmento H,
Villar E, Lima-Mendez G, Faust K, Sunagawa S, Claverie J-M, Moreau H, Desdevises Y,
Bork P, Raes J, de Vargas C, Karsenti E, Kandels-Lewis S, Jaillon O, Not F, Pesant S,
Wincker P, Ogata H. 2013. Exploring nucleo-cytoplasmic large DNA viruses in Tara
Oceans microbial metagenomes. *ISME J* **7**:1678–1695.
Höps W, Jeffryes M, Bateman A. 2018. Gene Unprediction with Spurio: A tool to identify
spurious protein sequences. *F1000Res* **7**:261.
Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, von Mering C, Bork P. 2017.
Fast Genome-Wide Functional Annotation through Orthology Assignment by eggNOG-

Mapper. *Mol Biol Evol* **34**:2115–2122.

Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR, Letunic I, Rattei T, Jensen LJ, von Mering C, Bork P. 2019. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res* **47**:D309–D314.

Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN, Hernsdorf AW, Amano Y, Ise K, Suzuki Y, Dudek N, Relman DA, Finstad KM, Amundson R, Thomas BC, Banfield JF. 2016. A new view of the tree of life. *Nat Microbiol* **1**:16048.

Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**:119–119.

Ivanova NN, Schwientek P, Tripp HJ, Rinke C, Pati A, Huntemann M, Visel A, Woyke T, Kyrpides NC, Rubin EM. 2014. Stop codon reassignments in the wild. *Science* **344**:909–913.

Jaroszewski L, Li Z, Krishna SS, Bakolitsa C, Wooley J, Deacon MA, Wilson AI, Godzik A. 2009. Exploration of uncharted regions of the protein universe. *PLoS Biol* **7**.

Jehl P, Sievers F, Higgins DG. 2015. OD-seq: outlier detection in multiple sequence alignments. *BMC Bioinformatics* **16**:269–269.

Kopf A, Bicak M, Kottmann R, Schnetzer J, Kostadinov I, Lehmann K, Fernandez-Guerra A, Jeanthon C, Rahav E, Ullrich M, Wichels A, Gerdts G, Polymenakou P, Kotoulas G, Siam R, Abdallah RZ, Sonnenschein EC, Cariou T, O'Gara F, Jackson S, Orlic S, Steinke M, Busch J, Duarte B, Caçador I, Canning-Clode J, Bobrova O, Marteinsson V, Reynisson E, Loureiro CM, Luna GM, Quero GM, Löscher CR, Kremp A, DeLorenzo ME, Øvreås L, Tolman J, LaRoche J, Penna A, Frischer M, Davis T, Katherine B, Meyer CP, Ramos S, Magalhães C, Jude-Lemeilleur F, Aguirre-Macedo ML, Wang S, Poulton N, Jones S, Collin R, Fuhrman JA, Conan P, Alonso C, Stambler N, Goodwin K, Yakimov MM, Baltar F, Bodrossy L, Van De Kamp J, Frampton DM, Ostrowski M, Van Ruth P, Malthouse P, Claus S, Deneudt K, Mortelmans J, Pitois S, Wallom D, Salter I, Costa R, Schroeder DC, Kandil MM, Amaral V, Biancalana F, Santana R, Pedrotti ML, Yoshida T, Ogata H, Ingleton T, Munnik K, Rodriguez-Ezpeleta N, Berteaux-Lecellier V, Wecker P, Cancio I, Vaulot D, Bienhold C, Ghazal H, Chaouni B, Essayeh S, Ettamimi S, Zaid EH, Boukhatem N, Bouali A, Chahboune R, Barrijal S, Timinouni M, El Otmani F, Bennani M, Mea M, Todorova N, Karamfilov V, Ten Hoopen P, Cochrane G, L'Haridon S, Bizsel KC, Vezzi A, Lauro FM, Martin P, Jensen RM, Hinks J, Gebbels S, Rosselli R, De Pascale F, Schiavon R, Dos Santos A, Villar E, Pesant S, Cataletto B, Malfatti F, Edirisinghe R, Silveira JAH, Barbier M, Turk V, Tinta T, Fuller WJ, Salihoglu I, Serakinci N, Ergoren MC, Bresnan E, Iriberri J, Nyhus PAF, Bente E, Karlsen HE, Golyshin PN, Gasol JM, Moncheva S, Dzhembekova N, Johnson Z, Sinigalliano CD, Gidley ML, Zingone A, Danovaro R, Tsiamis G, Clark MS, Costa AC, El Bour M, Martins AM, Collins RE, Ducluzeau A-L, Martinez J, Costello MJ, Amaral-Zettler LA, Gilbert JA, Davies N, Field D, Glöckner FO. 2015. The ocean sampling day consortium. *Gigascience* **4**:27.

Köster J. 2018. Reproducible data analysis with Snakemake. *F1000Res* **7**.

La Cono V, La Spada G, Arcadi E, Placenti F, Smedile F, Ruggeri G, Michaud L, Raffa C, De Domenico E, Sprovieri M, Mazzola S, Genovese L, Giuliano L, Slepak VZ, Yakimov MM. 2013. Partaking of Archaea to biogeochemical cycling in oxygen-deficient zones of meromictic saline Lake Faro (Messina, Italy). *Environ Microbiol* **15**:1717–1733.

Langfelder P, Zhang B, Horvath S. 2008. Defining clusters from a hierarchical cluster tree: the Dynamic Tree Cut package for R. *Bioinformatics* **24**:719–720.

Levins R. 1966. THE STRATEGY OF MODEL BUILDING IN POPULATION BIOLOGY. *Am Sci* **54**:421–431.

Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows–Wheeler transform.
Bioinformatics **26**:589–595.
Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of protein
or nucleotide sequences. *Bioinformatics* **22**:1658–1659.
Liu XL. 2017. Deep Recurrent Neural Network for Protein Function Prediction from Sequence.
bioRxiv. doi:10.1101/103994
Lloyd-Price J, Mahurkar A, Rahnavard G, Crabtree J, Orvis J, Hall AB, Brady A, Creasy HH,
McCracken C, Giglio MG, McDonald D, Franzosa EA, Knight R, White O, Huttenhower
C. 2017. Strains, functions and dynamics in the expanded Human Microbiome Project.
Nature **550**:61–66.
Lobb B, Kurtz DA, Moreno-Hagelsieb G, Doxey AC. 2015. Remote homology and the functions
of metagenomic dark matter. *Front Genet* **6**:1–12.
Ma Y, Madupu R, Karaoz U, Nossa CW, Yang L, Yooseph S, Yachimski PS, Brodie EL, Nelson
KE, Pei Z. 2014. Human papillomavirus community in healthy persons, defined by
metagenomics analysis of human microbiome project shotgun sequencing data sets. *J*
Virol **88**:4786–4797.
Macarthur RH. 1957. On the relative abundance of bird species. *Proc Natl Acad Sci U S A*
43:293–295.
Martiny AC, Treseder K, Pusch G. 2013. Phylogenetic conservatism of functional traits in
microorganisms. *ISME J* **7**:830–838.
Matsen FA, Kodner RB, Armbrust EV. 2010. pplacer: linear time maximum-likelihood and
Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*
Bioinformatics **11**:538.
Méheust R, Burstein D, Castelle CJ, Banfield JF. 2019. The distinction of CPR bacteria from
other bacteria based on protein family content. *Nat Commun* **10**:4173.
Mendler K, Chen H, Parks DH, Lobb B, Hug LA, Doxey AC. 2019. AnnoTree: visualization and
exploration of a functionally annotated microbial tree of life. *Nucleic Acids Res* **47**:4442–
4448.
Miklós I, Podani J. 2004. RANDOMIZATION OF PRESENCE–ABSENCE MATRICES:
COMMENTS AND NEW ALGORITHMS. *Ecology*. doi:10.1890/03-0101
Mitchell AL, Almeida A, Beracochea M, Boland M, Burgin J, Cochrane G, Crusoe MR, Kale V,
Potter SC, Richardson LJ, Sakharova E, Scheremetjew M, Korobeynikov A, Shlemov A,
Kunyavskaya O, Lapidus A, Finn RD. 2020. MGnify: the microbiome analysis resource in
2020. *Nucleic Acids Res* **48**:D570–D578.
Muller EEL. 2019. Determining Microbial Niche Breadth in the Environment for Better
Ecosystem Fate Predictions. *mSystems* **4**. doi:10.1128/mSystems.00080-19
NCBI Resource Coordinators. 2018. Database resources of the National Center for
Biotechnology Information. *Nucleic Acids Res* **46**:D8–D13.
Needham DM, Yoshizawa S, Hosaka T, Poirier C, Choi CJ, Hehenberger E, Irwin NAT, Wilken
S, Yung C-M, Bachy C, Kurihara R, Nakajima Y, Kojima K, Kimura-Someya T, Leonard
G, Malmstrom RR, Mende DR, Olson DK, Sudo Y, Sudek S, Richards TA, DeLong EF,
Keeling PJ, Santoro AE, Shirouzu M, Iwasaki W, Worden AZ. 2019. A distinct lineage of
giant viruses brings a rhodopsin photosystem to unicellular marine predators. *Proc Natl*
Acad Sci U S A **116**:20574–20583.
Olson DK, Yoshizawa S, Boeuf D, Iwasaki W, DeLong EF. 2018. Proteorhodopsin variability
and distribution in the North Pacific Subtropical Gyre. *ISME J* **12**:1047–1060.
Pachiadaki MG, Brown JM, Brown J, Bezuidt O, Berube PM, Biller SJ, Poulton NJ, Burkart MD,
La Clair JJ, Chisholm SW, Stepanauskas R. 2019. Charting the Complexity of the
Marine Microbiome through Single-Cell Genomics. *Cell* **179**:1623-1635.e11.
Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, Hugenholtz P.
2018. A standardized bacterial taxonomy based on genome phylogeny substantially

revises the tree of life. *Nat Biotechnol* **36**:996–1004.

Pasolli E, Asnicar F, Manara S, Zolfo M, Karcher N, Armanini F, Beghini F, Manghi P, Tett A,
Ghensi P, Collado MC, Rice BL, DuLong C, Morgan XC, Golden CD, Quince C,
Huttenhower C, Segata N. 2019. Extensive Unexplored Human Microbiome Diversity
Revealed by Over 150,000 Genomes from Metagenomes Spanning Age, Geography,
and Lifestyle. *Cell* **176**:649-662.e20.

Perdigão N, Rosa AC, O'Donoghue SI. 2017. The Dark Proteome Database. *BioData Min* **10**:1–
11.

Price MN, Wetmore KM, Waters RJ, Callaghan M, Ray J, Liu H, Kuehl JV, Melnyk RA, Lamson
JS, Suh Y, Carlson HK, Esquivel Z, Sadeeshkumar H, Chakraborty R, Zane GM, Rubin
BE, Wall JD, Visel A, Bristow J, Blow MJ, Arkin AP, Deutschbauer AM. 2018. Mutant
phenotypes for thousands of bacterial genes of unknown function. *Nature* **557**:503–509.

Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, Nielsen T, Pons N, Levenez F,
Yamada T, Mende DR, Li J, Xu J, Li Shaochuan, Li D, Cao J, Wang B, Liang H, Zheng
H, Xie Y, Tap J, Lepage P, Bertalan M, Batto J-M, Hansen T, Le Paslier D, Linneberg A,
Nielsen HB, Pelletier E, Renault P, Sicheritz-Ponten T, Turner K, Zhu H, Yu C, Li
Shengting, Jian M, Zhou Y, Li Y, Zhang X, Li Songgang, Qin N, Yang H, Wang Jian,
Brunak S, Doré J, Guarner F, Kristiansen K, Pedersen O, Parkhill J, Weissenbach J,
MetaHIT Consortium, Bork P, Ehrlich SD, Wang Jun. 2010. A human gut microbial gene
catalogue established by metagenomic sequencing. *Nature* **464**:59–65.

Quince C, Walker AW, Simpson JT, Loman NJ, Segata N. 2017. Shotgun metagenomics, from
sampling to analysis. *Nat Biotechnol* **35**:833–844.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**:841–842.

Remmert M, Biegert A, Hauser A, Söding J. 2012. HHblits: Lightning-fast iterative protein
sequence searching by HMM-HMM alignment. *Nat Methods* **9**:173–175.

Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Anderson IJ, Cheng J-F, Darling A, Malfatti S,
Swan BK, Gies EA, Dodsworth JA, Hedlund BP, Tsiamis G, Sievert SM, Liu W-T, Eisen
JA, Hallam SJ, Kyrpides NC, Stepanauskas R, Rubin EM, Hugenholtz P, Woyke T.
2013. Insights into the phylogeny and coding potential of microbial dark matter. *Nature*
499:431–437.

Ross A, Ward S, Hyman P. 2016. More Is Better: Selecting for Broad Host Range
Bacteriophages. *Front Microbiol* **7**:1352.

Rost B. 1999. Twilight zone of protein sequence alignments. *Protein Eng Des Sel* **12**:85–94.

Roumpeka DD, Wallace RJ, Escalettes F, Fotheringham I, Watson M. 2017. A Review of
Bioinformatics Tools for Bio-Prospecting from Metagenomic Sequence Data. *Front*
Genet **8**:23.

Rusch DB, Halpern AL, Sutton G, Heidelberg KB, Williamson S, Yooseph S, Wu D, Eisen JA,
Hoffman JM, Remington K, Beeson K, Tran B, Smith H, Baden-Tillson H, Stewart C,
Thorpe J, Freeman J, Andrews-Pfannkoch C, Venter JE, Li K, Kravitz S, Heidelberg JF,
Utterback T, Rogers Y-H, Falc n LI, Souza V, Bonilla-Rosso G, Eguiarte LE, Karl DM,
Sathyendranath S, Platt T, Bermingham E, Gallardo V, Tamayo-Castillo G, Ferrari MR,
Strausberg RL, Nealson K, Friedman R, Frazier M, Venter JC. 2007. The Sorcerer II
Global Ocean Sampling Expedition: Northwest Atlantic through Eastern Tropical Pacific.
PLoS Biol **5**:1–34.

Salazar G, Cornejo-Castillo FM, Borrull E, Díez-Vives C, Lara E, Vaqué D, Arrieta JM, Duarte
CM, Gasol JM, Acinas SG. 2015. Particle-association lifestyle is a phylogenetically
conserved trait in bathypelagic prokaryotes. *Mol Ecol* **24**:5692–5706.

Salazar G, Paoli L, Alberti A, Huerta-Cepas J, Ruscheweyh H-J, Cuenca M, Field CM, Coelho
LP, Cruaud C, Engelen S, Gregory AC, Labadie K, Marec C, Pelletier E, Royo-Llonch M,
Roux S, Sánchez P, Uehara H, Zayed AA, Zeller G, Carmichael M, Dimier C, Ferland J,

Kandels S, Picheral M, Pisarev S, Poulain J, Tara Oceans Coordinators, Acinas SG,
Babin M, Bork P, Bowler C, de Vargas C, Guidi L, Hingamp P, Iudicone D, Karp-Boss L,
Karsenti E, Ogata H, Pesant S, Speich S, Sullivan MB, Wincker P, Sunagawa S. 2019.
Gene Expression Changes and Community Turnover Differentially Shape the Global
Ocean Metatranscriptome. *Cell* **179**:1068-1083.e21.

Sanger F, Nicklen S, Coulson AR. 1977. DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* **74**:5463–5467.

Sberro H, Fremin BJ, Zlitni S, Edfors F, Greenfield N, Snyder MP, Pavlopoulos GA, Kyrpides NC, Bhatt AS. 2019. Large-Scale Analyses of Human Microbiomes Reveal Thousands of Small, Novel Genes. *Cell* **178**:1245-1259.e14.

Scales BS, Dickson RP, LiPuma JJ, Huffnagle GB. 2014. Microbiology, genomics, and clinical significance of the Pseudomonas fluorescens species complex, an unappreciated colonizer of humans. *Clin Microbiol Rev* **27**:927–948.

Schloss PD, Handelsman J. 2008. A statistical toolbox for metagenomics: assessing functional diversity in microbial communities. *BMC Bioinformatics* **9**:34.

Sirén K, Millard A, Petersen B, Gilbert MTP, Clokie MRJ, Sicheritz-Pontén T. 2021. Rapid discovery of novel prophages using biological feature engineering and machine learning. *NAR Genom Bioinform* **3**. doi:10.1093/nargab/lqaa109

Skewes-Cox P, Sharpton TJ, Pollard KS, DeRisi JL. 2014. Profile hidden Markov models for the detection of viruses within metagenomic sequence data. *PLoS One* **9**:e105067.

Spang A, Saw JH, Jørgensen SL, Zaremba-Niedzwiedzka K, Martijn J, Lind AE, van Eijk R, Schleper C, Guy L, Ettema TJG. 2015. Complex archaea that bridge the gap between prokaryotes and eukaryotes. *Nature* **521**:173–179.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**:1312–1313.

Steinegger M, Meier M, Mirdita M, Vöhringer H, Haunsberger SJ, Söding J. 2019a. HH-suite3 for fast remote homology detection and deep protein annotation. *BMC Bioinformatics* **20**:473.

Steinegger M, Mirdita M, Söding J. 2019b. Protein-level assembly increases protein sequence recovery from metagenomic samples manyfold. *Nat Methods* **16**:603–606.

Steinegger M, Salzberg SL. 2020. Terminating contamination: large-scale search identifies more than 2,000,000 contaminated entries in GenBank. *Genome Biol* **21**:115.

Steinegger M, Söding J. 2018. Clustering huge protein sequence sets in linear time. *Nat Commun* **9**:2542.

Steinegger M, Soding J. 2017. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat Biotechnol* **advance on**. doi:10.1038/nbt.3988

Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G, Djahanschiri B, Zeller G, Mende DR, Alberti A, Cornejo-Castillo FM, Costea PI, Cruaud C, d'Ovidio F, Engelen S, Ferrera I, Gasol JM, Guidi L, Hildebrand F, Kokoszka F, Lepoivre C, Lima-Mendez G, Poulain J, Poulos BT, Royo-Llonch M, Sarmento H, Vieira-Silva S, Dimier C, Picheral M, Searson S, Kandels-Lewis S, Tara Oceans coordinators, Bowler C, de Vargas C, Gorsky G, Grimsley N, Hingamp P, Iudicone D, Jaillon O, Not F, Ogata H, Pesant S, Speich S, Stemmann L, Sullivan MB, Weissenbach J, Wincker P, Karsenti E, Raes J, Acinas SG, Bork P. 2015. Ocean plankton. Structure and function of the global ocean microbiome. *Science* **348**:1261359.

The UniProt Consortium. 2017. UniProt: the universal protein knowledgebase. *Nucleic Acids Res* **45**:D158–D169.

Thomas AM, Segata N. 2019. Multiple levels of the unknown in microbiome research. *BMC Biol* **17**:48.

UniProt Consortium T. 2018. UniProt: the universal protein knowledgebase. *Nucleic Acids Res* **46**:2699.

1234 van Dongen S, Abreu-Goodger C. 2012. Using MCL to Extract Clusters from Networks In: van
1235   Helden J, Toussaint A, Thieffry D, editors. Bacterial Molecular Networks: Methods and
1236   Protocols. New York, NY: Springer New York. pp. 281–295.
1237 Vanhoutreve R, Kress A, Legrand B, Gass H, Poch O, Thompson JD. 2016. LEON-BIS: multiple
1238   alignment evaluation of sequence neighbours using a Bayesian inference system. *BMC*
1239   *Bioinformatics* **17**:271–271.
1240 Vorobev A, Dupouy M, Carradec Q, Delmont T, Annamale A, Wincker P, Pelletier E. 2020.
1241   Transcriptome reconstruction and functional analysis of eukaryotic marine plankton
1242   communities via high-throughput metagenomics and metatranscriptomics. *Genome Res*.
1243   doi:10.1101/gr.253070.119
1244 Wyman SK, Avila-Herrera A, Nayfach S, Pollard KS. 2018. A most wanted list of conserved
1245   microbial protein families with no known domains. *PLoS One* **13**:e0205749.
1246 Yooseph S, Li W, Sutton G. 2008. Gene identification and protein classification in microbial
1247   metagenomic sequence data via incremental clustering. *BMC Bioinformatics* **9**:1–13.
1248 Yooseph S, Sutton G, Rusch BD, Halpern LA, Williamson JS, Remington K, Eisen AJ,
1249   Heidelberg BK, Manning G, Li W, Jaroszewski L, Cieplak P, Miller SC, Li H, Mashiyama
1250   TS, Joachimiak PM, Van Belle C, Chandonia MJ, Soergel AD, Zhai Y, Natarajan K, Lee
1251   S, Raphael JB, Bafna V, Friedman R, Brenner ES, Godzik A, Eisenberg D, Dixon EJ,
1252   Taylor SS, Strausberg LR, Frazier M, Venter CJ. 2007. The Sorcerer II global ocean
1253   sampling expedition: Expanding the universe of protein families. *PLoS Biol* **5**:0432–
1254   0466.
1255 Žure M, Fernandez-Guerra A, Munn CB, Harder J. 2017. Geographic distribution at subspecies
1256   resolution level: closely related Rhodopirellula species in European coastal sediments.
1257   *ISME J* **11**:478–489.