Dorsolateral striatal circuits support broadly opponent aspects of action suppression and production

Bruno F. Cruz, Sofia Soares, Joseph J. Paton*
Champalimaud Research, Champalimaud Centre for the Unknown, Lisbon, PT.
*to whom correspondence should be addressed. joe.paton@research.fchampalimaud.org

Abstract

Imbalance between action suppression and production characterizes several neurological disorders of the basal ganglia (BG)[1]. Relatedly, two major circuits of the BG, the direct and indirect pathways, are hypothesized to function in opposition, promoting and suppressing actions, respectively[2]. Yet activity of neurons initiating the two pathways, striatal direct (dMSNs) and indirect (iMSNs) medium spiny neurons, appears to be positively correlated around movement, apparently contradicting direct-indirect functional opponency[3]. Here we show that while coactivation occurs during movement, action suppression produces systematic features of functional opponency between the two pathways. First, suppression produced elevated iMSN activity and diminished dMSN activity in dorsolateral striatum of mice. Second, as the need to suppress movements to one or the other side of the body evolved over time, so too did the relative levels of activity in the two hemispheres, and in opposite patterns in the two pathways. Lastly, optogenetic inhibition of dMSNs, but not iMSNs, slowed movements without affecting their likelihood, whereas inhibition of iMSNs, but not dMSNs, disrupted whether and when mice suppressed or produced particular actions. These data demonstrate large-scale opponency in the activity of neurons that initiate the direct and indirect pathways of the BG, but suggest distinct action-related functions of each pathway. Specifically, the data support a novel model of sensorimotor striatum wherein the direct pathway appears to be necessary for augmenting generalized movement vigor, and the indirect pathway for the proactive suppression of specific behaviors.

Introduction

Adaptive behavior involves a judicious combination of suppression and production of actions. A predator must suppress its urge to pounce until its prey is within reach, just as humans must suppress giving in to temptation to secure longer-term rewards.

The basal ganglia (BG) are a collection of subcortical structures that are thought to regulate the appropriate selection of actions depending on expected consequences[4,5]. In addition, the inability to balance action production and suppression is associated with disorders that involve the BG such as ADHD[6], Parkinson's, and Huntington's diseases[1]. Interestingly, two major BG pathways, the so-called direct and indirect pathways, possess anatomical and molecular characteristics consistent with promoting and suppressing actions, respectively[7,8]. These two pathways originate in the major input area of the BG, the striatum, at direct striatonigral medium spiny neurons (dMSNs) and indirect striatopallidal medium spiny neurons (iMSNs) that project directly or indirectly toward the output areas of the BG. While multiple lines of evidence suggest functional opponency between the two pathways, an apparent discordance between neural activity on the one hand, and anatomical and cell type-specific perturbation data on the other has led to ongoing debate regarding the rules that govern BG circuit function.

As predicted by anatomy[9], activating dMSNs can rapidly suppress, while activating iMSNs can rapidly enhance, the activity of inhibitory output neurons of the BG in the substantia nigra[10–12]. At a behavioral level, activation of dMSNs consistently produces opposite effects to those of activating iMSNs with respect to locomotion[11], ongoing motor sequence production[13,14], reinforcement[15,16], and value-based decisions[17]. However, the activity of dMSNs and iMSNs in sensorimotor striatum appears to be largely positively correlated around action initiation[3,18,19] or around transitions between actions[20]. Such observations of concurrent activation of the two pathways have been used to argue against the hypothesis that they functionally oppose each other[3,18].

A longstanding, and potentially reconciling, view of BG circuit function is that the two pathways might contribute to selection amongst various actions in a competitive manner[2,21,22]. In this view, action selection proceeds through combined promotion of motor programs by the direct pathway, and suppression of motor programs by the indirect pathway. Such a model predicts broad coactivation of the two pathways during action production even as they function in opposition to each other. Notably, this framework also predicts that sustained suppression of action should promote large-scale decorrelation or even anticorrelation between the two pathways. This possibility, to our knowledge, remains untested. Activity of iMSNs should be elevated to suppress action, while the activity of dMSNs should be limited until action is released. Such observations would naturally reconcile currently disparate interpretations of BG circuit function.

To test this hypothesis we employed a variant of an interval discrimination task[23,24] requiring a series of self-initiated and cued actions, and critically, a sustained period of action

suppression. This design allowed us to assess whether the two pathways exhibited broad coactivation during action production and opponency during action suppression. Furthermore, the task requires dynamic suppression of distinct, lateralized behaviors over time, an ideal situation to assess whether the two pathways exhibit any action-*specific* opponent signals during action suppression. During this behavior, we then recorded activity from dMSNs and iMSNs in the dorsolateral striatum of mice using fiber photometry. We found that both pathways displayed phasic activation during action production, as previously reported. However, during action suppression we observed multiple clear signatures of functional opponency. First, overall iMSN activity was sustained whereas dMSN activity fell to or below baseline levels. Second, we observed evidence of action-specificity in these opponent signals in the form of opposite patterns of inter-hemispheric dynamics in the two pathways. A shift in the relative balance of activity between the two hemispheres developed during a trial as the need to suppress movement to one side of the body waned, and the need to suppress movement to the opposite side of the body grew.

To assess the functional importance of the observed patterns of neural activity, we performed a series of optogenetic inhibition experiments. These experiments provided further support for functional opponency and at the same time revealed a double dissociation between the respective roles of the direct and indirect pathways in dorsolateral striatum. Optogenetic inhibition of dMSNs, but not iMSNs, produced a slowing of movement whereas optogenetic inhibition of iMSNs, but not dMSNs, disrupted the ability of mice to suppress action.

These findings demonstrate clear opponency in the endogenous activity of the two major BG pathways during behavior, bringing into alignment molecular, anatomical, optogenetic and neurophysiological data in support of functional opponency between the two circuits. The data also suggest a novel mode of functional distinction between the two pathways. The direct pathway in sensorimotor striatum appears necessary for augmenting action vigor but, surprisingly, not the specification of which action is taken. The indirect pathway however, appears critical for sustained suppression of specific behaviors. These results not only help to resolve ongoing debate, but provide new insight into how BG circuitry can contribute to distinct aspects of action production and suppression, with broad implications for understanding the neural mechanisms of both normal and pathological behavioral control.

Results

Production and proactive suppression of action

We trained mice on a variant of a two-alternative interval categorization task wherein subjects were required to suppress movements during interval presentation. Briefly, trials were self-initiated by the mouse inserting their snout into a centrally located initiation nose port, eliciting a brief auditory tone (Fig 1a). Mice were required to maintain their position in the initiation port (fixation) until a second auditory tone was delivered. This second tone was delivered at a delay that was randomly chosen from a set of 6 intervals, symmetric about 1.5s, and ranging from 0.6s

to 2.4s. After delivery of the second tone, animals were free to choose either of two choice ports
located at an equal distance to either side of the initiation port. Rewards were delivered for choices
to one side ("short" choice) if the presented interval was shorter than a 1.5s decision boundary,
and at the opposite choice port ("long" choice) if the interval was longer than 1.5s. Mice learned
to categorize interval stimuli much longer or shorter than the decision-boundary with high
accuracy (92.1±0.7%, 0.6s and 2.4s intervals, mean ± s.e.m. n = 14 mice), yet choices were more
variable for intervals nearer to the decision-boundary (Fig 1b). In addition, mice produced a
stereotyped movement profile over each trial (Fig 1c). Movement speed increased leading to trial
initiation, followed by a brief period of postural adjustment before animals settled into immobility
until the second tone was delivered. Immediately following second tone delivery, movement
speed increased again as animals executed their choices. If animals failed to maintain fixation in
the initiation port until the second tone, an error tone was immediately delivered and the trial was
terminated (36.5±2.1% of all trials, n=14 mice). We will refer to these trials as *broken fixations*
throughout the text. Interestingly, animals often entered a choice port even after breaking fixation
and aborting the trial (52.1±4.9% of broken fixation trials, n=14 mice). These choices were
executed with a similar timecourse as valid choices (Extended Data Fig. 1a) and were largely
"appropriate", toward the "short" choice port when breaking early in a trial, and toward the "long"
port when breaking late in a trial (Fig 1d, Extended Data Fig. 1c). The pattern of broken fixations
reflects the overall reward associated with the two choices over time, and not the likelihood of
second tone occurrence (Extended Data Fig. 1d). These data are consistent with animals
developing a dynamic motor plan that remains latent as long as it is successfully suppressed.
Failure to suppress this temptation led to premature execution of the planned action.
135

Opposite modulation of striatal direct and indirect pathways during action suppression

To probe the large-scale activity of the direct and indirect pathways for signs of functional
opponency during mobility and active suppression of lateralized movements, we recorded the
activity of dMSNs and iMSNs in the dorsolateral striatum during task performance. We combined
mouse lines expressing Cre recombinase in either dMSNs (D1-Cre EY217Gsat line) or iMSNs
(A2a-Cre, KG139Gsat line) with cre-dependent viral expression of the calcium indicator
GCaMP6f[25](Fig. 2a), using coordinates previously shown to contain coactive dMSNs and iMSNs
during movement[3] (Extended Data Fig. 2a). No significant differences in behavior were detected
between the two mouse lines (Extended Data Fig. 3). We then used fiber photometry[24,26](Fig. 2
a-d) to access the pooled activity of a local population of dMSNs or iMSNs in the dorsolateral
striatum. To determine whether gross differences in activity patterns were present across a trial,
we first examined the combined activity of neurons located in both hemispheres. In individual
animals, we observed that activity of either dMSNs or iMSNs increased around task epochs when
animals were required to take action, namely trial initiation and choice execution, consistent with
the commonly observed coactivation of the two pathways (Fig. 2 e,f). Indeed, across all animals,
though the time courses of activity appeared to differ slightly, the mean activity of the two
pathways was indistinguishable around trial initiation (Fig. 2 g,h, iMSN vs dMSN = 0.141 [-1.09,
1.37] zΔF/F, p = 0.8065; Effect Size, 95%[CI], p-value). However, activity in dMSNs and iMSNs

displayed marked differences during interval presentation, when mice were required to suppress
movement (Fig. 2 e,f). Across all animals, iMSN activity was elevated relative to dMSN activity
(Fig. 2 g,h, Extended Data Fig. 4a, iMSN>dMSN = 0.783 [-0.071, 1.64] zΔF/F, p = 0.0345,). This
opponent pattern grew as the period of sustained action suppression wore on, perhaps reflecting
the growing need to suppress action as expectation of cue delivery grew during a trial and as
mice became certain that a choice to the long port would ultimately be rewarded. Indeed, we
detected modest yet significant upward and downward deviations in the rate of change of activity
in dMSNs and iMSNs preceding broken fixations as compared to time-matched control periods
(Extended Data Fig. 4b,d), indicating that disturbances in activity patterns in the two pathways
were associated with the failure to successfully suppress action.
165
To assess whether the need to suppress action in general might grow over time during a
trial, we computed the probability that mice break fixation at each time bin within the delay period
conditioned on their not having broken fixation up to that point, a quantity known as the hazard
rate. In the context of this behavioral task, computing the hazard rate as opposed to the overall
probability of breaking fixation (Fig. 1d, Extended Fig. 1c) effectively controls for the fact that
animals experienced more instances of early time bins in the delay and thus had more
opportunities to break fixation early in the delay. The hazard rate of broken fixation behavior
demonstrates that after a brief dip in broken fixation around the decision boundary, the likelihood
of breaking fixation rises dramatically (Extended Data Fig 1b). We next examined the hazard rates
of broken fixation conditioned on subsequent choice, and found that the early mode in the overall
hazard rate of broken fixations was comprised of trials where the mice subsequently made a short
choice, whereas the late rise in the overall hazard was comprised of trials where mice
subsequently made a long choice (Fig. 3a). Notably, not only is the task requirement that mice
map earlier times in a trial onto action towards one side of the body and later times in a trial onto
action towards the other side of the body reflected in the probability of broken fixations and
subsequent choices over time, but the urge to break fixation in a particular direction is asymmetric.
The urge to break fixation and make long choices late in the trial appears to far outweigh the urge
to break fixation and make short choices early in the delay period. If MSN activity during
successfully completed trials acted to suppress these lateralized urges to act early and late, we
might expect differences in the time course of activity between the two hemispheres. Indeed, in
the hemisphere contralateral to the rewarded location for "long" stimuli (contra-long, CL, Fig. 3b)
iMSN and dMSN activity steadily increased and decreased throughout the delay period,
respectively (Fig. 3c, , Extended Data Fig. 4b, difference between pre and post decision boundary
mean activity: iMSN:CL = 0.423 [0.006 0.840]zΔF/F, p = 0.0462, dMSN:CL -0.535 [-1.017 -
0.054]zΔF/F, p = 0.0272).
191
In contrast, in the hemisphere contralateral to the rewarded location for "short" stimuli
(contra-short, CS, Fig. 3b) activity levels in both pathways were relatively constant as compared
to the CL hemisphere (Fig. 3c, , Extended Data Fig. 4b, iMSN:CS = -0.104 [0.521 0.313]zΔF/F, p
= 0.928, dMSN:CS -0.040[-0.052 0.442]zΔF/F, p = 0.999)). These data reflect a situation where
lateralized patterns of activity reflected the strength of urge to move contralaterally over time.
Thus, relative levels of activity between the two hemispheres varied over time, and in opposite
directions in the two pathways (Fig. 3d). Such observations may indicate that BG circuitry residing

in a particular hemisphere is preferentially recruited to suppress movements to the contralateral
direction when and to the degree that the animal is tempted to move in that direction.
201

Broadly opponent yet distinct functional contributions of striatal direct and indirect pathways to the control of action

To test whether the observed opponent patterns of neural activity in the two pathways
reflected true functional opponency or whether they amounted to simply a correlation with the
predictions of the opponent function hypothesis, we next performed a series of optogenetic
inhibition experiments. We combined the same Cre lines used to label MSNs in photometry
experiments with cre-dependent viral expression of the light-activated proton pump ArchT[27] and
implanted tapered optical fibers[28] in the dorso-lateral striatum to enable inhibition of iMSN or
dMSN activity during action suppression (Fig. 4a, b). We first characterized the effect of
photoinhibition on neuronal firing by performing extracellular electrophysiological recordings from
striatal neurons during a quiet awake state in the absence of a behavioral task (Fig. 4c, i, Extended
Data Fig. 5a-d). Illumination of striatal tissue produced rapid and sustained inhibition of firing in
both iMSN-ArchT and dMSN-ArchT mice, indicating similar levels of inhibition of the two
pathways. We then delivered green light to the fiber(s) on a random minority of trials (30%),
starting at trial initiation and ramping off over 250ms starting at either the second tone onset or
when the animal broke fixation, whichever occurred first (Fig. 4b). Bilateral optogenetic inhibition
of iMSNs produced a near-complete inability of mice to suppress movement during interval
presentation (Fig. 4j). Animals broke fixation on 35±6% (n=4 mice) of non-inhibited trials and on
86±7% of trials when iMSNs were inhibited bilaterally (Fig. 4i, mixed-effects model, Interaction
Genotype:Hemisphere:Laser,   p   < 10^{-4},   odds   ratio   iMSN:Bilateral:LaserOff   /
iMSN:Bilateral:LaserOn = 0.0585 [0.0362, 0.0945], P<0.05). However, levels of broken fixation
were unaffected by bilateral inhibition of dMSNs during equivalent trial epochs (Fig. 4d, odds ratio
dMSN:Bilateral:LaserOff / dMSN:Bilateral:LaserOn = 0.8177[0.6271, 1.0664], p = 0.3544). On
broken fixation trials followed by movement toward a choice port, the first 250ms of this movement
overlapped with the ramping off of the light stimulation. We therefore asked whether these
movements were affected by inhibition of either iMSNs or dMSNs. In contrast to the observed
effect of iMSN inhibition on action suppression, we found that dMSN inhibition, but not iMSN
inhibition, resulted in a significant increase in movement time during broken fixation trials (Fig. 4e,
k, iMSN:laserOff vs iMSN:laserOn = 7.82 [-482, 497.3] ms, P = 1; dMSN:laserOff vs
dMSN:laserOn = -630.30 [-1193, 97.6] ms, P = 0.021), consistent with a role specifically for the
direct pathway in augmenting the vigor of movements[29]. Furthermore, dMSN activity in the DLS
was specifically necessary for movement invigoration immediately in advance of movement
initiation, as movement times were not affected during valid trials, when the laser began ramping
off during the time it took animals to initiate their choice movement, nor in a subset of experiments
when inhibition was applied during execution of the choice movement, starting when animals left
the initiation port (Extended Data Fig. 6).
238
239

Given the observed interhemispheric dynamics in the photometry signals collected from
MSNs (Fig. 3), we next asked whether unilateral MSN inhibition would disrupt action suppression
preferentially at the times when activity in a given hemisphere appeared to be most engaged.
Indeed, while unilateral iMSN inhibition produced a more modest increase in broken fixations
overall (Fig. 4j), the timing of broken fixations was systematically related to the laterality of
inhibition. Mice were more likely to break fixation early or late when iMSNs contralateral to the
"short" or "long" choice were inhibited, respectively, as compared to non-inhibited trials (Fig. 4l).
Taking the difference in probability of breaking fixation between inhibited and control trials as a
function of time during the trial gives a measure of the time-course of the effect on action
suppression for each hemisphere. Superimposed, these two measures cross near the 1.5s
decision boundary (Fig. 4n, right, Extended Data Fig. 7a), mirroring the precise contingency
between reward and action over time during a trial. These data provide strong evidence that
proactive suppressive control is handed off from the indirect pathway of one hemisphere to its
counterpart on the opposite hemisphere around the time animals should be switching the direction
of their planned movement from one side of the body to the other. Once again, we observed no
consistent effects of inhibiting dMSNs on the timing of broken fixations (Fig. 4 h).
256
Lastly, given the widely assumed importance of BG circuits in action selection, we asked
whether inhibition of MSNs affected the probability that particular actions were executed. When
inhibiting iMSNs, we observed a consistent increase in the probability that an animal would
execute a choice to the port contralateral to the site of iMSN inhibition, as compared to non-
inhibition trials, after breaking fixation (Fig. 4l). This effect did not simply reflect the fact that
animals were more likely to make short or long choices after early or late broken fixations,
respectively, because it was present in broken fixations made both before and after the decision-
boundary (Extended Data Fig. 7c). Features of the choice movements following iMSN inhibition,
as quantified through high-speed tracking of the animals' nape, were indistinguishable from
choice movements performed in the absence of iMSN inhibition (Extended Data Fig. 8). Once
again, we observed no significant effect of unilateral dMSN inhibition on lateralized choice
behavior after broken fixations (Fig. 4f) (odds ratio iMSN:CL:LaserOff /iMSN:CL:LaserOn =
0.257[0.1430, 0.463], P<10^{-4} , iMSN:CS:LaserOff / iMSN:CS:LaserOn = 3.369[1.7069, 6.651],
P<10^{-4} , dMSN:CL:LaserOff / dMSN:CL:LaserOn = 0.917[0.4484, 1.875], p=1 dMSN,CS,LaserOff
/ dMSN,CS,LaserOn = 0.737[0.3782, 1.438], p = 0.87), even when dMSN inhibition was applied
during execution of the choice movement (Extended Data Figure 6). These data, together with
the observed inter-hemispheric dynamics of the endogenous activity of iMSNs, strongly suggest
that the indirect pathway of a given hemisphere is engaged to suppress contralateral movements
when those particular actions would be tempting and thus in greater need of suppression.
276

Discussion

278
279   Here we demonstrate for the first time, to our knowledge, clear activity signatures of large-
280 scale functional opponency between neurons initiating the two major BG pathways in the normal,

281 non-pathological state. In particular, we observe opposite patterns of activity in the two pathways
282 when movements must be proactively and persistently suppressed.
283
Action suppression can be broadly separated into two classes. Reactive suppression
involves stopping behavior in course when presented with an external stimulus. In contrast,
proactive suppression involves selectively inhibiting particular response tendencies using
advance knowledge. The behavioral context we studied requires proactive suppression of time
varying response tendencies to move to the left or right. When subjects were required to suppress
the urge to move in a given direction, on average iMSNs located contralaterally to that direction
exhibited higher levels of activity than iMSNs located ipsilaterally. Consistent with these data,
functional magnetic resonance imaging data in humans and electrophysiological data in non-
human primates suggests that iMSNs might be selectively engaged to proactively suppress
action[30–33]. Leveraging the genetic access to iMSNs and dMSNs afforded by the use of mice as
a model organism, our data clearly establishes not only that iMSNs in the DLS were selectively
engaged as animals suppressed particular lateralized response tendencies, but that iMSNs, and
not dMSNs, were necessary for it. In addition, successful action suppression relied on iMSNs in
a given hemisphere at different points in time depending on learned task demands, indicating that
specific subpopulations of iMSNs can be deployed to dynamically shape action suppression in
time. While multiple studies to date have observed coactivation of dMSNs and iMSNs around
movement[3,18,20], a finding we reproduce here, transient decorrelation of activity between the two
pathways has been reported around transitions between actions in a behavioral sequence[20].
Given our observations that activity in the two pathways is decorrelated or even anti-correlated
during proactive action suppression, we hypothesize that previously reported transient
decorrelations between the two pathways may arise when the brain must largely suppress the
production of actions at behavioral transitions.
306
The current data also demonstrate distinct contributions of dMSNs and iMSNs in the DLS
to aspects of motor function beyond proactive action suppression. Bradykinesia seen in PD
patients is thought to result primarily from the loss of dopamine neurons in the Substantia nigra
pars compacta (SNc)[1]. Prior studies have identified that both dopaminergic input to the striatum
and the activity of striatal neurons, in particular dMSNs, are important for invigorating movement[34].
However, it was not clear based on previous work whether iMSN activity is necessary for
invigorating movement. Here we show that inhibiting dMSN activity slowed movement, consistent
with previous work, yet this did not affect which action was selected. Conversely, we show that
iMSN inhibition did not lead to less vigorous movements, but instead disrupted the suppression
of actions, and that this disruption reflected at least some degree of action specificity in iMSN
function as it occurred alongside a change in the likelihood that lateralized actions were produced.
Based on the current data, it remains uncertain the degree to which iMSNs act on specific actions,
although in sensorimotor striatum we might expect iMSNs in a given location to act on actions
involving parts of the body represented in the somatosensory motor cortical areas that provide
input to that striatal location[35].
322
At first glance, our results may appear to conflict with studies demonstrating that
optogenetic activation of dMSNs is sufficient to alter action selection. However, sufficiency does

not imply necessity, and our experiments, by inhibiting iMSNs and dMSNs, represent a test of
whether activity in these two cell types is necessary for specific features of behavioral control. In
addition, many of the studies that have demonstrated an effect of optogenetic activation of dMSNs
on action selection have focused on dorso-medial striatum (DMS) [15,17], and here we focus on
DLS. However, our recordings from the striatum revealed that optogenetic activation of iMSNs or
dMSNs in DLS, unlike the optogenetic inhibition employed in our behavioral experiments, resulted
in clear inhibition of other MSNs (Extended data figure 5). Thus, behavioral effects of activating
dMSNs or iMSNs could potentially be influenced by inhibition of neurons of the non-targeted MSN
class. Careful titration of optogenetic activation to match physiologically normal ranges of activity
might thus represent a critical improvement on methods for relating the effects of optogenetic
activation of targeted cell types to their normal function during behavior[36].
336
Interestingly, recovery of function in an experimental model of PD has been found to be
associated with a return to near normal levels of iMSN but not dMSN activity, consistent with
some degree of primacy for the indirect pathway in action production[37]. Together with the data
presented here, these results suggest the intriguing possibility that the indirect pathway provides
an inhibitory "mask", specifying which actions not to produce, while the direct pathway provides a
gain signal, as opposed to an action selection signal[2], for commands that are pushed through the
mask. In this view, the selective function of the sensorimotor BG circuits on actions can be
predominantly produced through the indirect pathway, a novel proposal that should serve as a
basis for future experiments. In the present data, both pathways were more active around
movements toward the contralateral side of the recording site (Extended Data Figure 9),
suggesting that suppressing and promoting signals may be targeted towards nearby regions in
the space of possible actions, perhaps in part by mechanisms in other brain systems such as the
cortex, thalamus, and cerebellum[2,38].
350
Here we focus on MSNs in the dorsolateral striatum, sometimes termed sensorimotor
striatum due to the preponderance of inputs from sensory and motor cortical areas that terminate
there[35,39,40]. Though we targeted this specific region, our measure of neural activity, as in other
studies employing photometry, pooled within a class of MSNs with likely diverse tuning properties.
While previous work has identified correlations between MSN firing and kinematic or motivational
variables[20,41–43], the lack of conclusive information regarding the tuning properties of individual
MSNs has been a recurring issue in efforts to determine the functional importance of the direct
and indirect pathways. For example, experiments might well be blind to functional organization
wherein neurons in the two pathways controlling specific actions produce anti-correlated activity,
because the granularity with which actions are encoded by striatal neurons, and resultantly, how
to design a behavioral scenario that would isolate these representations, is beyond current
understanding.
363
We were able to circumvent the general problem of diverse and unknown selectivity of
neurons in two ways. First, by training animals to remain immobile for an extended period in the
task, we pushed the brain toward suppressing the majority of actions, a behavioral manipulation
we hypothesized to have a common effect on the neurons in a given cell class regardless of their
selectivity for actions. Second, the striatum in a given hemisphere shows enhanced functional

involvement in contralateral movements[11,44]. By training animals to perform a task wherein the
relative value of left/right lateralized movements was varied over time during a prolonged period
of action suppression, and observing or manipulating activity on one hemisphere at a time, we
demonstrate a circuit mechanism by which dorso-lateral striatal circuits of the BG can control
generalized movement vigor or the suppression of actions.
374
Adaptive behavior fundamentally involves the interplay of action promoting and action
suppressing mechanisms. The data presented here demonstrate that in sensorimotor striatum,
elements of the direct and indirect BG pathways can express opposite patterns of modulation and
be required for generally opponent yet distinct promoting and suppressing aspects of motor
function. Knowledge of how circuits in other regions of the striatum, the BG at large, or elsewhere
in the brain mediate this interplay represents a critical avenue toward a fundamental
understanding of animal behavior. Such knowledge also has the potential to inform the
engineering of artificial systems that can behave appropriately in complex environments, as well
as a mechanistic understanding and the design of effective therapies for neurological and
neuropsychiatric disease.
385
386

Data availability statement: The data and analysis code that support the findings of this
study are available from the corresponding author upon reasonable request.

Acknowledgements: We would like to thank Brian Lau, John Krakauer, and Bassam Atallah
for comments on versions of the manuscript and the entire Paton lab for feedback during the
course of this project. We would also like to thank the ABBE Facility and the Scientific Hardware,
Histopathology and Rodent Champalimaud Research Platforms for unparalleled technical
assistance, and Ben Zarov and Daniela Domingues for helping with the training of some of the
animals included in this study. This work was developed with the support from the research
infrastructure Congento, co-financed by Lisboa Regional Operational Programme (Lisboa2020),
under the PORTUGAL 2020 Partnership Agreement, through the European Regional
Development Fund (ERDF) and Fundação para a Ciência e Tecnologia (Portugal) under the
project LISBOA-01-0145-FEDER-022170. The work was funded by an HHMI International
Research Scholar Award to J.J.P (#55008745)., European Research Council Consolidator grant
(#DYCOCIRC - REP-772339-1) to J.J.P., a Bial bursary for scientific research to J.J.P.
(#193/2016), internal support from the Champalimaud Foundation, and a PhD fellowship
(PD/BD/105945/2014) from Fundação para a Ciência e a Tecnologia to B.F.C.

Author contributions: J.J.P. and B.F.C. conceived of the experiments and wrote the
manuscript. S.S. assisted in analysis of the fiber photometry data and revised the manuscript.
B.F.C. carried out all experiments and analyzed the data. J.J.P. supervised all aspects of the
project.

Competing interests: The authors declare no competing financial interests.

Methods

Animals:

Adult (over 2 months) male and female mice of A2a:cre (KG139) and D1:cre (EY217) lines [45] were used for this study under the protocol approved by the Champalimaud Foundation Animal Welfare Body (Protocol Number: 2017/013), the Portuguese Veterinary General Board (Direcção-Geral de Veterinária, project approval 0421/000/000/2018) and in accordance with the European Union Directive 2010/63/EEC. Mice were group housed prior to surgical procedures and singly housed following surgery in an inverted 12h dark/light cycle. Mice were maintained under water deprivation for all behavioral experiments (>80% body weight from baseline ad libitum period before deprivation).

Behavioral apparatus:

The behavioral box (20 x 17 x 19 cm), contained 3 nose ports and a speaker. The behavioral box consisted of 3 front walls (135 degree angle between the center and the side walls) 2 side walls and a back wall with a 90 degree angle between them. Each of the three front walls contained a nose port equipped with an infrared emitter/sensor pair to access port entry and exit times. The central nose port was defined as the trial initiation port, and choices were reported at the lateral nose ports. For correct trials, a 4-6 µL calibrated water reward was delivered using a solenoid valve. Tones were delivered through a speaker mounted on the center wall. All sensors and effectors in the behavioral box were read and controlled using a microprocessor (Arduino Mega 2560, Arduino) via a custom circuit board. The task was implemented by the microprocessor, which outputted data via a serial communication port to a desktop computer running custom Python-based software. High-speed video was acquired at 120fps and 640*480 pixel resolution (FL3-U3-13S2, FLIR).

Behavioral task:

Mice were trained to categorize interval durations as either short or long by performing right and left choices as previously described in Soares et al.[24]. Briefly, mice self-initiated trials by entering the central nose port, triggering the delivery of a pair of tones (7,500 Hz, 150 ms) separated by one of 6 randomly uniformly sampled selected intervals (0.6, 1.05, 1.26, 1.74, 1.95 and 2.4 s or 0.6, 1.26, 1.38, 1.62, 1.74, and 2.4 s. Stable performance was usually achieved after 3-4 months of training. Trial availability was not signaled to the animal but inter-trial onset interval was kept constant within each animal (7-8 s). Thus, initiation port entries before the point that a trial became available were ineffectual. After the first tone was presented, mice were required to maintain interruption of the center nose port IR beam until the second tone was delivered, we refer to this action as "fixation" throughout the text. If the mouse departed the port before the second tone, an error tone (150ms of white noise) was played and the next trial availability delayed (timeout). We refer to these trials as "broken fixations". To prevent incorrectly flagging

trials as broken fixations due to sporadic state transitions in the IR beam, we only counted a trial
as broken fixation after the beam had been continuously uninterrupted for 50 ms. After both tones
were played, mice reported their judgments by entering one of the two laterally located nose ports
over the next 10 seconds. For intervals shorter than a 1.5 s category boundary, responses were
reinforced at one of the lateral ports. For intervals longer than 1.5 s, responses were reinforced
at the opposite port. Incorrect responses were followed by a white noise burst (150 ms) and a
timeout (12-18s inter-trial onset interval). The short/long vs right/left contingencies were
counterbalanced across animals. Therefore, we adopt the nomenclature of Contra-short/Contra-
long hemisphere throughout the paper. Sessions typically lasted 2 hours.
Psychometric functions were fit using a 4-parameter logistic function:

$$P(x) = (u - l) \times \frac{e^{\frac{x-b}{s}}}{1 + e^{\frac{x-b}{s}}} + l,$$

456
457 where P is the performance of the animal on interval x, u and l the upper and lower
458 asymptote of the curve, respectively, b the bias parameter and s the slope parameter.
459
460 Unless otherwise stated, broken fixations before the earliest interval (0.6s) were excluded
461 from analysis, as these often reflected the failure of animals to properly settle in the initiation port.
462 To calculate hazard of breaking fixation at a particular time bin in trial, $H(k)$, we used the
463 following equation:

$$H(k) = \frac{B(k)}{\sum_{j=k}^{T} B(j) + \sum_{j=k}^{T} C(j)}, \quad t < k < t + \Delta t$$

464
465
466
467 where $B(k)$ is the number of broken fixations that occured at time bin k, $\sum_{j=k}^{T} B(j)$ the
468 sum of all broken fixations that occured in the time bins greater or equal to k, up until the longest
469 possible interval T (2.4 seconds), and $\sum_{j=k}^{T} C(j)$ the sum of all completed trials that occured
470 from time bin k until T.
471

Viral injections and fiber implantation:

All surgeries were performed with mice under isoflurane anesthesia (1-2% at 0.8 L/min).
We stereotaxically targeted the dorsolateral striatum (DLS, coordinates below) bilaterally for all
viral deliveries.
After achieving stable performance (usually >3 months), mice were allowed to regain
baseline weight and were subject to viral injection and fiber implantation in the same surgery.
Mice health was assessed daily and after at least 5 days, the water deprivation regime was
reinstated. To make sure subjects recovered their pre-surgery performance before data collection
sessions (photometry or optogenetics), they were gradually retrained in the task without and then

with fibers attached. Upon reaching stable performance (1-2 weeks), data collection began. We
alternated recorded/manipulated hemisphere every day.
For fiber photometry experiments, we injected 300nL of a mixture of two viruses: AAV1-
Syn-Flex-GCaMP6f (titer ~1 x 10^13 gc/mL; University of Pennsylvania Vector Core) and AAV1-
CAG-Flex-tdTomato (titer ~0.5 x 10^12 gc/mL; University of Pennsylvania Vector Core), at a 5:1
ratio, in DLS striatum (single injection, AP 0.5mm, ML 2.1mm, DV 2.6mm from pia) using an
automated microprocessor controlled microinjection pipette with micropipettes pulled from
borosilicate capillaries (Nanoject II, Drummond Scientific). Injections were performed at 0.2 Hz
with 2.3 nL injection volumes per pulse. For all injections, the micropipette was kept at the injection
site 10 min before withdrawal. After injection, we implanted, bilaterally, 2 fibres (MFC_200/245-
0.53_ZF1.25(G)_FLT, DORIC LENSES) 200 µm above the injection site.
For optogenetic inhibition experiments, we injected AAV5-CAG-FLEX-ArchT-tdTomato
(titer 10^13 gc/mL, Addgene) in the DLS. For each hemisphere, we made two injections (500nL,
AP 0.5mm, ML 2.1mm, DV 2.7mm and 3.1mm from bregma) and implanted one fibre (Lambda-B
fibre 200um core, 0.39NA, 1.5mm emitting length, Optogenix) near the injection site (AP 0.5mm,
ML 2.1mm, DV 3.5mm from bregma) .
497

Fiber photometry

The photometry apparatus was adapted from[26]. For all experiments, a single blue laser
was coupled to a patchcord (100 µm core diameter, 0.22 NA) and connected to a collimator
adapter (EFL 4.5 mm, NA 0.50) and a neutral density filter. Dichroic mirrors were fixed inside the
main unit, allowing for 473 nm light delivery and GCaMP6f and tdTomato fluorescence detection.
The 473 nm light was coupled into a patchcord (200 µm core diameter, 0.48 NA) using a lens
(EFL 4.5 mm, NA 0.50) and a rotary joint. The patchcord was mated to one of two chronically
implanted optical fibers (200 µm core diameter, 0.48 NA). Laser intensities at the patchcord tip,
before mating to the chronically-implanted fiber, were 15-40µW. For detection of GCaMP6f
fluorescence, light was collected by the lens, transmitted and reflected by the dichroics before
final filtering and focusing into a photodetector. For detection of tdTomato fluorescence, light was
collected by the lens and transmitted through all dichroics before final filtering and focusing into a
second photodetector. Photodetector output was digitized at 1 kHz (PCIe 6351, National
Instruments) and recorded using custom software in Bonsai[46].
512

Fiber photometry data analysis

All photometry data analysis was performed with custom MATLAB software. Raw data
was downsampled to 100Hz and low-pass filtered at 20Hz. Slow fluctuations were removed by
subtracting a fitted polynomial to the raw signal (order < 5). For each session, ΔF/F was calculated
for both channels as $\Delta F/F_t = (F_t - F_0) / F_0$, where F_0 was calculated as the 10th lower percentile
from the filtered signal. Similarly to[24], robust regression using GCaMP6f and tdTomato ΔF/F was
performed and the coefficient estimates were used to calculate a predicted GCaMP6f ΔF/F based
on the observed tdTomato ΔF/F. This predicted GCaMP6f ΔF/F was then subtracted to the

observed GCaMP6f ΔF/F to calculate the corrected ΔF/F. In order to compare across sessions,
each session's corrected ΔF/F was z-scored using the mean and standard deviation calculated
from a baseline period (5 to 2 seconds before trial onset). Signals for individual trials were then
re-zeroed by subtracting the average of a period of 5 to 2 seconds before trial onset.
In order to compare the rate of change of activity aligned to broken fixations to time-
matched valid trials (Extended Data figure 4c-d) we began by calculating the first derivative of the
corrected ΔF/F for each trial. For each broken fixation trial, we aligned the trace to the timestamp
of the broken fixation and, in order to compare to a time-matched valid trial, we cropped the
average trace of all valid trials to the same length and aligned to the time of the broken fixation.
We repeated this process for all broken fixation trials.
531

Acute recordings

To confirm the ability to inhibit medium spiny neurons, we performed acute recordings in
the dorsal striatum of untrained animals. Briefly, similarly to trained animals, we virally expressed
ArchT in D1-Cre or A2a-Cre animals. After 3-5 weeks of viral expression, we implanted a small
headpost and a ground pin contralateral to the recorded hemisphere. After allowing animals to
recover, a small round craniotomy (1.5mm diameter) was opened over the same coordinates as
the virus injection. Recordings were performed while animals were head-restrained using custom
built headbar holders and on top of a passively rotatable cylinder.
A silicon probe (ASSY 77-H2, Cambridge NeuroTech) with a tapered optical fibre, identical
to the one used for optogenetic inhibition during behavior, glued to the back was slowly lowered
into dorsal striatum. Electrophysiology and laser modulation data were acquired at 30kHz with
OpenEphys hardware[47] and Bonsai[46].
Every 10 to 25 seconds, an interval was drawn from the set [0.60,1.05,1.95, 2.40]s and
light was continuously delivered during that duration. The range of power used was identical to
that used during the task and, similarly, was ramped off for 250ms after the drawn delay had
elapsed.
In a second batch of animals, we tested the extent of excitation and inhibition when
activating medium spiny neurons using ChR2. To achieve ChR2 expression we virally expressed
ChR2 (AAV9-EF1a-double floxed-hChR2(H134R)-EYFP-WPRE-HGHpA) or we used double
transgenic mice (A2a-Cre or D1-Cre crossed with Ai32 mouse line[48]). Light intensity was set to
0.5-1mW at the end of the fiber. The remainder of the protocol was identical.
Electrophysiology data was sorted using Kilosort2 (github.com/MouseLand/Kilosort2) and
manually curated using Phy (github.com/cortex-lab/phy).
Cells that fired, on average, less than 1 / 2.4 spikes/s (detection limit during inhibition)
during the baseline were excluded from analysis.
To determine if a cell was modulated, we compared the firing rate in a baseline period
prior to light delivery (-3 to -1s) to the average firing rate during the stimuli. We computed a t-test
per cell and considered whether a cell was significantly modulated after correcting for multiple
comparisons ($p < 0.05$/(Total number of recorded units)).
To classify units as putative medium spiny neurons we used previously described
criteria[49,50] based on firing statistics and waveform duration. Briefly, we classified units as putative

medium spiny neurons if baseline firing rate was less than 10Hz, coefficient of variation greater
than 1.5 and waveform duration greater than 600ms.

Optogenetic manipulations during task performance:

A 556nm laser (500mW, Optoelectronics Technology) was used as a light source to
activate ArchT. Briefly, the output of the laser was aligned to an acousto-optic modulator (AOM
MTS110-A3-VIS, AA OPTO-ELECTRONIC) and fibre launched into a patchcord (200 μm core,
0.48NA). The output of the patchcord was then connected to a power splitting rotary joint
(FRJ_1x2i_FC-2FC_0.5, DORIC) and finally into 1 or 2 patch cords (200 μm core, 0.48NA) that
connected to the animal's implanted tapered fibres (Lambda-B fibre 200um core, 0.39NA, 1.5mm
emitting length, Optogenix).
The light intensity was controlled by modulating the AOM using a dedicated Arduino Mega
2560 board connected to a DAC board (MCP4725, Sparkfun). Regardless of the stimulation
duration, all protocols included a 250ms linear ramping off, designed to reduce the potential for
rebound excitation[51].
The inhibition protocol was applied on a randomly selected 30% of trials. Light was
continuously delivered, with the onset aligned to the initiation of a trial (centre nose port) and
lasting until second-tone delivery or the exit of the subject from the centre-poke, whichever
occurred first. For all analyses of optogenetic manipulation data, the first 15 trials of each session
were excluded. Laser power was set to be 23 to 31mW at the end of the fibre for all animals.
For a subset of direct pathway animals (n=3, Extended Data Figure 6), a third protocol
was used wherein light was turned at termination of the second tone up until choice or 400ms,
whichever occurred first.
Each day, we changed the location of the inhibition (CS or CL, n=4 A2a-cre, n=6 D1-cre)
and, after collecting data from the unilateral manipulation conditions, we silenced both
hemispheres simultaneously (n=4 A2a-Cre and n=4 D1-Cre). Data from 4/6 and 3/5 D1 and A2a-
Cre animals was acquired for all conditions (i.e. unilateral and bilateral). For single animals, unless
otherwise stated, all analyses were performed by concatenating all trials from all sessions of the
same manipulation condition.
To calculate the fraction of trials wherein animals broke fixation as a function of time, in
control versus manipulated trials (i.e. Fig. 4g and m), we took for each condition all trials wherein
animals broke fixation after 0.6s and calculated their probability density as a function of time over
all trials of that condition (control vs manipulated).
595

Movement trajectories

Offline tracking of mouse position was performed using DeepLabCut targeting the nape
of each animal [52].
To compare trajectories between different conditions (Extended Data Figure 8), we first
computed a null distribution of integrated euclidian distances over time within a reference
condition (non-stimulated broken fixation condition), using all pairwise trial combinations. We
then compared pairs of trials where one trial was taken from a test condition, and the other from
the reference condition, for all pairwise combinations, resulting in a distribution of integrated
euclidean distances between reference and test trials. To compute distances between positions

from trials of differing duration, the longer trial of the pair was cropped to match the duration of
the shorter. To compare across trial pairs of different duration, we normalized euclidean distances
by trial duration, resulting in a measure of the average euclidian distance between reference and
test trial per unit time. We then computed for each animal an area under the receiver operating
characteristic curve (auROC), and asked whether the distribution of auROCs was significantly
different from 0.5 using a two-tailed t-test.
611

Statistics

Statistical analyses were done using MATLAB and R. Unless otherwise stated, we used
mixed effects models to test for fixed effects across experiments while specifying random
intercepts per animal implemented using the R package *lme4*[53] using data from relevant single
trials. To test the significance of relevant main effects, we report Anova F-statistics with
Satterthwaite adjusted degrees of freedom for the linear mixed models and Wald chi-squared
tests for the generalized linear mixed models. We report marginal means and post hoc contrasts
(t-statistics), with Tukey correction for multiple comparisons, using the R package *emmeans*[54,55]
as (Effect Size + 95%[CI], p-value) throughout the paper. Summary of comparisons along with
additional details is included in Extended Data Table 1. All tests are two-tailed, unless otherwise
stated in the Extended Data Table 1.

Immunohistochemistry and microscopy:

Histological analysis was performed after all experiments to confirm optical fiber
placement and expression patterns of transgenes. Mice were administered with a lethal dose of
pentobarbital (Eutasil, 100 mg/kg intraperitoneally) and perfused transcardially with 4%
paraformaldehyde. The brains were removed from the skull, stored for 24 hours in 4%
paraformaldehyde, and then kept in PBS until sectioning. A vibratome or cryostat was used to
section the brain into 50 or 40 μm thick slices that were then immunostained with antibodies
against GFP (A-6455, 1:1000, Invitrogen) and tdTomato (ab125096, 1:1000, abcam). Finally, all
slices were incubated in DAPI. Images were acquired with a confocal microscope (LSM 710,
ZEISS) or a slide scanner (Axio Scan.Z1,ZEISS).
633
634

Figures

Figure 1

Figure 1. Mice learned to dynamically suppress lateralized actions.
a) Task and event diagrams. Subjects self-initiate each trial in a center nose port. After a variable delay a second tone is played and they are asked to categorize the presented interval as "short" or "long" by responding in one of two side ports. Between the two tones subjects are required to maintain position in the centre port - "fixation". **b)** Bottom: Psychometric fit to the performance of each mouse that underwent photometric recordings (n=14, light gray) and the fit to the overall average performance across mice (black). Top - Reward expectancy calculated from the overall performance of animals on a given stimuli (blue trace depicts the rectified psychometric fit at 1.5s) **c)** Animal's head speed as a function of time aligned on trial initiation for a single session of each mouse that underwent photometric recordings during trials wherein the longest interval (2.4 seconds) was delivered and a correct choice performed. Gray, mean of individual animals; black, average of all mice (n=14). Shaded region highlights period of immobility (0.6s to 2.4s post-trial initiation) **d)** Probability density functions of "broken fixations" over time during the immobility period (0.6s to 2.4s), contingent on subsequent choice at one of the side ports. See also: methods and Extended Data Figure 1. Error bars represent s.e.m. across animals.

Figure 2

Figure 2. **Medium spiny neurons of the direct and indirect pathway exhibited opposite patterns of overall activity during periods of action suppression. a)** Viral strategy used to record Ca²⁺ activity in dorsal striatum. **b)** Pattern of transgene expression in A2a-Cre (iMSN) or D1-Cre (dMSN) animals in sagittal section, ~2.1mm ML. Str-Striatum, GPe-Lateral globus pallidus, SNr-Substantia nigra pars reticulata. **c-d)** example of photometric traces in direct (c) and

indirect (d) pathway MSNs. **e,f)** Single-trial photometric data (z-scored, see methods) for all
correct trials of a single D1-cre **(e)** and A2a-cre **(d)** animal across all sessions aligned to trial
initiation (yellow arrow). Interval offset is represented as a vertical grey bar, where darker grey
represents longer intervals. Trials were further ordered within interval by reaction time. **g,h)**
Average activity (z-scored) across all animals of a given genotype (dMSN n=6, iMSN n=8).
Darkest trace represents activity during the longest interval within the interval set (2.4 seconds)
and lighter gray traces corresponding to a subset of shorter intervals. Colored segments of the
trace highlight period of immobility (0.6s to 2.4s post trial initiation) wherein average activity was
significantly different in the two pathways (0.6s to 2.4s, $p<0.05$). Error bars represent s.e.m.
across animals.

Figure 3

Figure 3. dMSNs and iMSNs exhibited opposite patterns of activity in a given hemisphere that reflected the dynamic suppression of a contra-lateral action. a) Hazard rate of broken fixations trials (see methods) wherein animals subsequently made a choice at the port corresponding to a short (green) or long (purple) choice. **b)** Schematic of labelling convention, the three large circles represent the three nose ports present in the behavioral task apparatus. The filled grey circle represents the trial initiation port, while the unfilled circles represent the two choice ports. **c)** Averaged normalized activity recorded from the hemisphere contra-lateral to a long (left panel) or short choice (right panel) port (z-scored) across all iMSN (red) and dMSN (blue) mice. Only correct completed trials were included. **d)** Average of all pairwise differences in immobility period activity of dMSNs and iMSNs between the two hemispheres, subtracting activity recorded in hemispheres contralateral to the "long" choice port from activity recorded in hemispheres contralateral to the "short" choice port (i.e., CL activity - CS activity). Error bars represent s.e.m. across animals.

    Figure 4

699
700
Figure 4. Optogenetic inhibition of iMSNs, but not dMSNs, disrupted action suppression and selection, whereas inhibition of dMSNs, but not iMSNs, slowed movement. a) Viral strategy and **b)** Protocol of optogenetic manipulation. Laser was turned on at trial onset and turned off at stimulus offset or broken fixation, whichever occurred first. **c)** Raster plot (top) and PSTH (bottom) of a single dMSN exhibiting fast and reliable inhibition, recorded from an untrained animal (see also: Extended Data Figure 5). **d)** Overall probability of breaking fixation during dMSN inhibition experiments. Colored and black dots represent data from laser-on and laser-off trials, respectively. Grey dots represent single animals. Data is shown for trials where manipulation was applied bilaterally (yellow), or unilaterally to the hemisphere either contra-lateral to the location of short choice port (CS, green) or contra-lateral to the long choice port (CL, purple). All broken

fixations were included. **e)** Cumulative distribution of choice times during dMSN inhibition
experiments (i.e. time to travel from the center port to a side port) after broken fixation trials for
manipulated (yellow) and non-manipulated (black) conditions of all animals. Dashed lines show
the median of the distributions. Inset shows the difference in medians of the two distributions for
single animals (manipulated - non-manipulated) (see also: supplementary Extended Data Figure
6). **f)** Probability of reporting at the "long choice" port after breaking fixation during dMSN inhibition
experiments. All trials wherein the mice reported a choice after breaking fixation were included.
g) Probability of breaking fixation as a function of elapsed time since trial initiation, for manipulated
trials (green and purple) and session matched non-manipulated trials (black outline) in sessions
wherein dMSNs in the hemisphere contra-lateral to a short choice (left panel) or contra-lateral to
a long choice (middle panel) (see methods). **h)** difference between dMSN manipulated and control
distributions of the probability of breaking fixation over time. Grey dots show single animals. **i-n)**
Same as c-h) but for A2a-Cre(iMSN) mice. All error bars represent s.e.m. (dMSN n=6 mice, iMSN
n=4 mice). n.s.: $p > 0.05$; *: $p < 0.05$, see Extended Data table 1.
725
726

 Extended Data Figure 1

**Extended Data Figure 1. Behavior in broken fixation trials is consistent with mice having
developed a dynamic latent motor plan. a)** Distribution of movement times, i.e. time taken for
the animal to leave the centre port and report its choice, in completed trials (black) and broken
fixation trials (green). Inset depicts the medians of movement times per animal for completed and
broken fixation trials (two-tailed paired t-test). **b)** Comparison of average head trajectories, during
a choice movement (-0.5 to 1.5 seconds relative to leaving the center port) , for sessions of a
single animal, for trials wherein animals chose the short (black) or long (gray) nose port on valid
(top, black outline) or broken fixation (bottom, green outline) trials. Thinner purple lines depict
single trials and circles represent the Long (Black), Initiation (Orange) and Short (Gray) nose
ports. (see also: Extended Data Figure 8) **c)** Probability density function of Broken fixation
occurrence as a function of time since first tone (green) and corresponding hazard rate (black full
line). Grey dashed lines represent times at which a second tone might occur. Black dashed line

743 represents the decision boundary (1.5s). **d)** Average overall performance of all animals used in
744 the photometric recordings (black) and average "pseudo-performance" calculated from broken
745 fixation trials. To calculate the performance in broken fixation trials, we binned the times at which
746 animals aborted the trial and calculated the proportion of reports at the "long choice" port over all
747 reports. All error bars represent s.e.m. across animals (n = 14).
748

Extended Data Figure 2

Extended Data Figure 2. Histological reconstruction of sites of fiber implantation for
photometry (a) and optogenetic experiments (b). Animals are colored by their genotype
according to the legend. For optogenetics experiments, the DV coordinate is shown as the
deepest position the tapered fiber was observed in histological slices. Coronal slice (AP = +0.5
from bregma) adapted from [56].

Extended Data Figure 3

759
760

Extended Data Figure 3. No significant differences in behavior were detected between genotypes. A2a-Cre and D1-Cre single animals, included in the photometry experiments, are shown in red and blue, respectively. **a-e)** Single animal psychometric curve **(a)** fits and respective parameters **(b-e)** (see Methods for further details). **f)** Overall probability of breaking fixation (all trials included). **g)** Percentage of trials wherein animals attempted to make a choice after breaking fixation (all trials included). **h)** Probability of reporting at the "long choice" port after breaking fixation contingent on whether the animal aborted before (<-1.5s) or after (>1.5s) the decision boundary. **j)** Left, Hazard of breaking fixation in time for single animals (thin curves) and overall averages within genotype (thick lines). Right, differences between the hazard of breaking fixation after and before the decision boundary. **k)** Mean velocity during the delay period from correct trials of the longest interval (2.4 seconds). Data from Figure 1c. P-values correspond to unpaired t-tests between genotypes.Error bars represent S.E.M. across animals sharing the same genotype.

Extended Data Figure 4. a) Average activity across both hemispheres during the immobility period for each mouse (data points) relative to baseline, and across animal mean (bar). Activity for dMSNs is on the left and in blue, and iMSNs on the right and in red (baseline: mean activity -5s to -2s relative to trial initiation, n=8 mice iMSN, n=6 mice dMSN, only correct trials were included). **b)** Difference between average activity after the 1.5s decision boundary, and before the 1.5s decision boundary (>1.5s - <1.5s) for activity recorded in the hemisphere contra-lateral to the side of the long choice port (CL, contra-long, left panel) and for activity recorded in the hemisphere contra-lateral to the short choice port (CS, contra-short, right panel). n.s.: p>0.05, *: p<0.05. **c)** Rate of change (derivative of ΔF/F (z)) preceding broken-fixations (dashed line) and time-matched valid trials (full line, see methods) from D1-cre (blue) or A2a-Cre (red) animals. Insets represent the difference between the overall average activity of Broken fixation and valid trials from -0.4 to -0.05 s. Grey dots depict single animals. **d)** Difference between the broken fixation and valid traces plotted in c for each genotype. Shaded areas show times wherein the activity between the rate of change of broken fixations is significantly greater (yellow) or smaller (green) than valid trials (two-tailed paired t-test) for direct (blue, top) and indirect pathway (red, bottom) recorded animals.

Extended Data Figure 5

Extended Data Figure 5. *In vivo* **ArchT activation produces fast and reversible inhibition of**
putative medium spiny neurons (pMSN), whereas ChR2 activation produces mostly

excitation but also inhibition of pMSNs. a) Identification of putative medium spiny neurons
based on firing statistics and waveform duration (see methods). Green data points indicate
pMSNs. **b)** Changes in firing rate during light delivery and baseline period versus the baseline
firing rate of all recorded units (Including non pMSN units) from A2a-Cre (red) and D1-Cre (blue)
animals. Significantly negative or positively modulated cells (see Methods) are shown as closed
and open circles, respectively. Maximum theoretical inhibition is plotted as a grey dashed line (-
ΔFR = Baseline FR). **c)** Distribution of changes in firing rate during the period of light delivery for
putatively labeled MSN units. **d)** Overall average peristimulus time histogram (PSTH) of all
negatively modulated cells, putatively labeled as MSNs, recorded from A2a-Cre (n=15 cells) and
D1-Cre (n=32 cells) mice. All units were z-scored (see methods). Error bars represent s.e.m.
across neurons. Insets show the pattern of transgene expression in A2a-Cre (iMSN) or D1-Cre
(dMSN) animals in sagittal section, ~2.1mm ML. Str-Striatum, GPe-Lateral globus pallidus, SNr-
Substantia nigra pars reticulata.
e-f) Same as b-c) but for animals expressing ChR2. **g-h)** Overall average peristimulus time
histogram (PSTH) of all positively (g) and negatively (h) modulated cells, putatively labeled as
MSNs, recorded from A2a-Cre (n=78 and 9 cells) and D1-Cre (n=112 and 29 cells) mice.

Extended Data Figure 6

Extended Data Figure 6 - Manipulation-induced changes in vigor and action selection
depended on MSN type. a) Cartoon depicts the three different manipulated trial types:
Movement Time (Delay), laser was ramped off as the second tone is played (n=6 and 5 animals,
for D1-Cre and A2a-Cre, respectively). BrokenFix Choice Time (Delay), laser was ramped off as
the animal leaves the centre port causing a broken fixation (n=6 and 5 animals). MovementTime
(Decision) laser was turned on as the second tone is played until the animal either performs its
choice or 400ms elapse , whichever occurs first (n=3 D1-Cre animals).**b)** Differences in single
animal's median choice time between inhibited and non-inhibited trials. Error bars represent
S.E.M. across animals sharing the same genotype. **c)** Differences in the choice probability relative
to the site of unilateral stimulation (Contralateral Choice) between manipulated (n = 5 D1-cre and
n=4 A2a-cre) and non-manipulated trials * p < 0.05 (see Extended Data Table 1).

 Extended Data Figure 7

Extended Data Figure 7. Single animal distributions of the effect of unilateral optogenetic inhibition on broken fixation behavior. Same data as in figures 4h and 4n but pooled of periods before (<1.5s) or after (>1.5s) the decision boundary. Single dots represent single animals, bars represent means across animals, green: inhibition performed on the hemisphere contralateral to the side of the short choice port, purple: inhibition performed on the hemisphere contralateral to the long choice port, for dMSN inhibition experiments **a)** or iMSN inhibition experiments **b)**. n.s.: p>0.05, *: p<0.05. **c)** Bias to report a contra-lateral choice after inhibition of iMSNs is not explained by the tendency of mice to make particular choices after breaking fixation early or late in the delay. Each panel, one for manipulations performed in each hemisphere, depicts the data shown in Fig. 4k further split by whether fixation was broken before or after the 1.5s decision boundary.

Extended Data Figure 8

842
843

Extended Data Figure 8. Indirect pathway inhibition did not affect movement trajectories.
a) Example trajectories for a single animal aligned to center-out for choices to the "Long port"
(red) or "Short pot" (blue) for completed trials (Blue and Green outline) and Broken fixation trials
(Orange and Purple outline). Trials are further broken down by whether the indirect pathway was
inhibited (Green and Purple) or not (Blue and Orange). b) Distribution of pairwise euclidean
distances (each trial was cropped from center-out to decision, and the pairwise euclidean distance
between two trials was calculated taking into consideration only the shortest trial) between all
conditions and the non-inhibited broken fixation condition. Inset represents the area under the
curve of the receiver operating characteristic curve (auROC) calculated between the tested
distributions and the non-inhibited broken fixation condition. c-d) Same as a-b) but for the period
between trial initiation and center out. Error bars represent S.E.M. across animals.
* $p < 0.05$ (two-tailed paired t-test).

Extended Data Figure 9

858
859
860 **Extended Data Figure 9. Both direct and indirect pathways are more active during contra-**
861 **lateral movements.** Photometry signal aligned to leaving the center-port during a near boundary
862 interval (1.74s). Same dataset as Figure 3. Error bars represent S.E.M. across animals sharing
863 the same genotype.

Extended Data Table 1

Summary of linear mixed model comparisons.

Figure	Formula (Wilkinson Notation)	Test	Reported Relevant Independent Variable or Contrast	Dependent variable	Effect size (Effect[95%CI])	Statistic	P-value (corrected)	# Mice
2g,2h	Genotype*Hemisphere + (1|Animal)	Linear mixed effects model to test whether the activity during the delay period (averaging the activity of the longest trial of 2.4seconds from 0.6 to 2.4 seconds) in the indirect pathway is greater than in the direct pathway. Additionally, if the activity of the two pathways is significantly different at trial initiation (averaging from 0 to 0.6s).						
		One-tailed Post-Hoc, Tukey correction	Genotype (iMSN > dMSN), averaged over Hemisphere	Mean ΔF/F [0.6:2.4s]	0.749[-0.065,1.56] (Averaged over hemisphere)	t(12.5) = 2.004	0.0341	8 iMSN, 6 dMSN
		One-tailed Post-Hoc, Tukey correction	Genotype (iMSN vs dMSN)	Mean ΔF/F [0:0.6]	0.102[-1.11,1.31] (Averaged over hemisphere)	t(12.1) = 0.034	0.8573	8 iMSN, 6 dMSN
3a,b	Genotype*Hemisphere + (1|Animal)	Linear mixed effects model to test difference in activity between the period before and after the decision boundary. We took the average activity in the period before the 1.5s decision boundary [0.6:1.5[and after the decision boundary [1.5,2.4], for each longest interval trial (2.4s) and used its difference as a dependent variable.						
		Mixed Anova with Satterthwaite's approximated d.f.	Genotype:Hemisphere	Mean ΔF/F		F(1, 10.953) = 22.28	<0.001	8 iMSN, 6 dMSN
		Post-Hoc, Tukey correction	iMSN:CL vs 0		0.4228[0.006, 0.840]	t(12.5)=2.949	0.0462	
		Post-Hoc, Tukey correction	dMSN:CL vs 0		-0.5354[-1.017,-0.054]	t(12.5)=-3.265	0.0272	
		Post-Hoc, Tukey correction	iMSN:CS vs 0		-0.1043[-0.521, 0.313]	t(12.5)=-0.430	0.9275	
		Post-Hoc, Tukey correction	dMSN:CS vs 0		-0.0395[-0.521,0.442]	t(12.5)=-0.212	0.9988	
4d,i	Genotype*Hemisphere*Laser+ (1|Animal)	Generalized linear mixed-model for the probability of breaking fixation of a given trial as a function of Genotype, Hemisphere and manipulation. Contrasts are reported as odds ratios.						

		Wald chi-square test	Genotype:Hemisphere:Laser	Broken Fixation ~[0,1]		$X^2_{(2)}$ = 169.5609	$<10^{-4}$	6dMSN, 4iMSN
		Post-Hoc, Tukey correction	iMSN:Bilateral:LaserOff vs iMSN:Bilateral:LaserOn		0.0585[0.0362, 0.0945]	t(asymp) = -19.361	$<10^{-4}$	4
		Post-Hoc, Tukey correction	iMSN:CL:LaserOff vs iMSN:CL:LaserOn		0.7510[0.5784, 0.9750]	t(asymp) = -3.584	0.0176	4
		Post-Hoc, Tukey correction	iMSN:CS:LaserOff vs iMSN:CS:LaserOn		0.6245[0.4809,0.8110]	t(asymp) = -5.888	$<10^{-4}$	4
		Post-Hoc, Tukey correction	dMSN:Bilateral:LaserOff vs dMSN:Bilateral:LaserOn		0.8177[0.6271, 1.0664]	t(asymp) = -2.477	0.3544	4
		Post-Hoc, Tukey correction	dMSN:Bilateral:LaserOff vs dMSN:Bilateral:LaserOn		0.9052[0.7121,1.1505	t(asymp) = -1.358	0.9712	6
		Post-Hoc, Tukey correction	dMSN:Bilateral:LaserOff vs dMSN:Bilateral:LaserOn		0.8687[0.6801, 1.1096]	t(asymp) = -1.879	0.7723	6
4g,I	Genotype*Hemisphere*Laser+ (1|Animal)	Generalized linear mixed-model. Same model and analysis as 4d,i, but splitting data by broken fixations before [0.6:1.5[or after [>1.5,2.4] decision boundary. Contrasts are reported as odds ratios.						
		Wald test	Genotype:Hemisphere:Laser	Broken Fixation ~[0,1]		$X^2_{(1)}$ = 9.9902	0.0016	6dMSN, 4iMSN
		Post-Hoc, Tukey correction	iMSN:CL:LaserOff vs iMSN:CL:LaserOn		0.409[0.283, 0.589]	t(asymp) = -7.41	$<10^{-4}$	4
	Broken fixations BEFORE the decision boundary (<1.5s)	Post-Hoc, Tukey correction	iMSN:CS:LaserOff vs iMSN:CS:LaserOn		0.904[0.608, 1.342]	t(asymp) = -0.776	0.9943	4
		Post-Hoc, Tukey correction	dMSN:CL:LaserOff vs dMSN:CL:LaserOn		1.046[0.724, 1.511]	t(asymp) = 0.370	1.0000	6
		Post-Hoc, Tukey correction	dMSN:CS:LaserOff vs dMSN:CS:LaserOn		1.054[0.719, 1.544]	t(asymp) = 0.414	0.9999	6
		Wald test	Genotype:Hemisphere:Laser	Broken Fixation ~[0,1]		$X^2_{(1)}$ = 12.9339	0.0003	6dMSN, 4iMSN
	Broken fixations AFTER the decision boundary (>1.5s)	Post-Hoc, Tukey correction	iMSN:CL:LaserOff vs iMSN:CL:LaserOn		1.060[0.748, 1.501]	t(asymp) = 0.503	0.9997	4
		Post-Hoc, Tukey correction	iMSN:CS:LaserOff vs		0.485[0.349, 0.674]	t(asymp) = -6.655	$<10^{-4}$	4

			iMSN:CS:LaserOn				
		Post-Hoc, Tukey correction	dMSN:CL:Laser Off vs dMSN:CL:Laser On		0.734[0.530, 1.018]	t(asymp) = -2.868	0.0791	6
		Post-Hoc, Tukey correction	dMSN:CS:Laser Off vs dMSN:CS:Laser On		0.739[0.533, 1.026]	t(asymp) = 2.796	0.0958	6
4f,k	Genotype*Hemisphere*Laser+ (1|Animal)	Generalized linear mixed-model testing the probability of reporting a long choice after breaking fixation, as a function of Genotype, Hemisphere and manipulation. Contrasts are reported as odds ratios.						
		Wald test	Genotype:Hemisphere:Laser	Choice "long" ~[0,1]		$X^2_{(1)}$ = 40.4507	$<10^{-4}$	6dMSN, 4iMSN
		Post-Hoc, Tukey correction	iMSN:CL:LaserOff vs iMSN:CL:LaserOn		0.257[0.1430, 0.463]	t(asymp) = -7.003	$<10^{-4}$	4
		Post-Hoc, Tukey correction	iMSN:CS:LaserOff vs iMSN:CS:LaserOn		3.369[1.7069, 6.651]	t(asymp) = 5.414	$<10^{-4}$	4
		Post-Hoc, Tukey correction	dMSN:CL:Laser Off vs dMSN:CL:Laser On		0.917[0.4484, 1.875]	t(asymp) = -0.368	1.000	6
		Post-Hoc, Tukey correction	dMSN:CS:Laser Off vs dMSN:CS:Laser On		0.737[0.3782, 1.438]	t(asymp) = -1.382	0.8659	6
Fig.4e,j	Genotype*Laser + (1|Animal)	Linear mixed effects model to test the differences in the time animals take to report their choice at one of the side ports after breaking fixation, as a function of genotype and manipulation.						
		Mixed Anova with Satterthwaite's approximated d.f.	Genotype:Laser	BrokenFixation Mov. Time		F(1,2454.96) = 4.8377	0.02794	5 iMSN, 6 dMSN
		Post-Hoc, Tukey correction	iMSN:LaserOff vs iMSN:LaserOn		7.82 [-482, 497.3]	t(2454.77) = 0.041	1.0000	5
		Post-Hoc, Tukey correction	dMSN:LaserOff vs dMSN:LaserOn		-630.3[-1193,97.6]	t(2455.11) = -2.880	0.0209	6
Ext. Data Fig.6	Genotype*Laser + (1|Animal)	Linear mixed effects model to test the differences in the time animals take to report their choice at one of the side ports after successfully completing a trial between control and manipulated trials during the delay period						
	Movement Time(Delay)	Mixed Anova with Satterthwaite's approximated d.f.	Genotype:Laser	Mov. Time		F(1,14272.7) = 2.481	0.115	5 iMSN, 6 dMSN
		Post-Hoc, Tukey correction	iMSN:LaserOff vs iMSN:LaserOn		14.0 [-12.5, 40.5]	t(14273.21) = 1.353	0.529	5

		Post-Hoc, Tukey correction	dMSN:LaserOff vs dMSN:LaserOn		-6.3[-- 26.0, 13.4]	t(14271.67) = -0.820	0.845	6
Ext. Data Fig.6	Laser + (1|Animal)	Linear mixed effects model to test the differences in the time animals take to report their choice at one of the side ports after successfully completing a trial between control and manipulated trials during the decision epoch. (this experiment was only performed in D1-Cre Animals)						
	Movement Time(Decisio n)	Mixed Anova with Satterthwaite's approximated d.f.	Laser	Mov. Time		F(5194.0) = 2.481	0.70	3 dMSN
		Post-Hoc, Tukey correction	LaserOff vs LaserOn		-2.8 [- 16.9, 11.3]	t(14273.21) = -0.390	0.70	3 dMSN

867
868

References

1.   Albin, R. L., Young, A. B. & Penney, J. B. The functional anatomy of basal ganglia
disorders. *Trends Neurosci.* **12,** 366–375 (1989).
2.   Mink, J. W. The basal ganglia: focused selection and inhibition of competing motor
programs. *Prog. Neurobiol.* **50,** 381–425 (1996).
3.   Cui, G., Jun, S. B., Jin, X., Pham, M. D., Vogel, S. S., Lovinger, D. M. & Costa, R. M.
Concurrent activation of striatal direct and indirect pathways during action initiation. *Nature*
494, 238–242 (2013).
4.   Schultz, W. The Primate Basal Ganglia Between the Intention and Outcome of Action.
Functions of the Cortico-Basal Ganglia Loop 31–48 (1995). doi:10.1007/978-4-431-68547-
0_3
5.   Doya, K. What are the computations of the cerebellum, the basal ganglia and the cerebral
cortex? *Neural Netw.* **12,** 961–974 (1999).
6.   Barkley, R. A. Behavioral inhibition, sustained attention, and executive functions:
constructing a unifying theory of ADHD. *Psychol. Bull.* **121,** 65–94 (1997).
7.   Gerfen, C. R. & Surmeier, D. J. Modulation of striatal projection systems by dopamine.
Annu. Rev. Neurosci. **34,** 441–466 (2011).
8.   Alexander, G. E. & Crutcher, M. D. Functional architecture of basal ganglia circuits: neural
substrates of parallel processing. *Trends Neurosci.* **13,** 266–271 (1990).
9.   Smith, Y., Bevan, M. D., Shink, E. & Bolam, J. P. Microcircuitry of the direct and indirect
pathways of the basal ganglia. *Neuroscience* **86,** 353–387 (1998).
10.   Deniau, J. M. & Chevalier, G. Disinhibition as a basic process in the expression of striatal
functions. II. The striato-nigral influence on thalamocortical cells of the ventromedial
thalamic nucleus. *Brain Res.* **334,** 227–233 (1985).
11.   Kravitz, A. V., Freeze, B. S., Parker, P. R. L., Kay, K., Thwin, M. T., Deisseroth, K. &
Kreitzer, A. C. Regulation of parkinsonian motor behaviours by optogenetic control of basal
ganglia circuitry. *Nature* **466,** 622–626 (2010).
12.   Freeze, B. S., Kravitz, A. V., Hammack, N., Berke, J. D. & Kreitzer, A. C. Control of basal
ganglia output by direct and indirect pathway projection neurons. *J. Neurosci.* **33,** 18531–
18539 (2013).
13.   Tecuapetla, F., Jin, X., Lima, S. Q. & Costa, R. M. Complementary Contributions of Striatal
Projection Pathways to Action Initiation and Execution. *Cell* **166,** 703–715 (2016).
14.   Sippy, T., Lapray, D., Crochet, S. & Petersen, C. C. H. Cell-Type-Specific Sensorimotor
Processing in Striatal Projection Neurons during Goal-Directed Behavior. *Neuron* **88,** 298–
305 (2015).
15.   Kravitz, A. V., Tye, L. D. & Kreitzer, A. C. Distinct roles for direct and indirect pathway
striatal neurons in reinforcement. *Nat. Neurosci.* **15,** 816–818 (2012).
16.   Yttri, E. A. & Dudman, J. T. Opponent and bidirectional control of movement velocity in the
basal ganglia. *Nature* **533,** 402–406 (2016).
17.   Tai, L.-H., Lee, A. M., Benavidez, N., Bonci, A. & Wilbrecht, L. Transient stimulation of
distinct subpopulations of striatal neurons mimics changes in action value. *Nat. Neurosci.*
15, 1281–1289 (2012).
18.   Tecuapetla, F., Matias, S., Dugue, G. P., Mainen, Z. F. & Costa, R. M. Balanced activity in
basal ganglia projection pathways is critical for contraversive movements. *Nat. Commun.* **5,**
4315 (2014).
19.   Barbera, G., Liang, B., Zhang, L., Gerfen, C. R., Culurciello, E., Chen, R., Li, Y. & Lin, D.-T.
Spatially Compact Neural Clusters in the Dorsal Striatum Encode Locomotion Relevant

Information. *Neuron* **92,** 202–213 (2016).

20. Markowitz, J. E., Gillis, W. F., Beron, C. C., Neufeld, S. Q., Robertson, K., Bhagat, N. D.,
Peterson, R. E., Peterson, E., Hyun, M., Linderman, S. W., Sabatini, B. L. & Datta, S. R.
The Striatum Organizes 3D Behavior via Moment-to-Moment Action Selection. *Cell* **174,**
44–58.e17 (2018).

21. Denny-Brown, D. & Yanagisawa, N. The role of the basal ganglia in the initiation of
movement. *Res. Publ. Assoc. Res. Nerv. Ment. Dis.* **55,** 115–149 (1976).

22. Redgrave, P., Prescott, T. J. & Gurney, K. The basal ganglia: a vertebrate solution to the
selection problem? *Neuroscience* **89,** 1009–1023 (1999).

23. Gouvêa, T. S., Monteiro, T., Motiwala, A., Soares, S., Machens, C. & Paton, J. J. Striatal
dynamics explain duration judgments. *Elife* **4,** (2015).

24. Soares, S., Atallah, B. V. & Paton, J. J. Midbrain dopamine neurons control judgment of
time. *Science* **354,** 1273–1277 (2016).

25. Chen, T.-W., Wardill, T. J., Sun, Y., Pulver, S. R., Renninger, S. L., Baohan, A., Schreiter,
E. R., Kerr, R. A., Orger, M. B., Jayaraman, V., Looger, L. L., Svoboda, K. & Kim, D. S.
Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* **499,** 295–300
(2013).

26. Matias, S., Lottem, E., Dugué, G. P. & Mainen, Z. F. Activity patterns of serotonin neurons
underlying cognitive flexibility. *Elife* **6,** (2017).

27. Han, X., Chow, B. Y., Zhou, H., Klapoetke, N. C., Chuong, A., Rajimehr, R., Yang, A.,
Baratta, M. V., Winkle, J., Desimone, R. & Boyden, E. S. A high-light sensitivity optical
neural silencer: development and application to optogenetic control of non-human primate
cortex. *Front. Syst. Neurosci.* **5,** 18 (2011).

28. Pisanello, F., Mandelbaum, G., Pisanello, M., Oldenburg, I. A., Sileo, L., Markowitz, J. E.,
Peterson, R. E., Della Patria, A., Haynes, T. M., Emara, M. S., Spagnolo, B., Datta, S. R.,
De Vittorio, M. & Sabatini, B. L. Dynamic illumination of spatially restricted or large brain
volumes via a single tapered optical fiber. *Nat. Neurosci.* **20,** 1180–1188 (2017).

29. Turner, R. S. & Desmurget, M. Basal ganglia contributions to motor control: a vigorous
tutor. *Curr. Opin. Neurobiol.* **20,** 704–716 (2010).

30. Majid, D. S. A., Cai, W., Corey-Bloom, J. & Aron, A. R. Proactive selective response
suppression is implemented via the basal ganglia. *Journal of Neuroscience* **33,** 13259–
13269 (2013).

31. Watanabe, M. & Munoz, D. P. Presetting basal ganglia for volitional actions. *J. Neurosci.*
30, 10144–10157 (2010).

32. Ford, K. A. & Everling, S. Neural activity in primate caudate nucleus associated with pro-
and antisaccades. *J. Neurophysiol.* **102,** 2334–2341 (2009).

33. Amita, H. & Hikosaka, O. Indirect pathway from caudate tail mediates rejection of bad
objects in periphery. *Sci Adv* **5,** eaaw9297 (2019).

34. Panigrahi, B., Martin, K. A., Li, Y., Graves, A. R., Vollmer, A., Olson, L., Mensh, B. D.,
Karpova, A. Y. & Dudman, J. T. Dopamine Is Required for the Neural Representation and
Control of Movement Vigor. *Cell* **162,** 1418–1430 (2015).

35. Hintiryan, H., Foster, N. N., Bowman, I., Bay, M., Song, M. Y., Gou, L., Yamashita, S.,
Bienkowski, M. S., Zingg, B., Zhu, M., Yang, X. W., Shih, J. C., Toga, A. W. & Dong, H.-W.
The mouse cortico-striatal projectome. *Nat. Neurosci.* **19,** 1100–1114 (2016).

36. Coddington, L. T. & Dudman, J. T. The timing of action determines reward prediction
signals in identified midbrain dopamine neurons. *Nat. Neurosci.* **21,** 1563–1573 (2018).

37. Parker, J. G., Marshall, J. D., Ahanonu, B., Wu, Y.-W., Kim, T. H., Grewe, B. F., Zhang, Y.,
Li, J. Z., Ding, J. B., Ehlers, M. D. & Schnitzer, M. J. Diametric neural ensemble dynamics
in parkinsonian and dyskinetic states. *Nature* **557,** 177–182 (2018).

38. Park, J., Coddington, L. T. & Dudman, J. T. Basal Ganglia Circuits for Action Specification.
Annu. Rev. Neurosci. **43,** (2020).

39. Wall, N. R., De La Parra, M., Callaway, E. M. & Kreitzer, A. C. Differential innervation of
direct- and indirect-pathway striatal projection neurons. *Neuron* **79,** 347–360 (2013).
40. Hunnicutt, B. J., Jongbloets, B. C., Birdsong, W. T., Gertz, K. J., Zhong, H. & Mao, T. A
comprehensive excitatory input map of the striatum reveals novel functional organization.
Elife **5,** (2016).
41. Lau, B. & Glimcher, P. W. Action and outcome encoding in the primate caudate nucleus. *J.*
Neurosci. **27,** 14502–14514 (2007).
42. Rueda-Orozco, P. E. & Robbe, D. The striatum multiplexes contextual and kinematic
information to constrain motor habits execution. *Nat. Neurosci.* **18,** 453–460 (2015).
43. Klaus, A., Martins, G. J., Paixao, V. B., Zhou, P., Paninski, L. & Costa, R. M. The
Spatiotemporal Organization of the Striatum Encodes Action Space. *Neuron* **96,** 949
(2017).
44. Schwarting, R. K. & Huston, J. P. The unilateral 6-hydroxydopamine lesion model in
behavioral brain research. Analysis of functional deficits, recovery and treatments. *Prog.*
Neurobiol. **50,** 275–331 (1996).
45. Gerfen, C. R., Paletzki, R. & Heintz, N. GENSAT BAC cre-recombinase driver lines to study
the functional organization of cerebral cortical and basal ganglia circuits. *Neuron* **80,** 1368–
1383 (2013).
46. Lopes, G., Bonacchi, N., Frazão, J., Neto, J. P., Atallah, B. V., Soares, S., Moreira, L.,
Matias, S., Itskov, P. M., Correia, P. A., Medina, R. E., Calcaterra, L., Dreosti, E., Paton, J.
J. & Kampff, A. R. Bonsai: an event-based framework for processing and controlling data
streams. *Front. Neuroinform.* **9,** 7 (2015).
47. Siegle, J. H., López, A. C., Patel, Y. A., Abramov, K., Ohayon, S. & Voigts, J. Open Ephys:
an open-source, plugin-based platform for multichannel electrophysiology. *J. Neural Eng.*
14, 045003 (2017).
48. Madisen, L., Mao, T., Koch, H., Zhuo, J.-M., Berenyi, A., Fujisawa, S., Hsu, Y.-W. A.,
Garcia, A. J., 3rd, Gu, X., Zanella, S., Kidney, J., Gu, H., Mao, Y., Hooks, B. M., Boyden, E.
S., Buzsáki, G., Ramirez, J. M., Jones, A. R., Svoboda, K., Han, X., Turner, E. E. & Zeng,
H. A toolbox of Cre-dependent optogenetic transgenic mice for light-induced activation and
silencing. *Nat. Neurosci.* **15,** 793–802 (2012).
49. Benhamou, L., Kehat, O. & Cohen, D. Firing pattern characteristics of tonically active
neurons in rat striatum: context dependent or species divergent? *J. Neurosci.* **34,** 2299–
2304 (2014).
50. Yael, D., Zeef, D. H., Sand, D., Moran, A., Katz, D. B., Cohen, D., Temel, Y. & Bar-Gad, I.
Haloperidol-induced changes in neuronal activity in the striatum of the freely moving rat.
Front. Syst. Neurosci. **7,** 110 (2013).
51. Chuong, A. S., Miri, M. L., Busskamp, V., Matthews, G. A. C., Acker, L. C., Sørensen, A. T.,
Young, A., Klapoetke, N. C., Henninger, M. A., Kodandaramaiah, S. B., Ogawa, M.,
Ramanlal, S. B., Bandler, R. C., Allen, B. D., Forest, C. R., Chow, B. Y., Han, X., Lin, Y.,
Tye, K. M., Roska, B., Cardin, J. A. & Boyden, E. S. Noninvasive optical inhibition with a
red-shifted microbial rhodopsin. *Nat. Neurosci.* **17,** 1123–1129 (2014).
52. Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W. & Bethge, M.
DeepLabCut: markerless pose estimation of user-defined body parts with deep learning.
Nat. Neurosci. **21,** 1281–1289 (2018).
53. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software, Articles* **67,** 1–48 (2015).
54. Searle, S. R., Speed, F. M. & Milliken, G. A. Population Marginal Means in the Linear
Model: An Alternative to Least Squares Means. *Am. Stat.* **34,** 216–221 (1980).
55. Lenth, R. Least-Squares Means: The R Package lsmeans. *Journal of Statistical Software,*
Articles **69,** 1–33 (2016).
56. Franklin, K. B. J. & Paxinos, G. The mouse brain in stereotaxic coordinates 3rd edn. (2008).