Lateral Hypothalamic GABAergic neurons encode and potentiate sucrose's palatability

Aketzali Garcia, Alam Coss, Jorge Luis-Islas, Liliana Puron-Sierra, Monica Luna, Miguel Villavicencio, Ranier Gutierrez[*]

Laboratory of Neurobiology of Appetite; Department of Pharmacology, CINVESTAV, 07360. Mexico City, Mexico.

[*] Correspondence:
Ranier Gutierrez
ranier@cinvestav.mx

ORCID: https://orcid.org/0000-0002-9688-0289

Keywords: LHA[Vgat+] neurons, feeding circuits, palatability, sucrose

This PDF file includes:

Number of pages: 56

Number of figures: 12

Abstract

Sucrose is attractive to most species in the animal kingdom, not only because it induces a sweet taste sensation, but also for its positive palatability or likability. Although palatability is such an important sensory attribute, it is currently unknown which cell-types encode and modulate sucrose's palatability. Studies in mice have shown that GABAergic LHA$^{\text{Vgat+}}$ neurons evoke voracious eating; however, it is not known whether these neurons would be driving consumption by increasing palatability. Using optrode recordings, we measured sucrose's palatability while VGAT-ChR2 transgenic mice performed a brief access sucrose test. We found a subpopulation of LHA$^{\text{Vgat+}}$ neurons that their activity correlated with the increase in licking responses evoked by sucrose concentrations. Optogenetic gain of function experiments where mice were able to choose among available water, 3%, or 18% sucrose solution, uncovered that opto-stimulation of LHA$^{\text{Vgat+}}$ neurons consistently promoted higher intake of the most palatable stimuli (18% sucrose). In contrast, if they self-stimulated near the less palatable stimulus, some VGAT-ChR2 mice preferred water over 18% sucrose. Unexpectedly, activation of LHA$^{\text{Vgat+}}$ neurons increased quinine intake but only during water deprivation, since in sated animals, they failed to promote quinine intake or to tolerate an aversive stimulus. Conversely, these neurons promoted overconsumption of sucrose when it was the nearest stimuli. We conclude that a subpopulation of LHA$^{\text{Vgat+}}$ neurons increases tastant's intake by potentiating its palatability.

Introduction

The Lateral Hypothalamic Area (LHA) has been regarded as the "feeding center" since its lesion results in hypophagia and subsequent death [1,2]. It is part of a neural circuit related to feeding and reward [3,4] as rats are willing to press a lever to deliver electrical intra-cranial self-stimulation (ICSs), and if food is available, it also promotes feeding [3,5–7]. Moreover, if a sweet tastant is available, the rate of electrical ICSs is further increased, whereas bitter compounds decreased ICSs rates [8,9], suggesting an interaction between ICSs and taste palatability. In this regard, and because of its connections with different cortical and subcortical gustatory regions [10–13], the LHA is anatomically located to processes and broadcast taste palatability information [14,15]. Pioneering electrophysiological studies have recorded gustatory-evoked responses in the LHA [16,17]. One recent study uncovered two functional populations of LHA neurons: one that was activated by palatable tastants, e.g., sucrose, and the other by aversive tastants, like quinine [18]. However, the genetic identity of the LHA cell-type(s) involved in processing palatability-related information remains elusive, and to unveil their identity is a goal of this study.

The LHA is currently viewed as a hub of various cell-types [19] grossly divided into two larger populations related to feeding: the glutamatergic ($LHA^{Vglut2+}$) and GABAergic neurons (LHA^{Vgat+}) [20]. Activation of $LHA^{Vglut2+}$ neurons leads to reduced food intake and is aversive [21]. In contrast, stimulation of LHA^{Vgat+} neurons is rewarding and produces voracious eating of both foods with nutritional value [22,23] and those without calories or even gnawing behavior towards an irrelevant biological stimulus [23]. Moreover, LHA^{Vgat+} neurons enhance the salience of the nearest stimuli and induce reward via stimulation of their projections to the ventral tegmental area (VTA) [24]. In addition, the evoked feeding is mediated by modulation of terminals reaching a region adjacent to the locus coeruleus [25]. On the contrary, inhibition of GABAergic LHA^{Vgat+} cell somas is aversive and stops

feeding [22,23]. However, the role of LHA^{Vgat+} neurons in processing sucrose's palatability remains to

be determined.

This study identified a new role of LHA^{Vgat+} neurons in encoding and enhancing sucrose's palatability.

We found that a subpopulation of LHA GABAergic neurons increases the consumption of the most

palatable and proximal tastant. They accomplish this by potentiating the palatability of nearby

gustatory stimuli. For aversive stimuli, the effect of these neurons is different. In a single bottle test,

we found that water deprivation increased the tolerance for bitter compounds and gated a window of

opportunity within which the activation of GABAergic neurons is sufficient to reassign the negative

hedonic value of quinine temporarily and promotes its intake. Nevertheless, in a three-option

preference test, these neurons failed to develop a preference to quinine when sucrose was available.

Moreover, optogenetic activation of LHA^{Vgat+} neurons evoked many hallmark behaviors resembling

those seen in LHA electrical stimulation. In this regard, and consistent with their positive role in

palatability, our results could explain why electrical ICSs are further facilitated by the presence of

sweet solutions [8,9]. We also found that LHA^{Vgat+} neurons are the common neural substrate for both

reward and feeding since after repeated self-stimulation, the more the VGAT-ChR2 mice opto-self-

stimulated, the stronger the laser-induced licking they exhibited. We conclude that a subpopulation of

GABAergic LHA^{Vgat+} neurons combines stimulus-proximity and palatability-related information to

enhance nearby energy-rich foods' palatability.

Materials and methods

Animal subjects

We used 36 VGAT-ChR2 mice from both sexes and 8-16 weeks old (number ID 014548; Jackson Laboratory, Sacramento, CA, USA). 21 wildtype (WT) littermates were used as a control, from both sexes. Mice were individually housed in laboratory home cages and maintained in a temperature-controlled ($22 \pm 1° C$) room with 12:12 h light-dark cycle, and unless otherwise mentioned, Chow food (PicoLab Rodent Diet 20, MO, USA) and water was available *ad libitum*. For experiments with water restriction, after behavioral testing, mice were allowed to drink water for 1 h daily. All procedures were performed with the approval of the CINVESTAV Animal Care and Use Committee. Experiments were conducted at the same time of the day from 11:00 AM to 2:00 PM.

Surgical procedures

All mice were anesthetized with ketamine (100 mg/kg, i.p.) and xylazine (8 mg/kg, i.p.), and placed into a stereotaxic apparatus (Stoelting, IL, USA). Lidocaine (0.05 mL) was applied subcutaneously under the skin's head as an analgesic, and ophthalmic ointment (hydrocortisone, neomycin, and polymyxin-B) was used periodically to maintain eye lubrication. The antibiotic enrofloxacin (0.4 mL/kg) was injected for 3 days as a post-surgery recovery.

For experiments with opto-stimulation of LHA^{Vgat+} cell somas, a single multimode optical fiber 200 μm core diameter with a 0.39 NA (FT200UMT; Thorlabs, NJ, USA) was implanted unilaterally targeting the LHA using the following coordinates: AP: -1.25 mm, ML: ± 1 mm, DV: -4.9 mm, from bregma. For electrophysiology recordings, a custom-made optrode was unilaterally implanted

counterbalanced for both hemispheres (AP: -1.25 mm, ML: ± 1 mm, from bregma; DV: -5.2 mm

ventral to dura). The optrode comprises an array of 16 tungsten wires formvar coated (35 µm diameter)

surrounding and protruding 1 mm the single multimode optical fiber tip. No significant differences

were found between hemispheres, so data were pooled together (data not shown). To allow for

recovery, all experiments began seven days after surgery.

Histology and Immunofluorescence

To verify optrode and fiber optic location, at the end of the experiments, mice were sacrificed by an

overdose of pentobarbital (150 mg/kg) and perfused with PBS, followed by 4% paraformaldehyde.

Brains were fixed overnight in 4% paraformaldehyde and gradually replaced in a gradient of

concentrations until reaching 30% sucrose. For histology, brain slides (40 µm) were cut with a cryostat

(Thermo Scientific HM525 NX) and images were taken with a Nikon eclipse e200 microscope and a

Progress Gryphax microscope camera, using a 10x objective.

Free-floating brain slides (40 µm) were blocked with 1% BSA, 0.2% Triton in PBS for 30 min. Then,

they were washed with 0.2% Triton in PBS three times every 10 min, followed by the incubation with

the primary antibodies: mouse anti-GAD 67 primary antibody (Millipore, Mab5406, 1:1000 dilution)

and rabbit anti-GFP primary antibody (Invitrogen, A11122, 1:1000 dilution). Incubation took place

overnight at 4°C. The next day, brain slides were washed with 0.2% Triton in PBS three times every

10 min, then incubated with the secondary antibodies Alexa 647 goat anti-mouse (Invitrogen, A21235,

1:500 dilution) and Alexa 488 donkey anti-rabbit (Invitrogen, A21206, 1:500 dilution) during 90 min

at room temperature. Afterward, we applied DAPI to stain the nuclei. Brain slides were put on slides

with a mounting medium for fluorescence (Vectashield), and images were taken with a Leica confocal

microscope using a 63x objective.

Gustatory stimuli

Sucrose reagent-grade chemical quality (Sigma-Aldrich, Mexico City, Mexico) was used in the

following concentrations: no sucrose, 3, 10, and 18 wt%. Quinine hydrochloride dihydrate (QHCl)

reagent-grade was used at 0.004 wt% (Sigma-Aldrich, Mexico City, Mexico). Artificial Saliva

contained (mM): 4 NaCl, 10 KCl, 6 NaHCO$_3$, 6 KHCO$_3$, 0.5 CaCl$_2$, 0.5 MgCl$_2$, 0.24 K$_2$HPO$_4$, 0.24

KH$_2$PO$_4$ [26]. We added 0.05 mM HCl to adjust to pH 7. The solutions were dissolved in distilled

water, maintained under refrigeration, and used at room temperature. We also used 20 mg chocolate

pellets (Bio-Serv, NJ, USA), Chow food (PicoLab Rodent Diet 20, MO, USA), a high-fat diet with

45% kcal% fat (Research Diets, NJ, USA), granulated sugar cube, and cork.

Optogenetic stimulation

A 473 nm laser intensity was modulated by a DPSS system (OEM laser, UT, USA). Laser pulses were

synchronized with behavioral experiments with Med Associates Inc. software and TTL signal

generator (Med Associates Inc., VT, USA). The optical power at the patch cord was 15 mW, and it

was measured with an optical power meter (PM20A, Thorlabs, NJ, USA). This delivers between 10 to

12.6 mW at the tip of the fiber optic, depending on each fiber's efficiency. To explore the best

stimulation parameters for LHA GABAergic neurons, the laser was turned "on" at different frequencies

2.5, 5, 10, 20, and 50 Hz semi-randomly, with a 10 ms pulse width. For the other experiments, the laser

was fixed at 50 Hz. The laser was 2 s on and 4 s off, with 10 ms pulse width and a duty cycle of 50%.

Behavioral Assays

Pre-stimulation task

Mice were placed in an operant conditioning chamber with a central port, a door blocked the access to

the sipper during the first 15 min. In this pre-stimulation period, each mouse was opto-stimulated with

all pre-stimulation parameters following a Latin square design (described in Table 1). Then, the door

was opened and a 10% sucrose solution was available during the next 15 min. During the pre-

stimulation period, the laser was turned on for 2 s and off for 4 s.

Table 1. Pre-stimulation protocol schemes:

Laser on	Sucrose access
0 min	15 min
5 min	15 min
10 min	15 min
15 min	15 min

Open loop stimulation task

Mice were opto-stimulated by alternating 5 min off and 5 min on blocks across the session [24]. During

opto-stimulated blocks, the laser was turned "on" regardless of the mice's behavior and spatial position

in the chamber. The front panel of an operant conditioning chamber (Med Associates Inc, VT, USA)

was equipped with one or three ports, each with a sipper where individual licks were registered by a

contact lickometer (Med Associates Inc, VT, USA). To determine the best stimulation parameters to

induced feeding behavior, a group of naïve mice was tested with one single port, and in each session a

different opto-stimulation frequency (0, 2.5, 5, 10, 20, and 50 Hz) was delivered daily. Mice had free

access to a single sucrose 10% solution in the central port.

To determine whether opto-stimulation of LHA GABAergic neurons bias the intake towards the most

palatable stimuli available. A different group of mice was tested in an open loop stimulation task, in a

box equipped with three ports, where mice had free access to water (central sipper), a 3% sucrose

solution (left sipper), and 18% sucrose solution (right sipper). Lateral sippers were counterbalanced

across animals. This task comprised of 4 baseline days (with no opto-stimulation, data not shown) and

11 test days.

For the open loop stimulation sessions, with three ports, we video tracked the mouse's distance and

head angle relative to each of the three sippers. The sippers' position and three anatomical animal's

points, the nose, the neck, and the tail's base, were extracted using the DeepLabCut Python package

[27,28]. We define the mouse position as the coordinates (x and y) for the nose, whereas the head

direction was a 2D vector going from the neck to the nose. For each video frame, the relative distance

to each sipper was calculated (in pixels), and the relative angle between the head direction and position

of each sipper (in degrees). An angle of 0º means the mouse is looking at the sipper, and 180º means

facing in the opposite direction. For distance measuring, 100 pixels correspond to approximately 6 cm.

The mouse position and direction were extracted from the train of stimulation's first laser pulse, and

the following 6 s window was analyzed to find out if the mice initiated a licking behavior at any of the

3 sippers. Then, we built a two-dimensional array containing the probability of licking any sipper given

the mouse's distance and angle during the first laser of each opto-stimulation train.

Closed-loop stimulation task

The next eleven sessions, animals were tested in the same operant conditioning chamber described above, but this time opto-self-stimulation was delivered when the subject made a head entry in the central port (containing a sipper filled with water). The laser was turned on by 2 s with a time out of 4 s post-stimulation. Lateral ports delivered sucrose 3 and 18%. After this and on subsequent days, the central port stimulus was replaced across sessions following the protocols described below.

Table 2. Closed-loop stimulation protocol

Stimuli in central port	Sessions tested
Water	Last 5 sessions of the closed-loop (Figure 7)
Extinction (No opto-stimulation)	4
Water	3
Empty sipper	3
Water	3
0.004% Quinine	4
Airpuff	4
Unavoidable airpuff	3
Sucrose 18% (Water was placed in the sucrose 18% port)	4

Closed-loop stimulation task (one single bottle test: quinine)

Mice were put in a custom-made box with 18 x 13 x 7.5 cm internal dimensions. The front wall was equipped with one single sipper (a contact lickometer with V-shape registered individual licks). 0.004% quinine or 18% sucrose were tested with sated and water-deprived mice during 3 and 2 sessions, respectively. Opto-self-stimulation (2 s on with a time out of 4 s) was delivered when mice made a head entry in the central port. Water-deprived mice had access to *ad libitum* water for 1 h after the session finished.

Open loop stimulation in an open field with Chow, high-fat diet, or granulated sugar cube

Animals were placed in the circular arena's center (50 cm of diameter). Three or two food plates were

equidistant to each other, and each contained either Chow, high-fat diet, or granulated sugar cube. Mice

were opto-stimulated by alternating 5 min block with no-laser and 5 min laser block across the session.

In open loop, the laser was turned "on" regardless of the mice's behavior and spatial position in the

arena. All sessions were recorded and analyzed with bonsai software (https://open-ephys.org/bonsai).

We calculated the mouse' centroid at each frame and used that information to create a heatmap of its

position. The distance of the mouse's centroid from each stimulus was computed. An interaction

occurred when the distance was below 60 mm, and the animal stayed there for at least 1 s. Food plates

were weighted at the beginning and the end of each session.

Closed-loop in an open-field

In a circular arena, four stimuli were located equidistant to each other, either wood cork, Chow,

chocolate pellets, or a sipper filled with a 10% sucrose solution. During the session, a homemade

computer vision program tracked in real-time the position of the subject. Each mouse needs to enter

the designated area to receive opto-self-stimulation (2 s laser on and 4 s off). Mice need to leave and

re-enter the designated area to begin a new trial.

Brief access test

To evaluate sucrose's palatability, we used a brief access test in water-deprived mice. For this task, we

used a licking spout consisting of independent stainless-steel needles (20-gauge) cemented together

with dental acrylic at the sipper tube's tip [29]. Each needle was connected to a solenoid valve (Parker,

Ohio, USA) by a silicon tube. The drop (~2 μL) was calibrated before each session and maintained by

using a pressurized control system. The sequence of a trial is as follows. At the beginning of the task,

the sipper is extended in a lick position. To start a new trial, the subject had to introduce its head into

the central port to turn on the laser, then elicited a dry lick to start the reward epoch (7s). During this

period, a tastant's drop was delivered every lick. Once the epoch was finished, the sipper was retracted

for 3 s as inter-trial interval, and then the sipper was re-extended into the central port. Tastant solutions

(water, sucrose 3%, and sucrose 18%) were delivered in a semi-random order. The laser was turned on

using a closed-loop protocol. Briefly, each head entry triggers the laser on for 2s (or 7s). The time of

stimulation changed between sessions, either 2 s or 7s, in order to test whether licking could be evoked

for a more extended period. At the end of the session, mice had access to *ad libitum* water for 1 h.

Protocols are described below:

Table 3. Brief access taste protocol

Laser on	Time on	# Sessions
All trials	2 s	6
Water trials	2 s	3
Sucrose 3% trials	2 s	3
Sucrose 18% trials	2 s	3
All trials	7 s	4
Water trials	7 s	3
Sucrose 3% trials	7 s	3
Sucrose 18% trials	7 s	3
Mock laser (sucrose 3% trials) *	7 s	3

*For mock laser, mice were connected to a false fiber optic, whereas the real laser was glued outside

the skull to emit blue light.

In addition, in the same behavioral box, a brief access test was performed while the LHA was recorded,

followed by optogenetic tagging in the same session. Tastant solutions (artificial saliva, water, sucrose

3%, and 18 %) were delivered in a semi-random order. We operationally defined sucrose's palatability

as the enhancement of hedonically positive oromotor responses evoked by increasing sucrose

concentrations [30,31].

Electrophysiology

Neural activity was recorded using a Multichannel Acquisition Processor System (Plexon, Dallas TX,

USA) interfaced with Med Associates to record behavioral events simultaneously [32]. Extracellular

voltage signals were first amplified by an analog head-stage (Plexon, HST/16o25-GEN2-18P-2GP-

G1), then amplified and sampled at 40 kHz. Raw signals were band-pass filtered from 154 Hz to 8.8

kHz and digitalized at 12 bits resolution. Only single neurons with action potentials with a signal to

noise ratio of 3:1 were analyzed [32]. Action potentials were isolated online using a voltage-time

threshold window and three principals' components contour templates algorithm. Furthermore, off-

line spike sorting was performed (Plexon, offline sorter), and only single units with stable waveforms

across the entire session were included in the analyses [32].

Neuronal classification

Laser activated neurons (pLHA^{Vgat}) were detected by using two methods: 1) a Kruskal-Wallis to

compare firing rates during a baseline (-1 to -0.04 s), against the activity during the presentation of

laser (0 – 2 s, aligned to laser onset). 2) Cumsum statistic [33] to obtain the first bin (1 ms resolution)

after the first laser pulse that significantly increased the firing rate above baseline (from -20 to 0 ms).

Only neurons showing a significant value in both tests were considered laser-activated pLHA^{Vgat}.

We used a Kruskal-Wallis test to compare firing rates during the baseline (-1 to -0.04 s) period, against

the activity during the presentation of laser (0 to 2 s) aligned to laser onset for all frequency tested.

Neurons that significantly increased their firing rate during the laser stimulation were named

"Activated," and neurons that decreased their activity were named "Inhibited" or non-LHA^{Vgat} neurons.

Neurons with no significant modulation were classified as "unmodulated."

Optotagging stimulation protocol

To identify LHA^{Vgat+} neurons related to sucrose's palatability once the brief access test was finished,

each mouse with an optrode was tested in the optotagging task, and the laser was turned on 2 s at 50

Hz. ChR2-expressing cells LHA^{Vgat+} were those with an action potential evoked within $\leq$ 15 ms latency

[34], as measured by a cumsum statistic described in [33].

Palatability-responsive neurons

We used a best-window analysis to identify neurons whose firing rate correlated with palatability-

induced oromotor responses, following a previous study [29]. This method has been used to detect

palatability-related activity in a brief access test. Briefly, during each recording session, a palatability-

index (PI) was calculated. The PI reflects the overall appetitive oromotor responses elicited by each

tastant delivered in the session. It was computed by averaging the lick rate (number of licks/ 7s) during

the entire reward epoch, including all the trials per session. In mice, the PI takes values between 0-8

Hz for sucrose stimuli [35]. 0 Hz means animals ultimately rejected a solution in all trials, whereas 8

Hz indicates, they licked continuously during the entire reward epoch, thus reflecting a greater

palatability response elicited by the tastant. Then, the firing rate was calculated for a variety of time

centers (from 0.25 to 6.75 s with 0.5 s steps), and multiple window sizes (from 0.5 to 7 s, 0.5s steps),

such that each window was estimated as the center ± (window size/2). We identified the windows

where the mean firing rate was significantly different for at least one tastant delivered (i.e., tastants,

using a Kruskal-Wallis test at an alpha of 0.05). Next, we computed Pearson's correlation coefficient

(r; the alpha level at 0.05) between both the PI and the firing rate on a trial-by-trial basis. The window

with the largest absolute Pearson's correlation coefficient was selected as the "best-window." Thus,

for the "best-window" and each statistical test (i.e., Kruskal-Wallis test and Pearson's correlation

coefficient), a permutation assay was used for multiple-testing correction. This analysis was

accomplished by shuffling the different tastant delivered 20,000 times (using the Matlab function

"shuffle"). A corrected p-value was obtained using the following formula $p= (k+1)/(n+1)$, where k is

the times in which a permuted distribution leads to a p-value smaller than the original p-value and n is

the number of repetitions. Only time-windows with p's < 0.05 in both tests (Kruskal-Wallis and

Pearson Correlation) were considered as statistically significant. Thus, the "best-window" is where the

firing rate maximally correlated with the oromotor responses elicited by sucrose's palatability on a

trial-by-trial basis. Results were qualitatively similar if we used the lick bout duration rather than the

lick rate to compute the PI (data not shown) [29].

Data Analysis

Data analysis was performed using MATLAB (The MathWorks Inc., Natick, MA) and GraphPad Prism

(La Jolla, CA, USA). We used the mean ± sem and the α level at 0.05.

Results

Initially, we determined the optimal stimulation parameters to activate LHA GABAergic neurons. This

was accomplished by implanting an optrode in naïve mice that constitutively expressed ChR2 in

GABAergic neurons expressing the gene for the vesicular γ-aminobutyric acid transporter (VGAT-ChR2 mice) [36] (Figure 1A-B). We found that a laser stimulation of 50 Hz (10 ms width) evoked the strongest LHA neuronal responses (Figure 1C-F) [37]. Unless otherwise mentioned, this frequency was used for all subsequent experiments.

LHA neurons encode sucrose's palatability

In rats, previous electrophysiological studies suggest that ensembles of LHA neurons process palatability related information [17,18,38,39]. Based on these results, we then asked whether LHA neurons could provide information about sucrose's palatability by using a behavioral task with only palatable stimuli (i.e., artificial saliva (AS), water, sucrose 3 wt%, and sucrose 18 wt%). We operationally defined sucrose's palatability as the enhancement of hedonically positive oromotor responses triggered by increasing sucrose concentrations [29–31]. Specifically, oromotor responses include an increase in the lick rate or bout size. Thus, this definition of palatability should not be confused with a conscious hedonic feeling of pleasant taste that humans experience [40,41]. To this end, taste responses were recorded from a total of 284 LHA neurons, while VGAT-ChR2 mice performed a brief access test (Figure 2A). To uncover palatability-responsive neurons, we analyzed neuronal activity using a previously described "best-window analysis" [29] (see Materials and Methods). This strategy detects the optimal interval, where the firing rate best correlates with the sucrose's palatability index (PI) computed as the average lick rate, on a session by session basis, evoked by each gustatory stimulus in the Reward epoch (Figure 2B). Figure 2C depicts the responses of two representative LHA neurons: as expected, we found one whose activity increased as the sucrose concentration increased (positive Pearson's correlation, r = 0.87; Figure 2C *left*), and the other was negatively associated with palatability oromotor responses evoked by sucrose (r = -0.64; Figure 2C

right). Figure 2D shows the normalized activity exhibiting either positive (59/284 neurons, 21%) or

negative correlations (76/284, 27%) with the PI. Thus, these data confirmed that the LHA mice's

neurons exhibit palatability-related responses, and this activity can be either correlated or anti-

correlated with the sucrose solution's PI.

Dynamic tracking of palatability over the entire course of a behavioral session

Having demonstrated that LHA neurons encode sucrose's palatability, we then explored whether LHA

palatability-related neurons can dynamically adjust their responses to track the changes in sucrose's

palatability across the session when the animals would approach satiation. We found that the lick rate

declined across the session as a function of sucrose's palatability (Figure 2E [RM ANOVA, main effect

of stimuli F $_{(3, 402)}$ = 163.7, p < 0.0001,effect of time F $_{(9, 1206)}$ = 100.4, p < 0.0001]), and significant

stimuli by time interaction F $_{(27, 3618)}$ = 16.54, p < 0.0001]). Briefly, early in the session, animals licked

more for all stimuli, but as the session progressed, the lick rate declined especially for AS and water

but not for sucrose (3% and 18%), indicating that they tracked palatability responses rather than satiety

(Figure 2E). Similar to what has been found in the Nucleus Accumbens Shell (NAcSh) [29], a brain

region that sends outputs to LHA and is involved in feeding and palatability [29,42], we also found

that the LHA neuron's firing rate (either increasing or decreasing neurons) adjust across the session to

reflect sucrose's palatability (Figure 2F). These data indicate that whithin the session as the animals

approach satiety, the responses of the Palatability related-neurons tracked the rapid decline in the lick

rate mainly for the two less palatable stimuli (i.e., AS and water) but not for the higher sucrose-

concentration (sucrose 3% and 18%), suggesting these neurons are tracking palatability and not satiety.

A subpopulation of opto-identified LHA^{Vgat+} neurons encodes sucrose's palatability

To identify LHA^{Vgat+} neurons from our recordings, we opto-stimulated the same mice recorded in the brief access test seen in Figure 2. We verified the stability of the waveforms between tasks (i.e., brief access test and optotagging), and thus we only included single units with stable waveforms in the analysis (Supplementary Figure 1). Figure 3A displays the setup and the normalized (Z-score) activity of laser-activated neurons (48%; 137/284) (for details, see Material and Methods). Since these neurons may include ChR2 expressing cells and other LHA neurons modulated by indirect polysynaptic feedback of afferent fibers from other areas in the brain [43], we decided to name them putative LHA^{Vgat} (pLHA^{Vgat}) neurons. Note that pLHA^{Vgat} neurons comprise the ensemble recruited by the optogenetic stimulation of LHA GABAergic neurons, suggesting they convey similar information. Therefore, we next evaluated whether pLHA^{Vgat} neurons were correlated with sucrose's palatability. In total, 50% (69/137) of laser-activated neurons exhibited responses that were significantly palatability related; specifically, 34% showed a positive correlation (46 out of 137), whereas 17% (23 out of 137) had a negative correlation with sucrose's palatability (Figure 3B, Positive vs. Negative chi-square test $_{(1, 274)}$ = 10.25, $p < 0.01$). Figure 3C shows the population activity of pLHA^{Vgat} neurons with either a positive (yellow) or negative (purple) correlation with the PI (red). Furthermore, from all recorded LHA neurons that encode sucrose palatability with a Positive correlation (n = 59, see Figure 2D), we found that more than 78% (46/59) belonged to the pLHA^{Vgat} (Figure 3D). In contrast, only 30% (23/76) of pLHA^{Vgat} neurons were Negative palatability related (Figure 3D, Positive vs. Negative chi-square test $_{(1, 135)}$ = 30.25, $p < 0.0001$). Thus, pLHA^{Vgat} tends to encode sucrose's palatability with a biased towards positive correlations.

To confirm the identity of LHA^{Vgat+} neurons, we searched for neurons in which a brief pulse of light-evoked an action potential with an early latency (< 15 ms), that would reflect the expression of ChR2 in their somas [34]. From the pLHA^{Vgat} population, in total, 52 % (71 / 137) neurons exhibited an early

response to blue light (Figures 3E-F), suggesting that these pLHA[Vgat] neurons were LHA[Vgat+] neurons.

We found that over half of the identified LHA[Vgat+] neurons, 55% (39/ 71), were involved in encoding

sucrose's palatability, and the remaining (45%) were unmodulated by sucrose. From the LHA neurons

related to sucrose's palatability, 31% of LHA[Vgat+] neurons positively correlated with sucrose's

palatability (22/71), and 24% were anticorrelated (17/71) (Figure 3G, Positive vs. Negative; n.s.). The

LHA[Vgat+] neuronal responses with positive and negative correlations with lick-related palatability

responses are seen in Figure 3H. Once again, we compared the proportion of opto-identified LHA[Vgat+]

neurons against all palatability-responsive neurons recorded in the LHA. We found a trend to encode

sucrose's palatability in a positive, rather than negative manner. That is, 37% (22/59) of LHA[Vgat+]

neurons belonged to the Positive palatability-related population, whereas only 22% (17/76) were

Negative palatability-related neurons (Figure 3I, Positive vs. Negative; n.s.). Altogether, these data

suggest that LHA[Vgat+] neurons tend to encode sucrose's palatability in a positive rather than in a

negative manner. More important, our data are in agreement with the idea that the LHA[Vgat+] neurons

comprise many heterogeneous subpopulations with different functional responses [22].

Non-LHA[Vgat] neurons negatively encode sucrose's palatability

We also found that 25% of LHA neurons exhibited a laser-induced inhibition (70/284, non-LHA[Vgat]

neurons, Figure 3J), and the remaining 27% (77/284) were unmodulated during blue light stimulation

(Figure 3J, black trace). Unlike pLHA[Vgat] neurons, only 7% (5 out of 70) of the non-LHA[Vgat] neurons

showed a positive correlation, whereas the vast majority 36% (25 out of 70), had a negative correlation

with sucrose's palatability (Figures 3K and 3L, Positive vs. Negative chi-square test $_{(1, 140)}$ = 16.97, p

< 0.0001). These results suggest that non- LHA[Vgat] neurons (perhaps glutamatergic Vglut2 neurons)

preferentially encode sucrose's palatability with negative correlations (Figure 3M, Positive vs.

Negative chi-square test $_{(1, 135)}$ = 11.46, p < 0.001).

Activation of LHA^{Vgat+} neurons drives sucrose intake

If a subpopulation of LHA GABAergic neurons preferentially encodes sucrose's palatability by

increasing firing rates, its optogenetic stimulation should promote increased sucrose intake. To

characterize its impact on sucrose intake, we used an open loop optogenetic stimulation involving 5

min blocks with laser and no-laser stimulation, while naïve VGAT-ChR2 mice had *ad lib* access to a

sipper filled with sucrose 10% (Figure 4*A*). We analyzed the intake as a function of laser frequency

and found that in the first three blocks with laser, 50 Hz stimulation induced a significant increase in

sucrose intake (Figure 4B; Blocks 2, 4, and 6) compared with the previous blocks of no-laser (Figure

4B; *p* < 0.001; Blocks 1, 3, and 5). Figure 4*C* shows a raster plot for licking responses of a

representative open loop session. At 50 Hz, we found that feeding (licking) was elicited within a 0.72

s ± 0.03 latency from laser onset, and it abruptly stopped (0.27 s ± 0.1) after laser offset (see blue dash

rectangles). VGAT-ChR2 mice exhibited an increase in the laser-bound feeding index (number of licks

in a 2.5 s window from laser onset divided by total licks in the session; hereafter named laser-bound

feeding) as the laser frequency increased (Figure 4*D;* two-way ANOVA, group by frequency

interaction; F $_{(5, 36)}$ = 12.86, *p* < 0.0001). Thus, to induce sucrose intake, a subset of LHA^{Vgat+} neurons

requires continued activation. Our data then confirmed that 50 Hz is the best stimulation frequency to

drive neuronal responses and sucrose intake.

LHA^{Vgat+} neurons do not induce a persistent hunger state

A previous study showed that brief optogenetic stimulation of "hunger-inducing" AgRP neurons before

food availability promotes consummatory behavior that persists for several minutes in the absence of

continued AgRP neuron activation [44,45]. Based on this study, we tested whether the pre-stimulation

of LHA^{Vgat+} neurons could evoke a similar hunger state. Unlike AgRP neurons [44,45], pre-stimulation

of LHA^{Vgat+} neurons failed to evoke and sustain subsequent sucrose intake (see Supplementary Figure

2). Our results suggest that feeding occurs while LHA^{Vgat+} neurons are active (Figure 4C), but they do

not induce a persistent hunger state as AgRP neurons do [45].

Open loop activation of LHA^{Vgat+} neurons drives intake of the highest sucrose concentration available

To test whether activation of LHA^{Vgat+} neurons promotes the intake of the most palatable available

stimuli, naïve WT and VGAT-ChR2 mice were opto-stimulated in an open loop protocol, with access

to three adjacent ports containing water, sucrose 3%, or sucrose 18% (Figure 5A). As expected, both

WT and VGAT-ChR2 mice preferred sucrose 18% over water and sucrose 3% (Figure 5B, [WT: two-

way ANOVA main effect tastants; $F_{(2, 1560)} = 543.3$, $p < 0.0001$; VGAT-ChR2: $F_{(2, 3144)} = 983$, $p <$

0.0001]). However, we found that during laser stimulation, sated VGAT-ChR2 mice increased their

intake of 18% sucrose but mainly during the laser-activated blocks compared with controls (Figure 5B-

right [two-way ANOVA group by laser block interaction $F_{(7, 1568)} = 31.86$, $p < 0.0001$]) and rarely

licked in the absence of stimulation (Figure 5C). In this regard, VGAT-ChR2 mice seem to counteract

the evoked sucrose intake by voluntarily restraining consumption in no-laser blocks (Blocks 3, 5, and

7), resulting in no significant differences between groups in the total intake of 18% sucrose (Figure

5C, see black squares, unpaired Student's t-test, $t_{(196)} = 1.512$, $p = 0.1320$). A between days analysis

revealed that the evoked 18% sucrose intake began from the first stimulation day (Figure 5D [RM

ANOVA laser blocks, $F_{(1, 22)} = 64.64$, $p < 0.0001$]), although sucrose consumption ramps up

throughout the days (Figure 5D [RM ANOVA main effect days, $F_{(10, 220)} = 2.169$, $p < 0.05$]). This

suggests that LHA^{Vgat+} neurons, rather than inducing hunger *per se*, induced a learning process that

potentiates the intake of the most palatable stimulus and confines consumption mainly in the presence

of opto-stimulation.

The small increase in water and sucrose 3% intake observed during opto-stimulation of LHA^{Vgat+}

neurons could be explained by random stimulation near those tastants (Figure 5B *right panel*).

Accordingly, we found in laser blocks 2, 4, and 6, a significant increase in water and sucrose 3% in the

VGAT-ChR2 mice compared with the WT mice that completely neglected those tastants (Figure 5B,

^ $p < 0.05$ for sucrose 3%; † $p < 0.05$ for water). We hypothesize that this additional intake can be

attributed to trials where the laser-activation occurred near these less palatable stimuli. To answer this,

we employed a videography analysis (Figure 6A-B). When opto-stimulation occurred in the distance

minor to 50 pixels (approximately 3 cm, see Materials and Methods) relative to water or sucrose 3%

ports, the licking probability increased significantly after opto-stimulation (Figure 6B). The angle of

the head was less informative (Figure 6C-D). Conversely and since mice are naturally attracted to the

most palatable tastant, they spent more time near the lateral port with 18% sucrose, increasing their

overconsumption (Figure 6C, Video 1). Moreover, we observed that for the sucrose 18% port, even

when the mouse position was twice as far (i.e., 6 cm) from sucrose 18% port, the licking probability

increased significantly. These data are consistent with a study showing that opto-stimulation of

LHA^{Vgat+} neurons promotes the intake of the nearest stimulus [24] but further demonstrates that the

most palatable stimulus has nearly twice the distance of attraction than the other less palatable options.

Open loop activation of LHA^{Vgat+} neurons also increases the time spent and drives the intake of the

most palatable solid food available

We also explored whether LHA GABA neurons could induce the intake of the most palatable solid

food available. Thus, using an open loop protocol, we opto-stimulate these neurons while mice choose

among different solid foods. We found that optogenetic activation of LHA^{Vgat+} neurons increased the

time spent near the most palatable food available. Mice spent more time near the high-fat diet than the

granulated sugar cube or the Chow food pellet (see Supplementary Figure 3 and Video 2). When mice

were able to choose between a sugar cube and a Chow pellet, activation of these neurons increased the

consumption of sugar cube over Chow (see Supplementary Figure 4 and Video 3). Our data suggest

that open loop stimulation of LHA GABAergic neurons promotes the intake of the most palatable food

available.

Closed-loop stimulation of LHA^{Vgat+} neurons drives the intake of the nearest appetitive stimuli

To further test the idea that LHA^{Vgat+} neurons could induce preference to a proximal (but less palatable)

stimulus over a distal one with higher hedonic value, we used a closed-loop stimulation protocol. That

is, the same mice (from Figure 5) were photostimulated, but here the laser was triggered by each head

entry in the central port (i.e., opto-self-stimulation). This closed-loop configuration guarantees that

stimulation only occurs proximal to water, the least palatable stimulus of the three in the box (Figure

7*A-right*). We found that activation of LHA^{Vgat+} neurons is rewarding since transgenic mice visited the

central port significantly more and performed a higher number of opto-self-stimulations than the WT

mice (Figure 7B, unpaired Student's t-test, $t_{(196)} = 12$, $p < 0.0001$). Surprisingly, the total number of

licks in the session seen in Figure 7C shows that during closed-loop stimulation, the VGAT-ChR2 mice

explored and consumed more water than the WT mice. Note that in both open loop and closed-loop

protocols, the WT and VGAT-ChR2 mice exhibited, at the end of the session, a similar number of licks for 18 wt% sucrose (Figure 7C [two-way ANOVA group by protocol interaction F $_{(1, 391)}$ = 2.172, p = 0.1414]). That is, the total licks for sucrose 18% were not significantly different between groups, suggesting that these neurons do not interfere with the overall attractiveness of sucrose. However, by counting the number of licks given when the laser was turned on (in a 2.5 s window from laser onset- see Figure 7D), we uncovered a striking change in preference. As noted, in the open loop, the VGAT-ChR2 mice exhibited a higher preference to sucrose 18 wt%, over both sucrose 3 wt% and water. Now, in the closed-loop protocol, the same transgenic mice consumed more water compared with WT ($p <$ 0.0001), and the consumption of water was about the same as sucrose 18% (p = 0.8731, n.s.; Figure 7D). The increase in water intake was selective to the photostimulation window since the transgenic mice only drank sucrose 18%, and neglected water when the laser was turned off (see arrow Figure 7E, p < 0.0001). Thus, LHA^{Vgat+} neurons could promote water intake, but only when it is the nearest stimulus. The sucrose preference index (Figure 7F) showed that during the open loop all VGAT-ChR2 mice preferred sucrose 18% over water (Figure 7F; values > 0.5 and near to 1), whereas in the closed-loop protocol, when water was the nearest stimulus, most mice (n = 9 out of 12), significantly diminished their sucrose preference (Figure 7F, see the drop in preference index; paired Student's t-test, t $_{(58)}$ = 7.98, $p <$ 0.0001). From these 9 animals, 6 exhibited a higher preference to water over sucrose 18% (Preference index values < 0.5; Video 4). Figure 7G shows the color-coded location of fiber optic tips in each VGAT-ChR2 mice plotted in Figure "7F." However, the location of the optical fibers does not explain the variability in preference. Perhaps variability is due to different behavioral strategies used by each mouse. Nevertheless, the large variability in the drop of sucrose's preference resembles the findings with sweet-induced facilitation by electrical ICSs [9]. These data suggest that LHA^{Vgat+} neurons drive consummatory behavior by integrating the proximity and the hedonic value of the stimulus.

526
527

Previous studies have shown that after repeated LHA bulk stimulation (either electrically or with

optogenetic targeting all cell-types together), subjects switched from exhibiting stimulus-bound

feeding to only self-stimulating, indicating that LHA evoked feeding and reward could represent two

distinct processes [37,46]. Consequently, we explored whether laser-bound feeding changed after

repeated optogenetic stimulation. In contrast to previous studies, we observed that laser-bound feeding

induced by LHA^{Vgat+} neurons strengthened across sessions (Figure 8A, First 3 days: r = 0.12, p = 0.69;

Last 5 stimulation days: r = 0.63, p < 0.05) and after repeated stimulation, both opto-self-stimulation

and laser-bound feeding exhibited a robust correlation (Figure 8B, First 3 days: r = 0.42, p = 0.16; Last

5 days: r = 0.61, p < 0.05), suggesting that LHA evoked-feeding (licking) involves a learning process

[47] and that LHA GABAergic neurons are a common neural substrate for feeding and reward.

The "stimulus-proximity effect" is not restricted to liquid tastants, and it also occurs with chocolate
and Chow pellets

The "proximity effect" evoked by LHA^{Vgat+} neurons was not restricted to liquid tastants, it also applied

to solid foods. In a real-time place preference arena with four stimuli (Figure 9A), when chocolate

pellets or Chow food were the designated foods near opto-self-stimulation of LHA^{Vgat+} neurons, mice

also increased the time spent near those foods (Figure 9B; Video 5) as well as their intake (Figure 9C).

Thus, these LHA GABAergic neurons reinforced the approach and exploration to any, if not the most

appetitive stimuli, that happened to be proximal to the opto-stimulation.

LHA^{Vgat+} neurons activation is rewarding

In the same WT and VGAT-ChR2 mice tested in the closed-loop configuration seen in Figure 7, we went on to show that mice visited the central water-port to opto-self-stimulate. To do this, we performed extinction sessions with the laser disconnected (Figure 10A). We observed a rapid decrease in the number of opto-self-stimulations (Figure 10B; Extinction phase, gray shadow). As expected, self-stimulation rapidly recovered when the laser was turned on again (Figure 10, see water after extinction). Thus, LHA^{Vgat+} neurons convey a hedonically positive, rewarding signal [22].

LHA^{Vgat+} neurons also promote licking an empty sipper

A previous study demonstrated that chemogenetic activation of LHA^{Vgat+} neurons increased gnawing to non-edible cork [23]. In the absence of other stimuli but cork, we also observed gnawing behavior in some mice, although we did not systematically study stereotypy (see Video 6). Instead, we further explored this idea by replacing water, in the central port, with an empty sipper (Figure 10, Empty sipper). Transgenic mice continued opto-self-stimulating at the same place as if water were still present in the central port. LHA^{Vgat+} stimulation evoked licking an empty sipper compared with WT mice (Figure 10C, unpaired Student's t-test, t $_{(40)}$ =2.532, $p < 0.05$). Thus, LHA^{Vgat+} neurons evoke an imminent urge to express consummatory behavior even in the form of dry licking an empty sipper, a non-biological relevant stimulus. This effect is perhaps mediated by an increase in the rewarding value of appetitive oromotor responses *per se*.

When sucrose is available, activation of LHA^{Vgat+} neurons neither promotes liquid intake of an

aversive bitter tastant nor tolerance of punishment, but it further increases sucrose consumption

To explore whether the "stimulus-proximity effect" was also applied to aversive stimuli, we replaced

the central stimulus (water) with aversive stimuli, including quinine or airpuffs (Figure 10*A*). Upon

quinine presentation, the number of licks given in the central port sharply decreased (Figure 10C,

green), but transgenic mice continued to self-stimulate and go to the lateral port to lick for sucrose 18%

(Figure 10D, green). During the airpuff delivery phase (Figure 10D, dark gray), we obtained similar

results than those in quinine stimulation. Next, we required the animal to lick, at least once, the central

sipper to trigger opto-stimulation (thus making the airpuffs unavoidable). Under this condition,

transgenic mice completely stopped self-stimulation (Figure 10B, cyan) and aborted 18% sucrose

intake from the lateral port (Figure 10D, cyan). These results suggest that, in sated mice, LHA^{Vgat+}

neurons did not induce quinine intake nor increase tolerance to airpuffs.

Finally, we explored whether proximity to the most palatable tastant further facilitated its

overconsumption. Thus, we exchanged the position of water and sucrose 18%. When sucrose 18% was

delivered in the central port, transgenic mice greatly overconsumed it (Figure 10C, pink). The intake

of sucrose 18% was higher than for all the other tastants tested previously ($p < 0.0001$). These results

collectively suggest that activation of LHA^{Vgat+} neurons is rewarding and promotes increased

consumption of the nearest stimulus even if it is not the most palatable (e.g., water and empty sipper).

If the nearest stimulus happens to be the most palatable (i.e., sucrose 18%), then LHA GABA neurons

further facilitated its consumption.

 LHA GABA neurons increased quinine intake but only in water-deprived mice

Having demonstrated that in the presence of sucrose, activation of these neurons failed to increase

quinine intake (Figure 10C green), with a new group of naïve mice, we then explored whether LHA^{Vgat+}

neurons could induce quinine intake when it was the only option. As expected, in sated mice, we found

that activation of LHA^{Vgat+} neurons did not affect quinine intake compared with WT (Figure 11A).

Surprisingly, it promoted a higher quinine intake when transgenic mice were water-deprived (Figure

11B, unpaired Student's t-test, t (16) = 3.623, p < 0.01), specifically during photostimulation window

(Figure 11B, unpaired Student's t-test, t (16) = 5.821, p < 0.0001), suggesting that activation of these

neurons is sufficient to increase the acceptance of bitter tastants but only during water deprivation. In

contrast, we found that regardless of homeostatic needs, activation of these neurons increased sucrose

18% intake relative to WT mice (Figure 11C-D).

Activation of LHA^{Vgat+} neurons enhances palatability

We next explored whether, at equal stimulus distance, these neurons could enhance palatability

responses. To do this, in a new group of water-deprived mice, we employed a brief access test again.

In this task, we delivered gustatory stimuli water, sucrose 3%, and sucrose 18% from the same sipper

tube, but in different trials. Figure 12*A* displays the trial's structure (same conventions as in Figure 2).

We found that WT mice increased their licking rate as a function of sucrose's concentration, reflecting

its palatability (Figure 12B, *left panel*). In contrast, during the 2 s of laser stimulation, the VGAT-ChR2

mice exhibited a greater lick rate for all three tastants. However, after the laser was turned "off," these

mice abruptly stopped licking for water and sucrose 3% (Figure 12*B, right panel*) compared with the

WT group. Moreover, when stimulation was paired with water trials, transgenic mice selectively

increased their lick rate to water (Figure 12C, blue line and arrow), even surpassing licking responses

evoked by the most palatable sucrose 18%. A similar enhancement of oromotor responses was

observed by pairing sucrose 3% trials with LHA^{Vgat+} opto-self-stimulation (Figure 12D, pink).

Likewise, the laser-bound feeding (licking rate) was strongest when sucrose 18% was paired with the

optogenetic stimulation (Figure 12E, red, see arrow).

Finally, we show that palatability responses could be artificially extended as long as LHA^{Vgat+} neurons

were continuously activated. For this, we photostimulated them for up to 7 s. We observed a similar

enhancement pattern, maintained for the duration of opto-stimulation (see Figure 12F-I see arrows),

suggesting that LHA^{Vgat+} neurons can adjust the enhancement of oromotor palatability responses by

simply sustaining its neuronal activity. We verified that using optrode recordings, that 7 s optogenetic

stimulation produced sustained LHA^{Vgat+} neuronal responses (Figure 12J). Moreover, transgenic mice

did not merely use the light as a cue to guide behavior since laser-stimulation with a mock optical fiber

failed to increase licking (Figure 12K). In sum, our data demonstrate that activation of LHA^{Vgat+}

neurons triggers a reinforcing signal that amplifies the positive hedonic value of proximal stimuli,

promoting overconsumption.

Discussion

LHA has historically been viewed as a critical center for feeding [1–3], although it also processes

sucrose's palatability related information [18,38,48]. In addition to caloric value, sucrose's palatability

is the affective or hedonic attribute of sweetness that determines whether to like it or not. Despite the

importance of palatability to promote overconsumption, the specific LHA cell-type(s) identity involved

in processing sucrose's palatability has remained elusive. Our results demonstrated that a

subpopulation of LHA^{Vgat+} GABAergic neurons encodes sucrose's palatability by exhibiting two

opposite modulatory patterns, either correlating positively or negatively with the palatability index,

with a bias towards a positive correlation. Furthermore, opto-stimulation of LHA^{Vgat+} cell somas

promoted approach and intake of the most salient available and palatable tastant. In contrast, opto-self-

stimulation promoted increased liquid intake of the less-attractive and proximal stimuli, despite having

more palatable but distal tastants available. These findings show that LHA^{Vgat+} neurons compute and/or

combine, at least, two types of information: one related to stimulus proximity and the other to

palatability that results in enhancing stimulus-saliency [24]. More important, these data uncovered a

new function of a subgroup of LHA^{Vgat+} neurons as physiological potentiators of sucrose's palatability.

Previous work has shown that LHA^{Vgat+} population contains many subpopulations with different

functional responses, with at least one ensemble responding to appetitive (approach) and other to

consummatory behaviors [22]. Furthermore, activation of these neurons is rewarding, induces

voracious eating [22,23], and promotes interaction of the nearest stimuli (either objects or other mice)

[24], suggesting that they play a role in multiple motivated behaviors. However, it is also known that

LHA connects and receives direct inputs from multiple cortical and subcortical gustatory regions [10–

13,49], and some electrophysiological studies report that LHA neurons respond to gustatory stimuli –

in particular to tastant palatability [16,17,38,48,50]. As noted, in rodents, palatability is operationally

defined as the enhancement of hedonically positive oromotor responses induced by stimulating the

tongue with ascending sucrose's concentrations (Berridge and Grill, 1983; Spector et al., 1998;

Villavicencio et al., 2018). In agreement with this definition, we found that a subpopulation LHA

palatability related-neurons tracked licking oromotor-related responses by increasing or decreasing

their activity in a sucrose-concentration dependent manner. Also, whithin a session, LHA neurons

tracked the sucrose's palatability rather than satiety or hunger signals. We showed that these LHA^{Vgat+}

neurons could function as enhancers of sucrose's palatability. Optogenetic activation of these neurons

can also enhance water's palatability if it is the nearest stimuli. We found an increased lick rate for

water during activation of these neurons as if the animal were sampling a high sucrose concentration.

Moreover, their activation promotes the intake of liquid sucrose (or solid granulated sugar cube). These

neurons also increased the consumption of other more palatable stimuli like high-fat pellets (see

Supplementary Figure 3; Video 2), similar to the other GABAergic neurons but in zona incerta [51]. It

is tempting to propose that LHA GABAergic neurons could mediate the supra-additive effects of

combining fats and carbohydrates such as sucrose [52], but this remains to be experimentally tested.

Although activation of LHA^{Vgat+} promotes substantial feeding behavior, it has become clear that

LHA^{Vgat+} neurons are not directly involved in evoking hunger [23,25,44], as AgRP neurons do [45]. In

agreement with these findings, we found that the intake induced by a subset of LHA^{Vgat+} convey a

positive valence signal that combines both stimulus proximity and palatability related information.

Thus, these neurons enhance the saliency of nearby hedonically positive stimuli, whether those stimuli

are sapid chemicals, as we show, or social cues, as in the approach behavior towards juvenile or female

intruders and new objects [24].

It is important to highlight that opto-self-activation of LHA^{Vgat+} neurons resembles many hallmark

behaviors evoked by LHA electrical stimulation. In particular, our results could shed some light on

why, at low-intensity electrical currents, lever pressing to deliver ICSs only occurs if food (or sucrose)

is in close proximity [6,7,53,54]. Consistent with its role in enhancing sucrose's palatability, a

subpopulation of GABAergic neurons could explain the long-standing puzzle of why sweet tastants

further potentiate the rate of LHA electrical ICSs [9]. We concluded that a subpopulation LHA^{Vgat+}

neurons could account for many, if not all, of these electrically induced phenomena. That said, we also

found differences between unspecific LHA stimulation and our targeted LHA^{Vgat+} stimulation. Unlike

electrical LHA stimulation, we found that the laser-bound feeding was observed in all tested VGAT-

ChR2 mice (n = 25, Figures 4-12). In contrast, to the high variability found in rats exhibiting LHA

electrically-induced feeding, one study reported that only 12 of 34 rats showed stimulus-bound feeding

[55]. Similarly, a large variability was observed when unspecific bulk optogenetic stimulation activated

all cell-types found in LHA, simultaneously [46]. These studies reported that after repeated LHA

stimulation (either electrically or with optogenetics), some subjects switched from exhibiting stimulus-

bound feeding to only self-stimulating, suggesting that these two processes were flexible and not

correlated [37,46]. In contrast, we found that repeated stimulation of LHA^{Vgat+} neurons increases laser-

induced feeding (licking). Likewise, the correlation between optogenetic self-stimulation and laser-

bound feeding increases through stimulation days (Figure 8). That is, the more the animals self-

stimulated, the stronger the evoked laser-bound licking was. Thus, LHA^{Vgat+} neurons are the common

neural substrate for evoking both feeding and reward. Though, it was recently shown that they do it by

using two projection pathways; reward via a VTA projection and feeding via the peri-Locus Coeruleus

nuclei [25].

Given that LHA is involved in reward and aversion [48], we next tested whether LHA^{Vgat+} neurons

could promote bitter tastants' intake. We found that opto-stimulation of LHA^{Vgat+} neurons failed to

promote quinine intake, a bitter tastant, or tolerance of an aversive airpuff when sucrose was also

available. Thus, these neurons play a minimal role in increasing the preference to a proximal but

aversive stimulus over distal sucrose. Furthermore, in sated mice and using a single bottle test, these

neurons also failed to increase quinine intake. Unexpectedly, during water deprivation, a copious

quinine consumption was observed. These results agree with previous findings that chemogenetic

inhibition of LHA^{Vgat+} neurons did not alter the quinine rejection responses. Thus, in sated mice, these

neurons are not necessary to express hedonically negative responses induced by bitter tastants [56].

However, they did not explore their sufficiency. Our results showed that water deprivation temporarily gated LHA^{Vgat+} neurons to promote quinine intake, and further demonstrated that their activation is sufficient to increase acceptance of an aversive tastant during water deprivation.

The LHA comprises multiple heterogeneous and overlapping populations based on the expression of genetic markers for neuropeptides, receptors, and proteins involved in the synthesis and vesicular packaging of neurotransmitters [57,58]. Thus, the LHA^{Vgat+} population can be further subdivided into GABA neurons expressing leptin receptor (LepRb) [58,59], or the neuropeptide galanin (Gal), or neurotensin (Nts) [58,60,61]. It is known that the activation of LHA-GABA-LepRb is rewarding [62], similar to LHA^{Vgat+} neurons. Likewise, LHA GABA-Gal expressing neurons are related to food reward behavior, but unlike LHA^{Vgat+}, these neurons do not promote food consumption [60]. From these subpopulations, only the LHA GABA-Nts neurons recapitulate some (but not all) of the behavioral effects reported here. Unlike LHA^{Vgat+} neurons, a previous study found that LHA GABA-Nts neurons do not increase chow intake. Instead, they promote the liquid intake of palatable tastants (water, NaCl, and sucrose). In sated mice, chemogenetic activation of GABA-Nts increased the intake of bitter quinine, albeit with lower magnitude, when it was the only liquid available to eat with chow food, a marked contrast to our findings with broad LHA GABAergic neurons activation. Although, they did not explore quinine intake, in the absence of chow food or water-deprived mice. However, they performed a two-bottle test and found that GABA-Nts neurons increased water intake over quinine, suggesting that these neurons cannot develop a preference to bitter tastants, similar to what we found for the LHA^{Vgat+} population. Also, similar to our findings, activation of LHA GABA-Nts induced water-drinking, that was further facilitated if the solution was sucrose [61]. Thus, it will be interesting to determine the role that LHA GABA-Nts neurons play in encoding and potentiating sucrose's palatability. It follows that the LHA contains nested functions encoded in each subpopulation (or cell-types) that are then recruited selectively to exert a more refined control over feeding and reward.

732 A caveat of this study is that we employed bacterial artificial chromosomes (BAC) transgenic strain

733 mouse, VGAT-ChR2, that constitutively expressed ChR2 in GABAergic neurons expressing the gene

734 for the VGAT [36]. In this model, we cannot rule out unintended activation of GABAergic terminals

735 from distal regions [63], which also occurs with classic electrical stimulation. Nevertheless, in the more

736 specific transgenic model, the Vgat-ires-Cre mice has been found a similar feeding-bound behavior for

737 chow food [25], as we shown here (see Chow in Figures 9B and C; and Video 5). Moreover, the VGAT-

738 ChR2 transgenic model affords important advantages such as a consistent expression of ChR2 [64] and

739 heritable transgene expression patterns across experimental cohorts [65], which increased

740 reproducibility across animals tested. It is also a more selective model to characterize GABAergic

741 neurons (excluding the glutamatergic component) and their effects recapitulating classical effects

742 observed with electrical LHA stimulation [3,8].

744 In summary, here we found that at least a subpopulation of LHA$^{\text{Vgat+}}$ neurons could be an important

745 hub that links stimulus-proximity and palatability related information to potentiate the palatability of

746 nearby energy-rich foods containing sucrose.

748 **Conflict of interests**

749 The authors declare no competing financial interests.

751 **Author Contributions**

752 A.G., A.C., J. LI., and R.G. designed research; A.G., A.C., J. LI., L.PS., and M.L. performed research;

753 M.V. contributed unpublished analytic tools; A.G., A.C., J. LI., and R.G. analyzed data; A.G. and R.G.

754 wrote the paper; all authors reviewed and approved manuscript.

Funding

756 This project was supported in part by Productos Medix 3247, Cátedra Marcos Moshinky, fundación

757 Miguel Alemán Valdés, CONACyT Grants Fronteras de la Ciencia 63 and 10862, and Problemas

758 Nacionales 464 (to R.G.).

Acknowledgments

Aketzali Garcia had a CONACyT doctoral fellowship, and data in this work is part of her doctoral

dissertation in the Posgrado en Ciencias Biomédicas of the Universidad Nacional Autónoma de

México. We want to thank Martin Vignovich and Professor Sidney A. Simon for helpful comments in

an early version of this manuscript. We thank Mario Gil Moreno for building homemade optrode, and

valuable help of Juan de Dios Rodriguez-Callejas during the acquisition of confocal microscope

images. We also thank Ricardo Gaxiola, Victor Manuel García Gómez, and Fabiola Hernandez Olvera

for invaluable animal care.

References

1. Anand BK, Brobeck JR. Localization of a "feeding center" in the hypothalamus of the rat.
Proceedings of the Society for Experimental Biology and Medicine Society for Experimental
Biology and Medicine. 1951;77: 323–4.

2. Teitelbaum P, Epstein AN. The lateral hypothalamic syndrome: recovery of feeding and
drinking after lateral hypothalamic lesions. Psychological review. 1962;69: 74–90.

3. Delgado JM, Anand BK. Increase of food intake induced by electrical stimulation of the lateral
hypothalamus. The American journal of physiology. 1953;172: 162–8.
doi:10.1152/ajplegacy.1952.172.1.162

4.   Olds J, Milner P. Positive reinforcement produced by electrical stimulation of septal area and
other regions of rat brain. J Comp Physiol Psychol. 1954;47: 419–427. doi:10.1037/h0058775

5.   Mogenson GJ, Stevenson JA. Drinking induced by electrical stimulation of the lateral
hypothalamus. Experimental neurology. 1967;17: 119–27. doi:10.1016/0014-4886(67)90139-2

6.   Coons EE, Cruce JA. Lateral hypothalamus: food current intensity in maintaining self-
stimulation of hunger. Science. 1968;159: 1117–9. doi:10.1126/science.159.3819.1117

7.   Mendelson J. Lateral hypothalamic stimulation in satiated rats: the rewarding effects of self-
induced drinking. Science. 1967;157: 1077–1079. doi:10.1126/science.157.3792.1077

8.   Phillips AG, Mogenson GJ. Effects of taste on self-stimulation and induced drinking. J Comp
Physiol Psychol. 1968;66: 654–660. doi:10.1037/h0026523

9.   Poschel BPH. Do biological reinforcers act via the self-stimulation areas of the brain?
Physiology & Behavior. 1968;3: 53–60. doi:10.1016/0031-9384(68)90031-0

10.   Berthoud H-R, Münzberg H. The lateral hypothalamus as integrator of metabolic and
environmental needs: from electrical self-stimulation to opto-genetics. Physiol Behav.
2011;104: 29–39. doi:10.1016/j.physbeh.2011.04.051

11.   Simerly RB. Anatomical Substrates of Hypothalamic Integration. In: The Rat Nervous System.
Academic Press; 2004.

12.   Berk ML, Finkelstein JA. Efferent connections of the lateral hypothalamic area of the rat: an
autoradiographic investigation. Brain Res Bull. 1982;8: 511–526. doi:10.1016/0361-
9230(82)90009-0

13.   Ter Horst GJ, Luiten PG. Phaseolus vulgaris leuco-agglutinin tracing of intrahypothalamic
connections of the lateral, ventromedial, dorsomedial and paraventricular hypothalamic nuclei
in the rat. Brain Res Bull. 1987;18: 191–203. doi:10.1016/0361-9230(87)90190-0

14.   Ferssiwi A, Cardo B, Velley L. Gustatory preference-aversion thresholds are increased by
ibotenic acid lesion of the lateral hypothalamus in the rat. Brain Res. 1987;437: 142–150.
doi:10.1016/0006-8993(87)91535-6

15.   Berridge KC, Valenstein ES. What psychological process mediates feeding evoked by electrical
stimulation of the lateral hypothalamus? Behav Neurosci. 1991;105: 3–14. doi:10.1037//0735-
7044.105.1.3

16.   Schwartzbaum JS. Electrophysiology of taste, feeding and reward in lateral hypothalamus of
rabbit. Physiol Behav. 1988;44: 507–526. doi:10.1016/0031-9384(88)90313-7

17.   Yamamoto T, Matsuo R, Kiyomitsu Y, Kitamura R. Response properties of lateral
hypothalamic neurons during ingestive behavior with special reference to licking of various
taste solutions. Brain Res. 1989;481: 286–297. doi:10.1016/0006-8993(89)90805-6

18.   Li JX, Yoshida T, Monk KJ, Katz DB. Lateral hypothalamus contains two types of palatability-
812   related taste responses with distinct dynamics. The Journal of neuroscience : the official journal
813   of the Society for Neuroscience. 2013;33: 9462–73. doi:10.1523/JNEUROSCI.3935-12.2013

814   19.   Stuber GD, Wise RA. Lateral hypothalamic circuits for feeding and reward. Nature
815   neuroscience. 2016;19: 198–205. doi:10.1038/nn.4220

816   20.   Gutierrez R, Fonseca E, Simon SA. The neuroscience of sugars in taste, gut-reward, feeding
817   circuits, and obesity. Cell Mol Life Sci. 2020. doi:10.1007/s00018-020-03458-2

818   21.   Jennings JH, Rizzi G, Stamatakis AM, Ung RL, Stuber GD. The inhibitory circuit architecture
819   of the lateral hypothalamus orchestrates feeding. Science. 2013;341: 1517–1521.
820   doi:10.1126/science.1241812

821   22.   Jennings JH, Ung RL, Resendez SL, Stamatakis AM, Taylor JG, Huang J, et al. Visualizing
822   hypothalamic network dynamics for appetitive and consummatory behaviors. Cell. 2015;160:
823   516–27. doi:10.1016/j.cell.2014.12.026

824   23.   Navarro M, Olney JJ, Burnham NW, Mazzone CM, Lowery-Gionta EG, Pleil KE, et al. Lateral
825   Hypothalamus GABAergic Neurons Modulate Consummatory Behaviors Regardless of the
826   Caloric Content or Biological Relevance of the Consumed Stimuli. Neuropsychopharmacology :
827   official publication of the American College of Neuropsychopharmacology. 2016;41: 1505–12.
828   doi:10.1038/npp.2015.304

829   24.   Nieh EH, Vander Weele CM, Matthews GA, Presbrey KN, Wichmann R, Leppla CA, et al.
830   Inhibitory Input from the Lateral Hypothalamus to the Ventral Tegmental Area Disinhibits
831   Dopamine Neurons and Promotes Behavioral Activation. Neuron. 2016;90: 1286–1298.
832   doi:10.1016/j.neuron.2016.04.035

833   25.   Marino RAM, McDevitt RA, Gantz SC, Shen H, Pignatelli M, Xin W, et al. Control of food
834   approach and eating by a GABAergic projection from lateral hypothalamus to dorsal pons. Proc
835   Natl Acad Sci USA. 2020. doi:10.1073/pnas.1909340117

836   26.   Zocchi D, Wennemuth G, Oka Y. The cellular mechanism for water detection in the mammalian
837   taste system. Nat Neurosci. 2017;20: 927–933. doi:10.1038/nn.4575

838   27.   Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, et al. DeepLabCut:
839   markerless pose estimation of user-defined body parts with deep learning. Nat Neurosci.
840   2018;21: 1281–1289. doi:10.1038/s41593-018-0209-y

841   28.   Nath T, Mathis A, Chen AC, Patel A, Bethge M, Mathis MW. Using DeepLabCut for 3D
842   markerless pose estimation across species and behaviors. Nat Protoc. 2019;14: 2152–2176.
843   doi:10.1038/s41596-019-0176-0

844   29.   Villavicencio M, Moreno MG, Simon SA, Gutierrez R. Encoding of Sucrose's Palatability in
845   the Nucleus Accumbens Shell and Its Modulation by Exteroceptive Auditory Cues. Front
846   Neurosci. 2018;12: 265. doi:10.3389/fnins.2018.00265

847   30.   Berridge KC, Grill HJ. Alternating ingestive and aversive consummatory responses suggest a
848   two-dimensional analysis of palatability in rats. Behavioral neuroscience. 1983;97: 563–73.

31. Spector AC, Klumpp PA, Kaplan JM. Analytical issues in the evaluation of food deprivation
and sucrose concentration effects on the microstructure of licking behavior in the rat.
Behavioral neuroscience. 1998;112: 678–94.

32. Gutierrez R, Simon SA, Nicolelis MA. Licking-induced synchrony in the taste-reward circuit
improves cue discrimination during learning. The Journal of neuroscience : the official journal
of the Society for Neuroscience. 2010;30: 287–303. doi:10.1523/JNEUROSCI.0855-09.2010

33. Gutierrez R, Carmena JM, Nicolelis MAL, Simon SA. Orbitofrontal ensemble activity monitors
licking and distinguishes among natural rewards. J Neurophysiol. 2006;95: 119–133.
doi:10.1152/jn.00467.2005

34. Buonomano DV. Timing of neural responses in cortical organotypic slices. Proc Natl Acad Sci
USA. 2003;100: 4897–4902. doi:10.1073/pnas.0736909100

35. Glendinning JI, Chyou S, Lin I, Onishi M, Patel P, Zheng KH. Initial licking responses of mice
to sweeteners: effects of tas1r3 polymorphisms. Chem Senses. 2005;30: 601–614.
doi:10.1093/chemse/bji054

36. Zhao S, Ting JT, Atallah HE, Qiu L, Tan J, Gloss B, et al. Cell type-specific channelrhodopsin-
2 transgenic mice for optogenetic dissection of neural circuitry function. Nature methods.
2011;8: 745–52.

37. Gigante ED, Benalioaud F, Zamora-Olivencia V, Wise RA. Optogenetic Activation of a Lateral
Hypothalamic-Ventral Tegmental Drive-Reward Pathway. PLoS ONE. 2016;11: e0158885.
doi:10.1371/journal.pone.0158885

38. Norgren R. Gustatory responses in the hypothalamus. Brain Res. 1970;21: 63–77.
doi:10.1016/0006-8993(70)90021-1

39. Schwartzbaum JS. Electrophysiology of taste, feeding and reward in lateral hypothalamus of
rabbit. Physiol Behav. 1988;44: 507–526. doi:10.1016/0031-9384(88)90313-7

40. Grill, Berridge KC. Taste reactivity as a measure of the neural control of palatability. Progress
in Psychobiology and Physiological Psychology. Orlando, FL.: Academic Press; 1985. pp. 1–
54.

41. Sclafani A. The Hedonics of Sugar and Starch. The hedonics of taste. New York, NY.:
Lawrence Erlbaum Associates, Inc; 1991. pp. 59–87.

42. O'Connor EC, Kremer Y, Lefort S, Harada M, Pascoli V, Rohner C, et al. Accumbal D1R
Neurons Projecting to Lateral Hypothalamus Authorize Feeding. Neuron. 2015;88: 553–564.
doi:10.1016/j.neuron.2015.09.038

43. Nieh EH, Matthews GA, Allsop SA, Presbrey KN, Leppla CA, Wichmann R, et al. Decoding
neural circuits that control compulsive sucrose seeking. Cell. 2015;160: 528–541.
doi:10.1016/j.cell.2015.01.003

44. Burnett CJ, Li C, Webber E, Tsaousidou E, Xue SY, Brüning JC, et al. Hunger-Driven
Motivational State Competition. Neuron. 2016;92: 187–201. doi:10.1016/j.neuron.2016.08.032

45.   Chen Y, Lin Y-C, Zimmerman CA, Essner RA, Knight ZA. Hunger neurons drive feeding
through a sustained, positive reinforcement signal. Elife. 2016;5. doi:10.7554/eLife.18640

46.   Urstadt KR, Berridge KC. Optogenetic mapping of feeding and self-stimulation within the
lateral hypothalamus of the rat. PLoS ONE. 2020;15: e0224301.
doi:10.1371/journal.pone.0224301

47.   Sharpe MJ, Marchant NJ, Whitaker LR, Richie CT, Zhang YJ, Campbell EJ, et al. Lateral
Hypothalamic GABAergic Neurons Encode Reward Predictions that Are Relayed to the Ventral
Tegmental Area to Regulate Learning. Current Biology. 2017;27: 2089-2100.e5.
doi:10.1016/j.cub.2017.06.024

48.   Ono T, Nakamura K, Nishijo H, Fukuda M. Hypothalamic neuron involvement in integration of
reward, aversion, and cue signals. J Neurophysiol. 1986;56: 63–79. doi:10.1152/jn.1986.56.1.63

49.   Rolls ET. Taste, olfactory, and food reward value processing in the brain. Prog Neurobiol.
2015;127–128: 64–90. doi:10.1016/j.pneurobio.2015.03.002

50.   Karádi Z, Oomura Y, Nishino H, Scott TR, Lénárd L, Aou S. Responses of lateral hypothalamic
glucose-sensitive and glucose-insensitive neurons to chemical stimuli in behaving rhesus
monkeys. J Neurophysiol. 1992;67: 389–400. doi:10.1152/jn.1992.67.2.389

51.   Zhang X, van den Pol AN. Rapid binge-like eating and body weight gain driven by zona incerta
GABA neuron activation. Science. 2017;356: 853–859. doi:10.1126/science.aam7100

52.   DiFeliceantonio AG, Coppin G, Rigoux L, Edwin Thanarajah S, Dagher A, Tittgemeyer M, et
al. Supra-Additive Effects of Combining Fat and Carbohydrate on Food Reward. Cell Metab.
2018;28: 33-44.e3. doi:10.1016/j.cmet.2018.05.018

53.   Valenstein ES, Cox VC, Kakolewski JW. Modification of Motivated Behavior Elicited by
Electrical Stimulation of the Hypothalamus. Science. 1968;159: 1119–1121.
doi:10.1126/science.159.3819.1119

54.   Valenstein ES, Phillips AG. Stimulus-bound eating and deprivation from prior contact with
food pellets. Physiology & Behavior. 1970;5: 279–282. doi:10.1016/0031-9384(70)90099-5

55.   Valenstein ES, Cox VC. Influence of hunger, thirst, and previous experience in the test chamber
on stimulus-bound eating and drinking. J Comp Physiol Psychol. 1970;70: 189–199.
doi:10.1037/h0028708

56.   Fu O, Iwai Y, Narukawa M, Ishikawa AW, Ishii KK, Murata K, et al. Hypothalamic neuronal
circuits regulating hunger-induced taste modification. Nat Commun. 2019;10: 1–14.
doi:10.1038/s41467-019-12478-x

57.   Bonnavion P, Mickelsen LE, Fujita A, de Lecea L, Jackson AC. Hubs and spokes of the lateral
hypothalamus: cell types, circuits and behaviour. J Physiol (Lond). 2016;594: 6443–6462.
doi:10.1113/JP271946

58.   Mickelsen LE, Bolisetty M, Chimileski BR, Fujita A, Beltrami EJ, Costanzo JT, et al. Single-
cell transcriptomic analysis of the lateral hypothalamic area reveals molecularly distinct

populations of inhibitory and excitatory neurons. Nat Neurosci. 2019;22: 642–656.
doi:10.1038/s41593-019-0349-8

59. Leinninger GM, Jo Y-H, Leshan RL, Louis GW, Yang H, Barrera JG, et al. Leptin acts via
leptin receptor-expressing lateral hypothalamic neurons to modulate the mesolimbic dopamine
system and suppress feeding. Cell Metab. 2009;10: 89–98. doi:10.1016/j.cmet.2009.06.011

60. Qualls-Creekmore E, Yu S, Francois M, Hoang J, Huesing C, Bruce-Keller A, et al. Galanin-
Expressing GABA Neurons in the Lateral Hypothalamus Modulate Food Reward and
Noncompulsive Locomotion. J Neurosci. 2017;37: 6053–6065. doi:10.1523/JNEUROSCI.0155-
17.2017

61. Kurt G, Woodworth HL, Fowler S, Bugescu R, Leinninger GM. Activation of lateral
hypothalamic area neurotensin-expressing neurons promotes drinking. Neuropharmacology.
2019;154: 13–21. doi:10.1016/j.neuropharm.2018.09.038

62. Giardino WJ, Eban-Rothschild A, Christoffel DJ, Li S-B, Malenka RC, de Lecea L. Parallel
circuits from the bed nuclei of stria terminalis to the lateral hypothalamus drive opposing
emotional states. Nat Neurosci. 2018;21: 1084–1095. doi:10.1038/s41593-018-0198-x

63. Thoeni S, Loureiro M, O'Connor EC, Lüscher C. Depression of Accumbal to Lateral
Hypothalamic Synapses Gates Overeating. Neuron. 2020. doi:10.1016/j.neuron.2020.03.029

64. Zeng H, Madisen L. Mouse transgenic approaches in optogenetics. Prog Brain Res. 2012;196:
193–213. doi:10.1016/B978-0-444-59426-6.00010-0

65. Ting JT, Feng G. Development of transgenic animals for optogenetic manipulation of
mammalian nervous system function: progress and prospects for behavioral neuroscience.
Behav Brain Res. 2013;255: 3–18. doi:10.1016/j.bbr.2013.02.037

Figures

Figure 1. The 50 Hz stimulation evoked the largest LHA neuronal responses. **(A)** Representative location of an optrode in LHA of VGAT-ChR2 mice. **(B)** Confocal images from LHA immunostained for GFP (green), GAD67 (red), and DAPI (blue). Arrows indicate the colocalization of ChR2-EYFP and GAD67 (see merged). **(C)** Schematic of optrode recordings in LHA during control trials (with no-laser) and opto-stimulation at 2.5, 5, 10, 20, and 50 Hz delivered randomly. In each trial, the laser had a cycle of 2 s on and 4 s off. **(D)** Representative raster plots of two neurons recorded in LHA, the first one was activated (left panel), and the second inhibited (right panel) during opto-stimulation. Spiking responses were aligned (Time = 0 s) to the first laser pulse. Black ticks depict action potentials and

blue marks laser pulses. For visualization purposes, it is only shown laser pulses in the first trial for each trial type. Below are the corresponding PSTHs (firing rates, Sp/s). Red and blue ascending gradients indicate the frequency of stimulation (see E for the color bar). Black trace represents activity during control trials without laser. Vertical black lines indicate laser onset (Time = 0 s) and offset (Time = 2 s), respectively. **(E)** Normalized Z-score population responses of LHA neurons recorded from n = 5 VGAT-ChR2 mice. Red and blue colors depict activated or inhibited responses, respectively, relative to baseline (-1 to 0 s), thickness, and gradient colors indicate higher stimulation frequencies. Black trace illustrates neural activity during control trials for all recorded neurons (n = 186). **(F)** Percentage of neurons recruited as a function of stimulation frequency.

Figure 2. LHA neurons process palatability-related information. **(A)** Brief access taste and optotagging setup depicting a behavioral box equipped with a computer-controlled retractile sipper that for each lick independently delivers a drop of artificial saliva (AS), water, sucrose 3%, or 18% concentrations. After a head entry (dashed line), the first dry lick (black line) enables the 7 s reward epoch. During this period, in each lick (red), a drop of the four tastants was randomly delivered. After the reward epoch, the sipper was retracted for an Intertrial Interval (ITI) of 3 s and then re-extended to start a new trial. **(B)** Average lick rate during the entire reward epoch, reflecting greater palatability as a function of sucrose concentration. **(C)** Representative raster plot of two LHA neurons recorded while mice performed a brief access test. Spikes are plotted as black ticks, whereas licks for each tastant are in color-coded (green, AS; blue, water; orange, sucrose 3%; red, sucrose 18%). Below are corresponding peristimulus time histograms (PSTHs) of lick rate (Licks /s) and firing rate (Spikes/s; Sp/s) as a function of trial type. Neuronal responses were aligned (Time = 0 s) to the first rewarded lick. The brown rectangles depict the "best" window with the maximum Pearson correlation coefficient between firing rates and sucrose's palatability index (PI). The left and right raster plots displayed two neurons with a positive and negative correlation, respectively. **(D)** Z-score normalized activity (relative to AS trials) for LHA neurons with either positive or negative correlation against PI (red circles). **(E)** Population PSTHs of the lick rates given in the Reward epoch (Licks/7s) as a function of trial type, divided into blocks of 10th percentiles of trials each. **(F)** Population PSTHs of the firing rate during the Reward epoch (spikes/ 7s) of the palatability-related neurons across the session for each trial type. *Left panel* depicts neurons that fired more to higher sucrose concentrations (positive sucrose's palatability correlation; (D) yellow), whereas, *right panel* illustrates neurons with decreasing firing rates as the sucrose concentration increased (i.e., negative correlation with sucrose's palatability; (D) purple).

1014

Figure 3. A subpopulation of LHA^Vgat+ neurons encodes sucrose's palatability. **(A)** To confirm the LHA^Vgat+ neurons' identity after the brief access test, we performed laser opto-stimulation in a cycle 2 s on (at 50 Hz) and 10 s off over 15 min session. In this phase, mice had no access to the sipper (*above panel*), *below* the normalized Z-score population activity of all 284 LHA neurons recorded from 3 VGAT-ChR2 mice. Responses were aligned to the laser onset (Time = 0 s), relative to baseline (-1 to 0 s) activity. The red line depicts the responses of activated putative pLHA^Vgat neurons. **(B)** Percentage of laser-activated pLHA^Vgat+ neurons whose activity (in the brief access test) also positively or negatively correlates with sucrose's PI. * $p < 0.01$, chi-square. **(C)** Z-score activity (relative to AS) for all activated pLHA^Vgat neurons and its correlation with the sucrose's PI. **(D)** Percentage of Positive or Negative laser-activated palatability-related neurons. * $p < 0.0001$, chi-square. **(E)** PSTH of identified (opto-tagged) LHA^Vgat+ neurons exhibiting early evoked responses (i.e., ChR2 expressing cells). **(F)** Histogram of opto-tagged LHA^Vgat+ neurons with evoked early responses (latencies < 15 ms) from laser onset. **(G-I)** same conventions as in B-D, but for opto-tagged LHA^Vgat+ neurons encoding sucrose's palatability. **(J-M)** Non-LHA^Vgat neurons that negatively encode sucrose's palatability. *J,* Blue and black traces correspond to Inhibited (Non-LHA^Vgat neurons) and Unmodulated responses, respectively. Same conventions as in (*A*). **(K-M)** same conventions as in B-D, * $p < 0.001$, chi-square.

1031

Figure 4. In sated VGAT-ChR2 mice, 50 Hz stimulation of LHA^{Vgat+} neurons induced the greatest intake of sucrose. **A,** Behavioral setup for open-loop stimulation, where the laser was turned on regardless of the mice's behavior. The box was equipped with a central sipper where mice had free access to 10 % sucrose. Open-loop stimulation comprised 5 min block without laser followed by opto-stimulation for 5 min, in a cycle 2 s on – 4 s off, during a 40 min session. VGAT-ChR2 (n = 3) and WT (n = 5) mice received stimulation of only one of the following frequencies in ascending order per session 0, 2.5, 5, 10, 20, and 50 Hz. **B,** Opto-stimulation of LHA^{Vgat+} neurons induced sucrose (10%) intake in a frequency-dependent manner that peaked at 50 Hz. * Denotes statistically significant difference ($p < 0.001$) between no-laser and laser stimulated blocks at 50 Hz. Besides mice were sated, the lower intake at 0 Hz could be due to the lack of experience drinking in the novel box. **C,** Representative raster plot of open-loop stimulation from a VGAT-ChR2 mouse at 50 Hz. Lick responses (red ticks) were aligned to laser onset (blue rectangles, opto-stimulation period). The laser bound index was the number of licks within 2.5 s from laser onset (dashed blue rectangles) divided by total licks. **D,** The laser bound feeding index of VGAT-ChR2 increased as a function of the laser frequency. * Indicates significant difference from WT mice ($p < 0.01$), using a Two-way ANOVA followed by the Holm Sidak test. In this and other figures, data represent mean ± SEM.

1048

Figure 5. Open loop stimulation of LHA[Vgat+] neurons biases consumption towards the most palatable tastant and reinforce intake during opto-stimulation. **(A)** Behavioral protocol. Sated mice (n = 6 WT and n = 12 VGAT-ChR2) were located in an operant box equipped with three ports. The left port contained sucrose 18%, the central port water, and the right port sucrose 3%, counterbalanced across individuals. For details of open loop, see Figure 4A. **(B)** *Left and right panel*, number of licks for each gustatory stimulus for WT and VGAT-ChR2 mice. Note that the VGAT-ChR2 mice increased consumption of the most palatable stimuli available (sucrose 18%) mainly in opto-stimulated blocks (blue rectangles), while it decreased intake in blocks without laser. **(C)** Mean licks for sucrose 18%, during blocks with laser on and off from WT and VGAT-ChR2 mice. Black squares show the total licks given for each group. **(D)** Licks for sucrose 18% during blocks with or without laser across days. * Indicates a significant difference ($p < 0.0001$) between laser and no-laser blocks. # $p < 0.0001$ statistically significant difference between sucrose 18% and the other stimuli. ^ $p < 0.01$ between WT and VGAT-ChR2 for sucrose 3%; † $p < 0.01$ higher intake of water of VGAT-ChR2 compared with WT;+ p < 0.0001 higher consumption of VGAT-ChR2 compared with WT for sucrose 18% during blocks with laser. Using a two-way ANOVA and RM ANOVA followed by the Holm Sidak test.

1064

Figure 6. Licking probability increased as a function of distance. **(A)** Schematics for video tracking (see Materials and Methods). We calculated the distance and the head angle between the mouse's nose and each of the three sippers containing sucrose 3%, water, or sucrose 18% for each laser onset. **(B)** It

depicts the probability that laser onset evoked licking as a function of the mouse's distance relative to each sipper. **(C)** Polar plots depict the licking probability after laser onset, given distance and angle of the mouse relative to a licking sipper. At the onset of each laser stimulation, we computed the head angle and distance relative to each sipper, and, in the following 6 s window, we tally if the mouse licked to any sipper. We normalized by dividing the number of times licking was elicited by the number of first lasers occurring on those coordinates. The drawing of the mice's head illustrates that if a mouse was facing towards the sipper, the angle is 0°, in a perpendicular position (90°), or 180° if it was looking at the opposite direction. **(D)** The probability of licking after laser stimulation versus the angle between head direction and the sipper. * Indicates significantly different (p < 0.001) from all other distances. One-way ANOVA followed by the Holm Sidak test.

Figure 7. If stimulation occurred near the less palatable stimulus, LHA^{Vgat+} neurons induced water intake despite sucrose. **(A)** For the open loop stimulation, the laser was activated (2s on, 4s off) regardless of behavior and mice position. Same mice and conventions as in Figure 5*A*. In the closed-

loop configuration, the laser was triggered by a head entry in the central port. **(B)** The number of laser activations during the closed-loop protocol. The VGAT-ChR2 mice self-stimulate more than the WT mice. Each dot and triangle represent a single mouse. Unpaired t-test. **(C)** The total number of licks for each stimulus given during the entire session. In the open loop both mice groups licked more sucrose 18% than water or sucrose 3%. However, in the closed-loop configuration, the VGAT-ChR2 consumed more water (0 % sucrose) than the WT mice. **(D)** The number of licks evoked during 2.5 s after laser onset for each gustatory stimulus. **(E)** The number of licks when laser was turned off for each gustatory stimulus. **(F)** The water-sucrose 18% preference index from VGAT-ChR2 mice during photostimulation was defined as the number of licks of 18% sucrose divided by total licks for sucrose 18% + water. Thus, values higher than 0.5 indicate sucrose 18%, and values lower than 0.5 prefer water. Some VGAT-ChR2 mice preferred water over sucrose. The horizontal black lines indicate mean preference in both stimulation protocols. Paired t-test. **(G)** Fiber optic location in the LHA of VGAT-ChR2 mice. * $p < 0.0001$ Indicates significant difference from WT mice and stimulation protocols. # $p < 0.0001$ between sucrose 18% and the other stimuli. Two-way ANOVA followed by the Holm Sidak test.

Figure 8. After repeated activation of LHA^{Vgat+} neurons, laser-bound feeding (in open vs. closed-loop) strengthened and exhibited a positive correlation. **(A)** Scatter plots showing the relationship between

laser-bound feeding (licks given during the 2.5 s window from laser onset relative to total licks in the
session) in both open loop and closed-loop tasks during the first 3 days (l*eft panel*) and the last 5 days
of opto-stimulation of open loop vs. opto-self-stimulation of the closed-loop protocols (*right panel*).
(B) Likewise, after repeated activation, both opto-self-stimulations and laser-bound feeding (licking)
showed a significant correlation. Scatter depicting the average number of opto-self-stimulations vs. the
laser-bound feeding observed in the closed-loop protocol.

1107

Figure 9. Stimulation of LHA^{Vgat+} neurons is rewarding and biases approach towards the nearest
stimulus. **(A)** Different stimuli (plates containing cork, Chow, chocolate pellets, or a sipper filled with
10% liquid sucrose) were presented simultaneously in a circular arena. In this task, when mice cross a
designated area (dashed circles), the laser was turned on (2 s on; 4 s off) in a 40 min session. Thus, a
mouse had to leave and re-enter the designated area to receive a new opto-self-stimulation. Only one
designated area was used per session, and it remained in the same position for up to 3 or 4 consecutive

days. **(B)** Heat map and time spent from a representative WT (left panel) and VGAT-ChR2 mice (right panels), when the designated area was either the center, the pellet, or the Chow (see red circles for the currently designated zone). Colorbar indicates the number of frames the subject was detected in a given pixel; higher values indicate it remained in the same place for a longer time. *Below*, bar graphs depict the time spent in seconds exploring the designated area (within a radius of 5 cm). Each dot and triangle represent a single individual. Though mice normally avoid exploring the center of an open field, opto-self-stimulation in the center zone increased the time transgenic mice spent exploring it, compared with WT. Transgenic mice spent more time exploring the chocolate pellets and the Chow food when they were the designated zones. **(C)** Intake of chocolate pellet and Chow. Activation of LHA^{Vgat+} neurons in the center zone increased the consumption of chocolate pellets and chow compared with WT. *Inset*: Spill from chocolate pellets and chow. * Indicates statistically significant difference ($p < 0.05$) from WT. + p < 0.05 shows a significant increase in consumption and spill during the session that a designated area was opto-self-stimulated relative to the sessions where the center was opto-self-stimulated. Unpaired Student *t*-test. # $p < 0.01$ indicates significant difference between designated area opto-stimulated and the other open field areas. One-way ANOVA followed by the Holm Sidak test. See Video 5.

Figure 10. Activating LHA^{Vgat+} neurons does not induce the intake of a bitter tastant, nor an unavoidable aversive stimulus, but it increases the consumption of sucrose. **(A)** Schematic of behavioral setup showing stimuli delivered at central port. The water stimulus in the central port was replaced by an empty sipper, quinine, or airpuff. In all phases, head-entry in the central port triggered

the laser, except in the airpuff unavoidable condition, where the first lick delivered both the laser and the airpuff. Finally, the airpuff in the central port was replaced by sucrose 18%. **(B)** The number of opto-self-stimulations given for each stimulus. The VGAT-ChR2 mice performed more self-stimulations compared with WT, except during extinction sessions and when the airpuff was unavoidable. **(C)** The number of licks given to the central port. The licks in the central port decreased when an aversive stimulus was present, such as quinine or airpuffs. In contrast, a non-edibles stimulus such as an empty sipper elicited more licks from VGAT-ChR2 mice ($p < 0.05$). Moreover, when sucrose 18% was in the central port, VGAT-ChR2 group increased its consumption substantially compared with WT. **(D)** The number of licks in the lateral port containing sucrose 18% during opto-self-stimulation. The intake of the lateral port of sucrose 3 % is not shown because it was neglectable. Each dot and triangle represent a single individual. ^ Denotes statistically significant difference ($p < 0.0001$) from WT. Unpaired-Student t-test. * $p < 0.05$ relative to WT group. # $p < 0.05$ between sucrose 18% from other stimuli delivered at the central port. Two-way ANOVA followed by the Holm Sidak test.

Figure 11. Water deprivation gates LHA GABAergic neurons to increase the quinine intake. **(A)** Sated mice had free access to one sipper filled with quinine. The laser was turned on in closed-loop protocol (*left panel*). In sated VGAT-ChR2 mice, activation of these neurons did not increase quinine intake (*central panels*) but continued opto-self-stimulating (*right panel*). **(B)** Water deprived transgenic mice consumed more quinine than in WT mice (*central panel*) when the laser was turned on. **(C-D)** Total licks, number of licks during photostimulation, and when the laser was turned off in the session (*central panels*). Number of opto-self-stimulations of 18% sucrose during sated and water deprivation condition (*right panel*). * *p* < 0.05 indicates significant differences compared with WT mice according to an unpaired- t-student test.

Figure 12. Activation of LHA GABA neurons enhances water and sucrose's palatability. **(A)** Closed-loop stimulation during the brief access test. *Left panel,* behavioral setup. The box contained a sipper that delivered a ~2 μL drop of either water, sucrose 3%, or 18% per lick. *Right panel*, schematics of the structure of a trial. A head entry in the port (dashed vertical line) triggered opto-self-stimulation, at 50 Hz, in some sessions for 2 s and others for 7 s (blue rectangles). The reward epoch begins with the first dry lick on the sipper (black tick) and always last 7s (gray rectangles). During the reward epoch, a drop of tastant was delivered in each lick (red marks). At the end of it, the sipper was retracted, and an Inter Trial Interval (ITS) of 3 s began. **(B-E)** Data of closed-loop for 2s stimulation in the Reward epoch. PSTH of lick rate of WT (left panel) and VGAT-ChR2 mice (*right panel*) aligned (Time = 0 s) to laser onset. Opto-self-stimulation was delivered during all trials (B) or during only water trials (C, see arrow), or sucrose 3% (D), or 18 % (E). **(F-I)** Licking responses but during 7 s opto-self-stimulation. Same conventions as in B-E. The vertical blue line indicates laser onset and gray line the end of the reward epoch. The horizontal blue line indicates opto-self-stimulation window. * Indicates significant difference (*p* < 0.05) from WT group. # *p* < 0.05 among stimuli. *J,* Normalized Z-score

population activity of 284 neurons recorded in LHA in VGAT-ChR2 mice (n = 3). The laser was delivered during 7 s at 50 Hz. Neuronal responses were aligned to laser onset (Time = 0 s). Red and blue colors indicate activated or inhibited laser-evoked responses, relative to baseline (-1 to 0 s). Black trace illustrates neurons unmodulated by the laser. **(K)** Mock laser stimulation did not affect palatability responses. PSTH of licking responses, opto-self-stimulation was delivered along with sucrose 3% trials during 7s (arrow). Note that both groups increased the lick rate as a function of palatability, same conventions as in B-E. To elicit mock laser stimulation, mice were connected to a mock fiber optic (with no-laser connected), whereas the real fiber was glued outside the mice's head to emit blue light.

Supplementary Material

1204

Supplementary Figure 1. Waveform stability in extracellular recordings of LHA neurons across the
brief-access test (BAT) and the optotagging session. **(A)** Representative raster plot of a LHA^{Vgat+}
neuron recorded during the optotagging session. Spiking responses were aligned (Time = 0 s) to the
first laser pulse. Black ticks depict action potentials and blue marks on top, laser pulses (at 50 Hz). For
visualization purposes, it is only shown laser pulses in the first trial. Below, PSTHs (Spikes /s, Sp/s).
Vertical black lines indicate laser onset (Time = 0 s) and offset (Time = 2 s), respectively. **(B)** *Inset*
upper left depicts average waveforms of a single neuron showed in A across the BAT (black), the
optotagging stimulation (red), and when the laser was turned "on" (blue). Right, each dot represents an
action potential during the entire session (same color-coded). Values are plotted in a three-dimensional
space computed with the first three principal components (PCs) of the waveforms. **(C-D)** Same
conventions as in (A-B). ***E,*** Histogram of the Pearson correlation coefficients of waveforms recorded
during BAT against those recorded in the optotagging session. Waveforms depict two examples, one
depicting the worst and the other the best correlation included in analysis.

1218

1219

1220

Supplementary Figure 2. Pre-stimulation of LHA^{VGAT+} neurons did not trigger a "hunger state" in that a sustained consummatory behavior is not observed. **(A)** Schematic of the pre-stimulation protocol and raster plot from VGAT-ChR2 (licks were plotted as red ticks) and WT (licks as light red ticks) mice. For this experiment, all mice underwent all pre-stimulation protocols following a Latin square design: 1) mice were pre-stimulated for either 0, 5, 10, or 15 min with no sucrose available. After pre-stimulation, mice received 15 min Free access to a sipper filled with sucrose 10%. **(B)** The average number of licks for sucrose as a function of pre-stimulation.

1232

Supplementary Figure 3. Activation of LHA$^{\text{Vgat+}}$ neurons increases the time spent near the most palatable High Fat Diet (HFD). **(A)** In a circular arena, three plates containing Chow (green), HFD (yellow), and granulated sucrose cube (blue) were available simultaneously. In a 20 min session, the laser was turned "on" following an open loop protocol (2 s on 4 s off during 5 min block with laser and 5 min with no-laser; see Figure 5A). **(B)** Average heat map and time spent in the open field. To create each heat map, we identified the centroid of the mouse at every video frame. Colorbar indicates the number of frames the subject was detected in a given pixel; higher values indicate the mice stayed in that place for a longer time. Time spent (below) was calculated as the distance between the mouse's centroid relative to each food plate. We quantify a food interaction when the subject was within 60 mm distance relative to the plate's center and remained there for at least 1 s. Each dot and triangle represent a single individual. **(C)** Intake of Chow, HFD, and granulated sucrose cube. * Denote statistically

1244   significant differences ($p < 0.0001$) relative to WT. # $p < 0.0001$ between a stimulus relative to the
1245   others. One-way ANOVA followed by the Holm Sidak test. See Video 2.

1246

1247

1248   **Supplementary Figure 4.** Activation of LHA^Vgat+ neurons promotes intake to the most palatable solid
1249   food. **(A)** In a circular arena, Chow pellet and granulated sucrose cube were presented simultaneously.
1250   The laser was turned "on" in an open loop protocol. **(B)** Mean heat map and time spent. The same
1251   conventions are in Supplementary Figure 4. **(C)** Intake of Chow and granulated sucrose cube. *

Statistically significant differences ($p < 0.001$) relative to WT. # $p < 0.0001$ difference between a stimulus relative to the others. One-way ANOVA followed by the Holm Sidak test. See Video 3.

Video 1. Open loop activation of LHA^{Vgat+} neurons. Water, 3 wt%, and 18 wt% sucrose was delivered in each port. Opto-stimulation occurred in 5 min block of no-laser and 5 min block with the laser.

Video 2. Open loop activation of LHA^{Vgat+} neurons promotes preference toward HFD over granulated sucrose and Chow pellet. At equidistant position, three foods were placed on a small plate, and each contained either HFD, Chow food, or granulated sucrose cube. Opto-stimulation occurred in 5 min block of no-laser and 5 min block of the laser for 20 min.

Video 3. VGAT-ChR2 mice prefer granulated sucrose over Chow food. Preference and intake of granulated sucrose cube vs. Chow pellet after open-loop activation of LHA GABAergic neurons.

Video 4. Closed-loop activation of LHA^{Vgat+} neurons. Mice opto-self-stimulate in each head entry in the central licking port (where water is the nearest stimulus). Some VGAT-ChR2 mice exhibited a preference for the most palatable stimulus available (sucrose 18%), whereas others preferred the nearest but less palatable stimulus (water) over sucrose 18%

Video 5. Real-time place preference in an open field task. At equidistant separation, four stimuli were placed on a small plate, and each contained either wood cork, Chow pellet, chocolate pellets, or a sipper a 10% sucrose solution. Mice were photostimulated by 2 s on and 4s off when crossing the designated area.

Video 6. In some VGAT-ChR2 mice, activation of LHA^{Vgat+} neurons induces gnawing to a cork. Wood cork was placed in a transparent box, whereas mice were opto-stimulated in an open loop protocol.