1 **EukProt: a database of genome-scale predicted proteins across the diversity of**
2 **eukaryotes**
3

4 Daniel J. Richter[1], Cédric Berney[2,3], Jürgen F. H. Strassert[4], Yu-Ping Poh[5], Emily K.
5 Herman[6], Sergio A. Muñoz-Gómez[7], Jeremy G. Wideman[5], Fabien Burki[8], Colomban de
6 Vargas[2,3]
7

1. Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra), Passeig Marítim de la Barceloneta 37-49,
08003 Barcelona, Spain
2. Sorbonne Université, CNRS, Station Biologique de Roscoff, UMR 7144, ECOMAP, 29680 Roscoff, France
3. Research Federation for the study of Global Ocean Systems Ecology and Evolution, FR2022/Tara GOSEE,
Paris, France
4. Department of Evolutionary and Integrative Ecology, Leibniz Institute of Freshwater Ecology and Inland
Fisheries (IGB), Müggelseedamm 301, 12587 Berlin, Germany
5. Biodesign Center for Mechanisms of Evolution, School of Life Sciences, Arizona State University, Tempe,
Arizona, United States of America
6. Department of Agricultural, Food and Nutritional Sciences, Faculty of Agricultural, Life, and Environmental
Sciences, University of Alberta, Edmonton, Alberta, Canada
7. Unité d'Ecologie, Systématique et Evolution, Université Paris-Saclay, Orsay, France
8. Department of Organismal Biology and Science for Life Laboratory, Uppsala University, Norbyvägen 18D,
Uppsala 75236, Sweden
22

23 **Database Availability**
24

28 **Abstract**
29

EukProt is a database of published and publicly available predicted protein sets selected to represent
the breadth of eukaryotic diversity, currently including 993 species from all major supergroups as well
as orphan taxa. The goal of the database is to provide a single, convenient resource for gene-based
research across the spectrum of eukaryotic life, such as phylogenomics and gene family evolution.
Each species is placed within the UniEuk taxonomic framework in order to facilitate downstream
analyses, and each data set is associated with a unique, persistent identifier to facilitate comparison
and replication among analyses. The database is regularly updated, and all versions will be
permanently stored and made available via FigShare. The current version has a number of updates,
notably 'The Comparative Set' (TCS), a reduced taxonomic set with high estimated completeness
while maintaining a substantial phylogenetic breadth, which comprises 196 predicted proteomes. We
invite the community to provide suggestions for new data sets and new annotation features to be
included in subsequent versions, with the goal of building a collaborative resource that will promote
research to understand eukaryotic diversity and diversification.
43

44 **Introduction**
45

Over the past 15 years, the discovery of diverse novel microbial eukaryotes, coupled with methods to
reconstruct phylogenies based on hundreds of protein-coding genes (known as phylogenomics
(Eisen, 2003)) have led to a remarkable reshaping in our understanding of the eukaryotic tree of life,
the proposal of new supergroups and the placement of enigmatic lineages in known supergroups
(Brown et al., 2018; Burki et al., 2012, 2020; Gawryluk et al., 2019; Janouškovec et al., 2017;
Kamikawa et al., 2014; Lax et al., 2018; Strassert et al., 2019; Tice et al., 2021; Yabuki et al., 2015).
The phylogenomic approach has also been used to investigate branching patterns within eukaryotic
supergroups, including to understand species evolutionary relationships with implications for the early

evolutionary events in the three best-studied eukaryotic lineages: land plants (Wickett et al., 2014),
animals (King & Rokas, 2017) and fungi (Kiss et al., 2019; Y. Li et al., 2021; Strassert & Monaghan,
2021). Furthermore, analyses in diverse eukaryotes have the potential to reveal new genes and
pathways for biological processes currently characterized only in these three well-studied lineages
and their parasites (del Campo et al., 2014; Richter & Levin, 2019). In the ocean alone, planetary-
scale metagenomics studies (Bork et al., 2015; Tara Oceans Coordinators et al., 2020) have already
unveiled an extreme diversity of eukaryotic genes (Carradec et al., 2018) whose phylogenetic origin
and ecological function are mostly unknown.

A critical prerequisite to all genomic studies is the underlying database of predicted proteins from
which orthologs are extracted or other sequence analyses are performed. Because no single source
contains the protein predictions for the complete set of eukaryotes that have been sequenced at a
genomic scale, each study needs to assemble their own database, reducing reproducibility among
analyses (due to the inclusion of different taxa, or different data sets for the same taxon), and
producing a significant barrier to new researchers entering the field (we note that an existing database
of genome-scale protein data sets, PhyloDB (https://github.com/allenlab/PhyloDB), contains 550
eukaryotic species with at least 500 proteins, as of version 1.076.) In addition, because the large
majority of the protein data sets from diverse eukaryotes are not included in major public databases
(see Table 1), researchers cannot easily access them via standard tools such as NCBI BLAST
(Sayers et al., 2020) in order to find the homologs of a protein of interest, nor are valuable annotations
such as protein domains (e.g., Pfam (El-Gebali et al., 2019), Interpro (Mitchell et al., 2019)) or gene
ontology (The Gene Ontology Consortium, 2019) easily accessible for searching.

To address this gap, we assembled a database of protein sequences – EukProt – either from newly
performed protein prediction or by retrieving available sequences from a comprehensive set of
species representing known eukaryotic diversity. We placed each species within a universal
eukaryotic taxonomic framework, UniEuk (Berney et al., 2017), in order to ensure that the
evolutionary relationships among data sets are accurately and consistently described. We also
provide estimated completion statistics for each predicted proteome based on BUSCO scores and
have selected a suggested subset of 196 proteomes for use in comparative genomics investigations.
EukProt is designed to prioritize ease of use, with unique, persistent identifiers assigned to each data
set and a standard system of nomenclature to facilitate repeatability of analyses. To this end, we have
followed FAIR principles (Findable, Accessible, Interoperable, Reusable) in its construction (Wilkinson
et al., 2016).

The EukProt Database

EukProt (currently in version 3; 22 November, 2021) contains 993 eukaryotic species from 5 different
sequence data-types (Figure 1): genome (375 species), single-cell genome (56), transcriptome (498),
single-cell transcriptome (47), and expressed sequence tag (EST; 17). The data sets were
downloaded from 37 different sources (Table 1), with the two principal sources being NCBI (Sayers et
al., 2020) and the Moore Microbial Eukaryote Transcriptome Sequencing Project (MMETSP) (Keeling
et al., 2014). The database contains broad taxonomic representation, with species from 34 of 36
major deeply-branching eukaryotic lineages defined in the UniEuk database (Berney et al., 2017).
Within these major groups, however, the representation remains uneven, which results from several
factors: the difficulty of discovering, culturing or sequencing species from many lineages; a bias
towards sequencing either macroscopic, multicellular species or unicellular species that are parasites
(e.g., Apicomplexa), photosynthetic (such as diatoms, which are part of the Ochrophyte lineage), or
are otherwise economically important (del Campo et al., 2014); and very uneven evolutionary
histories among groups, with some having undergone major diversification with many different
morphologies, and others having diversified to a lesser extent (or with only a few surviving lineages).
105

106 We have implemented in version 3 several changes proposed by the community of EukProt database
107 users after version 2 was released: we have included (i) the 18S ribosomal DNA sequence
108 corresponding to each data set (when available), (ii) a measurement of the completeness of the data
109 set as estimated by BUSCO (Manni et al., 2021; Simão et al., 2015), and (iii) the assignment of
110 unique EukProt identifiers to individual protein sequences.
111
112 The EukProt database is organized around 5 guiding principles:
113
Breadth of species phylogenetic diversity: the objective of the database is to include as many
lineages as possible from the broad diversity of eukaryotes. For the majority of lineages, at least one
representative from all sequenced genera is included. The exceptions are lineages with hundreds of
sequenced representatives, or lineages whose sequenced species are relatively closely related, for
which we included only a subset of species selected to represent phylogenetic diversity in order to
balance the overall size of the database versus broad representation of eukaryotes. These well-
sequenced lineages are Chloroplastida (containing the land plants and green algae), Metazoa
(animals), Amoebozoa, Fungi, and some parasites/pathogens (members of Apicomplexa,
Peronosporomycetes, Metakinetoplastina and Fornicata). Within animals, we also emphasized the
inclusion of marine planktonic species, as we anticipate these could serve as mapping targets for data
from large-scale metagenomic and metatranscriptomic ocean sequencing projects (e.g., *Tara* Oceans
(Carradec et al., 2018; Karsenti et al., 2011; Richter et al., 2019), Malaspinas (Acinas et al., 2021;
Duarte, 2015), Bio-GO-SHIP (Larkin et al., 2021) and GEOTRACES (Biller et al., 2018)).
127
Convenience of access: the full data set and its associated metadata can easily be downloaded
from FigShare. For 624 of the total 993 species in the database, publicly available protein sequences
were ready to be included directly. For the remaining 369 species, additional processing steps were
required to produce protein sequences for inclusion in the database. The most common
bioinformatics processing steps were *de novo* assembling and translating transcriptomes from raw
read data (120), translating mRNA sequences from publicly available assembled transcriptomes or
EST projects lacking protein predictions (102), merging independent sets of protein predictions for the
same species (85 species) and, newly added in version 3 of the database, producing *de novo* gene
predictions for unannotated genomes (40). We also provide transcriptome assemblies and GFF
annotation files for the latter two classes, as they are not publicly available. A full list of the different
types of actions is described in the Methods, and the actions taken for each species are available in
the database metadata.
140
A unified taxonomic framework: we placed all species into the taxonomic framework generated in
the UniEuk project (Berney et al., 2017), which is based on the most recent consensus eukaryotic
classification (Adl et al., 2019) and phylogenomic evidence. This will maximize interoperability of the
database with other resources built upon to the UniEuk taxonomic framework (e.g., EukBank and
EukMap (Berney et al., 2017), and EukRibo (Berney, Cédric, 2022)), and ensure that results arising
from EukProt can use a common set of identifiers to describe analyses at different taxonomic levels.
For example, this might include labeling groups in a phylogenetic tree, or summarizing read
placement data by taxonomic group. Knowledge of the taxonomy can also facilitate the detection of
mis-identified gene sequences within a data set, via comparison of the topology of gene trees to the
taxonomy. To assist in species identification and to provide a convenient link to EukRibo (a manually
curated database of 18S rDNA sequences, part of UniEuk, aimed at taxonomic annotation of
metabarcoding datasets), we also supply in version 3 the 18S sequence corresponding to the
sequenced strain (or, if an 18S for the sequenced strain is not available, an 18S sequence from the
same species; see Methods). In addition to the complete taxonomic pathway containing all
intermediate nodes, we provide for each species three sequentially nested groupings ("taxogroup2"
within "taxogroup1" within "supergroup"; see Methods) that can help to judge the redundancy of data
sets, or conversely their lack of representation, within different groupings. The taxogroup1 level is

used as the leaves in Figure 1. Finally, for each data set we include a list of previous species
identifications, if any, to prevent issues resulting from revisions to species names or lineages and
corrections of species misidentifications, which have occurred frequently with improvements in
sequencing and phylogenetic techniques and the discovery of new eukaryotic organisms.
162
Appropriate references to publications and data sources: to ensure that the researchers who
generated and provided each data set receive appropriate credit, we provide the DOI of the
publication describing each data set as well as the URL from which it was downloaded. The list of
URLs should also allow users of the database to download the original sequences for each data set
(although it is not possible to guarantee that the URLs for all data providers will be permanently
available).
169
Reusability, persistence and replicability: the database will be released in successive versions,
each of which will be permanently stored and accessible at FigShare. Thus, analyses using the
database will need only to specify which version was used, enabling follow-up analyses or replications
to begin with the identical database. In addition, each individual data set within the database is
assigned a unique, permanent identifier. When a new data set becomes available for a given species,
it is assigned a new unique identifier (and the identifier of the data set it replaces, if any, is indicated in
the database metadata). These and all other changes between versions of the database are recorded
as appropriate.
178
179 **'The Comparative Set' (TCS): A selected subset of EukProt for comparative genomics**
180 **investigations**
181
Comparative genomics investigations are crucial to bridge the gaps between cell biology and
biochemistry on one hand and diversity and evolution on the other. Classically, the set of species to
include has been chosen by the authors of each study. With the goal of providing some consistency in
future comparative genomics studies, we have selected a set of predicted proteomes with high
completeness to represent the broad phylogeny of eukaryotes. A total of 196 predicted proteomes
were selected as 'The Comparative Set' (TCS), abbreviated in honor of the late Tom Cavalier-Smith
(Bass, 2021; Richards, 2021; Roger, 2021; Saldarriaga, 2021). All major model systems were
included in this set as well as proteomes derived from high-quality genomes. Additional proteomes
were chosen to maximize phylogenetic diversity and based on their completeness as estimated by
BUSCO (Manni et al., 2021) scores. Updates to the TCS will be based on availability of new
proteomes from phylogenetically important lineages currently underrepresented, and on community
feedback. Please send any comments or suggestions for the TCS to Jeremy.Wideman@asu.edu. The
complete EukProt database and the TCS are available for download and also available for BLAST
searches at http://evocellbio.com/eukprot/ using Sequenceserver (Priyam et al., 2019).
196
197 **Growing the EukProt Database with Community Involvement**
198
The core functionality of the database is the distribution of genome-scale protein sequences across
the diversity of eukaryotic life and within the UniEuk framework. However, we anticipate that
numerous other features might be useful to the community, and we hope to involve the community in
suggesting and adding new features in successive versions. These may include information or
analyses on full data sets, such as the sequencing technology that was used (e.g., Illumina, PacBio),
or the estimation of potential contamination levels (e.g. from mixed cultures or prey species) with non-
target species, as inferred using systematic sequence homology searches. Additionally, we could
consider adding features on the level of individual protein sequences, such as protein domains from
Pfam (El-Gebali et al., 2019)/Interpro (Mitchell et al., 2019), gene ontology
(The Gene Ontology Consortium, 2019)/eggNOG (Huerta-Cepas et al., 2019).
209

210 As new genome-scale eukaryotic protein data sets become available, we plan to add them to the
211 database. As yet, we do not have a formal mechanism to accomplish this, and will instead depend on
212 monitoring the literature and assistance from the community. As an example, for version 3, we relied
213 on PhycoCosm (Grigoriev et al., 2021), MycoCosm (Grigoriev et al., 2014) and PhyloFisher (Tice et
214 al., 2021) to add data sets that became recently available.
215
216 In the longer term, we hope the standardization of our database provides a path towards including all
217 data sets in a major sequence repository such as NCBI/EBI/DDBJ, so that they can be more broadly
218 accessible and integrated into the suites of tools available at these repositories.
219
220
221 **Methods**
222
223 *Species and strain identity*
224
We determined species and strain identities by reading the publications that described the data sets,
consulting the literature for naming revisions, and comparing 18S ribosomal DNA sequences for each
data set to reference sequence databases (see below for a description of how we retrieved 18S
sequences). For species that were previously known by other names, we recorded these previous
names in the metadata for the data set, except in cases where a species was originally assigned to a
genus but not identified to the species level (e.g., *Goniomonas* sp., now identified as *Goniomonas*
avonlea, is not listed as a previous name). This exception was not followed when the GenBank record
description for the 18S sequence contained a different name than the one used in EukProt; in this
case this previous name is always listed in the previous names field, in order to avoid confusion when
retrieving 18S sequences from GenBank.
235
236 When no strain name was available for an MMETSP data set, we used the MMETSP ID as the strain
237 name (for example, EP00362 has the strain name MMETSP1317). For strains with alternative names,
238 we included these in a dedicated field for alternative strain names; this list is not necessarily
239 exhaustive (i.e., it is not guaranteed to contain all alternative names for a given strain).
240
241 *Supergroups and taxogroups from UniEuk*
242
The full taxonomic pathways provided for all species (which follow the framework developed in the
UniEuk project (Berney et al., 2017)) are not based on a fixed number of ranks, but on a free,
unlimited number of taxonomic levels, in order to match phylogenetic evidence as closely as possible.
This provides end-users more information and flexibility, but could also make it more difficult to
summarize results of downstream analyses. Therefore, we provide three additional fields
("supergroup", "taxogroup1" and "taxogroup2") to help end-users whenever it is useful to distribute
eukaryotic diversity into a fixed number of taxonomic categories of roughly equivalent phylogenetic
depth or ecological relevance. The groupings called "supergroups'' in the context of UniEuk resources
(36 recognized lineages so far, of which 34 are included in EukProt; *Microheliella* and *Meteora* are not
represented) consist of strictly monophyletic, deep-branching eukaryotic lineages of a phylogenetic
depth equivalent to some of the first proposed supergroups such as Opisthokonta, Alveolata,
Rhizaria, and Stramenopiles. UniEuk "supergroups" correspond roughly to the deepest phylogenetic
resolution of eukaryotic relationships achievable with ribosomal RNA genes, and not necessarily to
the currently recognised highest-level groupings of eukaryotes based on phylogenomic evidence
(these groupings are, of course, present in the complete taxonomic pathways). UniEuk "supergroups"
are therefore highly variable in relative diversity, ranging from lineages consisting of a single, orphan
genus (e.g. *Ancoracysta*, *Mantamonas*, *Palpitomonas*), to Opisthokonta as a whole. The "taxogroup1"
(of which there are 72 in EukProt) and "taxogroup2" (of which there are 198 in EukProt) levels allow
further subdivision of large supergroups into lineages of relatively equivalent evolutionary or

ecological relevance, based on current knowledge. These levels are more arbitrarily defined but are intended to represent strictly monophyletic groupings that match one of the levels in the complete taxonomic pathways. As an illustrative example, diatoms are in the Diatomeae taxogroup2, which is within the Ochrophyta taxogroup1, which is within the Stramenopiles supergroup. Small, ecologically and morphologically homogeneous supergroups are not subdivided further; in such cases the "taxogroup1" and "taxogroup2" levels are the same as the "supergroup" level. The same approach will be used in EukRibo, a manually-curated database of reference ribosomal RNA gene sequences developed in parallel (Berney, Cédric, 2022) to help users link analyses of different types of genetic data.

Merging strains from the same species

In general, we only included data from a single strain/isolate per species. However, when only a single transcriptome data set was available for a given strain of a species, and there were additional published transcriptome data sets for other strains of the same species, we combined them using CD-HIT (W. Li & Godzik, 2006) run with default parameter values, in order to guard against the possibility that a single transcriptome might lack genes expressed only in one condition or experiment. When multiple strains were merged to produce a species' data set (there were 41 such cases), this information is indicated in the metadata for the data set. When a data set is indicated as merged, but more than one strain name is not listed in the strain column, this indicates that the data set was distributed as merged and we were unable to determine the exact strains used.

Processing steps applied to publicly available data

All sequences within each FASTA file are assigned a unique, standardized identifier based on the data set's EukProt ID and on the type of data (protein or transcriptome); this identifier is prepended to the existing FASTA header, separated by a space. Illegal characters are removed from sequences. The following characters are permitted, as defined by NCBI BLAST (https://blast.ncbi.nlm.nih.gov/Blast.cgi?CMD=Web&PAGE_TYPE=BlastDocs&DOC_TYPE=BlastHelp) for nucleic acid sequences: `ACGTNUKSYMWRBDHV` and for protein sequences: `ABCDEFGHIKLMNPQRSTUVWYZX*`.

EukProt metadata indicate the additional steps applied to each data set after downloading from the data source (if any).

All software parameter values were default, unless otherwise specified below or in the metadata record for a given data set.

'assemble mRNA': *de novo* transcriptome assembly using Trinity v. 2.8.4, http://trinityrnaseq.github.io/ (Haas et al., 2013). We trimmed Illumina input reads for adapters and sequence quality using the built-in '--trimmomatic' option. We trimmed 454 input reads prior to running Trinity with Trimmomatic v. 0.3.9, http://www.usadellab.org/cms/?page=trimmomatic (Bolger et al., 2014) with the directives 'ILLUMINACLIP:[454 adapters FASTA file]:2:30:10 SLIDINGWINDOW:4:5 LEADING:5 TRAILING:5 MINLEN:25'. When the sequence library was described as stranded in the NCBI Sequence Read Archive, we used the corresponding 'SS_lib_type' option.

'translate mRNA': *de novo* translation of mRNA sequences with Transdecoder v. 5.3.0, http://transdecoder.github.io/. When the number of predicted protein sequences for a given species was both less than half of the input mRNA sequences and less than 15,000, we reduced the minimum predicted protein length to 50 (from the default of 100).

313 'CD-HIT': clustering of protein sequences to produce a non-redundant data set using CD-HIT v. 4.6,
314 http://weizhongli-lab.org/cd-hit/ (W. Li & Godzik, 2006). We used this tool principally to combine
315 protein predictions for different strains of the same species, but also to reduce the size of very large
316 predicted protein sets (>50,000 proteins) that showed evidence of redundancy.
317
'extractfeat', 'seqret', 'transeq', and 'trimseq': from the EMBOSS package v. 6.6.0.0,
http://emboss.sourceforge.net/ (Rice et al., 2000). We used extractfeat to produce coding sequences
(CDS) from genomes with gene annotations in EMBL format but without publicly available protein
sequences. We used seqret to convert FASTQ files to FASTA files. We used transeq to translate
CDS directly into proteins. We used trimseq to trim EST sequences before translation with
Transdecoder.
324
325 'gffread': to produce protein sequences from genomes with gene annotations in GFF format but
326 without publicly available protein sequences, using the command line option '-y' in version 0.12.3
327 (Pertea & Pertea, 2020).
328
'predict genes': we used EukMetaSanity https://github.com/cjneely10/EukMetaSanity (Neely et al.,
2021) to perform automated annotation of genome sequences lacking publicly available protein
predictions. We used the following parameters, as specified in (Alexander et al., 2021): --min_contig
500 --min_contig_in_predict 500 --max_contig 100000000. We used the parameter --
min_contig_in_predict 200 (as it matched the default minimum contig length in Trinity). By default, we
selected the proteins at Tier 2 (predictions supported by at least 2 sources). If Tier 2 produced fewer
than 15,000 predicted proteins, we instead selected Tier 1. All other parameter values were left at
their defaults. We did not perform gene prediction on unannotated genomes for which a transcriptome
was already available for the same species (under the assumption that the gene predictions of the
transcriptome would be of higher quality, due to potential errors in the gene annotation process).
339
340 *Retrieval of 18S sequences*
341
342 We searched for 18S sequences of at least 1,500 base pairs in length for each data set in the
343 following order. If no sequence, or only a partial sequence was retrieved in a given step, we moved on
344 to the next step. Details specific to each data set are included in the EukProt metadata.
345
1.   Search GenBank (Sayers et al., 2020) for an 18S sequence for the species and strain. When
more than one sequence was available for the strain, we chose the one containing the largest
quantity of high-quality nucleotides (after identifying and eliminating sequencing errors,
chimeras, etc.).
2.   If the data set was sequenced as part of the MMETSP, use the 18S sequence from MMETSP
(if it was provided in the project metadata, and also was not already present in GenBank).
3.   Search GenBank for an 18S sequence for another strain of the same species.
4.   Using the 18S sequence for the most closely related species in GenBank as a query, search
the contigs associated with the data set (either genome or transcriptome) using blastn
(Altschul, 1997).
5.   Assemble the 18S sequence from the raw sequence reads associated with the data set,
using phyloFlash version 3.4 (Gruber-Vodicka et al., 2020) with the SILVA database version
138.1 (Quast et al., 2012) and SPAdes assembler version 3.14.1 (Prjibelski et al., 2020), with
default parameter values except for the number of reads used, and with read length set to
match the read length of the data set. For each data set, we initially ran phyloFlash with all
sequence reads. If we observed that there was evidence for reads mapping to the 18S of the
target lineage, but that did not result in a contig assembled by SPAdes, we ran phyloFlash
again with only the first 10 million reads. If this run showed a similar result (reads mapping but
no contigs), we ran again with only the first 1 million reads. When all three runs (all reads, first

10 million reads, first 1 million reads) did not result in a SPAdes assembly, we selected the
reads mapping to the 18S of the target lineage and performed an assembly using CAP3
version date 02/10/15 (Huang, 1999) with 'relaxed' parameter values from (Close et al.,
2009): -p 75 -d 200 -f 250 -h 90.

A selected subset of EukProt for phylogenetic studies

Single-copy orthologs were identified in EukProt v3 proteomes using BUSCO v5.2.2 (Manni et al.,
2021) with the Eukaryota Odb10 database. BUSCO searches were run in protein mode with the
default E-value cutoff of 1e-3. These BUSCO scores were visualized using Plotly (Plotly Technologies
Inc., 2015) and are available at http://evocellbio.com/eukprot/. Single-copy BUSCO markers for taxa
with BUSCO scores >30 were aligned to single-copy BUSCO profile HMMs with hmmalign (HMMER
v3.3; (Eddy, 2020)), trimmed with ClipKit (-m gappy; (Steenwyk et al., 2020)), and concatenated into a
supermatrix with FASConCat-G v1.04 (Kück & Longo, 2014). A guide tree was first inferred from this
supermatrix with IQ-TREE v2.0.3 using the LG4X model (-fast mode; (Minh et al., 2020)). A second
phylogenetic tree was inferred using the LG+PMSF(C10)+F+G4 model and using the LG4X tree as a
guide (Wang et al., 2018). A first subset of taxa was automatically selected from the previously
inferred tree to maximize phylogenetic diversity using Treemmer v0.1 (Menardo et al., 2018). A few
taxa that were phylogenetically misplaced with high support, likely due to a high percentage of
contaminating sequences, were visually detected and manually removed. Furthermore, predicted
proteins from model organisms (e.g., *Homo sapiens*, *Saccharomyces cerevisiae*, *Toxoplasma gondii*,
Dictyostelium discoideum, etc.) were included in the TCS. In addition, phylogenetically important taxa
with sufficiently high BUSCO scores were also included. Since different lineages universally lack
particular BUSCOs, a singular score cut-off is not reasonable and therefore each taxon group was
assessed in relation to a model system with a trusted genome. For example, while *Giardia intestinalis*
and *Toxoplasma gondii* have low BUSCO scores (23.5 and 61.2, respectively), they are generally
accepted as complete genomes. Thus, species in these lineages (e.g., Fornicata, Preaxostyla,
Apicomplexa, Parabasalia, etc.) with comparable BUSCO scores were included in the TCS. Finally, in
taxonomically important lineages in which no model yet exists (e.g., Ancyromonads) proteome sets
with BUSCO scores ~50% or greater were included, or if no such representative was available, it was
excluded from the TCS (e.g., Tsukubomonads). We welcome suggestions from the community for
future iterations of the TCS.

The EukProt database distribution on FigShare

The database is distributed in a single archive containing five files. One file contains 993 protein data
sets, for species with either a genome (375), a single-cell genome (56), a transcriptome (498), a
single-cell transcriptome (47), or an EST assembly (17). A second file contains assembled
transcriptome contigs, for 126 species with publicly available mRNA sequence reads but no publicly
available assembly. The proteins predicted from these assemblies are included in the proteins file. A
third file contains GFF annotations for 40 genomes lacking publicly available predicted protein
annotations. Finally, the database metadata are distributed as two files: one file for the data sets
included in the current version of the database (993), and a second file for data sets not included
(163), accompanied by the reason they were not included. For example, when a data set cleaned with
a genome-scale decontamination process is described in a manuscript, but the decontaminated data
are not publicly available, it is not included in the database.

Acknowledgements

We thank Michelle Leger for helpful advice, Núria Ros i Rocher for suggestions on the figure, Łukasz
Sobala for feedback on improvements to the database, Frederick Matsen for helpful discussions,

Chris Neely and Harriet Alexander for advice on genome annotation with EukMetaSanity, and
members of the Multicellgenome Lab for discussions. This work was supported by the French
Government 'Investissement d'Avenir' program OCEANOMICS (ANR-11-BTBR-0008), the ABiMS
computing cluster at the Station Biologique de Roscoff, France, and received funding from the
European Research Council (ERC) under the European Union's Horizon 2020 research and
innovation programme (grant agreement No. 949745). DJR was supported by postdoctoral
fellowships from the Beatriu de Pinós programme of the Government of Catalonia's Secretariat for
Universities and Research of the Ministry of Economy and Knowledge, and from "la Caixa"
Foundation (ID 100010434), with the fellowship code LCF/BQ/PI19/11690008. CB was supported by
a grant from the Gordon and Betty Moore Foundation (GBMF5257 / UniEuk) and is grateful to the
International Society of Protistologists for additional support. JGW is supported by the National
Science Foundation under Grant No. DBI-2119963. FB was supported by a grant from Science for
Life Laboratory, and by grants from the Swedish Research Council VR (2017-04563 and 2021-04055)
and Formas (2017-01197). JFHS acknowledges support from the German Research Foundation
(DFG; Grant STR1349/2-1 Project No. 432453260).

432
433

References

Acinas, S. G., Sánchez, P., Salazar, G., Cornejo-Castillo, F. M., Sebastián, M., Logares, R., Royo-Llonch, M.,
Paoli, L., Sunagawa, S., Hingamp, P., Ogata, H., Lima-Mendez, G., Roux, S., González, J. M., Arrieta,
J. M., Alam, I. S., Kamau, A., Bowler, C., Raes, J., … Gasol, J. M. (2021). Deep ocean metagenomes
provide insight into the metabolic architecture of bathypelagic microbial communities. *Communications
Biology*, *4*(1), 604. https://doi.org/10.1038/s42003-021-02112-2
Adl, S. M., Bass, D., Lane, C. E., Lukeš, J., Schoch, C. L., Smirnov, A., Agatha, S., Berney, C., Brown, M. W.,
Burki, F., Cárdenas, P., Čepička, I., Chistyakova, L., del Campo, J., Dunthorn, M., Edvardsen, B., Eglit,
Y., Guillou, L., Hampl, V., … Zhang, Q. (2019). Revisions to the Classification, Nomenclature, and
Diversity of Eukaryotes. *Journal of Eukaryotic Microbiology*, *66*(1), 4–119.
https://doi.org/10.1111/jeu.12691
Alexander, H., Hu, S. K., Krinos, A. I., Pachiadaki, M., Tully, B. J., Neely, C. J., & Reiter, T. (2021). *Eukaryotic
genomes from a global metagenomic dataset illuminate trophic modes and biogeography of ocean
plankton* [Preprint]. Microbiology. https://doi.org/10.1101/2021.07.25.453713
Altschul, S. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs.
Nucleic Acids Research, *25*(17), 3389–3402. https://doi.org/10.1093/nar/25.17.3389
Bass, D. (2021, March 27). Professor Thomas Cavalier-Smith FRS FRSC. *Protistology UK*.
https://www.protistology.org.uk/t-c-s
Berney, C., Ciuprina, A., Bender, S., Brodie, J., Edgcomb, V., Kim, E., Rajan, J., Parfrey, L. W., Adl, S., Audic,
S., Bass, D., Caron, D. A., Cochrane, G., Czech, L., Dunthorn, M., Geisen, S., Glöckner, F. O., Mahé,
F., Quast, C., … de Vargas, C. (2017). *UniEuk*: Time to Speak a Common Language in Protistology!
Journal of Eukaryotic Microbiology, *64*(3), 407–411. https://doi.org/10.1111/jeu.12414
Berney, Cédric. (2022). *EukRibo: A 18S reference database* (1.0) [Data set]. Zenodo.
https://doi.org/10.5281/ZENODO.6327890
Biller, S. J., Berube, P. M., Dooley, K., Williams, M., Satinsky, B. M., Hackl, T., Hogle, S. L., Coe, A., Bergauer,
K., Bouman, H. A., Browning, T. J., De Corte, D., Hassler, C., Hulston, D., Jacquot, J. E., Maas, E. W.,
Reinthaler, T., Sintes, E., Yokokawa, T., & Chisholm, S. W. (2018). Marine microbial metagenomes
sampled across space and time. *Scientific Data*, *5*(1), 180176. https://doi.org/10.1038/sdata.2018.176
Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data.
Bioinformatics, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170
Bork, P., Bowler, C., de Vargas, C., Gorsky, G., Karsenti, E., & Wincker, P. (2015). Tara Oceans studies plankton
at planetary scale. *Science*, *348*(6237), 873–873. https://doi.org/10.1126/science.aac5605
Brown, M. W., Heiss, A. A., Kamikawa, R., Inagaki, Y., Yabuki, A., Tice, A. K., Shiratori, T., Ishida, K.-I.,
Hashimoto, T., Simpson, A. G. B., & Roger, A. J. (2018). Phylogenomics Places Orphan Protistan
Lineages in a Novel Eukaryotic Super-Group. *Genome Biology and Evolution*, *10*(2), 427–433.
https://doi.org/10.1093/gbe/evy014
Burki, F., Okamoto, N., Pombert, J.-F., & Keeling, P. J. (2012). The evolutionary history of haptophytes and
cryptophytes: Phylogenomic evidence for separate origins. *Proceedings of the Royal Society B:*

Biological Sciences, *279*(1736), 2246–2254. https://doi.org/10.1098/rspb.2011.2301

Burki, F., Roger, A. J., Brown, M. W., & Simpson, A. G. B. (2020). The New Tree of Eukaryotes. *Trends in*
Ecology & Evolution, *35*(1), 43–55. https://doi.org/10.1016/j.tree.2019.08.008

Carradec, Q., Pelletier, E., Da Silva, C., Alberti, A., Seeleuthner, Y., Blanc-Mathieu, R., Lima-Mendez, G., Rocha,
F., Tirichine, L., Labadie, K., Kirilovsky, A., Bertrand, A., Engelen, S., Madoui, M.-A., Méheust, R.,
Poulain, J., Romac, S., Richter, D. J., Yoshikawa, G., … Wincker, P. (2018). A global ocean atlas of
eukaryotic genes. *Nature Communications*, *9*(1), 373. https://doi.org/10.1038/s41467-017-02342-1

Close, T. J., Bhat, P. R., Lonardi, S., Wu, Y., Rostoks, N., Ramsay, L., Druka, A., Stein, N., Svensson, J. T.,
Wanamaker, S., Bozdag, S., Roose, M. L., Moscou, M. J., Chao, S., Varshney, R. K., Szűcs, P., Sato,
K., Hayes, P. M., Matthews, D. E., … Waugh, R. (2009). Development and implementation of high-
throughput SNP genotyping in barley. *BMC Genomics*, *10*(1), 582. https://doi.org/10.1186/1471-2164-
10-582

del Campo, J., Sieracki, M. E., Molestina, R., Keeling, P., Massana, R., & Ruiz-Trillo, I. (2014). The others: Our
biased perspective of eukaryotic genomes. *Trends in Ecology & Evolution*, *29*(5), 252–259.
https://doi.org/10.1016/j.tree.2014.03.006

Duarte, C. M. (2015). Seafaring in the 21St Century: The Malaspina 2010 Circumnavigation Expedition.
Limnology and Oceanography Bulletin, *24*(1), 11–14. https://doi.org/10.1002/lob.10008

Eddy, S. R. (2020). *HMMER: biosequence analysis using profile hidden Markov models*. http://hmmer.org/

Eisen, J. A. (2003). Phylogenomics: Intersection of Evolution and Genomics. *Science*, *300*(5626), 1706–1707.
https://doi.org/10.1126/science.1086292

El-Gebali, S., Mistry, J., Bateman, A., Eddy, S. R., Luciani, A., Potter, S. C., Qureshi, M., Richardson, L. J.,
Salazar, G. A., Smart, A., Sonnhammer, E. L. L., Hirsh, L., Paladin, L., Piovesan, D., Tosatto, S. C. E.,
& Finn, R. D. (2019). The Pfam protein families database in 2019. *Nucleic Acids Research*, *47*(D1),
D427–D432. https://doi.org/10.1093/nar/gky995

Gawryluk, R. M. R., Tikhonenkov, D. V., Hehenberger, E., Husnik, F., Mylnikov, A. P., & Keeling, P. J. (2019).
Non-photosynthetic predators are sister to red algae. *Nature*, *572*(7768), 240–243.
https://doi.org/10.1038/s41586-019-1398-6

The Gene Ontology Consortium. (2019). The Gene Ontology Resource: 20 years and still GOing strong. *Nucleic*
Acids Research, *47*(D1), D330–D338. https://doi.org/10.1093/nar/gky1055

Grigoriev, I. V., Hayes, R. D., Calhoun, S., Kamel, B., Wang, A., Ahrendt, S., Dusheyko, S., Nikitin, R., Mondo, S.
J., Salamov, A., Shabalov, I., & Kuo, A. (2021). PhycoCosm, a comparative algal genomics resource.
Nucleic Acids Research, *49*(D1), D1004–D1011. https://doi.org/10.1093/nar/gkaa898

Grigoriev, I. V., Nikitin, R., Haridas, S., Kuo, A., Ohm, R., Otillar, R., Riley, R., Salamov, A., Zhao, X.,
Korzeniewski, F., Smirnova, T., Nordberg, H., Dubchak, I., & Shabalov, I. (2014). MycoCosm portal:
Gearing up for 1000 fungal genomes. *Nucleic Acids Research*, *42*(D1), D699–D704.
https://doi.org/10.1093/nar/gkt1183

Gruber-Vodicka, H. R., Seah, B. K. B., & Pruesse, E. (2020). phyloFlash: Rapid Small-Subunit rRNA Profiling
and Targeted Assembly from Metagenomes. *MSystems*, *5*(5). https://doi.org/10.1128/mSystems.00920-
20

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., Couger, M. B., Eccles, D., Li,
B., Lieber, M., MacManes, M. D., Ott, M., Orvis, J., Pochet, N., Strozzi, F., Weeks, N., Westerman, R.,
William, T., Dewey, C. N., … Regev, A. (2013). De novo transcript sequence reconstruction from RNA-
seq using the Trinity platform for reference generation and analysis. *Nature Protocols*, *8*(8), 1494–1512.
https://doi.org/10.1038/nprot.2013.084

Huang, X. (1999). CAP3: A DNA Sequence Assembly Program. *Genome Research*, *9*(9), 868–877.
https://doi.org/10.1101/gr.9.9.868

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K., Cook, H., Mende, D. R.,
Letunic, I., Rattei, T., Jensen, L. J., von Mering, C., & Bork, P. (2019). eggNOG 5.0: A hierarchical,
functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502
viruses. *Nucleic Acids Research*, *47*(D1), D309–D314. https://doi.org/10.1093/nar/gky1085

Janouškovec, J., Tikhonenkov, D. V., Burki, F., Howe, A. T., Rohwer, F. L., Mylnikov, A. P., & Keeling, P. J.
(2017). A New Lineage of Eukaryotes Illuminates Early Mitochondrial Genome Reduction. *Current*
Biology, *27*(23), 3717-3724.e5. https://doi.org/10.1016/j.cub.2017.10.051

Kamikawa, R., Kolisko, M., Nishimura, Y., Yabuki, A., Brown, M. W., Ishikawa, S. A., Ishida, K., Roger, A. J.,
Hashimoto, T., & Inagaki, Y. (2014). Gene Content Evolution in Discobid Mitochondria Deduced from
the Phylogenetic Position and Complete Mitochondrial Genome of Tsukubamonas globosa. *Genome*
Biology and Evolution, *6*(2), 306–315. https://doi.org/10.1093/gbe/evu015

Karsenti, E., Acinas, S. G., Bork, P., Bowler, C., De Vargas, C., Raes, J., Sullivan, M., Arendt, D., Benzoni, F.,

Claverie, J.-M., Follows, M., Gorsky, G., Hingamp, P., Iudicone, D., Jaillon, O., Kandels-Lewis, S., Krzic,
U., Not, F., Ogata, H., … Wincker, P. (2011). A Holistic Approach to Marine Eco-Systems Biology. *PLoS*
Biology, *9*(10), e1001177. https://doi.org/10.1371/journal.pbio.1001177

Keeling, P. J., Burki, F., Wilcox, H. M., Allam, B., Allen, E. E., Amaral-Zettler, L. A., Armbrust, E. V., Archibald, J.
M., Bharti, A. K., Bell, C. J., Beszteri, B., Bidle, K. D., Cameron, C. T., Campbell, L., Caron, D. A.,
Cattolico, R. A., Collier, J. L., Coyne, K., Davy, S. K., … Worden, A. Z. (2014). The Marine Microbial
Eukaryote Transcriptome Sequencing Project (MMETSP): Illuminating the Functional Diversity of
Eukaryotic Life in the Oceans through Transcriptome Sequencing. *PLoS Biology*, *12*(6), e1001889.
https://doi.org/10.1371/journal.pbio.1001889

King, N., & Rokas, A. (2017). Embracing Uncertainty in Reconstructing Early Animal Evolution. *Current Biology*,
27(19), R1081–R1088. https://doi.org/10.1016/j.cub.2017.08.054

Kiss, E., Hegedüs, B., Virágh, M., Varga, T., Merényi, Z., Kószó, T., Bálint, B., Prasanna, A. N., Krizsán, K.,
Kocsubé, S., Riquelme, M., Takeshita, N., & Nagy, L. G. (2019). Comparative genomics reveals the
origin of fungal hyphae and multicellularity. *Nature Communications*, *10*(1).
https://doi.org/10.1038/s41467-019-12085-w

Kück, P., & Longo, G. C. (2014). FASconCAT-G: Extensive functions for multiple sequence alignment
preparations concerning phylogenetic studies. *Frontiers in Zoology*, *11*(1), 81.
https://doi.org/10.1186/s12983-014-0081-x

Larkin, A. A., Garcia, C. A., Garcia, N., Brock, M. L., Lee, J. A., Ustick, L. J., Barbero, L., Carter, B. R., Sonnerup,
R. E., Talley, L. D., Tarran, G. A., Volkov, D. L., & Martiny, A. C. (2021). High spatial resolution global
ocean metagenomes from Bio-GO-SHIP repeat hydrography transects. *Scientific Data*, *8*(1), 107.
https://doi.org/10.1038/s41597-021-00889-9

Lax, G., Eglit, Y., Eme, L., Bertrand, E. M., Roger, A. J., & Simpson, A. G. B. (2018). Hemimastigophora is a
novel supra-kingdom-level lineage of eukaryotes. *Nature*, *564*(7736), 410–414.
https://doi.org/10.1038/s41586-018-0708-8

Letunic, I., & Bork, P. (2019). Interactive Tree Of Life (iTOL) v4: Recent updates and new developments. *Nucleic*
Acids Research, *47*(W1), W256–W259. https://doi.org/10.1093/nar/gkz239

Li, W., & Godzik, A. (2006). Cd-hit: A fast program for clustering and comparing large sets of protein or
nucleotide sequences. *Bioinformatics*, *22*(13), 1658–1659. https://doi.org/10.1093/bioinformatics/btl158

Li, Y., Steenwyk, J. L., Chang, Y., Wang, Y., James, T. Y., Stajich, J. E., Spatafora, J. W., Groenewald, M.,
Dunn, C. W., Hittinger, C. T., Shen, X.-X., & Rokas, A. (2021). A genome-scale phylogeny of the
kingdom Fungi. *Current Biology*, *31*(8), 1653-1665.e5. https://doi.org/10.1016/j.cub.2021.01.074

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO Update: Novel and
Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of
Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology and Evolution*, *38*(10), 4647–4654.
https://doi.org/10.1093/molbev/msab199

Marron, A. O., Ratcliffe, S., Wheeler, G. L., Goldstein, R. E., King, N., Not, F., de Vargas, C., & Richter, D. J.
(2016). The Evolution of Silicon Transport in Eukaryotes. *Molecular Biology and Evolution*, *33*(12),
3226–3248. https://doi.org/10.1093/molbev/msw209

Menardo, F., Loiseau, C., Brites, D., Coscolla, M., Gygli, S. M., Rutaihwa, L. K., Trauner, A., Beisel, C., Borrell,
S., & Gagneux, S. (2018). Treemmer: A tool to reduce large phylogenetic datasets with minimal loss of
diversity. *BMC Bioinformatics*, *19*(1), 164. https://doi.org/10.1186/s12859-018-2164-8

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler, A., & Lanfear, R.
(2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era.
Molecular Biology and Evolution, *37*(5), 1530–1534. https://doi.org/10.1093/molbev/msaa015

Mitchell, A. L., Attwood, T. K., Babbitt, P. C., Blum, M., Bork, P., Bridge, A., Brown, S. D., Chang, H.-Y., El-
Gebali, S., Fraser, M. I., Gough, J., Haft, D. R., Huang, H., Letunic, I., Lopez, R., Luciani, A., Madeira,
F., Marchler-Bauer, A., Mi, H., … Finn, R. D. (2019). InterPro in 2019: Improving coverage, classification
and access to protein sequence annotations. *Nucleic Acids Research*, *47*(D1), D351–D360.
https://doi.org/10.1093/nar/gky1100

Neely, C. J., Hu, S. K., Alexander, H., & Tully, B. J. (2021). *The high-throughput gene prediction of more than*
1,700 eukaryote genomes using the software package EukMetaSanity [Preprint]. Bioinformatics.
https://doi.org/10.1101/2021.07.25.453296

Pertea, G., & Pertea, M. (2020). GFF Utilities: GffRead and GffCompare. *F1000Research*, *9*, 304.
https://doi.org/10.12688/f1000research.23297.2

Plotly Technologies Inc. (2015). *Collaborative data science*. Plotly Technologies Inc. https://plot.ly

Priyam, A., Woodcroft, B. J., Rai, V., Moghul, I., Munagala, A., Ter, F., Chowdhary, H., Pieniak, I., Maynard, L. J.,
Gibbins, M. A., Moon, H., Davis-Richardson, A., Uludag, M., Watson-Haigh, N. S., Challis, R.,

Nakamura, H., Favreau, E., Gómez, E. A., Pluskal, T., … Wurm, Y. (2019). Sequenceserver: A Modern
Graphical User Interface for Custom BLAST Databases. *Molecular Biology and Evolution*, *36*(12), 2922–
2924. https://doi.org/10.1093/molbev/msz185
Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A., & Korobeynikov, A. (2020). Using SPAdes De Novo
Assembler. *Current Protocols in Bioinformatics*, *70*(1). https://doi.org/10.1002/cpbi.102
Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O. (2012). The
SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic
Acids Research*, *41*(D1), D590–D596. https://doi.org/10.1093/nar/gks1219
Rice, P., Longden, I., & Bleasby, A. (2000). EMBOSS: The European Molecular Biology Open Software Suite.
Trends in Genetics, *16*(6), 276–277. https://doi.org/10.1016/S0168-9525(00)02024-2
Richards, T. (2021). Thomas Cavalier-Smith (1942–2021). *Nature*, *593*(7858), 190–190.
https://doi.org/10.1038/d41586-021-01250-1
Richter, D. J., & Levin, T. C. (2019). The origin and evolution of cell-intrinsic antibacterial defenses in eukaryotes.
Current Opinion in Genetics & Development, *58–59*, 111–122. https://doi.org/10.1016/j.gde.2019.09.002
Richter, D. J., Watteaux, R., Vannier, T., Leconte, J., Frémont, P., Reygondeau, G., Maillet, N., Henry, N., Benoit,
G., Fernàndez-Guerra, A., Suweis, S., Narci, R., Berney, C., Eveillard, D., Gavory, F., Guidi, L.,
Labadie, K., Mahieu, E., Poulain, J., … Tara Oceans Coordinators. (2019). *Genomic evidence for global
ocean plankton biogeography shaped by large-scale current systems* [Preprint]. Ecology.
https://doi.org/10.1101/867739
Roger, A. J. (2021). Thomas Cavalier-Smith (1942–2021). *Current Biology*, *31*(16), R977–R981.
https://doi.org/10.1016/j.cub.2021.07.009
Saldarriaga, J. (2021). Obituary for Professor Thomas Cavalier-Smith FRS, FRSC. *International Society for
Evolutionary Protistology*. https://www.isep-protists.com/post/obituary-for-professor-thomas-cavalier-
smith
Sayers, E. W., Beck, J., Brister, J. R., Bolton, E. E., Canese, K., Comeau, D. C., Funk, K., Ketter, A., Kim, S.,
Kimchi, A., Kitts, P. A., Kuznetsov, A., Lathrop, S., Lu, Z., McGarvey, K., Madden, T. L., Murphy, T. D.,
O'Leary, N., Phan, L., … Ostell, J. (2020). Database resources of the National Center for Biotechnology
Information. *Nucleic Acids Research*, *48*(D1), D9–D16. https://doi.org/10.1093/nar/gkz899
Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO: assessing
genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, *31*(19),
3210–3212. https://doi.org/10.1093/bioinformatics/btv351
Steenwyk, J. L., Buida, T. J., Li, Y., Shen, X.-X., & Rokas, A. (2020). ClipKIT: A multiple sequence alignment
trimming software for accurate phylogenomic inference. *PLOS Biology*, *18*(12), e3001007.
https://doi.org/10.1371/journal.pbio.3001007
Strassert, J. F. H., Jamy, M., Mylnikov, A. P., Tikhonenkov, D. V., & Burki, F. (2019). New Phylogenomic Analysis
of the Enigmatic Phylum Telonemia Further Resolves the Eukaryote Tree of Life. *Molecular Biology and
Evolution*, *36*(4), 757–765. https://doi.org/10.1093/molbev/msz012
Strassert, J. F. H., & Monaghan, M. T. (2021). *Phylogenomics unravels the early diversification of fungi* [Preprint].
Evolutionary Biology. https://doi.org/10.1101/2021.12.12.472261
Tara Oceans Coordinators, Sunagawa, S., Acinas, S. G., Bork, P., Bowler, C., Eveillard, D., Gorsky, G., Guidi, L.,
Iudicone, D., Karsenti, E., Lombard, F., Ogata, H., Pesant, S., Sullivan, M. B., Wincker, P., & de Vargas,
C. (2020). Tara Oceans: Towards global ocean ecosystems biology. *Nature Reviews Microbiology*.
https://doi.org/10.1038/s41579-020-0364-5
Tice, A. K., Žihala, D., Pánek, T., Jones, R. E., Salomaki, E. D., Nenarokov, S., Burki, F., Eliáš, M., Eme, L.,
Roger, A. J., Rokas, A., Shen, X.-X., Strassert, J. F. H., Kolísko, M., & Brown, M. W. (2021).
PhyloFisher: A phylogenomic package for resolving eukaryotic relationships. *PLOS Biology*, *19*(8),
e3001365. https://doi.org/10.1371/journal.pbio.3001365
Wang, H.-C., Minh, B. Q., Susko, E., & Roger, A. J. (2018). Modeling Site Heterogeneity with Posterior Mean Site
Frequency Profiles Accelerates Accurate Phylogenomic Estimation. *Systematic Biology*, *67*(2), 216–
235. https://doi.org/10.1093/sysbio/syx068
Wickett, N. J., Mirarab, S., Nguyen, N., Warnow, T., Carpenter, E., Matasci, N., Ayyampalayam, S., Barker, M.
S., Burleigh, J. G., Gitzendanner, M. A., Ruhfel, B. R., Wafula, E., Der, J. P., Graham, S. W., Mathews,
S., Melkonian, M., Soltis, D. E., Soltis, P. S., Miles, N. W., … Leebens-Mack, J. (2014).
Phylotranscriptomic analysis of the origin and early diversification of land plants. *Proceedings of the
National Academy of Sciences*, *111*(45), E4859–E4868. https://doi.org/10.1073/pnas.1323926111
Wilkinson, M. D., Dumontier, M., Aalbersberg, Ij. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-
W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I.,
Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., … Mons, B. (2016). The FAIR Guiding Principles for

scientific data management and stewardship. *Scientific Data*, *3*(1), 160018.

https://doi.org/10.1038/sdata.2016.18

Yabuki, A., Kamikawa, R., Ishikawa, S. A., Kolisko, M., Kim, E., Tanabe, A. S., Kume, K., Ishida, K., & Inagki, Y.

(2015). Palpitomonas bilix represents a basal cryptist lineage: Insight into the character evolution in

Cryptista. *Scientific Reports*, *4*(1). https://doi.org/10.1038/srep04641

Table 1. The 37 different web hosts from which data sets were downloaded. The count for
figshare.com includes 272 species from the MMETSP (Keeling et al., 2014) for which a procedure to
remove cross-contamination was applied (Marron et al., 2016). * indicates that the URLs for one or
more data sets point to servers that are no longer responding or are otherwise inaccessible.
658

Hostname	Number of Data Sets
ncbi.nlm.nih.gov	429
figshare.com	357
datadryad.org	93
jgi.doe.gov	22
ebi.ac.uk	17
rutgers.edu	10*
oist.jp	8
mendeley.com	7
yadi.sk	6
genomics.cn	3
genoscope.cns.fr	3
cngb.org	2
compagen.org	2*
nrifs.fra.affrc.go.jp	2
sb-roscoff.fr	2
ugent.be	2
veupathdb.org	2
aarnet.edu.au	1
algaegenome.org	1*
bioenergychina.org	1*
bitbucket.org	1
dataverse.scholarsportal.info	1
davidadlergold.com	1
drive.google.com	1
genomevolution.org	1
github.com	1
licebase.org	1
liv.ac.uk	1*

malab.cn	1
obs-vlfr.fr	1
ovgu.de	1
sciencemag.org	1
treegenesdb.org	1
ufl.edu	1
umontreal.ca	1*
usda.gov	1
zenodo.org	1

Figure 1. Distribution of 993 source data sets on the eukaryotic tree of life, separated by data set type, with taxonomy based on UniEuk (Adl et al., 2019; Berney et al., 2017). The position of the eukaryotic root, indicated with a dashed line, is currently unresolved. Group names shown in gray are not officially recognized. The figure was created with iTOL (Letunic & Bork, 2019).