1 **EukProt: A database of genome-scale predicted proteins across the diversity of**
2 **eukaryotes**
3

4 Daniel J. Richter[1*], Cédric Berney[2,3], Jürgen F. H. Strassert[4], Yu-Ping Poh[5], Emily K.
5 Herman[6], Sergio A. Muñoz-Gómez[7], Jeremy G. Wideman[5], Fabien Burki[8], Colomban de
6 Vargas[2,3]
7
1. Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra), Passeig Marítim de la Barceloneta 37-49,
08003 Barcelona, Spain
2. Sorbonne Université, CNRS, Station Biologique de Roscoff, UMR 7144, ECOMAP, 29680 Roscoff, France
3. Research Federation for the study of Global Ocean Systems Ecology and Evolution, FR2022/Tara GOSEE,
Paris, France
4. Department of Evolutionary and Integrative Ecology, Leibniz Institute of Freshwater Ecology and Inland
Fisheries (IGB), Müggelseedamm 301, 12587 Berlin, Germany
5. Biodesign Center for Mechanisms of Evolution, School of Life Sciences, Arizona State University, Tempe,
Arizona, United States of America
6. Department of Agricultural, Food and Nutritional Sciences, Faculty of Agricultural, Life, and Environmental
Sciences, University of Alberta, Edmonton, Alberta, Canada
7. Unité d'Ecologie, Systématique et Evolution, Université Paris-Saclay, Orsay, France
8. Department of Organismal Biology and Science for Life Laboratory, Uppsala University, Norbyvägen 18D,
Uppsala 75236, Sweden
22
23 *Corresponding author
24 Correspondence: daniel.j.richter@gmail.com

25
26 **ABSTRACT**
27

EukProt is a database of published and publicly available predicted protein sets selected to represent
the breadth of eukaryotic diversity, currently including 993 species from all major supergroups as well
as orphan taxa. The goal of the database is to provide a single, convenient resource for gene-based
research across the spectrum of eukaryotic life, such as phylogenomics and gene family evolution.
Each species is placed within the UniEuk taxonomic framework in order to facilitate downstream
analyses, and each data set is associated with a unique, persistent identifier to facilitate comparison
and replication among analyses. The database is regularly updated, and all versions will be
permanently stored and made available via FigShare. The current version has a number of updates,
notably 'The Comparative Set' (TCS), a reduced taxonomic set with high estimated completeness
while maintaining a substantial phylogenetic breadth, which comprises 196 predicted proteomes. A
BLAST web server and graphical displays of data set completeness are available at
http://evocellbio.com/eukprot/. We invite the community to provide suggestions for new data sets and
new annotation features to be included in subsequent versions, with the goal of building a
collaborative resource that will promote research to understand eukaryotic diversity and
diversification.

43
44 *Keywords*: eukaryotes, eukaryotic diversity, transcriptomes, genomes, single-cell genomes,
45 phylogenomics, UniEuk taxonomy, predicted proteins

Introduction

Over the past 15 years, the discovery of diverse novel microbial eukaryotes, coupled with methods to reconstruct phylogenies based on hundreds of protein-coding genes (frequently referred to as phylogenomics, although the term was originally proposed as the more general integration of genome analysis and evolutionary studies [Eisen, 2003]) have led to a remarkable reshaping in our understanding of the eukaryotic tree of life, the proposal of new supergroups and the placement of enigmatic lineages in known supergroups (Burki et al., 2012, 2020; Kamikawa et al., 2014; Yabuki et al., 2015; Janouškovec et al., 2017; Brown et al., 2018; Lax et al., 2018; Strassert et al., 2019; Gawryluk et al., 2019; Tice et al., 2021). The phylogenomic approach has also been used to investigate branching patterns within eukaryotic supergroups, including to understand species evolutionary relationships with implications for the early evolutionary events in the three best-studied eukaryotic lineages: land plants (Wickett et al., 2014), animals (King & Rokas, 2017) and fungi (Kiss et al., 2019; Li et al., 2021; Strassert & Monaghan, 2022). Furthermore, analyses in diverse eukaryotes have the potential to reveal new genes and pathways for biological processes currently characterized only in these three well-studied lineages and their parasites (del Campo et al., 2014; Richter & Levin, 2019). In the ocean alone, planetary-scale metagenomics studies (Bork et al., 2015; Tara Oceans Coordinators et al., 2020) have already unveiled an extreme diversity of eukaryotic genes (Carradec et al., 2018) whose phylogenetic origin and ecological function are mostly unknown.

A critical prerequisite to all genomic studies is the underlying database of predicted proteins from which orthologs are extracted or other sequence analyses are performed. Because no single source contains the protein predictions for the complete set of eukaryotes that have been sequenced at a genomic scale, each study needs to assemble their own database, reducing reproducibility among analyses (due to the inclusion of different taxa, or different data sets for the same taxon), and producing a significant barrier to new researchers entering the field. In addition, because the large majority of the protein data sets from diverse eukaryotes are not included in major public databases (see Figure 2), researchers cannot easily access them via standard tools such as NCBI BLAST (Sayers et al., 2020) in order to find the homologs of a protein of interest, nor are they automatically integrated into major public databases containing annotations such as protein domains (e.g., Pfam [El-Gebali et al., 2019], Interpro [Mitchell et al., 2019]) or gene ontology (The Gene Ontology Consortium, 2019).

To address this gap, we assembled a database of protein sequences – EukProt – either from newly performed protein prediction or by retrieving available sequences from a comprehensive set of species representing known eukaryotic diversity. We note that an existing database of genome-scale protein data sets, PhyloDB (https://github.com/allenlab/PhyloDB), contains 550 eukaryotic species with at least 500 proteins, as of version 1.076. PhyloDB was most recently updated in 2015; EukProt includes data made available since then, allowing it to provide more species with generally higher completeness and representing a greater phylogenetic breadth. In addition, we placed each species within a universal eukaryotic taxonomic framework, UniEuk (Berney et al., 2017), in order to ensure that the evolutionary relationships among data sets are accurately and consistently described. We also provide estimated completeness statistics for each predicted proteome based on BUSCO scores and have selected a suggested subset of 196 proteomes for use in comparative genomics investigations. EukProt is designed to prioritize ease of use, with unique, persistent identifiers assigned to each data set and a standard system of nomenclature to facilitate repeatability of analyses. To this end, we have followed FAIR principles (Findable, Accessible, Interoperable, Reusable) in its construction (Wilkinson et al., 2016).

Methods

Species and strain identity

We determined species and strain identities by reading the publications that described the data
sets, consulting the literature for naming revisions, and comparing 18S ribosomal DNA sequences for
each data set to reference sequence databases (see below for a description of how we retrieved 18S
sequences). For species that were previously known by other names, we recorded these previous
names in the metadata for the data set, except in cases where a species was originally assigned to a
genus but not identified to the species level (e.g., *Goniomonas* sp., now identified as *Goniomonas*
avonlea, is not listed as a previous name). This exception was not followed when the GenBank record
description for the 18S sequence contained a different name than the one used in EukProt; in this
case this previous name is always listed in the previous names field, in order to avoid confusion when
retrieving 18S sequences from GenBank.
When no strain name was available for an MMETSP data set, we used the MMETSP ID as the
strain name (for example, EP00362 has the strain name MMETSP1317). For strains with alternative
names, we included these in a dedicated field for alternative strain names; this list is not necessarily
exhaustive (i.e., it is not guaranteed to contain all alternative names for a given strain).
111
112   **Supergroups and taxogroups from UniEuk**
113
The full taxonomic pathways provided for all species (which follow the framework developed in the
UniEuk project [Berney et al., 2017]) are not based on a fixed number of ranks, but on a free,
unlimited number of taxonomic levels, in order to match phylogenetic evidence as closely as possible.
This provides end-users more information and flexibility, but could also make it more difficult to
summarize results of downstream analyses. Therefore, we provide three additional fields
("supergroup", "taxogroup1" and "taxogroup2") to help end-users whenever it is useful to distribute
eukaryotic diversity into a fixed number of taxonomic categories of roughly equivalent phylogenetic
depth or ecological relevance. The groupings called "supergroups" in the context of UniEuk resources
(36 recognized lineages so far, of which 34 are included in EukProt; *Microheliella* and *Meteora* are not
represented) consist of strictly monophyletic, deep-branching eukaryotic lineages of a phylogenetic
depth equivalent to some of the first proposed supergroups such as Opisthokonta, Alveolata,
Rhizaria, and Stramenopiles. UniEuk "supergroups" correspond roughly to the deepest phylogenetic
resolution of eukaryotic relationships achievable with ribosomal RNA genes, and not necessarily to
the currently recognized highest-level groupings of eukaryotes based on phylogenomic evidence
(these groupings are, of course, present in the complete taxonomic pathways). UniEuk "supergroups"
are therefore highly variable in relative diversity, ranging from lineages consisting of a single, orphan
genus (e.g., *Ancoracysta*, *Mantamonas*, *Palpitomonas*), to Opisthokonta as a whole. The
"taxogroup1" (of which there are 72 in EukProt) and "taxogroup2" (of which there are 198 in EukProt)
levels allow further subdivision of large supergroups into lineages of relatively equivalent evolutionary
or ecological relevance, based on current knowledge. These levels are more arbitrarily defined but
are intended to represent strictly monophyletic groupings that match one of the levels in the complete
taxonomic pathways. As an illustrative example, diatoms are in the Diatomeae taxogroup2, which is
within the Ochrophyta taxogroup1, which is within the Stramenopiles supergroup. Small, ecologically
and morphologically homogeneous supergroups are not subdivided further; in such cases the
"taxogroup1" and "taxogroup2" levels are the same as the "supergroup" level. The same approach will
be used in EukRibo, a manually-curated database of reference ribosomal RNA gene sequences
developed in parallel (Berney 2022) to help users link analyses of different types of genetic data.
141
142   **Merging strains from the same species**
143
In general, we only included data from a single strain/isolate per species. However, when only a
single transcriptome data set was available for a given strain of a species, and there were additional
published transcriptome data sets for other strains of the same species, we combined them using CD-
HIT (Li & Godzik, 2006) run with default parameter values, in order to guard against the possibility
that a single transcriptome might lack genes expressed only in one condition or experiment. When

multiple strains were merged to produce a species' data set (there were 41 such cases), this
information is indicated in the metadata for the data set. When a data set is indicated as merged, but
more than one strain name is not listed in the strain column, this indicates that the data set was
distributed as merged and we were unable to determine the exact strains used.
153
154 **Processing steps applied to publicly available data**
155
All sequences within each FASTA file are assigned a unique, standardized identifier based on the
data set's EukProt ID and on the type of data (protein or transcriptome); this identifier is prepended to
the existing FASTA header, separated by a space. Illegal characters are removed from sequences.
The following characters are permitted, as defined by NCBI BLAST
(https://blast.ncbi.nlm.nih.gov/Blast.cgi?CMD=Web&PAGE_TYPE=BlastDocs&DOC_TYPE=BlastHelp
) for nucleic acid sequences: `ACGTNUKSYMWRBDHV` and for protein sequences:
`ABCDEFGHIKLMNPQRSTUVWYZX*`.
EukProt metadata indicate the additional steps applied to each data set after downloading from
the data source (if any).
All software parameter values were default (unless otherwise specified below or in the metadata
record for a given data set), as default parameter values are most likely to correspond to the options
recommended for general use by the authors of the software. Due to the large volume of data sets we
processed, and variability among them, we were not able to test parameter values beyond those
specified below.
170
'assemble mRNA': *de novo* transcriptome assembly using Trinity v. 2.8.4, http://trinityrnaseq.github.io/
(Haas et al., 2013). We trimmed Illumina input reads for adapters and sequence quality using the
built-in '--trimmomatic' option (whose default trimming settings are based on optimal trimming
parameters from [MacManes, 2014]). We trimmed 454 input reads prior to running Trinity with
Trimmomatic v. 0.3.9, http://www.usadellab.org/cms/?page=trimmomatic (Bolger et al., 2014) with the
directives 'ILLUMINACLIP:[454 adapters FASTA file]:2:30:10 SLIDINGWINDOW:4:5 LEADING:5
TRAILING:5 MINLEN:25' (corresponding to the default Trinity trimming parameters, but for single-end
reads). When the sequence library was described as stranded in the NCBI Sequence Read Archive,
we used the corresponding 'SS_lib_type' option.
180
181 'translate mRNA': *de novo* translation of mRNA sequences with Transdecoder v. 5.3.0,
182 http://transdecoder.github.io/. When the number of predicted protein sequences for a given species
183 was both less than half of the input mRNA sequences and less than 15,000, we reduced the minimum
184 predicted protein length to 50 (from the default of 100).
185
186 'CD-HIT': clustering of protein sequences to produce a non-redundant data set using CD-HIT v. 4.6,
187 http://weizhongli-lab.org/cd-hit/ (Li & Godzik, 2006). We used this tool principally to combine protein
188 predictions for different strains of the same species, but also to reduce the size of very large predicted
189 protein sets (>50,000 proteins) that showed evidence of redundancy.
190
'extractfeat', 'seqret', 'transeq', and 'trimseq': from the EMBOSS package v. 6.6.0.0,
http://emboss.sourceforge.net/ (Rice et al., 2000). We used extractfeat to produce coding sequences
(CDS) from genomes with gene annotations in EMBL format but without publicly available protein
sequences. We used seqret to convert FASTQ files to FASTA files. We used transeq to translate
CDS directly into proteins. We used trimseq to trim EST sequences before translation with
Transdecoder.
197
198 'gffread': to produce protein sequences from genomes with gene annotations in GFF format but
199 without publicly available protein sequences, using the command line option '-y' in version 0.12.3
200 (Pertea & Pertea, 2020).

'predict genes': we used EukMetaSanity https://github.com/cjneely10/EukMetaSanity (Neely et al., 2021) to perform automated annotation of genome sequences lacking publicly available protein predictions. We used the following parameters, as specified in (Alexander et al., 2021): --min_contig 500 --min_contig_in_predict 500 --max_contig 100000000. These settings were designed for generalized gene prediction on genomes expected to be sampled from across eukaryotic diversity, which is also the case for the data sets in EukProt. We used the parameter --min_contig_in_predict 200 (as it matched the default minimum contig length in Trinity). By default, we selected the proteins at Tier 2 (predictions supported by at least 2 sources). If Tier 2 produced fewer than 15,000 predicted proteins, we instead selected Tier 1. All other parameter values were left at their defaults. We did not perform gene prediction on unannotated genomes for which a transcriptome was already available for the same species (under the assumption that the gene predictions of the transcriptome would be of higher quality, due to potential errors in the gene annotation process).

Retrieval of 18S sequences

We searched for 18S sequences of at least 1,500 base pairs in length for each data set in the following order. If no sequence, or only a partial sequence was retrieved in a given step, we moved on to the next step. Details specific to each data set are included in the EukProt metadata.

1. Search GenBank (Sayers et al., 2020) for an 18S sequence for the species and strain. When more than one sequence was available for the strain, we chose the one containing the largest quantity of high-quality nucleotides (after identifying and eliminating sequencing errors, chimeras, etc.).
2. If the data set was sequenced as part of the MMETSP, use the 18S sequence from MMETSP (if it was provided in the project metadata, and also was not already present in GenBank).
3. Search GenBank for an 18S sequence for another strain of the same species.
4. Using the 18S sequence for the most closely related species in GenBank as a query, search the contigs associated with the data set (either genome or transcriptome) using blastn (Altschul, 1997).
5. Assemble the 18S sequence from the raw sequence reads associated with the data set, using phyloFlash version 3.4 (Gruber-Vodicka et al., 2020) with the SILVA database version 138.1 (Quast et al., 2012) and SPAdes assembler version 3.14.1 (Prjibelski et al., 2020), with default parameter values except for the number of reads used, and with read length set to match the read length of the data set. For each data set, we initially ran phyloFlash with all sequence reads. If we observed that there was evidence for reads mapping to the 18S of the target lineage, but that did not result in a contig assembled by SPAdes, we ran phyloFlash again with only the first 10 million reads. If this run showed a similar result (reads mapping but no contigs), we ran again with only the first 1 million reads. When all three runs (all reads, first 10 million reads, first 1 million reads) did not result in a SPAdes assembly, we selected the reads mapping to the 18S of the target lineage and performed an assembly using CAP3 version date 02/10/15 (Huang, 1999) with 'relaxed' parameter values from (Close et al., 2009): -p 75 -d 200 -f 250 -h 90.

A selected subset of EukProt for phylogenetic studies

Single-copy orthologs were identified in EukProt v3 proteomes using BUSCO v5.2.2 (Manni et al., 2021) with the Eukaryota Odb10 database. BUSCO searches were run in protein mode with the default E-value cutoff of 1e-3. These BUSCO scores were visualized using Plotly (Plotly Technologies Inc., 2015) and are available at http://evocellbio.com/eukprot/. Single-copy BUSCO markers for taxa with BUSCO scores >30 were aligned to single-copy BUSCO profile HMMs with hmmalign (HMMER v3.3; [Eddy, 2020]), trimmed with ClipKit (-m gappy; [Steenwyk et al., 2020]), and concatenated into a supermatrix with FASConCat-G v1.04 (Kück & Longo, 2014). A guide tree was first inferred from this

supermatrix with IQ-TREE v2.0.3 using the LG4X model (-fast mode; [Minh et al., 2020]). A second
phylogenetic tree was inferred using the LG+PMSF(C10)+F+G4 model and using the LG4X tree as a
guide (Wang et al., 2018). A first subset of taxa was automatically selected from the previously
inferred tree to maximize phylogenetic diversity using Treemmer v0.1 (Menardo et al., 2018). A few
taxa that were phylogenetically misplaced with high support, likely due to a high percentage of
contaminating sequences, were visually detected and manually removed. Furthermore, predicted
proteins from model organisms (e.g., *Homo sapiens*, *Saccharomyces cerevisiae*, *Toxoplasma gondii*,
Dictyostelium discoideum, etc.) were included in the TCS. In addition, phylogenetically important taxa
with sufficiently high BUSCO scores were also included. Since different lineages universally lack
particular BUSCOs, a singular score cut-off is not reasonable and therefore each taxon group was
assessed in relation to a model system with a trusted genome. For example, while *Giardia intestinalis*
and *Toxoplasma gondii* have low BUSCO scores (23.5 and 61.2, respectively), they are generally
accepted as complete genomes. Thus, species in these lineages (e.g., Fornicata, Preaxostyla,
Apicomplexa, Parabasalia, etc.) with comparable BUSCO scores were included in the TCS. Finally, in
taxonomically important lineages in which no model yet exists (e.g., Ancyromonads) proteome sets
with BUSCO scores ~50% or greater were included, or if no such representative was available, it was
excluded from the TCS (e.g., Tsukubomonads). We welcome suggestions from the community for
future iterations of the TCS.
271
272 **The EukProt database distribution on FigShare**
273
The database is distributed in five files. One file contains 993 protein data sets, for species with
either a genome (375), a single-cell genome (56), a transcriptome (498), a single-cell transcriptome
(47), or an EST assembly (17). A second file contains assembled transcriptome contigs, for 126
species with publicly available mRNA sequence reads but no publicly available assembly. The
proteins predicted from these assemblies are included in the proteins file. A third file contains GFF
annotations for 40 genomes lacking publicly available predicted protein annotations. Finally, the
database metadata are distributed as two files: one file for the data sets included in the current
version of the database (993), and a second file for data sets not included (163), accompanied by the
reason they were not included. For example, when a data set cleaned with a genome-scale
decontamination process is described in a manuscript, but the decontaminated data are not publicly
available, it is not included in the database.
285
286   **Results and Discussion**
287
288

Figure 1. Distribution of 993 source data sets on the eukaryotic tree of life, separated by data set type, with taxonomy based on UniEuk (Berney et al., 2017; Adl et al., 2019). The position of the eukaryotic root, indicated with a dashed line, is currently unresolved. Group names shown in gray are not officially recognized. The figure was created with iTOL (Letunic & Bork, 2019).

The EukProt database

EukProt (currently in version 3; 22 November, 2021) contains 993 eukaryotic species from 5 different sequence data-types (Figure 1): genome (375 species), single-cell genome (56), transcriptome (498), single-cell transcriptome (47), and expressed sequence tag (EST; 17). The data sets were downloaded from 37 different sources (Figure 2), with the two principal sources being NCBI (Sayers et al., 2020) and the Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP) (Keeling et al., 2014). The database contains broad taxonomic representation, with species from 34 of 36 major deeply-branching eukaryotic lineages defined in the UniEuk database (Berney et al., 2017). Within these major groups, however, the representation remains uneven, which results from several factors: the difficulty of discovering, culturing or sequencing species from many lineages; a bias towards sequencing either macroscopic, multicellular species or unicellular species that are parasites (e.g., Apicomplexa), photosynthetic (such as diatoms, which are part of the ochrophyte lineage), or are otherwise economically important (del Campo et al., 2014); and very uneven evolutionary histories among groups, with some having undergone major diversification with many different morphologies, and others having diversified to a lesser extent (or with only a few surviving lineages).

Figure 2. The 37 different web hosts from which data sets were downloaded. The count for figshare.com includes 272 species from the MMETSP (Keeling et al., 2014) for which a procedure to remove cross-contamination was applied (Marron et al., 2016). Individual URLs for each data set are

317　　listed in the metadata files available on FigShare. * indicates that the URLs for one or more data sets
318　　　　　　point to servers that are no longer responding or are otherwise inaccessible.
319
We have implemented in version 3 several changes proposed by the community of EukProt
database users after version 2 was released: we have included (i) the 18S ribosomal DNA sequence
corresponding to each data set (when available), (ii) a measurement of the completeness of the data
set as estimated by BUSCO (Simão et al., 2015; Manni et al., 2021), and (iii) the assignment of
unique EukProt identifiers to individual protein sequences.
The EukProt database is organized around 5 guiding principles:
326
Breadth of species phylogenetic diversity: the objective of the database is to include as many lineages
as possible from the broad diversity of eukaryotes. For the majority of lineages, at least one
representative from all sequenced genera is included. The exceptions are lineages with hundreds of
sequenced representatives, or lineages whose sequenced species are relatively closely related, for
which we included only a subset of species selected to represent phylogenetic diversity in order to
balance the overall size of the database versus broad representation of eukaryotes. These well-
sequenced lineages are Chloroplastida (containing the land plants and green algae), Metazoa
(animals), Amoebozoa, Fungi, and some parasites/pathogens (members of Apicomplexa,
Peronosporomycetes, Metakinetoplastina and Fornicata). Within animals, we also emphasized the
inclusion of marine planktonic species, as we anticipate these could serve as mapping targets for data
from large-scale metagenomic and metatranscriptomic ocean sequencing projects (e.g., *Tara* Oceans
[Karsenti et al., 2011; Carradec et al., 2018; Richter et al., 2019], Malaspinas [Duarte, 2015; Acinas et
al., 2021], Bio-GO-SHIP [Larkin et al., 2021] and GEOTRACES [Biller et al., 2018]).
340
Convenience of access: the full data set and its associated metadata can easily be downloaded from
FigShare. For 624 of the total 993 species in the database, publicly available protein sequences were
ready to be included directly. For the remaining 369 species, additional processing steps were
required to produce protein sequences for inclusion in the database. The most common
bioinformatics processing steps were *de novo* assembling and translating transcriptomes from raw
read data (120), translating mRNA sequences from publicly available assembled transcriptomes or
EST projects lacking protein predictions (102), merging independent sets of protein predictions for the
same species (85 species) and, newly added in version 3 of the database, producing *de novo* gene
predictions for unannotated genomes (40). We also provide transcriptome assemblies and GFF
annotation files for the latter two classes, as they are not publicly available. A full list of the different
types of actions is described in the Methods, and the actions taken for each species are available in
the database metadata.
353
A unified taxonomic framework: we placed all species into the taxonomic framework generated in the
UniEuk project (Berney et al., 2017), which is based on the most recent consensus eukaryotic
classification (Adl et al., 2019) and phylogenomic evidence. This will maximize interoperability of the
database with other resources built upon to the UniEuk taxonomic framework (e.g., EukBank and
EukMap [Berney et al., 2017], and EukRibo [Berney, 2022]), and ensure that results arising from
EukProt can use a common set of identifiers to describe analyses at different taxonomic levels. For
example, this might include labeling groups in a phylogenetic tree, or summarizing read placement
data by taxonomic group. Knowledge of the taxonomy can also facilitate the detection of mis-identified
gene sequences within a data set, via comparison of the topology of gene trees to the taxonomy. To
assist in species identification and to provide a convenient link to EukRibo (a manually curated
database of 18S rDNA sequences, part of UniEuk, aimed at taxonomic annotation of metabarcoding
data sets), we also supply in version 3 the 18S sequence corresponding to the sequenced strain (or, if
an 18S for the sequenced strain is not available, an 18S sequence from the same species; see
Methods). In addition to the complete taxonomic pathway containing all intermediate nodes, we
provide for each species three sequentially nested groupings ("taxogroup2" within "taxogroup1" within

"supergroup"; see Methods) that can help to judge the redundancy of data sets, or conversely their
lack of representation, within different groupings. The taxogroup1 level is used as the leaves in Figure
1. Finally, for each data set we include a list of previous species identifications, if any, to prevent
issues resulting from revisions to species names or lineages and corrections of species
misidentifications, which have occurred frequently with improvements in sequencing and phylogenetic
techniques and the discovery of new eukaryotic organisms.
375
Appropriate references to publications and data sources: to ensure that the researchers who
generated and provided each data set receive appropriate credit, we provide the DOI of the
publication describing each data set as well as the URL from which it was downloaded. The list of
URLs should also allow users of the database to download the original sequences for each data set
(although it is not possible to guarantee that the URLs for all data providers will be permanently
available).
382
Reusability, persistence and replicability: the database will be released in successive versions, each
of which will be permanently stored and accessible at FigShare. Thus, analyses using the database
will need only to specify which version was used, enabling follow-up analyses or replications to begin
with the identical database. In addition, each individual data set within the database is assigned a
unique, permanent identifier. When a new data set becomes available for a given species, it is
assigned a new unique identifier (and the identifier of the data set it replaces, if any, is indicated in the
database metadata). These and all other changes between versions of the database are recorded as
appropriate.
391
392 **'The Comparative Set' (TCS): A selected subset of EukProt for comparative genomics**
393 **investigations**
394
Comparative genomics investigations are crucial to bridge the gaps between cell biology and
biochemistry on one hand and diversity and evolution on the other. Classically, the set of species to
include has been chosen by the authors of each study. With the goal of providing some consistency in
future comparative genomics studies, we have selected a set of predicted proteomes with high
completeness to represent the broad phylogeny of eukaryotes. A total of 196 predicted proteomes
were selected as 'The Comparative Set' (TCS; Figure 3), abbreviated in honor of the late Tom
Cavalier-Smith (Saldarriaga, 2021; Bass, 2021; Richards, 2021; Roger, 2021). All major model
systems were included in this set as well as proteomes derived from high-quality genomes. Additional
proteomes were chosen to maximize phylogenetic diversity and based on their completeness as
estimated by BUSCO (Manni et al., 2021) scores. Updates to the TCS will be based on availability of
new proteomes from phylogenetically important lineages currently underrepresented, and on
community feedback. Please send any comments or suggestions for the TCS to
Jeremy.Wideman@asu.edu. The complete EukProt database and the TCS are available for download
and also available for BLAST searches at http://evocellbio.com/eukprot/ using Sequenceserver
(Priyam et al., 2019).
410
411
412

413
414 **Figure 3**. Distribution of 196 selected data sets of The Comparative Set (TCS) on the eukaryotic tree
415 of life, with relationships based on the UniEuk taxonomy (Berney et al., 2017; Adl et al., 2019). The
416 position of the eukaryotic root, indicated with a dashed line, is currently unresolved. Group names
417 shown in gray are not officially recognized. The figure was created with iTOL (Letunic & Bork, 2019).
418

Current limitations of the EukProt database

420
In the course of constructing successive versions of EukProt, we have observed several areas
that we believe to be current limitations of the database:
1. As can be observed in Figure 1, representation of species is highly uneven among taxonomic
groups. This is largely due to the difficulty in identifying, cultivating and sequencing species
from underrepresented groups; however, some representatives of key taxa have been
published but are not publicly accessible (a list of these species can be found in the "not
included" metadata table of our FigShare distribution).
2. We currently lack an automated, consistent method to identify and remove contaminated data
sets or individual contaminant sequences. Currently, contaminated data sets are identified by
end users and removed in subsequent versions.
3. Most species are not represented by genomes, which are the gold standard for providing
gene catalogs. Instead, due to technical limitations in cultivation and/or sequencing methods,

433		they are represented by transcriptomes from laboratory cultures or by single-cell genomes or
434		transcriptomes.
435	4.	Proteins smaller than 50 amino acids are not predicted with the settings we used for protein
436		prediction (but may be included in data sets for which we did not perform protein prediction).
437		We used the default settings for protein prediction in TransDecoder (see Methods), in order to
438		compromise between minimum protein length and total number of predicted proteins (as
439		proteomes predicted with protein sizes smaller than 50 amino acids are generally much
440		larger, which may significantly slow many downstream analyses). EukProt users interested in
441		proteins shorter than 50 amino acids (for the species on which we performed protein
442		predictions) would instead have to repeat protein predictions using their desired settings.
443		

Growing the EukProt database with community involvement

The core functionality of the database is the distribution of genome-scale protein sequences
across the diversity of eukaryotic life and within the UniEuk framework. However, we anticipate that
numerous other features might be useful to the community, and we hope to involve the community in
suggesting and adding new features in successive versions. These may include information or
analyses on full data sets, such as the sequencing technology that was used (e.g., Illumina, PacBio),
or the estimation of potential contamination levels (e.g., from mixed cultures or prey species) with
non-target species, as inferred using systematic sequence homology searches. Additionally, we
propose to use our GitHub as a flexible repository to disseminate community-contributed annotations
on the level of individual protein sequences, such as protein domains from Pfam (El-Gebali et al.,
2019)/Interpro (Mitchell et al., 2019), gene ontology (The Gene Ontology Consortium, 2019)/eggNOG
(Huerta-Cepas et al., 2019), which may be updated in between EukProt releases.

As new genome-scale eukaryotic protein data sets become available, we plan to add them to the
database. As yet, we do not have a formal mechanism to accomplish this, and will instead depend on
monitoring the literature and assistance from the community. As an example, for version 3, we relied
on PhycoCosm (Grigoriev et al., 2021), MycoCosm (Grigoriev et al., 2014) and PhyloFisher (Tice et
al., 2021) to add data sets that became recently available. Because the rate of genome sequencing is
increasing exponentially, we will prioritize adding species that are *incertae sedis* and those from
taxonomic groups with currently limited representation in EukProt, and only later would include new
proteomes from currently well sampled groups.

In the longer term, we hope the standardization of our database provides a path towards including
all data sets in a major sequence repository such as NCBI/EBI/DDBJ, so that they can be more
broadly accessible and integrated into the suites of tools available at these repositories.

Acknowledgements

472   We thank Michelle Leger for helpful advice, Núria Ros-Rocher for suggestions on the figure,
473 Łukasz Sobala and Aleix Obiol for feedback on improvements to the database, Frederick Matsen for
474 helpful discussions, Chris Neely and Harriet Alexander for advice on genome annotation with
475 EukMetaSanity, and members of the Multicellgenome Lab for discussions.

Data, script and code availability

479   The EukProt v3 database is available on FigShare
480 (https://doi.org/10.6084/m9.figshare.12417881.v3). Scripts used for data processing and for building
481 the tree in Figures 1 and 3, as well as annotations of protein data sets (e.g., BUSCO), are available
482 on GitHub via Zenodo (https://doi.org/10.5281/zenodo.7025267). A BLAST server is available at:
483 http://evocellbio.com/eukprot/

Conflict of interest disclosure

The authors declare they have no conflict of interest relating to the content of this article.

Funding

This work was supported by the French Government 'Investissement d'Avenir' program OCEANOMICS (ANR-11-BTBR-0008), the ABiMS computing cluster at the Station Biologique de Roscoff, France, and received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 949745). DJR was supported by postdoctoral fellowships from the Beatriu de Pinós programme of the Government of Catalonia's Secretariat for Universities and Research of the Ministry of Economy and Knowledge, and from "la Caixa" Foundation (ID 100010434), with the fellowship code LCF/BQ/PI19/11690008. CB was supported by grants from the Gordon and Betty Moore Foundation (GBMF5257 & GBMF8908 / UniEuk project) and is grateful to the International Society of Protistologists for additional support. JGW is supported by the National Science Foundation under Grant No. DBI-2119963. FB was supported by a grant from Science for Life Laboratory, and by grants from the Swedish Research Council VR (2017-04563 and 2021-04055) and Formas (2017-01197). JFHS acknowledges support from the German Research Foundation (DFG; Grant STR1349/2-1 Project No. 432453260).

References

Acinas SG, Sánchez P, Salazar G, Cornejo-Castillo FM, Sebastián M, Logares R, Royo-Llonch M, Paoli L, Sunagawa S, Hingamp P, Ogata H, Lima-Mendez G, Roux S, González JM, Arrieta JM, Alam IS, Kamau A, Bowler C, Raes J, Pesant S, Bork P, Agustí S, Gojobori T, Vaqué D, Sullivan MB, Pedrós-Alió C, Massana R, Duarte CM, Gasol JM (2021) Deep ocean metagenomes provide insight into the metabolic architecture of bathypelagic microbial communities. *Communications Biology*, **4**, 604. https://doi.org/10.1038/s42003-021-02112-2

Adl SM, Bass D, Lane CE, Lukeš J, Schoch CL, Smirnov A, Agatha S, Berney C, Brown MW, Burki F, Cárdenas P, Čepička I, Chistyakova L, del Campo J, Dunthorn M, Edvardsen B, Eglit Y, Guillou L, Hampl V, Heiss AA, Hoppenrath M, James TY, Karnkowska A, Karpov S, Kim E, Kolisko M, Kudryavtsev A, Lahr DJG, Lara E, Le Gall L, Lynn DH, Mann DG, Massana R, Mitchell EAD, Morrow C, Park JS, Pawlowski JW, Powell MJ, Richter DJ, Rueckert S, Shadwick L, Shimano S, Spiegel FW, Torruella G, Youssef N, Zlatogursky V, Zhang Q (2019) Revisions to the Classification, Nomenclature, and Diversity of Eukaryotes. *Journal of Eukaryotic Microbiology*, **66**, 4–119. https://doi.org/10.1111/jeu.12691

Alexander H, Hu SK, Krinos AI, Pachiadaki M, Tully BJ, Neely CJ, Reiter T (2021) *Eukaryotic genomes from a global metagenomic dataset illuminate trophic modes and biogeography of ocean plankton*. Microbiology. https://doi.org/10.1101/2021.07.25.453713

Altschul S (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research*, **25**, 3389–3402. https://doi.org/10.1093/nar/25.17.3389

Bass D (2021) Professor Thomas Cavalier-Smith FRS FRSC. *Protistology UK*. https://www.protistology.org.uk/t-c-s

Berney, Cédric (2022) EukRibo: a manually curated eukaryotic 18S rDNA reference database. https://doi.org/10.5281/ZENODO.6327890

Berney C, Ciuprina A, Bender S, Brodie J, Edgcomb V, Kim E, Rajan J, Parfrey LW, Adl S, Audic S, Bass D, Caron DA, Cochrane G, Czech L, Dunthorn M, Geisen S, Glöckner FO, Mahé F, Quast C, Kaye JZ, Simpson AGB, Stamatakis A, del Campo J, Yilmaz P, de Vargas C (2017) *UniEuk*: Time to Speak a Common Language in Protistology! *Journal of Eukaryotic Microbiology*, **64**, 407–411. https://doi.org/10.1111/jeu.12414

Biller SJ, Berube PM, Dooley K, Williams M, Satinsky BM, Hackl T, Hogle SL, Coe A, Bergauer K, Bouman HA, Browning TJ, De Corte D, Hassler C, Hulston D, Jacquot JE, Maas EW, Reinthaler T, Sintes E, Yokokawa T, Chisholm SW (2018) Marine microbial metagenomes sampled across space and time. *Scientific Data*, **5**, 180176. https://doi.org/10.1038/sdata.2018.176

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics, **30**, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Bork P, Bowler C, de Vargas C, Gorsky G, Karsenti E, Wincker P (2015) Tara Oceans studies
plankton at planetary scale. *Science*, **348**, 873–873. https://doi.org/10.1126/science.aac5605

Brown MW, Heiss AA, Kamikawa R, Inagaki Y, Yabuki A, Tice AK, Shiratori T, Ishida K-I, Hashimoto
T, Simpson AGB, Roger AJ (2018) Phylogenomics Places Orphan Protistan Lineages in a Novel
Eukaryotic Super-Group. *Genome Biology and Evolution*, **10**, 427–433.
https://doi.org/10.1093/gbe/evy014

Burki F, Okamoto N, Pombert J-F, Keeling PJ (2012) The evolutionary history of haptophytes and
cryptophytes: phylogenomic evidence for separate origins. *Proceedings of the Royal Society B:*
Biological Sciences, **279**, 2246–2254. https://doi.org/10.1098/rspb.2011.2301

Burki F, Roger AJ, Brown MW, Simpson AGB (2020) The New Tree of Eukaryotes. *Trends in Ecology*
& Evolution, **35**, 43–55. https://doi.org/10.1016/j.tree.2019.08.008

del Campo J, Sieracki ME, Molestina R, Keeling P, Massana R, Ruiz-Trillo I (2014) The others: our
biased perspective of eukaryotic genomes. *Trends in Ecology & Evolution*, **29**, 252–259.
https://doi.org/10.1016/j.tree.2014.03.006

Carradec Q, Pelletier E, Da Silva C, Alberti A, Seeleuthner Y, Blanc-Mathieu R, Lima-Mendez G,
Rocha F, Tirichine L, Labadie K, Kirilovsky A, Bertrand A, Engelen S, Madoui M-A, Méheust R,
Poulain J, Romac S, Richter DJ, Yoshikawa G, Dimier C, Kandels-Lewis S, Picheral M, Searson
S, Jaillon O, Aury J-M, Karsenti E, Sullivan MB, Sunagawa S, Bork P, Not F, Hingamp P, Raes J,
Guidi L, Ogata H, de Vargas C, Iudicone D, Bowler C, Wincker P (2018) A global ocean atlas of
eukaryotic genes. *Nature Communications*, **9**, 373. https://doi.org/10.1038/s41467-017-02342-1

Close TJ, Bhat PR, Lonardi S, Wu Y, Rostoks N, Ramsay L, Druka A, Stein N, Svensson JT,
Wanamaker S, Bozdag S, Roose ML, Moscou MJ, Chao S, Varshney RK, Szűcs P, Sato K,
Hayes PM, Matthews DE, Kleinhofs A, Muehlbauer GJ, DeYoung J, Marshall DF, Madishetty K,
Fenton RD, Condamine P, Graner A, Waugh R (2009) Development and implementation of high-
throughput SNP genotyping in barley. *BMC Genomics*, **10**, 582. https://doi.org/10.1186/1471-
2164-10-582

Duarte CM (2015) Seafaring in the 21St Century: The Malaspina 2010 Circumnavigation Expedition.
Limnology and Oceanography Bulletin, **24**, 11–14. https://doi.org/10.1002/lob.10008

Eddy SR (2020) HMMER: biosequence analysis using profile hidden Markov models.
http://hmmer.org/

Eisen JA (2003) Phylogenomics: Intersection of Evolution and Genomics. *Science*, **300**, 1706–1707.
https://doi.org/10.1126/science.1086292

El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M, Richardson LJ, Salazar
GA, Smart A, Sonnhammer ELL, Hirsh L, Paladin L, Piovesan D, Tosatto SCE, Finn RD (2019)
The Pfam protein families database in 2019. *Nucleic Acids Research*, **47**, D427–D432.
https://doi.org/10.1093/nar/gky995

Gawryluk RMR, Tikhonenkov DV, Hehenberger E, Husnik F, Mylnikov AP, Keeling PJ (2019) Non-
photosynthetic predators are sister to red algae. *Nature*, **572**, 240–243.
https://doi.org/10.1038/s41586-019-1398-6

The Gene Ontology Consortium (2019) The Gene Ontology Resource: 20 years and still GOing
strong. *Nucleic Acids Research*, **47**, D330–D338. https://doi.org/10.1093/nar/gky1055

Grigoriev IV, Hayes RD, Calhoun S, Kamel B, Wang A, Ahrendt S, Dusheyko S, Nikitin R, Mondo SJ,
Salamov A, Shabalov I, Kuo A (2021) PhycoCosm, a comparative algal genomics resource.
Nucleic Acids Research, **49**, D1004–D1011. https://doi.org/10.1093/nar/gkaa898

Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, Riley R, Salamov A, Zhao X, Korzeniewski
F, Smirnova T, Nordberg H, Dubchak I, Shabalov I (2014) MycoCosm portal: gearing up for 1000
fungal genomes. *Nucleic Acids Research*, **42**, D699–D704. https://doi.org/10.1093/nar/gkt1183

Gruber-Vodicka HR, Seah BKB, Pruesse E (2020) phyloFlash: Rapid Small-Subunit rRNA Profiling
and Targeted Assembly from Metagenomes (M Arumugam, Ed,). *mSystems*, **5**.
https://doi.org/10.1128/mSystems.00920-20

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B,
Lieber M, MacManes MD, Ott M, Orvis J, Pochet N, Strozzi F, Weeks N, Westerman R, William T,
Dewey CN, Henschel R, LeDuc RD, Friedman N, Regev A (2013) De novo transcript sequence
reconstruction from RNA-seq using the Trinity platform for reference generation and analysis.
Nature Protocols, **8**, 1494–1512. https://doi.org/10.1038/nprot.2013.084

Huang X (1999) CAP3: A DNA Sequence Assembly Program. *Genome Research*, **9**, 868–877.
https://doi.org/10.1101/gr.9.9.868

Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR,
Letunic I, Rattei T, Jensen LJ, von Mering C, Bork P (2019) eggNOG 5.0: a hierarchical,

functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Research*, **47**, D309–D314. https://doi.org/10.1093/nar/gky1085

Janouškovec J, Tikhonenkov DV, Burki F, Howe AT, Rohwer FL, Mylnikov AP, Keeling PJ (2017) A New Lineage of Eukaryotes Illuminates Early Mitochondrial Genome Reduction. *Current Biology*, **27**, 3717-3724.e5. https://doi.org/10.1016/j.cub.2017.10.051

Kamikawa R, Kolisko M, Nishimura Y, Yabuki A, Brown MW, Ishikawa SA, Ishida K, Roger AJ, Hashimoto T, Inagaki Y (2014) Gene Content Evolution in Discobid Mitochondria Deduced from the Phylogenetic Position and Complete Mitochondrial Genome of Tsukubamonas globosa. *Genome Biology and Evolution*, **6**, 306–315. https://doi.org/10.1093/gbe/evu015

Karsenti E, Acinas SG, Bork P, Bowler C, De Vargas C, Raes J, Sullivan M, Arendt D, Benzoni F, Claverie J-M, Follows M, Gorsky G, Hingamp P, Iudicone D, Jaillon O, Kandels-Lewis S, Krzic U, Not F, Ogata H, Pesant S, Reynaud EG, Sardet C, Sieracki ME, Speich S, Velayoudon D, Weissenbach J, Wincker P (2011) A Holistic Approach to Marine Eco-Systems Biology. *PLoS Biology*, **9**, e1001177. https://doi.org/10.1371/journal.pbio.1001177

Keeling PJ, Burki F, Wilcox HM, Allam B, Allen EE, Amaral-Zettler LA, Armbrust EV, Archibald JM, Bharti AK, Bell CJ, Beszteri B, Bidle KD, Cameron CT, Campbell L, Caron DA, Cattolico RA, Collier JL, Coyne K, Davy SK, Deschamps P, Dyhrman ST, Edvardsen B, Gates RD, Gobler CJ, Greenwood SJ, Guida SM, Jacobi JL, Jakobsen KS, James ER, Jenkins B, John U, Johnson MD, Juhl AR, Kamp A, Katz LA, Kiene R, Kudryavtsev A, Leander BS, Lin S, Lovejoy C, Lynn D, Marchetti A, McManus G, Nedelcu AM, Menden-Deuer S, Miceli C, Mock T, Montresor M, Moran MA, Murray S, Nadathur G, Nagai S, Ngam PB, Palenik B, Pawlowski J, Petroni G, Piganeau G, Posewitz MC, Rengefors K, Romano G, Rumpho ME, Rynearson T, Schilling KB, Schroeder DC, Simpson AGB, Slamovits CH, Smith DR, Smith GJ, Smith SR, Sosik HM, Stief P, Theriot E, Twary SN, Umale PE, Vaulot D, Wawrik B, Wheeler GL, Wilson WH, Xu Y, Zingone A, Worden AZ (2014) The Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP): Illuminating the Functional Diversity of Eukaryotic Life in the Oceans through Transcriptome Sequencing (RG Roberts, Ed,). *PLoS Biology*, **12**, e1001889. https://doi.org/10.1371/journal.pbio.1001889

King N, Rokas A (2017) Embracing Uncertainty in Reconstructing Early Animal Evolution. *Current Biology*, **27**, R1081–R1088. https://doi.org/10.1016/j.cub.2017.08.054

Kiss E, Hegedüs B, Virágh M, Varga T, Merényi Z, Kószó T, Bálint B, Prasanna AN, Krizsán K, Kocsubé S, Riquelme M, Takeshita N, Nagy LG (2019) Comparative genomics reveals the origin of fungal hyphae and multicellularity. *Nature Communications*, **10**. https://doi.org/10.1038/s41467-019-12085-w

Kück P, Longo GC (2014) FASconCAT-G: extensive functions for multiple sequence alignment preparations concerning phylogenetic studies. *Frontiers in Zoology*, **11**, 81. https://doi.org/10.1186/s12983-014-0081-x

Larkin AA, Garcia CA, Garcia N, Brock ML, Lee JA, Ustick LJ, Barbero L, Carter BR, Sonnerup RE, Talley LD, Tarran GA, Volkov DL, Martiny AC (2021) High spatial resolution global ocean metagenomes from Bio-GO-SHIP repeat hydrography transects. *Scientific Data*, **8**, 107. https://doi.org/10.1038/s41597-021-00889-9

Lax G, Eglit Y, Eme L, Bertrand EM, Roger AJ, Simpson AGB (2018) Hemimastigophora is a novel supra-kingdom-level lineage of eukaryotes. *Nature*, **564**, 410–414. https://doi.org/10.1038/s41586-018-0708-8

Letunic I, Bork P (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Research*, **47**, W256–W259. https://doi.org/10.1093/nar/gkz239

Li W, Godzik A (2006) Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics*, **22**, 1658–1659. https://doi.org/10.1093/bioinformatics/btl158

Li Y, Steenwyk JL, Chang Y, Wang Y, James TY, Stajich JE, Spatafora JW, Groenewald M, Dunn CW, Hittinger CT, Shen X-X, Rokas A (2021) A genome-scale phylogeny of the kingdom Fungi. *Current Biology*, **31**, 1653-1665.e5. https://doi.org/10.1016/j.cub.2021.01.074

MacManes MD (2014) On the optimal trimming of high-throughput mRNA sequence data. Frontiers in Genetics, 5, 13:1-7. https://doi.org/10.3389/fgene.2014.00013

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM (2021) BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes (J Kelley, Ed,). *Molecular Biology and Evolution*, **38**, 4647–4654. https://doi.org/10.1093/molbev/msab199

Marron AO, Ratcliffe S, Wheeler GL, Goldstein RE, King N, Not F, de Vargas C, Richter DJ (2016) The Evolution of Silicon Transport in Eukaryotes. *Molecular Biology and Evolution*, **33**, 3226–

3248. https://doi.org/10.1093/molbev/msw209

Menardo F, Loiseau C, Brites D, Coscolla M, Gygli SM, Rutaihwa LK, Trauner A, Beisel C, Borrell S,
Gagneux S (2018) Treemmer: a tool to reduce large phylogenetic datasets with minimal loss of
diversity. *BMC Bioinformatics*, **19**, 164. https://doi.org/10.1186/s12859-018-2164-8

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R (2020)
IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era (E
Teeling, Ed,). *Molecular Biology and Evolution*, **37**, 1530–1534.
https://doi.org/10.1093/molbev/msaa015

Mitchell AL, Attwood TK, Babbitt PC, Blum M, Bork P, Bridge A, Brown SD, Chang H-Y, El-Gebali S,
Fraser MI, Gough J, Haft DR, Huang H, Letunic I, Lopez R, Luciani A, Madeira F, Marchler-Bauer
A, Mi H, Natale DA, Necci M, Nuka G, Orengo C, Pandurangan AP, Paysan-Lafosse T, Pesseat
S, Potter SC, Qureshi MA, Rawlings ND, Redaschi N, Richardson LJ, Rivoire C, Salazar GA,
Sangrador-Vegas A, Sigrist CJA, Sillitoe I, Sutton GG, Thanki N, Thomas PD, Tosatto SCE, Yong
S-Y, Finn RD (2019) InterPro in 2019: improving coverage, classification and access to protein
sequence annotations. *Nucleic Acids Research*, **47**, D351–D360.
https://doi.org/10.1093/nar/gky1100

Neely CJ, Hu SK, Alexander H, Tully BJ (2021) The high-throughput gene prediction of more than
1,700 eukaryote genomes using the software package EukMetaSanity. *bioRxiv*.
https://doi.org/10.1101/2021.07.25.453296

Pertea G, Pertea M (2020) GFF Utilities: GffRead and GffCompare. *F1000Research*, **9**, 304.
https://doi.org/10.12688/f1000research.23297.2

Plotly Technologies Inc. (2015) Collaborative data science. https://plot.ly

Priyam A, Woodcroft BJ, Rai V, Moghul I, Munagala A, Ter F, Chowdhary H, Pieniak I, Maynard LJ,
Gibbins MA, Moon H, Davis-Richardson A, Uludag M, Watson-Haigh NS, Challis R, Nakamura H,
Favreau E, Gómez EA, Pluskal T, Leonard G, Rumpf W, Wurm Y (2019) Sequenceserver: A
Modern Graphical User Interface for Custom BLAST Databases. *Molecular Biology and Evolution*,
36, 2922–2924. https://doi.org/10.1093/molbev/msz185

Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A (2020) Using SPAdes De Novo
Assembler. *Current Protocols in Bioinformatics*, **70**. https://doi.org/10.1002/cpbi.102

Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO (2012) The
SILVA ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic Acids Research, **41**, D590–D596. https://doi.org/10.1093/nar/gks1219

Rice P, Longden I, Bleasby A (2000) EMBOSS: The European Molecular Biology Open Software
Suite. *Trends in Genetics*, **16**, 276–277. https://doi.org/10.1016/S0168-9525(00)02024-2

Richards T (2021) Thomas Cavalier-Smith (1942–2021). *Nature*, **593**, 190–190.
https://doi.org/10.1038/d41586-021-01250-1

Richter DJ, Levin TC (2019) The origin and evolution of cell-intrinsic antibacterial defenses in
eukaryotes. *Current Opinion in Genetics & Development*, **58–59**, 111–122.
https://doi.org/10.1016/j.gde.2019.09.002

Richter DJ, Watteaux R, Vannier T, Leconte J, Frémont P, Reygondeau G, Maillet N, Henry N, Benoit
G, Fernàndez-Guerra A, Suweis S, Narci R, Berney C, Eveillard D, Gavory F, Guidi L, Labadie K,
Mahieu E, Poulain J, Romac S, Roux S, Dimier C, Kandels S, Picheral M, Searson S, Tara
Oceans Coordinators, Pesant S, Aury J-M, Brum JR, Lemaitre C, Pelletier E, Bork P, Sunagawa
S, Karp-Boss L, Bowler C, Sullivan MB, Karsenti E, Mariadassou M, Probert I, Peterlongo P,
Wincker P, de Vargas C, Ribera d'Alcalà M, Iudicone D, Jaillon O, Tara Oceans Coordinators
(2012) Genomic evidence for global ocean plankton biogeography shaped by large-scale current
systems. *eLife*, **11**, e78129. https://doi.org/10.7554/eLife.78129

Roger AJ (2021) Thomas Cavalier-Smith (1942–2021). *Current Biology*, **31**, R977–R981.
https://doi.org/10.1016/j.cub.2021.07.009

Saldarriaga J (2021) Obituary for Professor Thomas Cavalier-Smith FRS, FRSC. *International Society*
for Evolutionary Protistology. https://www.isep-protists.com/post/obituary-for-professor-thomas-
cavalier-smith

Sayers EW, Beck J, Brister JR, Bolton EE, Canese K, Comeau DC, Funk K, Ketter A, Kim S, Kimchi
A, Kitts PA, Kuznetsov A, Lathrop S, Lu Z, McGarvey K, Madden TL, Murphy TD, O'Leary N,
Phan L, Schneider VA, Thibaud-Nissen F, Trawick BW, Pruitt KD, Ostell J (2020) Database
resources of the National Center for Biotechnology Information. *Nucleic Acids Research*, **48**, D9–
D16. https://doi.org/10.1093/nar/gkz899

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM (2015) BUSCO: assessing
genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, **31**,
3210–3212. https://doi.org/10.1093/bioinformatics/btv351

Steenwyk JL, Buida TJ, Li Y, Shen X-X, Rokas A (2020) ClipKIT: A multiple sequence alignment
trimming software for accurate phylogenomic inference (A Hejnol, Ed,). *PLOS Biology*, **18**,
e3001007. https://doi.org/10.1371/journal.pbio.3001007
Strassert JFH, Jamy M, Mylnikov AP, Tikhonenkov DV, Burki F (2019) New Phylogenomic Analysis of
the Enigmatic Phylum Telonemia Further Resolves the Eukaryote Tree of Life (B Shapiro, Ed,).
Molecular Biology and Evolution, **36**, 757–765. https://doi.org/10.1093/molbev/msz012
Strassert JFH, Monaghan MT (2022) Phylogenomics unravels the early diversification of fungi.
Current Biology, **32**, 3628-3635.e3. https://doi.org/10.1016/j.cub.2022.06.057
Tara Oceans Coordinators, Sunagawa S, Acinas SG, Bork P, Bowler C, Eveillard D, Gorsky G, Guidi
L, Iudicone D, Karsenti E, Lombard F, Ogata H, Pesant S, Sullivan MB, Wincker P, de Vargas C
(2020) Tara Oceans: towards global ocean ecosystems biology. *Nature Reviews Microbiology*.
https://doi.org/10.1038/s41579-020-0364-5
Tice AK, Žihala D, Pánek T, Jones RE, Salomaki ED, Nenarokov S, Burki F, Eliáš M, Eme L, Roger
AJ, Rokas A, Shen X-X, Strassert JFH, Kolísko M, Brown MW (2021) PhyloFisher: A
phylogenomic package for resolving eukaryotic relationships (A Hejnol, Ed,). *PLOS Biology*, **19**,
e3001365. https://doi.org/10.1371/journal.pbio.3001365
Wang H-C, Minh BQ, Susko E, Roger AJ (2018) Modeling Site Heterogeneity with Posterior Mean
Site Frequency Profiles Accelerates Accurate Phylogenomic Estimation. *Systematic Biology*, **67**,
216–235. https://doi.org/10.1093/sysbio/syx068
Wickett NJ, Mirarab S, Nguyen N, Warnow T, Carpenter E, Matasci N, Ayyampalayam S, Barker MS,
Burleigh JG, Gitzendanner MA, Ruhfel BR, Wafula E, Der JP, Graham SW, Mathews S,
Melkonian M, Soltis DE, Soltis PS, Miles NW, Rothfels CJ, Pokorny L, Shaw AJ, DeGironimo L,
Stevenson DW, Surek B, Villarreal JC, Roure B, Philippe H, dePamphilis CW, Chen T, Deyholos
MK, Baucom RS, Kutchan TM, Augustin MM, Wang J, Zhang Y, Tian Z, Yan Z, Wu X, Sun X,
Wong GK-S, Leebens-Mack J (2014) Phylotranscriptomic analysis of the origin and early
diversification of land plants. *Proceedings of the National Academy of Sciences*, **111**, E4859–
E4868. https://doi.org/10.1073/pnas.1323926111
Wilkinson MD, Dumontier M, Aalbersberg IjJ, Appleton G, Axton M, Baak A, Blomberg N, Boiten J-W,
da Silva Santos LB, Bourne PE, Bouwman J, Brookes AJ, Clark T, Crosas M, Dillo I, Dumon O,
Edmunds S, Evelo CT, Finkers R, Gonzalez-Beltran A, Gray AJG, Groth P, Goble C, Grethe JS,
Heringa J, 't Hoen PAC, Hooft R, Kuhn T, Kok R, Kok J, Lusher SJ, Martone ME, Mons A, Packer
AL, Persson B, Rocca-Serra P, Roos M, van Schaik R, Sansone S-A, Schultes E, Sengstag T,
Slater T, Strawn G, Swertz MA, Thompson M, van der Lei J, van Mulligen E, Velterop J,
Waagmeester A, Wittenburg P, Wolstencroft K, Zhao J, Mons B (2016) The FAIR Guiding
Principles for scientific data management and stewardship. *Scientific Data*, **3**, 160018.
https://doi.org/10.1038/sdata.2016.18
Yabuki A, Kamikawa R, Ishikawa SA, Kolisko M, Kim E, Tanabe AS, Kume K, Ishida K, Inagki Y
(2015) Palpitomonas bilix represents a basal cryptist lineage: insight into the character evolution
in Cryptista. *Scientific Reports*, **4**. https://doi.org/10.1038/srep04641
760