1
2
3 # Chronic Pb^{2+} Exposure Causes Hippocampal Network Hypersynchrony, Absence Seizures, and Sensorimotor Deficits
9 Abbreviated Title: LEAD EXPOSURE CAUSES HIPPOCAMPAL NETWORK HYPERSYNCHRONY
10
11
12
13

14 Nathan W. Schultheiss[1,5], Jennifer L. McGlothan[2,3], Tomás R. Guilarte[2,3*], Timothy A. Allen[1,2,4*]
15
[1]Department of Psychology, Florida International University, Miami, FL, 33199
[2]Department of Environmental Health Sciences, Robert Stempel College of Public Health, Florida International University, Miami,
FL, 33199
[3]Brain, Behavior & the Environment Program, Florida International University, Miami, FL, 33199
[4]Cognitive Neuroscience Program, Florida International University, Miami, FL, 33199
[5]Brain Institute, Nicklaus Children's Hospital, Miami, FL 33135

22
23
24
Number of figures: 5
Number of supplemental figures: 7
Number of supplemental tables: 1
Number of tables: 0
Number of text pages: 46
Number of Lines: 1071
Total character count (with spaces):
Title: 108 (12 words)
Abstract: 1944 (270 words)
Introduction: 7,597 (1,098 words)
Methods: 21419 (3,272 words)
Results: 22,882 (3,439 words)
Discussion: 9,033 (1,264 words)
References: 91
39

40 *Corresponding Authors:
41
42   Timothy A. Allen, PhD
43   Department of Psychology
44   email: tallen@fiu.edu
45
46   Tomás R. Guilarte, PhD
47   Department of Environmental Health Sciences
48   email: tguilart@fiu.edu
49
50   Address:
51   Florida International University
52   11200 SW 8th Street
53   Miami, FL, 33199

54
55
56 **Conflict of Interest:** The authors declare no competing financial interests.
57

Acknowledgements: This work was supported by R01 ES006189 and R01 ES035827 to T.R.G. and R01 MH113626 to T.A.A. We
thank the members of the Guilarte Lab and the Allen Lab for their assistance with experiments and for providing constructive
feedback on the manuscript. Acknowledging AI: ChatGPT-4-turbo (OpenAI) was used sparingly to help with snippets of MATLAB
scripts, SciSpace was used to help with literature searches, and Grammarly was used to edit text for punctuation and grammar. AI
was not used to generate content.

Keywords: hippocampus, gamma, schizophrenia, lead neurotoxicity, pre-pulse inhibition

Abstract

Chronic early-life lead (Pb^{2+}) exposure contributes to an array of cognitive and behavioral dysfunctions,

including impaired attention, memory, and intellectual abilities, in addition to increased social delinquency.

Notably, Pb^{2+} exposure is an environmental risk factor for adult psychopathologies, including schizophrenia

and epilepsy. Neurobiologically, Pb^{2+} is a potent N-methyl-D-aspartate receptor (NMDAR) antagonist, and

exposure during early life elicits a cascade of cellular neurotoxic effects that alter neurodevelopmental

trajectories. This includes reduced parvalbumin-expressing interneurons in the hippocampus (HC) and

altered synaptic transmission. Little is known, however, about the impact of chronic Pb^{2+} exposure on HC

network dynamics, which link cellular-molecular effects with cognitive-behavioral consequences. Here, we

tested the impact of chronic Pb^{2+} exposure on the HC local field potential (LFP) in freely behaving rats. We

found that Pb^{2+} exposure (1) caused a striking level of theta rhythmic hypersynchrony, (2) amplified fast

gamma synchronization, (3) disrupted behavioral modifications of theta and fast gamma, and (4)

exacerbated absence seizures appearing in the LFP as spike-wave discharges (SWDs) at theta

frequencies. Each of these rhythmic changes in the HC network was related to exploratory movements in

the open field. HC network alterations like these have also been linked to impaired prepulse inhibition of

the acoustic startle reflex (PPI). Thus, next, we tested the effect of Pb^{2+} exposure on PPI. We found that

adult males (PN50 and 120), but neither females nor juvenile males, showed reduced PPI, recapitulating

sex dependencies on PPI disruptions in schizophrenics. Altogether, these results are consistent with the

hypothesis that chronic early-life Pb^{2+} exposure causes dysfunction in the rhythmic network coordination of

the HC, limiting processing, and helping to account for cognitive deficits.

Introduction

Lead (Pb^{2+}) is a pervasive pollutant and potentially lethal toxin (Pb^{2+} tox profile for ATSDR, 2020) that

accounts for 0.6% of the total global cost of disease (Sample, 2021). Even low-level exposure to Pb^{2+}

during early childhood causes permanent neurological dysfunctions and may be a risk factor for

schizophrenia (SZ)(Guilarte et al., 2012) or epilepsy (Sasmaz et al., 2003). Since the late 1970s, public

health policy to eliminate Pb^{2+} from gasoline, paint, and other consumer products has helped lower blood

Pb^{2+} levels (BLL) in the general US population (Dignam et al., 2019). Still, a shocking proportion of children

and adults, particularly in disadvantaged urban communities, continue to be exposed to Pb^{2+} due to the

legacy of millions of tons of environmental contamination (Bernard & McGeehin, 2003; Yeter et al., 2020).

As recently as 2002, the healthcare cost-burden of Pb^{2+} exposure in the US was 2.2% of the total national

cost of healthcare (Wakefield, 2002); but numerous *ongoing* cases of significant Pb^{2+} exposure from

municipally-regulated resources, e.g. public water (Hanna-Attisha et al., 2016) and affordable housing

(Garnett & Iannuzzi, 2020), show that exposure risks are more extensive than previously appreciated

(https://www.reuters.com/investigates/special-report/usa-lead-testing/). Recent US estimates of children

with BLLs above the current Centers for Disease Control (CDC) reference level (10 μg/dL) exceed one-

quarter million (e.g., Egan et al., 2021). Global estimates exceed one-half billion children, as detailed in the

2020 UNICEF report entitled: *The Toxic Truth-Children's Exposure to Lead Pollution Undermines a*

Generation of Future Potential (UNICEF, 2020).

Childhood exposure to Pb^{2+} alters brain development, leading to impoverished cognitive abilities, as

well as social, emotional, and behavioral dysregulations that persist in adulthood (Lanphear et al., 2005;

2006; Canfield et al., 2003; Needleman et al., 2002; 2004; Miranda et al., 2007; Caito & Aschner, 2017).

Young children are disproportionately vulnerable to Pb^{2+} neurotoxicity (Canfield et al., 2003) because their

predisposition for exploratory behaviors increases exposures, and the immature blood-brain barrier is

permissive of Pb^{2+} penetration, leading to markedly higher brain concentrations (Barltrop, 1969; Rossouw

et al., 1987). Chelation therapy to reduce elevated BLLs can help mitigate the Pb^{2+} burden on the body, but

it does not affect cognitive or neuropsychological impairments (Rogan et al., 2001, Dietrich et al., 2004).

Notably, there are no proven therapies to ameliorate the neurotoxic damage caused by childhood Pb^{2+}

exposure once it has been sustained, and exposure prevention is the only available public health strategy

(CDC, 2021).

In the brain, Pb^{2+} is a potent antagonist of NMDARs at excitatory synapses in the HC and medial

prefrontal cortex (mPFC) (Guilarte & Miceli, 1993; Guilarte, 1997; 2023). Pb^{2+} binds to allosteric sites on

NMDARs (Gavazzo et al., 2008), causing receptor hypoactivity and reduced calcium influx (Hashemzadeh-

Gargari & Guilarte, 1999), diminishing kinase activity and nuclear CREB phosphorylation (Nihei et al.,

2000; Toscano et al., 2002; Toscano et al., 2005; Toscano & Guilarte, 2005); and attenuating synaptic

levels of brain-derived neurotrophic factor (BDNF) (Neal et al., 2010; Neal & Guilarte, 2010; Stansfield et

al., 2012). Significantly, developmental Pb^{2+} exposure alters the receptor-subunit composition of NMDARs

(Guilarte & McGlothan, 1998; Nihei et al., 2000), leading to impaired long-term potentiation (LTP) (Nihei et

al., 2000) and long-term depression (LTD) (Zhao et al, 1999) in the HC, reduced HC spine density (Zou et

al., 2023), and reduced entorhinal cortex inputs to the HC (Zou et a., 2023). This ultimately leads to HC-

dependent cognitive impairments (Kuhlmann et al., 1997; Jett et al., 1997; Munoz et al., 1988; Jaako-

Movits et al., 2005; McGlothan et al., 2008; Wang et al., 2016). It is unknown how Pb^{2+} impacts HC network

dynamics to mediate the effects of cellular pathologies on cognitive functions.

Beyond the direct effects of Pb^{2+} neurotoxicity, early-life Pb^{2+} exposure is further implicated in the

development of SZ (Opler et al., 2004; 2008). A current leading hypothesis proposes that Pb^{2+} exposure

escalates the risk for SZ through interactions of neurotoxic mechanisms with genetic and epigenetic factors

at critical developmental stages (Guilarte et al., 2012). Supporting this view, early-life Pb^{2+} levels of patients

diagnosed with schizophrenia have been found to significantly exceed those of control subjects, and

postmortem assays of confirmed SZ patients have shown selective elevation of Pb^{2+} levels (Modabbernia

et al., 2016). Additionally, Pb^{2+} exposure worsens behavioral, cognitive, and sensorimotor dysfunctions in

animal models of SZ (Abazyan et al., 2014).

Interestingly, at a cellular level, developmental Pb^{2+} exposure recapitulates a hallmark of schizophrenia

pathology by causing a reduction in the number of parvalbumin-expressing GABAergic interneurons

(PVGIs) in the HC and agranular mPFC (Hashimoto et al., 2003; Konradi et al., 2011; Stansfield et al.,

2015). These fast-spiking PVGIs are a critical component of mechanisms underlying gamma oscillations

(25-120 Hz) that coordinate neuronal assemblies. Gamma oscillations stem from reciprocal inhibitory-

excitatory (I-E) interactions of principal neurons nested within the local microcircuitry of inhibitory networks.

I-E interactions also give rise to the HC theta rhythm (6-10 Hz) that coordinates gamma oscillations nested

within each theta cycle (Tamura et al., 2017). Misbalance in these network dynamics possibly accounts for

comorbid evidence from electroencephalographic (EEG) showing systems-level Pb^{2+} neurotoxicity

disruptions (Kmiecik-Malecka et al., 2009; Govoni et al., 1988). In young children, low levels of Pb^{2+}

correlate to slow-wave activity in sensory-evoked EEG data and overall theta power (Otto et al., 1981;

Poblano et al., 2001). Additionally, Pb^{2+} exposure may also relate to some forms of epilepsy (Sasmaz et

al., 2003) and increased seizures (Arrieta et al., 2005), further implicating network-level physiological

disruptions as a critical level of analysis for chronic early life Pb^{2+} exposure.

Here, we explored the effects of Pb^{2+} exposure on network activity in the HC. To this aim, we performed

chronic *in vivo* recordings (silicon arrays) in a rodent model of chronic early-life Pb^{2+} exposure (CELLE)(see

Albores-Garcia et al. 2021). We evaluated delta, theta, and gamma frequency synchronization in HC

networks (Schultheiss et al. 2020). We show that Pb^{2+} exposure caused *hypersynchrony* of the HC theta

rhythm, an amplification of gamma oscillations, a disruption in behavior-related theta and gamma, and an

increase in SWD indicative of absence seizures. Next, we tested the same Pb^{2+} exposed rat model (males

and females) in PPI, a sensitive sensorimotor gating measure dependent on HC functions and related to

SZ. We found that only male rats had significantly impaired sensorimotor gating, consistent with the Pb^{2+}

exposure-induced HC network disruptions, SZ, and recapitulating sex differences in schizophrenics tested

on a human version of the PPI task (e.g., Mena et al., 2016). We propose that theta hypersynchrony, theta-

gamma activity patterns, and absence seizures represent key network dysfunction caused by early-life Pb^{2+}

neurotoxicity, and likely contribute to associated developmental neuropathologies, including schizophrenia

and childhood-onset epilepsies.

163   **Methods**

Subjects. Eleven adult Long-Evans rats (males) were implanted with commercially available silicon

probes (NeuroNexus) and used in the electrophysiological experiments. In one rat, the probe was located

in the lateral ventricle adjacent to the HC and was excluded. In another rat, tetrode signals were poor, and

the rat was also excluded, leaving nine rats with good implants. Five rats were chronically exposed to Pb^{2+}

(see below), and four were controls. Four additional adult male Long-Evans rats were implanted with

custom-built, dual-site implants for simultaneous recordings from the HC and mPFC. Two dual-site rats

were chronically exposed to Pb^{2+}, and two were controls. These rats were only used to look at the

relationship, if any, between HC SWDs and PFC activity. Two hundred thirty Long-Evans rats were used in

pre-pulse inhibition (PPI) experiments. These rats were evenly divided into twelve groups (18-21 per group)

corresponding to all combinations of (1) Pb^{2+} exposure and controls, (2) males and females, and (3) three

ages at the time of testing.

All rats were individually housed in a climate-controlled vivarium on a reversed 12hr light/dark cycle,

and experiments were conducted during dark cycle active periods. Rats had free access to food and water

in their home cages, but no food or water was available during behavioral sessions. Rats with dual-site

implants, but not rats in the other groups, received rewards (~¼ Fruit Loop) during behavioral sessions

delivered sporadically to augment rats' motivation for exploratory behavior.

Experiments were conducted per the Guide for the Care and Use of Laboratory Animals (National

Institutes of Health) following protocols approved by Florida International University's Institutional Animal

Care and Use Committee (FIU IACUC). Data from the six control rats used here had been previously

analyzed to establish the relationships of HC spectral modes to behaviors (Schultheiss et al., 2020). The

present study focused on the effects of Pb^{2+} exposure, and all analyses presented here are new.

Breeding and chronic Pb^{2+} exposure. Adult female rats (225-250g) (Charles River Laboratories) were

randomly assigned to a Prolab RMH-1000 diet containing 0 (control) or 1500ppm lead acetate (PbAc)

(Dyets, Inc., Bethlehem, PA). After 10-14 days of acclimation to the diet, each dam was paired with a non-

exposed adult male Long Evans rat (250-275g; Charles River Laboratories) to breed over 3 days. Litters

were culled to 10 pups on postnatal day one (PN1). Dams were maintained on their respective diet, and at

PN21 pups were weaned onto the same diet and maintained for the duration of the experiments. Blood

Pb^{2+} levels of rats used in PPI experiments (SuppTab1) were measured in samples obtained from cardiac

puncture under sodium pentobarbital anesthesia. Samples were prepared and measured following

protocols provided using the LeadCare Plus system (Magellan Diagnostics, N. Billerica, MA).

Electrodes. For most electrophysiological experiments reported here, silicon probes with 32 electrodes

arranged as eight tetrodes were surgically implanted to record LFPs from the dorsal HC. The eight tetrodes

were distributed across four probe shanks at tip-to-tetrode depths of 78 µm and 228 µm (NeuroNexus

A4X2-tet-5mm). Shanks were separated by 200 µm, giving each probe a total length of 0.67 cm, which was

oriented medial-lateral when implanted to sample along the proximal-distal axis near the CA1 subregion of

HC. Electrode impedances were measured to be 1.23 ± 0.32 MΩ (@1000Hz).

Custom-made dual-site implants housed 32 stainless steel wire electrodes (0.42 ± 0.25 MΩ) coupled to

an integrated electrode interface board within a 3-D printed implant body (Autodesk Inventor; 3D Systems

ProJet 1200). The electrode layout for these implants consisted of two grids, such that 18 electrodes

targeted HC (2 x 2.5 mm grid, four electrode rows, 0.5 mm spacing), and 14 electrodes were aligned to the

rostrocaudal axis of mPFC (4 x 1 mm grid, two rows, 0.5 mm spacing). HC sites were used in the primary

analyses, while mPFC sites were used in supplemental analyses.

Surgical procedures. Rats used in electrophysiological experiments weighed 600-700 g at the time of

surgery. General anesthesia was induced with isoflurane (5%) mixed with oxygen (0.8 L/min) that was

continuously delivered throughout surgery at 1-4% as needed. Body temperature was monitored

throughout surgery, and a Ringer's solution (5% dextrose) was given periodically to maintain hydration (1

ml increments every 60-90 minutes). Once the hair over the scalp was cut, rats were placed in a

stereotaxic device, and glycopyrrolate (0.5 mg/kg, s.c.) was administered to assist respiratory stability.

Ophthalmic ointment was then applied to the eyes. Four injections of marcaine were made to the scalp

(~0.1 ml, 7.5 mg/ml, s.c.), and a rostrocaudal incision was made, exposing the skull. The pitch of the skull

was leveled between bregma and lambda, two stainless steel ground screws were placed in the left

parietal bone, and five titanium support screws were anchored to the skull. For silicon probe implants

targeting dorsal HC, rectangular craniotomies were drilled to accommodate the probe shanks centered on

coordinates AP -3.2 mm, ML 2.7 mm. For dual-site implants, craniotomies were shaped to accommodate

the respective electrode grids targeting HC (AP -2.9 mm, ML 2.0 mm) and mPFC (AP 3.0 mm, ML

1.0 mm).

After removal of the dura, implants were lowered on the stereotaxic arm until the tips of the probe

shanks (or electrode wires) were just above the cortical surface. Then, the ground wire was attached to the

ground screws, and implants were lowered further until the electrodes reached a depth of ~2.8 mm below

the cortical surface. A thin layer of sodium alginate was applied in the craniotomy to protect the exposed

surface of the brain, and dental cement was used to secure the implant to the skull screws permanently.

Neosporin® was applied to the skin surrounding the implant, and flunixin (2.5 mg/kg, s.c.) was given for

analgesia. Before being returned to their home cages after surgery, rats were allowed to rest on a heating

pad until ambulatory. Neosporin® and flunixin were given again on the day following surgery, and rats were

monitored closely for five days to maintain body weight. Daily Baytril injections (s.c.) were also given for

five days to protect against postsurgical infection. A minimum of one week was reserved for recovery

before beginning experiments.

Behavioral setup for electrophysiological recordings. All electrophysiological experiments were

conducted in a large open-field environment (122 x 118 x 47 cm). A custom-made white cutting board

served as the floor of the behavioral arena, and the interior walls were lined with white vinyl. The arena was

mounted on posts 72 cm above the floor, and the exterior was wrapped in grounded Faraday shielding.

Two video cameras with a bird's eye view were used for behavioral tracking. These were mounted 28 cm

apart and 150 cm above the arena floor, providing ~4.6 pixel/cm resolution in recorded videos and

corresponding tracking data.

The entire behavioral apparatus was enshrouded with black curtains suspended from the ceiling and

extending to the floor. Small gaps at the Velcro® closures between the four curtain panels were used to

observe rats covertly to monitor that the head stage, cables, and elastic tethers were not tangled and to

confirm that rats were awake during periods of relative inactivity. All room lights were kept off during

recording sessions, and the white arena was illuminated dimly with red light from LED strips mounted at the

level of the video cameras above. During behavioral sessions, rats were placed on the arena floor and

recorded without any intervention except for infrequent behavioral checks or to adjust cables or

connections.

Electrophysiological recordings. Throughout each experiment, wideband neural data were acquired

with a Plexon OmniplexD recording system with A/D conversion at 40 kHz (16-bit). A digital head stage

processing subsystem (Intan, 32 channel) passed continuous wideband data referenced against the

ground screws (low cutoff = 0.7 Hz) to a hard drive using PlexControl data acquisition software. For

analysis of LFP recordings, wideband data were downsampled to 2 kHz. A maximum of one channel per

tetrode was included in our analyses, and several tetrodes across rats were excluded due to poor signal.

With these restrictions, 30 channels recorded from Pb^{2+}-exposed rats and 30 channels from control rats

were included in our analysis of HC LFP power spectra during running and stationary behaviors. These 60

channels were selected from among the 288 total recorded channels using a custom algorithm designed to

maximize signal quality by choosing for each tetrode the channel with the highest theta-delta ratio during

running (t-d$_{run}$) when theta would be expected to be strongest and delta weakest. Channel selection for

analyses of SWD (detailed below) was done manually through the inspection of several basic features,

such as the lack of evident noise in the voltage plots, the presence of separable theta and delta modes

throughout the session, and the ability to separate SWD-related theta and theta harmonics and running-

related theta and theta harmonics.

Behavioral tracking. The physical and software settings for the two video cameras used for behavioral

tracking were tuned independently to optimize detection of body contours on one camera and to optimize

tracking of head-mounted LEDs on the other. Using Cineplex software (Plexon), tracking data and video

frame captures (30 fps) were synchronized in real time to the simultaneously recorded neural data via a

central timing board. LED tracking data were collected online during experiments, and body contour

tracking data were derived offline from recorded videos.

LED tracking entailed tabulating the coordinates of two clusters of LED florets (green and blue) that

were offset ~2 cm laterally from the head stage at ~3 cm above the rat's head. Each LED cluster comprised

three florets oriented upward and slightly outward. This construction enabled consistent LED tracking

despite behaviors in which rats oriented their heads away from the level plane. The location of the head

was calculated as the midpoint between the two LED clusters for each time point. Body location was

calculated from difference images obtained by subtracting an image of the empty arena from each frame

capture during an experiment. Movement speeds were calculated from these body coordinates, except that

head location was used to calculate movement speeds in our seizure detection protocol (below).

Co-registration of LED and body tracking data. Following behavioral data acquisition, all analyses were

performed with custom-written MATLAB scripts (MathWorks, Natick, MA). First, the coordinate system for

camera one was mapped onto the coordinate system for camera two, enabling the integration of the two

types of tracking data. Next, for each timestep, we derived rats' head direction (HD) relative to the

environment (from the axis between the two head-mounted LED clusters); body direction (BD) relative to

the environment (from the midpoint between the LEDs to the center of the body); head angle (HA) relative

to the body (using the LED axis relative to BD); and the rotational velocity of the body (from changes in

BD).

Definitions of running and stationary bouts. Rats exhibit a wide range of behaviors in the open field,

including frequent rearing and grooming behaviors that incorporate fast body and head movements. These

movements manifest in tracking data as changes in body and LED coordinates, just as running does. To

isolate bouts of locomotion from other behaviors, we first required that movement speed be maintained

above 5 cm/sec for a minimum duration of 1.025 or 2.05 seconds (in parallel analyses) with a minimum

peak speed of 15 cm/sec. These minimum bout durations were chosen as a compromise between the

relatively fast timescale of rats' locomotor behaviors (e.g., rapid transitions between brief periods of

continuous running and interspersed pauses), the frequency resolution of power spectra, and the reduction

in number of oscillatory cycles (particularly for delta) contributing to amplitude and power estimates from

shorter time windows. We then eliminated all instances in which the head angle exceeded 35 degrees or

body rotational velocity exceeded 52.5 degrees per second at any time. Stationary bouts were required to

meet the same criteria for minimum duration, HA, and rotational velocity, and movement speeds were

required to remain below 5 cm/sec.

Measures of oscillatory synchronization. We analyzed delta-, theta-, and slow and fast gamma-

frequency synchronization (1-4 Hz, 6-10 Hz, 25-55 Hz, and 65-120 Hz, respectively) using short-time

power spectra (PSDs) of the HC LFP, as well as peak-to-trough measurements of individual oscillatory

cycles of the respective band-pass filtered LFPs. We also derived a continuous time series of

instantaneous amplitudes for each band by resampling cycle-by-cycle amplitude measures to 2 kHz. This

transform was applied in our analyses of speed modulation and cross-frequency correlations to mitigate the

vanishing likelihood that longer oscillatory cycles will elapse precisely within the time course of brief

behavioral bouts.

Boutwise analyses. We calculated the spectral content of HC recordings using Welch's method for

each stationary and running bout. Estimates of delta and theta power from PSDs for individual behavioral

bouts were defined as the mean values across the respective frequency ranges. Delta and theta peak

frequency and power estimates were measured directly from PSDs for behavioral bouts. For each

recording site, delta and theta power were defined as the mean values across all bouts of each type. To

determine whether Pb^{2+} exposure qualitatively impacts the behavioral modulation of power spectra, for

each rat, we subtracted the mean PSD for stationary bouts from the mean running spectrum. These

difference PSDs were then normalized either by the area of the mean stationary PSD, or by calculating the

percent change at each frequency. We also compiled the combined distributions of cycle amplitudes for

each frequency band across all rats in each group during all bouts of each type across all rats in each

group.

Sliding time-window analyses. To elaborate on bout-wise comparisons, we averaged movement

speeds and continuous amplitude time series for each frequency band using a two-second sliding window

(50% overlap) across each session for each rat. We then averaged oscillatory amplitudes within 5 cm/sec

movement speed bins up to 75 cm/sec. This approach, which is typical for evaluating speed modulation of

theta, ignores the distinction between running and movements during other behaviors, but it captures the

full nonlinear relationships of synchronization in each frequency band across the continuous range of

movement speeds. We calculated pairwise correlation coefficients between speed and amplitude

measures for each frequency band, and between bands.

Peak speed-triggered analyses. In a last series of analyses, we focused on the secondary regime of

running speed modulation by triggering cycle-by-cycle amplitudes on local peaks in running speed (>40

cm/sec) separated by at least 10 seconds. We then averaged amplitude measures within 0.5-second bins

spanning -9.75 to 9.75-second latencies. Because behavioral variability increases at greater temporal

offsets from speed peaks, the fidelity of relationships between time-binned cycle amplitudes and running

speed is maximal at the shortest latencies. We focused our initial statistical analyses on the 5 speed bins

spanning the 2.5-second window from -2.25 to 0.25-second latencies. We compared unnormalized cycle

amplitudes for each frequency band between Pb^{2+}-exposed rats and controls. We z-normalized cycle

amplitudes in two ways, in separate analyses, to address different effects of Pb^{2+} on synchronization. To

balance the contributions of each recording site to group-wise comparisons, we z-normalized oscillatory

amplitudes in each frequency band across the duration of each behavioral session. Next, we z-normalized

amplitudes across each peak-triggered analysis window separately before averaging within bins. By

eliminating the implicit assumption that amplitude measures are stable throughout behavioral sessions, this

approach had the effect of drawing running speed modulation of amplitudes into greater relief. That is,

changes in amplitude accompanying changes in speed were normalized relative to temporally local

amplitudes only, revealing a novel and reliable pattern of slow fluctuations in theta and gamma

synchronization that unfolded over the several seconds preceding and following speed peaks.

Absence seizure detection. Although seizures are rare and highly variable across rats, sessions, and

strains, rodents are a universal model for synchronized SWDs that dominate the LFP across the cortex

during non-convulsive seizures in the absence of epilepsy (e.g., van Luijelaar et al., 2014). SWDs occurred

overwhelmingly at times when rats were stationary, supporting the view that SWDs are the physiological

manifestations of 'absence' seizures. We estimated the incidence and duration of seizures by the

experimenter's hand measurements. Seizure parameters differed between the Pb^{2+}-exposed and control

groups, but biased judgment could manufacture the same effects. We developed a novel seizure detection

algorithm based on the frequency composition of the short time-windowed LFP to (1) obtain unbiased

estimates of the incidence of SWDs, and to (2) compare seizure parameters between Pb^{2+}-exposed rats

and controls. Using windows of two seconds (one-second overlap), we first derived power estimates within

a 2 Hz frequency range centered on the frequency of peak theta power (5-14 Hz) and within 2 Hz ranges

centered on the first and second harmonics of the theta-peak frequency. We took the z-scored values

across all time windows for each frequency band throughout each session. Negative values in these time

series were zeroed, yielding metrics of elevated theta and theta harmonic power. These metrics and the

time-averaged LFP amplitude were smoothed with a Gaussian, and their product was z-scored to give a

timeseries that was effective at identifying most SWDs of reasonable strength. We categorized detection

peaks as SWDs when they exceeded threshold values of 2 and only when movement speeds were below 5

cm/sec within the corresponding PSD windows. The incidence, strength, and duration of SWDs were then

compared between Pb^{2+}-exposed rats and controls.

PPI task. Pre-pulse inhibition of acoustic startle (PPI) was assessed in male and female rats at

PN28, PN50, and PN120 following a protocol adapted from Abazyan et al. (2014). Four identical SR-LAB

startle chambers (San Diego Instruments Inc., San Diego, CA, USA) were used to measure startle

responses. Rats were placed inside the startle chamber into a mounted Plexiglas® cylinder with an

accelerometer. Broadband background noise and acoustic stimuli were delivered via a speaker on the

chamber's ceiling. Sound levels were calibrated using a digital sound level meter, and accelerometer

sensitivities for each chamber were calibrated daily. The SR-LAB software and interface system controlled

the presentation of acoustic stimuli, and accelerometer measurements were recorded. Each experimental

session started with a five-minute period during which rats acclimatized to 68 dB of background noise that

was continuous throughout the session. Rats' average startle responses were calculated as the mean of 10

acoustic pulses (120dB, 100ms) presented at random intertrial intervals. Rats were left in the enclosure for

an additional five minutes before beginning PPI trials.

Each rat was exposed to the following trial types: pulse-alone (120 dB, 100 ms); pre-pulse alone

(90 dB, 20 ms); and pre-pulse–pulse combinations using one of five pre-pulse intensities: 74, 78, 82, 86,

and 90 dB. Pulse onsets lagged pre-pulse offsets by 80 ms. Each session consisted of six trials with each

pre-pulse intensity, presented in a pseudorandom order with random inter-trial intervals. Startle responses

were quantified as accelerometer measurements averaged over the 100 ms pulse presentations. PPI was

quantified as the percent change in startle amplitude on pre-pulse–pulse trials relative to pulse-alone trials.

Histology. After data collection, rats used in electrophysiological experiments were anesthetized with

isoflurane (5%) mixed with oxygen (800 ml/min), and marking lesions were made with a NanoZ (Plexon) to

deliver 10μA for 10s at each electrode location. An hour later, rats were transcardially perfused with 100 ml

phosphate-buffered saline (PBS), followed by 200 ml of 4% paraformaldehyde (PFA, pH 7.4; Sigma-

Aldrich, St. Louis, MO). Blood Pb^{2+} levels of these rats were measured in samples obtained from cardiac

puncture during the perfusion. Brains were post-fixed overnight in 4% PFA and then placed in a 30%

sucrose solution for cryoprotection. Frozen brains were cut on a Leica CM3050 S cryostat (40 µm; coronal)

and Nissl-stained. Images were acquired on an Olympus BX41 with a UPlanSApo 4x/0.16 objective under

brightfield illumination. Modest brightness, color, and contrast adjustments were made uniformly in macOS

Sonoma 14.5 using Preview. Lesion locations were determined by multiple raters using the microscope

directly, and then matched to the photomicrographs, which were marked using plates from the Paxinos &

Watson Atlas (2004).

Results

Electrophysiological recording sites. In nine rats, NeuroNexus probes were localized in the dorsal

HC, primarily in the stratum radiatum (rad) and stratum lacunosum (slm) layers of CA1, with some

electrode markings spanning the CA1-CA2-CA3 transition zone, and one electrode was deeper near DG

(SuppFig1A). In four rats, the implanted stainless-steel wire probes had electrode markings in the prelimbic

region (PL) of mPFC and the HC, mostly the slm layer of CA1, with some wires encroaching on CA2/CA3

subregions (SuppFig1B). For these probes, only medial sites in the mPFC were utilized. SupplFig1 shows

the histology for all rats.

Blood Pb^{2+} levels (BLLs). Post-mortem BLLs were more than 12-fold higher in the Pb^{2+} exposed

electrophysiology rats (five rats sampled: 29.7, 30.3, 38.3, 38.4, and 41.3 µg/dL) compared to controls (two

rats sampled: 2.8 and 2.8 µg/dL). These BLLs are at the higher end of the range typically observed in the

1500ppm CELLE paradigm, likely because these rats were much older than normally used at ~11 months.

SuppTab1 shows the BLLs for the PPI rats in all groups (ages PN28, PN50, and PN120). In the PPI Pb^{2+}

rats, BLLs were between 19 and 25 µg/dL, and the controls were all less than 1.9 µg/dL (limit of detection).

Grossly intact modes of HC synchronization and exploratory behavior in Pb^{2+} exposed rats. To

characterize any HC pathophysiology caused by chronic Pb^{2+} exposure, we evaluated the effects of

running and stationary modes on oscillatory synchronization in the HC. We first assessed the gross

spectral content of HC LFPs as rats freely explored a large 'open field' environment (Fig1A-D).

Spectrograms illustrating the time-frequency decomposition of HC LFPs were visually similar between

exposure groups (Fig1E and F). In both Pb^{2+}-exposed rats and controls, there were stronger theta rhythms

during running (green arrows) and elevated delta-band power when rats were stationary (cyan arrows).

This observation suggests that chronic Pb^{2+} exposure does not disrupt HC networks so severely as to

subvert the fundamental modes of oscillatory synchronization that coordinate neuronal ensembles and

cortico-HC circuit dynamics during behavioral episodes (Hyman et al., 2005; Jones & Wilson, 2005; Siapas

et al., 2005; Sirota et al., 2008; van der Meer & Redish, 2011).

Importantly, we did not observe any effects of Pb^{2+} exposure on several continuously sampled

measures of rats' locomotor behaviors in plots of continuous movement speed, spatial orientation (body

and head direction), and rotational velocity (SuppFig2); nor did we find differences in behavioral measures

for distinct bouts of running (number of bouts: $t_{(7)}$ = 0.271, p = 0.979; speed: $t_{(7)}$ = 0.186, p = 0.858; and

duration: $t_{(7)}$ = 0.172, p = 0.868), or during stationary bouts (number of bouts: $t_{(7)}$ = 0.261, p = 0.801; speed:

$t_{(7)}$ = 0.912, p = 0.392; duration: $t_{(7)}$ = 1.806, p = 0.114)(SuppFig2). These observations suggest no gross

changes in open-field behavior following Pb^{2+} exposure, which might have directly affected

electrophysiological activity.

Absence seizures in Pb^{2+}-exposed rats and controls. In addition to quasi-normal delta- and theta-

dominated network modes coupled to the locomotor stops and starts, during quiet immobility, we observed

sporadic instances of maximal theta synchronization accompanied by striking theta harmonics in both Pb^{2+}

(Fig1F) and control rats (Fig1E). LFP waveforms corresponding to these theta-harmonic events were

typical of SWDs and reflective of the neurophysiological manifestations of 'absence seizures'

(Fig1F)(Jafarian et al., 2020). SWD incidence was highly variable across rats and between recording

sessions for individual rats, but all rats exhibited SWD during at least one session, in line with previous

reports (e.g., Taylor et al., 2019; Rodgers et al., 2015; Letts et al., 2014; Pearce et al., 2014).

Theta rhythm hypersynchrony. Power spectra of HC LFPs confirmed that awake delta and theta

rhythmic network modes were superficially intact in Pb^{2+}-exposed rats (Fig2A), although there was

substantial variability across tetrodes. Representative PSDs across all eight tetrodes for a lead and control

rat are illustrated in Fig2A. Next, we ran a more detailed analysis, restricting PSD measures to periods of

running and stationary behaviors. Using this approach, we found that theta synchronization was strikingly

elevated in Pb^{2+}-exposed rats (30 channels) compared to controls (30 channels) during bouts of running (2-

sec windows: $t_{(58)}$ = 2.786, p = 0.007; 1-sec windows: $t_{(58)}$ = 3.177, p = 0.002) and stationary behaviors (2-

sec windows: $t_{(58)}$ = 2.056, p = 0.044; 1-sec windows: $t_{(58)}$ = 2.632, p=0.011) (Fig2B). Moreover, theta power

(dB: $t_{(58)}$ = 2.756, p = 0.008; %: $t_{(58)}$ = 1.711, p = 0.093, a significant trend) and theta-harmonics (dB: $t_{(58)}$ =

3.125, p = 0.003; %: $t_{(58)}$ = 3.106, p = 0.003) were also greater in Pb^{2+}-exposed rats compared to controls

during running, measured relative to stationary behaviors (Fig2C). Altogether, this suggests stronger theta

rhythmogenic mechanisms and/or stronger activation of those mechanisms in HC networks in Pb^{2+}

exposed rats. This also suggests that Pb^{2+} exposure-based augmentation of the difference in theta

synchronization between behavioral states may reflect an increased propensity of HC network activity to

self-organize theta rhythmicity given appropriate behavioral excitation, or it may reflect increased

susceptibility of HC to entrainment by theta rhythmic inputs recruited during running.

Delta power in Pb^{2+}-exposed rats was also elevated during 1-sec stationary bouts ($t_{(58)}$ = 2.047, p =

0.045) (Fig2B, inset), but this was neither significant nor a trend for 2-sec bouts ($t_{(58)}$ = 1.488, p = 0.142)

(Fig2B). Delta power also showed a significant trend of elevated levels in Pb^{2+} rats relative to controls

during running bouts of 1-sec windows, but not 2-sec windows (1-sec: $t_{(58)}$ = 1.798, p = 0.077; 2-sec: $t_{(58)}$ =

0.240, p = 0.812). There were no significant differences in delta between Pb^{2+} and control rats during

running measured relative to stationary behaviors (dB: $t_{(58)}$ = 0.374, p = 0.710; %: $t_{(58)}$ = 0.7850, p =

0.436)(Fig2C). Consistent with previous reports (Sakata et al., 2005), we also observed elevated delta

power at the onset or offset of SWDs (SupplFig3). This activity could reflect some aspect of behavior that

may be related to the onset of or escape from behavioral and cognitive arrest accompanying absence

seizures, or it may reflect a behavior-independent neurophysiological signature that is coupled to

transitions between seizure and non-seizure network states. Interestingly, linear fits to delta power across

individual recording sites from each rat further revealed a consistent distal-proximal gradient of increasing

delta power in control rats that was not evident in Pb^{2+}-exposed rats (SuppFig4).

Oscillatory amplitudes. Whereas delta or theta synchronization typically dominates the power

spectrum, gamma frequency synchronization is not well captured by power estimates taken from PSDs.

Gamma power is diminutive relative to slower frequencies, in part because gamma synchronization can

occur across a broad range of frequencies, while faster dynamics of power and frequency are generally

averaged across longer windows, and the network dynamics underlying gamma oscillations can vary in

scale from markedly localized network motifs to large-scale coherent circuits. Therefore, to evaluate the

impact of Pb^{2+} on gamma synchronization and to further explore delta and theta effects, we derived peak-

to-trough amplitudes of individual oscillatory cycles for each frequency band throughout each recording

session.

Modulation of oscillatory amplitudes by rats' movement speeds. Cycle amplitudes for delta, theta,

and fast gamma were related to movement speeds in control rats and Pb^{2+}-exposed rats (Fig2D). Each of

these relationships comprised: (1) a steep primary regime at slow movement speeds (~0-15 cm/sec)

mainly reflecting averaging across transitions between network modes at behavioral transitions between

immobility and running, and (2) a flatter secondary regime reflecting genuine running speed modulation of

oscillatory amplitudes. To demonstrate the relationships between amplitudes and speeds, we ran

correlations on each channel, counted the number of significant correlations across bands and groups, and

the mean Pearson's r as a measure of correlation strength. Results show that delta was negatively related

to running speed in 21/30 Pb^{2+}-exposed rats (mean r = -.508) and 29/30 controls (mean r = -0.727),

primarily reflecting a quick drop in the amplitude during the initial regime. Theta was positively correlated

with running speed in all 30/30 Pb^{2+}-exposed rats (mean r = 0.544) and 23/30 controls (mean r = 0.739).

Comparatively, slow gamma had very weak and inconsistent (positive and negative) relationships with

running speed, showing significant correlations in 9/30 Pb^{2+}-exposed rats (mean r = -.090) and 14/30

controls (mean r = -0.093). Lastly, fast gamma amplitudes had strong positive relationships with running

speeds, showing 21/30 Pb-exposed rats (mean r = 0.532) and 20/30 controls (mean r = 0.502).

Pb^{2+} amplifies delta, theta, and fast gamma synchronization. Theta cycle amplitudes, compiled

across rats in each group, were strikingly elevated in Pb^{2+}-exposed rats relative to controls during

stationary bouts ($t_{(114,635)}$ = 32.920, p = 1.45×10^{-236}) and running bouts ($t_{(102,653)}$ = 116.521, p < 0.001)

(Fig2E). Fast gamma-frequency oscillations were also markedly amplified during stationary bouts ($t_{(1,276,491)}$

= 168.199, p <0.001) and running ($t_{(1,127,377)}$ = 101.326, p < 0.001)(Fig2E). Interestingly, theta frequency

was substantially reduced in Pb^{2+}-exposed rats during stationary bouts (control: 7.159 ± 1.099 Hz, Pb:

6.837 ± 0.984 Hz; $t_{(842)}$ = 4.489, p = 8.153 x 10^{-6}) and running (control: 8.188 ± 0.439 Hz, Pb: 7.759 ±

0.602 Hz; $t_{(702)}$ = 10.769, p = 3.9 x 10^{-25}), potentially reflecting synchronization of larger neuronal

populations. Differences in delta-frequency cycle amplitudes between Pb^{2+}-exposed rats and controls were

small relative to those for theta and gamma (stationary: $t_{(41,741)}$ = 5.547, p = 2.932 x 10^{-8}; running, $t_{(34,211)}$ =

8.235, p = 1.860 x 10^{-16}). It is also important to consider that with such large numbers of individual cycle

amplitude measurements for each frequency band, statistical significance may not be all that informative.

However, these results generally show that Pb^{2+} increases cycle amplitude in the delta, theta, and fast

gamma bands, with the most substantial differences in the latter two bands (Fig2E).

Pb^{2+} exposure tightens oscillatory fidelity to behavior. Pairwise scatterplots of theta power, theta

frequency, and running speed for each bout type suggested that (1) covariation of theta power and

frequency was disrupted in Pb^{2+}-exposed rats relative to controls, and (2) running speed modulation of

theta frequency was confined to a narrower range of speeds; whereas (3) running speed modulation of

theta power occupied similar parameter spaces in Pb^{2+}-exposed rats and controls (SuppFig5).

Comparisons of distributions of correlation coefficients confirmed the expected positive correlations of theta

power and frequency to running speed for most recording sites in both Pb^{2+}-exposed rats and controls

(SuppFig6). Importantly, the correlation of theta power to running speed was elevated in Pb^{2+}-exposed rats

relative to controls ($t_{(58)}$ = 2.582; p = 0.012), potentially reflecting the weakening of HC processes underlying

spatial or mnemonic representations in favor of entrainment or tuning to locomotor parameters.

Correlations between running speed and theta frequency were similar in strength between groups ($t_{(58)}$ =

1.466; p = 0.178) despite the apparent restriction of speed modulation of theta to a narrower range of

speeds in Pb^{2-}-exposed rats (SuppFig5).

Behavioral modulation of delta, theta, and gamma oscillations. We have shown that Pb^{2+}-exposed

rats exhibited theta rhythm hypersynchrony and higher amplitude gamma frequency oscillations relative to

controls during stationary and running bouts, as well as increased correlations of oscillation parameters to

locomotor behavior (Fig2). To determine whether theta and gamma hypersynchrony during running were

related to functional differences in speed modulation of oscillatory amplitudes, we evaluated the amplitudes

of delta, theta, and fast gamma oscillations before and after peaks in running speed (>40 cm/sec) that were

separated by at least 10 seconds. Average peak-triggered amplitudes of delta, theta, and fast gamma

oscillations were elevated in Pb^{2+}-exposed rats relative to controls throughout the peak-triggered analysis

window (Fig3A). The non-zero average values of these z-scored amplitudes stem from the elevated

likelihood that an animal is running at times within the analysis window (recent or upcoming running

occurring within +/- 10 sec) relative to more remote times. As predicted by our bout-wise analyses of

oscillation amplitudes, delta cycle amplitudes were at their lowest near the times of running speed peaks.

In contrast, theta and fast gamma amplitudes were highest at or just preceding peaks in running speed

(Fig3A). The opposite pattern occurs with stop-triggered amplitudes (Fig3A).

Pb^{2+} selectively disrupts the behavioral modulation of fast gamma. We asked whether Pb^{2+}

exposure impacted the behavioral modulation of delta, theta, and fast gamma synchronization by

comparing z-normalized amplitudes in each frequency band at short latencies from running speed peaks.

Normalization across the duration of each recording session generated vectors representing the

proportionate change in oscillation amplitude with the behavioral parameter. It also balanced each

channel's contribution to group-wise comparisons despite possible differences in the amplitude or quality of

raw LFP signals. Fig3A shows speed peak-triggered averages of relative amplitudes for delta, theta, and

fast gamma. Normalized delta amplitudes were negative at all peak-latencies for both Pb^{2+}-exposed and

control rats, indicating that delta amplitudes near running speed peaks were smaller than the average

across all behaviors (Fig3B). In contrast, normalized amplitudes of theta and fast gamma oscillations were

positive at all latencies, indicating elevated synchronization during fast running that, for theta, returned to

near average levels immediately after speed peaks (Fig3B), but for fast gamma, returned to average levels

more slowly over several seconds. Using mixed-design ANOVAs, we compared normalized amplitudes of

the delta, theta, and fast gamma oscillations during the five-time bins spanning -2.25 sec and 0.25-sec

latencies between Pb^{2+}-exposed and control rats. We found a main effect of decreasing delta amplitudes

across latencies relative to speed peaks ($F_{(4, 44)}$ = 3.752, p = 0.015), but there was no effect of Pb^{2+}

exposure on delta synchronization relative to controls ($F_{(1, 44)}$ = 0.194, p = 0.673), nor an interaction effect

($F_{(4, 28)}$, = 0.174, p = 0.950). Interestingly, in this analysis, we found no difference across latencies for theta

amplitudes ($F_{(4, 44)}$ = 0.028, p = 0.872), no difference between Pb^{2+}-exposed and control rats ($F_{(1,44)}$ = 0.906,

p = 0.474), and no interaction effect ($F_{(4, 28)}$ = 1.851, p = 0.147). Notably, the ANOVA for fast gamma

revealed significant main effects of Pb^{2+} exposure ($F_{(4, 44)}$ = 7.119, p = 0.0321), latency ($F_{(4, 44)}$ = 6.684, p =

6.589 x 10^{-4}), but no interaction effect ($F_{(4, 28)}$ = 0.485, p = 0.746) indicating that whereas unnormalized fast

gamma oscillations were amplified in Pb^{2+}-exposed rats (Fig3A, right panel), the degree to which fast

gamma amplitude was modulated by running speed was diminished relative to controls (Fig3B, right

panel). Both effects can be seen in Fig3A, noting that although the raw amplitudes were larger in Pb^{2+}-

exposed rats, the range across which gamma amplitude varied as a function of peak latency was

compressed relative to smaller, more broadly modulated amplitudes in control rats.

Pb^{2+} disrupts slow dynamics of fast gamma. Lastly, we independently z-normalized amplitudes for

each frequency band across the 20-second triggering window for each peak in running speed included in

the analysis. This change to the normalization used in the previous analysis removes the assumption that

the range and scale of oscillatory amplitudes are stationary for a given frequency band throughout a

behavioral session. This analysis, more sensitive to temporal dynamics, revealed reliable amplifications of

theta and fast gamma oscillations at ~-4 sec latency from running speed peaks (Fig3C) that could reflect

behavioral or cognitive regularities occurring before a period of running. We asked whether chronic Pb^{2+}

exposure disrupted these slow temporal dynamics of theta or fast gamma amplitudes during running and

found that the elevation of gamma amplitude was selectively eliminated in Pb^{2+}-exposed rats while the

corresponding amplification of theta was preserved (Fig3C). We considered whether amplifying theta and

fast gamma oscillations before running speed peaks might be a spurious observation. To the contrary,

however, we consider its presence in both frequency bands and argue for it being a real effect. We also

argue that unless the effect were real, it would not be expected to show a relationship to running

parameters. However, we found that the early peaks in theta and fast gamma amplitudes predominantly

occurred in trials with fast peak speeds, supporting the possibility of cognitive activity or behavior that

reliably precedes fast-running peaks.

Pb^{2+} exposure exacerbates absence seizures. As noted earlier, both Pb^{2+}-exposed rats and

controls exhibited absence seizures while stationary. These events were characterized by SWDs in the

LFP (Fig4A), accompanied by maximal theta-frequency synchronization and several theta harmonics

appearing in spectrograms (Fig4B). In many sessions, seizures generated a considerable right tail in

distributions of theta power estimates during immobility (Fig4C). In pronounced cases theta power

distributions were bimodal (Fig4C, inset) such that: (1) the lowest theta values corresponded to quiet-

wakeful immobility; (2) intermediate and larger values corresponded to elevated theta during running; and

(3) theta values during SWDs were on the order of the maximal theta during fast running (Fig4D, left

panel). Recordings at sites exhibiting weaker theta during running often enabled easy detection of SWDs

(Fig4D, right panel) using a single threshold for theta power. Including additional terms for thresholding

power at theta harmonic frequencies further enabled the detection of SWDs (Fig4D, right, insets).

We recently characterized delta-dominated HC network activity during intermittent locomotion as a

fundamental mode of encoding or segmenting events within a behavioral episode (Schultheiss et al.,

2020). Seizures were uniformly distinct from these delta states during immobility, exhibiting minimal delta

power (Fig4E). Interestingly, in some of our data, strong delta synchronization was observed to flank

seizures in the spectrogram, occurring at seizure onset or offset (Fig4B, red arrowheads), which could aid

seizure prediction in a clinical setting.

Given the variability in the spectral content of recordings at different HC sites (as in Fig4D, left vs

right), discerning the spectral signature of SWDs could be a greater or lesser challenge. Using a novel

seizure-detection algorithm that integrates the self-normalized (z-scored over time) power at theta

frequency and at the first two harmonics of theta with the concurrent LFP signal amplitude, we were able to

detect high-reliability SWDs occurring during immobility (Fig4F), see Methods for algorithmic details. We

would argue these are likely to have been absence seizures. We found a broad range of values for which

the detected events could be visually confirmed by varying the detection threshold applied to this algorithm.

The range across which the algorithm performed well was bounded below by false positives, where the

algorithm identified other instances of theta harmonic activity, typically during running, in addition to SWDs.

We reduced most false positives by requiring the animal to be stationary during the theta-harmonic event.

The algorithm's functional range was bounded above by false negatives, where the algorithm exclusively

identified SWDs but missed some proportion of the true incidence. Pb^{2+}-exposed rats exhibited *normal*

SWDs with the same theta fundamental frequency as controls ($t_{(42)} = 0.027$, $p = 0.979$). However, Pb^{2+}

exposed rats had significantly more SWDs ($t_{(42)} = 2.428$. $p = 0.020$), trended toward greater strength ($t_{(247)} =$

1.801, $p = 0.073$), had significantly longer durations ($t_{(247)} = 4.294$, $p = 2.518 \times 10^{-05}$), and occurred during a

significantly greater proportion time spent stationary ($t_{(42)} = 2.162$, $p = 0.036$)(Fig4G). To further confirm the

validity of these SWDs as seizure-like events, we used ethosuximide (200 mg/kg) injections part of the way

through a session in some of the rats. We observed that ethosuximide eliminated the appearance of SWDs

at doses that did not alter gross behavior, supporting that these events are absence seizures (Fig4H).

A schizophrenia-like pattern of sensorimotor gating deficits caused by Pb^{2+} exposure. The

electrophysiology experiments predict that disrupted HC network hypersynchrony should disrupt HC-

related cognitive tasks. To examine this possibility, we measured PPI of the acoustic startle response

(ASR), a test of sensorimotor gating that is disrupted by HC damage (Pouzet et al., 1999; Bast et al.,

2001a; 2001b; 2003), and a canonical dysfunction linked to schizophrenia. Chronic Pb^{2+} exposure caused

significant impairments of PPI at all stimulus levels for both late adolescence (PN50) and young adults

(PN120) males (Mixed ANOVA: dB Levels: $F_{(3.12, 109.23)} = 53.49$, $p < 0.001$; Pb^{2+} vs. controls: $F_{(1,35)} = 11.32$,

$p = 0.002$; Interactions: $F_{(3.12, 109.23)} = 2.56$, $p = 0.069$) and PN120 (Mixed ANOVA: dB Levels: $F_{(2.88, 115.11)} =$

59.59, $p < 0.001$; Pb^{2+} vs. controls: $F_{(1,40)} = 9.30$, $p = 0.004$, Interaction: $F_{(2.88, 115.11)} = 2.00$, $p = 0.12$)(Fig5),

but neither juveniles at PN28 (Mixed ANOVA: dB Levels: $F_{(2.57, 89.83)} = 68.69$, $p < 0.001$; Pb^{2+} vs. controls:

$F_{(1,35)} = 0.54$, $p = 0.47$; Interaction: $F_{(2.57, 89.83)} = 2.56$, $p = 0.069$) nor females at any age exhibited any

deficits (Pb vs. Control p's > 0.10; Fig5)(SuppFig7). This pattern recapitulates the sex dependence of PPI

deficits exhibited in male and female SZ subjects (Mena et al., 2016; Takahashi et al., 2008; Ochoa et al.,

2012), supporting the view that our rodent model for studying the Pb^{2+} induced HC pathophysiology may

also be relevant for understanding aberrant HC network dynamics in SZ.

620   **Discussion**

Summary of Main Findings. Childhood Pb^{2+} intoxication alters the neurodevelopmental trajectory of the

brain, including the HC and PFC, and increases the risk of neurodevelopmental disorders, such as

schizophrenia (see Albores-Garcia et al., 2021) and epilepsy (Sasmaz et al., 2003). Here, we asked how

Pb^{2+} exposure impacts the HC at the network level using a rodent model of chronic early-life Pb^{2+} exposure

(CELLE) that presents with molecular and cellular correspondences to schizophrenia (Stansfield et al.,

Transl Psych 5: e522, 2015)**.** The prediction, based on established neurobiological deficits resultant of Pb^{2+}

exposure, including hypofunction of the NMDA receptor (Guilarte, 1997), reduced LTP (Nihei et al., 2000),

and reduced entorhinal cortex inputs (Zhou et al., 2023), is that chronic Pb^{2+} exposure should diminish HC

activity. This outcome would help explain Pb^{2+} exposure-induced cognitive deficits observed in CELLE rats

(McGlothan et al., 2008; Kuhlmann et al., 1997; Jaako-Movits et al., 2005) and potentially the cognitive and

intellectual deficits seen in children. Simply put, reduced HC activity would reduce HC function.

Alternatively, the loss of PVGI interneurons (Stansfield et al., 2015) and impaired LTD (Zhao et al., 1999)

predict network overexcitation in the HC. Instead of either prediction, we found that Pb^{2+} exposure caused

network *hypersynchrony,* which appears as changes in the coordinated rhythms that enable efficient

network processing. This suggests that Pb^{2+} exposure causes a rhythmic architectural shift in the HC

network, reshaping activity patterns in a way that diminishes function. Specifically, we found that Pb^{2+}

exposure resulted in four main HC network changes, including: (1) HC theta hypersynchrony, (2) gamma

oscillation amplification, (3) disruption of behavioral-related theta and gamma network modulations, and (4)

a striking exacerbation of absence seizures characterized by maximal theta synchrony during SWDs. We

also showed that HC SWDs corresponded to SWDs occurring in cortical mPFC sites. Lastly, we predicted

that aberrant HC hypersynchrony and SWDs caused by CELLE would impair sensorimotor gating because

it is disrupted by aberrant HC activity (Zhang et al., 2002). Thus, we tested Pb^{2+}-exposed rats on the PPI of

the acoustic startle reflex task. We found that PPI was disrupted in late adolescence and young adult male

rats but not female or juvenile males—a pattern that recapitulates PPI deficits in schizophrenia (Kumari et

al., 2004). Theoretically, these results suggest that HC hypersynchrony and SWDs cause behavioral

deficits by reducing the flexibility and processing efficiency of the HC network and causing behavioral

arrests (a.k.a., the blank stare) via intermittent absence seizures (Holmes et al., 1987; Taylor et al., 2019).

That is, hypersynchronous HC networks, the extreme of which is seizure-related activity, could be causing

some of the Pb^{2+} exposure-related PPI and cognitive deficits in CELLE rats and possibly in Pb^{2+} exposed

children (Kponee-Shovein et al., 2020).

How can hypersynchrony cause network dysfunction? Oscillations coordinate HC neuronal ensembles,

creating time windows for synaptic integration and encoding dimensions of ongoing, remembered, and

imagined future experiences, in turn guiding spatial navigation and decision-making processes (Buzsaki &

Watson, 2012; Buzsaki & Wang, 2012). In addition, HC encoding of the present location and possible

future trajectories comprise sequential neuronal activations that reiterate and update gamma-linked spatial

representations, which occur on every cycle of the theta rhythm (Lowet et al., 2023; Buzsaki & Wang,

2012). Whereas gamma-theta synchrony usually enables cognitive representations, any change (in any

direction) in the spatiotemporal structure of these neural representations and rhythmic mechanisms would

diminish neural computations and mnemonic processes. Our finding of theta-gamma hypersynchrony in

Pb^{2+} exposed rats argues that 'overly' structured neural activity may diminish HC function, presumably by

reducing the variability and flexibility of spiking activity patterns to a narrower range of less dynamic

network states. For example, information processing is lost if all neurons are synchronous (or

asynchronous). This suggests there must be an optimal point of rhythmic coordination to maximize

processing, which is assumed to be benchmarked by the control rats. In support of this notion, several

neuropsychiatric disorders are associated with hypersynchrony, including epilepsy (Katuri et al., 2023),

schizophrenia (Delgado-Sallant et al., 2021), Alzheimer's disease (Kazim et al., 2021), among others.

Theta and SWDs. HC SWDs reflect entrainment to a corticothalamic circuit oscillation initiated in the

cortex and amplified by intrathalamic circuits. The robust theta-band fundamental frequency of SWDs

suggests the involvement of normal theta rhythmogenic mechanisms, including but not limited to

interneuron generators and modulators (Amilhon et al., 2015), medial septal GABAergic input (Borhegyi et

al., 2004), and Huygens synchronization (Kocsis et al., 2021). We also found that theta frequency was

lower in Pb^{2+}-exposed rats during running and immobility. Slower frequency oscillations are generally

associated with synchronization of larger populations of neurons. Lower frequency theta could, therefore,

reflect the broader synchronization of HC networks, entraining a higher proportion of neurons and

restricting HC functional capacity by reducing the number and flexibility of network states available. Here,

the reduction in GABAergic diversity (Stansfield et al., 2015) (loss of PVGI cells) in chronic Pb^{2+} exposure,

specifically within the context of homeostatic regulation of the E/I balance, suggests that the theta-gamma

hypersynchrony observed may be due to an overrepresentation of the contribution of one inhibitory

interneuronal subtype to the network dynamics recruiting larger populations of neurons. It is provocative to

consider that the theta-band fundamental frequency of SWDs may reflect the pathological engagement of

theta rhythmogenic mechanisms that are normally the foundation of spatial encoding and episodic memory

systems. We argue that disrupted theta rhythmic coordination of HC networks is a key mechanism of

chronic Pb^{2+} exposure-related cognitive deficits.

Possible translational avenues to explore. Given our results, electrophysiological measurements

(e.g., EEG, MEG, deep electrodes) should be helpful in distinguishing endophenotypes in Pb^{2+} exposed

populations, as hypersynchrony and SWDs should be reflected, in some way, in these approaches (e.g.,

Kmiecik-Malecka et al., 2009). This strategy enables mappings of clinical features onto underlying disease

mechanisms based on the generalization that similar clinical presentations are likely to share similar

underlying pathophysiology. When distinct clusters among features can be identified, it serves the purpose

of identifying a mechanism of particular importance, because the manifestation of the disease depends on

it, or divides the disease into subtypes. The significance of this for distinguishing endophenotypes is that

the development of effective treatments of the complex cognitive effects of Pb^{2+} exposure may also exhibit

efficacy for cases of other diseases that share a hypersynchronous profile stemming from disruption of

GABAergic dynamics.

Theta hypersynchrony and SWDs suggest possible interventions for hypersynchronous neural activity

resulting from Pb^{2+} exposure, including (1) deep brain stimulation in a network hub to disrupt

hypersynchrony and circuit entrainment to rhythmic activity (Zangiabadi et al., 2019), (2) pharmacological

interventions to reduce excitability and rhythmogenesis (e.g., anticonvulsants), or (3) using the promising

BDNF-mimetic 7,8-dihydroxyflavone (7,8-DHF) which mitigates the negative impact of Pb^{2+-}exposure on

presynaptic terminals (Zhang et al., 2018). This suggests that 7,8-DHF might modify network dynamics with

a safe, inexpensive substance found in many fruits in the human diet (Guilarte, 2023). All these approaches

require further research on the efficacy and optimal timing of each intervention to reduce hyperexcitability

and enhance cognitive function in cases of chronic Pb^{2+} exposure, as well as their long-term impacts on

brain function and cognitive/behavioral health.

Conclusion. Chronic Pb^{2+} exposure during early brain development leads to significant alterations in HC

network activity, including theta-gamma hypersynchrony and absence seizures. Theoretically, aberrant

synchrony disrupts informational processing in the HC, resulting in diminished cognitive functions and

impaired sensorimotor gating. Interestingly, these deficits may manifest more subtly in children as blank

stares, distractibility, and reduced intellectual abilities in the classroom setting. If left untreated, such issues

may lead to disruptive behaviors in the short term and social delinquency in the long term.

References

1. Abazyan, B., Dziedzic, J., Hua, K., Abazyan, S., Yang, C., Mori, S., Pletnikov, M. V., & Guilarte, T. R. (2014). Chronic exposure of mutant DISC1 mice to lead produces sex-dependent abnormalities consistent with schizophrenia and related mental disorders: a gene-environment interaction study. Schizophrenia bulletin, 40(3), 575–584. https://doi.org/10.1093/schbul/sbt071

2. Amilhon, B., Huh, C. Y., Manseau, F., Ducharme, G., Nichol, H., Adamantidis, A., & Williams, S. (2015). Parvalbumin Interneurons of HC Tune Population Activity at Theta Frequency. Neuron, 86(5), 1277–1289.

3. Albores-Garcia, D., McGlothan, J. L., & Guilarte, T. R. (2021). Early-life lead exposure and neurodevelopmental disorders. *Current opinion in toxicology*, *26*, 22–27.

4. Arrieta, O., Palencia, G., García-Arenas, G., Morales-Espinosa, D., Hernández-Pedro, N., & Sotelo, J. (2005). Prolonged exposure to lead lowers the threshold of pentylenetetrazole-induced seizures in rats. *Epilepsia*, *46*(10), 1599–1602.

5. Barltrop, D. (1969). Transfer of lead to the human foetus. In: Mineral metabolism in pediatrics (Barltrop D, Burland W, eds). Oxford: Blackwell Scientific Publications.135-151.

6. Bast, T., Zhang, W. N., & Feldon, J. (2001a). Hyperactivity, decreased startle reactivity, and disrupted prepulse inhibition following disinhibition of the rat ventral HC by the GABA(A) receptor antagonist picrotoxin. Psychopharmacology, 156(2-3), 225–233.

7. Bast, T., Zhang, W. N., Heidbreder, C., & Feldon, J. (2001b). Hyperactivity and disruption of prepulse inhibition induced by N-methyl-D-aspartate stimulation of the ventral HC and the effects of pretreatment with haloperidol and clozapine. Neuroscience, 103(2), 325–335.

8. Bast, T., & Feldon, J. (2003). Hippocampal modulation of sensorimotor processes. Progress in neurobiology, 70(4), 319–345.

9. Bernard, S. M., & McGeehin, M. A. (2003). Prevalence of blood lead levels >or= 5 micro g/dL among US children 1 to 5 years of age and socioeconomic and demographic factors associated with blood of lead levels 5 to 10 micro g/dL, Third National Health and Nutrition Examination Survey, 1988-1994. *Pediatrics*, *112*(6 Pt 1), 1308–1313.

10. Borhegyi, Z., Sebestyén Varga, V., Szilágyi, N., Fabó, D., & Freund, T. F. (2004). Phase Segregation of Medial Septal GABAergic Neurons during Hippocampal Theta Activity. The Journal of Neuroscience, 24(39), 8470–8479.

11. Buzsáki G & Wang XJ (2012). Mechanisms of gamma oscillations. Annu Rev Neurosci. 2012;35:203-25.

12. Buzsáki G, Watson BO. (2012) Brain rhythms and neural syntax: implications for efficient coding of cognitive content and neuropsychiatric disease. Dialogues Clin Neurosci., 14(4):345-67.

13. Caito, S., & Aschner, M. (2017). Developmental Neurotoxicity of Lead. Advances in neurobiology, 18, 3–12.

14. Canfield, R. L., Henderson, C. R., Jr, Cory-Slechta, D. A., Cox, C., Jusko, T. A., & Lanphear, B. P. (2003). Intellectual impairment in children with blood lead concentrations below 10 microg per deciliter. The New England journal of medicine, 348(16), 1517–1526.

15. CDC (2021). 30th anniversary of the CDCs Childhood Lead Poisoning Prevention Program (CLPPP) https://www.cdc.gov/nceh/lead/about/30th-anniversary.html (Accessed on June 4, 2021)

16. Delgado-Sallent C, Nebot P, Gener T, Fath AB, Timplalexi M, Puig MV. Atypical, but Not Typical, Antipsychotic Drugs Reduce Hypersynchronized Prefrontal-Hippocampal Circuits during Psychosis-Like States in Mice: Contribution of 5-HT2A and 5-HT1A Receptors. Cereb Cortex. 2022 Aug 3;32(16):3472-3487.

17. Dietrich KN, Ware JH, et al. (2004) Effect of chelation therapy on the neuropsychological and behavioral development of lead-exposed children after school entry. Pediatrics, 114,19-26.

18. Dignam T, Kaufmann RB, LeStourgeon L, Brown MJ (2019). Control of lead sources in the United States, 1970-2017: Public health progress and current challenges to eliminating lead exposure. J Public Hlth Manag Pract 25: S13-S22.

19. Egan KB, Cornwell CR, Courtney JG, Ettinger AS. (2021) Blood Lead Levels in U.S. Children Ages 1-11 Years, 1976-2016. Environ Health Perspect.129(3):37003.

20. Gavazzo, P., Zanardi, I., Baranowska-Bosiacka, I., & Marchetti, C. (2008). Molecular determinants of Pb2+ interaction with NMDA receptor channels. *Neurochemistry international*, *52*(1-2), 329–337.

21. Govoni, S., Battaini, F., Fernicola, C., Coniglio, L., & Trabucchi, M. (1988). Central nervous system effects of lead: a study model in neuro.

22. Guilarte, T. R., Opler, M., & Pletnikov, M. (2012). Is lead exposure in early life an environmental risk factor for Schizophrenia? Neurobiological connections and testable hypotheses. Neurotoxicology, 33(3), 560–574.

23. Guilarte T. R. (1997). Pb2+ inhibits NMDA receptor function at high and low affinity sites: developmental and regional brain expression. Neurotoxicology, 18(1), 43–51.

24. Guilarte T. R. (2023). Invited Perspective: Challenging the Dogma of Lead Neurotoxicity. Environ Health Perspect, 31(8):81306.

25. Guilarte, T. R., & Miceli, R. C. (1992). Age-dependent effects of lead on [3H]MK-801 binding to the NMDA
receptor-gated ionophore: in vitro and in vivo studies. Neuroscience letters, 148(1-2), 27–30.
26. Guilarte, T. R., & McGlothan, J. L. (1998). Hippocampal NMDA receptor mRNA undergoes subunit specific
changes during developmental lead exposure. Brain research, 790(1-2), 98–107.
27. Hanna-Attisha, M., LaChance, J., Sadler, R. C., & Schnepp, A. C. (2016). Elevated blood lead levels in
children associated with the Flint drinking water crisis: A spatial analysis of risk and public health response.
American Journal of Public Health. 106. 283-290.
28. Hashemzadeh-Gargari H., Guilarte T. R. (1999). Divalent cations modulate N-Methyl-D-Aspartate Receptor
function at the glycine site. Journal Pharmacology & Experimental Therapeutics 290: 1356-1362.
29. Hashimoto, T., Volk, D. W., Eggan, S. M., Mirnics, K., Pierri, J. N., Sun, Z., Sampson, A. R., & Lewis, D. A.
(2003). Gene expression deficits in a subclass of GABA neurons in the prefrontal cortex of subjects with
schizophrenia. Journal of Neuroscience. 23(15), 6315–6326.
30. Holmes, G.L., McKeever, M. and Adamson, M. (1987), Absence seizures in children: Clinical and
electroencephalographic features. Ann Neurol., 21: 268-273.
31. Hyman, J. M., Zilli, E. A., Paley, A. M., & Hasselmo, M. E. (2005). Medial prefrontal cortex cells show dynamic
modulation with the hippocampal theta rhythm dependent on behavior. *Hippocampus*, *15*(6), 739–749.
32. Jaako-Movits, K., Zharkovsky, T., Romantchik, O., Jurgenson, M., Merisalu, E., Heidmets, L. T., &
Zharkovsky, A. (2005). Developmental lead exposure impairs contextual fear conditioning and reduces adult
hippocampal neurogenesis in the rat brain. Internat. J. of Dev. Neurosci., 23(7), 627–635.
33. Jafarian M, Esmaeil Alipour M, Karimzadeh F. Experimental Models of Absence Epilepsy. Basic Clin
Neurosci. 2020 Nov-Dec;11(6):715-726. doi: 10.32598/bcn.11.6.731.1. Epub 2020 Nov 1.
34. Jett, D. A., Kuhlmann, A. C., & Guilarte, T. R. (1997). Intrahippocampal administration of lead (Pb) impairs
performance of rats in the Morris water maze. Pharmacology, biochemistry, and behavior, 57(1-2), 263–269.
35. Jones, M. W., & Wilson, M. A. (2005). Phase precession of medial prefrontal cortical activity relative to the
hippocampal theta rhythm. *Hippocampus, 15*(7), 867–873.
36. Katuri, G., Rosa Jr, E., & Follmann, R. (2023). Phase Synchronization in Brain Collective Dynamics. In
Proceedings of the 14th ACM International Conference on Bioinformatics, Computational Biology, and Health
Informatics (BCB '23). Association for Computing Machinery, New York, NY, USA, Article 44, 1–7.
37. Kazim, S. F., Seo, J. H., Bianchi, R., Larson, C. S., Sharma, A., Wong, R. K. S., Gorbachev, K., & Pereira, A.
C. (2021). Neuronal Network Excitability in Alzheimer's Disease: The Puzzle of Similar versus Divergent
Roles of Amyloid β and Tau. 8(2).
38. Kmiecik-Małecka, E., Małecki, A., Pawlas, N., Woźniakova, Y., & Pawlas, K. (2009). The effect of blood lead
concentration on EEG, brain electrical activity mapping and psychological test results in children. Polish
Journal of Environmental Studies, 18(6), 1021–1027.
39. Kocsis B, Martínez-Bellver S, Fiáth R, Domonkos A, Sviatkó K, Schlingloff D, Barthó P, Freund TF, Ulbert I,
Káli S, Varga V, Hangya B. Huygens synchronization of medial septal pacemaker neurons generates
hippocampal theta oscillation. Cell Rep. 2022 Aug 2;40(5):111149.
40. Kponee-Shovein, K. Z., Weisskopf, M. G., Grashow, R., Rotem, R. S., Coull, B. A., Schnaas, L., Hernández-
Chávez, M. del C., Sánchez, B. N., Peterson, K. E., Hu, H., & Téllez-Rojo, M. M. (2020). Estimating the
causal effect of prenatal lead exposure on prepulse inhibition deficits in children and
adolescents. Neurotoxicology, 78, 116–126.
41. Kuhlmann, A. C., McGlothan, J. L., & Guilarte, T. R. (1997). Developmental lead exposure causes spatial
learning deficits in adult rats. Neuroscience letters, 233(2-3), 101–104.
42. Kumari V., Aasen I., Sharma T. (2004). Sex differences in prepulse inhibition deficits in chronic schizophrenia.
Schizophr Res., 69(2-3):219-35.
43. Lanphear BP, Hornung R, Khoury J, Yolton K, Baghurst P, Bellinger DC, Canfield RL, Dietrich KN,
Bornschein R, Greene T, Rothenberg SJ, Needleman HL, Schnaas L, Wasserman G, Graziano J, Roberts R.
(2005). Low-level environmental lead exposure and children's intellectual function: an international pooled
analysis. Environ Health Perspect. 113(7):894-9.
44. Lanphear BP, Hornung R, Khoury J, Yolton K, Dietrich KN. (2006). Lead and IQ in Children: Lanphear et al.
Respond. Environ Health Perspect. 114(2):A86–7.
45. Letts, V.A., Beyer, B.J. and Frankel, W.N. (2014), Hidden in plain sight: spike-wave discharges in mouse
inbred strains. Genes, Brain and Behavior, 13: 519-526.
46. Lowet, E., Sheehan, D. J., Chialva, U., De Oliveira Pena, R., Mount, R. A., Xiao, S., Zhou, S., Tseng, H.,
Gritton, H. J., Shroff, S. N., Kondabolu, K., Cheung, C., Piatkevich, K. D., Boyden, E. S., Merz, J., Hasselmo,
M. E., Rotstein, H. G., & Han, X. (2023). Theta and gamma rhythmic coding through two spike output modes
in the HC during spatial navigation. Cell Reports, 42(8), 112906.

47. McGlothan, J. L., Karcz-Kubicha, M., & Guilarte, T. R. (2008). Developmental lead exposure impairs
extinction of conditioned fear in young adult rats. Neurotoxicology, 29(6), 1127–1130.
48. Mena A, Ruiz-Salas JC, Puentes A, Dorado I, Ruiz-Veguilla M, De la Casa LG. (2016). Reduced Prepulse
Inhibition as a Biomarker of Schizophrenia. Front Behav Neurosci. 10:202.
49. Miranda, M. L., Kim, D., Galeano, M. A., Paul, C. J., Hull, A. P., & Morgan, S. P. (2007). The relationship
between early childhood blood lead levels and performance on end-of-grade tests. Environmental health
perspectives, 115(8), 1242–1247.
50. Modabbernia A, Velthorst E, Gennings C, De Haan L, Austin C, Sutterland A, Mollon J, Frangou S, Wright R,
Arora M, Reichenberg A. (2016). Early-life metal exposure and schizophrenia: A proof-of-concept study using
novel tooth-matrix biomarkers. Eur Psychiatry, 36:1-6.
51. Munoz, C., Garbe, K., Lilienthal, H., & Winneke, G. (1988). Significance of hippocampal dysfunction in low
level lead exposure of rats. Neurotoxicology and teratology, 10(3), 245–253.
52. Neal, A. P., Stansfield, K. H., Worley, P. F., Thompson, R. E., Guilarte TR (2010). Lead exposure during
synaptogenesis alters vesicular proteins and impairs vesicular release: potential role of NMDA receptor-
dependent BDNF signaling. Toxicological Sciences 116: 249-263.
53. Neal A. P. & Guilarte T. R. (2010). Molecular Neurobiology of lead: Effects on synaptic function. Molecular
Neurobiology 42: 151-160.
54. Needleman, H. L., McFarland, C., Ness, R. B., Fienberg, S. E., & Tobin, M. J. (2002). Bone lead levels in
adjudicated delinquents. A case control study. Neurotoxicology and teratology, 24(6), 711–717.
55. Needleman H. (2004). Lead poisoning. Annual review of medicine, 55, 209–222.
56. Nihei, M. K., Desmond, N. L., McGlothan, J. L., Kuhlmann, A. C., & Guilarte, T. R. (2000). N-methyl-D-
aspartate receptor subunit changes are associated with lead-induced deficits of long-term potentiation and
spatial learning. Neuroscience, 99(2), 233–242.
57. Ochoa S, Usall J, Cobo J, Labad X, Kulkarni J (2012). Gender differences in schizophrenia and first- episode
psychosis: a comprehensive literature review. Schizophrenia Research Treatment 2012: 916198.
58. Opler, M. G., Brown, A. S., Graziano, J., Desai, M., Zheng, W., Schaefer, C., Factor-Litvak, P., & Susser, E.
S. (2004). Prenatal lead exposure, delta-aminolevulinic acid, and schizophrenia. Environmental health
perspectives, 112(5), 548–552.
59. Opler, M. G., Buka, S. L., Groeger, J., McKeague, I., Wei, C., Factor-Litvak, P., Bresnahan, M., Graziano, J.,
Goldstein, J. M., Seidman, L. J., Brown, A. S., & Susser, E. S. (2008). Prenatal exposure to lead, delta-
aminolevulinic acid, and schizophrenia: further evidence. Environmental health perspectives, 116(11), 1586–
1590.
60. Otto, D. A., Benignus, V. A., Muller, K. E., & Barton, C. N. (1981). Effects of age and body lead burden on
CNS function in young children. I. slow cortical potentials. *Electroencephalography and clinical
neurophysiology*, *52*(4), 229–239.
61. Paxinos, G., and Watson, C. (2004). The rat brain in stereotaxic coordinates 5th Edition. (Massachusetts:
Academic Press).
62. Pearce, P. S., Friedman, D., LaFrancois, J. J., Iyengar, S. S., Fenton, A. A., MacLusky, N. J., Scharfman, H.
E. (2014). Spike–wave discharges in adult Sprague–Dawley rats and their implications for animal models of
temporal lobe epilepsy. Epilepsy & Behavior, 32, 121-131.
63. Poblano, A., Rothenberg, S. J., Schnaas, L., Elías, Y., & Cruz, M. L. (2001). Spatial distribution of EEG theta
activity as a function of lifetime lead exposure in 9-year-old children. *Neurotoxicology*, *22*(4), 439–446.
64. Rodgers, K. M., Dudek, F. E., & Barth, D. S. (2015). Progressive, seizure-like, spike-wave discharges are
common in both injured and uninjured sprague-dawley rats: implications for the fluid percussion injury model
of post-traumatic epilepsy J. Neurosci. 35 (24) 9194-9204.
65. Rogan WJ, Dietrich KN, et al. (2001) The effect of chelation therapy with succimer on neuropsychological
development in children exposed to lead. New England Journal Medicine 344: 1421-1426.
66. Rossouw, J., Offermeier, J., Van Rooyen, J. M. (1987). Apparent central neurotransmitter receptor changes
induced by low-level lead exposure during different developmental phases in the rat. Toxicol Appl Pharmacol.
91, 132-139.
67. Sakata, K., Woo, N. H., Martinowich, K., Greene, J. S., Schloesser, R. J., Shen, L., Lu, B. (2009) Critical role
of promoter IV-driven BDNF transcription in GABAergic transmission and synaptic plasticity in the prefrontal
cortex. PNAS 106: 5942-5947.
68. Sample, J.A. Childhood lead poisoning: Clinical manifestations and diagnosis.
https://www.uptodate.com/contents/childhood-lead-poisoning-clinical-manifestations-and-diagnosis.
(Accessed June 4, 2021)
69. Sasmaz, S., Uz, E., Pinar, T., Vural, H., Eiri, M., Ilihan, A., & Akyol, mer. (2003). Hair lead and cadmium
concentrations in patients with epilepsy and migraine. Neuroscience Research Communications, 32(2), 107–
114.

70. Schultheiss, N.W., Schlecht, M., Jayachandran, M., Brooks, D.R., McGlothan, J.L., Guilarte, T.R. & Allen, T.A.
(2020). Awake delta and theta-rhythmic hippocampal network modes during intermittent locomotion behaviors
in the rat. Behavioral Neuroscience.
71. Sengupta, S., Talidou, A., Lefebvre, J., & Skinner, F. K. (2024). Cell-type specific contributions to theta-
gamma coupled rhythms in the HC. bioRxiv.
72. Siapas, A. G., Lubenov, E. V., & Wilson, M. A. (2005). Prefrontal phase locking to hippocampal theta
oscillations. *Neuron*, *46*(1), 141–151.
73. Sirota, A., Montgomery, S., Fujisawa, S., Isomura, Y., Zugaro, M., & Buzsáki, G. (2008). Entrainment of
neocortical neurons and gamma oscillations by the hippocampal theta rhythm. *Neuron*, *60*(4), 683–697.
74. Stansfield, K. H., Pilsner, J. R., Lu, Q., Wright, R., Guilarte, T. R. (2012). Dysregulation of BDNF-TrkB
signaling in developing hippocampal neurons by lead: Implications for an environmental basis of
neurodevelopmental disorders. Toxicological Sciences.127: 277-295.
75. Stansfield, K. H., Ruby, K. N., Soares, B. D., McGlothan, J. L., Liu, X., & Guilarte, T. R. (2015). Early-life lead
exposure recapitulates the selective loss of parvalbumin-positive GABAergic interneurons and subcortical
dopamine system hyperactivity present in schizophrenia. Translational psychiatry, 5(3), e522.
76. Takahashi H, Iwase M, Ishii R, Ohi K, Fukumoto M, Azechi M, Ikezawa K, Kurimoto R, Canuet L, Nakahachi
T, like N, Tagami S, Morihara T, Okochi M, Tanaka T, Kazui H, Yoshida T, Tanimukai H, Yasuda Y, Kudo T,
Hashimoto R, Takeda M (2008). Impaired prepulse inhibition and habituation of acoustic startle response in
Japanese patients with schizophrenia. Neuroscience Research 62: 187- 194.
77. Tamura, M., Spellman, T.J., Rosen, A.M. et al. (2017). Hippocampal-prefrontal theta-gamma coupling during
performance of a spatial working memory task. Nat Commun 8, 2182.
78. Taylor, J. A., Reuter. J. D., Kubiak, R. A., Mufford, T. T., Booth, C. J., Dudek, F. E., & Barth, D. S. (2019).
Spontaneous recurrent absence seizure-like events in wild-caught rats. J. Neurosci., 39(24), 4829-4841.
79. Toscano, C. D., Hashemzadeh-Gargari, H., McGlothan, J. L., & Guilarte, T. R. (2002). Developmental Pb2+
exposure alters NMDAR subtypes and reduces CREB phosphorylation in the rat brain. *Brain research.*
Developmental brain research, *139*(2), 217–226.
80. Toscano, C. D., & Guilarte, T. R. (2005). Lead neurotoxicity: from exposure to molecular effects. *Brain*
research. Brain research reviews, *49*(3), 529–554.
81. Toscano, C. D., O'Callaghan, J. P., & Guilarte, T. R. (2005). Calcium/calmodulin-dependent protein kinase II
activity and expression are altered in the hippocampus of Pb2+-exposed rats. *Brain research*, *1044*(1), 51–
58.
82. UNICEF (2020), "The toxic truth: Children's exposure to lead pollution undermines a generation of future
potential", 2nd Edition.
83. Wakefield, J. The lead effect? (2002). Environ. Health Perspect. 110, A574– A580.
84. Wang, T., Guan, R. L., Liu, M. C., Shen, X. F., Chen, J. Y., Zhao, M. G., & Luo, W. J. (2016). Lead Exposure
Impairs HC Related Learning and Memory by Altering Synaptic Plasticity and Morphology During Juvenile
Period. Molecular neurobiology, 53(6), 3740–3752.
85. WHO (2010). https://www.who.int/ceh/publications/leadguidance.pdf
86. Yeter D, Banks EC, Aschner M (2020) Disparity in risk factor severity for early childhood blood lead among
predominantly African-American black children: the 1999 to 2010 US NHANES. Int J Env Res Public Health
17: 1552.
87. van der Meer, M. A., & Redish, A. D. (2011). Theta phase precession in rat ventral striatum links place and
reward information. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, *31*(8),
2843–2854.
88. van Luijtelaar G, Onat FY, Gallagher MJ. (2014). Animal models of absence epilepsies: what do they model
and do sex and sex hormones matter? Neurobiol Dis., 72 Pt B:167-79.
89. Zangiabadi N, Ladino LD, Sina F, Orozco-Hernández JP, Carter A, Téllez-Zenteno JF. Deep Brain
Stimulation and Drug-Resistant Epilepsy: A Review of the Literature. Front Neurol. 2019 Jun 6;10:601.
90. Zhang, W.-N., Bast, T., & Feldon, J. (2002). Prepulse inhibition in rats with temporary inhibition/inactivation of
ventral or dorsal HC. Pharmacology, Biochemistry and Behavior, 73(4), 929–940.
91. Zhang XL, McGlothan JL, Miry O, Stansfield KH, Loth MK, Stanton PK, Guilarte TR. (2018). From the Cover:
7,8-Dihydroxyflavone Rescues Lead-Induced Impairment of Vesicular Release: A Novel Therapeutic
Approach for Lead Intoxicated Children. Toxicol Sci.,161(1):186-195.
92. Zhao WF, Ruan DY, Xu YZ, Chen JT, Wang M, Ge SY. (1999). The effects of chronic lead exposure on long-
term depression in area CA1 and dentate gyrus of rat HC in vitro. Brain Res., 818(1):153-9.
93. Zou, R., Gu, X., Huang, C., Wang, H.-L., & Chen, X.-T. (2023). Chronic Pb exposure impairs learning and
memory abilities by inhibiting excitatory projection neuro-circuit of the HC in mice. Toxicology, 153717.
943

944
945

946
Figure 1. During exploratory navigation, HC LFPs display grossly similar spectral content in
***Pb^{2+}-exposed rats as controls*. A.** Representative spatial trajectory during free exploration of the square
(120 cm) 'open field' environment. Black lines represent the rat's body direction (BD as in **B**), and green
and blue dots represent head-mounted LEDs. **B**. Continuous measures of locomotor behavior, including
head direction relative to the environment (HD) and head angle (HA, derived from HD and BD). **C**.
Recording sites of 8 tetrodes located at two depths on each of four shanks of a silicon probe. Tracks left by
the shanks of the implanted probe are visible as disruptions of the CA1 cell layer. **D.** Representative LFP
traces (top) during running and immobility (bottom) showed corresponding periods of strong theta and
delta-band activity (e.g., as highlighted in cyan and green). **E & F**. Representative spectrograms for control
rats (**E**) and Pb^{2+}-exposed rats (**F, top left panel**) showed qualitatively similar patterns of speed modulation
(overlaid white traces) of theta (green arrows) and delta (cyan arrows). **F, top right panel**. Putative
absence seizures occurred in all Pb^{2+}-exposed rats. Seizure incidence was highly variable across rats and
sessions for each rat, i.e., the spectrograms shown in **F** illustrate sessions during which a single seizure
lasting 7 seconds and ~25 minutes of continuous seizing occurred (**top** and **bottom panels, respectively)**.
The red arrowhead in the **top left** panel indicates the seizure magnified in the **top right**. Green arrows
(**bottom panel**) indicate running accompanied by elevated theta power. Red arrows indicate seizure bouts
during pauses in the rat's intermittent locomotor behaviors.
964

Figure 2. Theta-frequency hypersynchrony occurs in rats exposed to Pb²⁺. **A.** Power spectral
densities (PSDs) of HC LFPs both for Pb^{2+}-exposed rats and for controls exhibited diminished delta and
heightened theta during running (**right**) relative to stationary bouts (**left**). Grey and red traces indicate
representative PSDs of simultaneous recordings from each site in the HC in individual rats. Thick black
traces (**left panels**) are PSDs from the site with the highest delta power during immobility, highlighting the
apparent tradeoff between delta and theta peaks across recording sites in each animal. **B**. Marked
elevation of theta power in Pb^{2+}-exposed rats relative to controls during two-second bouts of immobility
(**left**) and running (**right**) (1 sec bout **insets**). **C**. Stronger behavioral modulation of theta power in Pb^{2+}-
exposed rats than controls in terms of the absolute change (normalized by total power (0-25 Hz) (**left**) and
the relative change at each frequency (%) (**right**) between stationary and running bouts for control (black)
and Pb^{2+}-exposed rats (red). Enlargement of the low-frequency range of PSDs in the left and center panels
(**right**). **D.** Running speed modulation of delta, theta, slow, and high gamma. **E**. Similar patterns of
movement speed modulation of cycle-by-cycle amplitudes of theta and gamma oscillations (across all
behaviors) in Pb^{2+}-exposed rats and controls. Significance is indicated as follows: *, $p<0.05$; ** $p<0.01$; ***,
$p<0.001$.
981

Figure 3. Pb²⁺ exposure leads to disruptions in behavior-triggered high gamma activity. **A.** Running-
triggered speed- and stop-triggered speed transition for delta, theta, and high gamma amplitudes. **B.** Z-
normalized amplitudes for each frequency band across the 20-second triggering window for each peak in
running speed, emphasizing temporal dynamics disrupted at high gamma frequencies. **C.** Peak running
speed separated in 20 cm/sec bins, showing fast gamma disruptions (right-most red line plots).
Significance is indicated as follows: *, p<0.05.

Figure 4. Chronic Pb²⁺ exposure exacerbates absence seizures in the HC. A. SWD (SWDs) in the HC LFP from four Pb²⁺-exposed rats (see SupplFig3 for coincident mPFC SWDs). **B**. Spectrogram showing six seizures. SWD overlaid in white. **C**. Theta power distributions during immobility show evidence of

numerous seizure events creating a pronounced right tail or second peak at high theta power values. **D**.
The theta power to movement speed relationship shows cases where theta during seizures corresponds to
maximal theta during fast running (left). In other cases, weaker theta draws into relief the theta values
corresponding to SWDs (right). Inset distributions of theta and harmonic power, as in C. **E**. Delta
synchronizations, were minimal when power at the first harmonic of theta was elevated. **F**. Derivation of the
seizure detection algorithm. **G**. The theta fundamental frequency did not differ between groups, but Pb^{2+}-
exposed rats exhibited exacerbated seizure number, proportion of stationary time, and duration. There was
a trend toward significance in the strength of the seizures (p = 0.073). **H,** Sample spectrogram from one
Pb^{2+} rat treated with ethosuximide (200mg/kg, i.p.) eliminates seizure but not normal theta and delta
dominated running and stationary behaviors, respectively. Significance is indicated as follows: *, p<0.05;
***, p<0.001.
1005

1006

1007
1008
Figure 5. Prepulse inhibition was disrupted in adult male rats exposed to Pb^{2+}. There was no effect of
Pb^{2+} exposure at PN28, but there was marked inhibition of PPI in male rats at PN50 and PN120. Insert
graphs in each panel show the startle response at 120 dB. No effect on startle response was observed in
Pb^{2+}-exposed male rats relative to controls at any age (p's > 0.10). Thus, CELLE does not alter the
animal's sensitivity to a startle. However, CELLE presented with significant deficits in PPI in male rats
relative to controls at PN50 and PN120 (p's < 0.01).
1015

1016
1017

AGE	PN28		PN50		PN120	
SEX	Males	Females	Males	Females	Males	Females
Control	≤1.9 n=18	≤1.9 n=16	≤1.9 n=17	≤1.9 n=15	≤1.9 n=17	≤1.9 n=15
1500ppm PbAc	21.1 ± 0.9 n=18	20.9 ± 1.1 n=16	20.2 ± 1.1 n=17	22.1 ± 1.3 n=20	19.6 ± 1.3 n=17	24.3 ± 2.2 n=14
p-value	≤0.0001	≤0.0001	≤0.0001	≤0.0001	≤0.0001	≤0.0001

1018
1019 **Supplemental Table 1.** Blood Pb^{2+} levels (μg/dL) in male and female rats contributed to the data in Fig5
1020 and SupplFig7, respectively. Each value is the mean ± sem. Sample size n's = number of litters (one male
1021 and female animal per litter).
1022

1023

1024
1025

Supplemental Figure 1. Histological estimation of recording sites. Nissl-stained coronal brain sections are presented for all rats. The yellow arrows indicate electrode markings in each section. **A.** In nine rats, NeuroNexus probes (4 shanks) were implanted in the HC. Pb^{2+} exposed rats are in the upper left cluster, while controls are in the upper right cluster. **B.** In four rats, custom-made stainless-steel ($50\mu m$ tips) dual-site electrodes were implanted in the mPFC and HC. Pb^{2+} exposed rats are in the lower left cluster, while controls are in the lower right cluster. The cal bars are 1mm. Abbreviations: CA1, field CA1 of the HC; CA3, field CA3 of the HC; DG, dentate gyrus of the HC; and PL, prelimbic region of the mPFC.

1033

1034
Supplemental Figure 2. Precise delineation of stationary and running bouts. **A.** Head and body
direction (**top**) were used to derive rotational velocity and head angle (**middle**), which in turn were used to
identify bouts of running (boxed at **bottom**) while excluding rotational and reorienting behaviors (red points
and circles). **B.** Speed occupancy (**left**) and rotational velocity (**right**) measured throughout behavioral
sessions did not differ between control and Pb^{2+}-exposed rats, nor did bout-wise behavioral measures (**C**)
of running (**left**) and stationary bouts (**right**), including the numbers of bouts, median movement speeds,
and bout durations.
1042

1043

Prefrontal Cortical LFPs During a Spike-Wave Discharge in a Control Rat

Hippocampal LFPs During a Spike-Wave Discharge From a Control Rat

1044   ***Supplemental Figure 3. SWDs in a control rat.*** An example of an SWD recorded simultaneously in
1045   mPFC (above) and HC sites (below). The HC sites showed some variability in the LFP waveforms during
1046   the seizure and the continuation of theta oscillations following seizure offset. mPFC SWDs only showed up
1047   when HC SWDs did, not vice versa.
1048

1049

Supplemental Figure 4. Proximal-distal gradient of delta and theta power in HC. A. An apparent proximal-distal gradient in delta power across probe shanks in stationary control rats (**left**), was absent in Pb^{2+}-exposed rats (**right**). Black and red lines in A and B reflect the best fit to power estimates along each rat's dorsal and ventral rows of tetrodes (across shanks). **B.** Theta power at sites along the proximal-distal axis in control and Pb^{2+}-exposed rats during running bouts.

1055

1056

Supplemental Figure 5. Pb^{2+} exposure reduces theta frequency and disrupts running speed
modulation. **A.** Scatter plots of theta power and frequency taken from the power spectral density for each
recorded channel during each bout of stationary (**left**) and running behaviors (**right**) for all rats. Black and
green lines indicate means for control and Pb^{2+}-exposed rats, respectively. Dashed lines indicate the mean
theta frequencies for each group. **B.** Running speed modulation of theta frequency. **C.** Running speed
modulation of theta power.
1063

1064

Supplemental Figure 6. Running speed modulation of theta power is elevated in Pb²⁺-exposed rats.
A. Distributions of correlation coefficients between delta (blue) and theta (red) power estimates and running speed (one-sample t-tests, null = 0). Individual correlations were calculated across running bouts (1 sec) for each recorded channel from control rats (**top**) and Pb²⁺-exposed rats (**bottom**). **B.** As in A, for 2-sec running bouts. **C.** As in A, but comparing the delta and theta peak frequencies correlations to running speed. Significance as indicated.

Supplemental Figure 7. Prepulse inhibition is not disrupted in adult female rats exposed to Pb²⁺. No effect of Pb²⁺ exposure was found in female rats at any age (p's> 0.10). Insert graphs in each panel show the startle response at 120 dB. No effect on startle response was observed in Pb²⁺-exposed female rats relative to controls at any age (p's>0.10), nor was there an effect of Pb²⁺ relative to controls (p's > 0.10).