1
2
3 Chronic Lead Exposure Causes Theta and Gamma Hypersynchrony in the Hippocampus and Disrupts
4 Sensorimotor Gating

5
6
7
8
9 Abbreviated Title: LEAD EXPOSURE CAUSES HIPPOCAMPAL HYPERSYNCHRONY
10
11
12

13 Nathan W. Schultheiss[1], Jennifer L. McGlothan[2], Tomás R. Guilarte[2*], Timothy A. Allen[1,2,3*]
14
15 [1]Department of Psychology, Florida International University, Miami, FL, 33199
16 [2]Department of Environmental Health Sciences, Robert Stempel College of Public Health, Florida International University, Miami,
17 FL, 33199
18 [3]Cognitive Neuroscience Program, Florida International University, Miami, FL, 33199

19
20
21
Number of figures: 8
Number of supplemental figures: 1
Number of tables: 1
Number of text pages: 33
Total character count (with spaces):
Title: 111
Abstract: 1,847
Introduction: 5,678
Methods: 16,253
Results: 12,809
Discussion: 5,924
References: 9,792
Total: 59,514
35
36
37
38

39 *Corresponding Authors:
40
41     Timothy A. Allen, PhD
42     Department of Psychology
43     email: tallen@fiu.edu
44     Website: http://allenlab.fiu.edu/
45     Twitter: @AllenNeuroLab
46
47     Tomás R. Guilarte, PhD
48     Department of Environmental Health Sciences
49     email: tguilart@fiu.edu
50     website: https://stempel.fiu.edu/brain-behavior-environment-laboratory/
51     Twitter: @guilartelab
52
53     Address:
54     Florida International University
55     11200 SW 8th Street
56     Miami, FL, 33199

57
58

59 Conflict of Interest: The authors declare no competing financial interests.
60 Acknowledgements: This work was supported by R01 ES006189 to T.R.G. and R01 MH113626 to T.A.A.
61 Keywords: hippocampus, schizophrenia, lead neurotoxicity, pre-pulse inhibition

**ABSTRACT**

Chronic lead ($Pb^{2+}$) exposure from childhood contributes to an array of cognitive and behavioral

dysfunctions including impaired attention and intellectual ability, learning and memory deficits, and

delinquency. It is also an environmental risk factor for adult psychopathologies, most notably

schizophrenia. $Pb^{2+}$ is a potent N-methyl-D-aspartate receptor (NMDAR) antagonist and exposure during

early life elicits a cascade of cellular neurotoxic effects that alter the developmental trajectory leading to a

loss of parvalbumin-expressing interneurons in hippocampus and altered synaptic transmission. Little is

known, however, about the impact of chronic $Pb^{2+}$ exposure on hippocampal network dynamics which

serve as a link between cellular-molecular effects and cognitive-behavioral consequences of $Pb^{2+}$

neurotoxicity. Here, we tested the effects of chronic $Pb^{2+}$ exposure on local field potential activity (LFP) in

the hippocampus of freely-behaving rats. We demonstrate that $Pb^{2+}$ exposure causes striking

hypersynchrony in both the theta- and gamma-frequency bands, in addition to more modest changes in

delta. Next, we tested the effect of $Pb^{2+}$ exposure on prepulse inhibition of the acoustic startle reflex (PPI)

in male and female rats at different developmental timepoints. We found that adult males (PN50 and 120),

but neither females nor juvenile males, showed reduced PPI that could not be attributed to changes in the

startle reflex. This pattern recapitulates sex- and age-dependencies of PPI disruption in schizophrenic

patients. Overall, these results are consistent with the hypothesis that $Pb^{2+}$ is an environmental risk factor

for psychopathology in adulthood, especially those symptoms related to cognitive and sensory-motor

gating processes that depend on rhythmic coordination of network activity in the hippocampus.

**Introduction**

Environmental exposure to lead ($Pb^{2+}$) is a major health problem affecting millions of people

worldwide. $Pb^{2+}$ exposure in children has been linked to reduced intellectual abilities, increases in

delinquent and antisocial behaviors, and attention deficit and hyperactivity disorder (Needleman et al.,

2002; 2004; Miranda et al., 2007; Canfield et al., 2003; Caito & Aschner, 2017). Further, early life $Pb^{2+}$

exposure has been proposed as an environmental risk factor for adult psychopathology, most notably with

regard to schizophrenia (Opler et al., 2004; 2008). A current leading hypothesis is that schizophrenia is

best understood as the result of interactions between genetic and environmental factors (Guilarte et al.,

2012), with evidence that $Pb^{2+}$ exposure exacerbates cognitive and behavioral deficits in the DISC1 knock

out mouse model of schizophrenia (Abazyan et al., 2014).

Understanding the relationship between $Pb^{2+}$-exposure and mental health pathologies requires a

neurobiological approach to establishing the link between genetic and environmental factors. At the cellular

and molecular levels, much is known about $Pb^{2+}$ effects on the developing brain. $Pb^{2+}$ is a potent N-methyl-

D-aspartate (NMDA) receptor antagonist (Guilarte, 1997; Guilarte & Miceli, 1993; Guilarte & Miceli, 1992).

Low-to-moderate $Pb^{2+}$ levels in the brain (which has no endogenous function) leads to NMDAR

hypoactivity, which can alter the developmental trajectory of the NMDAR subunits (Guilarte & McGlothan,

1998; Nihei et al., 2000), and impair the maintenance synaptic long-term potentiation (Nihei et al., 2000). At

the cellular level, $Pb^{2+}$ exposure causes a reduction in the number of parvalbumin-labeled GABAergic

interneurons in the hippocampus and agranular medial prefrontal cortex (Stansfield et al., 2015), a hallmark

cellular pathology in schizophrenic brains (e.g., Hashimoto et al., 2003). At the level of the whole brain,

annual childhood blood $Pb^{2+}$ levels have been found to be inversely correlated with cortical gray matter

volume (Brubaker et al., 2010). Likewise, an electroencephalography (EEG) study of $Pb^{2+}$-exposure, as

measured by tooth $Pb^{2+}$, showed there are significant alterations in delta, theta and gamma bands related

to cognitive function, the severity of which corresponded to the exposure levels (Needleman, 1983).

However, the bridge between the cellular-molecular and cognitive-behavioral levels is not well

understood. Linking these levels requires animal models of $Pb^{2+}$ exposure allowing for invasive techniques

that can establish a causal link between $Pb^{2+}$ exposure, specific neurobiological mechanisms, and

behaviorally-relevant network consequences. Here, we performed chronic *in vivo* hippocampal recordings

from a well-established rodent model of $Pb^{2+}$ exposure during open field behaviors (Schultheiss et al.,

2020). This initial study of the rhythmic network effects of chronic $Pb^{2+}$ exposure focused on the

hippocampus because chronic $Pb^{2+}$-exposure causes reliable and severe deficits in hippocampal-

dependent spatial learning (Kuhlmann et al., 1997; Jett et al., 199; Munoz et al., 1988) which can be

ameliorated with environment enrichment (Cao et al., 2008; Guilarte et al., 2002), similar to other

hippocampal insults. Likewise, $Pb^{2+}$-exposure causes contextual but not cued fear conditioning deficits

(Jaako-Movits et al., 2005; McGlothan et al., 2008; Wang et al., 2016), in addition to fear extinction

impairments (McGlothan et al., 2008). These behavioral results are consistent with the central role of

hippocampal networks in mediating the effects of chronic $Pb^{2+}$-exposure in psychotic disorders.

Our recordings and analysis focused on the spectral content of local field potentials (LFPs) from the

hippocampus in of $Pb^{2+}$-exposed rats using multisite silicon probes during stationary and running

behaviors. LFPs are a good choice to examine network properties because they reflect the rhythmic

synchrony among neurons aggregating local and long-range synaptic potentials. We observed several key

abnormalities, most notably excessive hypersynchrony in the theta and gamma bands and abnormal

behavior-related changes the theta and gamma bands. Hypersynchrony is detrimental to local network

processing and has been observed in other disorders. For example, hippocampal hypersynchrony in the

theta and slow gamma ranges have also been observed in the LFP recordings of the freely-behaving

*Fmr1-KO* mouse, a genetic model of Fragile X syndrome which is characterized by intellectual disability

(Arbab et al., 2018).

Next, because over activation of the hippocampus is known to cause impaired sensory-motor gating

(Bast et al., 2001a,b; Bast et al. 2003), we followed the network finding up by looking at the consequences

of Pb2+-exposure on prepulse inhibition of the startle reflex (PPI). We found that $Pb^{2+}$-exposure caused

large and reliable sensory-motor gating deficits in male, but not female, adult rats in developmentally-

relevant pattern (PN50 and PN120, but not PN28). These effects could not be attributed to baseline startle

levels, and are in line with the $Pb^{2+}$ hypothesis of schizophrenia because PPI impairments are a hallmark

pathological phenotype in the disease.

Altogether the present findings show that $Pb^{2+}$ exposure leads to pathological theta and gamma

hypersynchrony in the hippocampus and causes sensory-motor gating deficits. These abnormal

hippocampal network effects may explain both the cognitive and the sensory-motor gating deficits seen in

adult psychopathologies resultant of Pb2+ exposure, and suggests targeting hippocampal network

restoration as a therapeutic target.

**Methods**

*Subjects.* Nine adult male Long-Evans rats were used in electrophysiological experiments. Five of

these were chronically exposed to lead ($Pb^{2+}$), and four were controls. Two hundred thirty rats were used in

PPI experiments. These rats were evenly divided into twelve groups (18-21 per group) corresponding to all

combinations of (1) $Pb^{2+}$ exposure and controls, (2) males and females, and (3) three ages at the time of

testing. All rats were individually housed in a climate-controlled vivarium on a reversed 12hr light/dark

cycle, and experiments were conducted during the dark cycle active periods. Rats had free access to food

and water when in their home cages, but no food or rewards were available during behavioral sessions. All

studies were carried out in strict accordance with the recommendations in the Guide for the Care and Use

of Laboratory Animals of the National Institutes of Health under protocols approved by Florida International

University Institutional Animal Care and Use Committee. Data from the four control rats in the

electrophysiological experiments were previously analyzed to establish the relationships of hippocampal

spectral modes to behaviors (Schultheiss et al., 2020). The present study focuses on the effects of $Pb^{2+}$

exposure and all analyses presented here are new.

*Breeding and Chronic $Pb^{2+}$ Exposure.* Prior to breeding, adult female rats (225-250 g) were

purchased from Charles River Laboratories and randomly assigned to a Prolab RMH-1000 diet containing

0 (control) or 1500 ppm lead acetate (PbAc) (Dyets, Inc., Bethlehem, PA). After 10-14 days of acclimation

to the diet, each dam was paired with a non-exposed adult male Long Evans rat (250-275 g; Charles River

Laboratories) to breed over 3 days. Litters were culled to 10 pups on postnatal day one (PN1). Dams were

maintained on their respective diet, and at PN21 male pups were weaned onto the same diet and

maintained for the duration of the experiments.

*Silicon probe implants.* All electrophysiological procedures were the same as we have previously

described (see Schultheiss et al., 2020 for details). Briefly, silicon probes with 32 recording electrodes

arranged as eight tetrodes were surgically implanted for recording local field potentials (LFPs) from dorsal

hippocampus (HC). The eight tetrodes were distributed across four shanks of the probe at tip-to-tetrode

depths of 78 μm and 228 μm (NeuroNexus A4X2-tet-5mm). Shanks were separated by 200 μm giving each

probe a total length of 0.67 cm which was oriented medial-lateral when implanted to sample along the

proximal-distal axis of the CA1 subregion of HC. Electrode impedances were $1.23 \pm 0.32$ MΩ.

*Surgical procedures.* The rats used in electrophysiological experiments weighed 600-700 g at the

time of surgery. General anesthesia was induced with isoflurane (5%) mixed with oxygen (0.8 L/min) that

was continuously delivered throughout surgery at 1-4% as needed. Body temperature was monitored

throughout surgery, and a Ringer's solution (5% dextrose) was given periodically to maintain hydration (1

ml increments every 60-90 minutes). Once the hair over the scalp was cut, rats were placed in a

stereotaxic device, and glycopyrrolate (0.5 mg/kg, s.c.) was administered to assist respiratory stability.

Ophthalmic ointment was then applied to the eyes. Four injections of marcaine were made to the scalp

(~0.1 ml, 7.5 mg/ml, s.c.), and a rostrocaudal incision was made exposing the skull. The pitch of the skull

was leveled between bregma and lambda, two stainless steel ground screws were placed in the left

parietal bone, and five titanium support screws were anchored to the skull. Rectangular craniotomies were

drilled to accommodate the probe shanks centered on coordinates A/P -3.24 mm, M/L 2.7 mm. After

removal of the dura, the probe was lowered on the stereotaxic arm until the tips of the probe shanks were

just above the cortical surface. Then, the ground wire was attached to the ground screws, and implants

were lowered further until the electrodes reached a depth of ~2.8 mm below the cortical surface. A thin

layer of sodium alginate was applied in the craniotomy to protect the exposed surface of the brain, and

dental cement was applied to permanently secure the implant to the skull screws. Neosporin® was applied

to the skin surrounding the implant, and flunixin (2.5 mg/kg, s.c.) was given for analgesia. Following

surgery, rats were allowed to rest on a heating pad until ambulatory, prior to being returned to their home

cages. Neosporin® and flunixin were given again on the day following surgery, and rats were monitored

closely for five days to ensure that body weight was maintained. Daily Baytril injections (s.c.) were also

given for five days to protect against postsurgical infection. A minimum of one week was reserved for

recovery prior to beginning experiments.

*Behavioral Setup.* All electrophysiological experiments were conducted in a large open field

environment (122 x 118 x 47 cm). A custom-made white cutting board served as the floor of the behavioral

arena, and the interior walls were lined with white vinyl. The arena was mounted on stilts 72 cm above the

floor and the exterior was wrapped in grounded Faraday shielding. Two video cameras were used for

behavioral tracking with a bird's eye view. These were mounted 28 cm apart from one another,150 cm

above of the arena floor, providing ~4.6 pixel/cm resolution in recorded videos and corresponding tracking

data.

The entire behavioral apparatus was enshrouded with black curtains suspended from the ceiling

and extending to the floor. Small gaps at the Velcro® closures between the four curtain panels were used to

observe rats covertly to monitor that the head-stage, cables, and elastic tethers were not tangled, and to

confirm that rats were awake during periods of relative inactivity. All room lights were kept off during

recording sessions, and the white arena was illuminated dimly with red light from LED strips mounted

above at the level of the video cameras.

During behavioral sessions rats were placed on the arena floor and recorded without any

intervention except for infrequent behavioral checks or to adjust cables or connections.

*Electrophysiological recordings.* Throughout each experiment wide-band neural data were acquired

with a Plexon OmniplexD recording system with A/D conversion at 40 kHz (16 bit). A digital head stage

processing subsystem (Intan, 32 channel) passed continuous wide-band data referenced against the

ground screws (low cutoff = 0.7 Hz) to a hard drive using PlexControl data acquisition software. For

analysis of LFP recordings, wideband data was down-sampled to 2 kHz. A maximum of one channel per

tetrode was included in our analyses, and twelve tetrodes were excluded for poor signal. As a result of

these restrictions, 30 channels recorded from $Pb^{2+}$-exposed rats were analyzed, and 30 channels from

control animals were analyzed. All of the selected channels were included in all analyses, providing a

robust sampling of hippocampal recording sites for each group.

*Behavioral tracking.* The physical and software settings for the two video cameras used for

behavioral tracking were tuned independently to optimize detection of body contours on one camera and to

optimize tracking of head-mounted LEDs on the other. LED tracking entailed tabulating the coordinates of

two clusters of LED florets (one green and one blue) that were offset ~2 cm laterally from the head-stage at

~3 cm above the rat's head. Each LED cluster comprised three florets oriented upward and slightly

outward. This construction enabled consistent LED tracking despite behaviors in which rats oriented their

heads away from the level plane. Body location was calculated from difference images obtained by

subtracting an image of the empty arena from each frame capture during an experiment. Movement

speeds were calculated from these body coordinates. The location of the head was calculated as the

midpoint between the two LED clusters for each timepoint. Using Cineplex software (Plexon), LED tracking

data was collected online during experiments, and body contour tracking was derived offline from recorded

videos. Video tracking frame captures (30 fps) were synchronized in real time to the simultaneously

recorded neural data via a central timing board (Plexon).

*Co-registration of LED and body tracking data.* Following acquisition, all analyses of behavioral data

were performed with custom-written Matlab scripts (MathWorks, Natick, MA). First, the coordinate system

for camera one was mapped onto the coordinate system for camera two, enabling integration of the two

types of tracking data. We then derived for each timestep the animal's head direction (HD) relative to the

environment (from the axis between the two head-mounted LED clusters); body direction (BD) relative to

the environment (from the midpoint between the LEDs to the center of the body); head angle (HA) relative

to the body (using the LED axis relative to BD); and the rotational velocity of the body.

*Definitions of running and stationary bouts.* Rats exhibit a wide range of behaviors that involve body

and head movements at substantial speeds. These movements appear in tracking data as changes in body

and LED coordinates, just as running does. To isolate bouts of locomotion from other behaviors including

grooming and rearing, we first required that movement speed be maintained above 5 cm/sec for a

minimum duration of 1.025 or 2.05 seconds with a minimum peak speed of 15 cm/sec. We then eliminated

all instances where head angle exceeded 35 degrees or body rotational velocity exceeded 52.5 degrees

per second at any time. Stationary bouts were required to meet the same minimum duration, HA, and

rotational criteria and not exceed a maximum movement speed of 5 cm/sec.

*Derivation of oscillatory metrics.* We evaluated the correspondence to animals' movement speeds

of LFP dynamics in the delta-, theta-, and slow and fast gamma-frequency bands (1-4 Hz, 6-10 Hz, 25-55

Hz, and 65-120 Hz, respectively) using amplitude timeseries for each band and using the power spectral

density of the LFP. We derived amplitude timeseries for each band by resampling to 2kHz the peak-to-

trough amplitudes of each oscillatory cycle of the respective band-pass filtered LFPs. The spectral content

of HC recordings was estimated using Welch's method for each stationary and running bout. The 1.025

and 2.05 second minimum durations required in our bout definitions were chosen as a compromise

between the relatively fast timescale of rats' locomotor behaviors, e.g. bouts of running and interspersed

pauses, and the reduction in number of oscillatory cycles (particularly for delta) contributing to amplitude

measures for shorter time windows.

Estimates of delta and theta power from the PSDs for individual behavioral bouts were defined as

the mean values across the respective frequency ranges. For each recording site, delta and theta power

estimates were defined as the medians of these values across all bouts of each type. The distributions of

these values for each group are plotted as insets in Figure 3.

Delta and theta peak frequencies and peak power estimates were measured directly from the PSDs

for behavioral bouts. Scatterplots of these values are shown in Figure 4. Correlation coefficients relating

these measures to running speed were derived for each recording site across all bouts of each type.

Distributions of these correlation coefficients for each group are shown in Figure 5.

We defined behavioral modulation of oscillatory metrics as the difference between running bouts

and stationary bouts. First, mean PSDs across bouts of each type were derived for each recording site.

Then behavioral modulation was calculated for each site as the subtraction of the mean stationary PSD

from the mean running PSD. Behavioral modulation for each site was then normalized either by the area of

the mean stationary PSD (as in Figure 6A) or by dividing by the mean stationary PSD to give the percent

change at each frequency (as in Figure 6B). Behavioral modulation of delta and theta were then defined as

the mean values across the respective bands for each recording site. Distributions of these values are

shown as scatterplots in Figure 6.

To further evaluate the effects of $Pb^{2+}$ exposure on oscillatory synchronization in each frequency

band, we collected peak-to-trough amplitude measurements for every cycle of band-pass filtered LFPs for

each recording site for all bouts of each type. The normalized distributions of these amplitude

measurements are plotted in Figure 7.

*PPI experiments*. Pre-pulse inhibition of acoustic startle (PPI) was assessed in male and female

rats at PN28, PN50, and PN120 following a protocol adapted from Abazyan et al. (2014). Four identical

SR-LAB startle chambers (San Diego Instruments Inc., San Diego, CA, USA) were used to measure startle

responses. Rats were placed inside the startle chamber into a mounted Plexiglas® cylinder with an

accelerometer. Broadband background noise and acoustic stimuli were delivered via a speaker on the

ceiling of the chamber. Sound levels were calibrated using a digital sound level meter, and accelerometer

sensitivities for each chamber were calibrated daily. Presentation of acoustic stimuli was controlled by the

SR-LAB software and interface system, and accelerometer measurements were recorded. Each

experimental session started with a five-minute period during which rats acclimatized to 68 dB background

noise that was continuous throughout the session. Next, rats' average startle responses were calculated as

the mean of 10 acoustic pulses (120dB, 100ms) presented at a random inter-trial intervals. Rats were left in

the enclosure for an additional five minutes prior to beginning PPI trials. Rats were exposed to the following

trial types: pulse-alone trial (120dB, 100ms); pre-pulse alone (90dB, 20ms); and four pre-pulse–pulse

combinations (20 ms pre-pulse presented 80 ms before the 120 dB, 100 ms pulse) using one of the five

pre-pulse intensities: 74, 78, 82, 86 and 90 dB. Each session consisted of six presentations of each trial

type presented in a pseudorandom order with random inter-trial intervals. The average response during the

100 ms recording window beginning at stimulus onset was used as the measure of startle responses. PPI

was quantified for each animal as the percent change in startle amplitude on pre-pulse–pulse trials relative

to pulse-alone trials.

*Blood $Pb^{2+}$ Levels.* Blood $Pb^{2+}$ levels of rats used in PPI experiments (Table 1) were measured in

samples obtained from cardiac puncture under sodium pentobarbital anesthesia. Samples were prepared

and measured following protocols provided using the LeadCare Plus system (Magellan Diagnostics, N.

Billerica, MA).

*Histology*. At the completion of data collection, rats used in electrophysiological experiments were

anesthetized with isoflurane (5%) mixed with oxygen (800 ml/min) and marking lesions were made with a

NanoZ (Plexon) to deliver 10μA for 10s at each electrode locations. About an hour later, rats were

transcardially perfused with 100 ml phosphate-buffered saline (PBS), followed by 200 ml of 4%

paraformaldehyde (PFA, pH 7.4; Sigma-Aldrich, St. Louis, MO). Blood $Pb^{2+}$ levels of these rats were

measured in samples obtained from cardiac puncture during the perfusion. Brains were post-fixed

overnight in 4% PFA and then placed in a 30% sucrose solution for cryoprotection. Frozen brains were cut

on a Leica CM3050 S cryostat (40 μm; coronal) and Nissl-stained. Marking lesions were mapped onto

plates of the Paxinos & Watson Atlas (2004).

307                  **Results**

Our analyses focused on relationships between locomotor behaviors and concurrent hippocampal

synchronization in delta-, theta-, and gamma-frequency bands to assess the impact of $Pb^{2+}$ exposure on

network dynamics in hippocampus and possible commonalities with the pathophysiology of schizophrenia.

For this we compared the spectral content of local field potential (LFP) recordings from hippocampus

(Figure 1A&B) in rats chronically exposed to $Pb^{2+}$ and controls.

*Overall Behavior*. Recordings were made throughout behavioral sessions during which rats freely

explored a large 'open field' arena (122 cm x 118 cm). No task was imposed during behavioral sessions; no

rewards were delivered; and environmental stimuli and experimenter contact were strictly minimized. Under

these conditions, we did not observe any differences between $Pb^{2+}$-exposed rats and controls in

continuous behavioral measures including movement speed, spatial orientation, and rotation velocity

(Supplemental Figure 1). Rats in both groups were more often stationary (~20-40% of session time) than

moving at any other speed, but all rats exhibited sporadic episodes of exploration yielding a consistent

peak in speed occupancy at ~7 cm/s. These active episodes typically comprised running, rearing, and

olfactory sampling behaviors, and were interspersed amongst periods of grooming and awake inactivity. To

sharpen our analysis, we isolated precisely-defined bouts of running and stationary behaviors, and

excluded rearing, grooming, and reorienting behaviors by imposing exclusion thresholds for speed,

rotation, and head angle as in Schultheiss et al. (2020). Bout-wise measures of running and stationary

behavior were also equivalent between $Pb^{2+}$-exposed rats and controls (Supplemental Figure 1D and 1E).

*Movement speed modulation of delta, theta, and gamma oscillations.* To provide initial insight into

relationships between neural rhythms and locomotor behavior, we derived instantaneous amplitude

timeseries for delta-, theta-, and gamma-frequency bands by resampling peak-to-trough measurements for

each cycle of the respective band-pass filtered LFP. The amplitudes of delta, theta, and fast gamma

oscillations were all consistently modulated by animals' movement speeds in both control and $Pb^{2+}$-

exposed rats (Figure 1C). Whereas theta and gamma were positively correlated to speed, delta was

negatively correlated to movement speed (Schultheiss et al., 2020). Figure 1C shows these relationships

for all recording sites in all rats (n = 30 per group), illustrating the qualitative similarity between groups.

Movement speed modulation of slow gamma amplitude was generally less prominent or absent, but some

recording sites in both control and $Pb^{2+}$-exposed animals exhibited negative correlations to speed.

As we have recently demonstrated, distinct modes of delta- and theta-dominated hippocampal

network activity can transition rapidly in tandem with the stops and starts of rats' locomotor trajectories. In

that study delta and theta power were orthogonal across modes such that strong delta synchronization

occurred when theta was weak and vice versus. This pattern of exclusivity between hippocampal modes

may reflect distinct neurophysiological processes or cognitive functions contributing to spatial navigation

and memory (Schultheiss et al., 2020). Thus, to minimize blurring the oscillatory dynamics of distinct

modes that accompany behaviors occurring in rapid succession, we focused our subsequent analyses on

hippocampal synchronization during the precisely-defined stationary and running bouts.

Power spectral densities (PSDs) of the HC LFP showed higher delta power during stationary bouts

and higher theta power during running bouts for both $Pb^{2+}$-exposed animals and controls. Figure 2 A and B

illustrate representative PSDs for two rats in each group, and show considerable variability across

recording sites from individual rats in the delta power exhibited during stationary bouts. Linear fits to delta

power across individual recording sites from each rat revealed a consistent distal-proximal gradient of

increasing delta power in control animals that was not evident in $Pb^{2+}$-exposed animals (Fig. 2C). No

consistent theta power gradient was observed during running bouts for either group (Fig. 2D).

$Pb^{2+}$-exposed rats exhibited marked elevation of theta power relative to controls during both

stationary bouts (1 sec: p=0.01, t=-2.6, df=58; 2 sec: p=0.04, t=-2.1, df=58) (Fig. 3A) and running bouts

(1 sec: p=0.002, t=-2.8, df=58; 2 sec: p=0.002, t=-3.2, df=58) (Fig. 3B). Delta power in $Pb^{2+}$-exposed rats

was also elevated relative to controls during two second stationary bouts (p=0.05, t=-2.1, df=58), but this

trend was weak for one second bouts that are generally more strongly impacted by behavioral variability

(p=0.14, t=-1.5, df=58). No differences in delta power between groups were observed during running.

We also found that theta frequency was substantially lower in $Pb^{2+}$-exposed rats compared to

controls during both stationary bouts (control: 7.2 ± 1.1, Pb: 6.8 ± 1.0; $p=8.1 \times 10^{-6}$, t=-4.5, df=842) and

running bouts (control: 8.2 ± 0.4, Pb: 7.8 ± 0.6; $p=3.9 \times 10^{-25}$, t=-10.77, df=702) (Fig.4A). Interestingly,

pairwise 2-D scatterplots of theta power, theta frequency, and running speed for each bout type suggest

that (1) covariation of theta power and theta frequency was disrupted in $Pb^{2+}$-exposed rats relative to

controls (Fig. 4A, right; green  vs black lines); and (2) running speed modulation of theta frequency was

confined to a narrower range of speeds (Fig. 4B, right; green  vs black lines); and in contrast, (3) running

speed modulation of theta power occupied similar parameter spaces in $Pb^{2+}$-exposed rats and controls

(Fig. 4C, right; green vs black lines). Taken together, these observations suggest that $Pb^{2+}$ exposure may

disrupt the fidelity of relationships between theta synchronization and locomotor behavior. To evaluate this

quantitatively, we focused on two second running bouts and compared distributions across recording sites

of the correlation coefficients between running speed and theta (and delta) measures. As expected, we

found positive correlations of theta power and frequency to running speed for the majority of recording sites

in both $Pb^{2+}$-exposed rats and controls (Fig. 5). Importantly, we also found that the fidelity of theta power to

running speed was heightened in $Pb^{2+}$-exposed rats relative to controls (p=0.0149, t=2.511, df=58),

potentially reflecting a reduction in the dimensionality of hippocampal representations in favor of locomotor

parameters, i.e. limiting the integration of multidimensional information arriving at the hippocampus from

widespread cortical regions. However, in contrast to the apparent restriction of running speed modulation of

theta frequency to a narrower range of speeds in $Pb^{2+}$-exposed rats (Fig. 4B), we found that correlations

between running speed and theta frequency were similar in strength between groups (Fig. 5C, top vs

bottom; p=0.192, t=-1.319, df=58).

Interestingly, delta power was unrelated or even positively correlated to running speed in control

animals across running bouts, but it was significantly negatively correlated to running speed in

$Pb^{2+}$exposed rats. Accordingly, comparison of running speed modulation of delta power showed a

significantly strengthened negative correlation in $Pb^{2+}$exposed rats (delta power, p=0.0026, t=3.143, df=58).

It is important to consider, however, that these correlations of delta parameters to running speed were

taken across running bouts when delta is often weak or absent. As a result, they may reflect theta-delta

interactions that are distinct from the differences between delta- and theta-dominated modes that parallel

behavioral state changes.

Next, we evaluated differences in behavioral modulation of the PSD between $Pb^{2+}$-exposed rats

and controls by subtracting mean PSDs for stationary bouts (normalized by the integral of the PSD) from

the corresponding mean PSDs for running bouts for each recording site. This approach yielded 'difference

spectra' that revealed stronger behavioral modulation of theta and the first harmonic of theta in $Pb^{2+}$-

exposed animals relative to controls (Fig. 7A). Augmentation of the difference in theta synchronization

between behavioral states may reflect increased susceptibility of HC to entrainment by theta rhythmic

inputs recruited during running or to self-organization of theta oscillations with hippocampal excitation. We

also generated difference spectra to capture the percent change of power at each frequency between

stationary and running bouts (Fig. 7B). Interestingly, when quantified in this way, behavioral modulation of

the theta harmonic was robustly elevated in $Pb^{2+}$-exposed rats relative to controls, whereas behavioral

modulation of fundamental theta did not differ significantly between groups. This analysis also revealed a

possible reduction of behavioral modulation of relative oscillatory power in the 40-60 Hz range.

*Effect of $Pb^{2+}$ exposure on oscillatory cycle amplitudes.* Our analyses so far have evaluated the

spectral content of HC LFPs during stationary and running bouts and have robustly shown that exposure to

$Pb^{2+}$ yields marked hypersynchrony and elevated behavioral modulation of the theta rhythm. Our approach

has been to compare PSDs directly as well as correlations between power spectra and locomotor behavior.

However, these analyses are not well suited to address potential effects of $Pb^{2+}$ on gamma-frequency

synchronization. To evaluate the impact of $Pb^{2+}$ on gamma-frequency synchronization and to further

explore delta and theta effects, for each frequency band we compiled the distributions of the peak-to-trough

amplitudes of individual cycles during stationary and running bouts (Fig. 8A). As expected, delta, theta, and

fast gamma amplitudes were all strongly modulated by behavior for control rats and for $Pb^{2+}$-exposed rats.

In contrast, slow gamma did not show differences in amplitude between bout types. Comparisons of $Pb^{2+}$-

exposed rats to controls demonstrated pronounced increases in cycle amplitudes for theta- and gamma-

frequency bands during stationary bouts (theta: $p=1.4 \times 10^{-236}$, t=-32.9, df=114,635; slow gamma: p=0, t=-

78.2, df=554,144; fast gamma: p=0, t=-168.2, df=1,276,491) and during running bouts (theta: p=0, t=-

116.5, df=102,653; slow gamma: p=0, t=-90.5, df=491,214; fast gamma: p=0, t=-101.32, df=1,127,377).

Although statistically significant, differences in delta cycle amplitudes between $Pb^{2+}$-exposed rats and

controls were small relative to those for theta and gamma (stationary: $p=2.9 \times 10^{-8}$, t=-5.5, df=41741;

running: $p=1.9 \times 10^{-16}$, t=8.2, df=34211). It is important to consider that with such large numbers of individual

cycle amplitude measurements, statistical significance may not be informative.

*Disruption of sensorimotor gating.* In a last series of experiments, we tested the effect of $Pb^{2+}$

exposure on PPI of acoustic startle, an assay of sensorimotor gating known to be impaired in adults with

schizophrenia. Figure 8 shows the pattern of results analyzed in males and females across development

(PN28, PN50, and PN120). We tested the baseline acoustic startle responses in all groups and found no

significant differences (all t-test p's >0.05), suggesting any differences observed in PPI reflect sensory-

motor gating.

*Males.* We found effects of Pb2+-exposure in adult males (PN50 and PN120), but not at PN28,

demonstrating a developmental trajectory for the emergence of PPI deficits that is similar to schizophrenia.

At PN28, a two-way ANOVA showed a main effect of dB level ($F_{2.57, 89.83}=68.89$, p<0.001), no main effect of

Pb2+-exposure ($F_{1,35}=0.54$, p=0.467), and no level x exposure interaction ($F_{2.57, 89.83}=2.56$, p=0.069). At

PN50, there was a main effect dB level ($F_{3.12, 109.23}=53.49$, p<0.001), a main effect of Pb2+-exposure

($F_{1,35}=11.32$, p<0.01), but no level x exposure interaction ($F_{3.12, 109.23}=1.34$, p=0.266). At PN120, there was a

main effect dB level ($F_{2.88, 115.11}=59.59$, p<0.001), a main effect of Pb2+-exposure ($F_{1,35}=9.30$, p<0.01), but

no level x exposure interaction ($F_{2.88, 115.11}=2.00$, p=0.120).

*Females.* We found no effects of Pb2+-exposure in female rats at any age. At PN28, a two-way

ANOVA showed a main effect of dB level ($F_{2.56, 92.28}$=42.74, p<0.001), no main effect of Pb2+-exposure

($F_{1,36}$=0.01, p=0.942), and no level x exposure interaction ($F_{2.56, 92.28}$=0.17, p=0.888). At PN50, there was a

main effect dB level ($F_{2.89, 101.26}$=90.26, p<0.001), no main effect of Pb2+-exposure ($F_{1,35}$=0.28, p=0.602),

and no level x exposure interaction ($F_{2.89, 101.26}$=0.61, p=0.606). At PN120, there was a main effect dB level

($F_{2.82, 104.17}$=57.96, p<0.001), no main effect of Pb2+-exposure ($F_{1,37}$=0.02, p=0.878), and no level x

exposure interaction ($F_{2.82, 104.17}$=0.61, p=0.597).

438                                              **Discussion**

439            We analyzed the spectral content of hippocampal LFPs and found that rats chronically exposed to

$Pb^{2+}$ exhibited hypersynchrony in the theta- and gamma-frequency bands both while stationary and during

running. Behavioral modulation of theta power was also stronger in $Pb^{2+}$-exposed rats relative to controls,

suggesting increased efficacy of the neural mechanisms for theta rhythmogenesis as well as elevated

baseline levels of theta synchronization. We also found that theta frequency was markedly lower in

$Pb^{2+}$-exposed animals than controls, and running speed modulation of theta frequency was restricted to a

narrow range of speeds. Thus, chronic $Pb^{2+}$ exposure increased theta synchrony while reducing theta

frequency and rendering it less flexible in response to changes in behavior. Slower frequency oscillations

are generally associated with synchronization of larger populations of neurons, and intrinsic coupling within

hippocampal networks and active membrane properties of hippocampal neurons are well suited to self-

organize theta-rhythmic activity (Goutagny et al., 2009, Stark et al., 2015). Importantly, parvalbumin

expressing interneurons (PVGIs) are major contributors to theta rhythmic synchronization of hippocampal

principal cells (Amilhon et al., 2015), but PVGIs are reduced in animals chronically exposed to $Pb^{2+}$. One

potential explanation of this inconsistency is that the reduction in number is compensated for by increased

local connectivity of the remaining PVGIs to maintain an appropriate excitatory-inhibitory balance within

hippocampal networks. Such a homeostatic function might endow the remaining PVGIs with stronger

influence over the spike timing of larger populations of synaptic targets leading to excessive synchrony.

Interestingly, ablation of NMDARs on PVGIs (presumably without reducing PVGI number) yields the

opposite pattern of effects on theta than $Pb^{2+}$ exposure, increasing theta frequency while reducing power,

and resulting in disrupted hippocampal representations and impaired memory (Korotkova et al., 2010).

In addition to theta hypersynchrony, we also found increased gamma-frequency synchronization in

$Pb^{2+}$-exposed animals. Elevated gamma activity in normal animals is integral to cognitive processes

including attention, perception, and memory encoding (e.g. Fell et al., 2001, Jutras et al., 2009, Yamamoto

et al., 2014, Fries et al., 2001), but $Pb^{2+}$-exposed animals show deficits in cognition and memory. One

possibility is that the elevated gamma synchronization seen in $Pb^{2+}$-exposed animals is in fact a

consequence of the excessive theta synchrony that we have described. Pastoll and colleagues (2013)

showed that theta-frequency excitation of layer II of medial entorhinal cortex *in vitro* was sufficient to elicit

well-timed gamma oscillations nested with theta cycles. This demonstrates that the local microcircuitry

responsible for gamma synchronization can be engaged by theta-rhythmic excitation in the absence of an

intact animal, and this mechanism could be strongly driven by theta hypersynchrony in $Pb^{2+}$-exposed

animals.

Importantly, hypersynchrony reflects reduced stochasticity in network dynamics which can be

detrimental to encoding capacity and flexibility. We further observed increased behavioral modulation of

theta power. It is important to note that this is not behavioral modulation of encoding but rather a greater

increase in theta synchronization that could reflect a further decrease in encoding capacity or flexibility

during running. For example, single neurons in parahippocampal structures commonly show conjunctive

representations of multiple spatial dimensions. When the spatial tuning of conjunctive grid-head direction

cells in medial entorhinal cortex is disrupted by inactivation of the medial septum, the fidelity of those cells

for head direction tuning is increased (Brandon et al., 2013). Here, we revealed greater fidelity of theta

power to running speed, which may also reflect a reduction in the dimensionality of hippocampal encoding.

This pattern of effects would be consistent with reduced integration of information types or modalities of

sensory input accompanying elevated entrainment of hippocampal populations to theta and stronger

encoding fidelity for running speed. This interpretation could also incorporate our finding of disrupted

sensorimotor gating resulting from $Pb^{2+}$ exposure. If the hippocampus is excessively involved in encoding

behavior, it may be less receptive to incoming sensory information. Furthermore, $Pb^{2+}$ neurotoxicity and

schizophrenia are both related to behavioral inflexibility or cognitive-behavioral perseveration consistent

with this general formulation.

Related to the hypersynchrony observed here, $Pb^{2+}$ exposure has been shown to increase the

likelihood of epilepsy six-fold (Chang et al., 2011; Clark et al., 2012). Epilepsy is defined by its

manifestation of neural hypersynchrony that can be either focal or global. Taking epilepsy as a model,

hypersynchrony can cause specific or general cognitive deficits corresponding to focal or global

appearances, in addition to the damaging effects of the seizure events on network architecture and

dynamics resulting from dysfunctional plasticity resulting from hypersynchrony.

Though a lesser effect, we also note that there were interesting correlations of delta parameters to

running speed in Pb2+-exposed rats when delta would normally be expected to be weak (as we previously

described, Schultheiss et al., 2020). However, in addition to typical orthogonality between delta- and theta-

dominated modes, we sometimes observed apparent correlations between frequency-bands occurring

within a given network mode (such as high delta *and* high theta). See Figure 5 insets in Schultheiss et al.,

(2020). This suggests that $Pb^{2+}$-exposure may also cause atypical cross-frequency couplings that are

detrimental to cognition and behavior.

## References

Abazyan, B., Dziedzic, J., Hua, K., Abazyan, S., Yang, C., Mori, S., Pletnikov, M. V., & Guilarte, T. R. (2014). Chronic exposure of mutant DISC1 mice to lead produces sex-dependent abnormalities consistent with schizophrenia and related mental disorders: a gene-environment interaction study. *Schizophrenia bulletin*, *40*(3), 575–584. https://doi.org/10.1093/schbul/sbt071

Amilhon, B., Huh, C. Y., Manseau, F., Ducharme, G., Nichol, H., Adamantidis, A., & Williams, S. (2015). Parvalbumin Interneurons of Hippocampus Tune Population Activity at Theta Frequency. Neuron, 86(5), 1277–1289. https://doi.org/10.1016/j.neuron.2015.05.027

Arbab, T., Battaglia, F. P., Pennartz, C., & Bosman, C. A. (2018). Abnormal hippocampal theta and gamma hypersynchrony produces network and spike timing disturbances in the Fmr1-KO mouse model of Fragile X syndrome. *Neurobiology of disease*, *114*, 65–73. https://doi.org/10.1016/j.nbd.2018.02.011

Bast, T., Zhang, W. N., & Feldon, J. (2001a). Hyperactivity, decreased startle reactivity, and disrupted prepulse inhibition following disinhibition of the rat ventral hippocampus by the GABA(A) receptor antagonist picrotoxin. *Psychopharmacology*, *156*(2-3), 225–233. https://doi.org/10.1007/s002130100775

Bast, T., Zhang, W. N., Heidbreder, C., & Feldon, J. (2001b). Hyperactivity and disruption of prepulse inhibition induced by N-methyl-D-aspartate stimulation of the ventral hippocampus and the effects of pretreatment with haloperidol and clozapine. *Neuroscience*, *103*(2), 325–335. https://doi.org/10.1016/s0306-4522(00)00589-3

Bast, T., & Feldon, J. (2003). Hippocampal modulation of sensorimotor processes. *Progress in neurobiology*, *70*(4), 319–345. https://doi.org/10.1016/s0301-0082(03)00112-6

Brandon, M. P., Bogaard, A. R., Schultheiss, N. W., & Hasselmo, M. E. (2013). Segregation of cortical head direction cell assemblies on alternating θ cycles. Nature neuroscience, 16(6), 739–748. https://doi.org/10.1038/nn.3383

Brubaker, C. J., Dietrich, K. N., Lanphear, B. P., & Cecil, K. M. (2010). The influence of age of lead exposure on adult gray matter volume. *Neurotoxicology*, *31*(3), 259–266. https://doi.org/10.1016/j.neuro.2010.03.004

Caito, S., & Aschner, M. (2017). Developmental Neurotoxicity of Lead. *Advances in*

*neurobiology*, *18*, 3–12. https://doi.org/10.1007/978-3-319-60189-2_1

Canfield, R. L., Henderson, C. R., Jr, Cory-Slechta, D. A., Cox, C., Jusko, T. A., & Lanphear, B. P.

(2003). Intellectual impairment in children with blood lead concentrations below 10 microg per deciliter. *The*

*New England journal of medicine*, *348*(16), 1517–1526. https://doi.org/10.1056/NEJMoa022848

Cao, X., Huang, S., & Ruan, D. (2008). Enriched environment restores impaired hippocampal long-

term potentiation and water maze performance induced by developmental lead exposure in

rats. *Developmental psychobiology*, *50*(3), 307–313. https://doi.org/10.1002/dev.20287

Chang, Y. T., Chen, P. C., Tsai, I. J., Sung, F. C., Chin, Z. N., Kuo, H. T., Tsai, C. H., & Chou, I. C.

(2011). Bidirectional relation between schizophrenia and epilepsy: a population-based retrospective cohort

study. *Epilepsia*, *52*(11), 2036–2042. https://doi.org/10.1111/j.1528-1167.2011.03268.x

Clarke, M. C., Tanskanen, A., Huttunen, M. O., Clancy, M., Cotter, D. R., & Cannon, M. (2012).

Evidence for shared susceptibility to epilepsy and psychosis: a population-based family study. *Biological*

*psychiatry*, *71*(9), 836–839. https://doi.org/10.1016/j.biopsych.2012.01.011

Fries, P., Reynolds, J. H., Rorie, A. E., & Desimone, R. (2001). Modulation of oscillatory neuronal

synchronization by selective visual attention. *Science* (New York, N.Y.), 291(5508), 1560–1563.

https://doi.org/10.1126/science.1055465

Goutagny, R., Jackson, J., & Williams, S. (2009). Self-generated theta oscillations in the

hippocampus. *Nature neuroscience*, *12*(12), 1491–1493. https://doi.org/10.1038/nn.2440

Guilarte, T. R., Opler, M., & Pletnikov, M. (2012). Is lead exposure in early life an environmental risk

factor for Schizophrenia? Neurobiological connections and testable hypotheses. *Neurotoxicology*, *33*(3),

560–574. https://doi.org/10.1016/j.neuro.2011.11.008

Guilarte T. R. (1997). Pb2+ inhibits NMDA receptor function at high and low affinity sites:

developmental and regional brain expression. *Neurotoxicology*, *18*(1), 43–51.

Guilarte, T. R., Miceli, R. C., Altmann, L., Weinsberg, F., Winneke, G., & Wiegand, H. (1993).

Chronic prenatal and postnatal Pb2+ exposure increases [3H]MK801 binding sites in adult rat

forebrain. *European journal of pharmacology*, *248*(3), 273–275. https://doi.org/10.1016/0926-6917(93)90054-t

Guilarte, T. R., & Miceli, R. C. (1992). Age-dependent effects of lead on [3H]MK-801 binding to the NMDA receptor-gated ionophore: in vitro and in vivo studies. *Neuroscience letters*, *148*(1-2), 27–30. https://doi.org/10.1016/0304-3940(92)90796-a

Guilarte, T. R., & McGlothan, J. L. (1998). Hippocampal NMDA receptor mRNA undergoes subunit specific changes during developmental lead exposure. *Brain research*, *790*(1-2), 98–107. https://doi.org/10.1016/s0006-8993(98)00054-7

Guilarte, T. R., Toscano, C. D., McGlothan, J. L., & Weaver, S. A. (2003). Environmental enrichment reverses cognitive and molecular deficits induced by developmental lead exposure. *Annals of neurology*, *53*(1), 50–56. https://doi.org/10.1002/ana.10399

Hashimoto, T., Volk, D. W., Eggan, S. M., Mirnics, K., Pierri, J. N., Sun, Z., Sampson, A. R., & Lewis, D. A. (2003). Gene expression deficits in a subclass of GABA neurons in the prefrontal cortex of subjects with schizophrenia. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, *23*(15), 6315–6326. https://doi.org/10.1523/JNEUROSCI.23-15-06315.2003

Jaako-Movits, K., Zharkovsky, T., Romantchik, O., Jurgenson, M., Merisalu, E., Heidmets, L. T., & Zharkovsky, A. (2005). Developmental lead exposure impairs contextual fear conditioning and reduces adult hippocampal neurogenesis in the rat brain. *International journal of developmental neuroscience : the official journal of the International Society for Developmental Neuroscience*, *23*(7), 627–635. https://doi.org/10.1016/j.ijdevneu.2005.07.005

Jett, D. A., Kuhlmann, A. C., & Guilarte, T. R. (1997). Intrahippocampal administration of lead (Pb) impairs performance of rats in the Morris water maze. *Pharmacology, biochemistry, and behavior*, *57*(1-2), 263–269. https://doi.org/10.1016/s0091-3057(96)00349-8

Korotkova, T., Fuchs, E. C., Ponomarenko, A., von Engelhardt, J., & Monyer, H. (2010). NMDA receptor ablation on parvalbumin-positive interneurons impairs hippocampal synchrony, spatial representations, and working memory. Neuron, 68(3), 557–569. https://doi.org/10.1016/j.neuron.2010.09.017

Kuhlmann, A. C., McGlothan, J. L., & Guilarte, T. R. (1997). Developmental lead exposure causes

spatial learning deficits in adult rats. *Neuroscience letters*, *233*(2-3), 101–104.

https://doi.org/10.1016/s0304-3940(97)00633-2

McGlothan, J. L., Karcz-Kubicha, M., & Guilarte, T. R. (2008). Developmental lead exposure

impairs extinction of conditioned fear in young adult rats. *Neurotoxicology*, *29*(6), 1127–1130.

https://doi.org/10.1016/j.neuro.2008.06.010

Miranda, M. L., Kim, D., Galeano, M. A., Paul, C. J., Hull, A. P., & Morgan, S. P. (2007). The

relationship between early childhood blood lead levels and performance on end-of-grade

tests. *Environmental health perspectives*, *115*(8), 1242–1247. https://doi.org/10.1289/ehp.9994

Munoz, C., Garbe, K., Lilienthal, H., & Winneke, G. (1988). Significance of hippocampal dysfunction

in low level lead exposure of rats. *Neurotoxicology and teratology*, *10*(3), 245–253.

https://doi.org/10.1016/0892-0362(88)90024-4

Needleman H. L. (1983). Lead at low dose and the behavior of children. *Acta psychiatrica*

*Scandinavica. Supplementum*, *303*, 26–37. https://doi.org/10.1111/j.1600-0447.1983.tb00939.x

Needleman, H. L., McFarland, C., Ness, R. B., Fienberg, S. E., & Tobin, M. J. (2002). Bone lead

levels in adjudicated delinquents. A case control study. *Neurotoxicology and teratology*, *24*(6), 711–717.

https://doi.org/10.1016/s0892-0362(02)00269-6

Needleman H. (2004). Lead poisoning. *Annual review of medicine*, *55*, 209–222.

https://doi.org/10.1146/annurev.med.55.091902.103653

Nihei, M. K., Desmond, N. L., McGlothan, J. L., Kuhlmann, A. C., & Guilarte, T. R. (2000). N-methyl-

D-aspartate receptor subunit changes are associated with lead-induced deficits of long-term potentiation

and spatial learning. *Neuroscience*, *99*(2), 233–242. https://doi.org/10.1016/s0306-4522(00)00192-5

Opler, M. G., Brown, A. S., Graziano, J., Desai, M., Zheng, W., Schaefer, C., Factor-Litvak, P., &

Susser, E. S. (2004). Prenatal lead exposure, delta-aminolevulinic acid, and schizophrenia. *Environmental*

*health perspectives*, *112*(5), 548–552. https://doi.org/10.1289/ehp.6777

Opler, M. G., Buka, S. L., Groeger, J., McKeague, I., Wei, C., Factor-Litvak, P., Bresnahan, M.,

Graziano, J., Goldstein, J. M., Seidman, L. J., Brown, A. S., & Susser, E. S. (2008). Prenatal exposure to

lead, delta-aminolevulinic acid, and schizophrenia: further evidence. *Environmental health*

*perspectives*, *116*(11), 1586–1590. https://doi.org/10.1289/ehp.10464

608             Paxinos, G., and Watson, C. (2004). The rat brain in stereotaxic coordinates 5<sup>th</sup> Edition.

609     (Massachusetts: Academic Press).

610             Schultheiss, N.W., Schlecht, M., Jayachandran, M., Brooks, D.R., McGlothan, J.L., Guilarte, T.R. &

611     Allen, T.A. (2020). Awake delta and theta-rhythmic hippocampal network modes during intermittent

612     locomotion behaviors in the rat. *Behavioral Neuroscience.*

613             Stansfield, K. H., Ruby, K. N., Soares, B. D., McGlothan, J. L., Liu, X., & Guilarte, T. R. (2015).

614     Early-life lead exposure recapitulates the selective loss of parvalbumin-positive GABAergic interneurons

615     and subcortical dopamine system hyperactivity present in schizophrenia. *Translational psychiatry*, *5*(3),

616     e522. https://doi.org/10.1038/tp.2014.147

617             Stark, E., Eichler, R., Roux, L., Fujisawa, S., Rotstein, H. G., & Buzsáki, G. (2013). Inhibition-

618     induced theta resonance in cortical circuits. *Neuron*, *80*(5), 1263–1276.

619     https://doi.org/10.1016/j.neuron.2013.09.033

620             Wang, T., Guan, R. L., Liu, M. C., Shen, X. F., Chen, J. Y., Zhao, M. G., & Luo, W. J. (2016). Lead

621     Exposure Impairs Hippocampus Related Learning and Memory by Altering Synaptic Plasticity and

622     Morphology During Juvenile Period. *Molecular neurobiology*, *53*(6), 3740–3752.

623     https://doi.org/10.1007/s12035-015-9312-1

624             Yamamoto J, Suh J, Takeuchi D, Tonegawa S. Successful execution of working memory linked to

625     synchronized high-frequency gamma oscillations. Cell. 2014;157(4):845-857.

626     https://doi:10.1016/j.cell.2014.04.009

**Figure 1. Movement speed modulation of delta, theta, and gamma oscillations in the hippocampal local field potential (LFP). A.** Eight recording sites (one tetrode each) located at two depths on each of four shanks of a silicon probe. Tracks left by the shanks of the implanted probe are visible as disruptions of the cell layer. **B.** Representative LFP traces recorded while an animal freely explored the behavioral arena (top) and the corresponding movement speed of the animal (bottom). **C.** Movement speed modulation of the amplitudes of delta, theta, and gamma oscillations (across all behaviors) in control rats and rats chronically exposed to lead ($Pb^{2+}$) in their diet.

**Figure 2. Power spectra during stationary and running bouts. A.** Diminished delta and heightened
theta during running (**right**) relative to stationary bouts (**left**) for two control rats. **B.** As in A for two Pb$^{2+}$-
exposed rats. **C.** An apparent proximal-distal gradient in delta power across probe shanks in stationary
control animals (**left**), is absent in Pb$^{2+}$-exposed animals (**right**). Black and red lines in C and D reflect the
best fits to power estimates along the dorsal and ventral rows of four tetrodes (across shanks) for each rat.
**D.** Theta power at sites along the proximal-distal axis in control and Pb$^{2+}$-exposed animals during running
bouts.

**Figure 3. Theta-frequency hypersynchrony in Pb$^{2+}$-exposed rats. A.** Elevated theta-frequency power in
Pb$^{2+}$ rats during stationary bouts (2 sec bouts at **top**, 1 sec bouts **below**). **B.** Marked increase in theta
power in Pb$^{2+}$-exposed rats during running bouts.

**Figure 4. Pb²⁺ exposure reduces theta frequency and disrupts running speed modulation. A.** Scatter plots of theta power and frequency taken from the power spectral density for each recorded channel during each bout of stationary (**left**) and running behaviors (**right**) for all rats. Black and green lines indicate means for control and Pb²⁺-exposed rats, respectively. Dashed lines indicate the mean theta frequencies for each group. **B.** Running speed modulation of theta frequency. **C.** Running speed modulation of theta power.

661
**Figure 5. Running speed modulation of theta power is elevated in Pb$^{2+}$-exposed animals. A.**
Distributions of correlation coefficients between delta (blue) and theta (red) power estimates and running
speed. Individual correlations were calculated across running bouts (1 sec) for each recorded channel from
control rats (**top**) and Pb$^{2+}$-exposed rats (**bottom**). **B.** As in A, for two second running bouts. **C.** As in A, but
comparing correlations of delta and theta peak frequencies to running speed.
667

**Figure 6. Pb²⁺-exposure elevates behavioral modulation of theta power. A.** Change in power spectra between stationary and running bouts (**top**) for control (black) and Pb²⁺-exposed rats (red) normalized by total power (0-25 Hz). Enlargement of the low frequency range of PSDs is shown below. Scatters of individual measures for each channel and rat and using one and two second bout definitions (**bottom**). **B.** Percent change in power at each frequency between stationary and running bouts (**top**) for control (black) and Pb²⁺-exposed rats (red). PSD enlargement (**middle**) and scatters (**bottom**) as in A.

**Figure 7. Pb²⁺ exposure amplifies theta and gamma-frequency synchrony. A.** Distributions of normalized cycle amplitudes for delta, theta, and gamma frequency bands during stationary (blue) and running bouts (green) in control (**top**) and Pb²⁺-exposed animals (**bottom**). **B.** Distributions of unnormalized cycle amplitudes for each frequency band during stationary (**top**) and running bouts (**bottom**) in control (black) and Pb²⁺-exposed animals (red).

683
**Figure 8. Prepulse inhibition in male and female rats chronically exposed to Pb$^{2+}$.** The results show
no effect of Pb$^{2+}$ exposure at PN28 but marked inhibition of PPI in male rats at PN50 and PN120. No effect
of Pb$^{2+}$ exposure was found in female rats at any age. Insert graphs in each panel are startle response at
120 dB. No effect on startle response was observed in Pb$^{2+}$-exposed male or female rats relative to
controls at any age. Thus, CELLE does not alter the sensitivity of the animal to a startle. However, CELLE
produces a significant deficit on PPI in male rats relative to controls (*p<0.05). Data are expressed as
mean ± sem. n=number of litters (one male and female animal per litter).
691
692

| AGE | PN28 | | PN50 | | PN120 | |
|---|---|---|---|---|---|---|
| SEX | Males | Females | Males | Females | Males | Females |
| Control | ≤1.9 n=18 | ≤1.9 n=16 | ≤1.9 n=17 | ≤1.9 n=15 | ≤1.9 n=17 | ≤1.9 n=15 |
| 1500ppm PbAc | 21.1 ± 0.9 n=18 | 20.9 ± 1.1 n=16 | 20.2 ± 1.1 n=17 | 22.1 ± 1.3 n=20 | 19.6 ± 1.3 n=17 | 24.3 ± 2.2 n=14 |
| p-value | ≤0.0001 | ≤0.0001 | ≤0.0001 | ≤0.0001 | ≤0.0001 | ≤0.0001 |

695 **Table 1:** Blood $Pb^{2+}$ levels (µg/dL) in male and female rats contributing to data represented in Figure 9. Each
696 value is the mean ± sem. n= number of litters (one male and female animal per litter).
697

**Supplemental Figure 1. Precise delineation of stationary and running bouts. A.** Schematic of the orientation measures used in definitions of behavioral bouts. The star indicates the rat's location in B when the image was captured. **B.** Representative locomotor trajectories showing thigmotaxis during the first 40 seconds of an experimental session (**left**) followed by runs through the interior of the open field arena during the subsequent epoch (**right**). Arrows indicate the rat's direction of running. As in A, blue and green points represent head-mounted LEDs used to derive orientation measures, and black line segments capture the rat's body orientation at each timepoint. **C.** Head and body direction (**top**) were used to derive rotational velocity and head angle (**middle**), which in turn were used to identify bouts of running (**bottom**) while excluding rotational and reorienting behaviors (red points and circles). **D.** Speed occupancy (**left**) and rotational velocity (**right**) measured throughout behavioral sessions did not differ between control and $Pb^{2+}$-exposed rats, nor did bout-wise behavioral measures (**F**) of running (**left**) and stationary bouts (**right**), including the numbers of bouts, median movement speeds, and bout durations.