1

Mendelian randomization analyses support causal relationships between

blood metabolites and the gut microbiome

Xiaomin Liu[1,2,8], Xin Tong[1,8], Yuanqiang Zou[1], Xiaoqian Lin[1,2], Hui Zhao[1,2], Liu Tian[1],

Zhuye Jie[1], Qi Wang[1,2], Zhe Zhang[1], Haorong Lu[3], Liang Xiao[1,4,5], Xuemei Qiu[1], Jin Zi[1],

Rong Wang[1], Xun Xu[1], Huanming Yang[1,6], Jian Wang[1,6], Yang Zong[1], Weibin Liu[1],

Yong Hou[1], Shida Zhu[1], Huijue Jia[1,7*], and Tao Zhang[1*]

[1]BGI-Shenzhen, Shenzhen, China.

[2]College of Life Sciences, University of Chinese Academy of Sciences, Beijing

100049, China;

[3]China National Genebank, BGI-Shenzhen, Shenzhen, China.

[4]Shenzhen Engineering Laboratory of Detection and Intervention of Human Intestinal

Microbiome, BGI-Shenzhen, Shenzhen, China.

[5]BGI-Qingdao, BGI-Shenzhen, Qingdao, China.

[6]James D. Watson Institute of Genome Sciences, Hangzhou, China.

[7]Shenzhen Key Laboratory of Human Commensal Microorganisms and Health

Research, BGI-Shenzhen, Shenzhen, China.

[8]These authors contributed equally to this work.

*e-mail: tao.zhang@genomics.cn, jiahuijue@genomics.cn

The gut microbiome has been implicated in a variety of physiological states, but controversy over causality remains unresolved. Here, we performed bidirectional Mendelian randomization (MR) analyses on 3,432 Chinese individuals with whole genome, whole metagenome, anthropometric, and blood metabolic trait data. We identified 58 causal relationships between the gut microbiome and blood metabolites, and replicated 43 of them. Increased relative abundances of fecal *Oscillibacter* and *Alistipes* were causally linked to decreased triglyceride concentration. Conversely, blood metabolites such as glutamic acid appeared to decrease fecal *Oxalobacter,* and members of *Proteobacteria* were influenced by metabolites such as 5-methyltetrahydrofolic acid, alanine, glutamate, and selenium. Two-sample MR with data from Biobank Japan partly corroborated results with triglyceride and with uric acid, and also provided causal support for published fecal bacterial markers for cancer and cardiovascular diseases. This study illustrates the value of human genetic information to help prioritize gut microbial features for mechanistic and clinical studies.

Metagenome-wide association studies (MWAS) using human stool samples, as well as animal models, especially germ-free mice, have pointed to a potential role of the gut microbiome in diseases such as cardiometabolic, autoimmune, neuropsychiatric disorders and cancer, with mechanistic investigations for diseases such as obesity, colorectal cancer and schizophrenia[1-4]. Twin-based heritability estimation and more recent metagenome-genome-wide association studies (M-GWAS) have questioned the traditional view of the gut microbiota as a purely environmental factor[5-9], although the extent of the genetic influence remains controversial[7,10]. Yet, all these published cohorts, except for human sequences in the metagenomic data of HMP (Human Microbiome Project), utilized array data for human genetics, and most of them had 16S rRNA gene amplicon sequencing for the fecal microbiota[5-9].

As the gut microbiome is considered to be highly dynamic, causality has been an unresolved issue in the field. Mendelian randomization (MR)[11] offers an opportunity to distinguish between causal and non-causal effects from cross-sectional data, without animal studies or randomized controlled trials. An early study used MR to look at the gut microbiota and ischemic heart disease[12]. Recently, a study used MR to confirm that increased relative abundance of bacteria producing the fecal volatile short-chain fatty acid (SCFA) butyrate was causally linked to improved insulin response to oral glucose challenge; in contrast, another fecal SCFA, propionate, was causally related to an increased risk of T2D[13]. However, both studies used genotype data, and it was not clear to what extent the genetic factors explained the microbial feature of interest.

In this study, we present a large-scale M-GWAS using whole genome and fecal microbiome, followed by bidirectional MR for the fecal microbiome and anthropometric features as well as blood metabolites. In a two-stage design from different cities in China, we identified 58 causal links from MR in the 4D-SZ discovery cohort of 2,002 individuals with high-depth whole-genome sequencing data (1,539 individuals with microbiome data for one-sample MR). We replicated 43 of the 58 causal effects using

low-depth whole-genome sequencing data of another 1,430 individuals (1,006

individuals with microbiome data for one-sample MR). In general, unidirectional

causal effects could be found both from the gut to the blood and from the blood to the

gut, but bidirectional effects were rarely detected. A few of the M-GWAS associations

with gut microbial functional modules, e.g. module for lactose/galactose degradation

and the *ABO* loci, reached study-wide significance, illustrating the power of shotgun

metagenomic data together with whole genome sequencing. The MR findings were

corroborated and extended by summary statistics from the Biobank Japan study, e.g.

causal effect of *Proteobacteria* on type 2 diabetes (T2D), congestive heart disease

and colorectal cancer, underscoring the significance of human genetic data to help

guide microbiome intervention studies.

Results

Fecal microbiome features associated with human genetic variants. We set out

to identify human genetic variants to be included as the randomizing layer of MR (**Fig.**

1). The 4D-SZ (multi-omics, with more time points to come, from Shenzhen, China)

discovery cohort consisted of high-depth whole-genome sequencing data from 2,002

blood samples (mean depth of 42×; **Supplementary Table 1** and **Supplementary**

Fig. 1a), out of which 1,539 individuals had metagenomic shotgun sequencing data

from stool samples (8.56 ± 2.28 Gb; **Supplementary Fig. 1b**). Fecal M-GWAS was

performed using 10 million common and low-frequency variants (minor allele

frequency (MAF) ≥ 0.5%) and 500 unique microbial features (Spearman's correlation

< 0.99; **Methods**). The M-GWAS was adjusted for age, gender, BMI, defecation

frequency, stool form, self-reported diet, lifestyle factors, and the first four principal

components from the genomic data to account for population stratification.

With this large cohort of whole genome and whole metagenome data, we

performed M-GWAS analysis and identified a total of 625 associations involving 548

independent loci for one or more of the 500 microbial features at genome-wide

significance ($P < 5 \times 10^{-8}$). With a more conservative Bonferroni-corrected study-wide

significant P value of 1.0×10^{-10} ($= 5 \times 10^{-8}/500$), we identified 28 associations with

fecal microbial features involving 27 genomic loci, of which 5 correlated with gut

bacteria and the other 22 associated with gut metabolic pathways (**Supplementary**

Table 2).

For MR, it was important for the genetic variants used to be representative of the

microbiome features (**Supplementary Fig. 2**), so a more suggestive P value of less

than 1×10^{-5} was used (**Supplementary Table 2**), as in previous MR studies[13,14].

Each microbial feature had an average of 44 genetic variants (**Fig. 2a** and

Supplementary Table 3). The corresponding genetic variants explained microbial

features to a median value of 24.9%, e.g. 45.5% of the microbial metabolic pathway

for succinate consumption and 44.6% of *Phascolarctobacterium succinatutens* (an

asaccharolytic, succinate-utilizing bacterium), while only 6.8% of genus *Edwardsiella*

(**Supplementary Table 3**). The phenotypic (relative abundance) variance of five

genera (*Bilophila*, *Oscillibacter*, *Faecalibacterium*, *Megasphaera* and *Bacteroides*)
could be explained over 35% by their corresponding independent genetic variants
(**Fig. 2b**). Thus, although human genetic associations (array data) have been
reported to explain only 10% or 1.9% of the gut microbiota[7,10], the suggestive
associations from the current M-GWAS study could be highly predictive of certain gut
taxa and functions.
For better confidence in these suggestive associations, we sequenced a
replication cohort of 1,430 individuals from multiple cities in China (also shotgun
metagenomic sequencing for stool samples, but about 8× whole-genome sequencing
for human genome; **Supplementary Fig. 1c,d**). Among the 22,293 independent
associations identified in the discovery cohort with $P < 10^{-5}$, 4,876 variants were not
available in the low-depth replication dataset and 87.6% of them were not common
variants (MAF < 0.05). We were able to replicate 2,324 of the remaining 17,417
independent associations in the same effect direction of minor allele ($P < 0.05$,
Supplementary Table 2), indicating that the associations were not random false
positives. The fraction of associations replicated in the same direction ($P < 0.05$) using
the suggestive cut-off of $P < 10^{-5}$ (2,324/17,417) was not lower than the more stringent
cut-offs (54/625 of the $P < 5 \times 10^{-8}$, and 2/28 of the $P < 10^{-10}$). Two well-replicated
signals from the study-wide threshold were rs1461780285 in the *ABO* blood group
associated with module MF0007: lactose and galactose degradation ($P_{discovery}$ = 2.10 ×
10^{-12} and $P_{replication}$ = 1.09 × 10^{-10}; **Supplementary Fig. 3a,b**) and rs142693490 near
the *LCORL* gene (implicated in spermatogenesis, body frame and height) associated
with MF0034: alanine degradation II ($P_{discovery}$ = 1.28 × 10^{-12} and $P_{replication}$ = 0.014;
Supplementary Fig. 3c,d). Genetic variant rs1461780285 is in strong linkage
disequilibrium (LD, r^2 = 0.99) with multiple SNPs (rs507666, rs532436, rs651007,
rs579459 and rs579459) in the *ABO* gene. These SNPs located in a block were
found to be associated with metabolites levels in both this study and previous studies,
especially for serum alkaline phosphatase levels (**Supplementary Table 4**). In
addition, we were able to replicate several previously reported microbial signals[6-10],
especially for rs12354611 and *Bacteroides stercoris* (P = 8.64 × 10^{-6}; **Supplementary**
Table 5 and **Supplementary Note**).
141
Blood metabolic traits associated with human genetic variants. On the other
hand, plasma metabolite levels are also known to associate with host genetic variants
(**Fig. 1**). We thus performed whole genome-wide association tests for each of the 112
metabolites, with log-transformed relative concentration. We identified a total of 174
associations involving 158 loci that independently associated with one or more of the
112 metabolites at genome-wide significance ($P < 5 \times 10^{-8}$). With a more conservative
Bonferroni-corrected study-wide significant P value of 4.5 × 10^{-10} (= 5 × 10^{-8}/112
metabolites), we identified 39 associations with metabolites involving 28 genomic loci
(**Supplementary Table 6**). These included previously well-established associations
such as the *UGT1A* family associated with serum total bilirubin[15,16] and *ASPG*
associated with asparaginate[15].
According to the suggestive threshold of $P < 10^{-5}$, we identified 6,541 metabolic

quantitative trait loci (mQTLs), of which 361 were associated with two or more
metabolites (**Supplementary Table 6**). The average number of genetic variants was
58 for each metabolic trait (**Fig. 3a** and **Supplementary Table 7**). The percentage of
variance explained by the corresponding genetic variants ranged from 13.3% (red
blood cell distribution) to as high as 48.3% (blood mercury concentration) and 45.9%
(blood alpha-fetoprotein value), with a median value of 28.6% (**Fig. 3b**). Among these,
268 variants or their proxy variants ($r^2 > 0.6$; distance < 1 Mb) have been reported in
the GWAS catalog[17] (**Supplementary Table 8**). Some variants were associated with
diseases in the GWAS catalog such as chronic kidney disease, Alzheimer's disease,
coronary artery disease, Crohn's disease, ovarian cancer, breast cancer and gastric
cancer.
Among the 6,541 suggestive mQTLs identified in the 4D-SZ discovery cohort with
$P < 10^{-5}$, 5,088 variants were covered by the replication dataset. 717 and 31 were
replicated at nominal ($P < 0.05$) and suggestive significance ($P < 10^{-5}$), respectively, in
the same effect direction of minor allele (**Supplementary Table 6**). Especially for the
174 genome-wide and 39 study-wide significant associations, we could replicate 51
and 29 associations in the same direction ($P < 0.05$), respectively. The top
associations confirmed by the low-depth genomes ($P < 4.5 \times 10^{-10}$ both in discovery
and replication cohorts) included: *FECH* associated with manganese; *UGT1A* family
associated with serum total bilirubin as well as direct and indirect (unconjugated)
bilirubin; *ASPG* associated with asparagine; *CPS1* associated with glycine; and
APOE associated with low density lipoprotein (**Supplementary Fig. 4**). Overall, the
accurate identification of genetic determinants and the high variance explained for
both microbial features and blood metabolites are optimal for MR analysis to
investigate causality.
179
From observational correlation to Mendelian randomization. As a prerequisite for
strong causality, we investigated the correlation between relative abundances of 500
unique fecal microbial features (taxa and functional modules) and 112 host metabolic
traits using multivariate linear regression. After adjustment for gender and age, we
observed 457 significant associations (false discovery rate (FDR) corrected $P < 0.05$;
Supplementary Table 9 and **Online Methods**). Three metabolites, glutamic acid,
5-methyltetrahydrofolic acid (5-methyl THF, active form of folic acid) and selenium,
were associated with the largest number of microbial features (58, 40 and 38,
respectively; **Supplementary Fig. 5**). These associations extend findings from
various studies and suggest quantitative relationships between gut microbial
taxa/functions and plasma metabolites.
As we were fortunate to have all the data in the same cohort, we first performed
one-sample MR analyses to identify causal relationships for the 457 observational
correlations in the discovery cohort consisting of 1,539 individuals with both metabolic
and microbiome traits. We found 58 significant causal effects, of which 17 showed
causal effects for gut microbial features on blood metabolic traits and the other 41
showed causal effects for blood metabolic traits on gut microbial features ($P < 1.09 \times$
$10^{-4} = 0.05/457$; **Fig. 4** and **Supplementary Table 10**). Only 4 of these were

bidirectional. By applying one-sample MR analyses to the replication dataset of 1,006
low-depth genomes as well as metabolic and microbiome traits from individuals in
different cities, we could replicate 43 of the 58 causal relationships (in the same
direction and $P < 0.05$; **Supplementary Table 10**), indicating that the effects were not
random false positives.
Moreover, we also used six different two-sample MR methods, which are more
commonly performed when only summary statistics are available from two different
cohorts, to analyze our data both in the discovery cohort (summary data for 2,002
samples with metabolic traits and 1,539 samples with microbial features) and the
replication cohort (summary data for 1,430 samples with metabolic traits and 1,006
samples with microbial features). The one-sample MR and the two-sample MR
analyses showed highly consistent results, and the Spearman's correlation for beta
coefficients between one-sample and two-sample MR reached 0.767 for the discovery
cohort ($P < 2.2 \times 10^{-16}$). The 58 causal associations identified by one-sample MR were
also significant in the two-sample MR analyses. An additional 14 causal associations
were identified by the two-sample MR analyses (**Supplementary Table 11**), possibly
due to the larger cohort size. We also examined the presence of horizontal pleiotropy
by using the MR-PRESSO Global test[18]. Only one causal association (the negative
effect of selenium on the abundance of *Methylobacillus flagellates*, $P_{\text{MR-PRESSOGlobaltest}} =$
0.01; **Supplementary Table 9**) showed pleiotropy, while all the other 71 causal
relationships showed no evidence of pleiotropy ($P > 0.05$). Thus, our MR analyses
identified robust causal relationships between blood metabolic traits and specific
features of the gut microbiome.
221
Effects of the gut microbiome on blood metabolic traits. As some of the
MR-identified relationships appeared linked, we performed hierarchical clustering for
the 12 microbial features and 8 blood metabolites involved in the 17 causal
relationships from the gut microbiome to blood metabolites, which formed two clusters.
One cluster involved decreasing the plasma levels of triglyceride and alanine by gut
microbial taxa or functional modules, and the other involved decreasing the levels of
5-methyltetrahydrofolic acid or progesterone, but increasing serum uric acid or
plasma glutamic acid by gut microbial features (**Fig. 4a**). Reassuringly, the species
Mobiluncus curtisii was clustered next to its corresponding genus *Mobiluncus*, and
modules including serine degradation and threonine degradation, sucrose
degradation and pectin degradation, were likewise next to one another.
The most significant causal effect was *Oscillibacter* on decreasing blood
triglyceride concentration (**Fig. 5a-c**), and to a lesser extent on lowering body-mass
index (BMI) and waist-hip ratio (WHR), whereas the effect with plasma alanine was
bidirectional. Using 134 genetic variants to construct a polygenic risk score (PRS)
(134 genetic variants and the constructed PRS explained 39.3% and 49.6% of the
phenotypic variance, respectively, **Fig. 3b** and **Supplementary Table 10**) for
one-sample MR analysis in the discovery cohort, we estimated that each 1-s.d.
increase in the abundance of *Oscillibacter* would generate a 0.261 mmol/L decrease
in triglyceride concentration ($P = 2.53 \times 10^{-10}$), a 0.161 kg/m² decrease in BMI ($P =$

1.33×10^{-4}) and 0.126 ratio decrease in WHR ($P = 2.73 \times 10^{-3}$). This causal

relationship was robust when four two-sample MR tests were performed ($P_{\text{GCTA-GSMR}} =$

4.34×10^{-11}, $P_{\text{Inverse\_variance\_weighted}} = 2.45 \times 10^{-15}$, $P_{\text{weighted-median}} = 1.22 \times 10^{-7}$ and $P_{\text{MR-Egger}}$

$= 1.35 \times 10^{-5}$) (**Fig. 5c**), and there was no evidence of horizontal pleiotropy

($P_{\text{MR-PRESSOGlobaltest}} = 0.18$; **Supplementary Table 11**). The reverse MR analysis

(testing the effect of genetic predictors of triglyceride on *Oscillibacter* abundance) was

significant but did not reach the multiple test corrected significance ($10^{-4} < P < 0.05$).

Oscilibacter is a Gram-negative Clostridial bacterium, phylogenetically close to

Oscillospira[19], which could produce valerate or butyrate. In addition, higher relative

abundance of *Alistipes* was also associated with decreased blood triglyceride

concentration ($P = 8.31 \times 10^{-8}$, **Fig. 5d**). At the species level, both *A. shahii* ($P = 1.37 \times$

10^{-6}) and *unclassified Alistipes sp. HGB5* ($P = 3.36 \times 10^{-5}$) showed negative effects on

blood triglyceride. The effects of both *Oscilibacter* and *Alistipes* for lowering blood

triglyceride concentration were confirmed in the replication cohort ($P = 3.39 \times 10^{-4}$ and

$P = 2.88 \times 10^{-4}$, respectively; **Supplementary Tables 10** and **11**). These findings

support the decrease in relative abundances of *Oscillibacter* and *Alistipes* in obese

individuals compared to individuals with normal BMI reported in previous studies[20-22],

suggesting that these bacteria are promising supplementation agents for individuals

of a suitable genetic background.

The gut microbiome potential for pectin degradation II (42.6% of the variance

explained by GRS) showed a handful of significant MR hits with blood traits (**Fig. 4a**),

including positive effects on alanine ($P = 8.57 \times 10^{-5}$) and serum uric acid ($P = 1.34 \times$

10^{-6}), and negative effects on progesterone ($P = 6.68 \times 10^{-7}$). *Bacteroidetes* and

Fusobacteria were the only two phyla that positively correlated with the abundance of

pectin degradation II (Spearman rank correlation, $\rho = 0.48$ and 0.15, respectively),

which included the two previously reported pectin-degrading species *Bacteroides*

thetaiotaomicron and *Fusobacterium varium*[23,24]. In the 4D-SZ cohort, *F. varium*

correlated with pectin degradation II (Spearman's correlation, $\rho = 0.12$) and increased

blood alanine ($P = 0.02$) and serum uric acid ($P = 0.04$); *B. thetaiotaomicron*

correlated with pectin degradation II (Spearman's correlation, $\rho = 0.21$) but showed no

detectable effect on alanine or uric acid ($P > 0.05$; **Supplementary Fig. 7a,b,d**).

Instead, *B. dorei,* the bacterial species most strongly correlated with pectin

degradation II (Spearman rank correlation, $\rho = 0.32$, **Supplementary Fig. 7c**),

positively contributed to alanine ($P = 0.05$) and serum uric acid levels ($P = 3.40 \times 10^{-4}$;

Supplementary Fig. 7d).

Effects of blood metabolites on gut microbial features. For the 41 causal

relationships from blood metabolic traits to gut microbial features (one-sample MR,

Supplementary Table 10), hierarchical clustering revealed two clusters, one mostly

involved decreasing abundance of bacteria by plasma alanine or glutamic acid, and

the other involved decreasing abundance of bacteria by selenium or

5-methyltetrahydrofolic acid (**Fig. 4b**). *F. prausnitzii* showed a negative effect on

plasma selenium (**Fig. 4a**), while plasma selenium showed negative effects on gut

Proteobacteria such as Enterobacteriaceae (e.g. *Escherichia coli*, $P = 3.79 \times 10^{-5}$),

Pseudomonas stutzeri (P = 1.06 × 10^{-6}), and modules such as arginine degradation II
(P = 2.65 × 10^{-6}), succinate conversion to propionate (P = 3.55 × 10^{-5}), and anaerobic
fatty acid beta oxidation (P = 9.71 × 10^{-5}) (**Fig. 4b**).
Bacteria from the phylum *Proteobacteria* were negatively affected by not only
selenium but also 5-methyltetrahydrofolic acid (**Fig. 4b**). We directly verified the effect
of 5-methyltetrahydrofolic acid on *Escherichia in vitro*. Supplementing
5-methyltetrahydrofolic acid in growth media indeed slowed down the growth of a
strain of *Escherichia coli* AM17-9 compared to lower concentrations or absence of
5-methyltetrahydrofolic acid (**Supplementary Fig. 8**).
A handful of bacteria were also affected by glutamic acid. The negative influence
of glutamic acid (48 variants with suggestive associations and the constructed PRS
explained 24.9% and 25.4% of the phenotypic variance, respectively) on the genus
Oxalobacter (P = 1.56 × 10^{-6}) may help explain the lower prevalence of *Oxalobacter* in
developed countries, besides the lower intake of oxalate and antibiotic use[25]. Whether
limiting glutamic acid could raise *Oxalobacter* and prevent kidney stones remains to
be tested. Glutamic acid negatively affected melibiose degradation (to glucose,
galactose, P = 2.05 × 10^{-5} from two-sample MR), but showed positive effects on
alanine degradation I (P = 5.46 × 10^{-5}), anaerobic fatty acid beta-oxidation (P = 9.36 ×
10^{-5}), and bidirectional positive effect on serine degradation (P = 6.85 × 10^{-7} for serine
degradation to glutamic acid and P = 9.90 × 10^{-6} for glutamic acid to serine
degradation, respectively).
307
Causal relationships between the gut microbiome and diseases. We further
investigated the effects of the 72 significant causal relationships (**Supplementary
Table 11**) involving 40 microbial features and 12 metabolic traits on diseases by
performing two-sample MR analysis using gut microbiome GWAS summary data in
this 4D-SZ cohort, together with blood quantitative traits and diseases GWAS
summary statistics from Biobank Japan[26] (**Fig. 1** and **Supplementary Table 12**),
given that Japanese people have a genetic architecture similar to Chinese. Only
routine blood parameters but no amino acids, hormones or microelements were
included in the Biobank Japan study. Thus, only five of the 72 causal associations,
involving triglyceride and serum uric acid were available for further investigation in the
Biobank Japan data. The relationship between unclassified Lachnospiraceae
bacterium 9_1_43BFAA and uric acid was reciprocal in the 4D-SZ cohort, and we
could replicate the causal effect of uric acid on increased unclassified
Lachnospiraceae bacterium 9_1_43BFAA abundance in the Japanese cohort,
whereas the reciprocal effect, i.e. potential effect of unclassified Lachnospiraceae
bacterium 9_1_43BFAA on uric acid was not replicated, possibly due to lack of
variants in the genotyped Japanese cohort (15 instead of 67, **Supplementary Table
13**). The other three associations were not replicated, possibly for the same reason.
For example, genus *Oscillibacter* had 135 variants with P < 10^{-5} in our summary data
but only 15 were available in the Biobank Japan summary data.
MR inference using our gut microbiome M-GWAS summary data and disease
GWAS summary statistics from Biobank Japan found that *Alistipes*, which showed

negative effects on blood triglyceride in the 4D-SZ cohort, lowered the risks of
cerebral aneurysm (**Supplementary Table 14**; $P = 4.61 \times 10^{-4}$) and hepatocellular
carcinoma ($P = 0.045$) in the Biobank Japan cohort. According to the genetic
associations we identified for *Proteobacteria*, we were able to see in Biobank Japan
disease data that *Proteobacteria* increased the risk of T2D (**Fig. 6a**; $P = 7.61 \times 10^{-4}$,
two-sample MR), congestive heart failure ($P = 0.003$) and colorectal cancer ($P =
0.047$). This is consistent with MWAS findings mainly for Enterobacteriaceae[1] and
suggest that the metabolites identified above (5-methyltetrahydrofolic acid, selenium)
might help prevent the diseases. Folic acid is indeed recommended for heart
diseases[27]. In addition, *Escherichia coli* increased the risk of urolithiasis (**Fig. 6b**; $P =
0.009$) and hepatocellular carcinoma ($P = 0.04$) but decreased interstitial lung disease
risk ($P = 0.007$). Similarly, *Salmonella enterica* increased prostate cancer risk but
decreased interstitial lung disease risk. The Pseudomonadales order was the only
microbial feature showing a positive effect on pulmonary tuberculosis. The denitrifying
bacteria *Achromobacter* increased the risk of atopic dermatitis ($P = 0.005$), gastric
cancer ($P = 0.008$), esophageal cancer ($P = 0.027$) and biliary tract cancer ($P = 0.034$).
Bacteroides intestinalis, which was reported to be relatively depleted in patients of
atherosclerotic cardiovascular disease[28], was found here to increase with potassium,
and *B. intestinalis* showed a negative effect on epilepsy. *Streptococcus parasanguinis*
had a positive effect on colorectal cancer and posterior wall thickness
(echocardiography), consistent with MWAS studies[1,28,29]. These results illustrated the
potential significance of the gut microbiome-blood metabolite relationships in
understanding and preventing cardiometabolic diseases and cancer.
353
354
355   **Discussion**
356
In summary, we identified many genetic loci associated with microbial features and
metabolic traits, and found 58 causal relationships between the gut microbiome and
blood metabolites using one-sample bidirectional MR. 43 out of the 58 one-sample
MR signals could be replicated in a low-depth genome sequenced cohort also from
China. Two-sample MR replicated the 58 causal relationships in the same direction
and identified an additional 14 causal relationships. Two-sample MR using summary
statistics from Biobank Japan identified effects of gut microbial features on diseases,
suggesting potential applications of microbiome intervention in cardiometabolic,
kidney and lung diseases, and cancer. While mechanistic investigations using
germ-free mice and reference bacterial strains have been popular, our data-driven
analyses underscore the clinical relevance of gut microbes that have not been
extensively cultured and characterized, e.g. *Oscilibacter* and *Alistipes* for lowering
triglyceride concentration and a number of disease risks, which may be particularly
relevant for East Asian regions undergoing rapid changes in lifestyle and disease
profiles.
Although associations between the gut microbiome and blood features such as
amino acids and vitamins have been known for some time, our MR analyses could

inspire more mechanistic and interventional studies. The unique data available from
the 4D-SZ cohort allowed appreciation of overlooked features such as selenium.
Interestingly, while selenium showed adverse effects on Gammaproteobacteria, *F.*
prausnitzii showed a negative effect on plasma selenium. Nitrogen is a limiting
resource for many ecosystems. In the modern human gut microbiome without high
intake of nitrite, proteins are probably the major source of nitrogen[30], and the
glutamate-glutamine reservoir is a key buffering mechanism for the inflammatory
potential of excess amines[31-34]. The increase in *Proteobacteria* and decrease in
Oxalobacteraceae observed in these Chinese individuals no more than 30 years old
on average could potentially explain susceptibility to cardiometabolic and kidney
diseases later in life. The bidirectional link between strontium and *Streptococcus*
parasanguinis implies an interplay between water source and cardiovascular
diseases[28,35].
Metabolism of polysaccharides that cannot be directly digested by the host is an
important function of the colonic microbiome. We found degradation of pectin (or
sucrose) to negatively affect progesterone level. This is an interesting possibility to
provide scientific support for traditional dietary advice for pregnant women to ensure
full-term pregnancy. Hyperuricemia and gout is a growing epidemic in East Asia, and
soft drinks containing fructose are a strong factor that is no less important than beer
and meat[36]. Gut microbial (*Bacteroides*, *Fusobacterium*) pectin degradation module
positively contributed to circulating levels of alanine and uric acid. Further studies on
the trans-kingdom metabolic flux of carbon and nitrogen would be necessary for
personalized management of uric acid and alanine levels.
For the nascent field of M-GWAS and microbiome MR, there are also a lot of
opportunities for methodological development by statistical experts. Low-frequency
microbes are common in an individual's gut and could play physiological or
pathological roles[1,37]. Our MR results for gut microbial species were supported by MR
for higher taxonomic units such as genus or phylum (**Fig. 4** and **Supplementary**
Table 11). Yet, the *P* values were sometimes more significant for the larger taxa,
suggesting similar functions contributed by other species. Functional redundancy in
the microbiome has been discussed ever since the beginning of the microbiome
field[38,39], and here we identified study-wide significant host genetic associations with
gut microbial functional modules, and causal effects of other gut microbial functional
modules on host levels of circulating metabolites. Distribution of the microbiome
taxonomic or functional data constitutes another layer of consideration, in addition to
the human allele frequencies. Gathering a more homogenous cohort could enable
identification of signals in a relatively small cohort, while corrections for comparing
different populations might involve host-microbiome interactions. As the gut
microbiome can be influenced by medication[40], and heritability for most traits is higher
in younger individuals[41], healthy young adults are probably preferable for M-GWAS
studies, while microbiome-drug interactions in older individuals could be an important
direction for MR studies.
Overall, our data-driven approach underscores the great potential of M-GWAS
and MR for a full picture of the microbiome, which can be mechanistically illuminating

and are poised to help focus intervention efforts to mitigate inflammation and prevent
or alleviate complex diseases.
420
421

Acknowledgements
We sincerely thank the support provided by China National GeneBank. We thank all
the volunteers for their time and for self-collecting the fecal samples using our kit. We
are very grateful to Professor Karsten Kristiansen (Department of Biology, University
of Copenhagen, Denmark; BGI-Qingdao, BGI-Shenzhen, China) for his support for
the joint PhD program.

Author Contributions
H.J. and T.Z. conceived and organized this study. J.W. initiated the overall health
project. X.X., H.Y., S.Z., Y.H., W.L. and Y. Zong contributed to organization of the
cohort sample collection and questionnaire collection. H.L. led the DNA extraction and
sequencing. X.Q., J.Z., and R.W. generated the metabolic data. X. Liu, T.Z., and X.T.
processed the whole genome data. Y. Zou, X. Lin, Z.Z., H.Z., L.T., Q.W., Z.J. and L.X.
processed the metagenome data. X. Liu and X.T. performed the Mendelian
randomization analyses. X. Liu and H.J. wrote the manuscript. All authors contributed
to data and text in this manuscript.
438

439   **Competing Interests**
440   The authors declare no competing interests.
441
442

References

1.   Wang, J. & Jia, H. Metagenome-wide association studies: fine-mining the microbiome.

Nat Rev. Microbiol. **14**, 508-522 (2016).

2.   Moschen, A. R. *et al.* Lipocalin 2 protects from inflammation and tumorigenesis

associated with gut microbiota alterations. *Cell Host Microbe* **19**, 455-469 (2016).

3.   Long, X. *et al. Peptostreptococcus anaerobius* promotes colorectal carcinogenesis

and modulates tumour immunity. *Nat. Microbiol.* **4**, 2319-2330 (2019).

4.   Zhu, F. *et al.* Transplantation of microbiota from drug-free patients with schizophrenia

causes schizophrenia-like abnormal behaviors and dysregulated kynurenine

metabolism in mice. *Mol. Psychiatry* **25**, 2905-2918 (2020).

5.   Liu, X. *et al.* A genome-wide association study for gut metagenome in Chinese adults

illuminates complex diseases. *Cell Discovery* **7**, 9 (2021).

6.   Bonder, M. J. *et al.* The effect of host genetics on the gut microbiome. *Nat. Genet.* **48**,

1407-1412 (2016).

7.   Wang, J. *et al.* Genome-wide association analysis identifies variation in vitamin D

receptor and other host factors influencing the gut microbiota. *Nat. Genet.* **48**,

1396-1406 (2016).

8.   Turpin, W. *et al.* Association of host genome with intestinal microbial composition in a

large healthy cohort. *Nat. Genet.* **48**, 1413-1417 (2016).

9.   Blekhman, R. *et al.* Host genetic variation impacts microbiome composition across

human body sites. *Genome Biol.* **16**, 191 (2015).

10.   Rothschild, D. *et al.* Environment dominates over host genetics in shaping human gut

microbiota. *Nature* **555**, 210-215 (2018).

11.   Burgess, S., Timpson, N. J., Ebrahim, S. & Davey Smith, G. Mendelian randomization:

where are we now and where are we going? *Int. J. Epidemiol.* **44**, 379-388 (2015).

12.   Yang, Q., Lin, S. L., Kwok, M. K., Leung, G. M. & Schooling, C. M. The roles of 27

genera of human gut microbiota in ischemic heart disease, type 2 diabetes mellitus,

and their risk factors: a Mendelian randomization study. *Am. J. Epidemiol.* **187**,

1916-1922 (2018).

13.   Sanna, S. *et al.* Causal relationships among the gut microbiome, short-chain fatty

acids and metabolic diseases. *Nat. Genet.* **51**, 600-605 (2019).

14.   Sakaue, S. *et al.* Trans-biobank analysis with 676,000 individuals elucidates the

association of polygenic risk scores of complex traits with human lifespan. *Nat. Med.*

26, 542-548 (2020).

15.   Shin, S. Y. *et al.* An atlas of genetic influences on human blood metabolites. *Nat.*

Genet. **46**, 543-550 (2014).

16.   Cox, A. J. *et al.* Association of SNPs in the *UGT1A* gene cluster with total bilirubin and

mortality in the Diabetes Heart Study. *Atherosclerosis* **229**, 155-160 (2013).

17.   MacArthur, J. *et al.* The new NHGRI-EBI Catalog of published genome-wide

association studies (GWAS Catalog). *Nucleic Acids Res.* **45**, D896-D901 (2017).

18.   Verbanck, M., Chen, C. Y., Neale, B. & Do, R. Detection of widespread horizontal

pleiotropy in causal relationships inferred from Mendelian randomization between

complex traits and diseases. *Nat. Genet.* **50**, 693-698 (2018).

19.   Katano, Y. *et al.* Complete genome sequence of *Oscillibacter valericigenes* Sjm18-20[T]

(=NBRC 101213[T]). *Stand. Genomic Sci.* **6**, 406-414 (2012).

20.   Thingholm, L. B. *et al.* Obese individuals with and without type 2 diabetes show

different gut microbial functional capacity and composition. *Cell Host Microbe* **26**,

252-264.e10 (2019).

21.   Hu, H. J. *et al.* Obesity alters the microbial community profile in Korean adolescents.

PLoS One **10**, e0134333 (2015).

22.   Tims, S. *et al.* Microbiota conservation and BMI signatures in adult monozygotic twins.

ISME J. **7**, 707-717 (2013).

23.   Noack, J., Dongowski, G., Hartmann, L. & Blaut, M. The human gut bacteria

Bacteroides thetaiotaomicron and Fusobacterium varium produce putrescine and

spermidine in cecum of pectin-fed gnotobiotic rats. *J. Nutr.* **130**, 1225-1231 (2000).

24.   Luis, A. S. *et al.* Dietary pectic glycans are degraded by coordinated enzyme

pathways in human colonic Bacteroides. *Nat. Microbiol.* **3**, 210-219 (2018).

25.   PeBenito, A. *et al.* Comparative prevalence of *Oxalobacter formigenes* in three human

populations. *Sci. Rep.* **9**, 574 (2019).

26.   Ishigaki, K. *et al.* Large-scale genome-wide association study in a Japanese

population identifies novel susceptibility loci across different diseases. *Nat. Genet.* **52**,

669-679 (2020).

27.   Huo, Y. *et al.* Efficacy of folic acid therapy in primary prevention of stroke among

adults with hypertension in China: the CSPPT randomized clinical trial. *JAMA* **313**,

1325-1335 (2015).

28.   Jie, Z. *et al.* The gut microbiome in atherosclerotic cardiovascular disease. *Nat.*

Commun. **8**, 845 (2017).

29.   Yachida, S. *et al.* Metagenomic and metabolomic analyses reveal distinct

stage-specific phenotypes of the gut microbiota in colorectal cancer. *Nat. Med.* **25**,

968-976 (2019).

30.   Reese, A. T. *et al.* Microbial nitrogen limitation in the mammalian large intestine. *Nat.*

Microbiol. **3**, 1441-1450 (2018).

31.   Petrus, P. *et al.* Glutamine links obesity to inflammation in human white adipose tissue.

Cell Metab. **31**, 375-390.e11 (2019).

32.   Liu, R. *et al.* Gut microbiome and serum metabolome alterations in obesity and after

weight-loss intervention. *Nat. Med.* **23**, 859-868 (2017).

33.   Choi, W. M. *et al.* Glutamate signaling in hepatic stellate cells drives alcoholic

steatosis. *Cell Metab.* **30**, 877-889.e877 (2019).

34.   Kang, D. J. *et al.* Gut microbiota drive the development of neuroinflammatory

response in cirrhosis in mice. *Hepatology* **64**, 1232-1248 (2016).

35.   Long, T. *et al.* Plasma metals and cardiovascular disease in patients with type 2

diabetes. *Environ. Int.* **129**, 497-506 (2019).

36.   Kuo, C. F., Grainge, M. J., Zhang, W. & Doherty, M. Global epidemiology of gout:

prevalence, incidence and risk factors. *Nat. Rev. Rheumatol.* **11**, 649-662 (2015).

37.   Li, J. *et al.* An integrated catalog of reference genes in the human gut microbiome. *Nat.*

Biotechnol. **32**, 834-841 (2014).

38.   Qin, J. *et al.* A human gut microbial gene catalogue established by metagenomic

sequencing. *Nature* **464**, 59-65 (2010).

39.   Human Microbiome Project Consortium. Structure, function and diversity of the

healthy human microbiome. *Nature* **486**, 207-214 (2012).

40.   Maier, L. *et al.* Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature*

555, 623-628 (2018).

41.   Polderman, T. J. *et al.* Meta-analysis of the heritability of human traits based on fifty

years of twin studies. *Nat. Genet.* **47**, 702-709 (2015).

Figure legends

Figure 1 | Study design and workflow. This schematic representation highlights, for
each step, the research question that we sought to answer, the analysis workflow, the
data used, and the generalized result. We first performed metagenome and
metabolome GWAS to detect genetic variants associated with microbial features and
metabolic traits, respectively, both in discovery and replication cohorts (Step 1). We
then performed observational analysis to identify which microbial feature (taxa, GMM)
correlated with metabolic traits in this cohort (Step 2). We used 2,545 samples with
information of both microbial features and metabolic traits. We observed 457
significant associations between 500 unique microbial features and 112
anthropometric and blood metabolic traits at an FDR adjusted $P < 0.05$. We then
estimated causal relationships for the 457 observational associations through
bidirectional MR analysis in discovery cohort (Step 3). One-sample BMR detected 58
causal associations between microbial features and blood metabolites after multiple
test correction ($P < 1.09 \times 10^{-4}$); two-sample BMR detected the same associations
and an additional 14 associations. As a validation, we replicated the discovered
causal relationships by using the same MR analysis in an independent replication
cohort (Step 4). Over half (43) of the 58 causal associations were replicated in the
same direction ($P < 0.05$). Finally, we used two-sample MR analysis to investigate the
effects of the identified 72 causal relationships on diseases from the Biobank Japan
study (Step 5).

Figure 2 | Independent genetic variants and their explained variance of
microbial features. a, Density plot showing the distribution of number of independent
genetic variants that associated with each microbial feature ($r^2 < 0.1$ and $P < 1 \times 10^{-5}$),
as calculated by GWAS analysis for a total of 500 unique microbial features. The
x-axis indicates the number of independent genetic variants for each microbial feature
(taxon or GMM). The *y*-axis indicates the number of microbial features under a given
number of independent predictors. **b**, Variance explained by the corresponding

independent genetic variants for each microbial feature. The polar bar plot indicates
how much the independent genetic variants of each common genus (appeared at
least 50% of samples) explained for their phenotypic variance (relative abundance of
each genus). Genera were classified according to their respective phylum, which are
marked with different colors. The h^2 was calculated using REML method in GCTA
tools.
575
Figure 3 | Independent genetic variants and their explained variance of
metabolic traits. a, Density plot showing the distribution of number of independent
genetic variants that associated with each metabolic trait ($r^2 < 0.1$ and $P < 1 \times 10^{-5}$),
as calculated by GWAS analysis for a total of 112 metabolic traits. The *x*-axis
indicates the number of independent genetic variants for each metabolic trait. The
y-axis indicates the number of metabolic traits under a given number of independent
predictors. **b**, Variance explained by the corresponding independent genetic variants
for each metabolic trait. The polar bar plot indicates how much the independent
genetic variants of each metabolic trait explained for their phenotypic variance. Each
metabolic trait was classified into different catalogs which were marked with different
colors. The h^2 was calculated using REML method in GCTA tools.
587
Figure 4 | Identifying 58 causal relationships for the microbial features and
metabolic traits. a, Plot showing the causal effects of 12 specific microbial features
on 8 metabolic traits involved in the 17 causal associations from gut microbiome to
blood metabolites. **b**, Plot showing the causal effects of 7 blood metabolites on 33
microbial features involved in 41 causal associations from blood metabolites to gut
microbiome. Cells marked with "**" represent 43 of the 58 associations identified in
discovery cohort that were also replicated in replication cohort, while "*" represent the
other 15 only significant in discovery cohort. Cells are colored according to the beta
coefficients from one-sample MR analysis, with red and blue corresponding to positive
and negative associations, respectively.
598
Figure 5 | Causal effects of genus *Oscillibacter* and *Alistipes* on decreasing
blood triglyceride concentration. a, Schematic representation of the MR analysis
results: genetic predisposition to higher abundance of *Oscillibacter* is associated with
decreased blood triglyceride concentration, and to a lesser extent for lowering body
mass index (BMI) and waist-hip ratio (WHR). **b**, Forest plot representing the effect of
per 1-s.d. increase in *Oscillibacter* abundance on blood triglyceride, BMI and WHR, as
estimated using observational and MR analysis, respectively. Observational
correlation analysis was performed using a multivariate linear model in a total of 2,545
samples (purple). One-sample MR analysis was carried out by using a PGS
constructed by up to 134 genetic predictors as an instrumental variable, as estimated
in discovery cohort (blue) and replication cohort (red), respectively. The beta
estimates and 95% confidence interval (CI) values as well as *P* values from both the
observational and one-sample MR analysis were listed. **c**,**d**, Forest plots representing
the MR estimates and 95% CI values of the causal effects of *Oscillibacter* (**c**) and

Alistipes (**d**) on triglyceride level, respectively, as estimated using a one-sample MR
and six different two-sample MR methods both in discovery cohort (blue) and
replication cohort (red). The *P* values calculated by each MR method are also listed.
616
Figure 6 | Causal effects of *Proteobacteria* and *Escherichia coli* on diseases. a,**b**,
Forest plots representing the MR estimates and 95% CI values of the causal effects of
Proteobacteria (**a**) and *Escherichia coli* (**b**) on diseases, as estimated using six
different two-sample MR approaches (GSMR, IVW, MR Egger, Weighted median,
Simple mode and Weighted mode). We show the estimated β values (95%CI) and *P*
values, as estimated by each MR approach by using the diseases' summary statistics
from Biobank Japan and the gut microbiome GWAS summary data from this
discovery cohort with high-depth WGS.
625
626
627   **Methods**
628
Study participants. All adult Chinese individuals included in this study were recruited
for a multi-omic study, with some volunteers having samples from as early as 2015,
which would constitute the time dimension in '4D'. The discovery cohort was recruited
during a physical examination from March to May in 2017 in the city of Shenzhen,
including 2,002 individuals with blood samples, of which 1,539 had fecal samples. All
these individuals were enlisted for high-depth whole genome and whole metagenomic
sequencing. For replication, blood samples were collected from 1,430 individuals, of
which 1,006 had fecal samples. The replication cohort was designed in the same
manner but organized at smaller scales in multiple cities (Wuhan, Qingdao, etc.) in
China. The protocols for blood and stool collection, as well as the whole genome and
metagenomic sequencing, were similar to our previous studies[5,42-44]. For blood
samples, buffy coat was isolated and DNA was extracted using HiPure Blood DNA
Mini Kit (Magen, Cat. no. D3111) according to the manufacturer's protocol. Feces
were collected with MGIEasy kit, and stool DNA was extracted in accordance with the
MetaHIT protocol as described previously[5,42]. The DNA concentrations from blood and
stool samples were estimated by Qubit (Invitrogen). 200 ng of input DNA from blood
and stool samples were used for library preparation and then processed for
paired-end 100-bp and single-end 100-bp sequencing, respectively, using
BGISEQ-500 platform[45].
The study was approved by the Institutional Review Boards (IRB) at
BGI-Shenzhen, and all participants provided written informed consent at enrolment.
650
High-depth whole genome sequence for discovery cohort. 2,002 individuals in
discovery cohort were sequenced to a mean of 42× for whole genome. The reads
were aligned to the latest reference human genome GRCh38/hg38 with BWA[46]
(version 0.7.15) with default parameters. The reads consisting of base quality <5 or
containing adaptor sequences were filtered out. The alignments were indexed in the

BAM format using Samtools[47] (version 0.1.18) and PCR duplicates were marked for
downstream filtering using Picardtools (version 1.62). The Genome Analysis Toolkit's
(GATK[48], version 3.8) BaseRecalibrator created recalibration tables to screen known
SNPs and INDELs in the BAM files from dbSNP (version 150). GATKlite (v2.2.15) was
used for subsequent base quality recalibration and removal of read pairs with
improperly aligned segments as determined by Stampy. GATK's HaplotypeCaller
were used for variant discovery. GVCFs containing SNVs and INDELs from GATK
HaplotypeCaller were combined (CombineGVCFs), genotyped (GenotypeGVCFs),
variant score recalibrated (VariantRecalibrator) and filtered (ApplyRecalibration).
During the GATK VariantRecalibrator process, we took our variants as inputs and
used four standard SNP sets to train the model: (i) HapMap3.3 SNPs; (ii) dbSNP build
150 SNPs; (iii) 1000 Genomes Project SNPs from Omni 2.5 chip; and (iv) 1000G
phase1 high confidence SNPs. The sensitivity threshold of 99.9% to SNPs and 99%
to INDELs were applied for variant selection after optimizing for Transition to
Transversion (TiTv) ratios using the GATK ApplyRecalibration command. After
applying the recalibration, there were 60,978,451 raw variants left, including 55 million
SNPs, and 6 million INDELs.
We applied a conservative inclusion threshold for variants: (i) mean depth >8×; (ii)
Hardy-Weinberg equilibrium (HWE) $P > 10^{-5}$; and (iii) genotype calling rate > 98%. We
demanded samples to meet these criteria: (i) mean sequencing depth > 20×; (ii)
variant calling rate > 98%; (iii) no population stratification by performing principal
components analysis (PCA) analysis implemented in PLINK[49] (version 1.9); and (iv)
excluding related individuals by calculating pairwise identity by descent (IBD, Pi-hat
threshold of 0.1875) in PLINK. Only 10 samples were removed in quality control
filtering. After variant and sample quality control, 1,992 individuals with 6.12 million
common (MAF ≥ 5%) and 3.90 million low-frequency (0.5% ≤ MAF < 5%) variants
from discovery cohort were left for subsequent analyses.
683
Low-depth whole genome sequence for replication cohort. 1,430 individuals in
the replication cohort were sequenced to a mean of 8× for whole genome. We used
BWA to align the whole genome reads to GRCh38/hg38 and used GATK to perform
variant calling by applying the same pipelines as for the high-depth WGS data. After
completing the joint calling process with CombineGVCFs and GenotypeGVCFs
options, we obtained 43,402,368 raw variants. A more stringent process in the GATK
VariantRecalibrator stage compared with the high-depth WGS was then used, and the
sensitivity threshold of 98.0% to both SNPs and INDELs was applied for variant
selection after optimizing for Transition to Transversion (TiTv) ratios using the GATK
ApplyRecalibration command. Further, we kept variants with < 10% missing genotype
frequency and minor allele count > 5. All these high-quality variants were then
imputed using BEAGLE 5[50] with the 1,992 high-depth WGS dataset as reference
panel. We retained only variants with imputation info. > 0.7 and obtained 10,905,418
imputed variants. We further filtered this dataset to keep variants with
Hardy-Weinberg equilibrium $P > 10^{-5}$ and genotype calling rate > 90%. Similar to what
we did for the discovery cohort, samples were required to have mean sequencing

depth > 6×, variant call rate > 98%, no population stratification and no kinship. Finally,
1,430 individuals with 5,884,439 high-quality common and low-frequency variants
(MAF ≥ 0.5%) from the replication cohort were left for subsequent analysis. To assess
the data quality, we sequenced 27 samples with both high-depth and low-depth WGS
data and then compared the 5,318,809 variants between them for each individual.
The average genotype concordance was 98.66% (**Supplementary Table 15**).
706
Metagenomic sequencing and profiling. High-quality whole metagenomic
sequencing was performed for 1,539 samples from the discovery cohort and 1,004
samples from the replication cohort with fecal samples available. The reads were
aligned to hg38 using SOAP2[51] (version 2.22; identity ≥ 0.9) to remove human reads.
The gene profiles were generated by aligning high-quality sequencing reads to the
integrated gene catalog (IGC) by using SOAP2 (identity ≥ 0.95) as previously
described[37]. The relative abundance profiles of phylum, order, family, class, genera
and species were determined from the gene abundances. To eliminate the influence
of sequencing depth in comparative analyses, we downsized the unique IGC mapped
reads to 20 million for each sample. The relative abundance profiles of gene, phylum,
order, family, class, genus and species were determined accordingly using the
downsized mapped reads per sample.
GMMs (gut metabolic modules) reflect bacterial and archaeal metabolism specific
to the human gut, with a focus on anaerobic fermentation processes[52]. The current set
of GMMs was built through an extensive review of the literature and metabolic
databases, inclusive of MetaCyc[53] and KEGG, followed by expert curation and
delineation of modules and alternative pathways. Finally, we identified 620 common
microbial taxa and GMMs present in 50% or more of the samples.
725
Metabolic trait profiling. Measurements of metabolic traits (anthropometric
characteristics and blood metabolites) were performed for all 3,432 individuals during
the physical examination in this study. The clinical tests, including blood tests and
urinalysis, were performed in licensed physical examination organization. The
anthropometric measurements such as height, weight, waistline and hipline were
measured by nurses. Age and gender were self-reported. The metabolites, i.e.
vitamins, hormones, amino acids and trace elements including heavy metals, were
chosen from a health management perspective. Measurements of blood metabolites
were performed as described in detail by Jie et al.[39]; blood amino acids were
measured by ultra high pressure liquid chromatography (UHPLC) coupled to an AB
Sciex Qtrap 5500 mass spectrometry (AB Sciex, US) with the electrospray ionization
(ESI) source in positive ion mode using 40 µl plasma; blood hormones were
measured by UHPLC coupled to an AB Sciex Qtrap 5500 mass spectrometry (AB
Sciex, US) with the atmospheric pressure chemical ionization (APCI) source in
positive ion mode using 250 µl plasma; blood trace elements were measured by an
Agilent 7700x ICP-MS (Agilent Technologies, Tokyo, Japan) equipped with an
octupole reaction system (ORS) collision/reaction cell technology to minimize spectral
interferences using 200 µl whole blood; water-soluble vitamins were measured by

UPLC coupled to a Waters Xevo TQ-S Triple Quad mass spectrometry (Waters, USA)
with the electrospray ionization (ESI) source in positive ion mode using 200 µl plasma;
fat-soluble vitamins were measured by UPLC coupled to an AB Sciex Qtrap 4500
mass spectrometry (AB Sciex, USA) with the atmospheric pressure chemical
ionization (APCI) source in positive ion mode using 250 µl plasma.
749
Observational correlation analyses. As many microbial features (taxonomies and
pathways) are highly correlated, we first performed a number of Spearman correlation
tests and kept only one member of pairs of bacteria or GMMs showing >0.99
correlation coefficient. This filtering resulted in a final set of 500 unique features (99
GMMs and 401 gut taxa) that were used for analyses. We correlated these 500
microbial features with 112 measured metabolic traits, including 9 anthropometric
measurements (BMI, WHR, etc.) and 103 blood metabolites (amino acids, vitamins,
microelements, etc.) in the 3,432 individuals. All metabolic traits and microbial
features were transformed using natural logarithmic function to reduce skewness of
distributions. For each phenotype, we excluded outlier individuals with more than four
standard deviations away from the mean. The metabolite measures were then
centered and scaled to mean of 0 and standard deviation of 1.
The relationship between metabolic traits and microbial features were evaluated
by multivariable linear regression analysis while adjusted for age and gender. After
achieving the raw P value, we used the *p.adjust()* function in R (v3.2.5)) to perform the
multiple test correction and calculated adjusted P values with the Benjamini–
Hochberg procedure. The results were considered significant when FDR adjusted P
value was < 0.05. The correlated microbial features and metabolic traits, raw P and
FDR adjusted P values, are included in **Supplementary Table 9**.
769
Clustering of microbiome-metabolite associations. To assess the association
clusters of 58 identified causal relationships involving the effects of 12 microbial
features on 8 metabolic traits and the effects of 7 metabolic traits on 33 microbial
features, we performed a hierarchical clustering analysis. Beta coefficients of
associations between the microbial features and metabolic traits from one-sample MR
analysis were used to construct distance matrices. Complete-linkage hierarchical
clustering was used to cluster the metabolites and microbiome traits from the distance
matrices using the 'hclust' function in R, and the results were visualized as a heatmap.
778
Genome-wide association analysis for microbial features. We tested the
associations between host genetic variants and gut bacteria using linear or logistic
model based on the abundance of gut bacteria. The abundance of bacteria with
occurrence rate over 95% in the cohort was transformed by the natural logarithm and
the outlier individual who was located away from the mean by more than four standard
deviations was removed, so that the abundance of bacteria could be treated as a
quantitative trait. Otherwise, we dichotomized bacteria into presence/absence
patterns to prevent zero inflation; the abundance of bacteria could then be treated as
a dichotomous trait. Next, for 10 million common and low-frequency variants (0.5% ≤

MAF < 5%) identified in the discovery cohort and 5.9 million common and
low-frequency variants identified in replication cohort, we performed a standard single
variant (SNP/INDEL)-based M-GWAS analysis via PLINK using a linear model for
quantitative trait or a logistic model for dichotomous trait. Given the effects of diet and
lifestyles on microbial features, we included age, gender, BMI, defecation frequency,
stool form, 12 diet and lifestyle factors, as well as the top four principal components
(PCs) as covariates for M-GWAS analysis in both the discovery and the replication
cohort.
796

Genome-wide association analysis for metabolic traits. For each of the 112
anthropometric and metabolic traits, the $\log_{10}$-transformed of the median-normalized
values was used as a quantitative trait. Samples with missing values and values
beyond 4 s.d. from the mean were excluded from association analysis. Each of the 10
million common and low-frequency variants identified in the discovery cohort and the
5.9 million common and low-frequency variants identified in the replication cohort was
tested independently using a linear model for quantitative trait implemented in PLINK.
Age, gender and the top four PCs were included as covariates.
805

Independent predictor and explained phenotypic variance. For each whole
genome-wide association result of microbial features and metabolic traits, we first
selected genetic variants that showed association at $P < 1 \times 10^{-5}$ and then performed
the linkage disequilibrium (LD) estimation with a threshold of LD $r^2 < 0.1$ for clumping
analysis to get independent genetic predictors. The P-value threshold of 1×10^{-5} was
used for selection of genetic predictors associated with microbial features by
maximizing the strength of genetic instruments and the amount of the average genetic
variance explained by the genetic predictors in an independent sample. For each
microbial feature, we got genetic instruments in discovery cohort using different P
thresholds, including 5×10^{-8}, 1×10^{-7}, 1×10^{-6} and 1×10^{-5}. We tested the strength
of these instruments under different P thresholds by checking whether they predicted
corresponding microbial features in an independent sample (**Supplementary Note**,
Supplementary Table 16 and **Supplementary Fig. 6**), and we observed that the
mean value of instrumental F statistics is 3.57 and on average only 0.28% phenotype
variance could be explained by instruments on microbial features when using 5×10^{-8}
as instrumental cut-off. Therefore, we used a more liberal threshold of $P < 1 \times 10^{-5}$ to
select the instruments for microbial features, and the instrumental mean F statistics
reached 51.4 (greater than 10), which indicates a strong instrument[54]. The average
phenotypic variance explained by instruments on microbial features was 22.6% for
the discovery cohort and 5.09% for the replication cohort (**Supplementary Fig. 2**).
For consistency, we used the same threshold and procedure for selecting genetic
predictors of metabolic traits in both the discovery and the replication cohort. The LD
estimation between variants was calculated in 2,002 samples for the discovery cohort
and in 1,430 samples for the replication cohort, respectively. For each phenotype, the
variance explained by the corresponding independent genetic predictors was
estimated using a restricted the maximum likelihood (REML) model as implemented

in the GCTA software[55]. We adjusted for age, gender and the top four PCs in the
REML analysis.

One-sample MR analysis. To investigate the causal effects between microbial
features and metabolic traits available from the same cohort, we first performed
one-sample bidirectional MR analysis in discovery cohort, which included 1,539
individuals with both metabolite and microbiome traits. We specified a threshold of $P <$
1×10^{-5} to select SNP instruments and LD $r^2 < 0.1$ threshold for clumping analysis to
get independent genetic variants for MR analysis. Then, an unweighted polygenic risk
score (PRS) was calculated for each individual using independent genetic variants
from GWAS data. Each SNP was recoded as 0, 1 and 2, depending on the number of
trait-specific risk increasing alleles carried by an individual. We performed
Instrumental variable (IV) analyses employing two-stage least square regression
(TSLS) method[56]. In the first stage, for each exposure trait, association between the
GRS and observational phenotype value was assessed using linear regression and
predicted fitted values based on the instrument were obtained. In the second stage,
linear regression was performed with outcome trait and genetically predicted
exposure level from the first stage. In both stages, analyses were adjusted for age,
gender and top four principal components of population structure. For each trait,
TSLS was performed using 'ivreg' command from the AER package in R. We
attempted to replicate the causal effects between traits in replication cohort with 1,004
individuals.

Two-sample MR analysis. To maximize the sample size in MR analysis and confirm
the causal effect between microbial features and metabolic traits, we also performed
two-sample bidirectional MR analysis using six different methods, including
genome-wide complex trait analysis-generalized summary Mendelian randomization
(GCTA-GSMR) approach[57] and five other methods (Inverse-variance weighting
(IVW)[58,59], MR–Egger regression[60,61], Weighted median[62], Mode-based estimate
(MBE)[63] including Simple mode and Weighted mode) implemented in "TwoSampleMR"
R package as a robust validation. A consistent effect across the six methods is less
likely to be a false positive. If the genetic variants have horizontally pleiotropic effects
but are independent of the effects of the genetic variants on the exposure, this is
known as balanced pleiotropy. If all the pleiotropic effects are biasing the estimate in
the same direction (directional pleiotropy), this will bias the results (with the exception
of the MR-Egger method). We used the MR-PRESSO (mendelian randomization
pleiotropy residual sum and outlier) Global test to estimate for the presence of
directional pleiotropy.
We first performed GWAS analysis for every trait and used summary statistics
data for MR analysis. Genetic variants with $P < 1 \times 10^{-5}$ and LD $r^2 < 0.1$ were selected
as instrumental variables.

In vitro growth experiments of *Escherichia coli*. To directly test the interactions
between *Escherichia coli* and 5-methyltetrahydrofolic acid, the anaerobic growth of a

strain *Escherichia coli* AM17-9 was characterized at different concentrations of
5-methyltetrahydrofolic acid. The *Escherichia coli* AM17-9, isolated from feces of a
male, was routinely grown in Luria-Bertani (LB) broth while supplementing
5-methyltetrahydrofolic acid with concentrations of 0, 1 and 2 ng/ml, respectively. The
normal concentration of 5-methyltetrahydrofolic acid in human blood ranged from 4.4
ng/ml to 32.8 ng/ml. The growth of *Escherichia coli* AM17-9 was inhibited when
supplementing 5-methyltetrahydrofolic acid from 0 to 2 ng/ml. The optical density at
600 nm (OD600) was measured at intervals of two hours using a microplate reader.
884
MR analyses for diseases in Biobank Japan. We downloaded summary statistics
data for 42 diseases and 59 blood quantitative traits in 212,453 Japanese
individuals[26] (http://jenger.riken.jp/en/result; **Supplementary Table 13**). By combining
these data and the gut microbiome GWAS summary data from discovery cohort with
high-depth WGS, we performed the two-sample bidirectional MR analysis to
investigate the causal effect between the exposure (40 microbial features and 12
metabolic traits that were involved in the 72 significant causal relationships; **Fig. 4**)
and the outcome (42 diseases from BioBank Japan) by applying the GSMR method
and the other five MR tests as described in the previous paragraph. For consistency,
genetic variants with $P < 1 \times 10^{-5}$ and LD r^2 <0.1 were also selected as instrumental
variables for phenotypes in the Biobank Japan study.
896
897 **Methods-only References**
898

42. Jie, Z. *et al.* A transomic cohort as a reference point for promoting a healthy human

gut microbiome. *Medicine in Microecology* **8**, 100039 (2021).

43. Zhu, J. *et al.* Over 50,000 metagenomically assembled draft genomes for the human

oral microbiome reveal new taxa. *Genomics, Proteomics & Bioinformatics*, doi:

10.1016/j.gpb.2021.05.001 (2021).

44. Liu, X. *et al.* Metagenome-genome-wide association studies reveal human genetic

impact on the oral microbiome. *bioRxiv*, doi:10.1101/2021.05.06.443017 (2021).

45. Fang, C. *et al.* Assessment of the cPAS-based BGISEQ-500 platform for

metagenomic sequencing. *Gigascience* **7**, 1-8 (2018).

46. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* **25**, 1754-1760 (2009).

47.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,

2078-2079 (2009).

48.   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for

analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297-1303

(2010).

49.   Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based

linkage analyses. *Am. J. Hum. Genet.* **81**, 559-575 (2007).

50.   Browning, B. L., Zhou, Y. & Browning, S. R. A one-penny imputed genome from

next-generation reference panels. *Am. J. Hum. Genet.* **103**, 338-348 (2018).

51.   Li, R. *et al.* SOAP2: an improved ultrafast tool for short read alignment. *Bioinformatics*

25, 1966-1967 (2009).

52.   Vieira-Silva, S. *et al.* Species-function relationships shape ecological properties of the

human gut microbiome. *Nat. Microbiol.* **1**, 16088 (2016).

53.   Caspi, R. *et al.* The MetaCyc database of metabolic pathways and enzymes and the

BioCyc collection of Pathway/Genome Databases. *Nucleic Acids Res.* **42**, D459-D471

(2014).

54.   Teumer, A. Common methods for performing Mendelian randomization. *Front.*

Cardiovasc. Med. **5**, 51 (2018).

55.   Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide

complex trait analysis. *Am. J. Hum. Genet.* **88**, 76-82 (2011).

56.   Permutt, T. & Hebel, J. R. Simultaneous-equation estimation in a clinical trial of the

effect of smoking on birth weight. *Biometrics* **45**, 619-622 (1989).

57.   Zhu, Z. *et al.* Causal associations between risk factors and common diseases inferred

from GWAS summary data. *Nat. Commun.* **9**, 224 (2018).

58.   Burgess, S., Butterworth, A. & Thompson, S. G. Mendelian randomization analysis

with multiple genetic variants using summarized data. *Genet. Epidemiol.* **37**, 658-665

(2013).

59.   Bowden, J. *et al.* A framework for the investigation of pleiotropy in two-sample

summary data Mendelian randomization. *Stat. Med.* **36**, 1783-1802 (2017).

60.   Bowden, J., Davey Smith, G. & Burgess, S. Mendelian randomization with invalid

instruments: effect estimation and bias detection through Egger regression. *Int. J.*

Epidemiol. **44**, 512-525 (2015).

61.   Bowden, J. *et al.* Assessing the suitability of summary data for two-sample Mendelian

randomization analyses using MR-Egger regression: the role of the I^2 statistic. *Int. J.*

Epidemiol. **45**, 1961-1974 (2016).

62.   Bowden, J., Davey Smith, G., Haycock, P. C. & Burgess, S. Consistent estimation in

Mendelian randomization with some invalid instruments using a weighted median

estimator. *Genet. Epidemiol.* **40**, 304-314 (2016).

63.   Hartwig, F. P., Davey Smith, G. & Bowden, J. Robust inference in summary data

Mendelian randomization via the zero modal pleiotropy assumption. *Int. J. Epidemiol.*

46, 1985-1998 (2017).

Data availability

The human reference hg38 datasets were publicly available from

http://hgdownload.soe.ucsc.edu/goldenPath/hg38/bigZips/. All summary statistics that

support the findings of this study including the associations between host genetics

and microbiomes, host genetics and metabolites are publicly available from
https://ftp.cngb.org/pub/CNSA/data2/CNP0000794/. The release of the data was
approved by the Ministry of Science and Technology of China (Project ID:
2020BAT1137). Individual-level data including host genetics, metagenomics and
metabolites have been uploaded to the GSA database
(https://ngdc.cncb.ac.cn/bioproject/browse/PRJCA005334). Access to individual-level
data has to be approved by corresponding authors (tao.zhang@genomics.cn,
jiahuijue@genomics.cn), and is subject to the policies and approvals from the Human
Genetic Resource Administration, Ministry of Science and Technology of the People's
Republic of China. The summary statistics data for 42 diseases and 59 blood
quantitative traits in 212,453 Japanese individuals are available from Biobank Japan
(http://jenger.riken.jp/en/result).
969
Code availability
The host genome reads were aligned to the latest reference human genome
GRCh38/hg38 with BWA (v.0.7.15; http://bio-bwa.sourceforge.net/). The alignments
were indexed in the BAM format using Samtools (v.0.1.18;
http://samtools.sourceforge.net/) and PCR duplicates were marked for downstream
filtering using Picardtools (v.1.62; http://broadinstitute.github.io/picard/). The genome
variants calling were performed using GATK (v3.8;
https://gatk.broadinstitute.org/hc/en-us). The low-depth data were imputed using
BEAGLE (v.5.0; https://faculty.washington.edu/browning/beagle/b5_0.html). The
metagenome reads were aligned to hg38 using SOAP (v.2.22;
http://soap.genomics.org.cn). Quality control, association analyses and genetic risk
score analyses were performed in PLINK (v1.90; http://zzz.bwh.harvard.edu/plink/).
Variance explained analysis was performed using REML model in GCTA (v.1.26.0;
https://cnsgenomics.com/software/gcta). One-sample MR analyses were performed
using TSLS method in AER package (v1.2-9;
https://www.rdocumentation.org/packages/AER/versions/1.2-9/topics/ivreg).
Two-sample MR analyses were performed in GSMR (v1.0.7;
http://cnsgenomics.com/software/gsmr/) and TwoSampleMR (v.0.5.6;
https://mrcieu.github.io/TwoSampleMR/). All statistics analyses and visualizations
were performed in R (v.3.2.5; https://www.r-project.org).
990
991

Step 1. What are the genetic variants associated with metabolic traits and microbial features?

Discovery cohort: 2,002 samples with anthropometric and blood metabolites and high-depth whole genome, 1,539 of which with gut metagenome; 10M common and low-frequency variants (MAF > 0.005; HWE P > 10⁻⁵; variants calling rate > 0.98)

Replication cohort: 1,430 samples with anthropometric and blood metabolites and low-depth whole genome, 1,006 of which with gut metagenome; 5.9M common and low-frequency variants (MAF > 0.005; HWE P > 10⁻⁵; variants calling rate > 0.98)

Metagenome and metabolome GWAS detected genetic variants associated with microbial features and metabolic traits, respectively.

Step 2. Which microbial features correlate with metabolic traits?

A total of 3,432 samples with anthropometric and blood metabolic traits and whole genome data, of which 2,545 samples with gut metagenome

Identifying 457 observationally significant associations between 500 unique microbial features and 112 anthropometric and blood metabolites at a FDR adjusted P < 0.05 using a multivariable linear regression analysis

Step 3. Do changes in microbial features causally affect metabolic traits or vice versa?

One sample BMR detected 58 causal associations between microbial features and blood metabolites after multiple test correction (P < 1.09 ×10⁻⁴); two sample BMR detected the same associations and an additional 14 associations

17 causal associations from gut microbiome to blood metabolites involving12 microbial features and 8 blood metabolites
Two taxa, genus Oscillibacter and Alistipes decreased triglyceride level.

41 causal associations from blood metabolites to gut microbiome involving 7 blood metabolites and 33 microbial features

Step 4. Can we replicate causal relationships?

43 of the 58 causal associations were replicated in the same directions and P < 0.05 in an independent sample of replication cohort, for example genus Oscillibacter and Alistipes consistently decreased triglyceride level.

Step 5. Whether the identified causal relationships have potential link to disease?

Metagenome GWAS   Disease and and blood quantitative traits GWAS

Trait X → Trait Y

Two sample MR using metagenome GWAS summary data in this study, together with disease and blood quantitative traits GWAS summary data from Biobank Japan study

We replicate the causal effect of uric acid on increased species unclassified Lachnospiraceae bacterium 9_1_43BFAA abundance

Some gut bacteria potentially linked to disease risk