Adjacent terrestrial landscapes impact the biogeographical pattern of soil *Escherichia coli* in produce

fields by modifying the importance of environmental selection and dispersal

Jingqiu Liao[a, b], Peter Bergholz[c], Martin Wiedmann[a#]

Department of Food Science, Cornell University, Ithaca, NY 14853[a]; Graduate Field of Microbiology,

Cornell University, Ithaca, NY 14853[b]; Department of Microbiological Sciences, North Dakota State

University, Fargo, ND 58105[c]

Running Head: Effects of landscape on the distribution of *E. coli*

Address correspondence to Martin Wiedmann, martin.wiedmann@cornell.edu

ABSTRACT High-quality habitats for wildlife (e.g., forest) provide essential ecosystem services while increasing species diversity and habitat connectivity. Unfortunately, presence of such habitats adjacent to produce fields may increase risk for contamination of fruits and vegetables by enteric bacteria, including *Escherichia coli*. *E. coli* survives in extra-host environments (e.g., soil) and could disperse across landscapes by wildlife. Understanding how terrestrial landscapes impact the distribution of soil *E. coli* is of importance in assessing the contamination risk of agricultural products. Here, using multi-locus sequence typing, we characterized 938 *E. coli* soil isolates collected from two watersheds with different landscape patterns in New York state, USA, and compared the distribution of *E. coli* and the influence of two ecological forces (environmental selection and dispersal) on the distribution between these two watersheds. Results showed that for the watershed with widespread produce fields, sparse forests, and limited interaction between the two land-use types, *E. coli* composition was significantly different between produce field sites and forest sites; this distribution was shaped by relatively strong environmental selection likely from soil phosphorus and slight dispersal limitation. For the watershed with more forested areas and stronger interaction between produce field sites and forest sites, *E. coli* composition between these two land-use types was relatively homogeneous; this distribution appeared to a consequence of wildlife-driven dispersal, inferred by competing models. Collectively, our results suggest that terrestrial landscape attributes could impact the biogeographic pattern of enteric bacteria by adjusting the importance of environmental selection and dispersal.

IMPORTANCE Understanding the ecology of enteric bacteria in extra-host environments is important to allow for development and implementation of strategies to minimize pre-harvest contamination of produce with enteric pathogens. Our findings suggest that watershed landscape is an important factor influencing the importance of ecological drivers and dispersal patterns of *E. coli*. For watersheds with widespread produce fields, *E. coli* appears to experience local adaptation, possibly due to exposure to

environmental stresses associated with agricultural activities. In contrast, for watersheds with high forest

coverage we found evidence for wildlife-driven dispersal of *E. coli*, which might facilitate more frequent

genetic exchange in this environment. Agricultural areas in such watersheds may have a higher risk of

produce contamination due to less environmental constraints and higher potential of dispersal of enteric

bacteria between locations. The significance of our research lies in exploring ecological principles

underlying the biogeographic pattern of enteric bacteria at the regional level, which can inform

agricultural, environmental and public health scientists that aim to reduce the risk of food contamination

by enteric bacteria.

KEYWORDS enteric bacteria, landscape, environmental selection, dispersal, wildlife

INTRODUCTION

Forests and other riparian buffers can provide ecological and agricultural benefits (e.g., reducing

soil erosion and leaching of chemical and fecal waste into surface water sources, providing habitat and

connective pathways for wildlife) as well as aesthetic benefits (1–3). While high quality habitats (e.g.,

forest) offer conservation services, they may also bring unintended consequences and may increase the

risk of pre-harvest contamination of produce crops. For one, extra-host environments, such as soil in

high quality habitats, can be a critical reservoir for enteric bacteria, leading to potential dispersal of

enteric bacteria to adjacent agricultural fields (4). Direct fecal deposition onto produce by wildlife is also

a potential pathway introducing enteric bacteria onto food crops (5). By providing wildlife movement

pathways, high quality habitats may facilitate the wildlife-driven dispersal of enteric bacteria through

riparian corridors to agricultural regions, possibly resulting in contamination of food crops (6–9).

Subsequent persistence and/or regrowth of pathogenic enteric bacteria introduced into fresh produce

fields would then further increase food safety risks (10). Since the survival of enteric bacteria and

movement of wildlife vary by land use types (11, 12), it is reasonable to hypothesize that watershed

landscape impacts the distribution of enteric bacteria.

Environmental selection and dispersal are two fundamental ecological forces that drive the

distribution of bacteria (13–16). The essential roles of environmental selection via abiotic (e.g., pH,

salinity) and biotic selective pressures on bacteria have been well documented in many local and even

global habitats (17–20). Environmental selection facilitates the genetic divergence of some

ecophysiological traits owing to their contribution to fitness benefits for adaptation of bacteria to diverse

habitats, such as those with different land cover types (21, 22). With the influence of environmental

selection, a high level of bacterial dissimilarity between locations (beta diversity) can be maintained in a

wide range of environments (15, 23, 24). The role of dispersal in driving the distribution of bacteria at

local as well as regional scales is evident since dispersal provides a mechanism for bacteria to colonize

new habitats (25, 26). The relative importance of dispersal in shaping bacterial distribution varies among

microbial taxa due to diversity in the capacity of bacteria to disperse via wind, water, and wildlife. For

example, bacteria with a long range dispersal capacity (e.g., *Polaromonas*) tend to exhibit a more global

distribution (27), while bacteria with a limited dispersal range (e.g., *Rhizobiaceae*, *Bradyrhizobiaceae*,

Xanthomonadaceae) tend to show more ecological specialization (15). Importantly, wildlife presence

and movement is fundamentally affected by the physical elements and features of land (15). Thus,

wildlife-driven dispersal of bacteria can be quantitatively predicted by landscape ecological methods

based on the relationship of wildlife behaviors and landscape characteristics (e.g., patchiness, land-use

interspersion, patch connectivity, patch diversity, and land-use interactions) (8, 28–30). Based on these

principles, investigating environmental selection and dispersal of enteric bacteria from and within

habitats with distinct landscape patterns has the power to elucidate the role of terrestrial landscapes in

impacting the distribution of enteric bacteria and assess the associated risk of preharvest contamination

of food by pathogenic enteric bacteria.

As a commensal or pathogenic enteric bacterium widespread in diverse habitats, *Escherichia coli*

primarily resides in the intestines of warm-blooded animals, and survives in extra-host environments

such as water, soil, sediments as well (31, 32). Soil is a habitat of particular interest for *E. coli*, since the

high chemical and physical heterogeneity of soil across different environments could pose multifarious

environmental selection pressures on *E. coli* (22, 23, 32). Observations that the prevalence of *E. coli*

varies by land cover types (e.g., deciduous forest, cropland, pasture) (22) also suggest that different land

uses could stimulate different types and intensities of selective pressures that act on *E. coli*. The key

edaphic variables influencing the growth of *E. coli* in soil are commonly recognized as pH and moisture

(23, 33), while some other soil properties such as organic matter and texture could also play a role (10).

In addition, wildlife, such as avian species and ruminant animals, could act as dispersal vehicles of *E.*

coli (34, 35). *E. coli* can also be transmitted between wildlife hosts through contact and can be deposited

in new locations (e.g., produce fields) by defecation, which often happens when wildlife forages for

food (36). Given the intensive interaction with both extra-host and host habitats, *E. coli* may be a useful

model to build predictive capabilities surrounding interactions between bacteria and agricultural

landscapes at the meter to kilometer scale. Such an understanding is particularly important for lands

where fresh fruits and vegetables are cultivated, as it could be used to develop better strategies for

minimizing pathogen introduction into preharvest environments.

We hypothesized that the importance of environmental selection and dispersal for the

distribution of *E. coli* is dependent on landscape and specifically hypothesized that (i) *E. coli* in

watersheds with higher coverage of agricultural environments is strongly driven by environmental

selection associated with agricultural activity, while (ii) *E. coli* in regions with higher coverage of

natural environment is largely influenced by wildlife-driven dispersal. To test our hypotheses, we

characterized 938 generic *E. coli* isolates obtained from soil samples collected from two watersheds -

Flint Creek and Hoosic River, both located in the New York State, using a hierarchical multi-locus

sequence typing (MLST) scheme. These two watersheds represent an interesting comparison between

one with widespread produce fields and limited interaction between produce fields and forest (Flint

Creek; 69% produce field, 12% forest by area; adjacency$_{produce|forest}$ = 23%; Fig. 1a) and one with heavily

forested areas and strong interaction between produce fields and forest (Hoosic River; 28% produce

field, 38% forest by area; adjacency$_{produce|forest}$ = 36%; Fig. 1b). Next, we investigated the distribution of

E. coli in these two watersheds, and assessed the relationship between *E. coli* distribution and soil

variables and the distance-decay relationship among *E. coli* populations. Last, we developed dispersal

models for four wildlife vehicle candidates (large nuisance wildlife species, small mammals, small

flocking insectivore/granivores, and migratory bird flocks) to quantify the importance of wildlife-driven

dispersal on the dispersal of *E. coli* in the two watersheds.

RESULTS

Distribution of *E. coli*. Soil samples from Flint Creek (predominated by produce fields) showed

considerably lower prevalence of *E. coli* than soil samples from Hoosic River (predominated by forests);

35% and 72% of soil samples, respectively, were positive for *E. coli* in these two watersheds. These

samples yielded 289 and 649 *E. coli* isolates, respectively. Based on initial 2-gene MLST (*mdh* and

uidA), a total of 138 isolates from Flint Creek and 277 isolates from Hoosic River were selected for

characterization by full 7-gene MLST (*aspC*, *clpX*, *icd*, *lysP*, *fadD*, in addition to *mdh* and *uidA*). The 7-

gene MLST generated 121 unique multilocus sequence types (ST) for Flint Creek and 191 unique ST for

Hoosic River (Table S1, Table S2). Analysis by goeBURST identified 96 and 108 *E. coli* clonal groups

for Flint Creek and Hoosic River, respectively, based on ST at single locus variant level (Fig. S1, Fig.

S2). For Flint Creek, forest sites had slightly higher mean richness of *E. coli* clonal groups than produce

field sites, but the difference was not significant ($p = 0.86$) (Fig. S3). In contrast, the mean richness of *E.*

coli clonal groups from produce field sites was significantly higher than forest sites in the Hoosic River

watershed ($p < 0.05$) (Fig. S3).

E. coli in the two watersheds displayed distinct distribution patterns. For Flint Creek, principal

coordinates analysis (PCoA) based on the dissimilarity of *E. coli* clonal groups clustered sampling sites

by land-use (Fig. 2a). Both permutational multivariate analysis of variance (PERMANOVA) test and

analysis of similarities (ANOSIM) test showed that this clustering by land-use was significant ($p < 0.05$;

Table S3, Table S4). By contrast, for the Hoosic River, the sampling sites were not significantly

clustered by land-use in PCoA (Fig. 2b) of *E. coli* clonal groups (PERMANOVA $p = 0.86$, ANOSIM p

$= 0.58$; Table S3, Table S4), indicating a more homogeneous composition of *E. coli* between produce

field sites and forest sites in this watershed. To test the robustness of these results and to assess the

effects of potential sampling bias on the *E. coli* diversity, we further repeated PCoA, PERMANOVA,

and ANOSIM analyses after excluding sites with a low number of *E. coli* clonal groups detected (≤ 3);

sites being excluded were Field 6, Field 8, and Forest F9 from Flint Creek. Results showed that

sampling sites from this subset were significantly clustered by land-use in the PCoA plot (Fig. S4) for *E.*

coli clonal groups from Flint Creek at least at the 0.1 level (PERMANOVA $p = 0.05$, ANOSIM $p = 0.08$;

Table S5). Slightly larger *p* values in PERMANOVA and ANOSIM tests using a subset of sites (as

comparted to *p*-values for the whole set of sites) are likely due to the reduced sample size in these

analyses.

The importance of soil variables driving the distribution of *E. coli*. We first employed

variance partitioning analysis (VPA) to quantify the contribution of soil variables and spatial variables to

the distribution of *E. coli* from Flint Creek and Hoosic River. After screening for high levels of

covariation, moisture, pH, sodium, phosphorus, barium, manganese, and antimony were included in

VPA (Table S6). VPA revealed that the soil variables individually explained 11.7% of the biological

variation of dissimilarity of *E. coli* clonal groups from Flint Creek, while spatial variables individually

explained 5.5% of this variation (Fig. 3a). The variation explained by spatially structured soil variables

was relatively low (0.1%) and about 83% of the variation was unexplained by selected variables (Fig.

3a). By contrast, for *E. coli* clonal groups from Hoosic River, about 99% of the variation of dissimilarity

could not be explained by selected variables (Fig. 3b). Individually, selected soil variables did not

explain any of the observed variation and spatial variables only explained 1% of the variation for the

Hoosic River watershed (Fig. 3b). These results indicate that environmental selection outweighs spatial

factors with regard to the importance for *E. coli* distribution in the Flint Creek watershed; *E. coli* from

this watershed appear to undergo much stronger environmental selection or a much slower rate of

dispersal than *E. coli* in the Hoosic River watershed.

To identify the key soil variables correlated with the dissimilarity of *E. coli* clonal groups, we

performed partial Mantel tests by correcting the correlation of geographic distance and biological

dissimilarity. For Flint Creek, partial Mantel tests showed that geographic distance-corrected

phosphorus and antimony concentrations in soil were significantly correlated with the dissimilarity of *E.*

coli clonal groups (correlation coefficient r = 0.18, $p < 0.05$ and r = 0.41, $p < 0.05$, respectively) (Fig.

3c). In addition, for sites in Flint Creek, we observed that the mean concentration of phosphorus was

significantly higher for soil samples from produce field sites as compared to samples from forest sites

(Mann-Whitney test $p < 0.05$, Fig. S5a). Also, in Flint Creek, the mean concentration of antimony was

higher for soil samples from produce field sites compared to samples from forest sites, but the difference

was not significant (Mann-Whitney test $p = 0.45$, Fig. S5b). In contrast, none of the soil variables were

found to be significantly correlated with the dissimilarity of *E. coli* clonal groups from Hoosic River (*p*

> 0.05) (Fig. 3c). This result was consistent with our observation, based on VPA, that soil variables

minimally contributed to the distribution of *E. coli* clonal groups in the Hoosic River watershed.

The relationship between the dissimilarity of *E. coli* and geographic distance. We first

assessed the relationship between dissimilarity of *E. coli* clonal groups and geographic distance using

Mantel tests. Results showed a weak correlation between the dissimilarity of *E. coli* clonal groups and

geographic distance for Flint Creek at the 0.1 level (Mantel correlation coefficient r = 0.16, *p* = 0.08;

Table S7). By contrast, no significant correlation at the 0.1 level was observed between the dissimilarity

of *E. coli* clonal groups and geographic distance for Hoosic River (r = 0.11, *p* = 0.12; Table S7). Linear

regression analysis further showed that the slope of the regression line of the linear relationship between

the dissimilarity of *E. coli* clonal groups and geographic distance for Flint Creek (slope = 3.4 $\times 10^{-3}$, R^2 =

0.027, Fig. 4a) was about 3 times steeper than that for Hoosic River (slope = 9.7 $\times 10^{-4}$, R^2 = 0.011, Fig.

4b), indicating a stronger distance-decay relationship in *E. coli* in Flint Creek than Hoosic River.

Overall, results of Mantel test and linear regression analysis along with the VPA suggest that spatial

factors play a more important role in driving the distribution of *E. coli* clonal groups in Flint Creek than

Hoosic River, and the dispersal of *E. coli* in Flint Creek was slightly limited, while *E. coli* in the Hoosic

River was more likely not constrained by dispersal limitation.

To evaluate the impact of land-use on the relationship between *E. coli* and geographic distance,

we conducted Mantel tests and assessed the distance-decay relationship for produce field sites and forest

sites separately within each watershed. For Flint Creek, Mantel tests showed that the correlation between

the dissimilarity of *E. coli* clonal groups and geographic distance was not significant at the 0.1 level; the

correlation coefficient for forest sites (r = 0.27) was slightly larger than that for produce field sites (r =

0.22) (Table S8). For Hoosic River, the dissimilarity of *E. coli* clonal groups and geographic distance for

forest sites was significantly and highly correlated ($p < 0.05$, correlation coefficient = 0.49), while there

was no significant correlation for produce field sites ($p = 0.56$) (Table S8). Consistently, based on the

linear regression analysis assessing the distance-decay relationship, the slope of the linear regression line

for forest sites in both Flint Creek and Hoosic River (slope = 7.0×10^{-3} and 4.3×10^{-3}, respectively) was

steeper than that for produce field sites (slope = 5.1×10^{-3} and -3.0×10^{-4}, respectively) (Fig. S6a and

S6b). These results suggest that dispersal limitation for *E. coli* tend to be weaker in produce field sites

than in forest sites, which implies that dispersal of *E. coli* may be more efficient in the produce fields.

Wildlife-driven dispersal of *E. coli*. Four common classes of wildlife vehicles (large nuisance

wildlife species, small mammals, small flocking insectivore/granivores, and migratory bird flocks) were

selected for identifying potential dispersal vehicles of *E. coli* (characteristics of these dispersal vehicles

are detailed in Table 1). By adjusting distances among sampled sites to account for movement

preferences of these four types of wildlife vehicles (*i.e.*, cost-distance or landscape resistance modeling),

we sought to assess whether dispersal associated with wildlife behavior explains the *E. coli* distribution

better than distance alone. As shown in Table 2, the predicted dispersal model was developed based on

the most-likely cost and attraction models selected for each wildlife vehicle according to their

characteristics. The predicted dispersal model for small mammals was defined to have a biological

riparian corridor effect, no proximity effect, absolute dispersal barriers effect, and no attraction

coefficient. The predicted dispersal model for large nuisance wildlife species was defined to have a

biological riparian corridor effect, strong proximity effect, porous dispersal barriers effect, and habitat

quality coefficient. The predicted dispersal model for migratory bird flocks was defined to have a

biological riparian corridor effect, weak proximity effect, no dispersal barriers effect, and area

independent coefficient. The predicted dispersal model for small flocking insectivore/granivores was

defined to have a biological riparian corridor effect, weak proximity effect, absolute dispersal barriers

effect, and area independent coefficient. Definition of these effects can be found in Table 1.

Mantel tests showed that none of these dispersal models significantly predicted the composition

of *E. coli* clonal groups in Flint Creek ($p > 0.05$), while two wildlife-driven dispersal models – dispersal

via migratory bird flocks (M459) and via small flocking insectivore/granivores (M556) – were found to

be significantly correlated with the dissimilarity of *E. coli* clonal groups in Hoosic River (correlation

coefficient r = 0.17 and r = 0.16, respectively; $p < 0.05$) (Fig. 5). In addition, the model for dispersal via

large nuisance wildlife species (M377) was marginally significantly correlated with the dissimilarity of

E. coli clonal groups (r = 0.14, $p = 0.056$), so the role of dispersal of *E. coli* by large nuisance wildlife

species cannot be absolutely excluded. Thus, migratory bird flocks, small flocking

insectivore/granivores, and large nuisance wildlife species were identified as potential dispersal vehicles

which were associated with the distribution of *E. coli* in Hoosic River. The observation that cost-

distance model correlated with the dissimilarity of *E. coli* clonal groups in Hoosic River better than

geographic distance alone (Mantel correlation coefficient r = 0.11, $p = 0.12$; Table S7) suggests some

dispersal among sites by the action of wildlife. Our results also suggest that wildlife-driven dispersal

played a more important role in shaping the distribution of *E. coli* in Hoosic River as compared to Flint

Creek.

DISCUSSION

E. coli has widely been used as an indicator of fecal contamination (37) and potential presence of

other pathogenic enteric bacteria in water (38). *E. coli* comprises a wide spectrum of phenotypes

including harmless commensal variants as well as distinct pathotypes with the capacity to either cause

intestinal or extraintestinal infections in humans and many animals (39). The fecal-oral transmission

route of *E. coli* often involves transient presence in extra-host habitats (e.g., surface water, soil, plant

surfaces), including produce fields (23). Therefore, understanding the ecology of *E. coli* in extra-host

habitats will not only provide an improved understanding of *E. coli* interaction with environment, but

will also benefit public health by providing knowledge that can be used to minimizing introduction of *E.*

coli and possibly other enteric pathogens into preharvest environments.

Environmental stressors such as limited availability of nutrients and water, presence of toxic

molecules, and large alterations in temperature and moisture can impose fitness cost on *E. coli* and other

microbes (40). Fragmented landscapes with smaller forest and grassland patches expose surface soil to

sunlight and greatly increase daily variation in soil conditions. Reduced forest and grassland cover could

also hinder the movement of wildlife, bringing negative demographic and genetic consequences (41).

Thus, in order to disperse to and survive new habitats, *E. coli* needs to overcome those barriers by

maintaining variable survival strategies such as evolving adaptive traits relying on dispersal to rescue

local populations. In this scenario, landscape structure imposes constraints on environmental selection

and dispersal, which is particularly essential for the dispersal of *E. coli* among different extra-host

habitats.

To quantitatively probe the importance of environmental selection and dispersal in driving the

distribution and composition of *E. coli* in soil under the impact of landscape, we compared the

biogeographic patterns of *E. coli* isolated from two watersheds with distinct landscape patterns (i.e.,

Flint Creek, an area with widespread produce fields and limited interaction between produce fields and

forest, and Hoosic River, a heavily forested area with strong interaction between produce fields and

forest). Our data specifically suggests that in the watershed with widespread produce fields and sparse

forest coverage, environmental selection, possibly caused by soil phosphorus, and slightly limited

dispersal may result in relative heterogeneous composition of *E. coli* between produce field and forest

sites and potential local adaptation in *E. coli*. In contrast, in the watershed with heavily forested areas,

no evidence of environmental selection was observed, and dispersal facilitated by wildlife such as

migratory bird flocks and small flocking insectivore/granivores may enhance the likelihood of genetic

exchange among *E. coli* populations, resulting in relative homogeneous composition between produce

field sites and forest sites in this watershed. This higher level of homogeneity is consistent with greater

interaction between produce fields and forests in the Hoosic River watershed (adjacency$_{produce|forest}$ =

36%) compared to Flint Creek (adjacency$_{produce|forest}$ = 23%).

Agricultural practice involving input of phosphorous in soil may enhance the selective

pressure on *E. coli*. Agricultural activities normally involve cultivation and soil amendments, which

could dramatically change soil organic matter and nutrient pools in comparison to undisturbed systems

(e.g., forest) (42). Consequently, long-term organic and chemical amendments could dramatically

impact the abundance, diversity, and composition of bacterial communities in soil of agricultural land

(43). This is because such alteration of soil properties could trigger selective pressures on bacteria,

sorting the individuals or traits that better cope with modified soil condition, which has been termed

"local adaptation" (23). For example, copper-amendment in agricultural soil has been found to

significantly increase the frequency of copper-resistant Gram-negative bacteria (44). Based on the

results of our study reported here, agricultural practices may have caused selective pressure on soil *E.*

coli, partially resulting in the distinct *E. coli* composition between produce fields sites and forest sites in

the Flint Creek watershed. Consistent with our findings, Dusek et al. (22) observed *E. coli* population

structures that differed between cropland and forest, with much lower prevalence of *E. coli* in cropland

than forest. The diverse lifestyles and phenotypes of *E. coli* strains were thought to be caused by

population expansion paired with differential niche adaptation under specific selective pressures in the

last 5 million years (39). Our findings suggest that agriculture-stimulated selective pressures may

contribute to *E. coli* diversification.

Besides directly yielding selective pressures on *E. coli*, soil property alterations caused by

agricultural activities may also indirectly impact the adaptation of *E. coli* by changing the interaction

with other microbial taxa in the community. A wealth of studies have shown that anthropogenic

activities in agricultural land greatly change biological soil characteristics including the diversity,

structure, and metabolic guilds of microbial communties (45–48). For example, *E. coli* has been

reported to exhibit bacteriocin-mediated competitive interactions (49) and cooperative interactions using

cross-feeding metabolic products with other taxa in the microbial community (50). Changes in microbial

community structure and composition caused by agricultural activities thus possibly generate selective

pressures on *E. coli* through altering the competition and cooperation behaviors with other microbial

taxa.

The high correlation between phosphorous and antimony and the dissimilarity of *E. coli*

composition observed in the watershed with widespread produce fields (Flint Creek) suggest that the

alteration in these two soil variables may change the structure of *E. coli* populations in Flint Creek.

Phosphorus is one of the soil variables well documented to dramatically change after the conversion of

undisturbed systems to agriculture (42, 51). The input of phosphorus in fertilizer and manure to

agricultural systems have been reported to often exceed the output in harvested crops (52). Phosphorus

is a critical nutrient for the growth of bacteria and is part of many biomolecules in bacterial cells (e.g.,

DNA, phospholipids, polyphosphates, and ATP). Phosphorus availability could act as an important

selective force driving divergence among bacterial populations. For example, Coleman et al. (21)

identified a number of genes encoding functions related to phosphorus acquisition and metabolism (e.g.,

alkaline phosphatase, a pathway for phosphonate utilization, upregulation during phosphorus-starvation

conditions) as significantly enriched in *Prochlorococcus* populations in locations with lower phosphate

concentrations in North Atlantic and North Pacific subtropical gyres. Antimony is a toxic metalloid

present widely at trace concentrations in natural soil (53, 54). Its concentration could be elevated or even

reach contamination threshold in agricultural lands due to human activities (55). For example,

application of lead arsenate pesticides in produce field can increase antimony concentration, since

antimony is present as a contaminant in the antimony- and arsenic-containing ores used for pesticide

manufacturing (56). A previous study has shown that increased antimony could prevent the growth of *E.*

coli, *Bacillus subtilis* and *Staphylococcus aureus*, and may affect nitrogen cycle in soil by changing

urease activity under neutral pH (57). However, the concentration of antimony detected in this studied

was relatively low across all sites, and for some sites the concentration was below detection limit, thus

the presence of antimony may not necessarily inhibit the growth of *E. coli* in soil studied here.

Therefore, it is most likely that phosphorus represents agriculture-related selective pressures on *E. coli,*

though other physical soil parameters may have also played a role. In addition, the fact that this

correlation was overserved in Flint Creek but not in Hoosic River suggests that *E. coli* populations found

in different watersheds and environments may differ in their adaptive traits (e.g., those associated with

phosphorous).

E. coli in a watershed with high forest coverage may experience very weak selective

pressure and a proximity effect of forest. In this study, environmental selection tended to be very

weak on *E. coli* in watershed with higher forest coverage (Hoosic River). This relatively week

environmental selection might be because, compared to produce fields, plant cover and shading in forest

could moderate perturbations in soil moisture, nutrients and temperature, thus providing more favorable

and stable conditions with fewer environmental stressors for *E. coli* (4, 22). As previously proposed (4),

soil in undisturbed temperate forests could act as potential habitat for long-period persistent, even

resident *E. coli* populations rather than acting as a transient habitat. Albeit *E. coli* may be exposed to

fewer or less intense stressors in undisturbed environments, as compared to disturbed ones, some factors

such as temperature, moisture and nutrients have been shown to be correlated with *E. coli* density in

forest (58, 59). Due to lack of niche differentiation caused by environmental selection, we observed

more homogeneous *E. coli* compositions between forest and produce field in watershed with higher

forest coverage. We also observed that *E. coli* was much more prevalent in the watershed with higher

forest coverage (72%) as compared to the watershed with lower forest coverage (35%), consistent with

previous findings by Dusek et al. (22).

The higher prevalence of *E. coli* in the watershed with higher forest coverage might be caused by

proximity effect, which proposes that the likelihood of *E. coli* isolation from surrounding sites such as

produce field increases with the proximity to forests (22). Such a proximity effect is formed by the

spread of *E. coli* out of forests into surrounding areas, given that forest is a vital sink for *E. coli* (4). In

addition, the large adjacency between forest and produce fields in the watershed with higher forest

coverage, which indicates strong direct interactions between the two land covers, may enhance the

proximity effect. Consistent with our findings, Dusek et al. (22) reported that *E. coli* was more prevalent

in a landscape with greater forest coverage; this study specifically showed that *E. coli* was most

prevalent in soils sampled in close proximity (0 to 38 m) of forests, but was up to 90% less prevalent

when forest cover in the 250m radius was less than 7%. In addition to *E. coli*, such proximity effects of

forest have also been reported for *Listeria monocytogenes* and other *Listeria* species. Weller et al. (60)

found that with a 100m increase in the distance of a sampling site from forests, the likelihood of *L.*

monocytogenes and other *Listeria* species isolation in croplands decreased by 14% and 16%,

respectively.

Watershed landscape could constrain or facilitate the dispersal of soil *E. coli* by influencing

the movement of wildlife host. Wildlife, which is thought to be an important vehicle for the dispersal

of foodborne pathogens between hosts and locations (36), could enable bacteria to overcome landscape

barriers and make the dispersal of bacteria more active. Several studies have specifically indicated that

wildlife is a major source of *E. coli* in surface waters and may contribute to the contamination of *E. coli*

in rural watersheds and produce fields by defecation (11, 36, 61, 62). Landscape connectivity (i.e., the

degree to which a landscape facilitates or prevents movement of organisms among resource patches) and

particular landscape elements such as the structure of habitat (e.g., riparian corridor, terrestrial land,

waterbody) have also previously been shown to influence dispersal of pathogens (63, 64) and the

movement of wildlife (41). Wildlife-dependent dispersal of *E. coli* would thus be indirectly impacted by

landscape. Our results showed that the dispersal of soil *E. coli* in the watershed with widespread produce

fields was slightly limited, while the dispersal of soil *E. coli* in the watershed with high forest coverage

tended to be largely facilitated by wildlife, highlighting the important role of watershed landscapes on

the dispersal of *E. coli*. These influences may consequently shape the spatial patterns of pathogenic *E.*

coli persistence and incidence (65). Consistent with our findings, Mechai et al. (66) showed evidence of

the impact of landscape connectivity on the dispersal patterns of *Borrelia burgdorferi*, particularly

rodent-associated strains, which is relevant to the spread of Lyme disease risk across locations.

Based on the above notions, our observation that the dispersal of soil *E. coli* in the watershed

with widespread produce fields was relatively limited could be explained by a combination of

environmental constraints (i.e., relatively strong environmental selection) in soil and poor connectivity

of agricultural areas, which may impede the movement of wildlife that disperses *E. coli* (41). By

contrast, the dispersal of soil *E. coli* in the watershed with high forest coverage tended to be promoted

by wildlife. This may be because forest exhibits better connectivity and could provide passage and

support to the movement of wildlife vehicles of *E. coli*. Besides wildlife, it is also possible that the

dispersal of *E. coli* was directly influenced by the landscape elements of the two watersheds. Forest and

most produce fields in Hoosic River, which is heavily forested, were both located in a floodplain. By

contrast, forest in Flint Creek was in a floodplain but produce fields were not. Since during periods of

high discharge, a floodplain normally experiences flooding, such events may facilitate the dispersal of *E.*

coli between forest and produce field particularly in the Hoosic River watershed. This hypothesis is

supported by a number of modelling studies, showing that the peak fecal bacteria levels during flooding

can be more than 20 or even 50 times higher than prior to flooding (67–69). Future studies on comparing

the distribution of *E. coli* before, during, and after flooding and assessing the correlation between

flooding-associated landscape factors (e.g., elevation and patchiness) and distribution of *E. coli* are

needed for an improved understanding of the impact of landscape on the microbial biogeography.

Migratory bird flocks, small flocking insectivore/granivores, and large nuisance wildlife were

identified, in our study here, as potential vehicles that disperse *E. coli* and affect the distribution of *E.*

coli in the watershed with high forest coverage. Migratory bird flocks (e.g., Canada goose) tend to have

low movement cost in all land-use types except for heavy urban development areas (70). Small flocking

insectivore/granivores (e.g., European starling) tend to have low movement cost in all land-use types

except for open water and heavy urban development areas (70). Large nuisance wildlife (e.g., white

tailed deer or feral swine) tend to have low movement cost in forests, scrublands, grasslands, wooded

wetlands and cultivated croplands, while they can have high movement cost to cross roads (71). Each of

these three classes of wildlife have been previously reported (34, 35, 72) to serve as dispersal vehicles of

E. coli and have been considered public health concerns in terms of agricultural contamination. For

example, a study of *E. coli* isolated from fecal samples of Canada geese over a year in Colorado, US

reported that the prevalence for *E. coli* ranged from 2% during the coldest months to 94% during the

warmest time of the year (72). European starlings, which are considered an invasive species in the

United States and a nuisance pest to agriculture, were proposed to be a potential suitable reservoir and

vector of *E. coli* O157:H7, and can carry and disseminate this human pathogen to cattle (34). In addition,

deer feces were reported to contaminate fresh strawberries, being responsible for an outbreak of *E. coli*

O157:H7 infections in Oregon (35). Forest is a relative stable environment with less disturbance of

anthropologic activities, thus serving as an ideal living habitat for wild animals (73). Forest may provide

easy transport pathways for small flocking insectivore/granivores and large nuisance wildlife to move

around and support high density of migratory birds, which increases the chance of colonized *E. coli*

dispersed from forest to adjacent produce fields.

Conclusion. By comparing the biogeographic patterns of *E. coli* isolated from two watershed

with distinct landscape characteristics in New York state, we showed that terrestrial landscape could

impact the distribution of *E. coli* by adjusting the importance of environmental selection and dispersal.

Environmental stress was identified as a possible strong contributor to local adaptation of *E. coli* in the

watershed with widespread produce fields. On the other hand, wildlife-driven dispersal, which could

facilitate genetic exchange, was identified as a major force in shaping *E. coli* populations in the

watershed with high forest coverage. As such, our findings not only highlight the critical role of

landscape in driving the biogeographic pattern of *E. coli* in perspective of ecology, but also open the

possibility that the evolutionary forces (e.g., positive selection, genetic drift, gene flow) driving the

diversification of *E. coli* vary by watershed landscape as well. In addition, our study suggests that due to

the less intense environmental stress, frequent wildlife-facilitated dispersal, and the proximity effect of

forest on *E. coli*, produce fields in watershed with high forest coverage may have higher risk in *E. coli*

contamination. This information can inform spatial modeling of food contamination risk associated with

produce fields in different watersheds, which can be used to modify pre-harvest product sampling

strategies and produce harvest methods to account for the spatial structure in contamination risk in a

produce field. Despite the contributions to the field of microbial ecology, our study has some non-

negligible limitations. First, we did not differentiate commensal and pathogenic *E. coli*; however,

biogeographic patterns and the impact of landscape may differ between the two groups. Second, we only

assessed the coverage of forest and produce field as one landscape attribute using a relatively small set

of sampling sites, while many other landscape attributes such as the size, diversity, and richness of patch

size may also be important to the biogeographic pattern and adaptation of *E. coli*. Third, wildlife

population/community structure, which could be strongly affected by land-use features (74), was not

included in the competing models. Future studies warrant more intensive sampling efforts, sequencing

techniques with higher discriminatory power (e.g., whole genome sequencing), and comprehensive

assessment of a wider range of high order landscape attributes and wildlife characteristics to better

understand the impact of the landscape on the biographic pattern and adaptation of *E. coli*. Such

methodology development, ideally validated by experimental data, could considerably improve

prediction of produce contamination risk based on the potential influence of landscape on the dispersal

of *E. coli* to produce field, benefit the development of trade-off risk assessments of food contamination,

and eventually help to decrease human exposure to pathogenic enteric bacteria.

MATERIALS AND METHODS

Study sites and soil collection. Two watersheds with different landscape patterns, Flint Creek

and Hoosic River, located in the New York state, were selected for this study based on topography and

land-cover composition. Flint Creek is an area with widespread vegetable and livestock production that

is sparsely forested (69% produce field, 12% forest by area), whereas the Hoosic River watershed is a

heavily forested area with interspersed produce production (28% produce field, 38% forest by area). Soil

sampling was carried out between September 4 and October 10, 2012 on 7 farms comprising 19 produce

field sites and in 16 forest sites (Fig. 1). For produce fields, two parallel 200 m transects were laid in

each field, perpendicular to the forest boundary. Along each transect, five soil samples (at approximately

5 cm-depth) were collected at 50 m intervals using sterile scoops (Fisher Scientific, Hampton, NH) and

sterile Whirl-Pak bags (Nasco, Fort Atkinson, WI). Latex gloves and disposable plastic boot covers

(Nasco, Fort Atkinson, WI) were worn for sample collection. Gloves and boot covers were changed

between each site, and gloves were disinfected with 70% ethanol prior to sample collection. A total of

278 soil samples were collected with 142 and 136 samples collected from the Flint Creek and the Hoosic

River watershed, respectively. All samples were transported to the Food Safety Lab at Cornell

University in an icebox. Samples were stored at $4 \pm 2°C$ in dark and processed within 24 h of collection.

Isolation of *E. coli*. *E. coli* were isolated from soil samples as previously described (23). Briefly,

8 g sieved soil was diluted 1:10 in EC medium with 4-methylumbelliferyl-β,D-glucuronide broth (EC-

MUG). To maximize genetic diversity among recovered *E. coli* isolates, the suspension was subdivided

among four 96-well microtiter plates for a total of 384 subsamples of approximately 180 uL each.

Microtiter plates were incubated at 37°C. Bacteria from fluorescent wells were isolated on EC-MUG

agar plates and were further tested with a standard biochemical assay for glutamate decarboxylase and

beta-glucuronidase activity. Isolates that were positive for these two tests were presumptively identified

as *E. coli*, which were confirmed by subsequent gene sequencing as detailed below. No *E. coli* isolates

were detected in samples from 3 produce sites (Field 7, Field 9, and Field 10), and 2 forest sites (Forest

F4 and Forest F18).

DNA extraction, MLST genotyping and clonal groups. Genomic DNA was extracted from *E.*

coli by alkaline lysis of biomass in 50 mM NaOH at 95°C. Two genes (*mdh* and *uidA*) were sequenced

first from all isolates. Then, only the unique two-gene sequence types from each sample were subjected

to additional sequencing in five additional genes (*aspC*, *clpX*, *icd*, *lysP*, *fadD*) by Sanger sequencing,

performed by the Cornell University Life Sciences Core Laboratories Center. Evaluation of sequence

read quality and assembly of forward and reverse reads were performed using Perl scripts, which

iterated runs of phred and CAP3, respectively. Sequences with a probability of error of > 0.005 (Q score

< 23) in terms of read quality were edited manually, where possible, or discarded. Assembled sequences

of each MLST locus were aligned and trimmed to standard base positions matching the *E. coli* K-12

sequence type from the STEC Center website (http://www.shigatox.net) (23). Alignments of assembled

sequences for isolates from Flint Creek and Hoosic River are available on GitHub

(https://github.com/pbergholz/Dispersal-cost-modeling). The clonal groups of *E. coli* strains for Flint

Creek and Hoosic River were determined based on MLST sequence types using the goeBURST analysis

program (75) at single locus variant level.

Remotely sensed data and soil property data. GPS coordinates of sites were imported into the

Geographical Resources Analysis Support System (GRASS) geographic information system (GIS)

environment. Map layers for land cover (National Land Cover Database [NLCD], 2006) and the digital

elevation model (DEM; Shuttle Radar Topography Mission, 1-arc-second data set) were acquired from

the U.S. Geological Survey (USGS) Earth Explorer geographical data bank

(http://earthexplorer.usgs.gov/). Map layers for soil characteristics were acquired from the U.S.

Department of Agriculture Soil Survey Geographic (SSURGO) database

(http://soils.usda.gov/survey/geography/ssurgo/). Road and hydrologic line graphs were obtained from

the Cornell University Geospatial Information Repository (CUGIR; http://cugir.mannlib.cornell.edu/).

Percent landcover and adjacency were estimated by using FRAGSTATS v. 3.3 to analyze

landcover within a 2 km buffer surrounding the Flint Creek and Hoosic River, respectively (76). All

land NLCD maps identified as pasturage was reclassified as cropland for our analyses. Percent

adjacency was calculated as the proportion pixels in the NLCD map that were adjacent forest and

produce field, compared to the total of non-self-adjacencies in 2 km buffer surrounding the waterway.

For example, adjacency$_{produce|forest}$ = 10% would indicate that 10% of the edges of produce fields abutted

forest in a given area.

Organic matter, moisture, pH, aluminium, arsenic, boron, barium, calcium, cadmium, cobalt,

chromium, copper, iron, potassium, magnesium, manganese, molybdenum, sodium, nickel, phosphorus,

lead, sulphur, strontium, and zinc content of soil samples were measured at Cornell Nutrient Analysis

Lab.

Distribution of *E. coli* clonal groups and its relationship with geographic location. The

Mann-Whitney test was used to determine if number of clonal groups differed significantly between soil

samples from produce field sites and forest sites for Flint Creek and Hoosic River. Principal coordinate

analysis (PCoA) was conducted using phyloseq package in R 3.6.0 to visualize the distribution of *E. coli*

clonal groups among sites, based on Bray-Curtis distance. The 95% confidence ellipse in the PCoA plot

assumes a multivariate normal distribution. Permutational multivariate analysis of variance

(PERMANOVA) (77) was employed using the adonis function in R 3.6.0's vegan package to test

whether the centroids and dispersion of sample groups as defined by land-use (produce field or forest)

are equivalent for both groups based on Bray-Curtis distance of *E. coli* clonal groups. PERMANOVA

test statistic (F) and *p*-value were obtained by 9,999 permutations. Analysis of similarities (ANOSIM)

(77) was employed using the anosim function in R 3.6.0's vegan package to test whether there is a

significant difference between two groups (produce field sties and forest sites) of sampling units based

on the Bray-Curtis distance of *E. coli* clonal groups. ANOSIM test statistic (R) and p-value were

obtained by 9,999 permutations. To test the sampling bias which possibly caused the large variation of *E.*

coli clonal groups observed across sampling sites within each watershed, sites with a number of clonal

groups ≤ 3 (Field 6, Field 8, and Forest F9) were excluded and this subset of samples were repeated for

PCoA, PERMANOVA, and ANOSIM analyses.

Mantel tests were performed using vegan package in R 3.6.0 to assess the relationship between

the biological dissimilarity of *E. coli* and geographic distance (9,999 permutations). Biological

dissimilarity of *E. coli* clonal groups was calculated in Bray-Curtis distance using vegan package in R

3.6.0. Geographic distance between isolates was calculated from latitude and longitude, using the geopy

module in Python 3.6.8. Linear regression analysis of biological dissimilarity of *E. coli* and geographic

distance was performed in R 3.6.0. Distance-decay relationship was inferred from the slope and R^2 of

the linear regression. A steeper slope with a larger R^2 value suggests stronger distance-decay

relationship.

The relationship between *E. coli* clonal groups and soil variables. Due to a lack of mineral

soil to measure soil property after combusting away the organic matter, Forest F6, Forest F11, and

Forest F16 were not included in analyses on the relationship between *E. coli* clonal groups and soil

variables. After screening for covariation, soil variables with low levels of covariation ($r < 0.7$ and $p <$

0.05 in Pearson's correlation analysis) were selected for variation partitioning analysis (VPA) using

vegan package in R 3.6.0 to quantify the relative contribution of the environment effect and the

geographical effect on the dissimilarity of *E. coli* clonal groups based on Bray-Curtis distance (78).

Principal coordinates of neighbor matrices (PCNM) were used to represent spatial patterns based on

GPS coordinates (79). By using 'ordistep' function in vegan package, a subset of PCNM variables

which significantly explained variation in the dissimilarity of *E. coli* clonal groups was included in VPA.

Four components of variations were calculated in VPA: i) pure contribution of environmental effect

($R^2_A - R^2_G$); ii) pure geographical effect ($R^2_A - R^2_E$); iii) spatially structured environmental effect ($R^2_G +$

$R^2_E - R^2_A$); and iv) unexplained effect ($1 - R^2_A$). R^2_A, R^2_G, and R^2_E represent variation of the dissimilarity

of *E. coli* clonal groups explained by all variables, spatial variables, and environmental variables,

respectively.

Partial Mantel test was performed to examine the correlation between environmental

dissimilarity of each soil variable and the dissimilarity of *E. coli* clonal groups independent of

geographical influence using vegan package in R 3.6.0 (9,999 permutations). Dissimilarity of *E. coli*

clonal groups was calculated in Bray-Curtis distance, and environmental dissimilarity was calculated in

Euclidian distance. Soil variables with a *p* value < 0.05 in partial Mantel tests were defined as key soil

variables. Mann-Whitney tests were further performed to determine if key soil variables differed

significantly between soil samples from produce field sites and forest sites.

Dispersal model formulation and selection. To predict the dispersal of *E. coli* across watershed

landscapes, multiple dispersal models were developed to describe landscape effects by integrating

remotely sensed and field-collected data into resistance surfaces for wildlife vehicles. Four common

classes of wildlife vehicles including (i) large nuisance wildlife species, (ii) small mammals, (iii) small

flocking insectivore/granivores, and (iv) migratory bird flocks were selected in this study.

Predicted dispersal among sites was calculated according to the equation below:

$$D_{i,j} = \frac{L_i \times A_j}{C_{i,j}}$$

Where $D_{i,j}$ is the dispersal rate among sites *i* and *j*, L_i is the *E. coli* load from the source site (i.e.,

starting point), A_j is the attraction (gravity) coefficient of the sink site (i.e., stopping point) to vehicle

and $C_{i,j}$ is the least-cost distance between sites *i* and *j*. *E. coli* load L_i expresses the expected mobility of

E. coli from these areas as a function of expected prevalence. Expected prevalence was inferred from

random forest analysis of *E. coli* prevalence based on sampling excursions. One load map was generated

per watershed. The attraction (gravity) coefficient A_j describes the tendency of a dispersal vehicle to

move towards an area on the landscape and expected residence-time of dispersal vehicle after they arrive

at a location. Attraction was primarily a function of percent favored land-cover for each of the vehicles

and interspersion of land-cover types. The least-cost distance $C_{i,j}$ describes the movement preferences of

a dispersal vehicle in terms of a friction surface (borrowed from circuit theory) that predicts resistance of

the landscape to movement of dispersal vehicles. The cost surfaces were a function of baseline

resistance (dependent on the dispersal vehicle), riparian corridor effect (i.e., the tendency of wildlife to

prefer movement through riparian forests), dispersal barrier effect (i.e., the strength of barriers to

movement, such as major road- and water-ways), and proximity effect (i.e., the strength and type of

edge interactions among forests, produce fields, pasturage, and urban areas). The least-cost distance was

measured as the distance along the path that accrued the least cumulative cost between pairs of

movement start and stop sites. The characteristics of the dispersal vehicles, *E. coli* load model (L_i),

attraction model (coefficient A_j), cost model (i.e., riparian corridor effect, dispersal barrier effect and

proximity effect) were shown in Table 1, which were summarized on the basis of published literature

(73–84). Based on these characteristics, the most-likely attraction model and cost model were selected

for each class of vehicle, generating the predicted dispersal models (Table 2).

For each of the predicted dispersal models for the four classes of wildlife vehicles, an association

matrix $D_{i,j}$ containing predicted dispersal rates along least cost paths among all pairs of sites was

generated. This was accomplished by using a set of scripts developed in the GRASS GIS ver. 6.4.3

programming environment; Perl scripts were used to automate calculations in the GIS; scripts are

available on GitHub (https://github.com/pbergholz/Dispersal-cost-modeling). Mantel tests were

employed to estimate the correlation between predicted dispersal models and biological dissimilarity of

E. coli clonal groups among sampled sites in each watershed using R version 3.6.0. Statistical

significance of model fits was estimated by 9,999 permutations. The wildlife vehicle for which the

predicted model had the highest significant correlation coefficient was deemed to represent the dominant

dispersal vehicle for *E. coli*.

Acknowledgements: This research was supported by the Center for Produce Safety (research agreement

number 201121642-01, representing a subcontract under Award Number SCB11072 from the California

Department of Food and Agriculture).

REFERENCES

1.   Berges SA, Moore LAS, Isenhart TM, Schultz RC. 2010. Bird species diversity in riparian buffers,

row crop fields, and grazed pastures within agriculturally dominated watersheds. Agrofor Syst

79:97–110.

2.   Borin M, Passoni M, Thiene M, Tempesta T. 2010. Multiple functions of buffer strips in farming

areas. Eur J Agron 32:103–111.

3.   Finder RA, Roseberry JL, Woolf A. 1999. Site and landscape conditions at white-tailed

deer/vehicle collision locations in Illinois. Landsc Urban Plan 44:77–85.

4.   Byappanahalli MN, Whitman RL, Shively DA, Sadowsky MJ, Ishii S. 2006. Population structure,

persistence, and seasonality of autochthonous *Escherichia coli* in temperate, coastal forest soil

from a Great Lakes watershed. Environ Microbiol 8:504–513.

5.   Mostaghimi S, Benham B, Wolfe ML. BSLC: A tool for bacterial source characterization for

watershed management. Appl Eng Agric.

6.   Kouznetsov MY, Roodsari R, Pachepsky YA, Shelton DR, Sadeghi AM, Shirmohammadi A,

Starr JL. 2007. Modeling manure-borne bromide and fecal coliform transport with runoff and

infiltration at a hillslope. J Environ Manage 84:336–346.

7.   Lowe WH, McPeek MA. 2014. Is dispersal neutral? Trends Ecol Evol. Elsevier Ltd.

8.   Oliver DM, Heathwaite AL, Fish RD, Chadwick DR, Hodgson CJ, Winter M, Butler AJ. 2009.

Scale appropriate modelling of diffuse microbial pollution from agriculture. Prog Phys Geogr

Earth Environ 33:358–377.

9.   Walsh CJ, Kunapo J. 2009. The importance of upland flow paths in determining urban effects on

stream ecosystems. J North Am Benthol Soc 28:977–990.

10.   Wang H, Zhang T, Wei G, Wu L, Wu J, Xu J. 2014. Survival of *Escherichia coli* O157:H7 in

soils under different land use types. Environ Sci Pollut Res 21:518–524.

11.   Harmel RD, Karthikeyan R, Gentry T, Srinivasan R. Effects of agricultural management, land use,

and watershed scale on *E. coli* concentrations in runoff and streamflow. Trans ASABE 53:1833–

1841.

12.   Foster DR, Motzkin G, Bernardos D, Cardoza J. 2002. Wildlife dynamics in the changing New

England landscape. J Biogeogr 29:1337–1357.

13.   Gibbons SM. 2017. Metapopulation theory provides new insight into microbial biogeography.

Environ Microbiol 19:849–850.

14.   Nemergut DR, Costello EK, Hamady M, Lozupone C, Jiang L, Schmidt SK, Fierer N, Townsend

AR, Cleveland CC, Stanish L, Knight R. 2011. Global patterns in the biogeography of bacterial

taxa. Environ Microbiol 13:135–144.

15.   Nemergut DR, Schmidt SK, Fukami T, O'Neill SP, Bilinski TM, Stanish LF, Knelman JE, Darcy

JL, Lynch RC, Wickey P, Ferrenberg S. 2013. Patterns and processes of microbial community

assembly. Microbiol Mol Biol Rev 77:342–356.

16.   Liao J, Cao X, Wang J, Zhao L, Sun J, Jiang D, Huang Y. 2017. Similar community assembly

mechanisms underlie similar biogeography of rare and abundant bacteria in lakes on Yungui

Plateau, China. Limnol Oceanogr 62:723–735.

17.   Yashiro E, Pinto-Figueroa E, Buri A, Spangenberg JE, Adatte T, Niculita-Hirzel H, Guisan A,

van der Meer JR. 2016. Local environmental factors drive divergent grassland soil bacterial

communities in the western Swiss Alps. Appl Environ Microbiol 82:6303–6316.

18.   Lozupone CA, Knight R. 2007. Global patterns in bacterial diversity. Proc Natl Acad Sci U S A

104:11436–11440.

19.   Semenov A V., Franz E, Van Overbeek L, Termorshuizen AJ, Van Bruggen AHC. 2008.

Estimating the stability of *Escherichia coli* O157:H7 survival in manure-amended soils with

different management histories. Environ Microbiol 10:1450–1459.

20.   Liao J, Wang J, Huang Y. 2015. Bacterial community features are shaped by geographic location,

physicochemical properties, and oil contamination of soil in main oil fields of China. Microb Ecol

70:380–389.

21.   Coleman ML, Chisholm SW. 2010. Ecosystem-specific selection pressures revealed through

comparative population genomics. Proc Natl Acad Sci U S A 107:18634–18639.

22.   Dusek N, Hewitt AJ, Schmidt KN, Bergholz PW. 2018. Landscape-scale factors affecting the

prevalence of *Escherichia coli* in surface soil include land cover type, edge interactions, and soil

pH. Appl Environ Microbiol 84:e02714-17.

23.   Bergholz PW, Noar JD, Buckley DH. 2011. Environmental patterns are imposed on the

population structure of *Escherichia coli* after fecal deposition. Appl Environ Microbiol 77:211–

219.

24.   Fierer N, Jackson RB. 2006. The diversity and biogeography of soil bacterial communities. Proc

Natl Acad Sci U S A 103:626–631.

25.   Forbes AE, Chase JM. 2002. The role of habitat connectivity and landscape geometry in

experimental zooplankton metacommunities. Oikos 96:433–440.

26.   Berga M, Östman Ö, Lindström ES, Langenheder S. 2015. Combined effects of zooplankton

grazing and dispersal on the diversity and assembly mechanisms of bacterial metacommunities.

Environ Microbiol 17:2275–2287.

27. Darcy JL, Lynch RC, King AJ, Robeson MS, Schmidt SK. 2011. Global distribution of

Polaromonas phylotypes - Evidence for a highly successful dispersal capacity. PLoS One

6:e23742.

28. Biek R, Real LA. 2010. The landscape genetics of infectious disease emergence and spread. Mol

Ecol 19:3515–3531.

29. Cushman SA, McKelvey KS, Hayden J, Schwartz MK. 2006. Gene flow in complex landscapes:

testing multiple hypotheses with causal modeling. Am Nat 168:486–99.

30. Storfer A, Murphy MA, Evans JS, Goldberg CS, Robinson S, Spear SF, Dezzani R, Delmelle E,

Vierling L, Waits LP. 2007. Putting the "landscape" in landscape genetics. Heredity (Edinb).

Nature Publishing Group.

31. Jang J, Hur HG, Sadowsky MJ, Byappanahalli MN, Yan T, Ishii S. 2017. Environmental

Escherichia coli: ecology and public health implications—a review. J Appl Microbiol 123:570–

581.

32. Nandakafle G, Seale T, Flint T, Nepal M, Venter SN, Brözel VS. 2017. Distribution of diverse

Escherichia coli between cattle and pasture. Microbes Environ 32:226–233.

33. Chandler DS, Craven JA. 1980. Relationship of soil moisture to survival of *Escherichia coli* and

Salmonella typhimurium in soils. Aust J Agric Res 31:547–555.

34. Kauffman MD, LeJeune J. 2011. European Starlings (*Sturnus vulgaris*) challenged with

Escherichia coli O157 can carry and transmit the human pathogen to cattle. Lett Appl Microbiol

53:596–601.

35. Laidler MR, Tourdjman M, Buser GL, Hostetler T, Repp KK, Leman R, Samadpour M, Keene

WE. 2013. *Escherichia coli* O157:H7 infections associated with consumption of locally grown

strawberries contaminated by deer. Clin Infect Dis 57:1129–1134.

36.   Avery SM, Moore A, Hutchison ML. 2004. Fate of *Escherichia coli* originating from livestock

faeces deposited directly onto pasture. Lett Appl Microbiol 38:355–359.

37.   Byappanahalli M, Fowler M, Shively D, Whitman R. 2003. Ubiquity and persistence of

Escherichia coli in a Midwestern Coastal Stream. Appl Environ Microbiol 69:4549–4555.

38.   Bolster CH, Haznedaroglu BZ, Walker SL. 2009. Diversity in cell properties and transport

behavior among 12 different environmental *Escherichia coli* isolates. J Environ Qual 38:465–472.

39.   Leimbach A, Hacker J, Dobrindt U. 2013. *E. coli* as an all-rounder: The thin line between

commensalism and pathogenicity. Curr Top Microbiol Immunol 358:3–32.

40.   Ramos JL, Gallegos MT, Marqués S, Ramos-González MI, Espinosa-Urgel M, Segura A. 2001.

Responses of gram-negative bacteria to certain environmental stressors. Curr Opin Microbiol

4:166–171.

41.   Shepard DB, Kuhns AR, Dreslik MJ, Phillips CA. 2008. Roads as barriers to animal movement in

fragmented landscapes. Anim Conserv 11:288–296.

42.   Compton JE, Boone RD. 2000. Long-term impacts of agriculture on soil carbon and nitrogen in

New England forests. Ecology 81:2314–2330.

43.   Chaudhry V, Rehman A, Mishra A, Chauhan PS, Nautiyal CS. 2012. Changes in bacterial

community structure of agricultural land due to long-term organic and chemical amendments.

Microb Ecol 64:450–460.

44.   Berg J, Tom-Petersen A, Nybroe O. 2005. Copper amendment of agricultural soil selects for

bacterial antibiotic resistance in the field. Lett Appl Microbiol 40:146–151.

45.   Jangid K, Williams MA, Franzluebbers AJ, Sanderlin JS, Reeves JH, Jenkins MB, Endale DM,

Coleman DC, Whitman WB. 2008. Relative impacts of land-use, management intensity and

fertilization upon soil microbial community structure in agricultural systems. Soil Biol Biochem

40:2843–2853.

46.   Buckley DH, Schmidt TM. 2001. The structure of microbial communities in soil and the lasting

impact of cultivation. Microb Ecol 42:11–21.

47.   Øvreås L, Torsvik V. 1998. Microbial diversity and community structure in two different

agricultural soil communities. Microb Ecol 36:303–315.

48.   Enwall K, Nyberg K, Bertilsson S, Cederlund H, Stenström J, Hallin S. 2007. Long-term impact

of fertilization on activity and composition of bacterial communities and metabolic guilds in

agricultural soil. Soil Biol Biochem 39:106–115.

49.   Majeed H, Gillor O, Kerr B, Riley MA. 2011. Competitive interactions in Escherichia coli

populations: The role of bacteriocins. ISME J 5:71–81.

50.   Harcombe W. 2010. Novel cooperation experimentally evolved between species. Evolution (N Y)

64:2166–2172.

51.   Cambardella CA, Elliott ET. 1994. Carbon and nitrogen dynamics of soil organic matter fractions

from cultivated grassland soils. Soil Sci Soc Am J 58:123–130.

52.   Sharpley AN. 1995. Soil phosphorus dynamics: agronomic and environmental impacts. Ecol Eng

5:261–279.

53.   Li J, Wang Q, Oremland RS, Kulp TR, Rensing C, Wang G. 2016. Microbial antimony

biogeochemistry: Enzymes, regulation, and related metabolic pathways. Appl Environ Microbiol

82:5482–5495.

54.   Steely S, Amarasiriwardena D, Xing B. 2007. An investigation of inorganic antimony species and

antimony associated with soil humic acid molar mass fractions in contaminated soils. Environ

Pollut 148:590–598.

55.   Tóth G, Hermann T, Da Silva MR, Montanarella L. 2016. Heavy metals in agricultural soils of

725   the European Union with implications for food safety. Environ Int 88:299–309.

726   56.   Wagner SE, Peryea FJ, Filby RA. 2003. Antimony impurity in lead arsenate insecticide enhances

727   the antimony content of old prchard soils. J Environ Qual 32:736–738.

728   57.   An YJ, Kim M. 2009. Effect of antimony on the microbial growth and the activities of soil

729   enzymes. Chemosphere 74:654–659.

730   58.   Carrillo M, Estrada E, Hazen TC. 1985. Survival and enumeration of the fecal indicators

731   Bifidobacterium adolescentis and *Escherichia coli* in a tropical rain forest watershed. Appl

732   Environ Microbiol 50:468–476.

733   59.   Tate RL. 1978. Cultural and environmental factors affecting the longevity of *Escherichia coli* in

734   Histosols. Appl Environ Microbiol 35:925–9.

735   60.   Weller D, Shiwakoti S, Bergholz P, Grohn Y, Wiedmann M, Strawn LK. 2016. Validation of a

736   previously developed geospatial model that predicts the prevalence of *Listeria monocytogenes* in

737   New York State produce fields. Appl Environ Microbiol 82:797–807.

738   61.   Casarez EA, Pillai SD, Mott JB, Vargas M, Dean KE, Di Giovanni GD. 2007. Direct comparison

739   of four bacterial source tracking methods and use of composite data sets. J Appl Microbiol

740   103:350–364.

741   62.   Parajuli PB, Mankin KR, Barnes PL. 2009. Source specific fecal bacteria modeling using soil and

742   water assessment tool model. Bioresour Technol 100:953–963.

743   63.   Tischendorf L, Fahrig L. 2000. On the usage and measurement of landscape connectivity. Oikos

744   90:7–19.

745   64.   Plantegenest M, Le May C, Fabre F. 2007. Landscape epidemiology of plant diseases. J R Soc

746   Interface 4:963–972.

65.   Thrall PH, Burdon JJ. 1997. Host-pathogen dynamics in a metapopulation context: The ecological

and evolutionary consequences of being spatial. J Ecol 85:743.

66.   Mechai S, Margos G, Feil EJ, Lindsay LR, Michel P, Kotchi SO, Ogden NH. 2018. Evidence for

an effect of landscape connectivity on *Borrelia burgdorferi sensu stricto* dispersion in a zone of

range expansion. Ticks Tick Borne Dis 9:1407–1415.

67.   Wilkinson J, Jenkins A, Wyer M, Kay D. 1995. Modelling faecal coliform dynamics in streams

and rivers. Water Res 29:847–855.

68.   Muirhead RW, Davies-Colley RJ, Donnison AM, Nagels JW. 2004. Faecal bacteria yields in

artificial flood events: Quantifying in-stream stores. Water Res 38:1215–1224.

69.   Bai S, Lung WS. 2005. Modeling sediment impact on the transport of fecal bacteria. Water Res

39:5232–5240.

70.   Shepard ELC, Wilson RP, Rees WG, Grundy E, Lambertucci SA, Vosper SB. 2013. Energy

landscapes shape animal movement ecology. Am Nat 182:298–312.

71.   Putman RJ. 1997. Deer and road traffic accidents: Options for management. J Environ Manage

51:43–57.

72.   Kullas H, Coles M, Rhyan J, Clark L. 2002. Prevalence of Escherichia coli serogroups and human

virulence factors in faeces of urban Canada geese (Branta canadensis). Int J Environ Health Res

12:153–162.

73.   Robinson SK, Yahner RH. 1997. Eastern deciduous forests: Ecology and wildlife conservation. J

Wildl Manage 61:1445.

74.   Coulon A, Guillot G, Cosson JF, Angibault JMA, Aulagnier S, Cargnelutti B, Galan M, Hewison

AJM. 2006. Genetic structure is influenced by landscape features: Empirical evidence from a roe

deer population. Mol Ecol 15:1669–1679.

75. Francisco AP, Bugalho M, Ramirez M, Carriço JA. 2009. Global optimal eBURST analysis of

multilocus typing data using a graphic matroid approach. BMC Bioinformatics 10:152.

76.   FRAGSTATS: Spatial Pattern Analysis Program for Categorical Maps.

77. Anderson MJ, Walsh DCI. 2013. PERMANOVA, ANOSIM, and the Mantel test in the face of

heterogeneous dispersions: What null hypothesis are you testing? Ecol Monogr 83:557–574.

78. Borcard D, Legendre P, Drapeau P. 1992. Partialling out the Spatial Component of Ecological

Variation. Ecology 73:1045–1055.

79. Dray S, Legendre P, Peres-Neto PR. 2006. Spatial modelling: a comprehensive framework for

principal coordinate analysis of neighbour matrices (PCNM). Ecol Modell 196:483–493.

80. Bruun M, Smith HG. 2003. Landscape composition affects habitat use and foraging flight

distances in breeding European starlings. Biol Conserv 114:179–187.

81. Minderman J, Reid JM, Hughes M, Denny MJH, Hogg S, Evans PGH, Whittingham MJ. 2010.

Novel environment exploration and home range size in starlings *Sturnus vulgaris*. Behav Ecol

21:1321–1329.

82. Getz LL, Mcguire B. 2008. Factors influencing movement distances and home ranges of the

short-tailed shrew (*Blarina brevicauda*). Northeast Nat 15:293–302.

83. Wang M, Grimm V. 2007. Home range dynamics and population regulation: An individual-based

model of the common shrew *Sorex araneus*. Ecol Modell 205:397–409.

84. Morales JM, Moorcroft PR, Matthiopoulos J, Frair JL, Kie JG, Powell RA, Merrill EH, Haydon

DT. 2010. Building the bridge between animal movement and population dynamics. Philos Trans

R Soc B Biol Sci 365:2289–2301.

85. Brandl MT. 2006. Fitness of human enteric pathogens on plants and implications for food safety.

Annu Rev Phytopathol 44:367–392.

86.   Blanchong JA, Samuel MD, Scribner KT, Weckworth B V, Langenberg JA, Filcek KB. 2008.

794   Landscape genetics and the spatial distribution of chronic wasting disease. Biol Lett 4:130–133.

795   87.   Somarelli JA, Makarewicz JC, Sia R, Simon R. 2007. Wildlife identified as major source of

796   Escherichia coli in agriculturally dominated watersheds by BOX A1R-derived genetic

797   fingerprints. J Environ Manage 82:60–65.

798   88.   Lam TTY, Ip HS, Ghedin E, Wentworth DE, Halpin RA, Stockwell TB, Spiro DJ, Dusek RJ,

799   Bortner JB, Hoskins J, Bales BD, Yparraguirre DR, Holmes EC. 2012. Migratory flyway and

800   geographical distance are barriers to the gene flow of influenza virus among North American

801   birds. Ecol Lett 15:24–33.

802   89.   Kreakie BJ, Fan Y, Keitt TH. 2012. Enhanced migratory waterfowl distribution modeling by

803   inclusion of depth to water table data. PLoS One 7:e30142.

804   90.   Carlson JC, Engeman RM, Hyatt DR, Gilliland RL, DeLiberto TJ, Clark L, Bodenchuk MJ, Linz

805   GM. 2011. Efficacy of European starling control to reduce *Salmonella enterica* contamination in

806   a concentrated animal feeding operation in the Texas panhandle. BMC Vet Res 7.

807   91.   Carlson JC, Franklin AB, Hyatt DR, Pettit SE, Linz GM. 2011. The role of starlings in the spread

808   of *Salmonella* within concentrated animal feeding operations. J Appl Ecol 48:479–486.

FIGURE LEGENDS

FIG 1 Sampling maps of (a) Flint Creek and (b) Hoosic River. Dots indicate the sampling sites within

each watershed. Map layers for land cover (National Land Cover Database [NLCD], 2006) were

acquired from the U.S. Geological Survey (USGS) Earth Explorer geographical data bank

(http://earthexplorer.usgs.gov/).

FIG 2 PCoA plots of *E. coli* clonal groups of sites in (a) Flint Creek (FC) and (b) Hoosic River (HR).

Green dots indicate forest sites; orange dots indicate produce field sites; Green circle indicates the 95%

confidence ellipse of forest sites; orange circle indicatedsthe 95% confidence ellipse of produce field

sites. *p* values of permutational multivariate analysis of variance (PERMANOVA) test and analysis of

similarities (ANOSIM) test are shown; "*" indicates the clustering of sampling sites by land-use is

significant at the 0.05 level. (a) PCo Axis 1 and 2 explained 18.7% and 14.9%, respectively, of the

variation of *E. coli* clonal groups for Flint Creek. (b) PCo Axis 1 and 2 explained 14.6% and 12.0%,

respectively, of the variation of *E. coli* clonal groups for Hoosic River.

FIG 3 Variance partitioning analysis (VPA) showing relative contributions of spatial factors (Geo.), soil

variables (Env.), and spatially structured environmental variables to the variations of the dissimilarity of

E. coli clonal groups based on Bray-Curtis distance for (a) Flint Creek (FC) and (b) Hoosic River (HR).

(c) Partial Mantel correlation between the dissimilarity of *E. coli* clonal groups and geographic distance-

correlated dissimilarity of soil variables for FC and HR. r is the Partial Mantel correlation coefficient.

Significant correlations at the 0.05 and 0.01 level are denoted by "*" and "**", respectively.

FIG 4 Linear relationship between biological dissimilarity of *E. coli* clonal groups and geographical

distance for (a) Flint Creek (FC) and (b) Hoosic River (HR). The biological dissimilarity of *E. coli*

clonal groups was calculated as Bray–Curtis distance. Geographical distance was calculated as the actual

physical distance. Linear regression line is in grey; shaded area indicates 95% confidence region; R^2

indicates the variability explained by fitted linear regression model and the formula of the linear

relationship is shown.

FIG 5 Mantel test result of wildlife-driven dispersal models and composition of *E. coli* clonal groups for

Flint Creek and Hoosic River. M106, M377, M459, and M556 on the x-axis are the identification

numbers of the predicted dispersal models for small mammals, large nuisance wildlife, migratory bird

flocks, and small flocking insectivore/granivores, respectively. Examples of each wildlife vehicle type

are shown. The description of the predicted dispersal model for each wildlife vehicle is detailed in Table

2. Models significant or marginally significant at the 0.05 level in Mantel tests are denoted by "*".

(A) Flint Creek

Forest F2
Field 2
Forest F1
Field 1
Field 6
Forest F6
Field 5   Forest F5
Forest F7
Field 7
Forest F8   Forest F9
Field 8   Field 9
Field 3
Forest F4
Field 4

N
0   0.45   0.9   1.8   2.7   3.6
Miles

(A) Flint Creek

Field 17   Forest F17
Forest F16   Field 18
Field 16   Forest F18

Field 14
Field 13
Forest F14   Forest F15
Forest F13   Field 15
Field 19
Field 10   Field 12
Forest F12   Forest F11
Field 11

N
0   1   2   4   6   8
Miles

(B) Hoosic River

(a) FC

Env.

11.7%

Geo.

5.5%

0.1%
Spatially structured Env.

Unexplained 82.7%

(b) HR

Geo.

Env.

1.0%

Unexplained 99.0%

(c)

Partial Mantel r

0.50
0.25
0.00
-0.25
-0.50

Soil variables

TABLE 1 Basic dispersal model characteristics

Class	Description
Dispersal vehicles[a]	
LT: Large nuisance wildlife species (e.g. "deer" or "feral swine")	Forests, scrublands, grasslands, wooded wetlands and cultivated croplands impose low movement costs, roads impose increased movement costs based on census category
ST: Small mammals (e.g. "shrews", "voles" and "mice")	Light urban development, grassland and scrublands impose low movement costs, roads impose increased movement costs based on census category
LB: Small flocking insectivore/granivores (e.g., "starlings")	Open water and heavy urban development are higher cost, all other movement costs are low
MB: Migratory bird flocks (e.g. "Canada goose")	Heavy urban development imposes higher cost, all other movement costs are low.
Riparian corridor (movement costs are reduced by half)	
Adjacency	All land parcels that overlap a 100 m zone around the main river/creek
Distance	Land within 100 m of the main river/creek
Biological	Land below the 50 yr flood height for the main river/creek and adjacent wetlands
None	No riparian corridor effect
Dispersal barriers	
Absolute	Major roads and waterways are absolute barriers (movement cost 40,000 per pixel)
Porous	Major roads and waterways are porous barriers (movement cost 200 per pixel)
None	No barrier effect
Proximity effects (specifics vary by vehicle)	
Strong	Nearness to high quality habitat substantially reduces movement cost
Weak	Nearness to high quality habitat somewhat reduces movement cost
None	No benefit of proximity to good cover
Attraction (gravity) coefficients (specifics vary by vehicle)	
Habitat Quality	Proximity, interspersion and area of high quality habitat increase the chances that *E. coli* will be deposited
Reduced Habitat Effect	Effect of high quality habitat is reduced by half
Area Independent	As habitat quality model, but area of high quality habitat does not impact the result
None	Attraction does not influence dispersal
Load (source) coefficient	
E. coli load estimation	Areas close to forest or pasture-class landcover are higher load.

[a]Conceptual models are based on published literature (80-91).

TABLE 2 The predicted dispersal model for each wildlife vehicle

Vehicles	Most-likely cost model[a]			Most-likely attraction model (coefficient)[b]
	Riparian corridor	Proximity effects	Dispersal barriers	
ST (Small mammals)	Biological	None	Absolute	None
LT (Large nuisance wildlife species)	Biological	Strong	Porous	Habitat Quality
MB (Migratory bird flocks)	Biological	Weak	None	Area Independent
LB (Small flocking insectivore/granivores)	Biological	Weak	Absolute	Area Independent

[a]Cost model was developed based on three factors, riparian corridor, proximity effects, and dispersal barriers. Definition of the three factors can be found in Table 1. The most-likely cost model was selected based on the characteristics of each wildlife vehicle (Table 1).

[b]Definition of classes of attraction model can be found in Table 1. The most-likely attraction model was selected based on the characteristics of each wildlife vehicle (Table 1).