Article

Recalled parental bonding interacts with oxytocin receptor gene polymorphism in modulating anxiety and avoidance in adult relationships

Ilaria Cataldo [1,†] , **Andrea Bonassi** [1,2,†] , **Bruno Lepri** [2] , **Jia Nee Foo** [3,4] , **Peipei Setoh** [5] , and **Gianluca Esposito** [1,4,5] ,*

1   Department of Psychology and Cognitive Science, University of Trento, Rovereto, Italy
2   Mobile and Social Computing Lab, Bruno Kessler Foundation, Trento, Italy
3   Human Genetics, Genome Institute of Singapore, Singapore, Singapore
4   Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore, Singapore
5   Psychology Program, School of Social Sciences, Nanyang Technological University, Singapore, Singapore
*   Correspondence: gianluca.esposito@ntu.edu.sg; gianluca.esposito@unitn.it
†   † These authors contributed equally to this work.

Abstract: Early interactions with significant individuals affect social experience throughout the
course of a lifetime, as a repeated and prolonged perception of different levels of care, independence
or control influences the modulation of emotional regulatory processes. As many factors play a role in
shaping the expectations and features of social interaction, in this study we consider the influence of
parental bonding and genetic allelic variation of oxytocin receptor gene polymorphism (rs53576) over
levels of experienced anxiety and avoidance in 313 young adults belonging to two different cultural
contexts, namely Italy and Singapore. Results highlight a major effect of maternal characteristics, care
and overprotection, with differences between the two cultural groups. Additionally, the interaction
between rs53576 and maternal overprotection suggest different environmental susceptibility in the
Italian sample and the Singaporean one. Implication in clinical work and future steps are described
in the conclusion.

Keywords: parental bonding; close relationship; social behavior; social development; ethnicity;
cross-cultural study, oxytocin receptor gene; gene-environment interaction; gene-culture interaction;
rs53576

1. Introduction

Fundamental works in disciplines related to developmental psychology have demonstrated that
early social interactions with caregivers influence different aspects of child development, such as
social relationships [1–3], academic performance [4], response to stress [5,6], individual well-being and
risk for psychopathology [7–10]. In addition, emotion-related skills in children like interpretation,
regulation, and communication are built on the emotional and social relationships they share with
their close ones [11]. Therefore it is likely that positive parenting practices, like higher levels of
warmth, care and sensitivity, will result in more adaptive emotion-regulation skills [12]. The quality
of parental bonding influences the development of top-down processes of emotional regulation,
which could potentially shape social interactions and physiological responses in individuals. The
ability to perceive typical social situations as non-threatening and risk-free also has implications for
the autonomic nervous system [13]. Generally, the literature points towards higher care and lower
over-protection as ideal parenting practices, since parental care allows individuals to be less restrained
in social interactions, while parental over-protection reduces the effectiveness of emotional regulation

and the ability to explore social situations. Consequently, adverse parenting practices are also relevant
to the manifestation of social anxiety traits in individuals [14].
According to the original theories on attachment by Bowlby and Ainsworth, attachment-specific
features were considered to extend beyond cultural boundaries or parenting practices – in other
words, Bowlby's and Ainsworth's original theory was considered to be culturally universal in the
sense that parental behavior that is sensitive to the child's needs will result in attachment security
across a diverse array of cultural and ecological contexts. Despite this assumption, the ecological
components of attachment have always been seen as an important factor that needs to be taken
into account, together with other features that might play a role in child rearing behavior, such as
socioeconomic status or ethnicity. In fact, one of the major concerns in attachment-related research
fields is the limitation due to results coming primarily from countries belonging to Western societies.
With regards to cultural studies, Ainsworth's canonical work in Uganda and Baltimore underlined that
processes related to maternal behavior reflect specific universal features that cross cultural, geographic
and linguistic boundaries [15]. In their research based on a sample of Japanese mothers, Behrens and
colleagues found that maternal attachment was predictive for the child's attachment classification
with a distribution comparable to global results [16]. In a review on parental sensitivity in ethnic
minority families, Mesman highlights the positive role of secure pattern of attachment on children's
development [17]. Central to the quality of parenting practices and attachment are the dimensions of
care and protection – encompassed in the construct of parental bonding [2]. Parental care refers to the
sensitivity of parents towards the child's needs (emotional, physical, psychological), while parental
over-protection refers to the excessive restrictions (emotional, physical, psychological) imposed on
the child. Partially inconsistent results have been described in a recent work on the link between
caregiver's warmth and levels of anxiety and avoidance in adult attachment in a Chinese population
[18].
53

1.1. Familial bonds in Italy and Singapore

In light of these considerations, it appears clear that, to have a proper understanding of
parental bonds, it is essential to take a wider look at the concept of familial nucleus according to
the country where these family units are immersed. In the current study we will focus on two
countries, one more Western-oriented (Italy) and another one representative of a non-Western
model (Singapore). In Italy, which is configured within individualistic countries, family is seen as
an expanded network of relationships and ties, where lots of attention is given to the child. The
relationship that forms between parent and child tends to be supportive and encouraging towards
independence, presenting high levels of emotional bonding [19]. On the other hand, Singapore
is described as a multicultural context, comprised of a majority of Chinese, Indian and Malay
families. In Singapore, which belongs to a collectivist culture, family is considered the first group a
person joins in his/her life. The family frame belongs to a collectivist scaffolding, like other Asian
countries, where interdependent relationships occurs on a social and familiar level, especially for
Chinese and Indian households. However, for all ethnicities, the extended family traditionally
used to live with the original family and parental control over the children was extended until
adulthood. With the sudden rise of technology and globalization, it is difficult, for the Singaporeans,
to maintain a multigenerational structure in families, and lots of people decide either to live alone
or not to have children. Furthermore, parental roles changed for those who decide to have babies,
especially in the last couple of decades: if the hierarchy once used to be based on a patriarchal
model, with the father having disciplinary power and the mother staying at home taking care of
family duties, now both parents share the care over their children. Although evidence coming from
different cultures and ethnicities have been reported and discussed, concerns about the application of
Western-based methodologies to non-Western environments still remains. In fact, attachment-related
mechanisms are, in turn, influenced by emotional communication and the way emotions are

expressed and experienced notably differs from country to country [20]. Furthermore, while
attachment dynamics have been profusely explored, little is known about the effect of processes that
elapse between childhood and early adulthood, in the manifestation of dynamics in adult relationships.
81

1.2. OXTr polymorphism rs53576

Considering the development of specific social characteristics as a result of the interaction
between biological and environmental factors, it appears clear that behavioral facets might present
a combination of genetic, environmental, and neurobiological factors. This involves the genetic
regulation of neurotransmitters that are notably involved in the modulation of social functioning,
such as oxytocin receptor gene (OXTr) [21–23]. Evidence in literature reports results about the
involvement of oxytocin receptor gene Single Nucleotide Polymorphisms (SNP) in modulating the
response to social stressors [24–26]. Within the polymorphic region of the OXTr, allelic variations
of the single nucleotide polymorphism rs53576 have been showed to be associated with different
social behaviors. Specifically, the G allele appears to be linked with more optimal social development,
which is described through higher levels of dispositional empathy [27,28], favourable prosocial
features [23,27,29,30], and greater autonomic reactivity to social stressors [31]. On the other hand,
A/A homozygotes have shown overall poorer social traits [32,33], reduced empathy accuracy [27],
and positive affect [34], all representing a less flexible social development. Although many results
highlight the association between specific variations of the genotype and their determined social
features, there are still some discrepancies in literature [35–37]. With specific regards to the interaction
with perceived parental warmth, higher levels of paternal care have been found to be associated
with increased heart rate response towards social cues for individuals with a G/G variation in an
Italian sample [29]. A study by Kim and colleagues compared a Korean and an American population
to investigate the interaction between perceived stress and OXTr rs53576 variations in modulating
the seek for emotional support, suggesting that, among Americans, G-carriers were seeking more
emotional social support compared to A/A homozygous, while the same difference was not shown in
the Korean sample. Moreover, they also compared Korean raised in America and Korean grown and
living in Korea, finding greater emotional social support seeking behavior in the first sample, hence
highlighting the relevance of the cultural environment [38].
107

1.3. Current Study: Hypotheses

The wider purpose of the current study is to deepen our understanding of how the interaction
between individual genetic features and the perception of parental warmth during childhood affects
levels of anxiety and avoidance in adult relationships. The aim is to analyze the differences occurring in
two different countries, like Italy and Singapore. The choice of these two contexts allows a comparison
of a Western country and a non-Western country, as environmental factors, such as the country
one belongs to, may affect on one hand the perception of parental features in childhood, and on
the other hand the formation of the behavioral constructs that drive social relationships. In order
to explore parents' transmitted behavior towards the child (assessed in terms of warmth/care and
overprotection), we used the Parental Bonding Instrument (PBI) by Parker and colleagues [2]. To
measure levels of anxiety and avoidance in adult relationships, we employed the revised version of
the Experience in Close Relationships (ECR-R) [39]. For the Singaporean sample, we used the original
versions in English, while for the Italian sample, participants completed the translated and validated
Italian versions [40,41]. We hence set two hypotheses on the Western (i.e. Italian) and Eastern (i.e.
Singaporean) sample respectively and independently from the cultural differences.

HP1 **gene*environment interaction on the Western sample:** we expect to find a statistically
significant effect of the interaction between OXTr polymorphism and parental bonding features
(assessed using the PBI scales) over the main features of adult social relationships (measured

with the ECR-R), with Italian A-carriers showing different rates of distress in terms of anxiety
and avoidance than G/G homozygous when they reported low levels of parental care and high
levels of parental overprotection;

HP2 **gene*environment interaction on the Eastern sample:** we expect to observe statistically
significant effect of the interaction between OXTr polymorphism and parental bonding features
(assessed using the PBI scales) over the main features of adult social relationships (measured
with the ECR-R), with Singaporean G-carriers exhibiting different rates of distress in terms of
anxiety and avoidance than A/A homozygous when they recalled low levels of parental care
and high levels of parental overprotection.

The predicted gene*environment interactions on the two separated samples could not however disclose
distinct pattern of adult attachment between Eastern and Western samples. Given the complexity of
the multiple relationship between genetic factors, cultural influences and early parental bonding on
adult attachment, potential gene*culture interactions are finally hypothesized at an observational level:

HP3 **culture*gene*environment interactions on the total sample:** we predict a differential
susceptibility of OXTr rs53576 to perceived parental bonding (assessed with the PBI subscales)
between the Western (i.e. Italian) and the Eastern (i.e. Singaporean) groups in explaining levels
of anxiety and avoidance (measured using the ECR index).

2. Materials and Methods

The research was approved by the Ethical Committee of Nanyang Technological University
(IRB-2015-08-020-01) and of University of Trento (Prot-2017-019) A). Informed consent was obtained
from all participants, and the study was conducted following the Declaration of Helsinki. Participants
were recruited through social media platforms (students' Facebook groups, for the Italian sample)
and web-based software research study participation and management systems (SONA System, for
the Singaporean sample) among non-parent students of University of Trento, Italy, and Nanyang
Technological University, Singapore. After informed consent was given, the online form of the
questionnaires were sent through email to the participants (links to Qualtrics for the Singaporean
sample, links to Google Modules for the Italian sample). Participants who gave consent for the
genetic part of the experiment were contacted to set an appointment to obtain a buccal mucosa sample
for genetic information. The genetic assessment was conducted on anonymized biosamples at the
Nanyang Technological University (Singapore) and at the Department of Neurobiology and Behavior
at Nagasaki University (Japan). Results from questionnaires were anonymized at the beginning of the
collection. The final group consisted of 313 participants that fully completed the two questionnaires
and provided the experimenters with DNA sample (Italian sample N = 97 (M = 38; F = 59); Mean age
= 23.01 (M = 23.58; F = 22.64), SD = 3.82 [range 18 35]; Singaporean sample N = 216; M = 73; F = 143;
Mean age = 21.51, SD = 1.83 [range 18 31]).

2.1. Questionnaires

Parental Bonding Instrument. The Parental Bonding Instrument (PBI) is a 50-item self-report
questionnaire that investigates participants' perception of both maternal (25 items) and paternal (25
items) care and protection in their first 16 years of life. It was developed by Parker, Tupling and Brown
in 1979 using factor analysis from parents self-reporting their childhood experiences, the results of
which yielded two factors: warmth/care (hereafter referred to as care) and overprotection [2]. The
care scale (PBI_Care) measures the degree to which a mother or a father was empathetic and caring
versus cold and indifferent. The overprotection scale (PBI_OverP) measures the extent to which a
parent was intrusive or, conversely, fostered independence in the subject. The measure has been shown
to have high reliability, stability over time and no association with social desirability, neuroticism or
extroversion [42].

Experience in Close Relationships - Revised. In order to evaluate anxiety and avoidance level in

close relationships, the Experience in Close Relationships-Revised was employed (ECR-R; [39,41]). The 36-item self-report questionnaire assesses two major dimensions of an individual's attachment style (anxiety and avoidance) in a sentimental relationship. Both the anxiety and avoidance subscales consisted of 18 items each, rated on a 7-point Likert scale ranging from 1 (Strongly disagree) to 7 (Strongly agree). The anxiety dimension measures insecurity, jealousy and fear of abandonment as opposed to feeling secure about the availability and responsiveness of romantic partners. The avoidance dimension measures the feeling of discomfort being close to others and tendency to refrain from attachment. Instead of assigning participants to an attachment style category, this scale yields two separate dimension scores for each participant.

2.2. Genotyping

This study adopted the same DNA derivation and genotyping procedure used by Bonassi and colleagues [43], using ACGT, Inc. (Wheeling, IL). DNA was extracted from each kit using the Oragene DNA purifying reagent, then concentrations were assessed through spectroscopy (NanoDrop Technologies, USA). Concentrations for each sample were magnified through polymerase chain reaction (PCR) for the OXTr gene rs53576 region target. The forward and reverse primers that were used were 5-GCC CAC CAT GCT CTC CAC ATC-3 and 5-GCT GGA CTC AGG AGG AAT AGG GAC-3. For this DNA region, the frequency for the allelic distributions (A/A, A/G, G/G) differ among populations belonging to different ethnic groups. Although a unique consensus is still not reached and may depend on the alleles' distributions within a specific sample, the A allele is usually combined with the same heterozygous group A/G in the Western samples, whereas the G allele is more commonly paired with the heterozygous group A/G in the Eastern samples [29,35,44,45]. In cross-cultural investigations analyzing different ethnic groups as a whole, reducing the number of genetic groups, A/A is maintained in contrast with the combined couples A/G and G/G [38,46]. Here, Western participants (e.g., all Italian native participants with Caucasian ethnicity) having at least one A allele (A/A homozygotes or A/G) were classified into a single A-carriers group. Complementary, Eastern participants (e.g., all Singaporean native participants with South East-Asian ethnicity, namely 184 Chinese, 15 Malays, 8 Indian, 3 Caucasian, 3 Filipino, 1 Vietnamese, 1 Gurkha, and 1 Korean) having at least one G allele (G/G homozygotes or A/G) were treated as a single G-carriers group. The averaged distribution of the different genotypes in the Caucasian population is 25-35% for A/A homozygotes and 65-75% for G-carriers[1], whereas the distribution in the Italian sample was 15% for A/A homozygous and 85% for G-carriers. Specifically, genotype frequencies were as follows: A/A = 14 (14.43%), A/G = 35 (36.08%), G/G = 48 (49.49%). In the present Italian sample, the less frequent homozygous group A/A was merged with the heterozygous group A/G (A-carriers = 51%; G/G = 49%) before the analysis. Participants' age ($t = 1.761$, $p = 0.383$) and gender $X^2(1) = 0.016$, $p = 0.899$) did not significantly differ between the G/G and A groups. As for the Singaporean sample, the allelic distribution in East-Asian population is 65-75% for A/A homozygotes and 25-35% for G-Carriers [2], whereas in our sample the distributions were 37% for A/A homozygotes and 63% for G-carriers. Specifically, genotype frequencies were: A/A = 80 (37.04%), A/G = 94 (43.52%), and G/G = 42 (19.44%). Participants' age ($t = 0.481$, $p = 0.631$) and gender $X^2(1) = 0.189$, $p = 0.663$) did not significantly differ between the A/A and G groups. Alleles frequency distribution and Hardy-Weinberg Equilibrium parameters are reported in Table 1. In the total sample, the distribution of the alleles was as follows: A/A = 94 (30.03%), A/G = 129 (41.21%), and G/G = 90 (28.75%). Since the genotype frequencies for the whole sample (Total Sample = Italian Sample + Singaporean Sample) did not satisfy the Hardy-Weinberg Equilibrium, analyses focused

---

[1] 1000 Genomes project, ensembl.org
[2] 1000 Genomes project, ensembl.org

218 on the three distinct allelic combinations within the frame of an exploratory approach. Participants'
219 gender ($X^2(2)$) = 0.008, p = 0.996) did not significantly differ between the three groups. Participants'
220 age did not significantly differ between A/A and A/G (t = -0.814, p = 0.883), A/A and G/G (t = 0.417,
221 p = 0.413), and G/G and A/G (t = -0.991, p = 0.323).

Genotype	A/A	A/G	G/G	X^2	*p-value*
rs53576$_{ITA}$	14	35	48	3.05	.08
rs53576$_{SNG}$	80	94	42	2.24	.13
rs53576$_{Total}$	94	129	90	9.64	.02**

Table 1. Hardy-Weinberg Equilibrium results for OXTr rs53576 in the Italian sample, in the Singaporean sample, and in the total sample. **p<.01

2.3. Analyses

224 Data were analyzed in R (Version 4.0.0). Statistical tests were differently planned for
225 hypothesis-driven and exploratory analysis. The hypothesis-driven tests were performed separately
226 on the Italian and the Singaporean sample, whereas the exploratory tests were conducted on the total
227 number of participants (Total sample = Italian sample + Singaporean sample). Overall, the statistical
228 analyses follow a common procedure of variables' evaluation and tests' application.

2.3.1. Preliminary Analyses

Before proceeding with the data analysis, we tested the internal validity of each questionnaire in
both groups (Italy, Singapore). The Cronbach's alpha coefficients ranged from 0.78 to 0.94, suggesting
an overall very good internal consistency. In Table 2, all the Cronbach's alpha values are listed, together
with the main descriptive statistics of each sample. Psychometric features of the two instruments are
compatible with those reported in a recent review on adult attachment assessment tools [47].
Univariate and multivariate distributions of PBI and ECR-R scores were inspected for normality
and the presence of extreme values. 76 observations (Italian sample = 27; Singaporean sample = 49)
were found as having a value higher/lower to 2 *SDs* above/below the mean. Any artificial restriction
(i.e. outliers' removal or treatment) was avoided to preserve the meaning associated to the scoring of
each extreme observation. A high reliability of the subscales (see Table 2) also excludes potential biases
due to extreme scores [48]. As observed in the Table 2, acceptable skewness and kurtosis values as
well as visual inspection via density and quantile-quantile plots proved that PBI and ECR-R subscales
were normal univariate distributions (see Table 2). Only the distribution of PBI Maternal Care skewed
right in the Italian sample (skewness = 1.15; kurtosis = 0.17), thus the variable was square-transformed
(skewness = -0.43; kurtosis = -0.88) before analysis. The same approach was adopted by previous
researches in the field [38]. Preliminary tests were made in each population group to exclude any
significant effect on the ECR-R dependent variables which could be attributed to participants' gender
and age for the exploratory analysis (see Figures S1 and S2 of the "Supplementary Materials"). A
differential Bonferroni correction was applied to each group in adjusting *alpha* (Italian sample: 4
repeated measures, corrected *alpha* = 0.0125; Singaporean sample: 4 repeated measures, corrected
alpha = 0.0125; Total sample: 6 repeated measures, corrected *alpha* = 0.008). No effects of gender
or age on the ECR-R dimensions were significant (see Table 3). However, a main effect of culture
was observed on adult anxiety, whereas the effect of culture on adult avoidance did not survive
the correction (see Table 3; see Figure S3 of the "Supplementary Materials"). This preliminary result
confirmed the necessity to inspect the potential interaction between culture, genotype and caregiving
behavior, as hypothesized by *HP3*.

Group	Subscale	Cronbach α	Skewness	Kurtosis	Mean (SD)	Median
Italy	Age	-	-	-	23.01 (3.82)	22
	PBI M_Care	0.93	-1.15*	0.71	26.33 (8.50)	28
	Squared PBI M_Care	-	-0.43	-0.88	764.70 (373.24)	784
	PBI M_OverP	0.78	0.72	0.18	16.40 (6.30)	16
	PBI F_Care	0.94	-0.49	-0.69	23.23 (9.30)	25
	PBI F_OverP	0.83	0.75	0.17	14.12 (7.01)	13
	ECR-R Anxiety	0.91	0.70	0.14	56.59 (20.10)	53
	ECR-R Avoidance	0.88	0.38	-0.39	35.12 (15.37)	33
Singapore	Age	-	-	-	21.51 (1.83)	21
	PBI M_Care	0.91	-0.66	-0.03	25.39 (6.99)	26
	PBI M_OverP	0.86	0.40	-0.01	15.09 (7.06)	15
	PBI F_Care	0.92	-0.28	-0.41	21.44 (8.14)	22
	PBI F_OverP	0.86	0.69	0.51	11.57 (6.63)	11
	ECR-R Anxiety	0.93	-0.04	-0.39	64.79 (20.61)	66.50
	ECR-R Avoidance	0.94	0.12	-0.51	40.31 (17.13)	40
Total	Age	-	-	-	21.97 (2.70)	22
	PBI M_Care	0.92	-0.86	0.36	25.68 (7.49)	27
	PBI M_OverP	0.83	0.45	0.09	15.50 (6.85)	15
	PBI F_Care	0.93	-0.32	-0.53	21.99 (8.54)	23
	PBI F_OverP	0.83	0.71	0.46	12.36 (6.84)	12
	ECR-R Anxiety	0.92	0.18	-0.44	62.28 (20.76)	62
	ECR-R Avoidance	0.92	0.22	-0.48	38.70 (16.75)	39

Table 2. Summary of Cronbach α, skewness and kurtosis for each subscale of the Parental Bonding Instrument and Experience in Close Relationships in the Italian sample, Singaporean sample and total sample. Statistics of age as continuous variable are also reported. * = squared transformation applied.

Sample	ECR-R subscale	Variable	t (df)	p-value	95% CI	r
Italy	Anxiety	Gender	-1.21 (95)	.23	[-13.31 ; 3.25]	
		Age	0.19 (95)	.85	[-7.38 ; 8.93]	- 0.03
	Avoidance	Gender	2.19 (95)	.03	[0.67 ; 13.12]	
		Age	-0.49 (95)	.62	[-7-78 ; 4.69]	0.19
Singapore	Anxiety	Gender	1.69 (214)	.09	[-0.84 ; 10.80]	
		Age	-0.03 (214)	.97	[-5.64 ; 5.45]	0.03
	Avoidance	Gender	-0.21 (214)	.83	[-5.39 ; 4.34]	
		Age	1.00 (214)	.32	[-2.27 ; 6.92]	-0.07
Total	Anxiety	Gender	.54 (311)	.59	[-3.50 ; 6.16]	
		Age	-0.14 (311)	.89	[-5.27 ; 4.57]	- 0.05
		Culture	3.24 (311)	.13*	[3.18 ; 13.02]	
	Avoidance	Gender	0.81 (311)	.42	[-2.29 ; 5.50]	
		Age	0.16 (311)	.87	[-3.65 ; 4.30]	0.01
		Culture	2.56 (311)	.01	[1.19 ; 9.18]	

Table 3. Summary of statistical values from Student's t-tests for gender and age over the ECR-R dependent variables anxiety and avoidance. Pearson correlation coefficients among each ECR-R variable and age are also reported. *$p<.008$

2.4. Statistical Analyses

Exploratory correlation matrices were generated for the Italian, Singaporean and Total samples to probe possible associations among the continuous variables (e.g., PBI and ECR-R dimensions). A series of hierarchical multiple regression (HMR) analyses was performed on the Italian and Singaporean sample separately to test the hypothesis that gene*environment interaction would predict the dependent variable (DV) adult anxiety (e.g., one series on the Italian sample, one series on the Singaporean sample). A series of HMR analyses was also conducted on the Italian and Singaporean sample separately to test the hypothesis that gene*environment interaction would predict the DV adult avoidance (e.g., one series on the Italian sample, one series on the Singaporean sample). In summary,

a six-steps HMR was conducted for each ECR-R subscale (namely anxiety and avoidance) as DV in reference to each sample (see the models applied to each step at the Table S1 of the "Supplementary Materials"). Allelic variation for OXTr was entered at stage 1 as a between-subjects factor of the regression to control for differences due to the genetics (Italian sample: G/G vs. A-carriers; Singaporean sample: A/A vs. G-carriers). Progressively, all the PBI subscales as continuous predictors were entered to each following step; specifically, maternal care was entered at step 2, maternal overprotection at step 3, paternal care was entered at step 4, paternal overprotection at step 5. In the final model (step 6) the interaction between OXTr polymorphisms variants and perceived parental bonding was tested.

A further series of HMR analyses was conducted on the Total sample to test the exploratory hypothesis that culture would modulate the interaction between OXTr and early caregiving behavior on adult anxiety and avoidance as unique DV (e.g., one series on the total sample for anxiety as DV, one series on the total sample for avoidance as DV). Therefore, in the total sample an initial step was added in order to include the cultural variable (Italian vs. Singaporean) in the predictors of the regression model, for a total number of seven steps (see Table S1 of the "Supplementary Materials"). Allelic groups for OXTr were here added at stage 2 as a between-subjects factors (Total sample: A/A vs. A/G vs. G/G). Successively, all the PBI dimensions were singularly entered to each following step. The final model (step 7) was here defined by the interaction between culture, OXTr polymorphisms variants and perceived parental bonding.

In order to compare the significance of each progressive model, the analysis of variance (ANOVA) was used to analyze the contribution of each predictor added at every step. As contrast coding system for the categorical variables (e.g., culture and OXTr as factors) entered in the linear models, the dummy coding was applied to compare each level of a given categorical variable to a fixed reference level. The factor OXTr presented two levels for the analysis on the Italian (reference level: A-carriers as 1; G/G as 2) and Singaporean (reference level: G/G as 1; A-carriers as 2) samples, but three levels were considered for the analysis on the Total sample (reference level: A/A as 1; A/G as 2; G/G as 3). The factor culture included two levels for the analysis developed on the Total sample (reference level: Singaporean as 1; Italian as 2).

Overall, the hierarchical regression allowed to identify the positive predictors of the dependent variable at different stages of increasing complexity. The final step was reached when all the variables defined by the hypotheses were included in the model. As described, the final models included the interaction between all the embedded variables (interactive model) as predicted by the hypotheses.

This type of regression is particularly targeted in research design with multiple variables (e.g., different contributing factors and/or continuous subscales), nested designs and unpredictable outcomes or exploratory analyses [49,50] and was already applied in previous studies of this field [38,51]. Before conducting the HMR analyses, the related assumptions (multicollinearity, i.e., variance inflation factor, homogeneity of variance, i.e., Bartlett's test) were tested and satisfied. The post-hoc statistical power was calculated for the final model of each series of HMR analysis of each sample using G*Power software (version 3.1) [52]. The magnitude of each effect for the hierarchical linear models were estimated by R, R^2 and ΔR^2.

For each sample, the level of significance for the effects emerging from the final step of each HMR analysis was adjusted as a function of the two HMR analysis conducted on anxiety and avoidance respectively (2 repeated measures for Italian, Singaporean and Total sample separately; corrected $alpha = 0.025$). Bonferroni's method was chosen instead of alternative techniques to apply a more conservative correction.

For each sample, the significant main and interaction effects obtained from the final step of each series of HMR analysis were graphically represented by scatterplots with linear models and barplots. A further assessment was advanced to inspect the direction of such effects. Pearson's r and Fisher's z tests evaluated the degree of association between the continuous independent variables and the dependent variables [53]. Variables related to PBI subscales were also divided into two groups (low vs. high PBI dimension) using the median split procedure for post-hoc tests. Post-hoc analyses were

computed for significant results on ECR-R variables adopting the Student's *t*-tests. Welch's *t*-tests were preferred rather than Student's *t*-tests when the variance of the comparing two groups from the normal population was different. A false discovery rate correction (e.g., Bonferroni's method) for the *t*-tests was considered to circumscribe the probability of committing a type I error. The magnitude of the significant effects for *t*-test was estimated by Cohen's *d*.

With regards to the significant main effects, the mean differences of the given ECR-R variable between low and high PBI subscales ($\alpha = 0.05$) were tested (e.g., see below the main effects for the Italian, Singaporean and Total sample). Concerning the significant interactions between genotype and environment, low vs. high PBI levels were compared to analyze the differences between the genetic carriers on the ECR-R dimension (e.g., see below the effects for the Singaporean sample given the corrected $\alpha = 0.025$ and the total sample given the corrected $\alpha = 0.017$). In the context of the significant interactions between culture and environment, low vs. high PBI levels were contrasted to probe the differences between the Italian and Singaporean individuals (corrected $\alpha = 0.025$) on the ECR-R subscale (e.g., see below the effects for the Total sample). In reference to the significant interactions between culture, genotype and environment, low vs. high PBI levels were contrasted to inspect the differences between the genetic carriers from the Italian and Singaporean samples (corrected $\alpha = 0.025$) on the ECR-R variable (e.g., see below the effects for the Total sample).

Results

Means and standard deviations of the continuous variables are shown in Table 2, and their intercorrelations were examined, as reported in Table 4 for the Italian and Singaporean sample respectively, and in Table S2 of the Supplementary Materials for the total sample (see "Supplementary Materials").

subscale	M_Care	M_Overp	F_Care	F_Overp	Anxiety	Avoidance
M_Care		-.29***	.43***	-.10	-.06	-.07
M_Overp	-.31*		-.17	.39***	.1	.05
F_Care	.49***	-.23		-.18	-.16	-.09
F_Overp	-.07	.39***	-.30*		.17	0.10
Anxiety	-.36**	.42***	-.37**	.20		.38***
Avoidance	-.25	.19	-.20	.08	.35**	

Table 4. Pearson correlation coefficients and significance values among questionnaire subscales of the Italian and Singaporean samples separately. Values above the diagonal refer to the Singaporean sample, while values under the diagonal refer to the Italian sample. As regards the Italian sample, the PBI maternal care was square-transformed before the analysis. Significance is adjusted for multiple tests (corrected *alpha* = 0.003). *p<.003 **p<.001 ***p<.0001

A six-steps HMR was conducted separately on the Italian and Singaporean sample for each ECR-R subscale (namely Anxiety and Avoidance) as DV (see Table S1 of the "Supplementary Materials"). A seven-steps HMR was conducted on the Total sample for each ECR-R subscale as DV (see Table S1 of the "Supplementary Materials").

HMRs with the final step showing significant effects (*p*<0.025) are explained below and reported in Table 5 for the Italian sample, in Table 6 for the Singaporean sample and in Table 7 for the Total sample. Otherwise, explanation and tabulation of the results obtained from the HMRs not presenting significant effects ($p > 0.025$) at the final step are presented in the Supplementary Materials (see "Supplementary Materials").

Italian Sample

Anxiety

Table 5 represents the series of HMR conducted on ECR-R anxiety of the Italian sample (total SS =
38793). At step 1, OXTr explained a not significant 1% of variance ($F(1,95) = 0.62$, RSS = 38597, $p= .43$).
Entering the maternal care variable at step 2 explains a significant additional 13.3% of variance ($F(1,94)$
= 16.35, RSS = 33425, $p< .0001$). After introducing maternal overprotection at step 3, both maternal
care and maternal overprotection were positive predictors, accounting for a significant 10% of variance
($F(1,93) = 12.55$, RSS = 29455, $p< .001$). The addition of paternal care variable to the regression model
(step 4) justified a further significant 4% of variation in anxiety ($F(1,92) = 4.85$, RSS = 27921, $p= .03$)
given the main effects of maternal overprotection and maternal care. No significant change in the
R^2 was observed when the final variable paternal overprotection was included at step 5 ($F(1,91)$
= 0.00, RSS = 27921, $p= .99$). At step 6, the hypothesized interaction between OXTr and paternal
bonding dimensions on anxiety was tested ($f^2 = .39$, power = .99). Here, a main effect of maternal
overprotection was found ($\beta = 1.11$, SE = 0.46, $t = 2.40$, $p< .02$). Although the final model computed the
highest proportion of explained variance in anxiety by the model ($R^2 = .29$), no significant difference
was detected compared to the previous step ($F(1,91) = 0.32$, RSS = 27517, $p= .87$).

Step	Variable	β	SE	t	R	R^2	ΔR^2
1					.07	.01	.01
	OXTr rs53576	2.84	4.09	0.69			
2					.37	.14	.13***
	OXTr rs53576	3.11	3.83	0.81			
	M_Care	-0.02***	0.01	-3.81			
3					.49	.24	.10***
	OXTr rs53576	3.36	3.62	0.93			
	M_Care	-0.01**	0.01	-2.73			
	M_OverP	1.08***	0.30	3.54			
4					.53	.28	.04*
	OXTr rs53576	4.06	3.55	1.14			
	M_Care	-0.01	0.01	-1.46			
	M_OverP	1.01**	0.30	3.39			
	F_Care	-0.50*	0.22	-2.25			
5					.53	.28	0
	OXTr rs53576	4.06	3.57	1.14			
	M_Care	-0.01	0.01	-1.42			
	M_OverP	1.01**	0.33	3.11			
	F_Care	-0.50*	0.23	-2.13			
	F_OverP	0.00	0.29	0.00			
6					.54	.29	.01
	OXTr rs53576	18.08	17.73	1.02			
	M_Care	-0.01	0.01	-0.72			
	M_OverP	1.12*	0.47	2.40			
	F_Care	-0.48	0.38	-1.29			
	F_OverP	0.20	0.44	0.45			
	OXTr rs53576 x M_Care	0.00	0.01	-0.34			
	OXTr rs53576 x M_OverP	-0.26	0.67	-0.39			
	OXTr rs53576 x F_Care	-0.01	0.49	-0.01			
	OXTr rs53576 x F_OverP	-0.46	0.61	0.76			

Table 5. Hierarchical multiple regression on ECR-R anxiety for the Italian sample. Note. SE = standard error of unstandardized coefficient. *$p<.05$; **$p<.01$; ***$p<.001$.

As indicated in the Table 4, maternal overprotection was also found positively associated with adult anxiety (t = 4.46, df = 95, r = 0.42, $p<$ .0001) (Figure 1). Specifically, the two-tailed post hoc Welch's t test ($\alpha = 0.05$) revealed that felt anxiety towards the partner was significantly different between participants who experienced low levels in maternal overprotection compared to those who had a past of maternal overprotection (t = -2.31, df = 71.3, $p=$ .024, $d= -0.49$) (Figure 1). Means and standard error values for this effect on avoidance are available from the Table S6 of the Supplementary Materials (see "Supplementary Materials").

Contrary to *HP1*, no main effect of genotype, or interactions between covariates and genotype were identified.

Figure 1. Effect on anxiety for the Italian sample. (**A**). Effect of early maternal overprotection on the adult anxiety. Correlation between the scores of anxiety and maternal overprotection. Blue circles = Italian participants. The red line represents the linear model for the Italian participants. r-value refers to Pearson's r correlation. (**B**). Comparison between the scores of ECR-R anxiety in low and high maternal overprotection (* p < 0.025).

Avoidance

Table S3 of the Supplementary Materials (see "Supplementary Materials") illustrates the series of HMR conducted on ECR-R avoidance for the Italian sample. The related description is reported in the Supplementary Results (see "Supplementary Materials").

Singaporean Sample

Anxiety

Table S4 of the Supplementary Materials (see "Supplementary Materials") shows the series of HMR conducted on ECR-R anxiety for the Singaporean sample. The related details are presented in the Supplementary Results (see "Supplementary Materials").

Avoidance

Table 6 presents the series of HMR performed on ECR-R avoidance for the Singaporean sample (total SS = 63074). Entering at step 1 the only genetic variable did not increase the explained portion of variance ($F(1,214)$ = 0.18, RSS = 63021, $p=$ .67). At step 2, maternal care was added to the regression model and accounted for a non-significant 1% of additional variance ($F(1,213)$ = 0.99,

RSS = 62733, *p*= .32). The inclusion of maternal overprotection at step 3 did not further improve the value of R^2 (*F*(1,212) = 0.15, RSS = 62689, *p*= .70). No change in the explained variance was even observed when introducing the paternal care dimension at step 4 (*F*(1,211) = 0.87, RSS = 62438, *p*= .35). At step 5, the addition of the subscale paternal overprotection justified an increased 1% of variation in avoidance which was not significant (*F*(1,210) = 1.59, RSS = 61976, *p*= .21). At step 6, the hypothesized (*HP2*) interaction between OXTr and paternal bonding dimensions on avoidance was tested (f^2 = .05, power = 0.67). The final model maximized the explained variance (R^2 = .05), but no significant difference was found compared to the previous step (*F*(4,206) = 1.74, RSS = 59956, *p*= .14).

Step	Variable	β	SE	*t*	R	R^2	ΔR^2
1					.03	.00	.00
	OXTr rs53576	1.02	2.42	0.43			
2					.07	.01	.01
	OXTr rs53576	0.86	2.42	0.35			
	M_Care	-0.17	0.17	-0.99			
3					.08	.01	.00
	OXTr rs53576	0.78	2.44	0.32			
	M_Care	-0.15	0.18	-0.84			
	M_OverP	0.07	0.17	0.38			
4					.10	.01	.00
	OXTr rs53576	1.00	2.45	0.41			
	M_Care	-0.08	0.19	-0.39			
	M_OverP	0.06	0.18	0.33			
	F_Care	-0.15	0.16	-0.92			
5					.13	.02	.01
	OXTr rs53576	0.95	2.45	0.39			
	M_Care	-0.09	0.19	-0.49			
	M_OverP	-0.03	0.19	-0.16			
	F_Care	-0.12	0.16	-0.73			
	F_OverP	0.24	0.19	1.25			
6					.22	.05	.03
	OXTr rs53576	-17.16	13.28	-1.29			
	M_Care	-0.83*	0.35	-2.37			
	M_OverP	0.14	0.35	0.40			
	F_Care	0.22	0.29	0.76			
	F_OverP	0.02	0.33	0.05			
	OXTr rs53576 x M_Care	1.03*	0.42	2.45			
	OXTr rs53576 x M_OverP	-0.16	0.42	-0.39			
	OXTr rs53576 x F_Care	-0.45	0.35	-1.27			
	OXTr rs53576 x F_OverP	0.30	0.41	0.73			

Table 6. Hierarchical multiple regression on ECR-R avoidance for the Singaporean sample. Note. SE =standard error of unstandardized coefficient. **p<.05*

At this final step, early maternal care was the strongest predictor of adult avoidance (β = -0.83, SE = 0.35, *t* = -2.37, *p*< .02). As reported in the Table 4, the negative correlation between maternal care and adult avoidance was not significant (*t* =-1.02, df = 214, *r* = -0.07, *p*= .30) (Figure 2). Specifically, the two-tailed post hoc Student's *t* test (α = 0.05) showed that the avoidance towards the partner was not significantly different between participants who reported low and high scores in maternal care (*t* = 0.78, df = 214, *p*= .44, *d*= 0.11) (Figure 2).

The significant two-way interaction between rs53576 and maternal overprotection also emerged for the adult avoidance (β = -0.83, SE = 0.35, *t* = -2.37, *p*< .02). The distribution of genotypes A/A vs. G-carriers was not significantly different between high vs. low maternal care (X^2 (1) = 1.19, *p*= .028).

Maternal care was positively associated with the avoidance for G-carriers (t = 0.51, df = 134, r = 0.04, p= .61), but negatively associated with the avoidance for A/A homozygotes (t = -2.38, df = 78, r = -0.26, p= .02), as predicted by *HP2* (Figure 2). Although only one Pearson's r was significant, Fisher's z test confirmed that the difference between the slopes for A/A and G-carriers was significant (z = -2.14, p= .03). The one tailed post-hoc Student's t test (corrected α = 0.025) revealed that adult avoidance was not significantly different between A/A and G-carriers when they experienced a history of low maternal caregiving (t = -1.01, df = 107, p= .84, d= -0.21) (Figure 2).

Means and standard error values for each effect on avoidance are available from the Table S7 of the Supplementary Materials (see "Supplementary Materials").

Total Sample: Italian and Singaporean participants

Anxiety

Table S5 of the Supplementary Materials (see "Supplementary Materials") represents the series of HMR conducted on ECR-R anxiety for the total sample. The statistical details are indicated in the Supplementary Results (see "Supplementary Materials").

Avoidance

Table 7 reports the series of HMR performed on ECR-R avoidance for the total sample (total SS = 87563). The regression revealed that at stage 1 culture contributed significantly to the model explaining a first 2% of variance (F(1,311) = 6.90, RSS = 85763, p= .009). Culture was significant at this step, as positive predictor. At step 2, introducing the genetic factor did not increase the variance of avoidance (F(2,309) = 0.03, RSS = 85745, p= .97), but a main effect of culture was still present. At step 3, maternal care was added to the regression model and accounted for a significant 1% of additional variance (F(1,308) = 4.42, RSS = 84592, p= .04). Here, the variable culture and the novel variable maternal care were the strongest predictors. At step 4, the improvement of 1% of variation in avoidance due to the inclusion of maternal overprotection in the model was not significant (F(1,307) = 1.03, RSS = 84325, p= .31). Only culture was a positive predictor at this stage. No change in the explained variance was observed when introducing the paternal care dimension at step 5 (F(1,306) = 1.79, RSS = 83858, p= .18). Culture was still a strong predictor at this stage. The addition of paternal overprotection variable to the regression model at step 6 justified a 1% of variation in avoidance which did not result statistically significant F(1,305) = 1.11, RSS = 83569, p= .29). However, a significant main effect of culture was identified at this stage. Finally, at step 7, the potential interaction between culture, OXTr and PBI dimensions on avoidance (*H3*) was inspected (f^2 = .17, power = 0.99). This interactive model increased the explained variance (R^2 = .16) significantly from the previous step (F(22,283) = 1.70, RSS = 73827, p= .028).

At this last step, one one-way effect, three two-way interaction effects and two three-way interaction effects were found. Means and standard error values for each effect on avoidance are available from the Table S8 of the Supplementary Materials (see "Supplementary Materials"). Firstly, the early maternal overprotection singularly predicted adult avoidance (β = 5.03, SE = 1.71, t = 2.95, p< .004). As evident from the Table S4 of the Supplementary Materials (see "Supplementary Materials"), the positive correlation between maternal overprotection and adult avoidance was not significant (t =-1.26, df = 311, r = 0.07, p= .21) (Figure 3**A**; blue circles = Italian and Singaporean participants merged; the red line represents the linear model for all the participants). The two-tailed post hoc Student's t test (α = 0.05) showed that the avoidance towards the partner was not significantly different between participants who reported low and high scores in maternal overprotection (t = -1.27, df = 311, p= .21, d= -0.14) (Figure 3**B**). Whereas the main effect of paternal care (p> .025) did not survive the magnitude of the Bonferroni correction (corrected $alpha$ = 0.025).

SINGAPOREAN SAMPLE

Figure 2. Effects on avoidance for the Singaporean sample. (**A**). Effect of early maternal care on the adult avoidance. Correlation between the reported avoidance and maternal care. Blue circles = Singaporean participants. The red line represents the linear model for the Singaporean participants. (**B**). Comparison between the scores of ECR-R avoidance in low and high maternal care. (**C**). Effect of the interaction between genotype and early maternal care on the adult avoidance. Correlations between the scores of anxiety and the recalled maternal care. Red triangles = A/A; blue circles = G-carriers. Lines depict the linear models for A/A homozygotes (red) and G-carriers (blue). (**D**). Contrast between the scores of ECR-R avoidance in A/A homozygotes (red) and G-carriers (blue) divided into low and high maternal care. *r*-values refer to Pearson's *r* correlations for both the panels (**A**) and (**C**).

As concerns the interaction effects, the significant two-way interaction between culture and maternal overprotection also emerged for the avoidance (β = -4.89, SE = 1.74, t = -2.81, p= .005). Maternal oveprotection positively correlated with the avoidance reported by both Italian (t = 1.91, df = 95, r = 0.19, p= .06) and Singaporean participants (t = 0.69, df = 214, r = 0.05, p= .49) (Figure 3**C**; red triangles = Italian participants; blue circles = Singaporean participants; lines depict the linear models for Italian-red and Singaporean-blue participants). However, the slopes for Italians and Singaporeans were not significantly different (z = 1.15, p= .13). The exploratory two-tailed Student's

Step	Variable	β	SE	t	R	R²	Δ R²
1					.14	.02	.02**
	Culture	5.19*	2.03	2.55			
2					.14	.02	.00
	Culture	5.05*	2.15	2.35			
	OXTr rs53576 A/G-A/A	0.21	2.27	0.09			
	OXTr rs53576 G/G-A/A	-0.39	2.59	-0.15			
3					.18	.03	.01*
	Culture	4.74*	2.15	2.21			
	OXTr rs53576 A/G-A/A	-0.01	2.27	-0.01			
	OXTr rs53576 G/G-A/A	-0.69	2.58	-0.27			
	M_Care	-0.26*	0.13	-2.05			
4					.19	.04	.01
	Culture	4.88*	2.15	2.27			
	OXTr rs53576 A/G-A/A	-0.12	2.27	-0.05			
	OXTr rs53576 G/G-A/A	-0.99	2.60	-0.38			
	M_Care	-0.22	0.13	-1.67			
	M_OverP	0.14	0.15	0.99			
5					.21	.04	.00
	Culture	4.75*	2.15	2.20			
	OXTr rs53576 A/G-A/A	0.15	2.28	0.07			
	OXTr rs53576 G/G-A/A	-0.58	2.62	-0.22			
	M_Care	-0.14	0.14	-0.99			
	M_OverP	0.13	0.15	0.88			
	F_Care	-0.16	0.12	-1.31			
6					.21	.05	.01
	Culture	5.11*	2.18	2.34			
	OXTr rs53576 A/G-A/A	0.26	2.28	0.12			
	OXTr rs53576 G/G-A/A	-0.53	2.62	-2.20			
	M_Care	-0.16	0.15	-1.11			
	M_OverP	0.07	0.16	0.42			
	F_Care	-0.14	0.13	-1.08			
	F_OverP	0.16	0.16	1.03			
7					.40	.16	.11*
	Culture	11.02	27.75	0.40			
	OXTr rs53576 A/G-A/A	14.40	29.92	0.48			
	OXTr rs53576 G/G-A/A	-14.19	28.76	-0.49			
	M_Care	-0.73	0.85	-0.87			
	M_OverP	5.03**	1.71	2.95			
	F_Care	-1.52*	0.75	-2.04			
	F_OverP	-1.71	0.87	-1.96			
	Culture x OXTr rs53576 A/G-A/A	-27.14	32.77	-0.83			
	Culture x OXTr rs53576 G/G-A/A	-12.77	33.42	-0.38			
	Culture x M_Care	-0.10	0.91	-0.11			
	Culture x M_OverP	-4.89**	1.74	-2.81			
	Culture x F_Care	1.74*	0.80	2.19			
	Culture x F_OverP	1.73	0.93	1.87			
	OXTr rs53576 A/G-A/A x M_Care	0.04	0.97	0.05			
	OXTr rs53576 G/G-A/A x M_Care	0.78	0.89	0.87			
	OXTr rs53576 A/G-A/A x M_OverP	-5.27**	1.79	-2.96			
	OXTr rs53576 G/G-A/A x M_OverP	-4.57**	1.76	-2.59			
	OXTr rs53576 A/G-A/A x F_Care	1.25	0.86	1.46			
	OXTr rs53576 G/G-A/A x F_Care	1.60*	0.80	2.01			
	OXTr rs53576 A/G-A/A x F_OverP	2.18*	1.06	2.06			
	OXTr rs53576 G/G-A/A x F_OverP	1.27	0.96	1.33			
	Culture x OXTr rs53576 A/G-A/A x M_Care	1.24	1.06	1.17			
	Culture x OXTr rs53576 G/G-A/A x M_Care	-0.11	1.03	-0.11			
	Culture x OXTr rs53576 A/G-A/A x M_OverP	4.91**	1.83	2.67			
	Culture x OXTr rs53576 G/G-A/A x M_OverP	4.79**	1.84	2.61			
	Culture x OXTr rs53576 A/G-A/A x F_Care	-2.06*	0.93	-2.22			
	Culture x OXTr rs53576 G/G-A/A x F_Care	-1.44	0.90	-1.60			
	Culture x OXTr rs53576 A/G-A/A x F_OverP	-1.88	1.13	-1.66			
	Culture x OXTr rs53576 G/G-A/A x F_OverP	-1.09	1.10	-0.99			

Table 7. Hierarchical multiple regression on ECR-R avoidance for the total sample. Note. SE = standard error of unstandardized coefficient. *$p<.05$ **$p<.01$

t test (corrected *alpha* $= 0.025$) revealed that adult avoidance was significantly different between
Italian and Singaporean participants when they experienced a history of low maternal overprotection
($t = -2.91$, df = 162, *p*= .004, *d*= -0.51) (Figure 3**D**), but not when they had a past of high maternal
overprotection ($t = -0.88$, df = 147, *p*= .38, *d*= -0.15).
Whereas the two-way interaction between culture and paternal care (*p*> .025) was not considered
significant once applied *alpha* correction for the linear models (corrected *alpha* $= 0.025$).
Two further significant two-way interactions between OXTr levels and maternal overprotection on
adult avoidance were detected (A/G-A/A: β = -5.27, SE = 1.71, *t* = -2.96, *p*= .003; G/G-A/A: β =
-4.57, SE = 1.76, *t* = -2.59, *p*= .009). The distribution of genotypes A/A vs. A/G vs. G/G was not
significantly different between high vs. low maternal overprotection (X^2 (2) = 3.75, *p*= .15). Maternal
overprotection was positively associated with avoidance for A/A ($t = 1.32$, df = 92, *r* = 0.14, *p*= .19)
and G/G homozygotes ($t = 1.91$, df = 88, *r* = 0.20, *p*= .06), but not for the heterozygotes ($t = -0.31$, df =
127, *r* = -0.03, *p*= .75) (Figure 3**E**; red triangles = A/A; green squares = A/G; blue circles = G/G; lines
represent the linear models for A/A homozygotes-red, A/G-green and G/G-blue). No difference
between the slopes for A/A and A/G (z = -1.24, *p*= .21), for A/A and G/G (z = -0.44, *p*= .66) and for
A/G and G/G (z = -1.67, *p*= .09) was observed. In the only condition of high maternal overprotection
(corrected α = 0.017) the exploratory two-tailed post-hoc Student's *t* tests proved that adult avoidance
was not significantly different between A/A and A/G ($t = 1.34$, df = 97, *p*= .18, *d*= 0.28), A/A and
G/G ($t = 0.58$, df = 87, *p*= 57, *d*= 0.12), A/G and G/G ($t = 0.90$, df = 108, *p*= 37, *d*= 0.17) (Figure 3**F**).
Considerably, the two two-way interaction respectively between OXTr and paternal care and between
OXTr and paternal overprotection on adult avoidance did not fall below the level of acceptability
(*p*> .025).
Concerning the potential three-way interaction expected for the observational *HP3*, two significant
three-way interactions between culture, OXTr levels and maternal overprotection on adult avoidance
were discovered (A/G-A/A: β = 4.91, SE = 1.83, *t* = 2.67, *p*< .008; G/G-A/A: β = 4.79, SE = 1.84, *t* =
2.61, *p*= .009). In the Italian group, maternal overprotection was positively associated with avoidance
for all the genetic groups (A/A: $t = 1.86$, df = 12, *r* = 0.48, *p*= .09; A/G: $t = 1.39$, df = 33, *r* = 0.24, *p*= .17;
G/G: $t = 0.73$, df = 46, *r* = 0.11, *p*= .47) (Figure 3**G**; red triangles = A/A; green squares = A/G; blue
circles = G/G; lines show the linear models for A/A homozygotes-red, A/G-green and G/G-blue).
No difference between the slopes for all the Italian genetic groups was found (A/A-A/G: z = 0.80,
p= .21); A/A-G/G: z = 1.23, *p*= .11; A/G-G/G: z = 0.58, *p*= .28. In the Singaporean group, maternal
overprotection was positively associated with avoidance for A/A ($t = 0.90$, df = 78, *r* = 0.10, *p*= .37) and
G/G ($t = 1.59$, df = 40, *r* = 0.24, *p*= .12), but the same PBI dimension was negatively associated with
avoidance for A/G ($t = -0.85$, df = 92, *r* = -0.09, *p*= .40) (Figure 3**H**; red triangles = A/A; green squares
= A/G; blue circles = G/G; lines show the linear models for A/A homozygotes-red, A/G-green and
G/G-blue). Fisher's *z*-tests confirmed a significant difference between slopes for the A/G-G/G groups
(z = 1.75, *p*= .04) but not for the A/A-A/G (z = 1.23, *p*= .11) and the A/A-G/G (z = -0.74, *p*= .23)
groups. In the condition of low maternal overprotection the two-tailed post-hoc Student's *t* tests
(corrected *alpha* $= 0.025$) confirmed that adult avoidance was not significantly different between A/A
and G/G both in the Italian ($t = -0.36$, df = 30, *p*= .72, *d*= -0.15) and Singaporean group ($t = 0.74$, df =
61, *p*= .47, *d*= 0.21) (Figure 3**I**). Likewise, in the condition of high maternal overprotection (corrected
alpha $= 0.025$) adult avoidance was not significantly different between A/A and G/G both in the
Italian ($t = 2.28$, df = 28, *p*= .03, *d*= 1.04) and Singaporean group ($t = -0.94$, df = 57, *p*= .35, *d*= -0.25)
(Figure 3**J**).
Nevertheless, the last three-way interaction between culture, OXTr and paternal care approached but
did not reach the significance (*p*> .025).

3. Discussion

Early interactions have been widely proven to have a crucial role in shaping psychological mechanisms that drive social interactions later in life. However it is fundamental to take into account several factors that occur in the development of these psychosocial processes, such as genetic and cultural contributions. Starting from the evidence in literature suggesting the key-role of oxytocin [54,55] and parental bonding in influencing social behavior [21,56], the main purpose of this study was to examine the effects of the different genetic susceptibility and early interactions on the levels of anxiety and avoidance experienced in adult relationships in two different cultural contexts. Within this multicultural framework, the Italian and Singaporean samples were respectively considered as representative of a Western-dominant culture and a Eastern-dominant culture. Specifically, we focused on recalled parental bonding, assessed in terms of parental care and overprotection. Additionally, we included in our analyses the effect of allelic variation of one OXTr gene SNP, namely rs53576. We were expecting to find effects for the interaction between the genetic and perceived caregiving variables, with different influences in the groups (tested in *HP1* and *HP2*). Moreover, we explored a more intertwined model, adding the influence of the cultural environment (tested in *HP3*).

3.1. Gene-by-environment interaction within the Western-oriented sample and the Eastern-oriented sample evaluated independently

With regards to the Italian sample, the first step considered only the OXTr variation in explaining the levels of anxiety and avoidance. Perceived maternal features were progressively entered in the model, both resulting as strong predictors for anxiety, with care displaying a negative association and overprotection a positive one. In the subsequent steps, recalled paternal characteristics were added to the model, and care dimension for the father resulted as a predictor for anxiety as well. This might reflect the diversified, but equally important, contributions that both parents produce in their offspring, leading to different outcomes as a consequence [14]. In the model of interaction tested at the last stage, recalled maternal overprotection appears to explain levels of anxiety, with a positive correlation: the greater the overprotection felt, the higher the levels of experienced anxiety. Here, Italian individuals with a past of perceived maternal overprotection reported higher levels of anxiety than those who recalled a less controlling relationship with their mothers. Perceived maternal control during childhood, assessed in terms of overprotection, seems to be the construct that better explained level of anxiety in adult relationships. This result is in line with the theoretical framework of the development of attachment [57] and, furthermore, it is also supported by other studies [14,58–61]. The results suggests that both parents, to different extents, contribute in explaining the formation of anxiety-related patterns in adult social behavior. With regards to the experienced avoidance adulthood, we found that maternal care contributed to the unique variance as a predictor, negatively associated. Although no other variables alone were proved robust predictors, the interaction between genotype and early parental bonding dimensions increased substantially the explained variance in adult avoidance. However, due to the choice of a highly conservative correction method, the effect of maternal care emerging from the interactive model tested on avoidance did not reach the significance level. Interestingly, in the Italian sample, features of the mother appear to influence the formation of psychological constructs of close relationships more than those of the father, highlighting the importance of the specific parental role in explaining outcomes in adult relationships [62,63]. Moreover, within the final stages of the HMR, no significant effect was found for the interaction between OXTr rs53576 and parental bonding dimension on anxiety or avoidance, not confirming the first hypothesis. Overall, these results suggest that the quality of familiar bonds in this population, where higher levels of expressed emotions often characterize the relationship between parents and offspring [19,64,65], could affect the experience of close relationships in adulthood more independently from the genetic sensitivity, invalidating the starting *HP1*.

Concerning the Singaporean sample, the development of HMR on this population overall reveals distinct patterns of parental boding and genotype on adult attachment as a potential effect of the

Eastern culture. Concerning the adult anxiety, recalled paternal care and overprotection contributed more significantly to the model when respectively entered. Although the interaction between genotype and parental features did not improve the model in terms of the total amount of variance, an interaction effect between genotype and maternal overprotection emerged at this final stage. Such effect was marginally significant, thus did not persist when the correction was applied.

Unlike the Italian sample, neither genotype nor the recalled parental features resulted as a significant predictor of adult avoidance across the different steps of the model whose variance remains stable. In the final step, where all the targeted variables were included, maternal care was the strongest predictor of avoidance in adult close relationships with a negative trend. Moreover, the interaction between OXTr variations and maternal care reached the statistical significance as well. In particular, the two variations followed a mutually opposite pattern of association, displaying a slight positive correlation for maternal care over avoidance levels for G-carriers and a negative one for A/A homozygous. In line with *HP2*, the present cross-interaction supports the assumption of a differential sensitivity to environmental influences among carriers of distinctive genetic phenotype in OXTr rs53576 [54,55,66]. While evidence suggests overall effects of OXTr rs53576 on prosocial mechanisms, such as empathy, or seeking for emotional support, outcomes on attachment style are not consistent. Our results may reflect that boundary condition plays a stronger role when it comes to adult attachment dimensions [27,36,67], thus caution is needed in addressing A/A variant as a risk factor for non-optimal patterns of attachment. Currently there are few studies examining the effects of the interaction of recalled parental bonding and OXTr rs53576 on adult attachment dimensions in Eastern populations, considering the ethnic group separately [27,36,68], advocating the need for a focus on the importance of biological differences and distinct allelic distributions.

We observed that the effects in the two groups differ, especially in the relationship between recalled maternal bonding and attachment dimension in adult relationship, reflecting the typical traits of familiar bonds: with regards to the Italian sample, where great attention is projected on the child, and levels of emotions expressed are high [19], the construct of overprotection was associated to anxiety in adult relationship. According to the traits depicting individualistic culture, the person is expected to be self-reliant and not depending on other people for support, which in return might be reflected in greater anxiety expressed or experienced, instead of avoidance [69]. On the other hand, in the Singaporean sample, where interdependence represents a typical trait in social groups, starting from the family, avoidant pattern of behavior can describe a greater form of distress in relationships, compared to Western-oriented groups [69,70].

3.2. Gene-by-culture interactions in the total sample

While gene-by-environment framework focuses predominantly on external factors that characterize individual experience, the gene–culture interaction field of research focuses on the socio-cultural context. Within this theoretical frame, variants conferring genetic susceptibility can amplify typical cultural tendencies [38], such as interdependence [71]. Adopting a HMR, we aimed to explore the influence of the cultural environment, and of its interaction with genetic variations sensitive to external context, such as perceived parental caregiving.

In reference to the HMRs developed on the adult anxiety, culture was the first variable added to the model, explaining a significant portion of variance. The subsequent addition of genotype did not increased the variation in anxiety, but this increase was true for the addition of maternal and paternal care as well as paternal overprotection at different steps, respectively. Finally, the interaction between culture, genotype and parental bonding dimensions maximized the variance on adult anxiety. Despite this achievement, the model of interaction did no not reveal consistent effects on the adult anxiety.

In the context of the HMRs advanced for the adult avoidance, culture still was a key variable that is able to rise the variance significantly. Discordantly from the significant effects of the numerous parental features on the variation in anxiety, only maternal care enlarged the amount of explained variation in avoidance. In accordance with the final model observed from the HMRs on anxiety, the

highest quantity of total variance was reached when the interaction between culture, genotype and parental bonding dimensions was verified on avoidance.

From the model of three-way interaction, multiple effects were observed. Firstly, maternal overprotection positively predicted adult avoidance when the cultural and genotype influences were not further examined. Secondly, the cultural component played a dominant role over experienced avoidance in adult close relationships. We observed that the effects on the two groups are different, especially in the relationship between culture and maternal overprotection. Specifically, maternal overprotection was positively associated with adult avoidance for both Western and Eastern samples. However, the Western-oriented participants reported lower mean levels of avoidance from the partner than the Eastern-oriented individuals. Mousavi and colleagues, although framing their exploration in a wider perception of anxiety, including social phobia which is related to avoidant patterns of behavior, provided evidence that the perception of parental control or overprotection was reported in a stronger extent in a Caucasian sample [72].

Thirdly, the contributing action of the genetic predisposition driven by OXTr rs53576 was also identified. In this regard, the significant interaction between genotype and maternal overprotection predicted the adult avoidance when the participants were conceived together and independently from the cultural belonging to the group. However, the proper interpretation of this effect faced several difficulties in the light of the different alleles' frequency between Western and Eastern samples.

Analyses referring to the gene-by-culture frameworks displayed an interesting result: the different levels of associations between the genetic carriers A/A, A/G and G/G and maternal overprotection on the adult avoidance for the Western-oriented and Eastern-oriented sample discloses peculiar attachment patterns as a function of a differential genetic susceptibility to the external environment [38,73]. Here, the less expressed variation (A/A for the Italian sample, G/G for the Singaporean group) was the one more directly correlated to greater levels of avoidance for higher levels of perceived maternal overprotection. Similar results were found by Kim and colleagues when exploring the role of the culture using a gene-by-environment approach between individualistic (American) and collectivist (Korean) populations in investigating emotion suppression [73]. Avoidance, likewise emotion suppression, can be described as a coping strategy to deal with stressful experiences [74]. Analogously to findings of Kim and colleagues, levels of avoidance were observed more among G/G carriers in the Singaporean sample, compared to those carrying two copies of the A variant. Conversely, in the Italian group, G/G carriers showed significant lower levels of avoidance compared to the A/A carriers. These findings suggest that the allelic variation retained to confer greater sensitivity to environmental input is correlated to more optimal outcomes involving social and emotional strategies, according the specific societal context [38,73]. Interestingly, in the Singaporean sample, people carrying the A/G variation displayed an opposite trend in terms of negative relationship with avoidance compared to both the positive associations of A/A and G/G carriers to the same ECR-R dimension. In literature, evidence reports results for the heterozygous variation together with the less expressed homozygous pair of alleles in order to correct skewed distributions [73], hence this outcome needs caution when it comes to interpretation; the trend deductible from our results is that A/G carriers in the Singaporean sample tend to show an opposite pattern compared to G/G carriers, hence it is arguable that their experienced avoidance does not resent of the levels of perceived maternal control.

4. Conclusions

In this study we focused on two different social environmental: the family, within witch the person experiences early relationships and builds the first behavioral patterns, and the cultural context, in which the grown person adopts a combined set of individual and shared criterion in relating to others. One of the strengths of this study is to provide a further step in comparing two distinct environments, each consisting of parenting strategies and family composition. Another important element considered is the distinction between maternal and paternal bonding. In fact, although nowadays the competence regarding child care is becoming more and more equal between the two

parents, fatherhood is an area that still under-explored, especially in reference to the temporal dynamics of attachment in adulthood. Avoidance might have been more strongly correlated as it could play a mediating role between recalled parental features and anxiety [75]. In our approach, we divided the different combination of the single variants for the Italian and Singaporean group, in order to maximize the effects of the genetic distribution in the different ethnic groups. This study also presents limitations that are important to consider, starting from the cross-sectional approach that highlights a weaker reliance on retrospective reporting of parental bonding, compared to a longitudinal one. Another point is that, although we utilised original (for Singaporean group) and validated translated (for Italian group) version of the questionnaires, some conceptual-understanding issues could have led to different interpretation of the items, hence to altered scoring that might have influenced the results. Additionally, the numerosity of the sample represents a challenge in gene-by-environment interaction studies. In fact, limited to the allelic distribution in the total sample which comprises both the Italian and the Singaporean sample, the frequency of A/A and G/G homozygous and A/G heterozygotes was unbalanced, and this may have influenced the outcomes. Finally, for both of the countries, the sample was composed by undergraduate/post-graduate students, mainly belonging to Social Science or Psychological fields, who happened to be familiar with the theories beneath the study. Implication of our finding in the clinical work could cover an expanding area of interest in line with the current biopsychosocial model of health and psychological wellness, that is the influence of cultural components and early interactions in understanding the development of social behavior and related disorders. This would allow practitioners in child developmental and parenting fields to formulate interventions that take into account also on the cultural features of the person, which is a relevant topic, due to the increasing mobility of single people and families. Overall, while this study represents a further step in understanding the development of bonding mechanisms in different cultural contexts, it is at the same time a starting point for further exploration where more modern approaches could be combined, in order to have a more comprehensive understanding of the factors that might affect the dynamics of attachment, in both early stages and subsequent phases of life. Future research should extend the assessment by integrating psychological mechanisms underlying social behavior and expectations, which may play a mediating role between childhood reminiscence of parental care and experienced anxiety or avoidance in adulthood. Moreover, adult social relationship cannot exclude online social interactions; exploring the interaction between genes and environmental contexts over the experience of relationships in a virtual domain would allow researchers have access to a more structured and complete view of human relationships [76,77].

Supplementary Materials: Supplementary Results: *Italian Sample: Avoidance*; *Singaporean Sample: Anxiety*; *Total Sample: Anxiety*. **Supplementary Tables**: *Table S1*: Models applied into the six-steps hierarchical multiple regressions on the Italian, Singaporean and Total sample; *Table S2*: Correlation matrix among Questionnaire Subscales across all participants; *Table S3*: Hierarchical multiple regression on adult avoidance for the Italian participants; *Table S4*: Hierarchical multiple regression on adult anxiety for the Singaporean participants; *Table S5*: Hierarchical multiple regression on adult anxiety for all the participants; *Table S6*: Means and Standard Errors of ECR-R anxiety by the independent variables for the Italian participants; *Table S7*: Means and Standard Errors of ECR-R avoidance by the independent variables for the Singaporean participants; *Table S8*: Means and Standard Errors of ECR-R avoidance by the independent variables for all the participants assessed. **Supplementary Figures**: *Figure S1*: Comparison between levels of gender and age on anxiety and avoidance for the Italian participants; *Figure S2*: Comparison between levels of gender and age on anxiety and avoidance for the Singaporean participants; *Figure S3*: Comparison between levels of gender, age and culture on anxiety and avoidance for all the participants assessed.

Author Contributions: Conceptualization, I.C., B.L. and G.E.; methodology, A.B., I.C., B.L. and G.E.; formal analysis, A.B. and J.N.F.; investigation, A.B., I.C., B.L. and G.E.; data curation, I.C.; writing–original draft preparation, I.C. and A.B.; writing–review and editing, I.C, A.B., B.L., P.S. and G.E.; visualization, A.B.; supervision, B.L. and G.E.; funding acquisition, G.E. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by the NAP Start-up Grant M4081597 (GE) from Nanyang Technological University Singapore, the Singapore Ministry of Education ACR Tier-1 Grant (GE, PS), Singapore Ministry of Education ACR Tier1 (GE), Singapore Ministry of Education Social Science Research Thematic Grant

(MOE2016-SSRTG-017, PS). The founder agencies had no role in the conceptualization, design, data collection, analysis, decision to publish, or preparation of the manuscript.

Acknowledgments: The authors would like to thank Lim Mengyu and Reena Koh ChengYee (Nanyang Technological University) for their assistance in editing and proofreading the article, and Benedetta Corea for her support in graphical design. Explanatory icons or illustrations placed at the left side of Figure 3 have been obtained from the open-source websites Freepik and Pixabay.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Abbreviations

The following abbreviations are used in this manuscript:

ECR-R	Experience in Close Relationships - Revised
PBI	Parental Bonding Instrument
OXTr	Oxytocin receptor gene
HMR	Hierarchical Multiple Regression

Appendix A. Ethics

The research was approved by the Ethics Committee of Nanyang Technological University (IRB-2015-08-020-01) and of University of Trento (Prot-2017-019). Informed consent was obtained from all participants, and the study was conducted following the Declaration of Helsinki. The genetic assessment was conducted on anonymized biosamples at the Nanyang Technological University (Singapore) and at the Department of Neurobiology and Behavior at Nagasaki University (Japan). Results from questionnaires were anonymized at the beginning of the collection.

Appendix B. Data Repository

The data of this study can be found in the NTU's Data repository (DR-NTU Data) at the following address: https://doi.org/10.21979/N9/7NT0EG

References

1. Bowlby, J. Attachment and loss: Volume II: Separation, anxiety and anger. In *Attachment and Loss: Volume II: Separation, Anxiety and Anger*; London: The Hogarth Press and the Institute of Psycho-Analysis, 1973; pp. 1–429.

2. Parker, G.; Tupling, H.; Brown, L.B. A parental bonding instrument. *British journal of medical psychology* **1979**, *52*, 1–10.

3. Zafiropoulou, M.; Avagianou, P.A.; Vassiliadou, S. Parental bonding and early maladaptive schemas. *Journal of Psychological Abnormalities in Children* **2014**, *3*, 2329–95251000110.

4. Shin, H.; Lee, J.; Kim, B.; Lee, S.M. Students' perceptions of parental bonding styles and their academic burnout. *Asia Pacific Education Review* **2012**, *13*, 509–517.

5. Francis, D.D.; Meaney, M.J. Maternal care and the development of stress responses. *Current opinion in neurobiology* **1999**, *9*, 128–134.

6. Hane, A.A.; Cheah, C.; Rubin, K.H.; Fox, N.A. The role of maternal behavior in the relation between shyness and social reticence in early childhood and social withdrawal in middle childhood. *Social Development* **2008**, *17*, 795–811.

7. Picardi, A.; Caroppo, E.; Fabi, E.; Proietti, S.; Di Gennaro, G.; Meldolesi, G.N.; Martinotti, G. Attachment and parenting in adult patients with anxiety disorders. *Clinical practice and epidemiology in mental health: CP & EMH* **2013**, *9*, 157.

8. Rogosch, F.A.; Cicchetti, D.; Aber, J.L. The role of child maltreatment in early deviations in cognitive and affective processing abilities and later peer relationship problems. *Development and psychopathology* **1995**, *7*, 591–609.

9. Shaver, P.R.; Mikulincer, M. Attachment-related psychodynamics. *Attachment & human development* **2002**, *4*, 133–161.

10. Reti, I.M.; Samuels, J.; Eaton, W.W.; Bienvenu Iii, O.; Costa Jr, P.; Nestadt, G. Adult antisocial personality traits are associated with experiences of low parental care and maternal overprotection. *Acta Psychiatrica Scandinavica* **2002**, *106*, 126–133.

11. Spinrad, T.L.; Gal, D.E. Fostering prosocial behavior and empathy in young children. *Current opinion in psychology* **2018**, *20*, 40–44.

12. Morris, A.S.; Silk, J.S.; Steinberg, L.; Myers, S.S.; Robinson, L.R. The role of the family context in the development of emotion regulation. *Social development* **2007**, *16*, 361–388.

13. Porges, S.W. Social engagement and attachment: a phylogenetic perspective. *Annals of the New York Academy of Sciences* **2003**, *1008*, 31–47.

14. Rork, K.E.; Morris, T.L. Influence of parenting factors on childhood social anxiety: Direct observation of parental warmth and control. *Child & Family Behavior Therapy* **2009**, *31*, 220–235.

15. Ainsworth, M.D.S. *Infancy in Uganda: Infant care and the growth of love.*; Johns Hopkins Press, 1967.

16. Behrens, K.Y.; Hesse, E.; Main, M. Mothers' attachment status as determined by the Adult Attachment Interview predicts their 6-year-olds' reunion responses: A study conducted in Japan. *Developmental Psychology* **2007**, *43*, 1553.

17. Mesman, J.; van IJzendoorn, M.H.; Bakermans-Kranenburg, M.J. Unequal in opportunity, equal in process: Parental sensitivity promotes positive child development in ethnic minority families. *Child Development Perspectives* **2012**, *6*, 239–250.

18. Sun, Q.W.; Ng, K.M.; Guo, L. The link between parental bonding and adult attachment in Chinese graduate students: Gender differences. *The Family Journal* **2010**, *18*, 386–394.

19. Laudani, C.; Guzzo, G.; Cascio, V.L.; Pace, U.; Cacioppo, M. Does a Mediterranean model of family functioning in the perception of Italian and Spanish adolescents exist? A cross-national study. *Mediterranean Journal of Social Sciences* **2014**, *5*, 377.

20. Keller, H. Universality claim of attachment theory: Children's socioemotional development across cultures. *Proceedings of the National Academy of Sciences* **2018**, *115*, 11414–11419.

21. Cataldo, I.; Azhari, A.; Lepri, B.; Esposito, G. Oxytocin receptors (OXTR) and early parental care: an interaction that modulates psychiatric disorders. *Research in developmental disabilities* **2018**, *82*, 27–38.

22. Walum, H.; Lichtenstein, P.; Neiderhiser, J.M.; Reiss, D.; Ganiban, J.M.; Spotts, E.L.; Pedersen, N.L.; Anckarsäter, H.; Larsson, H.; Westberg, L. Variation in the oxytocin receptor gene is associated with pair-bonding and social behavior. *Biological psychiatry* **2012**, *71*, 419–426.

23. Tost, H.; Kolachana, B.; Hakimi, S.; Lemaitre, H.; Verchinski, B.A.; Mattay, V.S.; Weinberger, D.R.; Meyer-Lindenberg, A. A common allele in the oxytocin receptor gene (OXTR) impacts prosocial temperament and human hypothalamic-limbic structure and function. *Proceedings of the National Academy of Sciences* **2010**, *107*, 13936–13941.

24. Feldman, R. Oxytocin and social affiliation in humans. *Hormones and behavior* **2012**, *61*, 380–391.

25. Zink, C.F.; Meyer-Lindenberg, A. Human neuroimaging of oxytocin and vasopressin in social cognition. *Hormones and behavior* **2012**, *61*, 400–409.

26. Domes, G.; Heinrichs, M.; Gläscher, J.; Büchel, C.; Braus, D.F.; Herpertz, S.C. Oxytocin attenuates amygdala responses to emotional faces regardless of valence. *Biological psychiatry* **2007**, *62*, 1187–1190.

27. Rodrigues, S.M.; Saslow, L.R.; Garcia, N.; John, O.P.; Keltner, D. Oxytocin receptor genetic variation relates to empathy and stress reactivity in humans. *Proceedings of the National Academy of Sciences* **2009**, *106*, 21437–21441.

28. Smith, K.E.; Porges, E.C.; Norman, G.J.; Connelly, J.J.; Decety, J. Oxytocin receptor gene variation predicts empathic concern and autonomic arousal while perceiving harm to others. *Social neuroscience* **2014**, *9*, 1–9.

29. Truzzi, A.; Poquérusse, J.; Setoh, P.; Shinohara, K.; Bornstein, M.H.; Esposito, G. Oxytocin receptor gene polymorphisms (rs53576) and early paternal care sensitize males to distressing female vocalizations. *Developmental psychobiology* **2018**, *60*, 333–339.

30. Senese, V.P.; Shinohara, K.; Esposito, G.; Doi, H.; Venuti, P.; Bornstein, M.H. Implicit association to infant faces: Genetics, early care experiences, and cultural factors influence caregiving propensities. *Behavioural brain research* **2017**, *325*, 163–172.

31. Norman, G.J.; Hawkley, L.; Luhmann, M.; Ball, A.B.; Cole, S.W.; Berntson, G.G.; Caciopppo, J.T. Variation in the oxytocin receptor gene influences neurocardiac reactivity to social stress and HPA function: a population based study. *Hormones and behavior* **2012**, *61*, 134–139.

32. Thompson, R.J.; Parker, K.J.; Hallmayer, J.F.; Waugh, C.E.; Gotlib, I.H. Oxytocin receptor gene polymorphism (rs2254298) interacts with familial risk for psychopathology to predict symptoms of depression and anxiety in adolescent girls. *Psychoneuroendocrinology* **2011**, *36*, 144–147.

33. Wu, N.; Li, Z.; Su, Y. The association between oxytocin receptor gene polymorphism (OXTR) and trait empathy. *Journal of affective disorders* **2012**, *138*, 468–472.

34. Lucht, M.J.; Barnow, S.; Sonnenfeld, C.; Rosenberger, A.; Grabe, H.J.; Schroeder, W.; Völzke, H.; Freyberger, H.J.; Herrmann, F.H.; Kroemer, H.; others. Associations between the oxytocin receptor gene (OXTR) and affect, loneliness and intelligence in normal subjects. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* **2009**, *33*, 860–866.

35. Apicella, C.L.; Cesarini, D.; Johannesson, M.; Dawes, C.T.; Lichtenstein, P.; Wallace, B.; Beauchamp, J.; Westberg, L. No association between oxytocin receptor (OXTR) gene polymorphisms and experimentally elicited social preferences. *PloS one* **2010**, *5*, e11153.

36. Li, J.; Zhao, Y.; Li, R.; Broster, L.S.; Zhou, C.; Yang, S. Association of oxytocin receptor gene (OXTR) rs53576 polymorphism with sociality: a meta-analysis. *PLoS One* **2015**, *10*, e0131820.

37. Bakermans-Kranenburg, M.J.; van IJzendoorn, M.H. Oxytocin receptor (OXTR) and serotonin transporter (5-HTT) genes associated with observed parenting. *Social cognitive and affective neuroscience* **2008**, *3*, 128–134.

38. Kim, H.S.; Sherman, D.K.; Sasaki, J.Y.; Xu, J.; Chu, T.Q.; Ryu, C.; Suh, E.M.; Graham, K.; Taylor, S.E. Culture, distress, and oxytocin receptor polymorphism (OXTR) interact to influence emotional support seeking. *Proceedings of the National Academy of Sciences* **2010**, *107*, 15717–15721.

39. Fraley, R.C.; Waller, N.G.; Brennan, K.A. An item response theory analysis of self-report measures of adult attachment. *Journal of personality and social psychology* **2000**, *78*, 350.

40. Scinto, A.; Marinangeli, M.G.; Kalyvoka, A.; Daneluzzo, E.; Rossi, A. The use of the Italian version of the Parental Bonding Instrument (PBI) in a clinical sample and in a student group: an exploratory and confirmatory factor analysis study. *Epidemiology and Psychiatric Sciences* **1999**, *8*, 276–283.

41. Busonera, A.; Martini, P.S.; Zavattini, G.C.; Santona, A. Psychometric properties of an Italian version of the experiences in close relationships-revised (ECR-R) scale. *Psychological reports* **2014**, *114*, 785–801.

42. Parker, G. The parental bonding instrument: psychometric properties reviewed. *Psychiatric developments* **1989**, *7*, 317–335.

43. Bonassi, A.; Ghilardi, T.; Truzzi, A.; Cataldo, I.; Azhari, A.; Setoh, P.; Shinohara, K.; Esposito, G. Dataset on genetic and physiological adults' responses to social distress. *Data in brief* **2017**, *13*, 742–748.

44. Wu, S.; Jia, M.; Ruan, Y.; Liu, J.; Guo, Y.; Shuang, M.; Gong, X.; Zhang, Y.; Yang, X.; Zhang, D. Positive association of the oxytocin receptor gene (OXTR) with autism in the Chinese Han population. *Biological psychiatry* **2005**, *58*, 74–77.

45. Cataldo, I.; Neoh, M.J.Y.; Chew, W.F.; Foo, J.N.; Lepri, B.; Esposito, G. Oxytocin receptor gene and parental bonding modulate prefrontal responses to cries: a NIRS Study. *Scientific Reports* **2020**, *10*, 1–11.

46. Butovskaya, M.; Rostovtseva, V.; Butovskaya, P.; Burkova, V.; Dronova, D.; Filatova, V.; Sukhodolskaya, E.; Vasiliev, V.; Mesa, T.; Rosa, A.; others. Oxytocin receptor gene polymorphism (rs53576) and digit ratio associates with aggression: comparison in seven ethnic groups. *Journal of physiological anthropology* **2020**, *39*, 1–15.

47. Ravitz, P.; Maunder, R.; Hunter, J.; Sthankiya, B.; Lancee, W. Adult attachment measures: A 25-year review. *Journal of psychosomatic research* **2010**, *69*, 419–432.

48. Zijlstra, W.P.; van der Ark, L.A.; Sijtsma, K. Outliers in questionnaire data: Can they be detected and should they be removed? *Journal of Educational and Behavioral Statistics* **2011**, *36*, 186–212.

49. Bradley, R.H.; Corwyn, R.F.; Burchinal, M.; McAdoo, H.P.; García Coll, C. The home environments of children in the United States Part II: Relations with behavioral development through age thirteen. *Child development* **2001**, *72*, 1868–1886.

50. Elgar, F.J.; Mills, R.S.; McGrath, P.J.; Waschbusch, D.A.; Brownridge, D.A. Maternal and paternal depressive symptoms and child maladjustment: The mediating role of parental behavior. *Journal of abnormal child psychology* **2007**, *35*, 943–955.

51. Kogan, A.; Saslow, L.R.; Impett, E.A.; Oveis, C.; Keltner, D.; Saturn, S.R. Thin-slicing study of the oxytocin receptor (OXTR) gene and the evaluation and expression of the prosocial disposition. *Proceedings of the National Academy of Sciences* **2011**, *108*, 19189–19192.

52. Faul, F.; Erdfelder, E.; Buchner, A.; Lang, A.G. Statistical power analyses using G* Power 3.1: Tests for correlation and regression analyses. *Behavior research methods* **2009**, *41*, 1149–1160.

53. Diedenhofen, B.; Musch, J. cocor: A comprehensive solution for the statistical comparison of correlations. *PloS one* **2015**, *10*, e0121945.

54. Ebstein, R.; Israel, S.; Lerer, E.; Uzefovsky, F.; Shalev, I.; Gritsenko, I.; Riebold, M.; Salomon, S.; Yirmiya, N. Arginine vasopressin and oxytocin modulate human social behavior. *Annals of the New York Academy of Sciences* **2009**, *1167*, 87–102.

55. Ebstein, R.P.; Knafo, A.; Mankuta, D.; Chew, S.H.; San Lai, P. The contributions of oxytocin and vasopressin pathway genes to human behavior. *Hormones and behavior* **2012**, *61*, 359–379.

56. Feeney, J.A. Early parenting and parental attachment: Links with offspring's attachment and perceptions of social support. *Journal of Family Studies* **2002**, *8*, 5–23.

57. Bowlby, J. Developmental psychiatry comes of age. *The American journal of psychiatry* **1988**.

58. Parker, G.; Lipscombe, P. Influences on maternal overprotection. *The British Journal of Psychiatry* **1981**, *138*, 303–311.

59. Raudino, A.; Murray, L.; Turner, C.; Tsampala, E.; Lis, A.; De Pascalis, L.; Cooper, P.J. Child anxiety and parenting in England and Italy: The moderating role of maternal warmth. *Journal of Child Psychology and Psychiatry* **2013**, *54*, 1318–1326.

60. Kiel, E.J.; Kalomiris, A.E. Emotional Development and Anxiety. In *Handbook of Emotional Development*; Springer, 2019; pp. 665–693.

61. Smout, A.; Lazarus, R.S.; Hudson, J.L. The Relationship Between Parenting and Anxiety in Emerging Adulthood. *Cognitive Therapy and Research* **2020**, *44*, 182–195.

62. Land, L.N.; Rochlen, A.B.; Vaughn, B.K. Correlates of adult attachment avoidance: Men's avoidance of intimacy in romantic relationships. *Psychology of Men & Masculinity* **2011**, *12*, 64.

63. Gallagher, B.; Cartwright-Hatton, S. The relationship between parenting factors and trait anxiety: Mediating role of cognitive errors and metacognition. *Journal of Anxiety Disorders* **2008**, *22*, 722–733.

64. Pellerone, M.; Ramaci, T.; Herrera López, M.; Craparo, G. The role of identity development and decision making process on adult attachment: a cross-national study in sicilian and andalusian adolescents. *Clinical Neuropsychiatry* **2017**, *14*.

65. Pace, U.; Cacioppo, M.; Cascio, V.L.; Guzzo, G.; Passanisi, A. Are there similar or divergent transitions to adulthood in a Mediterranean context? A cross-national comparison of Italy and Spain. *Europe's journal of psychology* **2016**, *12*, 153.

66. Belsky, J.; Pluess, M. Beyond diathesis stress: differential susceptibility to environmental influences. *Psychological bulletin* **2009**, *135*, 885.

67. Gong, P.; Wang, Q.; Liu, J.; Xi, S.; Yang, X.; Fang, P.; Wang, B.; He, L.; Guo, W.; Zhang, M. The OXTR polymorphisms are not associated with attachment dimensions: A three-approach study. *Psychoneuroendocrinology* **2020**, *120*, 104780.

68. Gillath, O.; Shaver, P.R.; Baek, J.M.; Chun, D.S. Genetic correlates of adult attachment style. *Personality and Social Psychology Bulletin* **2008**, *34*, 1396–1405.

69. Sümer, N.; Yetkili, O. Cultural aspects of attachment anxiety, avoidance, and life satisfaction: Comparing the US and Turkey. In *Close relationships and happiness across cultures*; Springer, 2018; pp. 165–184.

70. Friedman, M.; Rholes, W.S.; Simpson, J.; Bond, M.; DIAZ-LOVING, R.; Chan, C. Attachment avoidance and the cultural fit hypothesis: A cross-cultural investigation. *Personal Relationships* **2010**, *17*, 107–126.

71. Luo, S.; Ma, Y.; Liu, Y.; Li, B.; Wang, C.; Shi, Z.; Li, X.; Zhang, W.; Rao, Y.; Han, S. Interaction between oxytocin receptor polymorphism and interdependent culture values on human empathy. *Social cognitive and affective neuroscience* **2015**, *10*, 1273–1281.

72. Mousavi, S.E.; Low, W.Y.; Hashim, A.H. Perceived parenting styles and cultural influences in adolescent's anxiety: A cross-cultural comparison. *Journal of Child and family Studies* **2016**, *25*, 2102–2110.

73. Kim, H.S.; Sherman, D.K.; Mojaverian, T.; Sasaki, J.Y.; Park, J.; Suh, E.M.; Taylor, S.E. Gene–culture interaction: Oxytocin receptor polymorphism (OXTR) and emotion regulation. *Social Psychological and Personality Science* **2011**, *2*, 665–672.

74. Kashdan, T.B.; Barrios, V.; Forsyth, J.P.; Steger, M.F. Experiential avoidance as a generalized psychological vulnerability: Comparisons with coping and emotion regulation strategies. *Behaviour research and therapy* **2006**, *44*, 1301–1320.

75. Fulton, J.J.; Kiel, E.J.; Tull, M.T.; Gratz, K.L. Associations between perceived parental overprotection, experiential avoidance, and anxiety. *Journal of Experimental Psychopathology* **2014**, *5*, 200–211.

76. Bonassi, A.; Cataldo, I.; Gabrieli, G.; Foo, J.N.; Lepri, B.; Esposito, G. Oxytocin receptor gene polymorphisms and early parental bonding interact in shaping instagram social behavior. *International journal of environmental research and public health* **2020**, *17*, 7232.

77. Carollo, A.; Bonassi, A.; Cataldo, I.; Gabrieli, G.; Tandiono, M.; Foo, J.N.; Lepri, B.; Esposito, G. The relation between Oxytocin Receptor Gene polymorphisms, adult attachment and Instagram sociability: an exploratory analysis **2020**.

Figure 3. Effects on avoidance for the total sample. (**A**). Effect of early maternal overprotection on the adult avoidance. (**B**). Comparison between the scores of ECR-R avoidance in low and high maternal overprotection. (**C**). Effect of the interaction between culture and early maternal overprotection on the adult avoidance. (**D**). Contrast between the scores of ECR-R avoidance in Italian (red) and Singaporean (blue). (**E**). Effect of the interaction between genotype and early maternal overprotection on the adult avoidance. (**F**). Contrast between the scores of ECR-R avoidance in A/A homozygotes (red), A/G heterozygotes (green) and G/G homozygotes (blue) divided into low and high maternal overprotection. (**G**). Effect of the interaction between genotype and early maternal overprotection on the adult avoidance for the Italian participants. (**H**). Effect of the interaction between genotype and early maternal overprotection on the adult avoidance for the Singaporean participants. (**I**). Comparison between the scores of ECR-R avoidance in A/A homozygotes (red), A/G heterozygotes (green) and G/G homozygotes (blue) divided into Italian and Singaporean participants for low maternal overprotection. (**J**). Contrast between the scores of ECR-R avoidance in A/A homozygotes (red), A/G heterozygotes (green) and G/G homozygotes (blue) divided into Italian and Singaporean participants for high maternal overprotection. *r*-values refer to Pearson's *r* correlations. (* p < 0.025)