Two neuronal peptides encoded from a single transcript regulate mitochondrial complex III in *Drosophila*

Justin A. Bosch[1,*], Berrak Ugur[2,†], Israel Pichardo-Casas[1], Jorden Rabasco[1], Felipe Escobedo[1], Zhongyuan Zuo[2], Ben Brown[3], Susan Celniker[3], David A. Sinclair[1], Hugo Bellen[2,4-6], and Norbert Perrimon[1,6,*].

1) Department of Genetics, Blavatnick Institute, Harvard Medical School, Boston, MA.
2) Department of Molecular and Human Genetics, BCM, Houston, TX 77030, USA
3) Lawrence Berkeley National Laboratory, Berkeley, CA.
4) Jan and Dan Duncan Neurological Research Institute, Texas Children's Hospital, Houston, TX 77030, USA
5) Department of Neuroscience, BCM, Houston, TX 77030, USA
6) Howard Hughes Medical Institute

[†]Current address: Departments of Neuroscience and Cell Biology, Howard Hughes Medical Institute, Yale University School of Medicine, New Haven, Connecticut 06510, USA

*Corresponding authors:

Justin A. Bosch and Norbert Perrimon (lead contact)
Harvard Medical School
77 Avenue Louis Pasteur
Dept. of Genetics, NRB 336
Boston, MA 02115
617-432-7672
Email: perrimon@genetics.med.harvard.edu (lead contact)
jabosch@hms.harvard.edu

Summary:

Naturally produced peptides (<100 amino acids) are important regulators of physiology, development, and metabolism. Recent studies have predicted that thousands of peptides may be translated from transcripts containing small open reading frames (smORFs). Here, we describe two peptides in *Drosophila* encoded by conserved smORFs, Sloth1 and Sloth2. These peptides are translated from the same bicistronic transcript and share sequence similarities, suggesting that they encode paralogs. Yet, Sloth1 and Sloth2 are not functionally redundant, and loss of either peptide causes animal lethality, reduced neuronal function, impaired mitochondrial function, and neurodegeneration. We provide evidence that Sloth1/2 are highly expressed in neurons, imported to mitochondria, and regulate mitochondrial complex III assembly. These results suggest that phenotypic analysis of smORF genes in *Drosophila* can provide a wealth of information on the biological functions of this poorly characterized class of genes.

93 **Keywords:** smORF, peptide, paralogs, mitochondria, complex III, bicistronic
94 transcript, neurodegeneration, *Drosophila*, CRISPR/Cas9

Introduction

Naturally produced peptides are regulators of metabolism, development, and physiology. Well-known examples include secreted peptides that act as hormones (PEARSON et al. 1993), signaling ligands (KATSIR et al. 2011), or neurotransmitters (SNYDER AND INNIS 1979). This set of peptides are produced by cleavage of larger precursor proteins (FRICKER 2005), peptides can also be directly translated from a transcript with a small open reading frame (smORF) (COUSO AND PATRAQUIM 2017; PLAZA et al. 2017; HSU AND BENFEY 2018; YEASMIN et al. 2018). Due to their small size (<100 codons), smORFs have been understudied. For example, smORFs are underrepresented in genome annotations (BASRAI et al. 1997), are theoretically a poor target for EMS mutagenesis, and are often ignored in proteomic screens. Consequently, there is growing interest in this class of protein-coding gene as a potentially rich source of novel bioactive peptides (MUDGE et al. 2022).

A major obstacle in identifying smORFs that encode functionally important peptides is distinguishing them from the enormous number of smORFs present in the genome by chance (e.g. 260,000 in yeast) (BASRAI et al. 1997). Many groups have identified and categorized smORFs with coding potential using signatures of evolutionary conservation, ribosomal profiling, and mass spectrometry (SAGHATELIAN AND COUSO 2015; COUSO AND PATRAQUIM 2017; PLAZA et al. 2017). Together, these approaches suggest there may be hundreds, possibly thousands, of unannotated smORF genes. However, these "omics" methods do not tell us which smORFs encode peptides with important biological functions.

Functional characterization of smORF genes in cell lines and model organisms has the potential to confidently identify novel peptides. Historically, unbiased genetic screens and gene cloning led to the fortuitous identification and characterization of smORF peptides (e.g. POLARIS (CASSON et al. 2002), RpL41 (SUZUKI et al. 1990), Nedd4 (KUMAR et al. 1993), *Drosophila* pri/tal (GALINDO et al. 2007)). More recently, candidate bioinformatically-predicted smORF-encoded peptides (aka SEPs) have been targeted for characterization (e.g., DWORF (NELSON et al. 2016), Elabela/toddler (CHNG et al. 2013; PAULI et al. 2014), Myomixer (BI et al. 2017), Myoregulin (ANDERSON et al. 2015), and Sarcolamban (MAGNY et al. 2013), and Hemotin (PUEYO et al. 2016)). Collectively, these studies have been invaluable for assigning biological functions to smORF peptides. Therefore, continued functional characterization is needed to tackle the enormous number of predicted smORF peptides.

Here, through an effort to systematically characterize human-conserved smORF genes in *Drosophila* (in preparation), we identified two previously unstudied smORF peptides CG32736-PB and CG42308-PA that we named Sloth1 and Sloth2 based on their mutant phenotypes. Remarkably, both peptides are translated from the same transcript and share amino acid sequence similarity, suggesting that they encode paralogs. Loss of function analysis revealed that

each peptide is essential for viability, and mutant animals exhibit defective
neuronal function and photoreceptor degeneration. These phenotypes can be
explained by our finding that Sloth1 and Sloth2 localize to mitochondria and play
an important role in complex III assembly. Finally, we propose that both peptides
bind in a shared complex. These studies uncover two new components of the
mitochondria and demonstrate how functional characterization of smORFs will
lead to novel biological insights.
192
193 **Results**
194
195 ***sloth1* and *sloth2* are translated from the same transcript and are likely**
196 **distantly related paralogs**
197
Current gene annotations for *sloth1* and *sloth2* (aka *CG32736* and *CG42308*,
respectively) indicate that they are expressed from the same transcript (Flybase,
Figure 1A), known as a bicistronic (or dicistronic) gene (BLUMENTHAL 2004;
CROSBY *et al.* 2015; KARGINOV *et al.* 2017). For example, nearby transcription
start sites (Figure 1A) are predicted to only generate a single transcript (HOSKINS
et al. 2011). In addition, a full-length transcript containing both smORFs is
present in the cDNA clone RE60462 (GenBank Acc# AY113525), which was
derived from an embryonic library (STAPLETON *et al.* 2002), and we detected the
full-length bicistronic transcript by RT-PCR amplification from total RNA from 3rd
instar larvae, adult flies, and S2R+ cells (Supplemental Figure 1). In addition, the
encoded peptides Sloth1 and Sloth2 have subtle sequence similarity (27%), are
similar in size (79aa and 61aa, respectively), and each contain a predicted single
transmembrane domain (Figure 1B). While this type of gene structure is relatively
rare in eukaryotes (BLUMENTHAL 2004; KARGINOV *et al.* 2017), there are known
cases in *Drosophila* of multicistronic transcripts encoding smORF paralogs – the
pri/*tal* locus (GALINDO *et al.* 2007) and the *Sarcolamban* locus (MAGNY *et al.*
2013). Furthermore, it is well known that paralogs are often found adjacent to
each other in the genome due to tandem duplication (TAYLOR AND RAES 2004).
Therefore, we propose that *sloth1* and *sloth2* are paralogs translated from the
same transcript.
218
Sloth1 and Sloth2 closely resemble their human orthologs (SMIM4 and
C12orf73*)*, based on sequence similarity, similar size, and presence of a
transmembrane domain (Figure 1B). Like Sloth1 and Sloth2, SMIM4 and
C12orf73 also have subtle amino acid sequence similarity to each other (Figure
1B). In addition, *sloth1* and *sloth2* are conserved in other eukaryotic species
(Figure 1C). Remarkably, *sloth1* and *sloth2* orthologs in choanoflagelate, sea
squirt, and lamprey exhibit a similar bicistronic gene architecture as *Drosophila*
(Figure 1C, Supplemental File 1). In contrast, *sloth1* and *sloth2* orthologs in
jawed vertebrates (e.g. mammals) are located on different chromosomes (e.g.
human Chr.3 and Chr.12, respectively). Interestingly, we only found one ortholog
similar to *sloth2* in the evolutionarily distant *Plasmodium*, and two orthologs
similar to *sloth2* in *Arabidopsis*, which are located on different chromosomes

(Figure 1C). Therefore, we hypothesize that the *sloth1* and *sloth2* ORFs
duplicated from an ancient single common ancestor ORF and became unlinked
in animals along the lineage to jawed vertebrates.
234
We next investigated *sloth1* and *sloth2* translation parameters and efficiency,
since their ORFs are frameshifted relative to each other (Figure 1A) and they are
not separated by an obvious internal ribosome entry site (IRES) (VAN DER KELEN
et al. 2009). Remarkably, only five nucleotides separate the stop codon of the
upstream ORF (*sloth1*) and the start codon of the downstream ORF (*sloth2*)
(Figure 1A). Therefore, *sloth1* should be translated first and inhibit translation of
sloth2, similar to the functions of so-called upstream ORFs (uORFs) (THOMPSON
2012). However, *sloth1* has a non-optimal Kozak sequence 5' to the start codon
(*ACAC*ATG) and *sloth2* has an optimal Kozak (*CAAA*ATG) (CAVENER 1987).
Therefore, scanning ribosomes may occasionally fail to initiate translation on
sloth1, in which case they would continue scanning and initiate translation on
sloth2, known as "leaky scanning" translation (THOMPSON 2012).
247
To test this translation model, we constructed an expression plasmid with the
Renilla Luciferase (*RLuc*) reporter gene downstream of *sloth1* (*sloth1-RLuc*),
while retaining non-coding elements of the original transcript (5' UTR, Kozak
sequences, 5bp intervening sequence) (Figure 1D). By transfecting this reporter
plasmid into *Drosophila* S2R+ cells, along with a *Firefly Luciferase* (*FLuc*) control
plasmid, we could monitor changes in translation of the downstream ORF by the
ratio of RLuc/FLuc luminescence. Using derivatives of the reporter plasmid with
Kozak or ATG mutations, we found that translation of the downstream ORF
increased when translation of *sloth1* was impaired (Figure 1E). Reciprocally,
translation of the downstream ORF was decreased when *sloth1* translation was
enhanced with an optimal Kozak. These results suggest that *sloth1* inhibits
translation of *sloth2*, and that balanced translation of both smORFs from the
same transcript might be achieved by suboptimal translation of *sloth1*.
261
262 ***sloth1* and *sloth2* are essential in *Drosophila* with non-redundant function**
263
To determine if *sloth1* and *sloth2* have important functions in *Drosophila*, we
used in vivo loss of function genetic tools. We used RNA interference (RNAi) to
knock down the *sloth1-sloth2* bicistronic transcript. Ubiquitous expression of an
shRNA targeting the *sloth1* coding sequence (Figure 2A) lead to significant
knockdown of the *sloth1-sloth2* transcript in 3rd instar larvae (Figure 2B), as
determined by two different primer pairs that bind to either the *sloth1* or *sloth2*
coding sequence. Ubiquitous RNAi knockdown of *sloth1-sloth2* throughout
development lead to reduced number of adult flies compared to a control (Figure
2C). This reduced viability was largely due to adult flies sticking in the food after
they eclosed from their pupal cases (Figure 2D). Escaper knockdown flies were
slow-moving and had 30% climbing ability compared to control flies (Figure 2E).
RNAi knockdown flies also had short scutellar bristles (Figure 2F).
276

We confirmed our RNAi results using CRISPR/Cas9 to generate somatic
knockout (KO) flies. By crossing flies ubiquitously expressing Cas9 (*Act-Cas9*)
with flies expressing an sgRNA that targets the coding sequence of either *sloth1*
or *sloth2* (Figure 2A, Supplemental Figure 2A), the resulting progeny will be
mosaic for insertions and deletions (indels) that cause loss of function in somatic
cells (PORT *et al.* 2014; XUE *et al.* 2014). Both *sloth1* and *sloth2* somatic KO flies
had significantly reduced viability compared to controls (Figure 2G). Furthermore,
escaper adults had short scutellar bristles (Figure 2H) and frequently appeared
sluggish. Importantly, similar phenotypes were observed when targeting either
sloth1 or *sloth2*.
287
Next, we further confirmed our loss of function results using CRISPR/Cas9 in the
germ line to generate KO lines for *sloth1* and *sloth2*. These reagents are
particularly important to test if *sloth1* and *sloth2* have redundant function by
comparing the phenotypes of single and double null mutants. We generated four
KO lines (Figure 2A, Supplemental Figure 2A-C): 1) a frameshift indel in *sloth1*
(*sloth1*-KO), 2) a frameshift indel in *sloth2* (*sloth2*-KO), 3) a 552 bp deletion of
the *sloth1* and *sloth2* reading frames (dKO), and 4) a knock-in of the reporter
gene *Gal4* that removes *sloth1* and *sloth2* coding sequences (*Gal4-KI*). Since
sloth1 and *sloth2* are on the X-chromosome, we analyzed mutant hemizygous
male flies. All four mutant lines were hemizygous lethal, which were rescued by a
genomic transgene (Figure 2I,), ruling out off-target lethal mutations on the X-
chromosome. Like RNAi and somatic KO results, rare mutant adult escaper flies
had slower motor activity (Figure 2J) and short scutellar bristles (Figure 2K).
Furthermore, the short scutellar bristle phenotype and slower motor activity could
be rescued by a genomic transgene (Figure 2J, K).
303
The phenotypic similarity of single and double mutants suggests that *sloth1* and
sloth2 are not functionally redundant. However, since both ORFs are encoded on
the same transcript, it is unclear if mutating one ORF will affect the other. For
example, a premature stop codon can induce non-sense mediated decay of an
entire transcript (NICKLESS *et al.* 2017). To address this possibility, we performed
additional fly lethality rescue experiments. First, transheterozygous female flies
(*sloth1-KO*/+, *sloth2-KO*/+) were viable and had normal scutellar bristles.
Second, we created single ORF versions of a genomic rescue transgene –
{*Δsloth1-sloth2*} and {*sloth1-Δsloth2*} (Supplemental Figure 2A). We found that
sloth1-KO lethality could only be rescued by {*sloth1-Δsloth2*}, and vice versa,
sloth2-KO lethality could only rescued by {Δ*sloth1-sloth2*} (Figure 2L).
Furthermore, single ORF rescue transgenes were unable to rescue the lethality
of *dKO* and *Gal4-KI* lines (Figure 2L). Third, we used the Gal4/UAS system
(BRAND AND PERRIMON 1993) to rescue mutant lethality with ubiquitously
expressed cDNA transgenes. These results showed that single ORF KOs could
only be rescued by expression of the same ORF (Figure 2L). Similar results were
found by expressing cDNAs encoding the human orthologs (Figure 2L). In all,
these results show that both *sloth1* and *sloth2* are essential, have similar loss of

function phenotypes, are not functionally redundant with one another, and are
likely to retain the same function as their human orthologs.
324
325   **Loss of *sloth1* and *sloth2* leads to defective neuronal function and**
326   **degeneration**
327
Since loss of *sloth1* and *sloth2* caused reduced adult mobility and climbing
defects (Figure 2E, J), we speculated that the two peptides normally play an
important role in the brain or muscle. To determine where *sloth1* and *sloth2* are
expressed, we used the *Gal4-KI* line as an in vivo transcriptional reporter. *Gal4-*
KI mobility defects and lethality could be rescued by expressing the entire
bicistronic transcript (*UAS-sloth1-sloth2*) (Figure 2J, L)*, or coexpression of both
smORFs as cDNA (*UAS-sloth1* and *UAS-sloth2*) (Figure 2L). Thus, the *Gal4-KI*
line is likely an accurate reporter of *sloth1* and *sloth2* expression. By crossing
Gal4-KI flies with a *UAS-GFP* fluorescent reporter, we observed strong GFP
expression in larval (Figure 3A, B) and adult brains (Figure 3C). In addition, *Gal4-*
KI is expressed in motor neurons at the larval neuromuscular junction (NMJ)
(Figure 3D) and in larval brain cells that are positive for the neuronal marker Elav
(Figure 3E).
341
We then tested if *sloth1* and *sloth2* were important for neuronal function by
measuring neuronal electrical activity in *dKO* animals. Electrical recordings taken
from the larval NMJ showed that *dKO* motor neurons have normal excitatory
junction potential (EJP) under resting conditions at 0.75 mM Ca $^{2+}$ (Supplemental
Figure 3). However, under high frequency stimulation (10hz), *dKO* NMJs could
not sustain a proper response (Figure 4A), indicating that there is a defect in
maintaining synaptic vesicle pools. Importantly, this phenotype is rescued by a
genomic transgene. To test if a similar defect is present in the adults, we
assessed phototransduction and synaptic transmission in photoreceptors via
electroretinogram (ERG) recordings (Wᴜ ᴀɴᴅ Wᴏɴɢ 1977; Hᴀʀᴅɪᴇ ᴀɴᴅ Rᴀɢʜᴜ
2001). ERGs recorded from young (1-3 days old) *dKO* photoreceptors showed
an amplitude similar to that of genomic rescue animals (Figure 4B). However,
upon repetitive light stimulation, ERG amplitudes were significantly reduced
(Figure 4B), suggesting a gradual loss of depolarization. Similar results were
observed when young flies were raised in 24hr dark (Figure 4C). Moreover, ERG
traces also showed a progressive loss of "on" and "off" transients (Figure 4B, C),
which is indicative of decreased synaptic communication between the
photoreceptor and the postsynaptic neurons. ERG phenotypes are rescued by a
full-length genomic rescue transgene, but not by single ORF rescue transgenes
(Figure 4B, C). To test if loss of both *sloth1* and *sloth2* lead to
neurodegeneration, we aged the animals for 4-weeks in 12hr light/dark cycle or
constant darkness and recorded ERGs. Similar to young animals, aged animals
raised in light/dark conditions also displayed a reduction in ERG amplitude upon
repetitive stimulation (Figure 4E). These results indicate that both *sloth1* and
sloth2 are required for sustained neuronal firing in larval motor neurons and adult
photoreceptors. Interestingly, similar mutant phenotypes in the NMJ and

photoreceptors are known to be due to defects in ATP production (VERSTREKEN
et al. 2005; SANDOVAL *et al.* 2014; JAISWAL *et al.* 2015).
370
In addition to measuring neuronal activity, we analyzed *dKO* neurons for changes
in morphology and molecular markers. Confocal imaging of the NMJ in *dKO* 3rd
instar larvae did not reveal obvious changes in synapse morphology or markers
of synapse function (Supplemental Figure 4). In contrast, using transmission
electron microscopy (TEM) of sectioned adult eyes, we observed reduced
photoreceptor number and aberrant morphology such as enlarged
photoreceptors and thinner glia in *dKO* animals (Figure 5A-C), suggestive of
degeneration. These phenotypes were rescued by a genomic transgene, but not
with single ORF rescue constructs (Figure 5A-C, Supplemental Figure 5).
Furthermore, these phenotypes were similar between young and aged flies, as
well as aged flies raised in the dark (Figure 5A-C, Supplemental Figure 5). It is
known that mutations affecting the turnover of Rhodopsin protein (Rh1) can lead
to photoreceptor degeneration (ALLOWAY *et al.* 2000; JAISWAL *et al.* 2015). To test
if this mechanism is occurring in *dKO* photoreceptors, we imaged Rh1 protein
levels using confocal microscopy. We observed Rh1 accumulation in
degenerating *dKO* photoreceptors in 4 week aged flies exposed to light (Figure
5D). However, Rh1 accumulation was milder in 4 week aged flies raised in the
dark (Supplemental Figure 6). These results point out that light stimulation, and
hence activity, enhance degeneration due to Rh1 accumulation in *dKO* animals.
390
391 **Sloth1 and Sloth2 localize to mitochondria and their loss impairs normal**
392 **respiration and ATP production**
393
394 Mitochondrial dysfunction in *Drosophila* is known to cause phenotypes that are
395 reminiscent of loss of *sloth1* and *sloth2*, such as pupal lethality, reduced neuronal
396 activity, photoreceptor degeneration, and Rh1 accumulation in photoreceptors
397 (JAISWAL *et al.* 2015). Therefore, we investigated the possible role of Sloth1 and
398 Sloth2 in mitochondria.
399
Prior to our work, a large-scale study of human protein localization suggested
that SMIM4 and C12orf73 localize to mitochondria in cultured cells (THUL *et al.*
2017). SMIM4 has a predicted mitochondrial targeting sequence using MitoFates
(FUKASAWA *et al.* 2015) (0.842), but C12orf73, Sloth1, and Sloth2 do not (.0016,
0.016, 0.009, respectively). In addition, SMIM4 and Sloth1 are predicted to
localize to the mitochondrial inner membrane using DeepMito (0.93 and 0.73,
respectively), but C12orf73 and Sloth2 are not (0.66 and 0.49, respectively)
(SAVOJARDO *et al.* 2020). To test if Sloth1/2 localize to mitochondria in
Drosophila, we transfected S2R+ cells with Sloth1-FLAG or Sloth2-FLAG. Both
Sloth1 and Sloth2 proteins colocalized with the mitochondrial marker ATP5α
(Figure 6A). Furthermore, Sloth1-FLAG and Sloth2-FLAG were enriched in
mitochondrial fractions relative to cytoplasmic fractions (Figure 6B). Similar
results were observed using stable S2R+ cell lines that express streptavidin

binding peptide (SBP) tagged Sloth1 or Sloth2 under a copper inducible promoter
(*MT-Sloth1-SBP* and *MT-Sloth2-SBP*) (Figure 6C).
415
Next, we raised antibodies to Sloth1/2 to determine their endogenous
localization. Using two independently generated antibodies for each peptide,
immunolocalization in larval brains from wild-type or *sloth1/2* dKO animals
showed no overlapping signal with a mitochondrial marker and no clear signal
above background (Supplemental Figure 7). Furthermore, we did not detect
Sloth1 or Sloth2 bands of the expected molecular weight on western blots from
wild-type S2R+ whole cell lysates or isolated mitochondria using anti-Sloth1, anti-
Sloth2, anti-SMIM4, or anti-C12orf73 (Supplemental Figure 8A-C). In contrast,
anti-Sloth1 western blots of mitochondria isolated from 3rd instar larvae and adult
thoraxes showed a <15kDa band that is absent from *sloth1/2* KO or RNAi
samples (Supplemental Figure 8D), suggesting this band corresponds to
endogenous Sloth1. Unfortunately, anti-Sloth2 failed to detect a similar band
under the same conditions (Supplemental Figure 8D).
429
Since our Sloth1/2 antibodies may not be sensitive enough to detect the
endogenous peptides, we generated a stable S2R+ cell line expressing *sloth1/2*
transcript under a copper inducible promoter (*MT-sloth1/2*) and induced
expression for 16hrs. Anti-Sloth1 and anti-Sloth2 western blots of mitochondria
isolated from *MT-sloth1/2* cells detected <15kDa bands that did not appear in
wild-type S2R+ cells, and thus are likely Sloth1 and Sloth2 peptides translated
from the overexpressed *sloth1/2* transcript (Supplemental Figure 8B).
Furthermore, Sloth1 and Sloth2 were enriched in *MT-sloth1/2* mitochondrial
fractions relative to cytoplasmic fractions (Figure 6D), similar to the results
obtained with FLAG and SBP-tagged peptides (Figures 6B-C). Based on their
amino acid sequence, Sloth1 and Sloth2 are predicted to run at 9.3kDa and
6.7kDa, respectively. While Sloth1 does appear to run larger than Sloth2, both
peptides run ~2kDa larger than expected (Figure 6D).
443
A method of assaying defects in mitochondrial function is measuring cellular
oxygen consumption from live cells with a Seahorse stress test. Since this
typically involves assaying a monolayer of cells, we generated KO S2R+ cell
lines using CRISPR/Cas9. Compared to control cells, single KO and double KO
S2R+ cells (Supplemental Figure 9A, B) had reduced basal respiration (Figure
7A, B), ATP production (Supplemental Figure 9C), and proton leaks
(Supplemental Figure 9D). Results were similar for single KO and dKO lines.
These results suggest that both *sloth1* and *sloth2* are required to support normal
mitochondrial respiration in S2R+ cells.
453
Next, we assayed *sloth1* and *sloth2* mutant flies for defects in mitochondrial
function. ATP levels are an important indicator of mitochondrial function (KANN
AND KOVACS 2007; GOLPICH *et al.* 2017) and mutations in *Drosophila*
mitochondrial genes can lead to reduced ATP levels (JAISWAL *et al.* 2015).
Indeed, *dKO* larvae had ~60% ATP compared to control larvae, which was

rescued by a genomic transgene (Figure 7C). Impaired mitochondrial function
can also lead to cellular stress responses, such as increased expression of the
mitochondrial chaperone Hsp60 (PELLEGRINO *et al.* 2013). Western blot analysis
showed that *Drosophila* Hsp60 was elevated in lysates from mutant larval brains
compared to control, and this effect was rescued by a genomic transgene (Figure
7D). Finally, mitochondrial dysfunction can cause changes in mitochondrial
morphology and number (TREVISAN *et al.* 2018). There were no obvious changes
in mitochondrial morphology in mutant larval motor neurons (Supplemental
Figure 4, Supplemental Figure 9E), and adult mutant photoreceptors contained
mitochondria with normal cristae (Figure 7E). In contrast, mitochondrial number
was increased in mutant photoreceptors in aged animals (Figure 7E,
Supplemental Figure 10A) and decreased in mutant photoreceptors in young
animals (Figure 7F, Supplemental Figure 10B). In all, these data suggest that
Sloth1 and Sloth2 localize to mitochondria and are important to support
respiration and ATP production.
474
475 **Sloth1/2 regulate respiratory complex III assembly**
476
While our study was in preparation, two studies demonstrated that human
SMIM4 and C12orf73 are inner mitochondrial membrane peptides important for
complex III assembly and physically interact with complex III subunits (ZHANG *et*
al. 2020; DENNERLEIN *et al.* 2021). If Sloth1 or Sloth2 have similar roles in
Drosophila, this could explain why *sloth1/2* mutant flies have reduced ATP
production.
483
To test for a role in Sloth1/2 in respiratory complex assembly, we visualized the
relative abundance of individual complexes and subunits in wild-type vs *sloth1/2*
loss of function animals. First, we resolved native respiratory complexes using
blue native polyacrylamide gel electrophoresis (BN-PAGE). Using mitochondria
isolated from adult thorax, we identified the five respiratory complexes (CI, CII,
CIII, CIV, CV) based on molecular weight and a previous study that established
this protocol (GARCIA *et al.* 2017). Importantly, a ~600kDa band corresponding to
complex III was diminished in mitochondria isolated from thoraxes with *sloth1/2*
knockdown (Figure 8A). Similarly, the complex III band was diminished in
mitochondria isolated from *sloth1/2* knockout 3rd instar larvae (Figure 8B). This
change was rescued by a wild-type genomic transgene, but not single paralog
transgenes (Figure 8B). Next, we detected individual respiratory subunits by
SDS-PAGE and western blotting of isolated mitochondria. Using antibodies that
recognize UQCR-C2, the fly homolog of human complex III subunit UQCRC2, we
found that the ~40kDa band corresponding to UQCR-C2 was diminished in
mitochondria isolated from *sloth1/2* RNAi adult thoraxes (Figure 8C), as well as
sloth1/2 knockout 3rd instar larvae (Figure 8D).
501
To test whether Sloth1/2 physically interact with subunits of mitochondrial
complex III, we performed co-immunoprecipitation experiments in transfected
S2R+ cells. SMIM4 and C12orf73 interact with complex III subunits UQCC1 and

UQCRFS1, respectively (ZHANG *et al.* 2020; DENNERLEIN *et al.* 2021). Therefore,
we tested if Sloth1 or Sloth2 could immunoprecipitate the fly homologs CG10075
(dUQCC1) or RFeSP (dUQCRFS1). Using Sloth1-FLAG as bait, we detected
CG10075-HA (Figure 8E) and RFeSP-HA (Figure 8F) binding to anti-FLAG
beads. In contrast, Sloth2-FLAG pulled-down CG10075-HA and RFeSP-HA
weakly or was at background levels (Figure 8E,F). Together, these results
suggest that Sloth1/2 are required for proper complex III assembly, mediated
through physical interaction with complex III subunits.
513
514 **Sloth1 and Sloth2 act in a stoichiometric complex**
515
We speculated that Sloth1 and Sloth2 could physically interact, based on the
observation that both share the same loss of function phenotypes and subcellular
localization. Indeed, some paralogs bind to the same protein complex
(SZKLARCZYK *et al.* 2008) and there is a tendency for proteins in the same
complex to be co-expressed (PAPP *et al.* 2003). To confirm this putative
interaction between Sloth1 and Sloth2, we used co-immunoprecipitation and
western blotting. This revealed that Sloth1-FLAG could immunoprecipitate
Sloth2-HA (Figure 9A), and reciprocally Sloth2-FLAG (Figure 9B) could
immunoprecipitate Sloth1-HA. Interestingly, the levels of tagged peptide in cell
lysates were higher when the opposite peptide was overexpressed (Figure 9A,B).
Proteins in a complex commonly have important stoichiometry and unbound
proteins can be degraded to preserve this balance (PAPP *et al.* 2003; SOPKO *et*
al. 2006; VEITIA *et al.* 2008; PRELICH 2012; BERGENDAHL *et al.* 2019).
Furthermore, imbalanced protein complex stoichiometry can lead to
haploinsufficient or dominant negative phenotypes (PAPP *et al.* 2003; SOPKO *et*
al. 2006; VEITIA *et al.* 2008; PRELICH 2012; BERGENDAHL *et al.* 2019).
532
To test this possibility for Sloth1/2, we overexpressed either *sloth1* or *sloth2* in
vivo. Low-level ubiquitous overexpression (using *da-Gal4*) of either *UAS-sloth1*
or *UAS-sloth2* cDNA had no effect on adult fly viability (Figure 2L). To increase
expression levels, we used the strong ubiquitous driver *tub-Gal4*. Whereas
tub>sloth1 flies were viable as adults, *tub>sloth2* animals were 100% pupal lethal
(Figure 9C). However, *tub>sloth2* animals could be rescued to adulthood by co-
expression of *sloth1*. Importantly, this rescue was not due to dilution of the Gal4
transcription factor on two *UAS* transgenes, since co-expression of *UAS-*
tdtomato did not rescue *tub>sloth2* lethality. Finally, *tub-Gal4* overexpression of
the entire *sloth1-sloth2* bicistronic transcript resulted in viable adult flies. In all,
these results suggest that Sloth1 and Sloth2 interact in a complex where their
stoichiometric ratio is important for normal function.
545
546 **Discussion**
547
Here, we have assigned new functions to two previously uncharacterized smORF
peptides. Sloth1 and Sloth2 appear to be distantly-related paralogs, yet each is
important to support mitochondrial and neuronal function in *Drosophila*. We

propose a model where Sloth1 and Sloth2 peptides are translated from the same
transcript, imported into mitochondria where they interact with each other and
complex III to promote its assembly (Figure 10). Our results are supported by two
recent studies published during preparation of this manuscript, in which human
Sloth1 (SMIM4) and Sloth2 (C12orf73/Brawnin) were discovered as novel
mitochondrial complex III assembly factors in cultured human cells and zebrafish
(ZHANG *et al.* 2020; DENNERLEIN *et al.* 2021).
558
Muti-cistronic genes are relatively rare in eukaryotes, but some have been
characterized in *Drosophila* (GALINDO *et al.* 2007; MAGNY *et al.* 2013) and
mammals (KARGINOV *et al.* 2017). Similar to operons in prokaryotes, it is thought
that multicistronic transcripts allow for coordinated expression of proteins in the
same pathway or complex (KARGINOV *et al.* 2017). Indeed, the similarity of loss of
function phenotypes between *sloth1* and *sloth2* suggest that they function
together in the same pathway/complex. Interestingly, 44/196 annotated
bicistronic genes in *Drosophila* contain two ORFs with homology to each other
(Flybase, DIOPT), and a recent study suggests that human bicistronic genes
containing a smORF frequently encode physically interacting peptide/protein pair
(CHEN *et al.* 2020). Therefore, related peptides encoded on the same transcript
may be a prevalent phenomenon in eukaryotes. ORF translation in multicistronic
transcripts can occur by different mechanisms, such as re-initiation of translation,
IRES, or leaky ribosome scanning (VAN DER KELEN *et al.* 2009). Our data and
observations support leaky scanning, and we propose a model whereby both
peptides are translated because *sloth1* contains a non-optimal Kozak sequence.
575
The presence of *sloth1* and *sloth2* orthologs in many eukaryotic species suggest
that their function is likely broadly conserved. Indeed, we could rescue the
lethality of *sloth1* and *sloth2* mutant flies by expressing their human counterparts.
Interestingly, *Plasmodium* and *Arabidopsis* only have homologs with similarity to
sloth2. Perhaps *sloth2* maintained functions more similar to its common ancestor
with *sloth1*. We were unable to identify homologs in some eukaryotes such as
yeast, though their amino acid sequence may simply be too diverged for
detection using bioinformatic programs such as BLAST.
584
The physical interactions of Sloth1-Sloth2, Sloth1-RFeSP, and Sloth1-CG10075,
and complex III assembly defects in *sloth1/2* loss of function animals, suggest
that Sloth1/2 together regulate complex III assembly. Indeed, Sloth1 is
bioinformatically predicted to localize to the mitochondrial inner membrane
(DeepMito), and Sloth1 and Sloth2 have predicted transmembrane domains
(TMHMM 2.0), suggesting they interact with complex III at the inner membrane.
This is supported by data showing SMIM4 and C12orf73 are integral membrane
proteins in the mitochondrial inner membrane (ZHANG *et al.* 2020; DENNERLEIN *et*
al. 2021). In addition, our data suggests that Sloth1 and Sloth2 interact in a
stoichiometric manner, explaining why single mutants have the same phenotype
as double mutants. This is supported by the finding that SMIM4 protein levels are
dependent on the presence of C12orf73 and vice versa (DENNERLEIN *et al.* 2021).

Perhaps maintenance of the proper ratio of Sloth1/2 is an important factor for
optimal complex III assembly. Future experiments could address whether Sloth1
and Sloth2 directly bind each other, or if they require complex III subunits for
physical association.
601
Several observations and experiments suggest that Sloth1/2 peptides do not
have equivalent function. The two peptides have weak homology to each other
(27% identity) and Sloth1 has 18aa (30%) more than Sloth2, suggesting
divergence of function. Unlike Sloth1, Sloth2 does not have a clear
mitochondrial-targeting signal. Perhaps Sloth2 has a cryptic signal that is not
recognized by prediction software, or Sloth2 may be co-imported with Sloth1.
Furthermore, we could not detect robust immunoprecipitation of RFeSP or
CG10075 using Sloth2 as bait. Perhaps Sloth2 binds complex III indirectly
through Sloth1, or Sloth 2 binds a different complex III subunit. More likely is that
both Sloth1 and Sloth2 need to be present for binding to complex III, and the
endogenous Sloth1 present under conditions of Sloth2-FLAG overexpression is
insufficient for co-IP assays. Sloth2 may also be less stable than Sloth1, which
could potentially explain why were unable to detect endogenous Sloth1 using
anti-Sloth1 antibodies. Interestingly, only strong overexpression of Sloth2, and
not Sloth1, was lethal to flies. Future studies may elucidate the mechanism
explaining these functional differences in Sloth1/2.
618
Neurons have a high metabolic demand and critically depend on ATP generated
from mitochondria to support processes such as neurotransmission (VERSTREKEN
et al. 2005; KANN AND KOVACS 2007). Therefore, it is not unexpected that
neurodegenerative diseases are frequently associated with mitochondrial
dysfunction (GOLPICH *et al.* 2017). We find similar results in *Drosophila*, where
loss of *sloth1* and *sloth2* leads to defects in mitochondrial function, impaired
neuronal function, photoreceptor degeneration, and Rh1 accumulation in
photoreceptors. Despite finding that the *Gal4-KI* reporter was strongly expressed
in neurons and could rescue *sloth1/2* lethality, it is likely these peptides play
important roles in other cell types. For example, publicly available RNA-seq data
suggest that they are ubiquitously expressed (Flybase). In addition, neuronal
expression of *sloth1* or *sloth2* was unable to rescue mutant lethality (Figure 2L).
Furthermore, we observed *sloth1/2* loss of function phenotypes in dissected adult
thoraxes, which are composed of mostly muscle. At present, there are no
reported human disease-associated mutations in *SMIM4* and *C12orf73*.
Mutations in these genes might not cause disease, or they might cause lethality.
It is also possible that the lack of functional information on these genes has
hampered identification of disease-associated mutations.
637
638 There is great interest in identifying the complete mitochondrial proteome (CALVO
639 *et al.* 2016), so it is remarkable that Sloth1/2 have been largely missed in
640 proteomic or genetic screens for mitochondrial components. For example, they
641 are not present in bioinformatic and proteomic datasets of fly mitochondrial
642 proteins (SARDIELLO *et al.* 2003; CHEN *et al.* 2015), nor in genetic screens of

lethal mutations on the X-chromosome affecting nervous system maintenance (YAMAMOTO *et al.* 2014). It is possible that the small size of these peptides lead to this discrepancy; due to less frequent mutations in these ORFs, or fewer tryptic products for MS. It is also possible that these peptides form weak interactions with mitochondrial proteins, preventing their immunoprecipitation. Recently, human SMIM4 was identified in a proteomic screen (DENNERLEIN *et al.* 2021), human C12orf73 was identified in two proteomics screens (LIU *et al.* 2018; ANTONICKA *et al.* 2020) and a bioinformatic screen (ZHANG *et al.* 2020), and mouse SMIM4 was identified in a proteomics screen (BUSCH *et al.* 2019).

Our discovery of *sloth1* and *sloth2* highlights the effectiveness of loss of function genetics for identifying smORF genes with important biological functions. Recent technical advances such as genome engineering (e.g. CRISPR/Cas9) and massively parallel profiling have the potential to rapidly assign functions to many uncharacterized smORFs (GUO *et al.* 2018; CHEN *et al.* 2020). For example, investigation of uncharacterized smORF genes may yield additional important mitochondrial components. Indeed, there is a greater tendency for annotated human smORF peptides to localize to mitochondria (72/719, 10%) compared to the whole proteome (1228/20351, 6%) (UniProt). Interestingly, ~40 smORF peptides function at the human mitochondrial inner membrane (UniProt), such as the Complex III member UQCRQ (82aa) (USUI *et al.* 1990) and the recently described Mitoregulin/MoxI (56aa) that regulates the electron transport chain and fatty acid β-oxidation (MAKAREWICH *et al.* 2018; STEIN *et al.* 2018; CHUGUNOVA *et al.* 2019). Therefore, modulation of protein complexes in the inner mitochondrial membrane may be a common function of smORF peptides. As functional annotation of hundreds, perhaps thousands, of smORF genes is becoming easier, many new biological insights are likely to emerge from their analyses.

Acknowledgements:

We thank the TRiP and DRSC for help generating transgenic flies, Dr. Marcia Haigis for use of a Seahorse XF analyzer, Claire Hu, Tera Levin, and Dan Richter for bioinformatics help, Lucy Liu for assistance mounting larvae to image the NMJ, Thai LaGraff for help with qPCR, and Rich Binari and Cathryn Murphy for general assistance. We thank members of the BDGP for discussions. We also thank the HMS MicRoN (Microscopy Resources on the North Quad) Core. J.A.B. was supported by the Damon Runyon Foundation. This work was supported by NIH grants R01GM084947, R01GM067761, R24OD019847, and NHGRI HG009352 (S.E.C). N.P. is an investigator of the Howard Hughes Medical Institute.

Author Contributions:

Conceptualization, J.A.B., B.U., I.P., B.B., S.C., H.B., N.P.; Methodology, J.A.B., B.U., I.P., N.P.; Investigation, J.A.B., B.U., I.P., J.R., F.E., Z.Z.; Writing – Original Draft, J.A.B.; Writing – Review & Editing, J.A.B., B.U., I.P., J.R., F.E., Z.Z., S.C.,

689   H.B., N.P.; Supervision, B.B., S.C., D.A.S., H.B., N.P.; Funding acquisition,
690   J.A.B., B.B., S.C., N.P.
691

698
699   **Figure titles and legends:**
700
Figure 1: Bicistronic gene structure of the smORFs *sloth1* and *sloth2*. A.
Bicistronic gene model for *sloth1* and *sloth2*. Zoom in shows intervening
sequence (*GCAAA*) between *sloth1* stop codon and *sloth2* start codon. **B.**
Comparison of protein structure, amino acid length size, and amino acid percent
identity between *Drosophila* and Human orthologs. Shaded rectangle indicates
predicted transmembrane (TM) domain. **C.** Phylogenetic tree of *sloth1* and *sloth2*
orthologs in representative eukaryotic species. Linked gene structure (candidate
bicistronic transcript or adjacent separate transcripts) is indicated by a black line
connecting red and blue squares. **D.** Plasmid reporter structure of *pMT-sloth1-*
Rluc and derivatives. Kozak sequences upstream of start codon are underlined.
Mutations indicated with shaded grey box. pMT= Metallothionein promoter. RLuc
= Renilla Luciferase. **E.** Quantification of RLuc luminescence/Firefly Luciferase,
normalized to *pMT-sloth1-Rluc*, for each construct. Significance of mutant
plasmid luminescence was calculated with a T-Test comparing to *pMT-sloth1-*
Rluc. Error bars are mean with SEM. **** $P \leq 0.0001$. N=4 biological replicates.
716
Figure 2: *sloth1* and *sloth2* loss of function analysis. A. *sloth1-sloth2*
transcript structure with shRNA and sgRNA target locations, primer binding sites,
in/del locations, and knock-in Gal4 transgene. **B.** qPCR quantification of RNAi
knockdown of the *sloth1-sloth2* transcript. Significance of fold change knockdown
was calculated with a T-Test comparing to *da>attP40* for PD43265 and
PD43573. Error bars show mean with SEM. P-values *** $P \leq 0.001$. N=6. **C.**
Quantification of adult fly viability from *sloth1-sloth2* RNAi knockdown. Fly cross
schematic (left) and graph (right) with percentage of progeny with or without the
CyO balancer. Ratios of balancer to non-balancer were analyzed by Chi square
test, **** $P \leq 0.0001$. Sample size (N) indicated on graph. **D.** Pictures of fly food
vials, focused on the surface of the food. *da>shRNA* flies are frequently found
stuck in the fly food. **E.** Quantification of adult fly climbing ability after *sloth1* and
sloth2 RNAi. Significance calculated with a T-test, **** $P \leq 0.0001$. Error bars
show mean with SD. N=3 biological replicates. **F.** Stereo microscope images of
adult fly thorax to visualize the scutellar bristles. RNAi knockdown by *da-Gal4*
crossed with either *attP40* or *UAS-shRNA*[JAB200]. Arrowheads point to the two
longest scutellar bristles. **G.** Quantification of adult fly viability from *sloth1-sloth2*
somatic knockout. Fly cross schematic (left) and graph (right) with percentage of

progeny with or without the CyO balancer. Ratios of balancer to non-balancer
were analyzed by Chi square test, **** P≤0.0001. Sample size (N) indicated on
graph. **H.** (Left) Stereo microscope images of adult fly thorax to visualize the
scutellar bristles. Somatic knockout performed by crossing *Act-Cas9* to sgRNAs.
(Right) Quantification of the frequency of adult flies with at least one short
scutellar bristle after somatic KO of *sloth1* or *sloth2.* Sample sizes indicated on
graph. Arrowheads point to the two longest scutellar bristles. **I.** Quantification of
adult fly viability from *sloth1-sloth2* hemizygous knockout in males and rescue
with a genomic transgene or *UAS-sloth1-sloth2* transgene. Fly cross schematic
(left) and graph (right) with percentage of male progeny with or without the FM7c
balancer. Sample size (N) indicated on graph. **J.** Still images from video of adult
flies inside plastic vials. Images are 5 seconds after vials were tapped. Adult flies
climb upward immediately after tapping. All flies are males. Each vial contains 10
flies, except dKO, which contains 5 flies. **K.** Stereo microscope images of adult
male fly thorax to visualize the scutellar bristles. *attP40* is used as a negative
control. Arrowheads point to the two longest scutellar bristles. **L.** Hemizygous
mutant male genetic rescue experiments.
752
Figure 3. *sloth1-sloth2* are expressed in neurons A. Fluorescent stereo
microscope images of 3rd instar larvae expressing GFP with indicated genotypes.
B. Fluorescent compound microscope image of 3rd instar larval brain expressing
UAS-GFP. DAPI staining labels nuclei. **C.** Confocal microscopy of adult brain
with indicated genotypes. Anti-HRP staining labels neurons. **D.** Confocal
microscopy of the 3rd instar larval NMJ at muscle 6/7 segment A2 expressing
UAS-GFP. Anti-FasII staining labels the entire NMJ. **E.** Confocal microscopy of
the 3rd instar larval ventral nerve cord (VNC) expressing *Gal4-KI, UAS-GFP-nls*.
GFP-nls is localized to nuclei. Anti-Elav stains nuclei of neurons. Arrow indicates
example nuclei that expresses UAS-GFP and is positive for Elav.
763
Figure 4. *sloth1-sloth2* are important for neuronal function. A. Traces of
electrical recordings from 3rd instar larval NMJ in control, *dKO*, and
dKO+genomic rescue animals over 10 minutes under high frequency stimulation
(10 Hz). Graph on right is a quantification of the relative excitatory junction
potential (EJP) for indicated genotypes. Error bars show mean with SD. N ≥ 5
larvae per genotype. Significance for each genotype was calculated with a T-Test
comparing to control flies. **B-D.** Traces of electroretinogram (ERG) recordings
from adult eye photoreceptors upon repetitive stimulation with light (left) and
quantification of the relative ERG amplitude for indicated genotypes (right). Error
bars show mean with SD. N ≥ 6 larvae per genotype. ** P≤0.01, *** P≤0.001.
Significance for each genotype was calculated with a T-Test comparing to control
flies. **B.** Recordings were taken from 1-3 days post-eclosion animals that were
raised in a 12hr light/dark cycle. "On" and "Off" transients indicated by closed and
open arrowhead, respectively. **C.** Recordings were taken from 1-3 days post-
eclosion animals that were raised in a 24hr dark. **D.** Recordings were taken from
four week aged animals that were raised in a 12hr light/dark cycle.
780

Figure 5. Loss of *sloth1-sloth2* causes neurodegeneration. A-C. Transmission electron microscopy (TEM) images of sectioned adult eye photoreceptors (left) and quantification of photoreceptor number and aberrant photoreceptors (right). Scalebar is 2µm. Filled red arrows indicate dead or dying photoreceptors. Open red arrows indicate unhealthy photoreceptors. Error bars show mean with SD. N ≥ 8 ommatidium per genotype. **A.** 4 weeks old raised in a 12hr light/dark cycle. **B.** 3 days old raised in a 12hr light/dark cycle. **C.** 4 weeks old raised in 24hr dark. **D.** Confocal microscopy of adult eye photoreceptors stained with phalloidin (green) and anti-Rh1 (red). Animals were 4 weeks old and raised in a 12hr light/dark cycle. Arrowheads indicate photoreceptors with higher levels of Rh1.

Figure 6. Sloth1 and Sloth2 localize to mitochondria. A. Confocal microscopy of S2R+ cells transfected with Sloth1-FLAG or Sloth2-FLAG and stained with anti-FLAG (green) and anti-ATP5alpha (red). DAPI (blue) stains nuclei. **B-D.** SDS-PAGE and western blotting of S2R+ cellular fractions. WCL = Whole Cell Lysate, cyto. = cytoplasmic lysate, mito. = mitochondrial lysate. Mitochondrial control = ATP5alpha, cytoplasmic control = alpha-tubulin. Each lane loaded equal amounts of protein (15µg/lane). Blots were stripped and reprobed after detection of each antigen. **B.** Transfected Sloth1-FLAG or Sloth2-FLAG. **C.** Stable cells expressing copper-inducible Sloth1-SBP or Sloth2-SBP. **D.** Stable cells expressing copper-inducible Sloth1-SBP or Sloth2-SBP.

Figure 7. *sloth1-sloth2* are important for mitochondrial function. A. Seahorse mitochondrial stress report for wildtype S2R+ and dKO #1 cells. Error bars show mean with SD. N=6 for each genotype. **B.** Quantification of basal OCR (timepoint 3) in panel A and including data from single KO and additional dKO cell lines. Significance of KO lines was calculated with a T-test compared to S2R+. Error bars show mean with SD. **** P≤0.0001. N=6 for each genotype. **C.** Quantification of ATP levels in 3rd instar larvae. Error bars show mean with SEM. N = 3 experiments. **D.** Western blot from lysates of 3rd instar larval brains. **E-F.** TEM images of sectioned adult photoreceptors (left) and quantification of mitochondria number (right). Mitochondria are indicated with red dots. Error bars show mean with SD. Sample size indicated on graph. **E.** Adult flies are 4 weeks old and raised in a 12hr light/dark cycle. **F.** Adult flies are 3 days old and raised in a 12hr light/dark cycle.

Figure 8. Sloth1 and Sloth2 physically interact with complex III and regulate its assembly. A-B. Blue native PAGE gel of mitochondria isolated from **A.** 10 adult thoraxes and **B.** 10 whole 3rd instar larvae of indicated genotype. Bands corresponding to native respiratory complexes are indicated with arrowheads. **C-D.** SDS-PAGE and western blotting of mitochondria isolated from **C.** adult thorax and **D.** whole 3rd instar larvae of indicated genotype. Each lane loaded equal amount of protein (15µg). Blots were stripped and reprobed after detection of each antigen. **E-F.** Western blots from co-immunoprecipitation experiments in transfected S2R+ cells using Sloth1-FLAG and Sloth2-FLAG as bait and either **E.**

827 RFeSP-HA or **F.** CG10075-HA as prey. Blots were striped and reprobed after
828 detection of each antigen. Arrowheads indicated expected band, asterisks
829 indicate unknown bands.
830
Figure 9. Sloth1 and Sloth2 act in a stoichiometric complex. A-B. Western
blots from co-immunoprecipitation experiments in transfected S2R+ cells. **A-B.**
Immunoprecipitation using Sloth1-FLAG and Sloth2-FLAG as bait and either **A.**
Sloth1-HA or **B.** Sloth2-HA as prey. Blots were striped and reprobed after
detection of each antigen. Arrowheads indicated expected band, asterisks
indicate unknown bands. **C.** Developmental viability assay using *tub-Gal4* to
overexpress indicated transgenes throughout development. Crosses resulting in
no viable adults are scored as lethal (black box).
839
840 **Figure 10. Model for Sloth1 and Sloth2 bicistronic translation and function**
841 **in mitochondria**
842
843 **<u>Methods</u>**
844
845 **Molecular cloning**
846
847 Plasmid DNAs were constructed and propagated using standard protocols.
848 Briefly, chemically competent TOP10 *E.coli.* (Invitrogen, C404010) were
849 transformed with plasmids containing either Ampicillin or Kanamycin resistance
850 genes and were selected on LB-Agar plates with 100µg/ml Ampicillin or 50µg/ml
851 Kanamycin. Oligo sequences are in Supplemental File 2.
852
<u>*sloth1-sloth2* expression reporters:</u> *pMT-sloth1-RLuc* was constructed by Gibson
(NEB E2611) assembly of two DNA fragments with overlapping sequence, 1)
5'UTR, *sloth1* coding sequence, and intervening sequence (*GCAAA*) were
amplified from S2R+ genomic DNA. 2) Plasmid backbone was amplified from
pRmHa-3-Renilla (Zʜᴏᴜ *et al.* 2008), which contains a *Metallothionein* promoter
and coding sequence for Renilla luciferase. *pMT-sloth1-RLuc* derivatives were
constructed by a PCR-based site directed mutagenesis (SDM) strategy.
860
861 <u>shRNA expression vector for in vivo RNAi:</u> *pValium20-sloth1-sloth2* (aka *UAS-*
862 *shRNA*, or *JAB200*) was constructed by annealing complementary oligos and
863 ligating into *pValium20* (Nɪ *et al.* 2011) digested with NheI and EcoRI. See
864 Supplemental Figure 1 for location of target site.
865
<u>sgRNA expression vectors for CRISPR/Cas9:</u> Plasmids encoding two sgRNAs
were constructed by PCR amplifying an insert and ligating into *pCFD4* (Pᴏʀᴛ *et*
al. 2014) digested with BbsI. sgRNAs constructed: *pCFD4-sloth1* (aka JAB203),
pCFD4-sloth2 (aka GP01169), *pCFD4-sloth1-sloth2* (aka JAB205, for dKO). See
Supplemental Figure 1 for location of target sites.
<u>Gal4 HDR donor plasmid:</u> *pHD-sloth1-sloth2-Gal4-SV40-loxP-dsRed-loxP* was
assembled by digesting *pHD-DsRed-attP* (Gʀᴀᴛᴢ *et al.* 2014) with EcoRI/XhoI

and Gibson assembling with four PCR amplified fragments: 1) Left homology arm
from genomic DNA from *nos-Cas9[attP2]* flies. 2) *Gal4-SV40* from *pAct-FRT-*
stop-FRT3-FRT-FRT3-Gal4 attB (BOSCH *et al.* 2015). 3) *loxP-dsRed-loxP* from
pHD-DsRed-attP. 4) Right homology arm from genomic DNA from *nos-*
Cas9[attP2] flies.
878
Custom pEntr vectors: Construction of pEntr vectors (for Gateway cloning) was
performed by Gibson assembly of PCR amplified backbone from pEntr-dTOPO
(Invitrogen C4040-10) and PCR amplified gene coding sequence (when
appropriate, with or without stop codon). List of plasmids: *pEntr_sloth1* (from
S2R+ cDNA), *pEntr_sloth2* (from S2R+ cDNA), *pEntr_hSMIM4* (from IDT
gBlock)*, pEntr_hC12orf73* (from IDT gBlock), *pEntr_sloth1-sloth2* transcript (from
S2R+ cDNA), *pEntr_sloth1-sloth2* genomic (from S2R+ genomic DNA), and
pEntr_BFP (from *mTagBFP2*). Derivatives of *pEntr_sloth1-sloth2* genomic that
lack *sloth1* or *sloth2* coding sequence, or derivatives of *pEntr_sloth1* or
pEntr_sloth2 with or without only the N-terminal signal sequence, were generated
by PCR amplifying the plasmid and reassembling the linearized plasmid (minus
the desired sequence) by Gibson.
891
892 Custom gateway expression vectors: *pMT-GW-SBP* was constructed by
893 digesting *pMK33-SBP-C* (YANG AND VERAKSA 2017) and *pMK33-GW* (Ram
894 Viswanatha) with XhoI/SpeI and ligating the GW insert into digested *pMK33-*
895 *SBP-C* using T4 ligase.
896
Gateway cloning LR reactions: Gateway cloning reactions were performed using
LR Clonase II Enzyme mix (Invitrogen 11791-020). See Supplemental File 3 for
plasmids constructed by Gateway reactions. Additional plasmids obtained were
pEntr_RFeSP (DmCD00481962), *pEntr_CG10075* (DmCD00473802) (The FlyBi
Consortium; https://flybi.hms.harvard.edu/), *pAWF* and *pAWH* (Carnegie
Science/Murphy lab), *pWalium10-roe* (PERKINS *et al.* 2015), and *pBID-G* (WANG
et al. 2012).
904
905 **Fly genetics**
906
907 Flies were maintained on standard fly food at 25˚C. Wild-type (WT) or control
908 flies refers to *yw*. The *yv; attP40* strain is used as a negative control for
909 experiments involving an shRNA or sgRNA transgene inserted into *attP40*.
910
Fly stocks were obtained from the Perrimon lab collection, Bloomington Stock
center (indicated with BL#), or generated in this study (see below). Bloomington
Stocks: *yw* (1495), *yv; P{y[+t7.7]=CaryP}attP40* (36304), *yv,P{y[+t7.7]=nos-*
phiC31\int.NLS}X; P{y[+t7.7]=CaryP}attP40 (25709), *P{y[+t7.7]=nos-*
phiC31\int.NLS}X, y[1] sc[1] v[1] sev[21]; P{y[+t7.7]=CaryP}attP2 (25710),
w[1118]; Dp(1;3)DC166, PBac{y[+mDint2] w[+mC]=DC166}VK00033 (30299),
y[1] M{w[+mC]=Act5C-Cas9.P}ZH-2A w[*] (54590), *y[1] sc[*] v[1] sev[21];*
P{y[+t7.7] v[+t1.8]=nos-Cas9.R}attP2 (78782), *w[*]; P{w[+mC]=UAS-*

2xEGFP}AH2 (6874), *w[1118]; P{w[+mC]=UAS-GFP.nls}14* (4775), *y1 w*;*
P{tubP-GAL4}LL7/TM3, Sb1 Ser1 (5138), *MN-Gal4, UAS-mitoGFP* (42737), *MN-*
Gal4, UAS-nSybGFP (9263), *UAS-tdTomato* (92759), *elav-Gal4* (8760).
Perrimon Lab stocks: *w; da-Gal4, lethal/FM7-GFP.*
923
Transgenic flies using PhiC31 integration were made by injecting attB-containing
plasmids at 200ng/ul into integrase-expressing embryos that contained an attP
landing site (attP40 or attP2). Injected adults were outcrossed to balancer
chromosome lines to isolate transgenic founder flies and eventually generate
balanced stocks. *pCFD4-sloth1[attP40]* (aka JAB203), *pCFD4-sloth2[attP40]*
(aka GP01169), *pCFD4-sloth1-sloth2[attP40]* (aka JAB205, for dKO), *pValium20-*
sloth1-sloth2[attP40] (aka *UAS-shRNA*, or *JAB200*) lines were selected with
vermillion+. pWalium10-sloth1[attP2], pWalium10-sloth2[attP2], pValium10-
sloth2[attP40], pWalium10-hSMIM4[attP2], pWalium10-hC12orf73[attP2],
pWalium10-sloth1-sloth2transcript[attP2], pBID-{sloth1-sloth2}[attP40], pBID-
{Δsloth1-sloth2}[attP40], pBID-{sloth1-Δsloth2}[attP40] were selected with
white+.
936
937 *sloth1*-KO, *sloth2-KO,* and *dKO* fly lines were made by crossing sgRNA-
938 expressing transgenic lines to *nos-Cas9[attP2]* flies, outcrossing progeny to *FM7-*
939 *GFP* balancer flies, and screening progeny founder flies for deletions by PCR
940 and Sanger sequencing.
941
942 *Gal4-KI* flies were made by injecting sgRNA plasmid (JAB205) and *pHD-sloth1-*
943 *sloth2-Gal4-SV40-loxP-dsRed-loxP*, each at 200ng/ul, into embryos expressing
944 Cas9 in the germ line (*nos-Cas9*). Injected adults were outcrossed to *FM7-GFP*
945 flies, progeny were screened for RFP+ expression, and RFP+ founder lines were
946 confirmed by PCR for a correct knock-in.
947
Knockdown crosses were performed by crossing *da-Gal4* with *pValium20-sloth1-*
sloth2[attP40]/CyO (aka *UAS-shRNA*, or *JAB200*) or *attP40/CyO* as a negative
control. Quantification of viability was performed by counting the number of
progeny with or without the CyO balancer. A Chi-square test was used to
determine if the ratio of non-balancer flies (CyO⁻) to balancer flies (CyO⁺) was
significantly altered in shRNA crosses compared to control crosses. Data was
analyzed using Excel and Prism.
955
For climbing assays, *da-Gal4/shRNA or da-Gal4/attP40* adult progeny were aged
1 week after eclosion and 10 flies were transferred into empty plastic vials
without use of CO_2. Climbing ability was quantified by tapping vials and
recording the number of flies that climb to the top of the vial within 10 seconds,
using video analysis. Climbing assays with the same 10 flies were performed
three times and averaged. Three biological replicates were performed for each
genotype. A T-Test was used to calculate statistical significance. Data was
analyzed using Excel and Prism.
964

Somatic knockout crosses were performed by crossing *Act-Cas9* to
sgRNA[attP40]/CyO or *attP40/CyO* as a negative control. *Act-Cas9/sgRNA[attP40]* female and male progeny were analyzed for phenotypes.
Quantification of viability was performed by counting the number of progeny with
or without the *CyO* balancer. A Chi-square test was used to determine if the ratio
of non-balancer flies (*CyO*⁻) to balancer flies (*CyO*⁺) was significantly altered in
somatic knockout crosses compared to control crosses. Male and female
progeny were analyzed separately because they differ in the number of copies of
the endogenous *sloth1-sloth2* loci on the X-chromosome. Data was analyzed
using Excel and Prism.

Mutant and genomic rescue crosses were performed by crossing *mutant/FM7-GFP* females to genomic rescue constructs or *attP40* as a negative control.
mutant/Y hemizygous male progeny were analyzed for phenotypes.
Quantification of viability was performed by counting the number of *mutant/Y* vs
FM7GFP male progeny. Gal4/UAS rescue crosses were performed by crossing
mutant/FM7-GFP;; da-Gal4 females to *UAS-X* lines. Additionally, *Gal4-KI/FM7-GFP* females were crossed to *UAS-X*. Rare *sloth1*-KO, *sloth2-KO, dKO,* and
Gal4-KI hemizygous adult males normally die by sticking to the fly food after they
eclose. To collect these rare mutants for further analysis (scutellar bristle images,
climbing assays), we inverted progeny vials so that mutant adults fell onto the dry
cotton plug once they eclose.

Overexpression crosses were performed by crossing *tub-Gal4/TM3* females to
UAS-X lines. At least 100 *tub-Gal4/UAS-X* progeny were analyzed for
phenotypes.

Cell fractionation and mitochondrial isolation

To isolate mitochondria from S2R+ cells, cell pellets were resuspended in 1.1ml
hypotonic buffer (10 mM NaCl, 1.5 mM MgCl2, 10 mM Tris- HCl pH 7.5),
transferred to cold glass dounce on ice, and incubated for 10min to induce cell
swelling. Cells were homogenized with 10 strokes using pestle B (tight pestle),
followed by addition of 800µl of 2.5x homogenization buffer (525mM mannitol,
175 mM sucrose, 12.5 mM Tris-HCl pH 7.5 and 2.5 mM EDTA). Homogenates at
this step are considered whole cell lysate (WCL). WCL was centrifuged at 1,300g
for 5min at 4˚C, supernatant transferred to a new tube, repeated centrifugation.
Supernatant was transferred to a new tube and centrifuged at 17,000g for 15min
at 4˚C. Supernatant was removed (cytoplasmic fraction) and 2ml 1x
Homogenization buffer (210 mM mannitol, 70 mM sucrose, 5 mM Tris-HCl pH 7.5
and 1 mM EDTA) was added to the pellet. The centrifugation was repeated and
250µl 1x Homogenization buffer was added to the pellet (mitochondrial fraction).
For SDS-PAGE comparisons of cell fractions, WCL, cytoplasmic, and
mitochondria were lysed in RIPA buffer and protein concentration normalized by
BCA assay (Thermo Fischer, 23227).

Mitochondrial isolation from 7 day old adult thoraxes and whole 3rd instar larvae
was modified from (GARCIA *et al.* 2017). Briefly, dissected adult male thoraxes or
whole 3rd instar male larvae were placed into 100µl mitochondrial isolation buffer
(250mM Sucrose, 150mM MgCl2, 10mM Tris-HCl pH 7.4) on ice. Thoraxes were
ground using a blue pestle and a motorized pestle holder. 400µl mitochondrial
isolation buffer was added to homogenized thoraxes and samples were
centrifuged at 500g at 4°C for 5min to pellet debris and tissues. Supernatant was
transferred to a new tube and the centrifugation repeated. Supernatant was
transferred to a new tube and centrifuged at 5000g at 4°C for 5min to pellet
mitochondria. The mitochondrial pellet was washed 2x by adding 1ml
mitochondrial isolation buffer and repeating centrifugation at 5000g at 4°C for
5min. For BN-PAGE experiments, 10 thoraxes or 10 whole 3rd instar larvae were
used. For SDS-PAGE, 30 thoraxes or 30 whole 3rd instar larvae were used, and
mitochondria were lysed in RIPA buffer and protein concentration normalized by
BCA assay (Thermo Fischer, 23227).
1026
1027 **Blue Native PAGE (BN-PAGE) of mitochondrial respiratory complexes**
1028
Native mitochondrial respiratory complexes were visualized by Blue Native PAGE
(BN-PAGE) gels following the manufacturer's instructions protocols (Nativepage
12% Bis Tris Protein Gels, 1.0 mm, 15 well, Thermo Fisher Scientific
BN1003BOX). Mitochondrial pellets from 10 thoraxes or 10 larvae were
resuspended in 20ul sample buffer cocktail (5µl sample buffer, 8µl 5% digitonin,
7µl H20, 2µl 5% Coomassie G-250 sample additive). 15µl sample ran on each
lane.
1036
1037 **Cell culture**
1038
Drosophila S2R+ cells (YANAGAWA *et al.* 1998), or S2R+ cells stably expressing
Cas9 and a mCherry protein trap in *Clic* (known as PT5/Cas9) (VISWANATHA *et al.*
2018), were cultured at 25°C using Schneider's media (21720-024,
ThermoFisher) with 10% FBS (A3912, Sigma) and 50 U/ml penicillin strep
(15070-063, ThermoFisher). S2R+ cells were transfected using Effectene
(301427, Qiagen) following the manufacturer's instructions.
1045
For generating stable cell lines MT-Sloth1-SBP, MT-Sloth2-SBP, and MT-
Sloth1/2, S2R+ cells were seeded in 6-well plates and transfected with *pMK33*
expression plasmids (see Supplemental File 3). *pMK33* derived plasmids contain
a Hygromycin resistance gene and a *Metallothionein* promoter to induce gene
expression. After 4 days, transfected cells were selected with 200µg/ml
Hygromycin in Schneider's medium for approximately 1 month. For induction of
gene expression, cells were cultured with 500 µM CuSO4 in Schneider's medium
for 16hrs.
1054
For generating KO cell lines, S2R+Cas9 cells were transfected with *tub-GFP*
plasmid (gift of Steve Cohen) and an sgRNA-expressing plasmid (*pCFD4-*

sloth1[attP40] (aka JAB203), *pCFD4-sloth2[attP40]* (aka GP01169), or *pCFD4-*
sloth1-sloth2[attP40] (aka JAB205, for dKO)). 48hrs after transfection, cells were
resuspended in fresh media, triturated to break up cell clumps, and pipetted into
a cell straining FACS tube (352235 Corning). Single GFP+ cells were sorted into
single wells of a 96 well plate containing 50% conditioned media using an Aria-
594 instrument at the Harvard Medical School Division of Immunology's Flow
Cytometry Facility. Once colonies were visible by eye (3-4 weeks), they were
expanded and analyzed by PCR and Sanger sequencing.
1065
For co-immunoprecipitation experiments, S2R+ cells were transfected in 100mm
petri dishes. Four days after transfection, cells were resuspended and
centrifuged at 1000g for 10min at 4˚C. Cell pellets were washed once with ice-
cold 1x PBS, re-centrifuged, and flash frozen in liquid nitrogen. Cell pellets were
subjected to mitochondrial isolation (described above) and mitochondrial pellets
were flash frozen in liquid nitrogen. Mitochondrial pellets were resuspended in
250µl mitochondrial lysis buffer (~.5-1ug/ul final protein concentration), incubated
on ice for 30min and centrifuged at 13,000g for 10min at 4˚C. The supernatant
was incubated with 20µl magnetic anti-FLAG beads (Sigma-Aldrich M8823) for
2hr at 4˚C with gentle rocking. Beads were washed 3x in mitochondrial lysis
buffer using a magnetic stand and eluted for 30min at 4˚C with 20ul 3xFLAG
peptide diluted at 1mg/ml in mitochondrial lysis buffer. Mitochondrial lysis buffer:
50 mM Tris-HCl pH 7.4, 150 mM NaCl, 10% glycerol (v/v), 20 mM MgCl2, 1%
digitonin (v/w) (Sigma D141), protease inhibitor (Pierce 87786), and 2 mM PMSF
added immediately before use.
1081
To measure mitochondrial respiration in S2R+ cells, we performed a Mito Stress
Test on a Seahorse XFe96 Analyzer (Agilent, 103015-100). 50,000 cells were
seeded into Seahorse XF96 tissue culture microplates and incubated at 25˚C
overnight. 1hr before analysis, cell culture media was replaced with serum-free
Schneider's media and drugs were loaded into the Seahorse XFe96 Sensor
Cartridge (Final concentrations: Oligomycin 1µM, Bam15 .5µM, 1µM
Antimyzin/Rotenone "R/A"). Seahorse analysis was performed at room
temperature. Mitochondrial respiration recordings were normalized to cell number
using CyQUANT (Thermo Fisher C7026) fluorescence on a plate reader. Data
analysis was performed using Seahorse Wave Desktop Software 2.6, Excel, and
Prism. N=6 wells for each condition. Significance was calculated using a T-Test.
1093
To measure *MT-sloth1-RLuc* reporter expression, S2R+ cells were transfected in
white opaque-bottom 96 well plates with *MT-sloth1-RLuc* (or derivatives) and
MT-FLuc (Firefly Luciferase) (ZHOU *et al.* 2008) as an internal control. Briefly, to
each well, 10ng of plasmid mix was added, then 10µl Enhancer mix (.8µl
Enhancer + 9.2µl EC buffer), and was incubated for 2-5min at room temperature.
20µl of Effectene mix (2.5µl Effectene + 17.5µl EC buffer) was added and
incubated for 5-10min at room temperature. 150µl of S2R+ cells (at 3.3x10^5
cells/ml) was added gently to each well and incubated at 25˚C. After 3 days
incubation, 200µM CuSO4 was added. After 24 hours incubation, media was

gently removed from the wells by pipetting and cell luminescence was measured
using the Dual-Glo assay (Promega E2920). Two luminescence normalizations
were performed. First, for each sample, Renilla luminescence was normalized to
Firefly luminescence (Rluc/Fluc). Next, Rluc/Fluc ratios for each sample were
normalized to Rluc/Fluc ratios for wild-type *MT-sloth1-RLuc* (aka fold change
Rluc/Fluc to WT). For each genotype, N=4. Significance was calculated using a
T-test. Data was analyzed using Excel and Prism.
1110
1111 **Western blotting**
1112
Protein or cell samples were denatured in 2x SDS Sample buffer (100mM Tris-
CL pH 6.8, 4% SDS, .2% bromophenol blue, 20% glycerol, .58 M β-
mercaptoethanol) by boiling for 10 min. For western blots using glycine-based
gels (Figure 7D, Figure 8C-F, Figure 9A-B, Supplemental Figure 8A,B,D),
denatured proteins and Pageruler Prestained Protein Ladder (Thermo Fisher
Scientific 26616) were loaded into 4–20% Mini-PROTEAN TGX gels (Biorad
4561096) using running buffer (25 mM Tris, 192 mM glycine, 0.1% SDS, pH 8.3).
For western blots using tricine-based gels (Figure 6B-D, Supplemental Figure 8C
) (to improve resolution of small peptides), denatured proteins and Precision Plus
Protein™ Dual Xtra Prestained Protein Standards (Biorad 1610377) were loaded
into 16.5% Mini-PROTEAN® Tris-Tricine Gels (Biorad 4563066) using
Tris/Tricine/SDS Running buffer (Biorad 1610744). Gels were ran at 100-200V in
a Mini-PROTEAN Tetra Vertical Electrophoresis Cell (Biorad 1658004). Proteins
were transferred to Immobilon-FL PVDF (Millipore IPFL00010) in transfer buffer
(25 mM Tris, 192 mM glycine) using a Trans-Blot Turbo Transfer System (Biorad
1704150) (Standard SD program). Resulting blots were incubated in TBST (1x
TBS + .1% Tween20) for 20min on an orbital shaker, blocked in 5% non-fat milk
in TBST for 1 hour at room temperature, and incubated with primary antibody
diluted in blocking solution overnight at 4˚C. Blots were washed with TBST and
incubated in secondary antibody in blocking solution for 4 hours at room
temperature. Blots were washed in TBST before detection of proteins. HRP-
conjugated secondary antibodies were visualized using ECL (34580,
ThermoFisher). Blots were imaged on a ChemiDoc MP Imaging System
(BioRad)**.** Antibody complexes were reprobed by incubating blots with stripping
buffer (Thermo Scientific 46430) following the manufacturer's instructions, re-
blocked in 5% non-fat milk in TBST, and incubated with primary antibody
overnight as described.
1140
For western blots from larval brains, 3rd instar larval brains were dissected in ice
cold PBS buffer with protease and phosphatase inhibitors. 10 brains per
genotype were homogenized in RIPA buffer and protein concentration was
measured by BCA assay (Thermo Fischer, 23227). Equal amounts of protein
samples were mixed with 1X Sample buffer (BioRad, 161-0747), boiled for 5 min,
and loaded into 4-20% Mini-PROTEAN® TGX gel (Bio-Rad). Gels were then
transferred to nitrocellulose membranes using Bio-Rad Trans-Blot SD Semi-Dry
Transfer system. Western blots using anti-Hsp60 likely recognize Hsp60A, as

opposed to Hsp60B/C/D, because only Hsp60A is expressed in the larval brain (flyrnai.org/tools/dget/web).

Commercially available or published antibodies used for western blotting: rat anti-HA (1:2000, Roche 11867423001) (Figure 9A,B), chicken anti-HA (1:1000, ET-HA100, Aves) (Figure 8E,F), mouse anti-FLAG (1:1000, Sigma F1804), mouse anti-SBP (1:1000, Santa Cruz sc-101595), mouse anti-a-Tubulin (1:20000, Sigma T5168), rabbit anti-GFP (1:5000, Invitrogen A-6455), rabbit anti-Hsp60 antibody (Abcam ab46798), mouse anti-actin (MP Biomedicals 08691002), anti-actin Rhodamine (Biorad 12004163), rabbit anti-SMIM4 (1:10,000, HPA047771), anti-UQCR-C2 (1:1000, (MURARI *et al.* 2020)), anti-SdhA (1:1000, (MURARI *et al.* 2020)), rabbit anti-C12orf73 (1:1000, HPA038883), anti-mouse HRP (1:3000, NXA931, Amersham), anti-rat HRP (1:3000, Jackson 112-035-062), anti-rabbit HRP (1:3000, Amersham NA934), anti-chicken HRP (1:1000, Sigma SAB3700199), anti-mouse 800 (only used in Figure 8E,F to detect mouse anti-FLAG) (1:5000, Invitrogen A32730). Anti-Sloth1 and Anti-Sloth2 antibodies (1:1000) were raised in rabbits (Genscript, PolyExpress Silver Package). Epitopes used: Anti-Sloth1 #1: RRLLDSWPGKKRFGC, Anti-Sloth1 #2: CEQQHLQARAANNTN, Anti-Sloth2 #1: CHSTQVDPTAKPPES, Anti-Sloth2 #2: CYKPLEDLRVYIEQE

Molecular biology

S2R+ cell genomic DNA was isolated using QuickExtract (QE09050, Lucigen). Fly genomic DNA was isolated by grinding a single fly in 50µl squishing buffer (10 mM Tris-Cl pH 8.2, 1 mM EDTA, 25 mM NaCl) with 200µg/ml Proteinase K (3115879001, Roche), incubating at 37˚C for 30 min, and 95˚C for 2 minutes. PCR was performed using Taq polymerase (TAKR001C, ClonTech) when running DNA fragments on a gel, and Phusion polymerase (M-0530, NEB) was used when DNA fragments were sequenced or used for molecular cloning. DNA fragments were run on a 1% agarose gel for imaging or purified on QIAquick columns (28115, Qiagen) for sequencing analysis. Sanger sequencing was performed at the DF/HCC DNA Resource Core facility and chromatograms were analyzed using Lasergene 13 software (DNASTAR).

For RT-qPCR analysis of *sloth1-sloth2* RNAi knockdown, *da-Gal4* was crossed with *attP40* or *UAS-shRNA* and ten 3rd instar larvae progeny of each genotype were flash frozen in liquid nitrogen. Frozen larvae were homogenized in 600µl Trizol (Invitrogen 15596026) and RNA extracted using a Direct-zol RNA Miniprep kit (Zymo Research, R2050). cDNA was generated using the iScript Reverse Transcription Supermix (BioRad 1708840). cDNA was analyzed by RT-qPCR using iQ SYBR Green Supermix (BioRad 170-8880). qPCR primer sequences are listed in Supplemental File 2. Each qPCR reaction was performed with two biological replicates, with three technical replicates each. Data was analyzed using Bio-Rad CFX Manager, Excel, and Prism. Data from *sloth1-sloth2* specific primers were normalized to primers that amplify *GAPDH* and *Rp49*. Statistical

1195   significance was calculated using a T-Test.

1196   **Bioinformatic analysis**
1197

Protein similarity between fly and human Sloth1 and Sloth2 orthologs was
determined using BLASTP (blast.ncbi.nlm.nih.gov) by defining the percent amino
acid identity between all four comparisons. Homologs in other organisms and
their gene structure were identified using a combination of BLASTP, Ensembl
(www.ensembl.org), HomoloGene (www.ncbi.nlm.nih.gov/homologene), and
DIOPT (www.flyrnai.org/diopt). Protein accession numbers: Human *SMIM4*
NP_001118239.1, Human *C12orf73* NP_001129042.1, Mouse *SMIM4*
NP_001295020.1, Mouse *C12orf73 homolog* NP_001129039.1, Zebrafish
SMIM4 NP_001289975.1, Zebrafish *C12orf73 homolog* NP_001129045.1,
Lamprey *SMIM4* XP_032827557.1, Lamprey *C12orf73 homolog*
XP_032827559.1, *D.melanogaster CG32736* NP_727152.1, *D.melanogaster*
CG42308 NP_001138171.1, *Arabidopsis AT5G57080* NP_200518.1, *Arabidopsis*
AT4G26055 NP_001119059.1, *Plasmodium PF3D7_0709800* XP_002808771.1,
Choanoflagellate (*Salpingoeca urceolata*) *m.92763* (RICHTER *et al.* 2018),
Choanoflagellate (*Salpingoeca urceolata*) *sloth2* homolog is unannotated but
present in comp15074_c0_seq2 (RICHTER *et al.* 2018). Sea squirt (*C. intestinalis*)
sloth1 and *sloth2* homologs are unannotated but present in *LOC100183920*
XM_018812254.2. Genomic sequences for *sloth1/2* ORFs in *D.melanogaster*,
Lamprey, Choanoflagellate, and Sea squirt are shown in Supplemental File 1.
1217

1218   Amino acid sequence of fly and human Sloth1∕Sloth2 were analyzed for
1219   predicted domains using the following programs: MitoFates
1220   (http://mitf.cbrc.jp/MitoFates/cgi-bin/top.cgi), DeepMito
1221   (http://busca.biocomp.unibo.it/deepmito/), TMHMM 2.0
1222   (http://www.cbs.dtu.dk/services/TMHMM/).
1223

1224   Amino acid sequences were aligned using Clustal Omega
1225   (https://www.ebi.ac.uk/Tools/msa/clustalo/) and visualized using Jalview
1226   (https://www.jalview.org/).
1227

1228   **Imaging**
1229

1230   For imaging adult scutellar bristles, adult flies were frozen overnight and
1231   dissected to remove their legs and abdomen. Dissected adults were arranged on
1232   a white surface and a focal stack was taken using a Zeiss Axio Zoom V16. Focal
1233   stacks were merged using Helicon Focus 6.2.2.
1234

For imaging larval brains, wandering 3rd instar larvae were dissected in PBS and
carcasses were fixed in 4% paraformaldehyde for 20min. Fixed carcasses were
either mounted on slides in mounting medium (see below), or permeabilized in
PBT, blocked for 1hr in 5% normal goat serum (S-1000, Vector Labs) at room
temperature, and incubated with primary antibody (anti-Elav) overnight at 4˚C,
washed with PBT, incubated with secondary antibody (anti-mouse 633) for 4hr at

room temperature, washed with PBT and PBS, and incubated in mounting media
(90% glycerol + 10% PBS) overnight at 4˚C. Larval brains were dissected from
carcasses and mounted on a glass slide under a coverslip using vectashield (H-
1000, Vector Laboratories Inc.). Images of larval brains were acquired on a Zeiss
Axio Zoom V16 or a Zeiss 780 confocal microscope. Images were processed
using Fiji software.

1247

For imaging the larval NMJ, wandering 3rd instar larvae were dissected as
previously described (BRENT *et al.* 2009). Briefly, larvae were pinned to a
Sylgard-coated (Dow 4019862) petri dish, an incision was made along their
dorsal surface, their cuticle was pinned down to flatten the body wall muscles,
and were fixed in 4% paraformaldehyde for 20min. Fixed carcasses were
permeabilized in PBT, blocked for 1hr in 5% normal goat serum (S-1000, Vector
Labs) at room temperature, and incubated with primary antibody overnight at
4˚C, washed with PBT, incubated with secondary antibody for 4hr at room
temperature, washed with PBT and PBS, and incubated in mounting media (90%
glycerol + 10% PBS) overnight at 4˚C. Whole carcasses mounted on a glass
slide under a coverslip using vectashield (H-1000, Vector Laboratories Inc.).
Images of the NMJ were acquired on a Zeiss Axio Zoom V16 or a Zeiss 780
confocal microscope. Images were taken from muscle 6/7 segment A2. Images
were processed using Fiji software. Quantification of bouton number from NMJ
stained with anti-HRP and anti-Dlg1 was performed by manual counting of
boutons in an entire NMJ for wild-type (N=8) and dKO animals (N=7). A T-test
was used to determine significance.

1265

1266   For imaging whole larvae, wandering 3rd instar larvae were washed with PBS and
1267   heat-killed for 5min on a hot slide warmer to stop movement. Larvae were
1268   imaged using a Zeiss Axio Zoom V16 fluorescence microscope.

1269

For imaging the adult brain, ~1 week old adult flies were dissected in PBS and
whole brains were fixed in 4% paraformaldehyde for 20min. Fixed brains were
permeabilized in PBT, blocked for 1hr in 5% normal goat serum (S-1000, Vector
Labs) at room temperature, incubated with anti-HRP 647 overnight at 4˚C,
washed with PBT and PBS, and incubated in mounting media (90% glycerol +
10% PBS) overnight at 4˚C. Adult brains were mounted on glass slides under a
coverslip using vectashield (H-1000, Vector Laboratories Inc.). Images of adult
brains were acquired on a Zeiss 780 confocal microscope. Images were
processed using Fiji software.

1279

For confocal microscopy of adult photoreceptors, the proboscis was removed
and the head was pre-fixed with 4% formaldehyde in PBS for 30 min. After pre-
fixation, eyes were removed from the head and fixed an additional 15 minutes.
Fixed eyes were washed with PBS 3x for 10 min each and permeabilized in 0.3%
Triton X-100 in PBS for 15 min. Permeabilized, fixed samples were blocked in 1X
PBS containing 5% normal goat serum (NGS) and 0.1% Triton X-100 for 1 h
(PBT). Samples were incubated in primary antibody diluted in PBT overnight at

4°C, washed 3x with PBT, and incubated in secondary antibodies in NGS for 1hr
at room temp the next day. Following secondary antibody incubation, samples
were washed with PBS and were mounted on microscope slides using
vectashield. Samples were imaged with LSM710 confocal with 63X objective and
processed using Fiji software.
1292
S2R+ cells transfected with Sloth1-FLAG or Sloth2-FLAG were plated into wells
of a glass-bottom 384 well plate (6007558, PerkinElmer) and allowed to adhere
for 2 hours. Cells were fixed by incubating with 4% paraformaldehyde for 30min,
washed with PBS with .1% TritonX-100 (PBT) 3x 5min each, blocked in 5%
Normal Goat Serum (NGS) in PBT for 1hr at room temperature, and incubated in
primary antibodies diluted in PBT-NGS overnight at 4˚C on a rocker. Wells were
washed in PBT, incubated with secondary antibodies and DAPI and washed in
PBS. Plates were imaged on an IN Cell Analyzer 6000 (GE) using a 20x or 60x
objective. Images were processed using Fiji software.
1302
List of antibodies and chemicals used for tissue staining: rat anti-Elav (1:50,
DSHB, 7E8A10), goat anti-HRP 647 (1:400, Jackson Immunoresearch, 123-605-
021), mouse anti-ATP5α (1:500, Abcam, ab14748), DAPI (1:1000, Thermo
Fisher, D1306), rabbit anti-FLAG (1:1000, Sigma, F7425), mouse anti-FasII
(1:25, DSHB, 1D4), mouse anti-brp (1:25, DSHB, nc82), mouse anti-Dlg1 (1:250,
DSHB, 4F3), anti-mouse 633 (1:500, A-21052, Molecular Probes), mouse
monoclonal anti-Rh1 (1:50, DSHB 4C5), Phalloidin conjugated with
Alexa 488 (1:250, Invitrogen A12379).
1311
1312 **Transmission electron microscopy (TEM) of adult photoreceptors**
1313
TEM of *Drosophila* adult retinae were performed following standard electron
microscopy procedures using a Ted Pella Bio Wave processing microwave with
vacuum attachments. Briefly, whole heads were dissected in accordance to
preserve the brain tissue. The tissue was covered in 2% paraformaldehyde, 2.5%
Glutaraldehyde, in 0.1 M Sodium Cacodylate buffer at pH 7.2. After dissection,
the heads were incubated for 48hrs in the fixative on a rotator at 4˚C. The pre-
fixed heads were washed with 3X millipore water followed by secondary fixation
with 1% aqueous osmium tetroxide, and rinsed again 3X with millipore water. To
dehydrate the samples, concentrations from 25%–100% of Ethanol were used,
followed by Propylene Oxide (PO) incubation. Dehydrated samples are infiltrated
with gradual resin:PO concentrations followed by overnight infiltration with pure
resin. The samples were embedded into flat silicone molds and cured in the oven
at 62°C for 3-5 days, depending on the atmospheric humidity. The polymerized
samples were thin-sectioned at 48-50 nm and stained with 1% uranyl acetate for
14 minutes followed by 2.5% lead citrate for two minutes before TEM
examination. Retina were viewed in a JEOL JEM 1010 transmission electron
microscope at 80kV. Images were captured using an AMT XR-16 mid-mount 16
mega-pixel digital camera in Sigma mode. Three animals per genotype per
condition were used for TEM. At least 30 photoreceptors were used for organelle

quantifications. Quantification of photoreceptor number, number of aberrant
photoreceptors, and number of mitochondria per photoreceptor, was performed
in Prism. Significance was calculated using a T-Test.
1336
1337 **Electrical recordings**
1338
Intracellular Recording from Larval NMJ
3rd instar larval NMJ recordings were performed as described previously (UGUR
et al. 2017). Briefly, free moving larvae are dissected in HL3.1 buffer without
Ca^{2+}. Recordings were performed by stimulating the segmental nerve innervating
a hemisegment A3, Muscle 6/7 through a glass capillary electrode filled with
HL3.1 with 0.75 mM Ca^{2+}. There were no differences in input resistance, time
constant τ, and resting membrane potential among different genotypes tested.
Repetitive stimulations were performed at 10Hz and were reported relative to the
first excitatory junction potential (EJP). Data were processed with Mini Analysis
Program by Synaptosoft, Clampfit, and Excel. At least 5 animals were used per
each genotype per essay. Significance was calculated using a T-Test.
1350
Electroretinograms (ERGs)
ERGs were recorded according to (JAISWAL *et al.* 2015). Briefly, flies were
immobilized on a glass slide with glue. Glass recording electrodes, filled with 100
mM NaCl, were placed on the surface of the eye to record field potential. Another
electrode placed on the humerals served as a grounding electrode. Before
recording ERGs, flies were adjusted to darkness for three minutes. Their
response to light was measured in 1sec. intervals for 30 sec. To test if the flies
can recover from repetitive stimulation, we recorded ERGs after 30 sec. and
1min constant darkness following repetitive stimulation. Data were processed
with AXON-pCLAMP8.1. At least 6 animals were used per each genotype per
essay. Significance was calculated using a T-Test.
1362
1363 **Measurement of ATP levels from larvae**
1364
Ten 3rd instar larvae were snap frozen with liquid nitrogen in a 1.5 mL centrifuge
tube. Following freezing, samples were homogenized in 100 μl of 6 M guanidine-
HCl in extraction buffer (100 mM Tris and 4 mM EDTA, pH 7.8) to inhibit
ATPases, and boiled for 3 min. The samples were centrifuged to remove cuticle.
Supernatant was serially diluted with extraction buffer and protein concentration
was measured using a BCA kit (Thermo Fischer, 23227). For each genotype,
ATP levels were measured from equal protein amounts using an Invitrogen ATP
detection kit (Invitrogen, A22066) according to their protocol. N=3 experiments,
biological triplicates per genotype per experiment. Significance was calculated
using a T-Test.
1375
1376 **Supplemental Information titles and legends**
1377

Supplemental Figure 1: Related to Figure 1. A. Comparison of gene and transcript structure of the *sloth1* and *sloth2* open reading frames. A common primer pair is used to distinguish genomic from cDNA (transcript) template by PCR. Sequence of *sloth1-2* genomic and *sloth1-2* transcript region provided. **B.** DNA gel image of PCR fragments amplified from indicated template samples. Predicted spliced transcript containing both sloth1 and sloth2 open reading frames is amplified from cDNA generated from adult flies, 3rd instar larvae, and S2R+ cells.

Supplemental Figure 2: Related to Figure 2. A. Extended gene structure of *sloth1* and *sloth2* and genetic reagents. **B.** Sequence analysis of KO, dKO, and Gal4-KI alleles. **C.** (Left) Diagram of HDR knock-in of Gal4 into the *sloth1-sloth2* locus. (Right) DNA gel confirming Gal4 knock-in by PCR primers that flank the homology arms. Expected DNA fragment size in parenthesis.

Supplemental Figure 3. Related to Figure 4. Traces of electrical recordings from 3rd instar larval NMJ in *dKO*, and *dKO+genomic rescue* animals. Graph on right is a quantification of the excitatory junction potential (EJP) for indicated genotypes. Significance was calculated with a T-Test compared to the *yw* control sample. Error bars show mean with SD. N ≥ 5 larvae per genotype.

Supplemental Figure 4. Related to Figure 5. Confocal microscopy images of 3rd instar larval NMJ at muscle 6/7 segment A2. Antibodies or fluorescent proteins (green) mark synaptic components and anti-HRP (red) marks neurons. Comparison of wild-type to dKO. Graph shows quantification of synaptic bouton number by anti-Dlg1 staining. Significance of dKO bouton number was calculated with a T-test compared to WT. Error bars show mean with SD. N ≥ 7 NMJs (each from a different animal).

Supplemental Figure 5. Related to Figure 5. A-C. Transmission electron microscopy (TEM) images of sectioned adult eye photoreceptors from indicated genetic backgrounds with accompanying quantification of photoreceptor number and aberrant photoreceptors. Scalebar is 2μm. Filled red arrows indicate dead or dying photoreceptors. Open red arrows indicate unhealthy photoreceptors. Error bars show mean with SD. **A.** Animals were 4 weeks old and raised in a 12hr light/dark cycle. **B.** Animals were 1-3 days old and raised in a 12hr light/dark cycle. **C.** Animals were 4 weeks old and raised in the dark.

Supplemental Figure 6. Related to Figure 5. Confocal microscopy of adult eye photoreceptors stained with phalloidin (green) and anti-Rh1 (red). Animals were 4 weeks old and raised in the dark. Arrows indicate photoreceptors with higher levels of Rh1.

Supplemental Figure 7. Related to Figure 6. Confocal microscopy of 3rd instar larval brain with antibody staining. Anti-Sloth1 or Anti-Sloth2 (green), mitochondria labeled with anti-ATP5alpha (red), and nuclei labeled with DAPI

1424 (blue). Wild-type (*yw*) or *sloth1/2* KO. **A.** Zoom out of entire brain showing region
1425 imaged in panels B and C. Scale bar 100µm. **B.** Results using two independent
1426 anti-Sloth1 antibodies (#1 and #2). Scale bar 20µm. **C.** Results using two
1427 independent anti-Sloth2 antibodies (#1 and #2). Scale bar 20µm**.**

1428
Supplemental Figure 8. Related to Figure 6. SDS-PAGE and western blotting
using anti-Sloth1 and anti-Sloth2 antibodies of cell and mitochondrial lysates.
Two independent (#1 and #2) anti-Sloth1 and Anti-Sloth2 antibodies were tested.
Arrowheads indicated expected band, asterisks indicate unrelated band(s).
Tricine gels were used. **A.** S2R+ whole cell lysates isolated from indicated
genotypes. Rhodamine-Actin used as loading control. **B.** S2R+ mitochondrial
lysates isolated from indicated genotypes. Anti-ATP5alpha used as loading
control. Mitochondrial control = ATP5alpha, cytoplasmic control = alpha-tubulin.
C. S2R+ fractions isolated from wild-type S2R+ cells. WCL = Whole Cell Lysate,
cyto. = cytoplasmic lysate, mito. = mitochondrial lysate. Blots were stripped and
reprobed after detection of each antigen. **D.** Mitochondrial lysates isolated from
3rd instar larvae or adult thorax mitochondrial isolation of indicated genotypes.
"da>" indicates da-Gal4 crossed with attP40 (wild-type), RNAi (UAS-shRNA-
sloth1/2), OE (UAS-sloth1/2 transcript).

1443
Supplemental Figure 9. Related to Figure 7. A. Sequence analysis of single
KO S2R+ clones for *sloth1* (clone 2F8) and *sloth2* (clone 3A7). sgRNA and PAM
site indicated by grey boxes. **B.** PCR genotyping of four independently derived
single cell dKO S2R+ clones. **C-D.** Seahorse mitochondrial stress test
quantification of **C.** ATP production and **D.** Proton leak. Significance of KO lines
was calculated with a T-test compared to S2R+. Error bars show mean with SD.
** P≤0.01, *** P≤0.001, **** P≤0.0001. N=6 for each genotype. **E.** Confocal
images of 3rd instar larval ventral nerve cord (VNC), axon bundles, and
neuromuscular junction (NMJ). *MN-Gal4 UAS-mitoGFP* (*MN>mitoGFP*) (GFP)
expresses mitochondrial-localized GFP in motor neurons. Neurons are stained
with anti-HRP (magenta).

1455
1456 **Supplemental Figure 10. Related to Figure 7. A-B.** TEM images of sectioned
1457 adult photoreceptors. **A.** Adult flies are 4 weeks old and raised on a 12hr
1458 light/dark cycle. Mitochondria are indicated with red dots. **B.** Adult flies are 3 days
1459 old and raised in a 12hr light/dark cycle.

1460
1461 **Supplemental File 1.** Genomic sequence of *sloth1-sloth2* homologs in *D.*
1462 *melanogaster*, *S. urceolata*, *P. marinus*, and *C. intestinalis*

1463
1464 **Supplemental File 2.** Oligo and dsDNA sequences

1465
1466 **Supplemental File 3.** Gateway cloning plasmid list

1467
1468 **Supplemental File 4.** Raw gel and western images

1469

References:

Alloway, P. G., L. Howard and P. J. Dolph, 2000 The formation of stable rhodopsin-arrestin complexes induces apoptosis and photoreceptor cell degeneration. Neuron 28**:** 129-138.

Anderson, D. M., K. M. Anderson, C. L. Chang, C. A. Makarewich, B. R. Nelson *et al.*, 2015 A micropeptide encoded by a putative long noncoding RNA regulates muscle performance. Cell 160**:** 595-606.

Antonicka, H., Z. Y. Lin, A. Janer, M. J. Aaltonen, W. Weraarpachai *et al.*, 2020 A High-Density Human Mitochondrial Proximity Interaction Network. Cell Metab 32**:** 479-497 e479.

Basrai, M. A., P. Hieter and J. D. Boeke, 1997 Small open reading frames: beautiful needles in the haystack. Genome Res 7**:** 768-771.

Bergendahl, L. T., L. Gerasimavicius, J. Miles, L. Macdonald, J. N. Wells *et al.*, 2019 The role of protein complexes in human genetic disease. Protein Sci 28**:** 1400-1411.

Bi, P., A. Ramirez-Martinez, H. Li, J. Cannavino, J. R. McAnally *et al.*, 2017 Control of muscle formation by the fusogenic micropeptide myomixer. Science 356**:** 323-327.

Blumenthal, T., 2004 Operons in eukaryotes. Brief Funct Genomic Proteomic 3**:** 199-211.

Bosch, J. A., N. H. Tran and I. K. Hariharan, 2015 CoinFLP: a system for efficient mosaic screening and for visualizing clonal boundaries in Drosophila. Development 142**:** 597-606.

Brand, A. H., and N. Perrimon, 1993 Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. Development 118**:** 401-415.

Brent, J. R., K. M. Werner and B. D. McCabe, 2009 Drosophila larval NMJ dissection. J Vis Exp.

Busch, J. D., M. Cipullo, I. Atanassov, A. Bratic, E. Silva Ramos *et al.*, 2019 MitoRibo-Tag Mice Provide a Tool for In Vivo Studies of Mitoribosome Composition. Cell Rep 29**:** 1728-1738 e1729.

Calvo, S. E., K. R. Clauser and V. K. Mootha, 2016 MitoCarta2.0: an updated inventory of mammalian mitochondrial proteins. Nucleic Acids Res 44**:** D1251-1257.

Casson, S. A., P. M. Chilley, J. F. Topping, I. M. Evans, M. A. Souter *et al.*, 2002 The POLARIS gene of Arabidopsis encodes a predicted peptide required for correct root growth and leaf vascular patterning. Plant Cell 14**:** 1705-1721.

Cavener, D. R., 1987 Comparison of the consensus sequence flanking translational start sites in Drosophila and vertebrates. Nucleic Acids Res 15**:** 1353-1361.

Chen, C. L., Y. Hu, N. D. Udeshi, T. Y. Lau, F. Wirtz-Peitz *et al.*, 2015 Proteomic mapping in live Drosophila tissues using an engineered ascorbate peroxidase. Proc Natl Acad Sci U S A 112**:** 12093-12098.

Chen, J., A. D. Brunner, J. Z. Cogan, J. K. Nunez, A. P. Fields *et al.*, 2020 Pervasive functional translation of noncanonical human open reading frames. Science 367**:** 1140-1146.

Chng, S. C., L. Ho, J. Tian and B. Reversade, 2013 ELABELA: a hormone essential for heart development signals via the apelin receptor. Dev Cell 27**:** 672-680.

Chugunova, A., E. Loseva, P. Mazin, A. Mitina, T. Navalayeu *et al.*, 2019 LINC00116 codes for a mitochondrial peptide linking respiration and lipid metabolism. Proc Natl Acad Sci U S A 116**:** 4940-4945.

Couso, J. P., and P. Patraquim, 2017 Classification and function of small open reading frames. Nat Rev Mol Cell Biol 18**:** 575-589.

Crosby, M. A., L. S. Gramates, G. Dos Santos, B. B. Matthews, S. E. St Pierre *et al.*, 2015 Gene Model Annotations for Drosophila melanogaster: The Rule-Benders. G3 (Bethesda) 5**:** 1737-1749.

Dennerlein, S., S. Poerschke, S. Oeljeklaus, C. Wang, R. Richter-Dennerlein *et al.*, 2021 Defining the interactome of the human mitochondrial ribosome identifies SMIM4 and TMEM223 as respiratory chain assembly factors. Elife 10.

Fricker, L. D., 2005 Neuropeptide-processing enzymes: applications for drug discovery. AAPS J 7**:** E449-455.

Fukasawa, Y., J. Tsuji, S. C. Fu, K. Tomii, P. Horton *et al.*, 2015 MitoFates: improved prediction of mitochondrial targeting sequences and their cleavage sites. Mol Cell Proteomics 14**:** 1113-1126.

Galindo, M. I., J. I. Pueyo, S. Fouix, S. A. Bishop and J. P. Couso, 2007 Peptides encoded by short ORFs control development and define a new eukaryotic gene family. PLoS Biol 5**:** e106.

Garcia, C. J., J. Khajeh, E. Coulanges, E. I. Chen and E. Owusu-Ansah, 2017 Regulation of Mitochondrial Complex I Biogenesis in Drosophila Flight Muscles. Cell Rep 20**:** 264-278.

Golpich, M., E. Amini, Z. Mohamed, R. Azman Ali, N. Mohamed Ibrahim *et al.*, 2017 Mitochondrial Dysfunction and Biogenesis in Neurodegenerative diseases: Pathogenesis and Treatment. CNS Neurosci Ther 23**:** 5-22.

Gratz, S. J., F. P. Ukken, C. D. Rubinstein, G. Thiede, L. K. Donohue *et al.*, 2014 Highly specific and efficient CRISPR/Cas9-catalyzed homology-directed repair in Drosophila. Genetics 196**:** 961-971.

Guo, X., A. Chavez, A. Tung, Y. Chan, C. Kaas *et al.*, 2018 High-throughput creation and functional profiling of DNA sequence variant libraries using CRISPR-Cas9 in yeast. Nat Biotechnol 36**:** 540-546.

Hardie, R. C., and P. Raghu, 2001 Visual transduction in Drosophila. Nature 413**:** 186-193.

Hoskins, R. A., J. M. Landolin, J. B. Brown, J. E. Sandler, H. Takahashi *et al.*, 2011 Genome-wide analysis of promoter architecture in Drosophila melanogaster. Genome Res 21**:** 182-192.

Hsu, P. Y., and P. N. Benfey, 2018 Small but Mighty: Functional Peptides Encoded by Small ORFs in Plants. Proteomics 18**:** e1700038.

Jaiswal, M., N. A. Haelterman, H. Sandoval, B. Xiong, T. Donti *et al.*, 2015 Impaired Mitochondrial Energy Production Causes Light-Induced Photoreceptor Degeneration Independent of Oxidative Stress. PLoS Biol 13**:** e1002197.

Kann, O., and R. Kovacs, 2007 Mitochondria and neuronal activity. Am J Physiol Cell Physiol 292**:** C641-657.

Karginov, T. A., D. P. H. Pastor, B. L. Semler and C. M. Gomez, 2017 Mammalian Polycistronic mRNAs and Disease. Trends Genet 33**:** 129-142.

Katsir, L., K. A. Davies, D. C. Bergmann and T. Laux, 2011 Peptide signaling in plant
development. Curr Biol 21**:** R356-364.
Kumar, S., Y. Yoshida and M. Noda, 1993 Cloning of a cDNA which encodes a novel
ubiquitin-like protein. Biochem Biophys Res Commun 195**:** 393-399.
Liu, X., K. Salokas, F. Tamene, Y. Jiu, R. G. Weldatsadik *et al.*, 2018 An AP-MS- and
BioID-compatible MAC-tag enables comprehensive mapping of protein
interactions and subcellular localizations. Nat Commun 9**:** 1188.
Magny, E. G., J. I. Pueyo, F. M. Pearl, M. A. Cespedes, J. E. Niven *et al.*, 2013 Conserved
regulation of cardiac calcium uptake by peptides encoded in small open
reading frames. Science 341**:** 1116-1120.
Makarewich, C. A., K. K. Baskin, A. Z. Munir, S. Bezprozvannaya, G. Sharma *et al.*,
2018 MOXI Is a Mitochondrial Micropeptide That Enhances Fatty Acid beta-
Oxidation. Cell Rep 23**:** 3701-3709.
Mudge, J. M., J. Ruiz-Orera, J. R. Prensner, M. A. Brunet, F. Calvet *et al.*, 2022
Standardized annotation of translated open reading frames. Nat Biotechnol
40**:** 994-999.
Murari, A., S. K. Rhooms, N. S. Goparaju, M. Villanueva and E. Owusu-Ansah, 2020 An
antibody toolbox to track complex I assembly defines AIF's mitochondrial
function. J Cell Biol 219.
Nelson, B. R., C. A. Makarewich, D. M. Anderson, B. R. Winders, C. D. Troupes *et al.*,
2016 A peptide encoded by a transcript annotated as long noncoding RNA
enhances SERCA activity in muscle. Science 351**:** 271-275.
Ni, J. Q., R. Zhou, B. Czech, L. P. Liu, L. Holderbaum *et al.*, 2011 A genome-scale shRNA
resource for transgenic RNAi in Drosophila. Nat Methods 8**:** 405-407.
Nickless, A., J. M. Bailis and Z. You, 2017 Control of gene expression through the
nonsense-mediated RNA decay pathway. Cell Biosci 7**:** 26.
Papp, B., C. Pal and L. D. Hurst, 2003 Dosage sensitivity and the evolution of gene
families in yeast. Nature 424**:** 194-197.
Pauli, A., M. L. Norris, E. Valen, G. L. Chew, J. A. Gagnon *et al.*, 2014 Toddler: an
embryonic signal that promotes cell movement via Apelin receptors. Science
343**:** 1248636.
Pearson, R. K., B. Anderson and J. E. Dixon, 1993 Molecular biology of the peptide
hormone families. Endocrinol Metab Clin North Am 22**:** 753-774.
Pellegrino, M. W., A. M. Nargund and C. M. Haynes, 2013 Signaling the mitochondrial
unfolded protein response. Biochim Biophys Acta 1833**:** 410-416.
Perkins, L. A., L. Holderbaum, R. Tao, Y. Hu, R. Sopko *et al.*, 2015 The Transgenic
RNAi Project at Harvard Medical School: Resources and Validation. Genetics
201**:** 843-852.
Plaza, S., G. Menschaert and F. Payre, 2017 In Search of Lost Small Peptides. Annu
Rev Cell Dev Biol 33**:** 391-416.
Port, F., H. M. Chen, T. Lee and S. L. Bullock, 2014 Optimized CRISPR/Cas tools for
efficient germline and somatic genome engineering in Drosophila. Proc Natl
Acad Sci U S A 111**:** E2967-2976.
Prelich, G., 2012 Gene overexpression: uses, mechanisms, and interpretation.
Genetics 190**:** 841-854.

Pueyo, J. I., E. G. Magny, C. J. Sampson, U. Amin, I. R. Evans *et al.*, 2016 Hemotin, a
Regulator of Phagocytosis Encoded by a Small ORF and Conserved across
Metazoans. PLoS Biol 14**:** e1002395.
Richter, D. J., P. Fozouni, M. B. Eisen and N. King, 2018 Gene family innovation,
conservation and loss on the animal stem lineage. Elife 7.
Saghatelian, A., and J. P. Couso, 2015 Discovery and characterization of smORF-
encoded bioactive polypeptides. Nat Chem Biol 11**:** 909-916.
Sandoval, H., C. K. Yao, K. Chen, M. Jaiswal, T. Donti *et al.*, 2014 Mitochondrial fusion
but not fission regulates larval growth and synaptic development through
steroid hormone production. Elife 3.
Sardiello, M., F. Licciulli, D. Catalano, M. Attimonelli and C. Caggese, 2003
MitoDrome: a database of Drosophila melanogaster nuclear genes encoding
proteins targeted to the mitochondrion. Nucleic Acids Res 31**:** 322-324.
Savojardo, C., N. Bruciaferri, G. Tartari, P. L. Martelli and R. Casadio, 2020 DeepMito:
accurate prediction of protein sub-mitochondrial localization using
convolutional neural networks. Bioinformatics 36**:** 56-64.
Snyder, S. H., and R. B. Innis, 1979 Peptide neurotransmitters. Annu Rev Biochem
48**:** 755-782.
Sopko, R., D. Huang, N. Preston, G. Chua, B. Papp *et al.*, 2006 Mapping pathways and
phenotypes by systematic gene overexpression. Mol Cell 21**:** 319-330.
Stapleton, M., J. Carlson, P. Brokstein, C. Yu, M. Champe *et al.*, 2002 A Drosophila full-
length cDNA resource. Genome Biol 3**:** RESEARCH0080.
Stein, C. S., P. Jadiya, X. Zhang, J. M. McLendon, G. M. Abouassaly *et al.*, 2018
Mitoregulin: A lncRNA-Encoded Microprotein that Supports Mitochondrial
Supercomplexes and Respiratory Efficiency. Cell Rep 23**:** 3710-3720 e3718.
Suzuki, K., T. Hashimoto and E. Otaka, 1990 Yeast ribosomal proteins: XI. Molecular
analysis of two genes encoding YL41, an extremely small and basic ribosomal
protein, from Saccharomyces cerevisiae. Curr Genet 17**:** 185-190.
Szklarczyk, R., M. A. Huynen and B. Snel, 2008 Complex fate of paralogs. BMC Evol
Biol 8**:** 337.
Taylor, J. S., and J. Raes, 2004 Duplication and divergence: the evolution of new
genes and old ideas. Annu Rev Genet 38**:** 615-643.
Thompson, S. R., 2012 So you want to know if your message has an IRES? Wiley
Interdiscip Rev RNA 3**:** 697-705.
Thul, P. J., L. Akesson, M. Wiking, D. Mahdessian, A. Geladaki *et al.*, 2017 A
subcellular map of the human proteome. Science 356.
Trevisan, T., D. Pendin, A. Montagna, S. Bova, A. M. Ghelli *et al.*, 2018 Manipulation of
Mitochondria Dynamics Reveals Separate Roles for Form and Function in
Mitochondria Distribution. Cell Rep 23**:** 1742-1753.
Ugur, B., H. Bao, M. Stawarski, L. R. Duraine, Z. Zuo *et al.*, 2017 The Krebs Cycle
Enzyme Isocitrate Dehydrogenase 3A Couples Mitochondrial Metabolism to
Synaptic Transmission. Cell Rep 21**:** 3794-3806.
Usui, S., L. Yu and C. A. Yu, 1990 The small molecular mass ubiquinone-binding
protein (QPc-9.5 kDa) in mitochondrial ubiquinol-cytochrome c reductase:
isolation, ubiquinone-binding domain, and immunoinhibition. Biochemistry
29**:** 4618-4626.

Van Der Kelen, K., R. Beyaert, D. Inze and L. De Veylder, 2009 Translational control
of eukaryotic gene expression. Crit Rev Biochem Mol Biol 44: 143-168.
Veitia, R. A., S. Bottani and J. A. Birchler, 2008 Cellular reactions to gene dosage
imbalance: genomic, transcriptomic and proteomic effects. Trends Genet 24:
390-397.
Verstreken, P., C. V. Ly, K. J. Venken, T. W. Koh, Y. Zhou et al., 2005 Synaptic
mitochondria are critical for mobilization of reserve pool vesicles at
Drosophila neuromuscular junctions. Neuron 47: 365-378.
Viswanatha, R., Z. Li, Y. Hu and N. Perrimon, 2018 Pooled genome-wide CRISPR
screening for basal and context-specific fitness gene essentiality in
Drosophila cells. Elife 7.
Wang, J. W., E. S. Beck and B. D. McCabe, 2012 A modular toolset for recombination
transgenesis and neurogenetic analysis of Drosophila. PLoS One 7: e42102.
Wu, C. F., and F. Wong, 1977 Frequency characteristics in the visual system of
Drosophila: genetic dissection of electroretinogram components. J Gen
Physiol 69: 705-724.
Xue, Z., M. Ren, M. Wu, J. Dai, Y. S. Rong et al., 2014 Efficient gene knock-out and
knock-in with transgenic Cas9 in Drosophila. G3 (Bethesda) 4: 925-929.
Yamamoto, S., M. Jaiswal, W. L. Charng, T. Gambin, E. Karaca et al., 2014 A drosophila
genetic resource of mutants to study mechanisms underlying human genetic
diseases. Cell 159: 200-214.
Yanagawa, S., J. S. Lee and A. Ishimoto, 1998 Identification and characterization of a
novel line of Drosophila Schneider S2 cells that respond to wingless
signaling. J Biol Chem 273: 32353-32359.
Yang, L., and A. Veraksa, 2017 Single-Step Affinity Purification of ERK Signaling
Complexes Using the Streptavidin-Binding Peptide (SBP) Tag. Methods Mol
Biol 1487: 113-126.
Yeasmin, F., T. Yada and N. Akimitsu, 2018 Micropeptides Encoded in Transcripts
Previously Identified as Long Noncoding RNAs: A New Chapter in
Transcriptomics and Proteomics. Front Genet 9: 144.
Zhang, S., B. Reljic, C. Liang, B. Kerouanton, J. C. Francisco et al., 2020 Mitochondrial
peptide BRAWNIN is essential for vertebrate respiratory complex III
assembly. Nat Commun 11: 1312.
Zhou, R., I. Hotta, A. M. Denli, P. Hong, N. Perrimon et al., 2008 Comparative analysis
of argonaute-dependent small RNA pathways in Drosophila. Mol Cell 32: 592-
599.
1688

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

A

sloth1-2 genomic
781bp

Forward / sloth1 / sloth2 / Reverse

sloth1-2 transcript
635bp

Forward / Reverse

Forward   AATCGAACAGCTGATTGCTG
Reverse   CTTAAATAGCTTTAGTCACCAATAACTTG

Key
<u>Underlined</u> – primer binding site
lowercase – intron
UPPERCASE – exon
RED – sloth1 coding sequence
BLUE – sloth 2 coding sequence

sloth1-2 genomic
    781bp

<u>AATCGAACAGCTGATTGCTG</u>CGAACCGGAACAAATGGAAATTGTATCGTGAGgcaagtggagtttcccctttacttttggcaaat
aataaataaacaaaggaacaagcctaaacattttcaattaaaccatatacagAACTAACGCACACATGTGACGGAGGCAATACAC
AAACACGGCACCTTTGAATCTCGCCTTAAAATTGGCGAAACCAACACGGAATTATATAACCGCCGGCTGAAAACACATGAGTCCG
TACAGCGGATCCGTGCGTCGTCTGCTGGACAGTTGGCCAGGAAAGAAGCGCTTCGGTGTCTACCGCTTCCTGCCGCTCTTCTTTT
TACTGGGCGCCGGCCTGGAATTCTCCATGATCAATTGGACAGTGGGCGAGACCAATTTCTgtgagactgctacgcttaaaaccctt
actttatttactaatacggaatcttttccatgcagACCGCACTTTTAAGCGCCGCCAGGCGAAGAACTACGTGGAAGAGCAGCA
GCATCTGCAGGCGCGGAGCCGCGAATAACACCAACTAAGCAAAATGCCCGCCGGAGTTTCCTGGGGGCCAGTACCTGAAATTCCTCG
GCTGTGCCCTGGCATCCATGATGGCCGGATCGCAGGCTGTTCACCTTTACTATAAGCCTCTGGAGGACTTGCGCGTCTACATCGA
ACAGGAGCAACACAGCACACAGGTGGATCCCACCGCAAAGCCACCGGAATCTGCATAACACTGTGTACTAGA<u>CAAGTTATTGGTG
ACTAAAGCTATTTAAG</u>

sloth1-2 transcript
    635bp

<u>AATCGAACAGCTGATTGCTG</u>CGAACCGGAACAAATGGAAATTGTATCGTGAGAACTAACGCACACATGTGACGGAGGCAATACAC
AAACACGGCACCTTTGAATCTCGCCTTAAAATTGGCGAAACCAACACGGAATTATATAACCGCCGGCTGAAAACACATGAGTCCG
TACAGCGGATCCGTGCGTCGTCTGCTGGACAGTTGGCCAGGAAAGAAGCGCTTCGGTGTCTACCGCTTCCTGCCGCTCTTCTTTT
TACTGGGCGCCGGCCTGGAATTCTCCATGATCAATTGGACAGTGGGCGAGACCAATTTCTACCGCACTTTTAAGCGCCGCCAGGC
GAAGAACTACGTGGAAGAGCAGCAGCATCTGCAGGCGCGAGCCGCGAATAACACCAACTAAGCAAAATGCCCGCCGGAGTTTCCT
GGGGCCAGTACCTGAAATTCCTCGGCTGTGCCCTGGCATCCATGATGGCCGGATCGCAGGCTGTTCACCTTTACTATAAGCCTCT
GGAGGACTTGCGCGTCTACATCGAACAGGAGCAACACAGCACACAGGTGGATCCCACCGCAAAGCCACCGGAATCTGCATAACAC
TGTGTACTAGA<u>CAAGTTATTGGTGACTAAAGCTATTTAAG</u>

B

bp	ladder	H20	adult flies gDNA	adult flies cDNA	larvae cDNA	S2R+ cDNA	*UAS-sloth1/2* plasmid

1650
1000
850 — ◄ 781bp genomic
650 — ◄ 635bp transcript
500
400
300

Figure 1-figure supplement 1

Figure 4-figure supplement 1

Figure 5-figure supplement 1

Figure 5-figure supplement 2

4weeks old (raised in 24hr dark)

Figure 5-figure supplement 3

A

DAPI anti-ATP5alpha

100μm

B

| anti-sloth1 #1 | | anti-sloth1 #2 | |
| *yw* | *sloth1/2* | *yw* | *sloth1/2* |

ATP5α Sloth1or2 DAPI

20μm

anti-Sloth1or2

anti-ATP5alpha

C

| anti-sloth2 #1 | | anti-sloth2 #2 | |
| *yw* | *sloth1/2* | *yw* | *sloth1/2* |

ATP5α Sloth1or2 DAPI

20μm

anti-Sloth1or2

anti-ATP5alpha

Figure 6-figure supplement 1

Figure 6-figure supplement 2

Figure 7-figure supplement 1

Figure 7-figure supplement 2