Arctic charr phenotypic responses to abrupt temperature change: an

insight into how cold water fish could respond to extreme climatic events.

Oliver E. Hooker[12], Colin E. Adams[1], Louise Chavarie[1*]

[1] Scottish Centre for Ecology and the Natural Environment, IBAHCM, Rowardennan, Loch

Lomond, Glasgow G63 0AW, UK

[2] PR statistics, 3/1 128 Brunswick Street, Glasgow, G1 1TF

Running title: An insight into how cold water fish could respond to extreme climatic

events.

Key words: mean phenotype, phenotypic variance, phenotypic variability, evolutionary

ecology, morphology, temperature

*Corresponding author:

Louise.Chavarie@glasgow.ac.uk

ABSTRACT

Phenotypic plasticity, the ability of an organism to express multiple phenotypes in response to the prevailing environmental conditions without genetic change, may occur as a response to anthropogenic environmental change. Arguably, the most significant future anthropogenic environment change is contemporary climate change. Given that increasing climate variability is predicted to pose a greater risk than directional climate change, we tested the effect of a water temperature differential of 4 ºC on the Arctic charr phenotypic response within a generation. We demonstrate that Arctic charr phenotype can respond rapidly and markedly to an environmental cue. The plastic response to different temperature regimes comprised a shift in the mean phenotype coupled with a reduction in the between-individual phenotypic variation in the expressed head shape. The magnitude of shape difference was cumulative over time but the rate of divergence diminished as fish became larger. Individuals raised in the elevated temperature treatment expressed a phenotype analogous to a benthivorous ecotype of this species rather than that of the parental pelagic feeding form. The response of cold-water freshwater species to temperature change is likely to be an interaction between the capacity of the organism for phenotypic plasticity, the speed of mean change in the environment (e.g., temperature), and the degree of short interval variation in the environment.

INTRODUCTION

In nature, the expression of intraspecific, discrete, alternative phenotypes can result from modulation by the environment through plasticity (Adams, Woltering & Alexander, 2003) or from the emergence of evolutionary groups with diverging gene pools (Wu & Ting, 2004), or a combination of both (Nosil, 2012). Phenotypic plasticity is the ability of a single genotype to express multiple alternative phenotypes in response to different environmental conditions (Pigliucci & Preston, 2004). Plasticity itself is a trait, present across a broad range of taxa (Lubchenco & Cubit, 1980; Corno & Jürgens, 2006; Berg & Ellers, 2010). It can be instantaneous, anticipatory or delayed, permanent or reversible, adaptive or non-adaptive, beneficial or harmful, passive, discrete, continuous, and generational (Whitman & Agrawal, 2009). It is known to facilitate the expression of novel phenotypic traits (Skúlason *et al.*, 2019) upon which selection may then act (Ghalambor *et al.*, 2007). Thus, phenotypic plasticity has an important role in the evolutionary processes of organisms that exhibit plasticity because variation fuels evolution (Schoener, 2011).

Identifying pathways along which phenotypic plasticity can arise is still a major topic for inquiry in evolutionary ecology, since the patterns and processes that underlie variation are multifaceted and highly variable. Despite the complex nature of phenotypic plasticity, there are some abiotic factors that are known to be ubiquitous in driving variation in ecosystems, one of which is temperature (Mcphee, Noakes & Allendorf, 2012; Noble, Stenhouse & Schwanz, 2018). Many plants and animals have the capacity for a plastic response to temperature, as it is unlikely that rapid environmental change would induce an equally rapid genetic response (Orizaola & Laurila, 2009; O'Dea *et al.*, 2016; Merilä & Hendry, 2014). A rapid change in temperature could induce greater levels of phenotypic variation among individuals within a population by exposing previously hidden (cryptic) genetic variation or by inducing new epigenetic changes (O'Dea *et al.*, 2016). Accordingly, the response of an organism to

70   temperature can include variation in the mean expressed phenotype but it can also include

71   change in the variance of phenotypes expressed in a population (O'Dea *et al.*, 2019).

72   As ectotherms, fish are particularly sensitive to change in temperature (Neuheimer *et*

al., 2011). A phenotypic trait in fish that is strongly linked with temperature is growth (van der

Have & de Jong, 1996; Kingsolver, Izem & Ragland, 2004). However, the subsequent effect

of temperature and induced growth rate heterogeneity (e.g., early developmental and juvenile

growth) on the modulation of morphology is not a mechanism fully understood, despite being

observed to affect morphology modulation in a number of species (Olsson, Svanbäck & Eklöv,

2006; Jacobson, Grant & Peres-Neto, 2015; Heino, 2014). A few studies have looked at the

potential for temperature change to result in heterochrony in the developmental and ontogenetic

processes that modulate the expression of functional traits (Parmesan, 2006; Charmantier *et*

al., 2008), whilst others have attempted to predict the morphological outcome of temperature

induced plasticity (Mcphee, Noakes & Allendorf, 2012). However, the ecological

consequences that result from temperature induced phenotypic change has received little

attention ectotherms at higher latitudes (Ramler *et al.*, 2014; Burggren, 2018). As the

expression of some discrete functional phenotypes in some species is plastic, and by definition

modulated by variation in the natural environment, it is presumed that they will also respond

to anthropogenic modifications of that environment. The most obvious and far reaching

anthropogenic modification of the contemporary environment is that of climate change. The

northern regions are warming at a rate twice the rest of the planet and given that a rise in water

temperature is predicted to occur, cold-water freshwater species will be foremost affected by

climate change (Solomon, 2007; Poesch *et al.*, 2016; Heino, Virkkala & Toivonen, 2009).

92   The Arctic charr (*Salvelinus alpinus*) has a circumpolar distribution, is the most

93   northern species of freshwater fish, and displays highly variable phenotypes (Klemetsen,

2013). The species is ideal to explore how phenotypic diversity is realized within and among generations as it expresses a high degree of phenotypic plasticity and rapid intra-specific divergence, and as a result of its range and habitat use, it experiences diverse selective environments, which facilitate tracking the link between ecological processes and evolutionary outcomes (Adams, Woltering & Alexander, 2003; Garduño-Paz & Adams, 2010; Elmer, 2016; Chavarie et al., 2010). Given that increasing climate variability is predicted to pose a greater risk than directional climate change (Vasseur et al., 2014), we tested the effect of an elevated water temperature of 4ºC on the short-term, plastic, phenotypic expression in an Arctic charr population that displays continuous variation. The induced temperature variation in this study is comparable in magnitude of year on year of extreme temperature fluctuations predicted by climate change models that Arctic charr populations are likely to experience (an interannual 4 ºC temperature difference is within the predicted range for the near future in northern areas; Woelders et al., 2018; Alley et al., 2003; Heino, Virkkala & Toivonen, 2009). We predict that Arctic charr individuals reared in warmer water will express greater phenotypic variation than fish at a control temperature (see Figure 1C) but that they will not display a significant shift in the mean phenotype expressed due to the short-term exposure to elevated temperatures (Figure 1B and D) (O'Dea et al., 2019). Overall, we aimed to: 1) quantify the among-individual morphological variation of the head of a unimodal Arctic charr population exposed to different temperature regimes during early development, 2) determine the allometry trajectories that result in any differences in expressed phenotype as evidence of differing developmental pathways in groups exposed to different temperature and 3) examine if at different temperature treatments, head shape among individuals is more analogous to the expressed phenotypes in plankton feeding (the parental phenotype) or macro-benthos feeding ecotypes in the wild.

MATERIALS and METHODS

Fish collection and rearing

Eggs from nine separate female Arctic charr were each fertilised by a single male to

create nine full sibling crosses, from a morphologically unimodal, plankton feeding population

that inhabits Loch Clair, Scotland. Fish were caught during spawning time from the in-flowing

River Coulin (57°32.648'N, 5°19.125'W). Fertilised eggs were water hardened and transferred

to incubation facilities at the Scottish Centre for Ecology and the Natural Environment, Loch

Lomond. Eggs were acclimatised over a two-hour period to the new water supply and placed

in mesh baskets suspended in a holding tank in a constant temperature room maintaining a

water temperature of 4 $^{\circ}$C ($\pm$ 0.5 $^{\circ}$C). Fertilisation success was greater than 95% for all family

groups.

Experimental procedure

To control for differences in development due to different temperatures, the number of

degree-days (dd) were used as a measure of developmental rate. Degree-days are the

cumulative count of the water temperature for a known period of time in days. Embryos reached

the eyed stage after 212dd at which point they were raised to a temperature of 6 $^{\circ}$C ($\pm$ 0.5 $^{\circ}$C)

for a further 89dd, before being separated into two temperature treatment groups (n=480 per

group), an ambient and an elevated temperature treatment. Developing embryos were held in

equal numbers in eight replicate tanks per treatment group (n=60 per tank). Water temperatures

were then either lowered by 2 $^{\circ}$C to 4 $^{\circ}$C or raised by 2 $^{\circ}$C to 8 $^{\circ}$C ($\pm$ 0.5 $^{\circ}$C) respectively.

Embryos exposed to the ambient temperature (4 $^{\circ}$C) began hatching after 367dd, the hatching

period lasted 73dd (total developmental time to 100% hatch 440dd). Those exposed to the

elevated temperature (8 $^{\circ}$C) began hatching after 388dd, the hatching period lasted 50dd (total

incubation time to 100% hatch 438dd). There was no significant difference in hatching rate

between temperatures or replicates. When hatching was complete, temperatures were raised by

3 $^{\circ}$C; the ambient temperature treatment to 7 $^{\circ}$C and the elevated temperature treatment to 11

$^{\circ}$C ($\pm$ 0.5 $^{\circ}$C). Fish became partially dependant on exogenous food at 505dd for the ambient

temperature treatment and 504dd for the elevated temperature treatment. When the yolk sack was fully exhausted, 689dd for the ambient temperature treatment and 686dd for the elevated temperature treatment, temperatures were raised a further 2 $^{\circ}$C for both treatments to 9 $^{\circ}$C (ambient temperature treatment) and 13 $^{\circ}$C (elevated temperature treatment). Both temperature treatments remained within $\pm$ 0.5 $^{\circ}$C. Fish were fed four times a day to satiation at three hour intervals ($\pm$ 0.5 hours) using a standard commercial 3mm hatchery sinking pellet.

Data collection

Adult Arctic charr used as brood stock were killed using a Schedule 1 method (UK Home Office Licence Number PPL 70/8794), photographed in a lateral position on the left side before spawning. Lateral view photographs of juveniles killed using a Schedule 1 method and under licence were taken at 700dd (N=130), 1000dd (N=80) and 1400dd (N=60) using a Cannon EOS 350D digital camera, for geometric morphometric analysis. Nine consistently identifiable landmarks on the head (Figure A1) were digitised in two dimensions on each fish image using TPSdig2 (Rohlf, 2006a) and TPSutil (Rohlf, 2006b).

Data analysis

Prior to geometric morphometric analysis, landmark data were subject to a Procrustes superimposition using MorphoJ (Klingenberg, 2011) to remove variation in the data created by size, position and orientation (Mitteroecker, 2009). The mean shape configuration was then computed and the variation around this mean calculated (Dryden & Mardia, 1998).

Following this, a single, pooled within-group regression of Procrustes co-ordinates on log centroid size was conducted in MorphoJ for samples collected at 700dd, 1000dd and 1400dd's. The residuals from this regression provide a measure of shape free from allometric scaling (Klingenberg & McIntyre, 1998) associated with early ontogeny. The residuals from this regression were subsequently used for all further morphometric analysis.

A single Discriminant Function Analysis (1000 permutations) was conducted in

MorphoJ to compare geometric morphometric data from three developmental stages (700dd,

1000dd and 1400dd) to test if the magnitude of shape difference between groups changed over

time. Procrustes distance and Mahalanobis distance were used as pairwise measures of the

magnitude of shape difference. Within each temperature treatment (ambient and warm), the

polynomial relationship between discriminant function scores and size (represented by centroid

size) were used to illustrate allometric trajectories. To decide whether allometric trajectories

between temperature treatments were parallel, convergent, divergent, or common (see Figure

A2), the slopes of the regressions were examined

Using the scores generated in the Discriminant Function Analysis as a measure of

shape, a generalised linear mixed effect model, fitted by maximum likelihood (using the

software R 3.1 for Windows (Team, 2014) and the package lme4) was used to describe the

effect of temperature, exposure time (the number of dd at each developmental stage), and fish

size (measured as centroid size) with replicate (tank) as a random effect, on the expression of

shape. Models were simplified by removing the highest order, least significant terms,

sequentially (Crawley, 2012). Terms that were removed from the model without significantly

increasing model deviance using likelihood ratio tests (χ^2) were discarded. *Post hoc* pairwise

comparisons using T-test assessed if centroid size (related to body size) differed between

ambient and elevated treatments at each sampling time (700, 1000, and 1400dd).

A Canonical Variate Analysis (1000 permutations) executed in MorphoJ was used to

establish which group had been affected by the temperature treatment by comparing the head

shape of individuals raised in ambient or elevated temperature conditions in the laboratory to

their parents.

A variance ratio test was conducted using the software R 3.1 for Windows on all raw

Procrustes coordinates and used to compare within-group phenotypic variation between

groups.

To establish which temperature regime had stimulated expression of head shape closest

to a typical plankton and macro-benthos feeding specialists from the wild, head shape of

juveniles at 1400dd was compared against the head shape of two known ecologically divergent

sympatric polymorphic groups of Arctic charr that show morphological and dietary

specialisations on either plankton or macro-benthos feeding resources, one from Loch Rannoch

(Perthshire, Scotland) (Adams *et al.*, 1998) and one from Loch Dughaill (Strathcarron,

Scotland) (Hooker *et al.*, 2016a) using a Canonical Variate Analysis (1000 permutations) in

MorphoJ. The same nine consistently identifiable landmarks used for analysing laboratory

raised individuals were digitised in two dimensions on existing images of fish from Loch

Rannoch and Loch Dughaill.

RESULTS

The Discriminant Function Analysis (DFA) showed that head shape differed across all

sampling periods between ambient and elevated temperature exposed fish (Procrustes distance

= 0.028; P<0.01). The magnitude of shape differences however increased over time (at 700dd,

Procrustes distance = 0.036; P<0.01; at 1000dd, Procrustes = 0.040; P<0.01; at 1400dd,

Procrustes distance = 0.061; P<0.01; Figure 2).

Discriminant Function Analysis of shape only, correctly assigned 73.8% of ambient

temperature fish and 82.8% of elevated temperature fish (from all sampling periods combined)

to the correct temperature exposure. In general, fish exposed to elevated temperature had a

more rounded head and sub-terminal mouth than ambient temperature fish (Figure A3).

Individuals in both temperature treatments displayed allometric trajectories with polynomial

relationships of head shape on size differing from 0 (P<0.05); centroid size explained 55% of

variation in Discriminant Function Score of head shape in the ambient temperature treatment

group (R^2 =0.55) and 57% in the elevated temperature treatment group (R^2=0.57). Regression

slopes indicated that allometric trajectories were divergent (Figure 2 and Figure A2).

219   Using discriminant function scores as a measure of shape in a mixed model,

temperature had a highly significant effect on head shape (likelihood ratio χ^2 = 39.81, p<0.01)

(Table 1). There was a significant interaction between Exposure time and Centroid Size

(likelihood ratio χ^2 = 7.23, p = 0.01) (Table 1). The interaction between exposure (number of

degree-days) and fish size (centroid size) was negative. Thus, the difference in morphology

between fish raised on each temperature regime was greater in larger fish, but the rate of

divergence decreased as fish became larger; indicating that speed of divergence caused by

different temperature regimes is greater during early ontogenetic stages. Pairwise comparisons

of centroid size between temperature treatments at each sampling time, showed a difference in

size at 700dd (t = 1.98, p < 0.01) and at 1400dd (t = 2.0, p < 0.01), but no size differences were

found at 1000dd (P < 0.05). At 700dd and 1400dd, individuals raised at ambient temperature

were larger than individuals exposed to elevated temperature.

231   The variance ratio test found the phenotypic variation within treatment groups was

232   higher in ambient temperature fish compared with elevated temperature fish (p = <0.01). For

233   ambient temperature, variance in Procrustes scores (SS distance/(n-1)) was 0.0046 at 700dd,

234   0.0032 at 1000dd, and 0.0018 at 1400dd. For elevated temperature, variance of Procrustes

235   scores (SS distance/(n-1)) was 0.0038 at 700dd, 0.0039 at 1000dd, and 0.0016 at 1400dd.

236   Canonical Variate Analysis (CVA) found that the head shape of fish exposed to the

ambient temperature treatment was not significantly different to that of their parents

(Procrustes distance = 0.014; *P*=0.38), however elevated temperature fish were significantly

different (Figure A4; Procrustes distance = 0.044; *P*=<0.001).

Using a CVA to compare experimental (Coulin) charr from both temperature groups to

benthic and pelagic foraging specialists from Loch Rannoch and Loch Dughaill, we found that

ambient temperature exposed fish expressed a phenotype closer to the pelagic ecotype and fish

exposed to elevated temperature displayed a phenotype closer to benthic ecotype for both

comparisons (Figure 3 and Table 2). CV1 mostly captures shape differences between fish of

different origin, experimental charr and Loch Rannoch charr (Figure 3A) and experimental and

Loch Dughaill charr (Figure 3B). CV2 however captures within group shape differences

between fish raised at ambient or elevated temperatures, benthic and pelagic ecotypes from

Loch Rannoch (Figure 3A) and benthic and pelagic ecotypes Loch Dughaill (Figure 3B).

DISCUSSION

Variation in climate can occur at different temporal scales (e.g. diurnally, intra-

seasonally or inter-annually), and extreme temperature anomalies are now occurring more

regularly and are predicted to increase in frequency in the future (Scheepens, Deng & Bossdorf,

2018). Accordingly, in this study we demonstrate clear differences of an environmentally

contingent expressed phenotype in an ectotherm subjected to a severe climate event. Our results

demonstrate a strong and immediate influence of temperature on the direction, magnitude, and

variation in the expression of phenotypic plasticity in a cold-water specialist species where the

population in the wild displays continuous variation (i.e., unimodal phenotype; Klemetsen,

2010; Klemetsen, 2013). Elsewhere studies on plasticity during development have shown

responses to alternative environmental inputs, resulting in dramatic morphological differences

among individuals (Morris, 2014; West-Eberhard, 2003). Here, offspring of wild Arctic charr

raised at elevated temperatures expressed a mean phenotype different to their siblings raised at

an ambient temperature but also to that of their parents. The magnitude of this morphological response mimicking the level of divergence observed in wild Arctic charr populations that express sympatric foraging specialist ecotypes showing a polymodal phenotype distribution (Skulason & Smith, 1995). When compared to polymorphic populations that show distinct and stable foraging specialisms, individuals raised in the elevated temperature treatment expressed a phenotype analogous to a benthivorous ecotype rather than that of their parental pelagic form (Adams *et al.*, 1998; Hooker *et al.*, 2016b). The head of fish exposed to elevated temperature was shorter, more robust, with a rounder snout; features that are typical of littoral macro-benthos feeding specialists (Figure A3) in several post-glacial fishes (Robinson & Parsons, 2002; Bolnick & Lau, 2008). In contrast, ambient temperature exposed fish expressed an elongated and pointed head phenotype, which is more typical of fish showing pelagic planktivory in the wild (Schluter, 2000; Taylor, 1999).

Exposure to different environmental conditions is known to have asymmetrical impacts on the stability of developmental pathways during early life-stages (Robinson & Wardrop, 2002; Svanbäck *et al.*, 2009; Lazić *et al.*, 2013). Considering that temperature is one of the most widely reported environmental factors driving plastic responses in fish (Pittman *et al.*, 2013), Arctic charr individuals subjected to higher temperature developed at a faster rate (Campbell, 2019). Developmental stability is known to limit the phenotypic expression from a single genotype, thus it can modulate the magnitude and structure of phenotypic variation among individuals and cohorts (Lazić *et al.*, 2015; Willmore *et al.*, 2006). In some cases, the expressed phenotype of an organism can be significantly altered by a single short-term environmental perturbation of the timing or rate during development; such events are thought to be a route to rapid phenotypic novelty, the raw material for selection (Parsons *et al.*, 2011). Several mechanisms, such as heterochrony, are known to drive the plastic expression of phenotypes through the effects of temperature exposure on developmental pathways (Georga

& Koumoundouros, 2010; Angilletta, Steury & Sears, 2004; Ramler *et al.*, 2014).

Heterochrony can disrupt the onset and termination of developmental processes and the rate at

which these processes occur. Developmental pace in structurally important features, such as

bone metabolism, are fundamentally linked to the form of morphological plasticity examined

in this experiment (Campinho, Moutou & Power, 2004; Sfakianakis *et al.*, 2004). Therefore,

temperature effects mediated through heterochrony has the potential to result in skeletal

development of different craniofacial features in Arctic charr (Parsons & Albertson, 2009;

Campbell, 2019).

Early development represents a critical life-stage for the emergence of phenotypic

variation. During ontogeny, the environment is likely to have a greater influence on a range of

traits and the magnitude of the variation of expression in a population (Gillooly *et al.*, 2001;

Ackerly & Ward, 2016; Imsland *et al.*, 2005; Parsons, Skúlason & Ferguson, 2010). The

thermal history experienced by an individual during early development is known to be a key

factor defining adult phenotype (Georga & Koumoundouros, 2010; Angilletta, Steury & Sears,

2004; Ramler *et al.*, 2014). It is particularly during early development that environmental

signals can influence developmental processes through epigenetic mechanisms, influencing the

expressed phenotype (Duclos, Hendrikse & Jamniczky, 2019). This observed effect on the

phenotype in this study was cumulative, with divergence between treatment groups increasing

over time. However, the rate of divergence decreased with time, suggesting that the effect of

temperature on the expression of phenotypic differences occurred predominant at early life

stages. This indicates that the timing of exposure to different environmental conditions by

individuals is important in the initiation of a plastic response and its magnitude.

Temperature exposure in early life is known to result in epigenetic effects on traits such

as growth, with consequences that can range from short-term to transgenerational (Burton &

Metcalfe, 2014). Our results suggest that temperature-induced variation in developmental

pathways results in divergent plastic allometric trajectories in the early development of Arctic charr. Allometry, can been seen as a canalising process and an interacting agent that can steer morphological variation (Franklin *et al.*, 2018; Simonsen *et al.*, 2017). However, the opposite is also true, whereas allometric trajectories can deviate from the "usual" shape-size relationship (e.g., whereas size increases, shape changes under a fixed allometric relationship), deviation from this fixed allometry relationship could serve as additional measure of development instability (Lazić *et al.*, 2015). Consequently, growth trajectories manifested as allometric scaling provide the potential to express different phenotypes, which increase the potential for novelty (Frankino *et al.*, 2005). Altogether, both heterochrony and allometry are involved in determining the degree of phenotypic variation generated during growth and morphological development. In this study, the expression of phenotypic variation observed appears to be influenced by temperature with concomitant developmental effects (Lazić *et al.*, 2015; Westneat, Wright & Dingemanse, 2015).

The potential of Arctic charr to display phenotypic variation is considerable, whether expressed variation is continuous (e.g., unimodal) or discontinuous (e.g., polymodal) (Klemetsen, 2013). Populations that express high level of variability in natural conditions show higher means and greater variance in the plastic expression of phenotype in response to different temperatures (O'Dea *et al.*, 2019; Reed *et al.*, 2010). Although our study also showed a significant plastic response resulting in a shift in the mean phenotype expressed, there was also a decrease in variation of the phenotypes expressed among individuals raised at higher temperature treatment, which contrasts with previous work. Increased phenotypic variance in response to elevated temperatures has been suggested to be the reflection of a reduction in genotype precision (e.g., canalization and developmental stability), this lack of precision resulting in a spreading of reaction norms (O'Dea *et al.*, 2019; Campbell, 2019). Our opposing result could be the result of a non-linear reaction norm of head shape on size. In this scenario

a specific range of temperature exposures (either colder or warmer) has a different magnitude

effect on phenotypic mean and variation (see description of how this may occur in Figure 1 in

Ramler *et al.*, 2014). Another, not mutually exclusive possibility, is that despite both treatments

being sampled after the same number of degree-days (a proxy for growth opportunity;

Neuheimer & Taggart, 2007) had elapsed, individual responses in the two temperature groups

was partly the result of a size effect. For an unknown reason, in our experiment, growth

opportunity and the actual realized body growth were uncoupled. Contrary to the predictions

of Neuheimer and Taggart (2007), the effect of time and temperature are thus not necessary

equal in the promotion of body size and anatomical structures (e.g., heterochrony and

allometry). Consequently, if variance is cumulative with the increase of size (Gillooly *et al.*,

2002), this could partly explain the discrepancies between our predictions and findings because

ambient temperature treatment individuals were generally larger by the end of the experiment.

Broader implications

Environmentally mediated variation in phenotypic expression is a fundamental attribute

of many organisms, which can result in both short and long-term ecological evolutionary

consequences for species that display plasticity. The disjunctive nature of the freshwater

environment that lacustrine Arctic charr occupy means that, in the absence of migratory routes

for range adjustment, this species will particularly need to cope to abrupt changing novel

conditions (O'Dea *et al.*, 2016; Collins *et al.*, 2013). High levels of plasticity is one way in

which they might do this. Yet, the role of phenotypic plasticity as a potential mitigator of

climate change is contentious, with both positive and negative effects possible. We demonstrate

that the expressed phenotypic mean in Arctic charr can respond to an environmental cue rapidly

and with a magnitude that is highly likely to have functional consequences (i.e. temperature-

mediated shift from a pelagic-like towards a benthic-like phenotype). In the context that even

subtle differences in individual's morphology can influence feeding efficiency and

specialization (Garduño-Paz & Adams, 2010), the potential of a fitness-buffering effect of

plasticity can easily fluctuate, depending if environmental cues and selective filters become

decoupled or not (Reed *et al.*, 2010; O'Dea *et al.*, 2016). Consequently, important questions

arise when extrapolating our results into the wild. Would such plastic phenotypic change result

in an "overshoot" of the optimum phenotype (e.g., would such processes produce phenotype-

environment mismatches) or would it buffer against rapid fluctuations in the environment?

While a plasticity response occurs within a single cohort, another important question at

hand is what would be the consequences to subsequent generations (e.g., transgenerational

plasticity effects)? The high magnitude of the short temporal environmental oscillations, for

instance extreme environmental events, can induce a greater set of phenotypic responses than

a small or slower change (e.g., our tested 4 ℃ variation scenario; Donelson *et al.*, 2018). More

gradual or stepwise change across generations have the potential to produce different

phenotypic results compared to a single large change within a cohort (e.g., reproductive

capacity of individuals subjected to a 3℃ elevated temperature within a generation compared

with a 1.5℃ elevation over two generations; Donelson *et al.*, 2016). In reality the probability

of occurrence of alternative effects of phenotypic plasticity (positive or negative) on responses

to temperature change is likely to be an interaction between the capacity of the organism for

phenotypic plasticity (both within-generation and transgenerational plasticity), the speed of

mean change in the environment (e.g., temperature), and the degree of short interval variation

in the environment (Reed *et al.*, 2010; Donelson *et al.*, 2018).

The last component emerging from this experiment with evolutionary consequences for

a population in the face of changing selection regimes by affecting population dynamics is the

phenotypic variance (Chevin, Lande & Mace, 2010; Reed *et al.*, 2010). In this study, we

showed a decrease in variation in expressed phenotype at elevated temperature. The decrease

in the variation in the phenotypes expressed among individuals raised in the elevated

temperature treatment could reduce the range of phenotypes for selection to act upon if this

effect occurs in the wild. Accordingly, phenotypic variance has the potential to be a very

important component providing a population with scope for recovery from rapid (possibly

transient) environmental change. Consequently, a reduction or an increase of expressed

phenotypic variance under climate change scenarios has important consequences at the

population level should not be underestimated.

CONCLUSION

The ecological and evolutionary consequence of rising temperature on Arctic charr will

depend on many factors (e.g., population specific lethal limits or growth optima, interspecific

dynamics, and changing disease risk; Crozier & Hutchings, 2014). Here, we demonstrate the

capacity of Arctic charr for a rapid plastic response to the effect of a severe temperature

fluctuation in effect over periods of less than a single year. The parental phenotype mean was

shifted within a generation, but also individuals showed a marked decline in in the expression

of intra-populaton phenotypic variation. Such variation has been recently shown to have multi-

generational effects on survival in another ectotherm the minnow *Phoxinus phoxinus* (Raffard

et al., 2019). Temperature varies in a stochastic manner resulting in year-to year variation

around a coarse-scale temporal trend. Inter-annual temperature variation can modify a cohort's

phenotype in a population, consolidating the role of plasticity in species resilience to rapid

environmental change, as genetic changes are not expected to be capable of responding over

such short time scales (Merilä & Hendry, 2014).

Phenotypic plasticity is thus likely to be particularly important for population

persistence in new and changing environments and where the pace of change is relatively rapid

(Morris, 2014). Contemporary temperature-driven divergence in Nordic freshwater fishes have

been reported (Kavanagh *et al.*, 2010). Whether this flexible response will be common within

and among cold-water species is unknown. Nonetheless, the fact remains that the majority of

landlocked Arctic charr populations are currently planktivorous (Maitland & Adams, 2018).

By confirming the potential of a rapid and marked change in morphology with higher

temperature, we provide evidence that a substantial portion of Arctic charr populations

worldwide are likely to be subjected to phenotype expression changes toward a benthic-like

ecotype under climate change scenarios. A similar effect may also occur in other salmonids

and other cold-water species.

ACKNOWLEDGMENTS

The authors would like to acknowledge Travis van Leeuwen, Luc Bussierre, Alex Lyle,

Jennifer Dodd, Martin Hughes, Peter Cunningham (Wester Ross Fisheries Trust) and The

Coulin Estate where fish were collected.

CONFLICT OF INTEREST:

We declare we have no competing interests

DATA AVAILABILITY

Upon acceptance, data will be available in dryad.

REFERENCE

Ackerly K.L. & Ward A.B. (2016) How temperature-induced variation in musculoskeletal anatomy
affects escape performance and survival of zebrafish (Danio rerio). *Journal of Experimental*
Zoology Part A: Ecological Genetics and Physiology, **325,** 25-40.
Adams C.E., Fraser D., Huntingford F.A., Greer R.B., Askew C.M. & Walker A.F. (1998) Trophic
polymorphism amongst Arctic charr from Loch Rannoch, Scotland. *Journal of Fish Biology,* **52,**
1259-1271.
Adams C.E., Woltering C. & Alexander G. (2003) Epigenetic regulation of trophic morphology through
feeding behaviour in Arctic charr, Salvelinus alpinus. *Biological Journal of the Linnean Society,*
78, 43-49.
Alley R.B., Marotzke J., Nordhaus W.D., Overpeck J.T., Peteet D.M., Pielke R.A., Pierrehumbert R.T.,
Rhines P.B., Stocker T.F., Talley L.D. & Wallace J.M. (2003) Abrupt Climate Change. *Science,*
299, 2005.
Angilletta M.J., Jr., Steury T.D. & Sears M.W. (2004) Temperature, Growth Rate, and Body Size in
Ectotherms: Fitting Pieces of a Life-History Puzzle1. *Integrative and Comparative Biology,* **44,**
498-509.

Berg M.P. & Ellers J. (2010) Trait plasticity in species interactions: a driving force of community
dynamics. *Evolutionary Ecology,* **24,** 617-629.
Bolnick D.I. & Lau O.L. (2008) Predictable patterns of disruptive selection in stickleback in postglacial
lakes. *American Naturalist,* **172**.
Burggren W. (2018) Developmental phenotypic plasticity helps bridge stochastic weather events
associated with climate change. *The Journal of Experimental Biology,* **221,** jeb161984.
Burton T. & Metcalfe N.B. (2014) Can environmental conditions experienced in early life influence
future generations? *Proceedings of the Royal Society B: Biological Sciences,* **281,** 20140311.
Campbell C.S. (2019) *The effects of climate change on phenotypic plasticity in Arctic charr*, University
of Glasgow.
Campinho M.A., Moutou K.A. & Power D.M. (2004) Temperature sensitivity of skeletal ontogeny in
Oreochromis mossambicus. *Journal of Fish Biology,* **65,** 1003-1025.
Charmantier A., Mccleery R.H., Cole L.R., Perrins C., Kruuk L.E.B. & Sheldon B.C. (2008) Adaptive
Phenotypic Plasticity in Response to Climate Change in a Wild Bird Population. *Science,* **320,**
800.
Chavarie L., Dempson J.B., Schwarz C., Reist J., Power G. & Power M. (2010) Latitudinal variation in
growth among Arctic charr in eastern North America: evidence for countergradient variation?
Hydrobiologia, **650,** 161-177.
Chevin L.-M., Lande R. & Mace G.M. (2010) Adaptation, plasticity, and extinction in a changing
environment: towards a predictive theory. *PLoS Biology,* **8,** e1000357-e1000357.
Collins M., Knutti R., Arblaster J., Dufresne J.-L., Fichefet T., Friedlingstein P., Gao X., Gutowski W.J.,
Johns T. & Krinner G. (2013) Long-term climate change: projections, commitments and
irreversibility. In: *Climate Change 2013-The Physical Science Basis: Contribution of Working*
Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change pp.
1029-1136. Cambridge University Press.
Corno G. & Jürgens K. (2006) Direct and Indirect Effects of Protist Predation on Population Size
Structure of a Bacterial Strain with High Phenotypic Plasticity. *Applied and Environmental*
Microbiology, **72,** 78.
Crawley M.J. (2012) *The R book,* John Wiley & Sons.
Crozier L.G. & Hutchings J.A. (2014) Plastic and evolutionary responses to climate change in fish.
Evolutionary Applications, **7,** 68-87.
Donelson J.M., Salinas S., Munday P.L. & Shama L.N.S. (2018) Transgenerational plasticity and climate
change experiments: Where do we go from here? *Global Change Biology,* **24,** 13-34.
Donelson J.M., Wong M., Booth D.J. & Munday P.L. (2016) Transgenerational plasticity of reproduction
depends on rate of warming across generations. *Evolutionary Applications,* **9,** 1072-1081.
Dryden I.L. & Mardia K.V. (1998) Statistical shape analysis: Wiley series in probability and statistics.
New York, NY: John Wiley & Sons, Ltd.
Duclos K.K., Hendrikse J.L. & Jamniczky H.A. (2019) Investigating the evolution and development of
biological complexity under the framework of epigenetics. *Evolution & Development,* **21,** 276-
293.
Elmer K.R. (2016) Genomic tools for new insights to variation, adaptation, and evolution in the
salmonid fishes: a perspective for charr. *Hydrobiologia***,** DOI:10.1007/s10750-10015-12614-
10755.
Frankino W.A., Zwaan B.J., Stern D.L. & Brakefield P.M. (2005) Natural Selection and Developmental
Constraints in the Evolution of Allometries. *Science,* **307,** 718.
Franklin O.D., Skúlason S., Morrissey M.B. & Ferguson M.M. (2018) Natural selection for body shape
in resource polymorphic Icelandic Arctic charr. *Journal of Evolutionary Biology,* **31,** 1498-1512.
Garduño-Paz M.V. & Adams C.E. (2010) Discrete prey availability promotes foraging segregation and
early divergence in Arctic charr, Salvelinus alpinus. *Hydrobiologia,* **650,** 15-26.
Georga I. & Koumoundouros G. (2010) Thermally induced plasticity of body shape in adult zebrafish
Danio rerio (Hamilton, 1822). *Journal of Morphology,* **271,** 1319-1327.

Ghalambor C.K., Mckay J.K., Carroll S.P. & Reznick D.N. (2007) Adaptive versus non-adaptive
phenotypic plasticity and the potential for contemporary adaptation in new environments.
Functional Ecology, 21, 394-407.
Gillooly J.F., Brown J.H., West G.B., Savage V.M. & Charnov E.L. (2001) Effects of size and temperature
on metabolic rate. *Science, 293,* 2248-2251.
Gillooly J.F., Charnov E.L., West G.B., Savage V.M. & Brown J.H. (2002) Effects of size and temperature
on developmental time. *Nature, 417,* 70-73.
Heino J., Virkkala R. & Toivonen H. (2009) Climate change and freshwater biodiversity: detected
patterns, future trends and adaptations in northern regions. *Biological Reviews, 84,* 39-54.
Heino M. (2014) Quantitative traits. In: *Stock identification methods* pp. 59-76. Elsevier.
Hooker O.E., Barry J., Van Leeuwen T.E., Lyle A., Newton J., Cunningham P. & Adams C.E. (2016a)
Morphological, ecological and behavioural differentiation of sympatric profundal and pelagic
Arctic charr (Salvelinus alpinus) in Loch Dughaill Scotland. *Hydrobiologia, 783,* 209-221.
Hooker O.E., Barry J., Van Leeuwen T.E., Lyle A., Newton J., Cunningham P. & Adams C.E. (2016b)
Morphological, ecological and behavioural differentiation of sympatric profundal and pelagic
Arctic charr (*Salvelinus alpinus*) in Loch Dughaill Scotland. *Hydrobiologia*, DOI:
10.1007/s10750-10015-12599-10750.
Imsland A.K., Foss A., Folkvord A., Stefansson S.O. & Jonassen T.M. (2005) The interrelation between
temperature regimes and fish size in juvenile Atlantic cod (Gadus morhua): effects on growth
and feed conversion efficiency. *Fish Physiology and Biochemistry, 31,* 347.
Jacobson B., Grant J.W.A. & Peres-Neto P.R. (2015) The interaction between the spatial distribution of
resource patches and population density: consequences for intraspecific growth and
morphology. *Journal of Animal Ecology, 84,* 934-942.
Kavanagh K.D., Haugen T.O., Gregersen F., Jernvall J. & Vøllestad L.A. (2010) Contemporary
temperature-driven divergence in a Nordic freshwater fish under conditions commonly
thought to hinder adaptation. *BMC Evolutionary Biology, 10,* 350.
Kingsolver J.G., Izem R. & Ragland G.J. (2004) Plasticity of Size and Growth in Fluctuating Thermal
Environments: Comparing Reaction Norms and Performance Curves1. *Integrative and
Comparative Biology, 44,* 450-460.
Klemetsen A. (2010) The charr problem revisited: exceptional phenotypic plasticity promotes
ecological speciation in postglacial lakes. *Freshwater Reviews, 3,* 49-74.
Klemetsen A. (2013) The most variable vertebrate on earth. *Journal of Ichthyology, 53,* 781-791.
Klingenberg C.P. (2011) MorphoJ: an integrated software package for geometric morphometrics.
Molecular Ecology Resources, 11, 353-357.
Klingenberg C.P. & Mcintyre G.S. (1998) GEOMETRIC MORPHOMETRICS OF DEVELOPMENTAL
INSTABILITY: ANALYZING PATTERNS OF FLUCTUATING ASYMMETRY WITH PROCRUSTES
METHODS. *Evolution, 52,* 1363-1375.
Lazić M.M., Carretero M.A., Crnobrnja-Isailović J. & Kaliontzopoulou A. (2015) Effects of Environmental
Disturbance on Phenotypic Variation: An Integrated Assessment of Canalization,
Developmental Stability, Modularity, and Allometry in Lizard Head Shape. *The American
Naturalist, 185,* 44-58.
Lazić M.M., Kaliontzopoulou A., Carretero M.A. & Crnobrnja-Isailović J. (2013) Lizards from urban areas
are more asymmetric: using fluctuating asymmetry to evaluate environmental disturbance.
PloS one, 8, e84190-e84190.
Lubchenco J. & Cubit J. (1980) Heteromorphic Life Histories of Certain Marine Algae as Adaptations to
Variations in Herbivory. *Ecology, 61,* 676-687.
Maitland P.S. & Adams C.E. (2018) *Arctic Charr in the Lochs of Scotland: An Assessment of Distribution
and Status,* Fast-Print Publishing.
Mcphee M.V., Noakes D.L. & Allendorf F.W. (2012) Developmental rate: A unifying mechanism for
sympatric divergence in postglacial fishes? *Curr. Zool., 58,* 21-34.

Merilä J. & Hendry A.P. (2014) Climate change, adaptation, and phenotypic plasticity: the problem and
the evidence. *Evolutionary Applications,* **7,** 1-14.
Mitteroecker P. (2009) The Developmental Basis of Variational Modularity: Insights from Quantitative
Genetics, Morphometrics, and Developmental Biology. *Evolutionary Biology,* **36,** 377-385.
Morris M.R. (2014) Plasticity-mediated persistence in new and changing environments. *International
journal of evolutionary biology,* **2014**.
Neuheimer A.B. & Taggart C.T. (2007) The growing degree-day and fish size-at-age: the overlooked
metric. *Canadian Journal of Fisheries and Aquatic Sciences,* **64,** 375-385.
Neuheimer A.B., Thresher R.E., Lyle J.M. & Semmens J.M. (2011) Tolerance limit for fish growth
exceeded by warming waters. *Nature Climate Change,* **1,** 110-113.
Noble D.W., Stenhouse V. & Schwanz L.E. (2018) Developmental temperatures and phenotypic
plasticity in reptiles: a systematic review and meta-analysis. *Biological Reviews,* **93,** 72-97.
Nosil P. (2012) *Ecological speciation,* Oxford University Press.
O'dea R.E., Lagisz M., Hendry A.P. & Nakagawa S. (2019) Developmental temperature affects
phenotypic means and variability: A meta-analysis of fish data. *Fish and Fisheries,* **20,** 1005-
1022.
O'dea R.E., Noble D.W., Johnson S.L., Hesselson D. & Nakagawa S. (2016) The role of non-genetic
inheritance in evolutionary rescue: epigenetic buffering, heritable bet hedging and epigenetic
traps. *Environmental Epigenetics,* **2**.
Olsson J., Svanbäck R. & Eklöv P. (2006) Growth rate constrain morphological divergence when driven
by competition. *Oikos,* **115,** 15-22.
Orizaola G. & Laurila A. (2009) Microgeographic variation in temperature-induced plasticity in an
isolated amphibian metapopulation. *Evolutionary Ecology,* **23,** 979.
Parmesan C. (2006) Ecological and Evolutionary Responses to Recent Climate Change. *Annual Review
of Ecology, Evolution, and Systematics,* **37,** 637-669.
Parsons K.J. & Albertson R.C. (2009) Roles for Bmp4 and CaM1 in Shaping the Jaw: Evo-Devo and
Beyond. *Annual Review of Genetics,* **43,** 369-388.
Parsons K.J., Sheets H.D., Skúlason S. & Ferguson M.M. (2011) Phenotypic plasticity, heterochrony and
ontogenetic repatterning during juvenile development of divergent Arctic charr (Salvelinus
alpinus). *Journal of Evolutionary Biology,* **24,** 1640-1652.
Parsons K.J., Skúlason S. & Ferguson M. (2010) Morphological variation over ontogeny and
environments in resource polymorphic arctic charr (Salvelinus alpinus). *Evolution &
Development,* **12,** 246-257.
Pigliucci M. & Preston K. (2004) *Phenotypic integration: studying the ecology and evolution of complex
phenotypes,* Oxford University Press.
Pittman K., Yúfera M., Pavlidis M., Geffen A.J., Koven W., Ribeiro L., Zambonino-Infante J.L. & Tandler
A. (2013) Fantastically plastic: fish larvae equipped for a new world. *Reviews in Aquaculture,*
5, S224-S267.
Poesch M.S., Chavarie L., Chu C., Pandit S.N. & Tonn W. (2016) Climate Change Impacts on Freshwater
Fishes: A Canadian Perspective. *Fisheries,* **41,** 385-391.
Raffard A., Cucherousset J., Santoul F., Di Gesu L. & Blanchet S. (2019) Intraspecific variation alters
ecosystem and next-generation performance as much as temperature. *bioRxiv,* 332619.
Ramler D., Mitteroecker P., Shama L.N.S., Wegner K.M. & Ahnelt H. (2014) Nonlinear effects of
temperature on body form and developmental canalization in the threespine stickleback.
Journal of Evolutionary Biology, **27,** 497-507.
Reed T.E., Waples R.S., Schindler D.E., Hard J.J. & Kinnison M.T. (2010) Phenotypic plasticity and
population viability: the importance of environmental predictability. *Proceedings of the Royal
Society B: Biological Sciences,* **277,** 3391-3400.
Robinson B.W. & Parsons K.J. (2002) Changing times, spaces, and faces: tests and implications of
adaptive morphological plasticity in the fishes of northern postglacial lakes. *Canadian Journal
of Fisheries and Aquatic Sciences,* **59,** 1819-1833.

Robinson B.W. & Wardrop S.L. (2002) Experimentally Manipulated Growth Rate in Threespine
Sticklebacks: Assessing Trade Offs with Developmental Stability. *Environmental Biology of*
Fishes, **63,** 67-78.
Scheepens J.F., Deng Y. & Bossdorf O. (2018) Phenotypic plasticity in response to temperature
fluctuations is genetically variable, and relates to climatic variability of origin, in Arabidopsis
thaliana. *AoB PLANTS,* **10**.
Schluter D. (2000) *The ecology of adaptive radiation,* OUP Oxford.
Schoener T.W. (2011) The Newest Synthesis: Understanding the Interplay of Evolutionary and
Ecological Dynamics. *Science,* **331,** 426.
Sfakianakis D.G., Koumoundouros G., Divanach P. & Kentouri M. (2004) Osteological development of
the vertebral column and of the fins in Pagellus erythrinus (L. 1758). Temperature effect on
the developmental plasticity and morpho-anatomical abnormalities. *Aquaculture,* **232,** 407-
424.
Simonsen M.K., Siwertsson A., Adams C.E., Amundsen P.-A., Præbel K. & Knudsen R. (2017) Allometric
trajectories of body and head morphology in three sympatric Arctic charr (Salvelinus alpinus
(L.)) morphs. *Ecology and Evolution,* **7,** 7277-7289.
Skúlason S., Parsons K.J., Svanbäck R., Räsänen K., Ferguson M.M., Adams C.E., Amundsen P.-A.,
Bartels P., Bean C.W., Boughman J.W., Englund G., Guðbrandsson J., Hooker O.E., Hudson
A.G., Kahilainen K.K., Knudsen R., Kristjánsson B.K., Leblanc C.a.L., Jónsson Z., Öhlund G., Smith
C. & Snorrason S.S. (2019) A way forward with eco evo devo: an extended theory of resource
polymorphism with postglacial fishes as model systems. *Biological Reviews,* **0**.
Skulason S. & Smith T.B. (1995) Resource polymorphisms in vertebrates. *Trends in Ecology & Evolution,*
10, 366-370.
Solomon S. (2007) *Climate change 2007-the physical science basis: Working group I contribution to the*
fourth assessment report of the IPCC, Cambridge University Press.
Svanbäck R., Mario Pineda-Krch A.M.D., Krch M., Doebeli M., Associate Editor: Claire De M. & Editor:
Donald L.D. (2009) Fluctuating Population Dynamics Promotes the Evolution of Phenotypic
Plasticity. *American Naturalist,* **174,** 176-189.
Taylor E.B. (1999) Species pairs of north temperate freshwater fishes: Evolution, taxonomy, and
conservation. *Reviews in Fish Biology and Fisheries,* **9,** 299-324.
Team R.C. (2014) R: A language and development for statistical computing.
Van Der Have T.M. & De Jong G. (1996) Adult Size in Ectotherms: Temperature Effects on Growth and
Differentiation. *Journal of Theoretical Biology,* **183,** 329-340.
Vasseur D.A., Delong J.P., Gilbert B., Greig H.S., Harley C.D.G., Mccann K.S., Savage V., Tunney T.D. &
O'connor M.I. (2014) Increased temperature variation poses a greater risk to species than
climate warming. *Proceedings of the Royal Society B: Biological Sciences,* **281,** 20132612.
West-Eberhard M.J. (2003) *Developmental plasticity and evolution,* Oxford University Press.
Westneat D.F., Wright J. & Dingemanse N.J. (2015) The biology hidden inside residual within-individual
phenotypic variation. *Biological Reviews,* **90,** 729-743.
Whitman D.W. & Agrawal A.A. (2009) What is phenotypic plasticity and why is it important. *Phenotypic*
plasticity of insects: Mechanisms and consequences, 1-63.
Willmore K.E., Zelditch M.L., Young N., Ah-Seng A., Lozanoff S. & Hallgrímsson B. (2006) Canalization
and developmental stability in the Brachyrrhine mouse. *Journal of Anatomy,* **208,** 361-372.
Woelders L., Lenaerts J.T.M., Hagemans K., Akkerman K., Van Hoof T.B. & Hoek W.Z. (2018) Recent
climate warming drives ecological change in a remote high-Arctic lake. *Scientific Reports,* **8,**
6858.
Wu C.-I. & Ting C.-T. (2004) Genes and speciation. *Nature Reviews Genetics,* **5,** 114-122.

Table 1. Parameter estimates from the minimum adequate generalized linear mixed model describing the effect of temperature, exposure and centroid size on shape DF scores).

	Estimate	SE	X^2	P
Intercept (Ambient)	-6.84	1.91	-	-
Exposure time	0.002	0.002	-	-
Elevated temperature	2.36	0.21	39.81	<0.01
Centroid size	1.63	0.48	-	-
Exposure time : Centroid Size	-0.0012	0.0005	7.23	<0.01

Table 2 Results from the CVA comparing ambient and elevated temperature raised fish to wild benthic and pelagic ecotypes from Loch Rannoch and Loch Dughaill.

Comparison	Degree of difference	Location	Procrustes distance	P value
Pelagic - Ambient	Less	Rannoch	0.157	<0.01
Pelagic - Elevated	More	Rannoch	0.158	<0.01
				<0.01
Benthic - Ambient	More	Rannoch	0.111	<0.01
Benthic - Elevated	Less	Rannoch	0.095	<0.01
				<0.01
Pelagic - Ambient	Less	Dughaill	0.121	<0.01
Pelagic - Elevated	More	Dughaill	0.149	<0.01
				<0.01
Benthic - Ambient	More	Dughaill	0.152	<0.01
Benthic - Elevated	Less	Dughaill	0.129	<0.01

Figure 1. Hypothetical expressed phenotype frequency distributions shown as reaction norms resulting from an environmental change, such as temperature. Black lines represent expressed phenotypes in a population subjected to exisiting conditions and grey line represent population expression of phenotype subjected to the new environmental conditions. In panel a), the population doesn't express any change in phenotypic mean or in phenotypic variation when exposed to the new environmental conditions. In panel b), the population shows a shift in the expressed phenotypic mean but not in phenotypic variation when exposed to the new environmental regime. In panel c), population the phenotypic mean stays the same after the environmental condition changes, but the variation in expressed phenotypic increases in the population exposed to the new environmental regime. In panel d), the population (ecotype) subjected to the new environmental conditions shift both in phenotype mean and in phenotypic variation. Direction and magnitude of variation in panel b-d are not absolute, and will be species and location dependent as well of the strength and time of exposure to the environmental variables.

Figure 2. Individual Discriminant Function Scores (DF1) for head shape of Arctic charr of different sizes (represented by centroid size). Three sample periods are given, 700dd are represented by circles, 1000dd by triangles and 1400dd by squares. Ambient exposed fish are denoted with open symbols; elevated temperature fish are denoted with closed symbols. Polynomial regressions for both temperature treatments are showing allometric trajectories toward phenotypic divergence of head shape changes (see Figure A2).

Figure 3. Canonical Variate Analysis of 1400dd ambient and elevated temperature exposed
Arctic charr and sympatric polymorphic populations of Arctic charr from; A) Loch Rannoch,
B) Loch Dughaill. Elipses represent 95% confidence limits for means. Polymorphic charr
populations are represented by squares and experimental Arctic charr by circles. Open symbols
denote wild plankton feeding Arctic charr and Arctic charr raised at an ambient temperature.
Closed symbols denote benthic feeding Arctic charr and Arctic charr raised at an elevated
temperature. X and Y axis percentages denotes the amount of variation for each canonical
variate. Wire frames on CV1 are scaled at -8 and +8, wire frames for CV2 are scaled at -4 and
+4. C) and D) show mean (+/- 95% CL) for CV2 scores.

 Appendix

 Figure A1. Nine landmarks were used to characterise the shape of the head; Landmark (LM)
 1, the tip of the nose; LM2, the most posterior part of the upper jaw; LM3, edge of cranium
 directly above the eye; LM4, edge of cranium at the central point between LM3 and LM1 at a
 90 degree angle; LM5-8, most upper, posterior, lower and anterior parts of eye respectively;
 LM9, most posterior part of the gill operculum.

742 Figure A2. Possible pattern of allometric trajectories between temperature treatments of this
743 study, figure is modified from Simonsen *et al.* (2017). Allometric trajectory patterns can be a)
744 parallel, b) divergent, c) convergent, or d) common.

757
758   A

759
760   B

Figure A3. Individuals sampled at 1400dd, from the group a) raised at elevated temperature, with a morphology analogous to benthic feeding wild ecotype and b) raised at ambient temperature, with a morphology analogous to plankton feeding wild ecotype (parental population).

767 Figure A4. CV1 and CV2 scores derived from Canonical Variate Analysis of: parent fish (grey
768 squares), ambient temperature exposed fish sampled at 1400dd (open circles) and elevated
769 temperature exposed fish sampled at 1400dd (closed circles) with 95% confidence elipses of
770 centroid. Percentage on X and Y axes denotes the amount of variation explained by each
771 canonical variate.