A 3-hydroxyacyl-CoA dehydratase (DEH) involved in very long chain fatty acid synthesis is crucial for oocyst mitotic division but not cell size during parasite transmission in *Plasmodium*

David S. Guttery[1,2,*,+] Rajan Pandey[1,3,+], David J. P. Ferguson[4], Richard J. Wall[1,5], Declan Brady[1], Dinesh Gupta[3], Anthony A. Holder[6], and Rita Tewari[1,*]

Running title: DEH is essential for sporogony

[1]School of Life Sciences, Queens Medical Centre, University of Nottingham, Nottingham, UK

[2]The Leicester Cancer Research Centre, College of Life Sciences, University of Leicester, Leicester LE2 7LX, UK

[3]Translational Bioinformatics Group, International Centre for Genetic Engineering and Biotechnology, Aruna Asaf Ali Marg, New Delhi-110067, India

[4]Department of Biological and Medical Sciences, Faculty of Health and Life Science, Oxford Brookes University, Oxford, UK

[5]Wellcome Trust Centre for Anti-Infectives Research, School of Life Sciences, University of Dundee, Dundee DD1 5EH, UK

[6]The Francis Crick Institute, London NW1 1AT, UK

+contributed equally

*Corresponding author

E-Mail: dsg6@le.ac.uk

E-mail: rita.tewari@nottingham.ac.uk

Funding: This project was funded by MRC Investigator Award and MRC project grants to RT [G0900109, G0900278, MR/K011782/1]. RP was supported by Newton Bhabha PhD Placement Program (BT/IN/UK/DBT-BC/2015-16). AAH is supported by the Francis Crick Institute (FC010097) which receives its core funding from Cancer Research UK (FC010097), the UK Medical Research Council (FC010097), and the

Competing interests: The authors have declared that no competing interests exist.

Abstract

Lipid metabolism is a crucially important area of biochemistry, and cells use fatty acids for membrane biosynthesis, energy storage and the generation of signaling molecules. A 3-hydroxyacyl-CoA dehydratase – DEH – is a key member of this process. Here we further characterized in-depth the location and function of DEH, applying *in-silico* analysis*,* live cell imaging, reverse genetics and ultrastructure analysis using the mouse malaria model *Plasmodium berghei*. We show that DEH is evolutionarily conserved across eukaryotic species, with a single DEH in *Plasmodium* spp. and up to three orthologs in the other eukaryotes studied. Using DEH-GFP live-cell imaging, we observed strong GFP fluorescence throughout the life-cycle, with areas of localized expression in the cytoplasm and a circular ring pattern around the nucleus that colocalized with ER markers. Functional analysis showed that while Δ*deh* mutants undergo normal oocyst cell growth during early stages of sporogony, endomitotic cell division and sporozoite development are completely ablated, blocking parasite transmission from mosquito to vertebrate host. Ultrastructure analysis confirmed degeneration of Δ*deh* oocysts, and a complete lack of sporozoite budding. Overall, we show that DEH is evolutionarily conserved, localizes to the ER and plays a crucial role in sporogony, potentially through its involvement in fatty acid production.

Introduction

Malaria remains one of the world's deadliest infectious diseases. Caused by apicomplexan parasites belonging to genus *Plasmodium,* malaria is responsible for great socio-economic loss to affected countries. According to WHO reports, there were 212 million clinical cases of malaria infection and 429000 deaths in 2015 (Organisation, 2018), and growing resistance against existing drugs has further intensified this problem. Hence, there is a growing need to identify new biological pathways and proteins essential for parasitic growth and development in human hosts, which could act as suitable drug targets. *Plasmodium* parasites have a complex life cycle and require two hosts to complete the life cycle: vertebrates (during asexual stages) and invertebrates (during sexual stages) (Aly *et al.*, 2009). The disease is transmitted to vertebrate hosts by infected female *Anopheles* mosquitoes, which inject sporozoites into the dermis of the vertebrate host during a blood meal. The parasite enters the circulation, and once it invades the liver, and subsequently erythrocytes, undergoes several rounds of atypical closed mitotic cell division through multiple rounds of DNA replication and asynchronous nuclear division (termed schizogony) to produce merozoites that invade erythrocytes. During this period of cyclic asexual proliferation in the blood stream, a subpopulation undergoes gametocytogenesis to develop into male and female gametocytes, which are transmitted to a mosquito during its blood meal. Gamete development, fertilization and zygote formation occur in the mosquito midgut, leading to the differentiation of an infective ookinete, which undergoes meiosis, and penetrates the midgut wall to develop into an oocyst on the basal surface of the midgut, where further rounds of closed mitotic cell division occur. Thousands of sporozoites develop within each oocyst, and then egress into the haemocoel to invade the salivary glands and begin a new life cycle.

Lipid metabolism includes essential cellular processes that use fatty acids (FAs) in membrane biosynthesis, energy storage and the generation of signaling molecules. FA elongation and very long chain fatty acid (VLCFA) synthesis occurs in two stages, both consisting of a four-step cyclic process that results in addition of two carbons to the chain with each cycle. In humans, the process involves condensation of acyl-CoA with

malonyl-CoA to produce 3-ketoacyl-CoA (catalyzed by one of seven FA elongases), reduction of 3-ketoacyl-CoA by a 3-ketoacyl-CoA reductase (KAR) to 3-hydroxyacyl-CoA, dehydration of 3-hydroxyacyl-CoA to 2,3-trans-enoyl-CoA (catalysed by one of four 3-hydroxyacyl-CoA dehydratase isoenzymes: HACD1-4), and finally reduction to an acyl-CoA with two additional carbon chain units by 2,3-trans-enoyl-CoA reductase (TER) (Kihara, 2012). HACD1-4 were initially annotated as PTPLA, PTPLB, PTPLAD1 and PTPLAD2, respectively due to their similarities to the yeast Phs1 gene product (Ikeda *et al.*, 2008). However, they were renamed to reflect their function as 3-hydroxyacyl Co-A dehydratases and their relatedness. HACD1 and HACD2 genes restored growth of yeast SAY32 Phs1-defective cells, indicating that they are functional homologues of Phs1, i.e. 3-hydroxyacyl-CoA dehydratases. Further, studies have indicated that HACD1 has an essential role in myoblast proliferation and differentiation (Lin *et al.*, 2012), with HACD1-deficient cell lines displaying S-phase arrest, compromised G2/M transition and retarded cell growth. Studies of the *Arabidopsis thaliana* Phs1 homologue PASTICCINO2 or PAS2 showed the protein has an essential role in VLCFA synthesis (Bach *et al.*, 2008), as well as being essential during cell division, proliferation and differentiation (Bellec *et al.*, 2002). Further, *Arabidopsis* PAS2 complements Phs1 function in a yeast mutant defective for FA elongation (Morineau *et al.*, 2016). PAS2 interacted with FA elongase subunits in the endoplasmic reticulum (ER) and in its absence 3-hydroxyacyl-CoA accumulates, as expected from loss of a dehydratase involved in FA elongation. Similarly, in the yeast *Saccharomyces cerevisiae* VLCFA synthesis is also catalyzed in the ER by a multi-protein elongase complex, following a similar reaction pathway as mitochondrial or cytosolic fatty acid synthesis (Tehlivets *et al.*, 2007).

In Apicomplexans, the process of fatty acid synthesis and assembly into more complex molecules is critical for their growth and development, while also determining their ability to colonize the host and to cause disease. They acquire lipids through *de novo* synthesis and through scavenging from the host (Mazumdar *et al.*, 2007), and simple components like mosquito-derived lipids determine within-host *Plasmodium* virulence by shaping sporogony and metabolic activity, affecting the quantity and quality of

sporozoites, respectively (Costa *et al.*, 2018). Fatty acid synthesis (FAS) occurs in the apicoplast via the type II FAS (FASII) pathway, followed by fatty acid elongation (FAE) on the cytoplasmic face of the ER through the elongase (ELO) pathway (Ramakrishnan *et al.*, 2012, Ramakrishnan *et al.*, 2013). Studies on whether FAS is essential in *Plasmodium* are contradictory. In *Plasmodium yoelii* the FASII enzymes are only essential during liver stages (Yu *et al.*, 2008, Vaughan *et al.*, 2009); whereas in *Plasmodium falciparum,* genetic disruption of the FASII enzymes FabI and FabB/F results in complete abolition of sporogony (van Schaijk *et al.*, 2014). Specifically, day 17 to day 23 after mosquito feeding, FabB/F mutant oocysts appeared to degenerate, and expression of the marker GFP was barely detectable, confirming that VLCFA synthesis is crucial for commencement of sporogony. The enzymatic steps of the ELO process are similar to those in the FASII pathway in the apicoplast (Tarun *et al.*, 2009); however, the growing chain is held by CoA instead of acyl carrier protein (ACP). In *Toxoplasma gondii,* the activity of the ELO-pathway is considered an alternative route to FASII-independent ^{14}C-acetate incorporation (Bisanz *et al.*, 2006) and is engaged in conventional elongation rather than *de novo* synthesis (Mazumdar *et al.*, 2007). Indeed, *P. falciparum* parasites with no functional FASII pathway can still elongate fatty acids; possibly because of the activity of the ELO pathway (Yu *et al.*, 2008).

In a genome-wide study of *Plasmodium berghei* (Pb) protein phosphatases, we identified 30 phosphatase genes together with one for a predicted protein tyrosine phosphatase-like protein, PbPTPLA, which was shown to be essential for sporozoite formation and completion of the parasite life cycle, but not fully characterized (Guttery *et al.*, 2014). However, despite the original annotation as an inactive PTP-like protein (Andreeva *et al.*, 2008, Wilkes *et al.*, 2008, Guttery *et al.*, 2014, Pandey *et al.*, 2014), more recent functional studies indicate that it is a key component of the VLCFA elongation cycle – more specifically the ELO pathway as a 3-hydroxyacyl-CoA dehydratase (DEH) (Stanway *et al.*, 2019). Therefore, to investigate further the role of DEH in *Plasmodium* development, we performed an in-depth genotypic and phenotypic analysis of the protein, using *in-silico,* genetic manipulation and cell biological techniques. We show evolutionary conservation of DEH in the model organisms

examined here. Furthermore, we show that PbDEH is located at the ER and is essential for cell division and parasite budding within oocyst but not cell growth of oocysts, thereby blocking parasite transmission.

Methods

Ethics statement

All animal work was performed following ethical approval and was carried out under United Kingdom Home Office Project Licence 40/3344, in accordance with the UK 'Animals (Scientific Procedures) Act 1986' and in compliance with 'European Directive 86/609/EEC' for the protection of animals used for experimental purposes. Six- to eight-week old female Tuck-Ordinary (TO) (Harlan) or Swiss Webster (Charles River) outbred mice were used for all experiments.

Identification of conserved domains and evolutionary lineage

The deduced amino acid sequence of PBANKA_134650 (PbPTPLA) now classified as DEH in the manuscript, was retrieved from PlasmoDB (release 27) (Aurrecoechea *et al.*, 2009). Conserved domains in PbPTPLA (DEH) were identified using the Conserved Domain Database (CDD) (Marchler-Bauer *et al.*, 2011), the Simple Modular Architecture Research Tool (SMART) (Schultz *et al.*, 1998) and Protein Family Database (PFAM) (Finn *et al.*, 2008). The deduced amino acid sequence and individual conserved domains were used as BLAST (BLASTP) queries to identify orthologues in PlasmoDB and NCBI protein databases. OrthoMCL database (version 5) was used to identify and retrieve *P. berghei* orthologues (Table S1) (Li *et al.*, 2003). Multiple sequence alignment was performed for the retrieved sequences using ClustalW (Larkin *et al.*, 2007). ClustalW alignment parameters included gap opening penalty (GOP) 10 and gap extension penalty (GOE) 0.1 for the pairwise sequence alignment; GOP 10 and GOE 0.2 was used for multiple sequence alignment. A gap separation distance cut-off of 4 and Gonnet protein weight matrix was used for the alignments. Residue-specific penalty and hydrophobic penalties were used, whereas end gap separation and negative matrix were excluded in the ClustalW alignments. The phylogenetic tree was inferred using the neighbor-joining method, computing the evolutionary distances using the Jones-Taylor-

Thornton (JTT) model for amino acid substitution with the Molecular Evolutionary Genetics Analysis software (MEGA 6.0) (Tamura *et al.*, 2013). Gaps and missing data were treated using a partial deletion method with 95% site-coverage cut-off and 1000 bootstrap replicates to generate a phylogenetic tree. iTOL was used to visualize the phylogenetic tree (Letunic *et al.*, 2019). The STRING database was used to identify PTPLA interacting proteins (Franceschini *et al.*). For STRING search, we used a cut-off of 0.70 for the parameters of neighborhood, gene fusion, co-occurrence, co-expression, experiments, databases and text mining results. Predotar (Small *et al.*, 2004) was used for inferring Pf and PbDEH subcellular localization.

Generation of transgenic parasites and genotype analysis

Details of GFP-tagged PTPLA (termed DEH-GFP in this study) and *deh* (PBANKA_1346500) knockout (KO) parasite lines (Δ*deh* in this study) are given in (Guttery *et al.*, 2014). For this study, the KO construct was transfected into the GFPCON wild-type line (Janse *et al.*, 2006), with 3 clones produced by serial dilution.

Parasite development in the mosquito

Anopheles stephensi mosquitoes (3–6 days old) were allowed to feed on anaesthetized mice infected with either wild type or mutant parasites at comparable gametocytemia as assessed by blood smears. Mosquitoes were dissected post-blood meal, on the days indicated. For midgut and salivary gland sporozoites, organs from 10-20 mosquitoes were pooled and homogenized, and released sporozoites were counted using a haemocytometer. For oocyst counts, midguts taken at day 7, 14 and 21 post-infection were harvested, mounted on a slide and oocysts counted using phase or fluorescence microscopy. To quantify sporozoites per oocyst (the ratio of number of sporozoites to number of oocysts), an equal number of mosquitoes from the same cage were used to count the number of oocysts and sporozoites. This number varied among experiments but at least 20 mosquitoes were used for each count. For light microscopy analysis of developing oocysts, at least 20 midguts were dissected from mosquitoes on the indicated days and mounted under Vaseline-rimmed cover slips. ER tracker (ThermoFisher) was used to perform co-localization studies according to manufacturer's

instructions. Images were collected with an AxioCam ICc1 digital camera fitted to a Zeiss AxioImager M2 microscope using a 63x oil immersion objective. Statistical analyses were performed using Graphpad prism software.

Electron Microscopy

The guts from mosquitoes harvested at 14 and 21 days post-infection were dissected and fixed in 2.5% glutaraldehyde in 0.1 M phosphate buffer and processed for electron microscopy (Guttery *et al.*, 2012).

Subcellular fractionation of parasite lysates and detection of DEH

Immunoprecipitation and subcellular fractionation of lysates containing GFP tagged DEH was performed as described previously (Guttery *et al.*, 2014). WT-GFP was used as the control protein in all experiments. In summary, cells from mouse blood infected with the DEH-GFP-expressing parasite were pelleted and then lysed in hypotonic buffer (10 mM Tris-HCl pH 8.4, 5 mM EDTA) containing protease inhibitors (Roche), freeze/thawed twice, incubated for 1 hr at 4°C and then centrifuged at 100,000 g for 30 min. The supernatant was collected as the soluble protein fraction (cytosol). The pellet was resuspended and washed in carbonate solution (0.1M Na_2CO_3, pH 11.0) containing protease inhibitors (Roche), and after incubation for 30 min at 4°C the sample was centrifuged again at 100,000 g for 30 min. The supernatant was saved as the peripheral membrane protein fraction (PMF). The residual pellet was solubilized in 4% SDS and 0.5% Triton X-100 in PBS, to form the integral membrane protein fraction (IMF). Samples from these three fractions, containing equal amounts of protein, were then analyzed by western blot using anti-GFP antibody.

Results

Phylogenetic analysis reveals that DEH is highly conserved among eukaryotes

Genome-wide analysis showed DEH is present in all the eukaryotic organisms studied here, which includes apicomplexans, yeast, fungi, plants, nematodes, insects, birds and mammals. The number of encoded DEH proteins was shown to vary from one to three in the studied organisms, with *Plasmodium* spp coding for a single DEH. Intriguingly,

both *Arabidopsis thaliana* and *Oryza sativa* encode three DEHs (*PAS2* and 2 *HACD* isozymes) each, as compared to two (HACD1 and HACD2) in *Homo sapiens* (plus two sharing relatively weak similarity – HACD3 and HACD4) and two in *Mus musculus*. Phylogenetic analysis using the neighbor joining method, clustered organisms based on their evolutionary relatedness (Figure S1, Table S1). In addition, the phylogenetic analysis suggests that gene duplication in non-chordata, chordata and plants where there are multiple DEH copies, may have happened from a single DEH gene independently to perform specific functions after divergence during evolution, based on the clustering of all DEH isoforms in the same cluster.

Plasmodium DEH does not contain the canonical PTPLA CXXGXXP motif and is predicted *in silico* to interact with factors associated with FAE

P. berghei (PBANKA_1346500) and *P. falciparum* (Pf; PF3D7_1331600) DEH genes are annotated as PTPLA (pfam04387) (Andreeva *et al.*, 2008, Wilkes *et al.*, 2008, Guttery *et al.*, 2014, Pandey *et al.*, 2014), the criterion for PTPLA being the presence of a PTP active site motif (CXXGXXR) but with arginine replaced by proline (CXXGXXP). However, CLUSTALW alignment of *Pb* and *Pf* protein sequences with the human and mouse HACD1 and HACD2 shows this motif is absent (Figure S2), indicating that *Plasmodium* DEHs cannot be classified as PTP-like proteins. Furthermore, STRING database analysis predicts that PfDEH interacts with FAE and FAS proteins, and many other proteins with an ER location (Figure S3). These proteins include 3-oxo-5-α-steroid 4-dehydrogenase (PBANKA_09127; PF3D7_1135900), Stearoyl-CoA δ-9 desaturase (PBANKA_1110700; PF3D7_0511200), putative long chain polyunsaturated fatty acid elongation enzyme (ELO-B, PBANKA_0104700; PF3D7_0605900 - involved in the FAE pathway) as well as β-hydroxyacyl-(Acyl-carrier-protein) dehydratase (FabZ), involved in stage 3 of fatty acid synthesis in the FASII pathway (Stanway *et al.*, 2019). In addition, interactome analysis revealed DEH interaction with a putative ER membrane protein, Acetyl-CoA transporter protein (PF10_0360).

DEH is expressed throughout the *Plasmodium* life-cycle stages and localized to the ER

To determine the expression profile and location of PbDEH, we used a single homologous recombination strategy to tag the 3' end of the endogenous *deh* locus with sequence coding for GFP (Guttery *et al.*, 2014), and then analyzed blood and mosquito stages of the life-cycle for GFP. Strong GFP fluorescence was observed throughout the life-cycle, with areas of localized expression in the cytoplasm and a circular ring formation around the nucleus (Figure 1). Predotar analysis (Small *et al.*, 2004) predicted an ER localization for both PbDEH and PfDEH. Colocalization with ER tracker confirmed the DEH-GFP location at the ER, in all parasite stages analyzed (Figure 2A), with subcellular fractionation of blood stage parasites confirming its integral membrane location (Figure 2B).

PbDEH is essential for mitotic cell division but not cell size of *Plasmodium* during oocyst development

Previous comparison of Δ*deh* and WT parasite lines highlighted the non-essential role of this gene for blood stage development (Guttery *et al.*, 2014). In this study we confirmed it is not essential during asexual blood stages, or for zygote development (Figure 3A). However, while the overall number of oocysts observed in Δ*deh* and WT lines was not significantly different (Guttery *et al.*, 2014), there was a significant reduction in Δ*deh* GFP-expressing oocysts beginning at day 7 and continuing through to day 21 post-infection (Figure 3B, C), with many appearing to be degenerating. Analysis of oocyst size revealed a small decrease from day 10 onwards in Δ*deh* lines compared to WT (Figure 3D), and by day 21 the vast majority of Δ*deh* oocysts expressed GFP no longer, and in the few that did GFP was present at very low levels or in fragmented patterns. However, it is important to note that Δ*deh* oocysts that continued to express GFP and showed faint DAPI staining of DNA, were similar in size to WT oocysts; whereas the vast majority of oocysts that reduced in size did not express GFP or stain with DAPI, suggesting they were dead. Analysis of salivary glands from mosquitoes infected with Δ*deh* parasites revealed no sporozoites, in contrast with the normal numbers of sporozoites in glands from WT-parasite infected mosquitoes (Figure 3E). Representative examples of Δ*deh* oocyst morphology and lack of sporozoite

development at all stages are shown in Figures 3F and G, highlighting oocyst degeneration, fragmented GFP expression and failure to form sporozoites.

Ultrastructure analysis confirmed oocyst degeneration and apoptotic nuclear chromatin condensation in Δ*deh* lines

To investigate further the marked differences in oocyst morphology and complete lack of sporozoite formation, we used electron microscopy to compare Δ*deh* and WT lines at 10-, 14- and 21-days post-infection. At 10 days, the oocysts of Δ*deh* and WT parasites were of a similar size. However, Δ*deh* oocysts showed numerous cytoplasmic vacuoles (Figure 4a) with evidence of dilatation of the nuclear membranes (Fig 4b), in contrast to WT oocysts which were completely filled with cytoplasm with numerous nuclear and mitochondrial profiles (Figure 4g, h). At fourteen days post-infection the WT parasite showed a mixture of early, mid and late stage oocysts with sporozoites at various stages of development (Figure 4i, j). In contrast, the majority of Δ*deh* oocysts exhibited advanced stages of degeneration with increased cytoplasmic vacuolation, dilitated nuclear membranes, and evidence of mitochondrial abnormalities (Fig 4c, d). There was little evidence of retraction of the plasmalemma from the oocyst wall and no evidence that sporozoite inner membrane complex (IMC) formation had been initiated in any of the oocysts. At day 21 post-infection, all Δ*deh* oocysts were in an advanced stage of degeneration - almost completely vacuolated with a few nuclei appearing to have undergone apoptotic-like nuclear chromatin condensation (Fig 4e, f), in contrast to WT oocysts, which were mostly mature with numerous fully formed and free sporozoites (Fig 4k, l) although a few degenerate oocysts were observed.

Discussion

Lipid metabolism is essential for cellular function, and includes critical pathways for FA synthesis and elongation. DEH is a 3-hydroxyacyl-CoA dehydratase involved in VLCFA synthesis, which interacts with several elongase units, is located at the ER (Beaudoin *et al.*, 2009, Morineau *et al.*, 2016) and has an essential role during development, differentiation, and maintenance of a number of tissue types (Li *et al.*, 2000, Bellec *et*

al., 2002, Pele *et al.*, 2005). In this study, we examined the location and function of *Plasmodium* DEH using *in silico,* genetic manipulation and cell biological techniques.

In our previous phosphatome study, a putative, catalytically inactive, PTP-like protein with an essential role during sporogony was identified (Guttery *et al.*, 2014), which had been classified as a putative PTPLA by others (Andreeva *et al.*, 2008, Wilkes *et al.*, 2008, Pandey *et al.*, 2014) based on high sequence similarity and e-score values. However, a recent genome-wide functional screen in *P. berghei* showed that PbPTPLA has an essential role in lipid metabolism, specifically during the ELO pathway as a 3-hydroxyacyl dehydratase (DEH) (Stanway *et al.*, 2019). The specific criterion for a PTP-like protein is the presence of a CXXGXXP motif (i.e. the CXXGXXR motif of PTPs, but with the arginine replaced by proline). However, we show here that this motif is not present in either *P. falciparum* or *P. berghei* protein and this, along with its proven function in lipid metabolism (Stanway *et al.*, 2019), suggests that the classification as a phosphatase-like protein is erroneous. Our *in silico* interactome analysis suggests that PfDEH interacts with a number of proteins involved in lipid metabolism, confirming previous functional findings (Stanway *et al.*, 2019) and adding further weight to its annotation as a 3-hydroxyacyl-CoA dehydratase.

Studies in mammalian systems have suggested that the ER-bound DEH catalyzes the third of four reactions in the long-chain FA elongation cycle (Ikeda *et al.*, 2008). Our detailed GFP-based localization analyses showed that the protein is expressed strongly throughout the life-cycle, with protein expression at localized areas in the cytoplasm and as a circular ring-like structure around the nucleus. *In-silico* analysis using Predotar and microscopy-based co-localization using ER tracker confirmed the ER location, consistent with previous studies suggesting a role in FAS in *Plasmodium* (Stanway *et al.*, 2019)*.* Phenotypic analysis of DEH function throughout the life cycle confirmed the results of our previous study (Guttery *et al.*, 2014), highlighting that it is essential for oocyst maturation and sporozoite development, but dispensable for asexual blood stage development (Bushell *et al.*, 2017). Time-course analysis at days 7, 14 and 21 after mosquito infection showed that while early-stage Δ*deh* oocysts were comparable in size

to WT oocysts, they begin to degenerate at an early stage of development, with a significant decrease in GFP-expressing oocysts even at day 7 post-infection, and as seen previously in other FAE-critical mutant parasites (Stanway *et al.*, 2019). Ultrastructure analysis confirmed that at 14 days post-infection, Δ*deh* oocysts were at an advanced state of degeneration, with no evidence of sporozoite development. Retraction of the oocyst plasmalemma (the parasite plasma membrane) from the oocyst capsule is a crucial first stage in sporozoite development, where sporoblast formation is followed by thousands of sporozoites budding off into the space delineated by the capsule (Aly *et al.*, 2009). A model of this process is detailed in (Burda *et al.*, 2017). Our study suggests that initiation of mitosis, which results in sporozoite development, does not even commence in Δ*deh* oocysts, since retraction of the plasmalemma and initiation of daughter IMC formation is ablated. The phenotype is similar to that of a cyclin-3 mutant (Roques *et al.*, 2015), with defects leading to abnormalities in membrane formation, vacuolation and subsequent cell death during the later stages of sporogony. However, in contrast to the cyclin-3 mutant, oocyst growth was not affected but sporogony was completely ablated in Δ*deh* parasites, and no transmission was observed in bite-back experiments. This suggests that the parasite cannot scavenge VLCFA from its mosquito host environment, and that DEH (and therefore the ELO pathway) is critical for oocyst mitotic maturation and differentiation. The cells were unable to progress further to form additional lobes and start sporozoite budding in its absence, although the oocyst size was not grossly affected, suggesting that two independent processes drive oocyst formation and sporogony, respectively.

While FASII activity is exclusively in the apicoplast (Shears *et al.*, 2015), our study showed that DEH-GFP is located at the ER, suggesting that it is an active component of the ELO pathway. The genes involved in the ELO pathway include members of the ELO family (1, 2 and 3), of which *ELO2* and *ELO3* are involved in keto- and enoyl-reduction (Kohlwein *et al.*, 2001). In yeast, the dehydratase step is carried out by the DEH-like homologue, Phs1, which has also been characterized as a cell cycle protein with mutants defective in the G2/M phase (Yu *et al.*, 2006). Gene knockout studies for any ELO proteins are few, with a single genome-wide functional analysis showing that the *P.*

berghei homologue of PF3D7_0605900 (a putative long chain polyunsaturated fatty acid elongation enzyme) is dispensable during the asexual blood stages (Bushell *et al.*, 2017). In addition, the comprehensive analysis of FAE in *Plasmodium* by Stanway et al. (Stanway *et al.*, 2019) showed that mutants of a ketoacyl-CoA reductase (KCR) have an identical phenotype to our DEH gene knockout lines, with normal development of ookinetes and oocysts gradually disappearing over the course of development, resulting in the complete ablation of sporogony. In contrast, ELO-A (stage 1 of VLCFA synthesis) mutants were critical for liver stage development. This suggests that reduction of ketoacyl-CoA to hydroxyacyl-CoA and subsequent dehydration of hydroxyacyl-CoA to enoyl-CoA (i.e. stages 2 and 3 of VLCFA synthesis) are the most crucial stages for oocyst maturation and sporogony; whereas the lack of a phenotype during sporogony of the ELO-A deletion may suggest functional redundancy and/or a compensatory mechanism such as an overlapping specificity for condensation of malonyl-CoA by either ELO-B or ELO-C, as suggested in *Trypanosoma brucei* (Lee *et al.*, 2006) and *Toxoplasma gondii* (Ramakrishnan *et al.*, 2012).

In conclusion, our PbDEH analysis using various *in-silico, in-vitro and in-vivo* approaches provides important insights into the crucial role DEH plays during VLCFA synthesis, and how disruption of the gene can affect parasite development in the mosquito. Future studies will elucidate further how lipid metabolism in *Plasmodium* can be explored as a viable target for therapeutic intervention.

Figure legends:

Figure 1: DEH-GFP protein expression in stages of the parasite life cycle

Expression of DEH-GFP in rings, trophozoites, schizonts, gametocytes, zygotes, ookinetes, oocysts and sporozoites. P28, a cy3-conjugated antibody which recognizes P28 on the surface of zygotes, and ookinetes was used as a marker of the sexual stages. Note that the female gametocyte in this figure has not been activated, and is not expressing P28. Scale bar = 5 µm.

Figure 2: Co-localization of DEH-GFP and ER tracker

(A) Analysis of DEH-GFP localization using ER tracker in merozoites, zygotes, ookinetes and sporozoites. Scale bar = 5 µm. (B) Anti-GFP western blot for subcellular localization of DEH-GFP. PMF – Peripheral membrane protein fraction, IMF – Integral membrane protein fraction.

Figure 3: Phenotypic analysis of Δ*deh* lines

(A) Ookinete conversion as a percentage in Δ*deh* and WT lines. Ookinetes were identified using the marker P28 and defined as those cells that successfully differentiated into elongated 'banana shaped' ookinetes. Bar is the mean ± SEM. n = 3 independent experiments. (B) Total number of GFP-positive oocysts per infected mosquito, including normal and smaller oocysts, at 7, 10, 14, and 21 days post-infection for Δ*deh* and WT parasite lines. Bar is the mean ± SEM. n = 3 independent experiments (20 mosquitoes for each). P <0.001 for all time points. (C) Oocysts (10x magnification) of Δ*deh* and WT lines at 7, 10, 14 and 21 days post-infection. Scale bar = 100 µm. (D) Individual Δ*deh* and WT oocyst diameters (µm) at 7, 10, 14 and 21 days post-infection. Horizontal line indicates the mean from 3 independent experiments (20 mosquitoes for each) of Δ*deh* and WT. *p <0.05, **p <0.01, ***p <0.001. (E) Total number of sporozoites per mosquito from 21 days post-infection salivary glands for Δ*deh* and WT lines. Three independent experiments, n = 20 mosquitoes for each replicate. *** p <0.001. (F) Oocysts (63x magnification) in Δ*deh* and WT lines. Images show DIC, GFP at 7, 10, 14 and 21 dpi. Scale bar = 20 µm. (G) Representative examples of Δ*deh* and

WT oocysts (63x magnification) at 21 dpi showing fragmented GFP and Hoechst staining. Scale bar = 20 µm.

Figure 4: Ultrastructure analysis of oocyst development in Δ*deh* lines.

Electron micrographs of Δ*deh* (a-f) and WT (g-l) parasites at 10 days (a, b, g, h), 14 days (c, d, i, j) and 21 days (e, f, k, l) post infection. Bars represent 10 µm (a, c, e, g, i, k) and 1 µm (b, d, f, h, j, l). (a) Low power image of an early oocyst showing lucent area made up numerous vacuoles (V). (b) Detail from a similar stage to that in (a) showing part of the cytoplasm containing a nucleus with a nuclear pole (NP). Note the lucent area due to the separation of the inner (inm) and outer (onm) nuclear membranes. (c) Low power image of a mid-stage oocyst showing nuclear swelling (N) and increased numbers of lucent cytoplasmic vacuoles (V). (d) Detail part of the cytoplasm showing a swollen nucleus (N), membrane bound lucent vacuoles (V) and mitochondria (Mi) with vesicles embedded in electron dense material. (e) Low power image of a late stage oocyst with abnormal nuclei (N) and the cytoplasm filled with electron lucent vacuoles. (f) Detail from the central region of (e) showing the peripheral location of electron dense chromatin (Ch) typical of apoptotic changes, while the cytoplasm consists of numerous vacuoles (V). (g) Low power image of an early oocyst (end of growth phase) in which the cytoplasm completely fills the oocyst and contains many nuclear profiles (N). (h) Detail of the peripheral cytoplasm limited by the plasmalemma (P) containing mitochondria (M) and nuclei (N). OW – oocyst wall. (i) Mid stage oocyst showing the surface formation of numerous sporozoites (Sp). N – nucleus. (j) Detail showing partially formed sporozoites (Sp) budding from the surface of the cytoplasmic mass. N – nucleus, R – rhoptry. (k) Mature oocysts containing large number of fully form sporozoites (Sp). (l) Detail of cross sections through mature sporozoites (Sp) containing rhoptries (R) and micronemes (MN).

Figure S1: Phylogenetic analysis of DEH homologues across different species

Phylogenetic analysis for PbDEH/PTPLA orthologs was performed using the neighbor joining method with MEGA6 software. Genome wide DEH/PTPLA analysis shows its presence in all organisms studied, including species from all major eukaryotic phyla.

The analysis shows the clustering of organisms based on their evolutionary relatedness. Blue – Chordata; Orange – Non-chordata; Purple – Nematoda; Pink - Ameobozoa; Red – Apicomplexa, Euglenozoa and Metamonada; Green – Plants and Algae; Yellow - Fungi and Black - Others.

Figure S2: ClustalW alignment of *Plasmodium* DEH with human, mouse and other apicomplexan homologues

ClustalW alignment of *Homo sapiens* (Hs), *Mus musculus* (Ms), *Plasmodium* (Pf, Pb), *Toxoplasma* (Tg) and *Cyclospora* (Cc) homologues of DEH. For apicomplexans, the proteins are currently annotated as PTPLA. Highlighted in red is the CXXGXXP motif that defines PTP-like proteins.

Figure S3: Predicted interacting partners of PfDEH

STRING analysis of PfDEH for potential interacting partners. The table on the right gives the annotation and combined score.

Table S1: Organisms and protein sequences used for phylogenetic analysis.

Author contributions: Conceived and designed the experiments: RT, RP and AAH. Performed the experiments: RT, RJW, RP, DJPF and DSG. Analyzed the data: RT, AAH, RP, RJW, DJPF and DSG. Contributed reagents/materials/analysis tools: DJPF, AAH and RT. Wrote the paper: RP, DSG, RJW, AAH and RT. Performed the functional and GFP tagging experiments: RP, RJW, DSG and RT. Phylogenetics analysis: RP and DSG. RP and DG performed database searches, sequence-based analysis, and other bioinformatics analysis. Electron microscopy experiment: DJPF.

Acknowledgements:
We thank Dr. Aline Freville and Ms. Jessica Stock, University of Nottingham, for their help in animal work and western blot development, respectively.

References

Aly, A.S.I., Vaughan, A.M. and Kappe, S.H.I. (2009). Malaria Parasite Development in the Mosquito and Infection of the Mammalian Host. *Annual review of microbiology* **63,** 195-221.

Andreeva, A.V. and Kutuzov, M.A. (2008). Protozoan protein tyrosine phosphatases. *Int J Parasitol* **38,** 1279-1295.

Aurrecoechea, C., Brestelli, J., Brunk, B.P., Dommer, J., Fischer, S., Gajria, B.*, et al.* (2009). PlasmoDB: a functional genomic database for malaria parasites. *Nucleic acids research* **37,** D539-543.

Bach, L., Michaelson, L.V., Haslam, R., Bellec, Y., Gissot, L., Marion, J.*, et al.* (2008). The very-long-chain hydroxy fatty acyl-CoA dehydratase PASTICCINO2 is essential and limiting for plant development. *Proc Natl Acad Sci U S A* **105,** 14727-14731.

Beaudoin, F., Wu, X., Li, F., Haslam, R.P., Markham, J.E., Zheng, H.*, et al.* (2009). Functional characterization of the Arabidopsis beta-ketoacyl-coenzyme A reductase candidates of the fatty acid elongase. *Plant Physiol* **150,** 1174-1191.

Bellec, Y., Harrar, Y., Butaeye, C., Darnet, S., Bellini, C. and Faure, J.D. (2002). Pasticcino2 is a protein tyrosine phosphatase-like involved in cell proliferation and differentiation in Arabidopsis. *Plant J* **32,** 713-722.

Bisanz, C., Bastien, O., Grando, D., Jouhet, J., Marechal, E. and Cesbron-Delauw, M.F. (2006). Toxoplasma gondii acyl-lipid metabolism: de novo synthesis from apicoplast-generated fatty acids versus scavenging of host cell precursors. *Biochem J* **394,** 197-205.

Burda, P.C., Schaffner, M., Kaiser, G., Roques, M., Zuber, B. and Heussler, V.T. (2017). A Plasmodium plasma membrane reporter reveals membrane dynamics by live-cell microscopy. *Sci Rep* **7,** 9740.

Bushell, E., Gomes, A.R., Sanderson, T., Anar, B., Girling, G., Herd, C.*, et al.* (2017). Functional Profiling of a Plasmodium Genome Reveals an Abundance of Essential Genes. *Cell* **170,** 260-272 e268.

Costa, G., Gildenhard, M., Eldering, M., Lindquist, R.L., Hauser, A.E., Sauerwein, R.*, et al.* (2018). Non-competitive resource exploitation within mosquito shapes within-host malaria infectivity and virulence. *Nat Commun* **9,** 3474.

Finn, R.D., Tate, J., Mistry, J., Coggill, P.C., Sammut, S.J., Hotz, H.R.*, et al.* (2008). The Pfam protein families database. *Nucleic Acids Res* **36,** D281-288.

Franceschini, A., Szklarczyk, D., Frankild, S., Kuhn, M., Simonovic, M., Roth, A.*, et al.* STRING v9.1: protein-protein interaction networks, with increased coverage and integration. *Nucleic Acids Res* **41,** D808-815.

Guttery, D.S., Ferguson, D.J.P., Poulin, B., Xu, Z., Straschil, U., Klop, O.*, et al.* (2012). A Putative Homologue of CDC20/CDH1 in the Malaria Parasite Is Essential for Male Gamete Development. *PLoS Pathog* **8,** e1002554.

Guttery, David S., Poulin, B., Ramaprasad, A., Wall, Richard J., Ferguson, David J., Brady, D.*, et al.* (2014). Genome-wide Functional Analysis of Plasmodium Protein Phosphatases Reveals Key Regulators of Parasite Development and Differentiation. *Cell Host & Microbe* **16,** 128-140.

Ikeda, M., Kanao, Y., Yamanaka, M., Sakuraba, H., Mizutani, Y., Igarashi, Y. and Kihara, A. (2008). Characterization of four mammalian 3-hydroxyacyl-CoA

dehydratases involved in very long-chain fatty acid synthesis. *FEBS Lett* **582,** 2435-2440.

Janse, C.J., Franke-Fayard, B., Mair, G.R., Ramesar, J., Thiel, C., Engelmann, S*., et al.* (2006). High efficiency transfection of Plasmodium berghei facilitates novel selection procedures. *Molecular and biochemical parasitology* **145,** 60-70.

Kihara, A. (2012). Very long-chain fatty acids: elongation, physiology and related disorders. *J Biochem* **152,** 387-395.

Kohlwein, S.D., Eder, S., Oh, C.S., Martin, C.E., Gable, K., Bacikova, D. and Dunn, T. (2001). Tsc13p is required for fatty acid elongation and localizes to a novel structure at the nuclear-vacuolar interface in Saccharomyces cerevisiae. *Mol Cell Biol* **21,** 109-125.

Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H*., et al.* (2007). Clustal W and Clustal X version 2.0. *Bioinformatics* **23,** 2947-2948.

Lee, S.H., Stephens, J.L., Paul, K.S. and Englund, P.T. (2006). Fatty acid synthesis by elongases in trypanosomes. *Cell* **126,** 691-699.

Letunic, I. and Bork, P. (2019). Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic acids research* **47,** W256-W259.

Li, D., Gonzalez, O., Bachinski, L.L. and Roberts, R. (2000). Human protein tyrosine phosphatase-like gene: expression profile, genomic structure, and mutation analysis in families with ARVD. *Gene* **256,** 237-243.

Li, L., Stoeckert, C.J., Jr. and Roos, D.S. (2003). OrthoMCL: identification of ortholog groups for eukaryotic genomes. *Genome Res* **13,** 2178-2189.

Lin, X., Yang, X., Li, Q., Ma, Y., Cui, S., He, D*., et al.* (2012). Protein tyrosine phosphatase-like A regulates myoblast proliferation and differentiation through MyoG and the cell cycling signaling pathway. *Mol Cell Biol* **32,** 297-308.

Marchler-Bauer, A., Lu, S., Anderson, J.B., Chitsaz, F., Derbyshire, M.K., DeWeese-Scott, C*., et al.* (2011). CDD: a Conserved Domain Database for the functional annotation of proteins. *Nucleic Acids Res* **39,** D225-229.

Mazumdar, J. and Striepen, B. (2007). Make it or take it: fatty acid metabolism of apicomplexan parasites. *Eukaryot Cell* **6,** 1727-1735.

Morineau, C., Gissot, L., Bellec, Y., Hematy, K., Tellier, F., Renne, C*., et al.* (2016). Dual Fatty Acid Elongase Complex Interactions in Arabidopsis. *PloS one* **11,** e0160631.

Organisation, W.H. (2018) World Malaria Report 2018. World Health Organization, pp. 166.

Pandey, R., Mohmmed, A., Pierrot, C., Khalife, J., Malhotra, P. and Gupta, D. (2014). Genome wide in silico analysis of Plasmodium falciparum phosphatome. *BMC Genomics* **15,** 1024.

Pele, M., Tiret, L., Kessler, J.L., Blot, S. and Panthier, J.J. (2005). SINE exonic insertion in the PTPLA gene leads to multiple splicing defects and segregates with the autosomal recessive centronuclear myopathy in dogs. *Hum Mol Genet* **14,** 1417-1427.

Ramakrishnan, S., Docampo, M.D., Macrae, J.I., Pujol, F.M., Brooks, C.F., van Dooren, G.G*., et al.* (2012). Apicoplast and endoplasmic reticulum cooperate in fatty acid

biosynthesis in apicomplexan parasite Toxoplasma gondii. *J Biol Chem* **287,** 4957-4971.

Ramakrishnan, S., Serricchio, M., Striepen, B. and Butikofer, P. (2013). Lipid synthesis in protozoan parasites: a comparison between kinetoplastids and apicomplexans. *Prog Lipid Res* **52,** 488-512.

Roques, M., Wall, R.J., Douglass, A.P., Ramaprasad, A., Ferguson, D.J., Kaindama, M.L.*, et al.* (2015). Plasmodium P-Type Cyclin CYC3 Modulates Endomitotic Growth during Oocyst Development in Mosquitoes. *PLoS Pathog* **11,** e1005273.

Schultz, J., Milpetz, F., Bork, P. and Ponting, C.P. (1998). SMART, a simple modular architecture research tool: identification of signaling domains. *Proc Natl Acad Sci U S A* **95,** 5857-5864.

Shears, M.J., Botte, C.Y. and McFadden, G.I. (2015). Fatty acid metabolism in the Plasmodium apicoplast: Drugs, doubts and knockouts. *Molecular and biochemical parasitology* **199,** 34-50.

Small, I., Peeters, N., Legeai, F. and Lurin, C. (2004). Predotar: A tool for rapidly screening proteomes for N-terminal targeting sequences. *Proteomics* **4,** 1581-1590.

Stanway, R.R., Bushell, E., Chiappino-Pepe, A., Roques, M., Sanderson, T., Franke-Fayard, B.*, et al.* (2019). Genome-Scale Identification of Essential Metabolic Processes for Targeting the Plasmodium Liver Stage. *Cell* **179,** 1112-1128 e1126.

Tamura, K., Stecher, G., Peterson, D., Filipski, A. and Kumar, S. (2013). MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol* **30,** 2725-2729.

Tarun, A.S., Vaughan, A.M. and Kappe, S.H. (2009). Redefining the role of de novo fatty acid synthesis in Plasmodium parasites. *Trends Parasitol* **25,** 545-550.

Tehlivets, O., Scheuringer, K. and Kohlwein, S.D. (2007). Fatty acid synthesis and elongation in yeast. *Biochim Biophys Acta* **1771,** 255-270.

van Schaijk, B.C., Kumar, T.R., Vos, M.W., Richman, A., van Gemert, G.J., Li, T.*, et al.* (2014). Type II fatty acid biosynthesis is essential for Plasmodium falciparum sporozoite development in the midgut of Anopheles mosquitoes. *Eukaryot Cell* **13,** 550-559.

Vaughan, A.M., O'Neill, M.T., Tarun, A.S., Camargo, N., Phuong, T.M., Aly, A.S.*, et al.* (2009). Type II fatty acid synthesis is essential only for malaria parasite late liver stage development. *Cell Microbiol* **11,** 506-520.

Wilkes, J.M. and Doerig, C. (2008). The protein-phosphatome of the human malaria parasite Plasmodium falciparum. *BMC Genomics* **9,** 412.

Yu, L., Pena Castillo, L., Mnaimneh, S., Hughes, T.R. and Brown, G.W. (2006). A survey of essential gene function in the yeast cell division cycle. *Mol Biol Cell* **17,** 4736-4747.

Yu, M., Kumar, T.R., Nkrumah, L.J., Coppi, A., Retzlaff, S., Li, C.D.*, et al.* (2008). The fatty acid biosynthesis enzyme FabI plays a key role in the development of liver-stage malarial parasites. *Cell Host Microbe* **4,** 567-578.

DIC	Hoechst	GFP	P28	Merge	
					Ring
					Trophozoite
					Schizont
					Female Gametocyte
					Male Gametocyte
					Zygote
					Ookinete
					Oocyst
					Sporozoite

10 Days | **14 Days** | **21 Days**