Gut cancer increases the risk for *Drosophila* to be preyed upon by hunting spiders

David Duneau[1,2*] and Nicolas Buchon[3]

[1]Université Toulouse 3 Paul Sabatier, CNRS, IRD; UMR5174 EDB (Laboratoire Évolution &

Diversité Biologique); Toulouse, France.

[2]Instituto Gulbenkian de Ciência, Rua da Quinta Grande 6, P-2780, Oeiras, Portugal

[3]Cornell Institute for Host Microbe Interactions and Disease, Department of Entomology, Cornell

University, Ithaca, NY 14853, USA.

* Corresponding author: David Duneau. david.duneau@gmail.com

Keywords: *Drosophila*, cancer, infection, bacteria, spider, predation

Abstract

Predators are thought to prey on individuals that are in poor physical condition, although the evidence supporting this is ambiguous. We tested if sick individuals where more predated using *Drosophila melanogaster* flies as manipulable preys. We asked whether hunting spiders, trapped from the wild, would selectively prey upon flies with compromised health (i.e. chronically infected or cancerous) versus healthy flies under laboratory conditions. Flies chronically infected with the bacterium *Providencia rettgeri*, a natural *Drosophila* pathogen, were not selectively preyed upon by neither jumping spiders, nor small wolf spiders. We suggest that this result support the hypothesis that chronic infection is associated with reduced symptoms notably to avoid the predation of their host. We then induced colon cancer in some of the flies and asked whether the presence of cancer led to selective predation; there is little evidence for this, even in vertebrates. As the cancer developed, the incidence of predation by jumping spiders on the afflicted flies increased. We conclude that disease can have different lethal consequences through predation, even in invertebrate species, and that cancer is a factor in selective predation. Our results may explain why early tumors, but not metastasized cancers, are commonly detected in organisms in the wild, as cancer bearing individuals are rapidly eliminated due to the strong selective pressure against them.

Introduction

In vertebrates, individuals that are injured, or diseased are often preyed upon, as are juveniles who have not reached adult speed or reflex (Mesa et al. 1994; Møller 2008; Genovart et al. 2010). This is especially true when a predator's preferred type of prey is difficult to catch at the prime of its performance (Temple 1987), however, there are exceptions (Penteriani et al. 2008). Although this conventional wisdom of selective predation is largely accepted, only a few studies have tested it empirically. Selective predation can only be said to exist when the relative frequencies of the types of prey in a predator's diet differ from the frequencies of the prey in the environment (Chesson 1978). Hence, to test the selective prey hypothesis in the wild, data on the types of prey sought by many of the predators and the health status of the whole prey population would be determined. This is obviously difficult to obtain. Furthermore, a prey individual may suffer from different conditions which influence each other and lead to spurious effects. For example, exhaustion from care for large brood increases susceptibility to parasites (Oppliger et al. 1996). Thus, the fittest hosts are the ones classified as being sick. Although those information may be vitally important in the context of eco-evolutionary dynamics (Brunner et al. 2017, 2019), the topic has been subjected more to speculation than to empirical study.

Probably the most studied question regarding selective predation is whether predators prey more on individuals with suboptimal conditions over healthy individuals. Even if hosts predators may avoid infected prey to avoid the risk of getting sick themselves (Goren and Ben-Ami 2017), it has been suggested, based on field observations, that infections could increase the vulnerability to predation (Møller and Erritzøe 2000; Duffy et al. 2005; Murray et al. 2006; Møller and Nielsen 2007; Miller et al. 2008; Møller et al. 2010; Adelman et al. 2017; Gooding et al. 2020) and there is some experimental support for this (Johnson et al. 2006; DeBlieux and Hoverman 2019; Gallagher et al. 2019). Consequently, by mediating selective predation, parasites can mediate the relationship between hosts and their prey (Hall et al. 2005; Møller 2008). For instance, removing predators can reduce vertebrate prey populations (Sih et al. 1985), which seems counterintuitive, as predators are expected to remove preys. However, mathematical models predict that virulent parasites may be selected against if

predation disproportionately removes them before they can be transmitted to the host (Packer et al.

2003). Without this check on virulent parasites by the predator, more prey may die from lethal

infections than from predation. Hence, if selective predation for infected individuals is a common

pattern, it may also be a major driver of host and parasite evolution and cannot be ignored any longer

(Møller 2008).

It is also relevant to consider that not all sickness come from infections and that non-infectious

diseases could also increase the risk of being predated. Hence, if it is accepted that parasites may

evolve in response to the predator of their hosts, one can imagine that non-infectious diseases, when

they have a genetic basis, could also evolve in response to the selective pressure imposed by predators.

To our knowledge, few to nothing is known about the potential selective predation of prey that are sick

due to a non-infectious disease. Mutations can occur during the cell replication and division required

to grow and maintain multicellular animals, and certain types of mutation, said oncogenic mutations,

can lead to the formation of tumors (National Cancer Institute 2020*a*). In most cases, tumors are

benign (Bissell and Hines 2011), but they can be malignant, and even fatal (National Cancer Institute

2020*b*). Clinical categorization of tumors has been established in unchallenging, even 'optimal'

environments (i.e. in the laboratory or clinic), usually to the point of death via organ failure. The

impact of tumor progression on animal survival under more natural conditions has largely been

neglected. However, depending on the relationship between tumor development and the health of the

prey, predation could affect the selection of oncogenic mutations and thus the risk to develop cancer.

To our knowledge, the possible interactions between a predator and a prey that is afflicted with cancer

are poorly understood and have not been studied to date (Vittecoq et al. 2013).

Here, we induced chronic bacterial infection and colon cancer in prey to understand their role

in increasing the likelihood of being predated. We use the fruit fly *Drosophila melanogaster,* already a

well-characterized model for infectious and non-infectious disease, as a novel model for sick prey. We

assess the predation of flies by hunting spiders, using flies chronically infected by a wild-caught

bacterial pathogen (*Providencia rettgeri*), and have cancer, under laboratory-controlled conditions.

These conditions allow for the ideal experimental setting to study predation and the role of prey well-

being in a population context.

Materials and methods

Prey and predators.

We used *Drosophila melanogaster* flies as prey in this study as they are tractable to genetic induction

of tumours, are well-defined models for bacterial infection, and allow for control of environmental and

genetic differences. Only one sex (males) were used, to avoid sex effects. We induced chronic

infection by injecting 23 nL of a suspension of few thousand cells of the bacterium *Providencia*

rettgeri (strain *Dmel*, isolated from wild-caught *Drosophila melanogaster;* Juneja and Lazzaro 2009)

in phosphate-buffered saline (PBS) into the abdomen of flies (Canton S line). The same volume of

sterile PBS without bacteria was injected into another group of flies that served as the control.

Individuals surviving for three days or more after injection of the bacteria have an established chronic

infection, as we have previously demonstrated (Duneau et al. 2017).

To induce tumors in the flies, we took advantage of signaling pathways that regulate cell growth in

mammals, which have a conserved function in *Drosophila*, the EGFR and Wnt pathways (Rudrapatna

et al. 2012; Millburn et al. 2016; Mirzoyan et al. 2019; Villegas 2019). We used targeted induction of

these pathways (Rasv12 for EGFR, APC RNAi for the Wnt pathway) to induce two degrees of

malignancy in the digestive tract of flies. This type of cancer is appropriate for this study, as it does

not overtly affect physical performance including locomotion, and will thus allow for the study of the

interaction of oncogenic phenomena and predation (see Dawson et al. 2018 for locomotion assays in

gut cancerous flies in another genetic background). To generate tumors in the gut of *Drosophila*, flies

esg^{TS} (*esgGal4*, *Gal80^{ts}*, *UAS-GFP*) were crossed to either background control flies (Trip stock

35786), flies carrying a *UAS-Ras^{v12}* (Bloomington 64196) or flies carrying a combination of *UAS-*

Ras^{v12} (Bloomington 641926) and *UAS-APC-RNAi* (Bloomington Trip stock 28582) combined. The

expression of Ras^{v12} in progenitor cells stimulates an accelerated turnover and a mild accumulation of

epithelial cells in the gut, akin to a dysplastic tissue (Buchon et al. 2010; Houtz et al. 2019). The

overexpression of Ras^{v12} combined with the knock-down of *APC* leads to increased proliferation, and

accumulation of progenitor cells and epithelial cells in the gut, reminiscent of an intestinal

disseminated tumor (Wang et al. 2013). F1 flies from these crosses were raised at 18°C (Gal4 off,

transgene expression off, normal development and emergence), and 3 days after emergence were

switched to 29°C (Gal4 on, transgene expression on) for 10 or 20 days before performing the

experiment. As in (Wang et al. 2013), crosses with WT flies showed mild tissue renewal (restricted

GFP signal), flies with expressing Ras^{v12} showed disseminated GFP in their gut and flies expressing

Ras^{v12} and *APC RNAi* showed gut enlargement and GFP accumulation.

We used two families of hunting spiders as predators: jumping spiders of several species from

the family of Salticidae (see illustration in Fig. 1) and small wolf spiders from the family of Lycosidae

(likely *Pardosa lugubris*). Both jumping and wolf spiders hunt by wandering rather than by using

webs as traps.

Jumping spiders have the sharpest vision of any arthropod, even surpassing many vertebrates

(Land and Nilsson 2012). They hunt during the day and rely on this astonishing vision when catching

a prey. Similar to the way a cat stalks its prey, jumping spiders turn towards their prey, directed by a

pair of lateral eyes that provide a nearly panoramic field of view with the ability to discern motion

(Zurek and Nelson 2012). Then, the spiders track, approach, and jump on the prey. The vision

accuracy is conferred by two pairs of forward-facing eyes that also provide a precise perception of

depth (Zurek et al. 2010; Nagata et al. 2012).

In absence of detectable effect on selective predation of infected preys, we decided to test our

hypothesis with another type of hunting spider species with different hunting behavior. Wolf spiders

are nocturnal hunters and their eyes function mainly as low-light movement detectors. Their vision is

not as acute as that of jumping spiders (Land and Nilsson 2012). The use of visual cues in prey

detection and orientation have only been well-studied in jumping spiders, and empirical evidence

suggests that wolf spiders rely more on vibrations to capture their prey than on vision (Lizotte and

Rovner 1988). Hence, the differences between these two types of hunting spiders primarily lie in their

ability to prey on flies that are at rest. In an environment where there is no place to hide, jumping

spiders would likely prey equally well on moving and non-moving prey, and selective predation in this

case should depend largely on the prey's capacity to escape.

All spiders in the study were obtained from the wild in the Cornell University campus (Ithaca,

NY, USA) or in the campus of the University of Toulouse 3 (Toulouse, France).

Predation trials

Predation trials were performed by incubating five healthy and five sick flies at 20°C with a spider in a

round plastic box (11.5cm x 8.5cm) containing a wet piece of cotton or foam. The flies were first

added to the box to settle before the introduction of the predators. After about 30 min, spiders were put

in the box. The trials generally lasted for about 4 hours or were stopped when 50% of the flies had

been preyed upon. This goal was sometimes exceeded as some spiders ate faster than expected.

Individual spiders have never been used twice as predators within an experiment.

To determine the number of surviving flies that were infected, we ground the remaining flies

at the end of the predation period and suspended the mixture in 250 µL of Luria-Bertani (LB) medium.

A droplet of five microliters of the resulting suspension was spread on a petri dish filled with agar

containing LB medium, and the plates were incubated overnight at 37°C to assess for the presence of

P. rettgeri. We calculated the proportion of infected flies at the end of predation period by determining

the number of droplets in which *P. rettgeri* were present. For a control comparison, we performed the

same process with infected and healthy flies, but did not expose them to predatory spiders. The

number of surviving flies that had cancer were counted by dissecting all surviving flies and observing

their guts under a fluorescent microscope, which reveals the presence/absence of cancer.

Statistical analysis

The use of the Fisher exact test or a chi-squared test would have permitted us to test for every trial

regardless of whether the number of flies from a particular category was more or less preyed upon than

those from another category. However, we used a hypergeometric distribution instead of a Chi-

squared distribution because, as flies were preyed upon, the population size decreased. The *p*-value for

each experimental trial was added up and compared to the summed *p*-value obtained by simulating

random trials. We assessed the overall *p*-value by determining the number of simulated trials required

to have an observed result (summed *p*-value observed) as probable as the simulation (summed *p*-value

simulated). Hence, a *p*-value of 0.05 indicated that 20 trials would be sufficient to have the same result

as the empirical data by chance, whereas a *p*-value of 0.0001 meant that 10,000 trials would be

required to achieve the same result as the empirical data by chance. We used the Manly Alpha Index

as the preference index to illustrate the selective predation on sick individuals (index 0: preference for

healthy individuals, index 0.5: no preference, index 1: preference for sick individuals). It was

calculated as log (total flies sick / flies sick but not eaten) / [(log(total flies sick / flies sick but not

eaten) + log(total flies healthy / flies healthy but not eaten)] as established by Manly (1972).

Results

Predation of chronically infected flies by hunting spiders. We tested whether chronic infection of

the flies by *P. rettgeri* affected the risk of being preyed upon by the spiders. The flies had been

infected for several days before exposure to the spiders, and their immune response was expected to be

activated (Chambers et al. 2019). Despite using two predator species, we did not detect selective

predation by either of the hunting spider types (Fig. 2A and B for jumping and wolf spiders,

respectively).

Predation of flies with cancer by hunting spiders. We asked if cancer of the gut, which we induced,

affected the risk for predation of flies by the hunting spiders. Hyperplasia is the enlargement of an

organ or tissue caused by increased cell division and is often an initial stage in the development of

cancer. We observed that flies in this early hyperplastic stage ($esg^{ts} > UAS\text{-}Ras^{v12}$) did not show an

increased likelihood of getting preyed upon by the jumping spiders, which are active diurnal hunters

(Fig. 3A). However, jumping spiders did catch more successfully flies with an advanced stage of

cancer ($esg^{ts} > UAS\text{-}Ras^{v12}, UAS\text{-}APC\text{-}IR$) (compared with healthy flies, and the preference seemed to

increase with the length of time after the tumor had been induced (Fig. 3B).

Discussion

Many diseases do not kill their hosts and are considered benign in optimal conditions, such as in a clinic or laboratory. However, it is very likely that non-lethal illnesses still increase the risk of mortality through indirect interactions with other environmental factors. Indeed, being sick may reduce foraging and mating abilities, increase the risk for lethal superinfections, or of being preyed upon. Even though many biological questions are presently being addressed using an integrative approach, the question of whether a non-lethal disease is truly benign in natural conditions seems to have been overlooked in many cases. We hope to generate interest in evaluating the lethality of diseases when their interaction with other risks, such as predation, is considered. Chronic infection by a wild-caught *Drosophila* pathogen did not significantly affect the risk of being preyed upon in the laboratory. In contrast, we observed that while the presence of early stage tumors did not change predation risk, the presence of advanced cancer did compared to age-matched controls.

Little is known about selective predation in individuals with non-infectious diseases. The prevailing opinion that general health, which can be affected by non-infectious sickness, correlates with the risk of being preyed upon, is largely observational (Hoey and McCormick 2004; Genovart et al. 2010). Nonetheless, although we should not be jumping to hasty conclusions, our results on the effects of non-infectious disease on selective predation suggests a stronger selective pressure than natural chronic infections.

Cancers are common in multicellular organisms and can occur whenever cells evade the normal cell checkpoints that control cell division, proliferation, and apoptosis, thus leading to uncontrolled cell division (Aktipis et al. 2015). Individuals always have some tumors, most of which never become lethal (Bissell and Hines 2011; Abu-Helil and van der Weyden 2019). Our understanding of oncogenic phenomena in wild populations of animals is limited, in particular whether tumors that are clinically non-lethal affect the well-being of organisms in the wild (McAloose and Newton 2009; Vittecoq et al. 2013; Pesavento et al. 2018). If tumor formation is inevitable, it is likely that multicellular organisms have evolved ways to control and tolerate them, perhaps resulting in a

trade-off between surviving the tumor and surviving other perils, such as infections or predation

(Pavard and Metcalf 2019). In some cases, selection for tumor tolerance may have arisen from

selection for tolerance to other factors, perhaps explaining why early stages of cancer, such as the

colon hyperplasia that we induced in *Drosophila*, do not change the likelihood of being preyed upon.

In our study, when the tumor developed into a cancer, predation of the flies increased, suggesting that

uncontrolled tumors strongly increase the likelihood of being preyed upon. This conclusion supports

the idea that individuals with uncontrolled tumors will suffer the risk of increased predation compared

with healthy individuals. Predation is a critical force shaping natural selection (Wade and Kalisz

1990). By removing prey genotypes susceptible to tumors, predators are likely to have selected for the

tumor tolerance largely observed in nature (Bissell and Hines 2011). Furthermore, by removing

individuals with advanced tumors, predation may explain why cancerous animals are rarely recorded

in autopsies of wild animal populations (Vittecoq et al. 2013).

The progression of cancer can, in many ways, be compared with infectious diseases. However,

if both reduce viability and survival, infectious diseases are, unlike oncogenic processes, under

selective pressure to ensure transmission of the pathogen to another host. For both non-infectious

diseases and cancer, selective predation could increase the selection against the sickness, as diseases

that are moderate and non-lethal could lead to a higher risk of the host being preyed upon, resulting in

mortality rates that are similar to those of infectious diseases. With infectious diseases, there may be

additional consequences. First, selective predation could affect the transmission of pathogens,

spreading them through predator feces over long distances. Second, it might affect epidemics, either

preventing them by removing parasite spreaders from the host population (Duffy et al. 2005) or by

increasing their likelihood of occurrence by dispersing parasites instead of containing them in dead

hosts (Strauss et al. 2016). An additional consideration is that predation may affect the evolution of

parasite virulence. It is generally assumed that predation can reduce virulence because hosts with

rapidly proliferating parasites would suffer more from the infection and may be eaten before

transmitting the parasite to another host, whereas the less rapidly proliferating parasite would impact

less host health and would be transmitted before the host is eaten (Møller et al. 2010). However, lower

virulence could evolve also through lower pathogenicity, that is by inducing less damage to the host,

and not only through reduced proliferation. The most important factor being that bacteria can

proliferate within their host as long as the symptoms do not increase the risk of predation. The bacteria

studied here, *P. rettgeri*, was obtained from the wild and is prevalent in populations of *D.*

melanogaster (Juneja and Lazzaro 2009). Despite being chronic and the fact that the pathogen can

trigger an immune response through the course of the infection (Duneau et al. 2017; Chambers et al.

2019), the chronic infection did not increase the predation in our study. This result supports the

hypothesis that chronic infections are selected for such that the symptoms that enhance predation are

reduced.

It is reasonable to assume that the success of predators is not simply due to favorable luck. In

fact, selective predation may be more the rule than the exception, although it is not understood what

makes a prey susceptible. The value of an integrative approach is recognized by most biologists and

dictates that we attempt to understand the role of predation in the evolution of diseases, infectious or

not. On the one hand, selective predation on sick individuals is likely to counter-select against

parasites or genetic diseases more strongly than if selection occurred upon disease-mediated host

death. On the other hand, in the same way that some parasites evolved to manipulate their host to

increase transmission to intermediate hosts, parasites may have evolved to make their hosts less

conspicuous to predation, for example, by interfering with the search for mates or by lessening the

effects of the disease on the host; for example, by reducing symptoms. Because cancer progression

generally results in the death of the organism, we cannot expect the incidence of lethal cancer to have

evolved in the same way as parasites. We argue that selective predation on prey with cancer may be

one of the reasons why tumors are commonly tolerated in animals, where advanced cancer is only

rarely detected.

Acknowledgements

262 We thank Violette Chiara, Jean-Baptiste Ferdy, Raphaël Jeanson, and Daniel Zurek for thoughtful

263 discussion; Pierrick Blanchard, Dieter Ebert, Simon Fellous, Sabine Noebel, and Jennifer Regan for

their comments on the manuscript; Cole Gilbert for providing the jumping spiders used in the cancer trials; and Christian Faucher for laboratory support. DD was supported by the French Laboratory of Excellence project 'TULIP' (ANR-10-LABX-41; ANR-11-IDEX-0002–02) and the LIA BEEG-B (Laboratoire International Associé-Bioinformatics, Ecology, Evolution, Genomics and Behaviour) (CNRS).

References

Abu-Helil, B., and L. van der Weyden. 2019. Metastasis in the wild: investigating metastasis in non-laboratory animals. Clinical and Experimental Metastasis 36:15–28.

Adelman, J. S., C. Mayer, and D. M. Hawley. 2017. Infection reduces anti-predator behaviors in house finches. Journal of Avian Biology 48:519–528.

Aktipis, C. A., A. M. Boddy, G. Jansen, U. Hibner, M. E. Hochberg, C. C. Maley, and G. S. Wilkinson. 2015. Cancer across the tree of life: Cooperation and cheating in multicellularity. Philosophical Transactions of the Royal Society B: Biological Sciences 370.

Bissell, M. J., and W. C. Hines. 2011. Why don't we get more cancer? A proposed role of the microenvironment in restraining cancer progression. Nature Medicine 17:320–329.

Brunner, F. S., J. M. Anaya-Rojas, B. Matthews, and C. Eizaguirre. 2017. Experimental evidence that parasites drive eco-evolutionary feedbacks. Proceedings of the National Academy of Sciences 114:3678–3683.

Brunner, F. S., J. A. Deere, M. Egas, C. Eizaguirre, and J. A. M. Raeymaekers. 2019. The diversity of eco☐evolutionary dynamics: Comparing the feedbacks between ecology and evolution across scales. Functional Ecology 33:7–12.

Buchon, N., N. A. Broderick, T. Kuraishi, and B. Lemaitre. 2010. Drosophila EGFR pathway coordinates stem cell proliferation and gut remodeling following infection. BMC Biology 8.

Chambers, M. C., E. Jacobson, S. Khalil, and B. P. Lazzaro. 2019. Consequences of chronic bacterial

infection in *Drosophila melanogaster*. PloS one 14:e0224440.

Chesson, J. 1978. Measuring preference in selective predation. Ecology 59:211–215.

Dawson, E. H., T. P. M. Bailly, J. Dos Santos, C. Moreno, M. Devilliers, B. Maroni, C. Sueur, et al.

2018. Social environment mediates cancer progression in *Drosophila*. Nature Communications

9:3574.

DeBlieux, T., and J. Hoverman. 2019. Parasite-induced vulnerability to predation in larval anurans.

Diseases of Aquatic Organisms 135:241–250.

Duffy, M. A., S. R. Hall, A. J. Tessier, and M. Huebner. 2005. Selective predators and their parasitized

prey: Are epidemics in zooplankton under top-down control? Limnology and Oceanography 50:412–

420.

Duneau, D., J.-B. Ferdy, J. Revah, H. C. Kondolf, G. A. Ortiz, B. P. Lazzaro, and N. Buchon. 2017.

Stochastic variation in the initial phase of bacterial infection predicts the probability of survival in *D.*

melanogaster. eLife 6:e28298.

Gallagher, S. J., B. J. Tornabene, T. S. DeBlieux, K. M. Pochini, M. F. Chislock, Z. A. Compton, L.

K. Eiler, et al. 2019. Healthy but smaller herds: Predators reduce pathogen transmission in an

amphibian assemblage. Journal of Animal Ecology 88:1613–1624.

Genovart, M., N. Negre, G. Tavecchia, A. Bistuer, L. Parpal, and D. Oro. 2010. The young, the weak

and the sick: Evidence of natural selection by predation. PLoS ONE 5:9774.

Gooding, E. L., M. R. Kendrick, J. F. Brunson, P. R. Kingsley-Smith, A. E. Fowler, M. E. Frischer,

and J. E. Byers. 2020. Black gill increases the susceptibility of white shrimp, *Penaeus setiferus*

(Linnaeus, 1767), to common estuarine predators. Journal of Experimental Marine Biology and

Ecology 524:151284.

Goren, L., and F. Ben-Ami. 2017. To eat or not to eat infected food: a bug's dilemma. Hydrobiologia

798:25–32.

Hall, S. R., C. E. Cáceres, M. A. Duffy, and C. E. Cáceres. 2005. Selective predation and productivity

jointly drive complex behavior in host-parasite systems. American Naturalist 165:70–81.

Hoey, A. S., and M. I. McCormick. 2004. Selective predation for low body condition at the larval-

juvenile transition of a coral reef fish. Oecologia 139:23–29.

Houtz, P., A. Bonfini, X. Bing, and N. Buchon. 2019. Recruitment of Adult Precursor Cells Underlies

Limited Repair of the Infected Larval Midgut in Drosophila. Cell Host and Microbe 26:412-425.e5.

Johnson, P. T. J., D. E. Stanton, E. R. Preu, K. J. Forshay, and S. R. Carpenter. 2006. Dining on

disease: How interactions between infection and environment affect predation risk. Ecology 87:1973–

1980.

Juneja, P., and B. P. Lazzaro. 2009. *Providencia sneebia sp. nov.* and *Providencia burhodogranariea*

sp. nov., isolated from wild *Drosophila melanogaster*. International Journal of Systematic and

Evolutionary Microbiology 59:1108–1111.

Land, M. F., and D.-E. Nilsson. 2012. Animal eyes. (Oxford University Press, ed.) (Oxford Ani.).

Oxford.

Lizotte, R. S., and J. S. Rovner. 1988. Nocturnal capture of fireflies by lycosid spiders: visual versus

vibratory stimuli. Animal Behaviour 36:1809–1815.

Manly, B. F. J. 1972. Tables for the analysis of selective predation experiments. Researches on

Population Ecology 14:74–81.

McAloose, D., and A. L. Newton. 2009. Wildlife cancer: A conservation perspective. Nature Reviews

Cancer 9:517–526.

Mesa, M. G., T. P. Poe, D. M. Gadomski, and J. H. Petersen. 1994. Are all prey created equal? A

review and synthesis of differential predation on prey in substandard condition. Journal of Fish

Biology 45:81–96.

Millburn, G. H., M. A. Crosby, L. S. Gramates, S. Tweedie, W. Gelbart, N. Perrimon, C. Extavour, et

al. 2016. Fly Base portals to human disease research using *Drosophila* models. DMM Disease Models

and Mechanisms 9:245–252.

Miller, M. W., H. M. Swanson, L. L. Wolfe, F. G. Quartarone, S. L. Huwer, C. H. Southwick, and P.

M. Lukacs. 2008. Lions and prions and deer demise. PLoS ONE 3.

Mirzoyan, Z., M. Sollazzo, M. Allocca, A. M. Valenza, D. Grifoni, and P. Bellosta. 2019. *Drosophila*

melanogaster: A model organism to study cancer. Frontiers in Genetics 10:1–16.

Møller, A. P. 2008. Interactions between interactions: Predator-prey, parasite-host, and mutualistic

interactions. Annals of the New York Academy of Sciences 1133:180–186.

Møller, A. P., and J. Erritzøe. 2000. Predation against birds with low immunocompetence. Oecologia

122:500–504.

Møller, A. P., J. Erritzøe, and J. Tøttrup Nielsen. 2010. Predators and microorganisms of prey:

goshawks prefer prey with small uropygial glands. Functional Ecology 24:608–613.

Møller, A. P., and J. T. Nielsen. 2007. Malaria and risk of predation: A comparative study of birds.

Ecology 88:871–881.

Murray, D. L., J. R. Cary, and L. B. Keith. 2006. Interactive effects of sublethal nematodes and

nutritional status on snowshoe hare vulnerability to predation. The Journal of Animal Ecology 66:250.

Nagata, T., M. Koyanagi, H. Tsukamoto, S. Saeki, K. Isono, Y. Shichida, F. Tokunaga, et al. 2012.

Depth perception from image defocus in a jumping spider. Science 335:469–471.

National Cancer Institute. 2020*a*. NCI Dictionary of Cancer Terms.

———. 2020*b*. NCI Dictionary of Cancer Terms.

Oppliger, A., P. Christe, and H. Richner. 1996. Clutch size and malaria resistance. Nature 381:565–

565.

Packer, C., R. D. Holt, P. J. Hudson, K. D. Lafferty, and A. P. Dobson. 2003. Keeping the herds

healthy and alert: Implications of predator control for infectious disease. Ecology Letters 6:797–802.

Pavard, S., and C. Metcalf. 2019. Trade-offs between mortality components in life history evolution:

the case of cancers. Human Evolutionary Demography 1–26.

Penteriani, V., M. Del Mar Delgado, P. Bartolommei, C. Maggio, C. Alonso-Alvarez, and G. J.

Holloway. 2008. Owls and rabbits: Predation against substandard individuals of an easy prey. Journal

of Avian Biology 39:215–221.

Pesavento, P. A., D. Agnew, M. K. Keel, and K. D. Woolard. 2018. Cancer in wildlife: patterns of

emergence. Nature Reviews Cancer 18:646–661.

Rudrapatna, V. A., R. L. Cagan, and T. K. Das. 2012. *Drosophila* cancer models. Developmental

Dynamics 241:107–118.

Sih, A., P. Crowley, M. McPeek, J. Petranka, and K. Strohmeier. 1985. Predation, competition, and

prey communities: A review of field experiments. Annual Review of Ecology and Systematics

16:269–311.

Strauss, A. T., M. S. Shocket, D. J. Civitello, J. L. Hite, R. M. Penczykowski, M. A. Duffy, C. E.

Cáceres, et al. 2016. Habitat, predators, and hosts regulate disease in *Daphnia* through direct and

indirect pathways. Ecological Monographs 86:393–411.

Temple, S. A. 1987. Do predators always capture substandard individuals disproportionately from

prey populations. Ecology 68:669–674.

Villegas, S. N. 2019. One hundred years of *Drosophila* cancer research: No longer in solitude. DMM

Disease Models and Mechanisms 12.

Vittecoq, M., B. Roche, S. P. Daoust, H. Ducasse, D. Missé, J. Abadie, S. Labrut, et al. 2013. Cancer:

A missing link in ecosystem functioning? Trends in Ecology and Evolution. Elsevier Current Trends.

Wade, M. J., and S. Kalisz. 1990. The causes of natural selection. Evolution 44:1947–1955.

Wang, C., R. Zhao, P. Huang, F. Yang, Z. Quan, N. Xu, and R. Xi. 2013. APC loss-induced intestinal

tumorigenesis in *Drosophila*: Roles of Ras in Wnt signaling activation and tumor progression.

Developmental Biology 378:122–140.

Zurek, D. B., and X. J. Nelson. 2012. Hyperacute motion detection by the lateral eyes of jumping

spiders. Vision Research 66:26–30.

Zurek, D. B., A. J. Taylor, C. S. Evans, and X. J. Nelson. 2010. The role of the anterior lateral eyes in

the vision-based behaviour of jumping spiders. Journal of Experimental Biology 213:2372–2378.

Figure legends

Figure 1. Photograph of a jumping spider consuming *Drosophila melanogaster* flies in the laboratory.

Figure 2. Predation by hunting spiders on a population of *Drosophila melanogaster* Canton S, of either healthy flies or flies chronically infected with *P. rettgeri*. A. Predation by jumping spiders (Hypergeometric test: n= 27 trials, *p*-value > 0.05, Median pref. index = 0.64) B. Predation by wolf spiders (Hypergeometric test: n= 30 trials, *p*-value > 0.05, Median pref. index = 0.43). Each dot represents the proportion of individuals eaten over five flies and the lines connect the groups of individuals from a same trial. Each trial represents ten flies (five from each type) presented to one spider at once. Preference index is calculated as in Manly (1972) and ranges from 0 to 1, from a preference to healthy flies to a preference for sick flies, respectively.

Figure 3. Predation by hunting spiders in a population of *Drosophila melanogaster* that were either healthy or in which colon tumours (hyperplasia) had been induced. Tumour size increased with induction time, during which uncontrolled cell division occurred. A. Uncontrolled cell division was triggered for 10 or 20 days in $esg^{TS} > Ras^{v12}$ flies to lead to hyperplasia (respectively with n= 17 and 15 trials). In both cases, individuals with hyperplasia were not differently predated upon compared with the healthy individuals (Hypergeometric test *p*-value: 10 days > 0.05; 20 days > 0.05, Median pref. index = 0.5). B. Fast uncontrolled cell division was triggered for 10 or 20 days in $esg^{TS} > Ras^{v12}$, *APC-RNAi* flies to lead to cancer (respectively with n= 16 and 20 trials). In both cases, individuals with cancer were more likely to be predated upon than individuals who did not have cancer (Hypergeometric test *p*-value: 10 days = 0.025; 20 days = 0.002). This pattern tended to be stronger as the tumour grew larger (Median pref. index: 10 days = 0.64; 20 days = 0.8). Each dot represents the proportion of individuals eaten over five flies and the lines connect the groups of individuals from a same trial. Each trial represents ten flies (five from each type) presented to one spider at once. Preference index is calculated as in Manly (1972) and ranges from 0 to 1, from a preference to healthy flies to a preference for sick flies, respectively.

Figure 1: Photograph of a jumping spider consuming Drosophila melanogaster flies in the laboratory.

Figure 2: Predation by hunting spiders on a population of *Drosophila melanogaster* Canton S, of either healthy flies or flies chronically infected with *P. rettgeri*. A. Predation by jumping spiders (Hypergeometric test: n= 27 trials, p-value > 0.05, Median pref. index = 0.64) B. Predation by wolf spiders (Hypergeometric test: n= 30 trials, p-value > 0.05, Median pref. index = 0.43). Each dot represents the proportion of individuals eaten over five flies and the lines connect the groups of individuals from a same trial. Each trial represents ten flies (five from each type) presented to one spider at once. Preference index is calculated as in Manly (1972) and ranges from 0 to 1, from a preference to healthy flies to a preference for sick flies, respectively.

Figure 3: Predation by hunting spiders in a population of *Drosophila melanogaster* that were either healthy or in which colon tumours (hyperplasia) had been induced. Tumour size increased with induction time, during which uncontrolled cell division occurred. A. Uncontrolled cell division was triggered for 10 or 20 days in esgTS > Rasv12 flies to lead to hyperplasia (respectively with n= 17 and 15 trials). In both cases, individuals with hyperplasia were not differently predated upon compared with the healthy individuals (Hypergeometric test p-value: 10 days > 0.05; 20 days > 0.05, Median pref. index = 0.5). B. Fast uncontrolled cell division was triggered for 10 or 20 days in esgTS > Rasv12, APC-RNAi flies to lead to cancer (respectively with n= 16 and 20 trials). In both cases, individuals with cancer were more likely to be predated upon than individuals who did not have cancer (Hypergeometric test p-value: 10 days = 0.025; 20 days = 0.002). This pattern tended to be stronger as the tumour grew larger (Median pref. index: 10 days = 0.64; 20 days = 0.8). Each dot represents the proportion of individuals eaten over five flies and the lines connect the groups of individuals from a same trial. Each trial represents ten flies (five from each type) presented to one spider at once. Preference index is calculated as in Manly (1972) and ranges from 0 to 1, from a preference to healthy flies to a preference for sick flies, respectively.