DNA methylation analysis of amplicons from individuals exposed to maternal

tobacco use during pregnancy, and offspring conduct problems in childhood and

adolescence

Alexandra J. Noble[1], John F. Pearson[2], Alasdair D. Noble[3] Joseph M. Boden[4], L.
John Horwood[4], Martin A. Kennedy[2], Amy J. Osborne[1]

[1]School of Biological Sciences, University of Canterbury, Christchurch, New Zealand

[2]Department of Pathology and Biomedical Sciences, University of Otago,
Christchurch, New Zealand

[3]AgResearch, Lincoln Research Centre, Christchurch, New Zealand

[4]Department of Psychological Medicine, University of Otago, Christchurch, New
Zealand

Abstract

Maternal tobacco smoking during pregnancy is a large driver of health inequalities and a higher prevalence of conduct problem has been observed in exposed offspring. Further, maternal tobacco use during pregnancy can also alter offspring DNA methylation. However, currently, limited molecular evidence have been found to support this observation. Thus we aim to examine the association between maternal tobacco use in pregnancy and whether offspring Conduct problems is mediated by tobacco exposure-induced via DNA methylation differences. Understanding the etiology of the causal link will be crucial in the early identification and treatment of CP in children and adolescents. DNA was sourced from the Christchurch Health and Development Study, a longitudinal birth cohort studied for over 40 years in New Zealand. Bisulfite-based amplicon sequencing of 10 loci known to play a role in neurodevelopment, or with associations with CP phenotypes, was undertaken. We identified nominally significant differential DNA methylation at specific CpG sites in *CYP1A1*, *ASH2L* and *MEF2C* in individuals with Conduct problems who were exposed to tobacco *in utero*. We conclude that environmentally-induced DNA methylation differences could play a role in the observed link between maternal tobacco use during pregnancy and childhood/adolescent Conduct problems However, larger sample sizes are needed to produce an adequate amount of power to investigate this interaction further.

Introduction

Mothers who smoked tobacco during pregnancy have a higher prevalence of offspring developing a conduct disorder phenotype compared to mothers who did not smoke (Wakschlag et al. 1997). This association has been proven in several different cohort studies and the observation have remained following adjustment for various other confounding factors, for example, socio economic status, maternal age, substance abuse, parental anti-social personality, and maladaptive parenting (Wakschlag et al. 1997; Joelsson et al. 2016). However, there is limited molecular evidence to suggest a link between *in utero* tobacco exposure and offspring conduct disorder, thus a direct link between *in utero* tobacco exposure and Conduct problems (CP) remains elusive. Previously we conducted a pilot study assessing differential DNA methylation in a small cohort of individuals who were exposed to tobacco *in uter*o, with sub-groups of individuals defined as having high conduct disorder scores (Noble et al. 2021). We found nominally significant DNA methylation changes with this interaction in several genes associated with neurodevelopment (cite epic in utero paper). Due to the limitations of using a small sample size combined with an array containing a large number of loci, results were underpowered, therefore observations were unable to reach genome wide significance. However, the biological relevance of these nominally significant CpG loci to the CP phenotype, combined with further research which has suggested an epigenetic link between *in utero* tobacco exposure and ADHD (Sengupta et al. 2017), implies that the link between DNA methylation and CP development in tobacco-exposed offspring should be investigated more fully.

Here, we will further pursue this hypothesis, by exploring differential methylation in genes that have known roles during *in utero* neurodevelopment and CP phenotypes, to understand whether DNA methylation may help explain the relationship between *in utero* tobacco exposure and development of CP in offspring. We applied a targeted approach via bisulfite-based amplicon sequencing (BSAS) of regions of genes involved in neurodevelopment. Amplicon sequencing has the ability to interrogate a region of the genome, therefore specifically targeting consecutive CpG sites in a row. We then assessed differential methylation in the DNA of participants from the Christchurch Health and Development Study (CHDS) whose mothers consumed tobacco during pregnancy, with high and low CP scores, and compared

this to controls who were not exposed. This approach allows us to specifically ask whether DNA methylation at genes involved in neurodevelopment and CP phenotypes are specifically differentially methylated in the DNA of offspring with CP, who were exposed to tobacco *in utero*. A significant interaction here would provide further support for a role for DNA methylation in the link between *in utero* exposure and CP development, something which has so far proved elusive.

Methods

Sample

A sub-group of individuals from the CHDS were selected for this study (Table 1). This longitudinal study originally included 97% of all the children (N= 1265) born in the Christchurch, New Zealand urban region during a period in mid-1977 and has been studied at 24 time points from birth to age 40 (N= 987 at age 30). All participants were aged between 28-30 when blood samples and DNA was extracted.

For the subsets studied in this report, CHDS participants were chosen based on their *in utero* tobacco exposure status, their adult smoking status, and their CP scores. Group 1 consisted of individuals who were exposed *in utero* to tobacco smoke, and never smokers at the time blood samples were taken (N= 32). Group 2 consisted of individuals who were exposed *in utero* to tobacco smoke and were themselves regular smokers at the time the blood was taken (N= 32). Group 3 consisted of individuals who were not exposed to tobacco *in utero*, and never smokers at the time blood was taken (N= 32). *In utero* tobacco exposure was defined as 10+ cigarettes per day throughout pregnancy. Within each group of 32, 16 individuals were selected with a 'high' score on a measure of childhood CP at age 7-9 years and 16 with a 'low' score. Severity of childhood CP was assessed using an instrument that combined selected items from the Rutter and Conners child behaviour checklists (Rutter M 1970; Conners 1970, 1969; Fergusson, Horwood, and Lloyd 1991) as completed by parents and teachers at annual intervals from 7-9 years. Parental and teacher reports were summed and averaged over the three years (Fergusson, Horwood, and Ridder 2005) to derive a robust scale measure of the extent to which the child exhibited conduct disordered/oppositional behaviours (mean (SD)=

50.1(7.9) ; range 41-97). For the purposes of this report a 'high' score was defined

as falling into the top quartile of the score distribution (scores >53) and a 'low' score

was defined as scores < 46.

Bisulfite-based amplicon sequencing

Bisulfite-based amplicon sequencing (BSAS) was carried out as described (Noble et

al. 2020). DNA was extracted from whole blood samples using the Kingfisher Flex

System (Thermo Scientific, Waltham, MA USA). DNA was quantified via nanodrop

(Thermo Scientific, Waltham, MA USA). Bisulfite treatment was carried out using the

EZ DNA Methylation-Gold kit (Zymo Research, Irvine, CA, USA) as per the

manufacturer's instructions. DNA samples were then diluted to a final concentration

of 100 ng/µl.

Amplicons for sequencing (Table 2 and Supplementary Table 1) were picked based

upon several criteria: i) previously published differential DNA methylation in response

to *in utero* tobacco smoking; ii) known associations with *in utero* brain development,

and; iii) known associations with CP phenotypes. Primers were then designed to

flank the CpG sites of interest, ~350 base pairs (bp) in total, or to amplify the

promoter region of the gene if a specific CpG site was not known. Multiple pairs of

primers were designed to amplify larger regions.

Bisulfite-converted DNA was amplified via PCR, using KAPA Taq HotStart DNA

Polymerase (Sigma, Aldrich) under the following conditions: 95 °C for 10 min, 95 °C

for 30 sec, 59 °C for 20 sec, 72 °C for 7 min, and held at 4 C° using the Mastercycler

Nexus (Eppendorf, Australia). This was then cycled a total of 40 times. PCR

products were purified with the Zymo DNA Clean & Concentrator Kit™ (Zymo

Research, Irvine, CA, USA).

Following PCR, DNA was cleaned up with Agencourt® AMPure® XP beads

(Beckman Coulter) and washed with 80% ethanol and allowed to air-dry. DNA was

then eluted with 52.5 µl of 10 mM Tris pH 8.5 before being placed back into the

magnetic stand. Once the supernatant had cleared, 50 µl was aliquoted for the

experiment. DNA samples were quantified using the Quant-iT™ PicoGreen™

dsDNA Assay kit (Thermo Fisher) using the FLUROstar® Omega (BMG Labtech).

Samples were processed using the Illumina MiSeq™ 500 cycle Kit V2 and
sequenced on the Illumina MiSeq™ system by Massey Genome Service
(Palmerston North). Illumina MiSeq™ sequences were trimmed using SolexaQA++
software (Cox, Peterson, and Biggs 2010) and aligned to FASTA bisulfite converted
reference sequences using the package Bowtie2 (version 2.3.4.3) Each individual
read was then aligned to all reference sequences using the methylation-specific
package Bismark (Krueger and Andrews 2011).

Statistics

Differential DNA methylation was assessed using the package edgeR (Chen et al.).
Coverage level was set to greater or equal to "8" across unmethylated and
methylated counts, as recommended by (Chen et al.). Two models were used – the
first was a bivariate model, to assess differences between the *in utero* exposed to
tobacco compared to the non-exposed control group (model 1).

$$Y \sim U + AS + e$$

161 The second was a multiple regression to assess the interaction term *in utero*
162 maternal smoke exposure and offspring conduct problem score (high or low, model
163 2).

$$Y \sim U + C + AS + U{:}C + e$$

Where, *Y* is defined as the methylation M ratio, *U* is the exposed/unexposed *in utero*
to maternal smoking, *C* is conduct problem score with high conduct problem score <
53 and low conduct problem core < 46, *AS* adult smoker/non-smoker and *e* is the
unexplained variation or error tem.

This model was fitted with both ANOVA parameters and with contrasts between *in*
utero exposure groups (exposed – non-exposed) within CP score levels. Top tables
were constructed using the topTags function in edgeR, Log fold change, average log
counts per million, and in some cases F statistic and were calculated and nominal
significance was given for $P < 0.05$, these were then corrected using FDR. Scatter
plots with the inclusion of confidence intervals were constructed from log
transformed normalised methylated and unmethylated counts.

Results

Here we assessed DNA methylation within 10 separate genes (Table 2). DNA sequence data for 15 amplicons from these 10 genes (Supplementary Table 1) was generated, comprising a total of 280 CpG sites. These CpG sites included a combination of sites previously identified as differentially methylated, as well as amplification of all CpGs within the promoter region of genes associated with *in utero* neurodevelopment and CP phenotypes (Table 2). Differential methylation across these CpG sites was calculated to address whether any were specifically differentially methylated in individuals with CP, in response to *in utero* tobacco exposure.

Quantification of DNA methylation at previously reported CpG sites in response to *in utero* exposure to tobacco

Initially, we attempted to validate in our cohort (age ~28-30 years) five CpG sites which have been previously reported to be differentially methylated in the DNA of cord blood from newborns, and whole blood from children and adolescents (ages newborn to 17) in response to *in utero* tobacco exposure (Table 1). Data were partitioned into those individuals exposed *in utero*, and those who were not (model 1), to assess whether or not BSAS could detect previously reported CpG sites (Table 3).

AHRR (cg05575921) displayed a 3.1% decrease in DNA methylation between exposed and non-exposed individuals, at a nominal P value of 0.02. This site has been previously identified as hypomethylated in adult tobacco smokers, as well as in postnatal cord blood samples between *in utero* tobacco-exposed and non-exposed individuals. The probe cg05549655 in the gene *CYP1A1* displayed a 5.19% increase in DNA methylation in the *in utero* exposed group, however, this site did not reach nominal statistical significance in our cohort. Cg09935388 and cg09662411 in *GFI1* were unable to be replicated as differentially methylated between the exposed and the non-exposed groups (no significant change in β values). Both CpG sites show

hypomethylation, supporting previous observations of differential methylation within
this gene. *CNTNAP2* (cg2594950) was similarly unable to be validated in our cohort
using the method BSAS.

Differentially methylated CpGs under the interaction of *in utero* tobacco
exposure and CP

Differential methylation dependent on both *in utero* exposure and CP score was
found at 10 loci in six genes at nominal significance level, none were significant after
correcting for false discovery rate (Table 4).

Of these CpG sites, five of the 10 CpG sites were found in the following genes:
CYP1A1, *GFI1*, ASH2L *and GRIN2b*. Differential methylation between *in utero*
exposed and non-exposed associated with for high conduct scores. No nominal
significance from the interaction observed in association with low conduct scores.
The top three CpG sites with nominal significance under the interaction are displayed
in Figure 1. Here, differential methylation is found in response to high CP score and
no differences are seen between the exposed and non-exposed low CP groups
(Figure 1).

Discussion

In utero tobacco exposure is known to alter DNA methylation at the genome-wide
level in offspring (Joubert, Haberg, et al. 2012; Joubert, Felix, Yousefi, Bakulski,
Just, Breton, Reese, Markunas, Richmond, Xu, et al. 2016) (Joubert Bonnie et al.
2012; Richmond et al. 2014). The later-life implications of these tobacco-induced
DNA methylation changes are unclear, however, an association between *in utero*
tobacco exposure and CP has previously been observed (Sengupta et al. 2017).
Given the complex etiology of CP phenotypes (Acosta, Arcos-Burgos, and Muenke
2004; Beaver et al. 2007; Salvatore and Dick 2018) and the vast array of
socioeconomic variables associated with tobacco use (Lantz et al. 1998), proving a

causal link between maternal smoking and offspring CP is inherently challenging.
Previously we quantified tobacco-induced DNA methylation changes that associate
with CP phenotypes in offspring exposed to tobacco *in utero* (via maternal smoking)
using the Illumina EPIC array, with results indicating that methylation was altered at
genes that may have roles in neurodevelopment and CP phenotypes. However, due
to a combination of a comparatively small sample size relative to the number of loci
on the array, only nominal significance was observed. The data suggested a role for
DNA methylation in the link between exposure and CP, so here we chose a panel of
genes with known roles in these things, to see if methylation is changed at these loci
too. Due to our small sample size we need to try an alternative approach. We've
shown that BSAS previously is very good for targeting differential methylated so here
we use BSAS at some targeted genes to see if we can detect differential methylation
specific to the interaction between high CP score and exposure?

Validation of previously identified differentially methylated CpG from *in utero*
tobacco exposure

First, we asked whether differentially methylated CpGs that have been previously
associated with *in utero* tobacco exposure were supported by this cohort. Here, we
present validation of differential methylation of a CpG site within the gene *AHRR*
(cg05575921). *AHRR* is a well-defined tobacco smoking gene, which is consistently
represented in tobacco methylation data. *AHRR* has previously been found to be
differentially methylated in response to *in utero* tobacco exposure (Richmond,
Simpkin, Woodward, Gaunt, Lyttleton, McArdle, Ring, Smith, Timpson, Tilling, Davey
Smith, et al. 2015; Joubert, Håberg, Nilsen, Wang, Vollset, Murphy, Huang, Hoyo,
Midttun, Cupul-Uicab, et al. 2012; de Vocht et al. 2015). Four other CpG sites
investigated here due to previous association with *in utero* tobacco exposure were
not differentially methylated in our data. However, the direction of methylation
change was supported at all five sites investigated (Tehranifar et al. 2018; Rauschert
et al. 2019; Rotroff et al. 2016). We suggest that further investigation in a larger
cohort may lead to nominal significance at the sites in *CYP1A1*, *CNTNAP2,* and
GFI1.

Identification of *in utero* exposure-related differentially methylated CpG sites that are specific to individuals with high CP scores

Epidemiological data suggests that there is an increased association between *in utero* tobacco exposure and behavioural disorder in children and adolescents (Carter et al. 2008; Mick et al. 2002). Thus, here, we investigated DNA methylation changes induced by *in utero* tobacco exposure as a potential molecular mechanism of dysfunction that could link the phenotypic trait of CP to maternal tobacco use during pregnancy. We therefore analysed DNA methylation patterns within our gene panel in response to *in utero* tobacco exposure and its interaction with CP status. A total of 10 CpG sites in seven genes were found to display nominal significance in DNA methylation in response to *in utero* tobacco exposure and CP in this cohort (Table 4).

In the 10 CpG sites we identified under the interaction, *CYP1A1* showed greater magnitude differential methylation in high CP scores (exposed *in utero* vs. non-exposed with high CPS), with reduced reversed or no evidence of differential methylation at the same sites with low CP score. This indicates that within the observed nominal methylation changes the interaction was being driven in the high CP score group. One gene (*ASH2L*), contained three nominally significantly differentially methylated CpG sites, and *CYP1A1* and *MEF2C* both had two.

CYP1A1 (Cytochrome P450 family 1 subfamily A member 1) is a well-established marker for *in utero* tobacco smoke exposure (Richmond et al. 2018; Lee Ken et al. 2015; Richmond et al. 2014; Tehranifar et al. 2018). Neither of the two sites we observed under the interaction have probes at these locations on the Illumina array system, thus emphasising a benefit of amplicon sequencing compared to an array-based method. Variant differences in *CYP1A1* have previously been associated with child behavioural problems at age 2, from prenatal maternal environmental tobacco smoke (Hsieh et al. 2010). This highlights the need for this gene to be further investigated for its role in the development of conduct problems following *in utero* tobacco exposure.

Three CpG sites from the gene *ASH2L* (ASH2 like histone lysine methyltransferase complex subunit) showed in consistent levels of differential methylation in response to *in utero* tobacco exposure and CP, with two displaying hyper- and one hypomethylation. *ASH2L* has been found to interact with *MEF2C* (Myocyte enhancer

factor 2C) to mediate changes in histone 3 lysine 4 trimethylation (H3K4me3) (Jung et al.). Here, we detected nominal significance at two CpG sites within *MEF2C* (chr5, 88179596 and 88179541). Both of these sites were associated with a greater level of hypomethylation in participants who were exposed to tobacco *in utero* with high CP scores in this cohort, although not at the FDR significance level. *MEF2C* plays a role in neural crest formation during development, where tissue-specific inactivation of the gene results in embryonic lethality (Verzi et al. 2007). Further, *MEF2* interacts with oxytocin, which is affiliated with prosocial behaviours (Kosfeld et al. 2005; Zak, Stanton, and Ahmadi 2007). Alterations to oxytocin have been shown to change the morphology of neurons via *MEF2A* (Meyer et al. 2018; Meyer et al. 2020). Functional roles of the gene in relation to early neuronal development still remain unclear, however it is thought to play a crucial role (Harrington et al. 2016). Recent research in animal models suggests that nicotine-dependent induction of the *ASH2L* and *MEF2C* complex during development induces alterations that could lead to fundamental changes in the brain (Jung et al.).

While we cannot assert causality, our targeted approach shows that *in utero* tobacco exposure may be altering methylation at CpG sites associated with neural phenotypes which persist into adulthood and are then associated with increased risk of high CP.

Conclusion

Here we have presented preliminary data to suggest that the association between maternal tobacco use during pregnancy and the development of CP in children and adolescents may in part be mediated by altered DNA methylation, induced by *in utero* tobacco exposure during development, at genes that have roles in *in utero* brain development and CP phenotypes. We acknowledge the limitations of this study described above, however, the data presented here are suggestive of a role for DNA methylation in the link between *in utero* tobacco exposure and offspring CP. Our findings should stimulate further study using larger sample sizes.

Abbreviations

CP   Conduct problems

CHDS Christchurch health and development study

BSAS Bisulfite based amplicon sequencing

SIDS   Sudden infant death syndrome

ADHD Attention-deficit hyperactivity disorder

DOHaD Developmental origins of human health and disease

GFI1   Growth Factor Independent one transcriptional repressor

CPS   Conduct disorder score

AHRR   Aryl hydrocarbon receptor repressor

ASH2L ASH2 like histone lysine methyltransferase complex subunit

BDNF Brain-derived neurotrophic factor,

CNTNAP2 Contactin associated protein 2

CYP1A1 Cytochrome P450 Family 1 Subfamily A Member 1

DUSP6 Dual specificity phosphatase 6

GRIN2b Glutamate Ionotropic Receptor NMDA Type Subunit 2B

MEF2C Myocyte enhancer factor 2C

PRDM8 PR/SET Domain 8

FC Fold change

CPM   Counts per million

FDR   False discovery rate

Funding

Funding for this study came from the Maurice and Phyllis Paykel Trust. CHDS was
funded by the Health Research Council of New Zealand (Programme Grant 16/600).
CMRF supplied funding for the manuscript to be written.

Availability of data

Upon request.

Contributions

AJN-molecular lab work, data analysis, and major contributor to manuscript. JFP-
study design, data analysis, and major contributor to manuscript. ADN- data

analysis. JMB and LJH study design, provided DNA samples via CHDS. MAK- study
design and over view. AJO- study design, molecular lab work, major contributor to
manuscript and source of funding. All authors read and approved the final
manuscript.

Acknowledgements

Not applicable

Ethics declarations

All aspects of the study were approved by the Southern Health and Disability Ethics
Committee, under application number CTB/04/11/234/AM10 "Collection of DNA in
the Christchurch Health and Development Study".

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

References

Acosta, Maria Teresa, Mauricio Arcos-Burgos, and Maximilian Muenke. 2004. 'Attention deficit/hyperactivity disorder (ADHD): Complex phenotype, simple genotype?', *Genetics in Medicine*, 6: 1-15.

Beaver, Kevin M., John Paul Wright, Matt DeLisi, Anthony Walsh, Michael G. Vaughn, Danielle Boisvert, and Jamie Vaske. 2007. 'A gene × gene interaction between DRD2 and DRD4 is associated with conduct disorder and antisocial behavior in males', *Behavioral and Brain Functions*, 3: 30.

Carter, Sarnia, Janis Paterson, Wanzhen Gao, and Leon Iusitini. 2008. 'Maternal smoking during pregnancy and behaviour problems in a birth cohort of 2-year-old Pacific children in New Zealand', *Early Human Development*, 84: 59-66.

Chen, Yunshun, Bhupinder Pal, Jane E. Visvader, and Gordon K. Smyth. 2017. 'Differential methylation analysis of reduced representation bisulfite sequencing experiments using edgeR', *F1000Res*, 6: 2055-55.

Conners, C. Keith. 1969. 'A teacher rating scale for use in drug studies with children', *The American Journal of Psychiatry*, 126: 884-88.

———. 1970. 'Symptom patterns in hyperkinetic, neurotic, and normal children', *Child development*, 41: 667-82.

Cox, Murray P., Daniel A. Peterson, and Patrick J. Biggs. 2010. 'SolexaQA: At-a-glance quality assessment of Illumina second-generation sequencing data', *BMC Bioinformatics*, 11: 485.

de Vocht, Frank, Andrew J Simpkin, Rebecca C Richmond, Caroline Relton, and Kate Tilling. 2015. 'Assessment of offspring DNA methylation across the lifecourse associated with prenatal maternal smoking using Bayesian Mixture Modelling', *International journal of environmental research and public health*, 12: 14461-76.

Demontis, Ditte, Raymond K Walters, Joanna Martin, Manuel Mattheisen, Thomas Damm Als, Esben Agerbo, Rich Belliveau, Jonas Bybjerg-Grauholm, Marie Bækved-Hansen, and Felecia Cerrato. 2017. 'Discovery of the first genome-wide significant risk loci for ADHD', *bioRxiv*: 145581.

Fergusson, D. M., L. J. Horwood, and M. Lloyd. 1991. 'Confirmatory factor models of attention deficit and conduct disorder', *Journal of child psychology and psychiatry, and allied disciplines*, 32: 257-74.

Fergusson, D. M., L. J. Horwood, and E. M. Ridder. 2005. 'Show me the child at seven: the consequences of conduct problems in childhood for psychosocial functioning in adulthood', *Journal of child psychology and psychiatry, and allied disciplines*, 46: 837-49.

Harrington, Adam J., Aram Raissi, Kacey Rajkovich, Stefano Berto, Jaswinder Kumar, Gemma Molinaro, Jonathan Raduazzo, Yuhong Guo, Kris Loerwald, Genevieve Konopka, Kimberly M. Huber, and Christopher W. Cowan. 2016. 'MEF2C regulates cortical inhibitory and excitatory synapses and behaviors relevant to neurodevelopmental disorders', *eLife*, 5: e20059.

Hsieh, Chia-Jung, Suh-Fang Jeng, Yi-Ning Su, Hua-Fang Liao, Wu-Shiun Hsieh, Kuen-Yuh Wu, and Pau-Chung Chen. 2010. 'CYP1A1 Modifies the Effect of Maternal Exposure to Environmental Tobacco Smoke on Child Behavior', *Nicotine & Tobacco Research*, 12: 1108-17.

Jiao, S. S., L. L. Shen, C. Zhu, X. L. Bu, Y. H. Liu, C. H. Liu, X. Q. Yao, L. L. Zhang, H. D. Zhou, D. G. Walker, J. Tan, J. Götz, X. F. Zhou, and Y. J. Wang. 'Brain-derived neurotrophic factor protects against tau-related neurodegeneration of Alzheimer's disease'.

Joelsson, Petteri, Roshan Chudal, Ardesheer Talati, Auli Suominen, Alan S. Brown, and Andre Sourander. 2016. 'Prenatal smoking exposure and neuropsychiatric comorbidity of ADHD: a finnish nationwide population-based cohort study', *BMC Psychiatry*, 16: 306.

Joubert, B. R., J. F. Felix, P. Yousefi, K. M. Bakulski, A. C. Just, C. Breton, S. E. Reese, C. A. Markunas, R. C. Richmond, C. J. Xu, L. K. Kupers, S. S. Oh, C. Hoyo, O. Gruzieva, C. Soderhall, L. A. Salas, N. Baiz, H. Zhang, J. Lepeule, C. Ruiz, S. Ligthart, T. Wang, J. A. Taylor, L. Duijts, G. C. Sharp, S.

A. Jankipersadsing, R. M. Nilsen, A. Vaez, M. D. Fallin, D. Hu, A. A. Litonjua, B. F. Fuemmeler, K. Huen, J. Kere, I. Kull, M. C. Munthe-Kaas, U. Gehring, M. Bustamante, M. J. Saurel-Coubizolles, B. M. Quraishi, J. Ren, J. Tost, J. R. Gonzalez, M. J. Peters, S. E. Haberg, Z. Xu, J. B. van Meurs, T. R. Gaunt, M. Kerkhof, E. Corpeleijn, A. P. Feinberg, C. Eng, A. A. Baccarelli, S. E. Benjamin Neelon, A. Bradman, S. K. Merid, A. Bergstrom, Z. Herceg, H. Hernandez-Vargas, B. Brunekreef, M. Pinart, B. Heude, S. Ewart, J. Yao, N. Lemonnier, O. H. Franco, M. C. Wu, A. Hofman, W. McArdle, P. Van der Vlies, F. Falahi, M. W. Gillman, L. F. Barcellos, A. Kumar, M. Wickman, S. Guerra, M. A. Charles, J. Holloway, C. Auffray, H. W. Tiemeier, G. D. Smith, D. Postma, M. F. Hivert, B. Eskenazi, M. Vrijheid, H. Arshad, J. M. Anto, A. Dehghan, W. Karmaus, I. Annesi-Maesano, J. Sunyer, A. Ghantous, G. Pershagen, N. Holland, S. K. Murphy, D. L. DeMeo, E. G. Burchard, C. Ladd-Acosta, H. Snieder, W. Nystad, G. H. Koppelman, C. L. Relton, V. W. Jaddoe, A. Wilcox, E. Melen, and S. J. London. 2016. 'DNA Methylation in Newborns and Maternal Smoking in Pregnancy: Genome-wide Consortium Meta-analysis', *Am J Hum Genet*, 98: 680-96.

Joubert, B. R., S. E. Håberg, R. M. Nilsen, X. Wang, S. E. Vollset, S. K. Murphy, Z. Huang, C. Hoyo, Ø Midttun, L. A. Cupul-Uicab, P. M. Ueland, M. C. Wu, W. Nystad, D. A. Bell, S. D. Peddada, and S. J. London. 2012. '450K epigenome-wide scan identifies differential DNA methylation in newborns related to maternal smoking during pregnancy', *Environ Health Perspect*, 120: 1425-31.

Joubert, B. R., S. E. Haberg, R. M. Nilsen, X. Wang, S. E. Vollset, S. K. Murphy, Z. Huang, C. Hoyo, O. Midttun, L. A. Cupul-Uicab, P. M. Ueland, M. C. Wu, W. Nystad, D. A. Bell, S. D. Peddada, and S. J. London. 2012. '450K epigenome-wide scan identifies differential DNA methylation in newborns related to maternal smoking during pregnancy', *Environ Health Perspect*, 120: 1425-31.

Joubert, Bonnie R, Janine F Felix, Paul Yousefi, Kelly M Bakulski, Allan C Just, Carrie Breton, Sarah E Reese, Christina A Markunas, Rebecca C Richmond, and Cheng-Jian Xu. 2016. 'DNA methylation in newborns and maternal smoking in pregnancy: genome-wide consortium meta-analysis', *The American Journal of Human Genetics*, 98: 680-96.

Joubert, Bonnie R, Siri E Håberg, Roy M Nilsen, Xuting Wang, Stein E Vollset, Susan K Murphy, Zhiqing Huang, Cathrine Hoyo, Øivind Midttun, and Lea A Cupul-Uicab. 2012. '450K epigenome-wide scan identifies differential DNA methylation in newborns related to maternal smoking during pregnancy', *Environmental health perspectives*, 120: 1425.

Joubert Bonnie, R., E. Håberg Siri, M. Nilsen Roy, Xuting Wang, E. Vollset Stein, K. Murphy Susan, Zhiqing Huang, Cathrine Hoyo, Øivind Midttun, A. Cupul-Uicab Lea, M. Ueland Per, C. Wu Michael, Wenche Nystad, A. Bell Douglas, D. Peddada Shyamal, and J. London Stephanie. 2012. '450K Epigenome-Wide Scan Identifies Differential DNA Methylation in Newborns Related to Maternal Smoking during Pregnancy', *Environmental Health Perspectives*, 120: 1425-31.

Jung, Yonwoo, Lawrence S. Hsieh, Angela M. Lee, Zhifeng Zhou, Daniel Coman, Christopher J. Heath, Fahmeed Hyder, Yann S. Mineur, Qiaoping Yuan, David Goldman, Angelique Bordey, and Marina R. Picciotto. 'An epigenetic mechanism mediates developmental nicotine effects on neuronal structure and behavior'.

Kosfeld, Michael, Markus Heinrichs, Paul J. Zak, Urs Fischbacher, and Ernst Fehr. 2005. 'Oxytocin increases trust in humans', *Nature*, 435: 673-76.

Krueger, Felix, and Simon R. Andrews. 2011. 'Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications', *Bioinformatics*, 27: 1571-72.

Lantz, Paula M., James S. House, James M. Lepkowski, David R. Williams, Richard P. Mero, and Jieming Chen. 1998. 'Socioeconomic Factors, Health Behaviors, and MortalityResults From a Nationally Representative Prospective Study of US Adults', *JAMA*, 279: 1703-08.

Lee Ken, W. K., Rebecca Richmond, Pingzhao Hu, Leon French, Jean Shin, Celine Bourdon, Eva Reischl, Melanie Waldenberger, Sonja Zeilinger, Tom Gaunt, Wendy McArdle, Susan Ring,

Geoff Woodward, Luigi Bouchard, Daniel Gaudet, Davey Smith George, Caroline Relton, Tomas Paus, and Zdenka Pausova. 2015. 'Prenatal Exposure to Maternal Cigarette Smoking and DNA Methylation: Epigenome-Wide Association in a Discovery Sample of Adolescents and Replication in an Independent Cohort at Birth through 17 Years of Age', *Environmental Health Perspectives*, 123: 193-99.

Li, Liang, Xiangbin Ruan, Chang Wen, Pan Chen, Wei Liu, Liyuan Zhu, Pan Xiang, Xiaoling Zhang, Qunfang Wei, Lin Hou, Bin Yin, Jiangang Yuan, Boqin Qiang, Pengcheng Shu, and Xiaozhong Peng.

Meyer, M., I. Berger, J. Winter, and B. Jurek. 2018. 'Oxytocin alters the morphology of hypothalamic neurons via the transcription factor myocyte enhancer factor 2A (MEF-2A)', *Mol Cell Endocrinol*, 477: 156-62.

Meyer, Magdalena, Kerstin Kuffner, Julia Winter, Inga D. Neumann, Christian H. Wetzel, and Benjamin Jurek. 2020. 'Myocyte Enhancer Factor 2A (MEF2A) Defines Oxytocin-Induced Morphological Effects and Regulates Mitochondrial Function in Neurons', *International Journal of Molecular Sciences*, 21: 2200.

Mick, Eric, Joseph Biederman, Stephen V. Faraone, Julie Sayer, and Seth Kleinman. 2002. 'Case-Control Study of Attention-Deficit Hyperactivity Disorder and Maternal Smoking, Alcohol Use, and Drug Use During Pregnancy', *Journal of the American Academy of Child & Adolescent Psychiatry*, 41: 378-85.

Noble, Alexandra J, John F Pearson, Joseph M Boden, L. John Horwood, Martin A Kennedy, and Amy J Osborne. 2021. 'Hypomethylation in FASTKD1 detected in the association between in utero tobacco exposure and conduct problem in a New Zealand longitudinal study': 2021.04.08.438710.

Noble, Alexandra, John Pearson, Joseph Boden, John Horwood, Neil Gemmell, Martin Kennedy, and Amy Osborne. 2020. 'A validation of Illumina EPIC array system with bisulfite-based amplicon sequencing', *bioRxiv*: 2020.05.25.115428.

Rauschert, Sebastian, Phillip E. Melton, Graham Burdge, Jeffrey M. Craig, Keith M. Godfrey, Joanna D. Holbrook, Karen Lillycrop, Trevor A. Mori, Lawrence J. Beilin, Wendy H. Oddy, Craig Pennell, and Rae-Chi Huang. 2019. 'Maternal Smoking During Pregnancy Induces Persistent Epigenetic Changes Into Adolescence, Independent of Postnatal Smoke Exposure and Is Associated With Cardiometabolic Risk', *Frontiers in genetics*, 10: 770-70.

Richmond, R. C., A. J. Simpkin, G. Woodward, T. R. Gaunt, O. Lyttleton, W. L. McArdle, S. M. Ring, Adac Smith, N. J. Timpson, K. Tilling, G. D. Smith, and C. L. Relton. 2015. 'Prenatal exposure to maternal smoking and offspring DNA methylation across the lifecourse: findings from the Avon Longitudinal Study of Parents and Children (ALSPAC)', *Human molecular genetics*, 24: 2201-17.

Richmond, Rebecca C, Matthew Suderman, Ryan Langdon, Caroline L Relton, and George Davey Smith. 2018. 'DNA methylation as a marker for prenatal smoke exposure in adults', *International Journal of Epidemiology*, 47: 1120-30.

Richmond, Rebecca C., Andrew J. Simpkin, Geoff Woodward, Tom R. Gaunt, Oliver Lyttleton, Wendy L. McArdle, Susan M. Ring, Andrew D. A. C. Smith, Nicholas J. Timpson, Kate Tilling, George Davey Smith, and Caroline L. Relton. 2015. 'Prenatal exposure to maternal smoking and offspring DNA methylation across the lifecourse: findings from the Avon Longitudinal Study of Parents and Children (ALSPAC)', *Human molecular genetics*, 24: 2201-17.

Richmond, Rebecca C., Andrew J. Simpkin, Geoff Woodward, Tom R. Gaunt, Oliver Lyttleton, Wendy L. McArdle, Susan M. Ring, Andrew D.A.C. Smith, Nicholas J. Timpson, Kate Tilling, George Davey Smith, and Caroline L. Relton. 2014. 'Prenatal exposure to maternal smoking and offspring DNA methylation across the lifecourse: findings from the Avon Longitudinal Study of Parents and Children (ALSPAC)', *Human Molecular Genetics*, 24: 2201-17.

Riva, Valentina, Marco Battaglia, Maria Nobile, Francesca Cattaneo, Claudio Lazazzera, Sara Mascheretti, Roberto Giorda, Chantal Mérette, Claudia Émond, and Michel Maziade. 2015.

'GRIN2B predicts attention problems among disadvantaged children', *European child &*
adolescent psychiatry, 24: 827-36.

Rotroff, Daniel M, Bonnie R Joubert, Skylar W Marvel, Siri E Håberg, Michael C Wu, Roy M Nilsen, Per
M Ueland, Wenche Nystad, Stephanie J London, and Alison Motsinger-Reif. 2016. 'Maternal
smoking impacts key biological pathways in newborns through epigenetic modification in
Utero', *BMC genomics*, 17: 976.

Rutter M, Tizard J, Whitmore K. 1970. 'Education, Health and Behaviour', *London: Longmans*.

Rzehak, Peter, Richard Saffery, Eva Reischl, Marcela Covic, Simone Wahl, Veit Grote, Annick
Xhonneux, Jean-Paul Langhendries, Natalia Ferre, Ricardo Closa-Monasterolo, Elvira Verduci,
Enrica Riva, Piotr Socha, Dariusz Gruszfeld, Berthold Koletzko, and group European
Childhood Obesity Trial Study. 2016. 'Maternal Smoking during Pregnancy and DNA-
Methylation in Children at Age 5.5 Years: Epigenome-Wide-Analysis in the European
Childhood Obesity Project (CHOP)-Study', *PLOS ONE*, 11: e0155554.

Salvatore, Jessica E., and Danielle M. Dick. 2018. 'Genetic influences on conduct disorder',
Neuroscience & Biobehavioral Reviews, 91: 91-101.

Sengupta, Sarojini M., Alicia K. Smith, Natalie Grizenko, and Ridha Joober. 2017. 'Locus-specific DNA
methylation changes and phenotypic variability in children with attention-deficit
hyperactivity disorder', *Psychiatry Research*, 256: 298-304.

Skogstrand, Kristin, Christian Munch Hagen, Nis Borbye-Lorenzen, Michael Christiansen, Jonas
Bybjerg-Grauholm, Marie Bækvad-Hansen, Thomas Werge, Anders Børglum, Ole Mors,
Merethe Nordentoft, Preben Bo Mortensen, and David Michael Hougaard. 'Reduced
neonatal brain-derived neurotrophic factor is associated with autism spectrum disorders'.

Suter, Melissa, Adi Abramovici, Lori Showalter, Min Hu, Cynthia Do Shope, Michael Varner, and
Kjersti Aagaard-Tillery. 2010. 'In utero tobacco exposure epigenetically modifies placental
CYP1A1 expression', *Metabolism-Clinical and Experimental*, 59: 1481-90.

Tehranifar, Parisa, Hui-Chen Wu, Jasmine A. McDonald, Farzana Jasmine, Regina M. Santella, Irina
Gurvich, Julie D. Flom, and Mary Beth Terry. 2018. 'Maternal cigarette smoking during
pregnancy and offspring DNA methylation in midlife', *Epigenetics*, 13: 129-34.

van Otterdijk, Sanne D, Alexandra M Binder, and Karin B Michels. 2017. 'Locus-specific DNA
methylation in the placenta is associated with levels of pro-inflammatory proteins in cord
blood and they are both independently affected by maternal smoking during pregnancy',
Epigenetics, 12: 875-85.

Verzi, Michael P., Pooja Agarwal, Courtney Brown, David J. McCulley, John J. Schwarz, and Brian L.
Black. 2007. 'The Transcription Factor MEF2C Is Required for Craniofacial Development',
Developmental Cell, 12: 645-52.

Wakschlag, L. S., B. B. Lahey, R. Loeber, S. M. Green, R. A. Gordon, and B. L. Leventhal. 1997.
'Maternal smoking during pregnancy and the risk of conduct disorder in boys', *Arch Gen*
Psychiatry, 54: 670-6.

Zak, Paul J., Angela A. Stanton, and Sheila Ahmadi. 2007. 'Oxytocin increases generosity in humans',
PLOS ONE, 2: e1128-e28.

Table 1 - CHDS subsets selected for analysis. The range of conduct problem scores in each category is indicated in brackets. A score of 53 or more is the top quartile for CP, a score of 60 or more the top decile for CP.

		Group 1 Exposed *in utero* and never smokers	Group 2 Exposed *in utero* and a regular smoker	Group 3 Not *exposed in utero* and never smokers
		n= 32	n= 32	n= 32
Sex				
	Male	69%	72%	60%
	Female	31%	28%	40%
Tobacco smoking status at the time of blood collection				
	Never	100%	0%	100%
	Regular	0%	100%	0%
Conduct problem score (CPS)				
	Low CPS (<46)	n= 16 (42-46)	n= 16 (42-46)	n= 16 (41-43)
	High CPS (>53)	n= 16 (53-75)	n= 16 (60-85)	n= 16 (53-68)

591   *Table 2 -* Genes selected to investigate the link between in utero tobacco exposure
592   and CP.

Gene	Function	Significance
AHRR (Joubert, Håberg, Nilsen, Wang, Vollset, Murphy, Huang, Hoyo, Midttun, and Cupul-Uicab 2012; Richmond, Simpkin, Woodward, Gaunt, Lyttleton, McArdle, Ring, Smith, Timpson, Tilling, Smith, et al. 2015; de Vocht et al. 2015; van Otterdijk, Binder, and Michels 2017; Rotroff et al. 2016)	Mediates toxicity of dioxin (found in cigarette smoke)	Hypomethylated in tobacco smokers and their offspring
ASH2L (Li et al.)	Histone lysine methyltransferase	Associated with schizophrenia
BDNF (Skogstrand et al. ; Jiao et al.)	Nerve growth factor	Promotes neuronal survival. Implicated in neurodegenerative disease
CNTNAP2 (Joubert, Felix, Yousefi, Bakulski, Just, Breton, Reese, Markunas, Richmond, and Xu 2016; Richmond, Simpkin, Woodward, Gaunt, Lyttleton, McArdle, Ring, Smith, Timpson, Tilling, Smith, et al. 2015; Rzehak et al. 2016)	Neurexin family – functions in vertebrate nervous system	Implicated in schizophrenia, autism, ADHD, intellectual disability. Hypomethylated in offspring of maternal smoking
CYP1A1 (Suter et al. 2010; Rotroff et al. 2016; Joubert, Håberg, Nilsen, Wang, Vollset, Murphy, Huang, Hoyo, Midttun, and Cupul-Uicab 2012; Richmond, Simpkin, Woodward, Gaunt, Lyttleton, McArdle, Ring, Smith, Timpson, Tilling, Smith, et al. 2015;	Monooxygenase – expression is induced by hydrocarbons found in cigarette smoke	Hypomethylated in offspring of maternal smoking

Gene	Function	Notes
de Vocht et al. 2015; van Otterdijk, Binder, and Michels 2017)		
DUSP6 (Demontis et al. 2017)	Protein phosphatase, cellular proliferation and differentiation	Regulates neurotransmitter homeostasis
GFI1 (Rotroff et al. 2016; Joubert, Håberg, Nilsen, Wang, Vollset, Murphy, Huang, Hoyo, Midttun, and Cupul-Uicab 2012; van Otterdijk, Binder, and Michels 2017)	Zinc finger protein - transcriptional repressor	Part of a complex that controls histone modifications and gene silencing. Hypermethylated in offspring of maternal smoking
GRIN2B (Riva et al. 2015)	Glutamate receptor – expressed early in the brain and is required for normal brain development	Mutations associated with autism, ADHD, schizophrenia
MEF2C (Demontis et al. 2017)	MEF2C is associated with hippocampal-dependent learning and memory	MEF2C is crucial for normal neuronal development. Associated with ADHD
PRDM8 (Joubert, Felix, Yousefi, Bakulski, Just, Breton, Reese, Markunas, Richmond, and Xu 2016)	Histone methyltransferase - Controls expression of genes involved in neural development and neuronal differentiation	Hypomethylated in offspring of maternal smoking

Table 3 - Previously reported CpG sites showing differential DNA methylation in response to *in utero* tobacco exposure, and their average methylation values in individuals from this cohort.

Gene	Illumina ID	Exposed *in*	Non-	β difference	P value

		utero β value	exposed in utero β value		
AHRR	cg05575921	72.287	75.448	-3.161	0.022
CNTNAP2	cg2594950	3.845	3.860	-0.014	0.991
CYP1A1	cg05549655	26.894	21.699	5.195	0.425
GFI1	cg09935388	75.151	75.330	-0.582	0.055
GFI1	cg09662411	95.837	97.400	-1.583	0.274

626 *Table 4 -* CpG sites where differential methylation between conduct problem scores
627 differs with *in utero* exposure at $P < 0.05$. Log Fold Change (FC) and P values
628 (unadjusted) from log ratio tests for the effect on normalized methylation ratios of: (1)
629 interaction between *in utero* exposure and Conduct Problem Score, (2) *In utero*
630 exposed - non-exposed contrast within Low CPS and (3) within High CPS

 participants. Loci with nominally significant (P< 0.05) interaction shown, all FDR P
632 values > 0.05.

Gene	CpG location	Interaction[1]		Low CPS[2]		High CPS[3]	
		Log FC	P value	Log FC	P value	Log FC	P value
CYP1A1	Chr15, 75019290	-2.013	0.010	0.344	0.493	-1.669	0.005
GFI1	Chr1, 92947705	-0.957	0.011	0.002	0.992	-0.955	0.001
ASH2L	Chr8, 37962878	1.257	0.024	-0.447	0.253	0.811	0.042
MEF2C	Chr5, 88179596	-1.679	0.040	0.678	0.174	-1.000	0.122
DUSP6	Chr12, 89746588	-1.444	0.041	0.864	0.107	-0.580	0.204
ASH2L	Chr8, 37962657	-0.199	0.042	0.052	0.455	-0.147	0.033
CYP1A1	Chr15, 75019127	-1.221	0.045	0.403	0.319	-0.819	0.072
ASH2L	Chr8, 37962901	1.250	0.046	-0.561	0.205	0.688	0.121
GRIN2b	Chr12, 14133359	2.711	0.048	0.121	0.903	2.832	0.004
MEF2C	Chr5, 88179541	-1.336	0.050	0.615	0.139	-0.720	0.190