Title

Developmental, cellular, and biochemical basis of transparency in the glasswing butterfly

Greta oto

Authors

Aaron F. Pomerantz[1,2*], Radwanul H. Siddique[3,4], Elizabeth I. Cash[5], Yuriko Kishi[6,7],

Charline Pinna[8], Kasia Hammar[2], Doris Gomez[9], Marianne Elias[8], Nipam H. Patel[1,2,6*]

Affiliations

1. Department Integrative Biology, University of California Berkeley, Berkeley, CA
   94720.
2. Marine Biological Laboratory, Woods Hole, MA 02543.
3. Image Sensor Lab, Samsung Semiconductor, Inc., 2 N Lake Ave. Ste. 240, Pasadena,
   CA 91101, USA.
4. Department of Medical Engineering, California Institute of Technology, Pasadena, CA
   91125.
5. Department of Environmental Science, Policy, & Management University of California,
   Berkeley, Berkeley, CA 94720.
6. Department Molecular Cell Biology, University of California Berkeley, Berkeley, CA
   94720.
7. Department of Biology and Biological Engineering, California Institute of Technology,
   Pasadena, CA.
8. ISYEB, 45 rue Buffon, CP50, Paris, CNRS, MNHN, Sorbonne Université, EPHE,
   Université des Antilles, France.
9. CEFE, 1919 route de Mende, Montpellier, CNRS, Univ Montpellier, Univ Paul Valéry
   Montpellier 3, EPHE, IRD, France.

* Corresponding author. Email: pomerantz_aaron@berkeley.edu, npatel@mbl.edu

Abstract (150 words)

Numerous species of Lepidoptera have transparent wings, which often possess scales of altered morphology and reduced size, and the presence of membrane surface nanostructures that dramatically reduce reflection. Optical properties and anti-reflective nanostructures have been characterized for several 'clearwing' Lepidoptera, but the developmental basis of wing transparency is unknown. We apply confocal and electron microscopy to create a developmental time-series in the glasswing butterfly, *Greta oto*, comparing transparent and non-transparent wing regions. We find that scale precursor cell density is reduced in transparent regions, and cytoskeletal organization differs between flat scales in opaque regions, and thin, bristle-like scales in transparent regions. We also reveal that sub-wavelength nanopillars on the wing membrane are wax-based, derive from wing epithelial cells and their associated microvillar projections, and demonstrate their role in enhancing-anti-reflective properties. These findings provide insight into morphogenesis of naturally organized micro- and nanostructures and may provide bioinspiration for new anti-reflective materials.

Introduction

The wings of butterflies and moths (Lepidoptera) have inspired studies across a variety of scientific fields, including evolutionary biology, ecology, and biophysics (*1–3*). Lepidopteran wings are generally covered with rows of flat, partially overlapping scales that endow the wings with colorful patterns. Adult scales are chitin-covered projections that serve as the unit of color for the wing. Each scale can generate color through pigmentation via molecules that selectively absorb certain wavelengths of light, structural coloration, which results from light interacting with the physical nanoarchitecture of the scale, or a combination of both pigmentary and structural coloration (*4, 5*). Cytoskeletal dynamics, including highly organized F-actin filaments during scale cell development, play essential roles in wing scale elongation and prefigure aspects of scale ultrastructure (*6, 7*).

In contrast to typical colorful wings, numerous species of butterflies and moths possess transparent wings that allow light to pass through, so that objects behind them can be distinctly seen (Fig. 1A-H, *8–10*). This trait has been interpreted as an adaptation in the context of camouflage, in which some lineages evolved transparent wings as crypsis to reduce predation (*11–13*). Transparency results from the transmission of light across the visible spectrum through a material, in this case the chitin membrane, without appreciable absorption or reflection. Levels of reflection are largely determined by the differences in refractive indices between biological tissues and the medium, and a larger difference results in higher surface reflection. Our knowledge on mechanisms underlying transparency in nature is primarily from aquatic organisms, which are frequently transparent, aided by the close match between the refractive indices of their aqueous tissue and the surrounding media — water (*14*). By contrast, transparency is rare and more challenging to achieve on land, primarily due to the large difference between the

refractive indices of terrestrial organism's tissue ($n = \sim$1.3-1.5) and air ($n = 1$), which

results in significant surface reflection (*9, 15, 16*).

Nevertheless, some organisms have evolved morphological innovations that

overcome the challenges of terrestrial transparency, notably in the form of anti-reflective

nanostructures. Early studies elucidated highly-ordered sub-wavelength nanostructures

(termed 'nipple arrays') on the corneal surface of insect eyes (*17*). These structures were

found to generally be ~150-250 nm in height and spaced ~200 nm apart, which reduces

reflection across a broad range of wavelengths by creating a smoother gradient of

refractive indices between air and chitin (*18*). Nanostructure arrays have also been

identified on the wings of cicadas, which help to reduce surface reflection over the visible

spectrum (*19*).

Some lepidopterans possess modified wing scales that allow light to reach the

wing surface, which is composed of chitin and has some inherent transparency, but due to

the high refractive index of chitin, $n = 1.56$ (*20*), the wing surface reflects light. For

example, the butterfly *Methona confusa* (Nymphalidae: Ithomiini) has exposed wing

membrane that lacks nanostructures on the surface, and as a result, the wing is somewhat

transparent, but retains a high degree of reflectivity (Fig. 1A-C). Conversely, the longtail

glasswing, *Chorinea faunus* (Riodinidae), contains small, widely spaced scales and dome-

shaped chitin nanoprotuberances on the membrane that generate anti-reflective properties

(Fig. 1D-F) (*21*). The hawkmoth, *Cephonodes hylas* (Sphingidae), has nude wings due to

deciduous scales that fall out upon eclosion, and possesses anti-reflective nanostructures

on its wing surface that morphologically resemble insect corneal nipple arrays (*9*). Nipple

array nanostructures have also been characterized in transparent wing regions of the tiger

moth *Cacostatia ossa* (Erebidae) (*22*). Finally, the glasswing butterfly *Greta oto*

(Nymphalidae: Ithomiini) contains thin, vertically oriented scales, allowing the wing

surface to be exposed, along with nanopillars that coat the surface. These irregularly

arranged nanopillars feature a random height and width distribution and enable

omnidirectional anti-reflective properties (Fig. 1G-I) (*10, 23*). More recent studies have

explored aspects of structural diversity, optical properties, phylogenetic distribution, and

ecological relevance of transparency within a wide range of butterflies and moths,

highlighting that transparency has evolved multiple times independently and may present

evolutionary benefits (*13, 24, 25*).

Lepidoptera are proving to represent an excellent group to investigate

transparency on land, but the developmental processes underlying wing transparency are

currently unknown. This presents a gap in our understanding of lepidopteran wing

evolution and diversification, as transparent butterflies and moths contain multitudes of

intriguing scale modifications and sub-wavelength cuticular nanostructures (*24, 25*). We

therefore set out to explore the development of wing transparency in the glasswing

butterfly *Greta oto*, which belongs to a diverse tribe (~393 species) of predominantly

transparent neotropical butterflies (*26*). We applied confocal and transmission electron

microscopy to compare wing development, scale cytoskeletal organization, and

membrane surface nanostructures between clear and opaque wing regions. Using

chemical treatments, scanning electron microscopy, and gas chromatography–mass

spectrometry, we found that nanostructures on the wing membrane surface are made of

two layers: a lower layer of chitin-based nipple-like nanostructures, and an upper layer of

wax-based nanopillars composed predominantly of long-chain *n*-alkanes. Finally, by

removing the wax-based nanopillars, we demonstrate their role in dramatically reducing

reflection on the wing surface via optical spectroscopy and analytical simulations.

Results

Scale measurements in clear and opaque wing regions of adult *Greta oto*

We investigated features of scale density, scale morphology, and the amount of wing

surface exposed in wings of adult *Greta oto*. We focused on two adjacent regions within

the forewing for consistency: a clear region within the discal cell and an opaque region

that consists mainly of black scales near the M2-M3 crossvein. (Fig 1G,J). The clear wing

region contained two types of alternating scale morphologies: bristle-like scales and

narrow, forked scales, while within the opaque wing region, scale morphologies

resembled 'typical' butterfly pigmented scales: flat and ovoid with serrations at the tips

(Fig1. K,L). The mean density of scales (± SD) in the adult wing were significantly lower

within the clear region (107 ± 19 scales per mm^2) compared to the opaque region (395 ±

23 scales per mm^2) (Student's t-test, P < 0.001, n = 3 individuals, Fig. 1M). In the clear

region, forked scales were significantly smaller in size (498 ± 39 μm^2) compared to the

bristle-like scales (831 ± 183 μm^2), while in the opaque region, scales were the largest

(3467 ± 382 μm^2) (ANOVA test, n = 3 individuals, Fig. 1N). Finally, the amount of

exposed wing membrane was significantly different between wing regions, with an

average of 83.1% ± 0.76 and 2.4% ± 3.4 exposed membrane in the clear and opaque

regions, respectively (Student's t-test, P < 0.001, n = 3 individuals, Fig. 1O).

Morphogenesis and cytoskeletal organization of developing scale cells

To investigate developmental processes of wing and scale development, we performed

dissections of *G. oto* pupae at different time points (Fig. 2). As in other species of

Lepidoptera, the early pupal wing consisted of a thin bilayer of uniform epithelial tissue

and by 16 hours after pupal formation (APF) numerous epidermal cells had differentiated

to produce sensory organ precursor (SOP) cells, which could be identified by

fluorescently labelling tissue with DAPI (Fig. 2B,C) as the SOP's are larger than, and

positioned slightly basal to, the rest of the epidermal cells. The SOPs are precursors to the

scale and socket cells and are organized into parallel rows. At this early stage of wing

development, we observed that the clear wing region harbored a lower density of SOP

cells relative to the opaque wing region (Fig. 2B,C). We can therefore infer that early into

wing development, SOP cell patterning is differentially regulated between clear and

opaque regions, which impacts the adult wing scale density and the amount of wing

membrane surface exposed in different parts of the wing.

Next, we investigated cellular and cytoskeletal organization during scale growth

in clear and opaque wing regions, using simultaneous confocal imaging of fluorescently

labeled scale cell membrane (wheat germ agglutinin; WGA), and F-actin (phalloidin)

(Fig. 2D-I). We found that general aspects of scale development in *G. oto* follow those

previously reported in several butterfly and moth species by *(6)*, with some notable

distinctions for modified scale growth in the clear wing regions of *G. oto*.

By 30 hours APF, the SOP cells have divided to produce the scale and socket cells

(Fig. 2D,E). The scale cell body lies internally within the wing, while the socket cell

associated with each scale cell lies in a more superficial position. At this pupal stage, the

morphological development of wing scale projections has begun, and the scale cells

develop as small buds containing short, densely packed parallel F-actin filaments.

Phalloidin staining showed the appearance of these small cylindrical buds containing F-

actin filaments, and WGA staining showed outlines of the membrane as the scale

outgrowths begin to project and elongate beyond the wing surface. At this stage, budding

scales in the clear wing region appeared morphologically similar to the unspecialized

opaque scales: roughly elongated balloon-shaped with numerous small actin rods fanning

out from the pedicel to the apical tip of the scale. In the clear region, early scale

projections showed alternating sizes. In the opaque region similar budding scales at a

higher density were found, with larger buds corresponding to future cover scales, and

smaller, shorter buds corresponding to future ground scales (Fig. 2D,E).

By 48 hours APF, scale cell extensions have grown and elongated (Fig. 2F,G).

The actin filaments have reorganized into smaller numbers of thick, regularly spaced

bundles along the proximal–distal axis of the scale just under the surface of the cell

membrane. At the base of the scales, fluorescent staining indicated that F-actin bundles

are tightly packed, while in more distal regions we could see an asymmetric distribution

of F-actin, with larger actin bundles in the adwing (facing the wing membrane) side of the

scales (movie S1). At this stage, scales in different regions of the wing had also started to

take on dramatically different morphologies. Scales in the clear region had elongated in a

vertical orientation and obtained two types of alternating morphologies: short and

triangular, or long and bristle-like outgrowths (Fig. 2F). The wings of other butterfly

species contain alternating ground and cover scales, in which the ground scales are

typically smaller in size than the cover scales, consistent with our observations of the

opaque regions of *G. oto* (Fig. 2F). Based on scale size and position, we interpret that

within the clear wing region of glasswing butterflies, the larger bristle-like scales are

modified cover scales and smaller forked scales are modified ground scales (Fig. 2F). In

the opaque region, scales have taken on a round and flattened morphology, similar to

what has been described in other colorful butterfly and moth species, with the ground

scales being shorter and wider than the cover scales (Fig. 2G).

By 60 hours APF, scale projections are even more elongated (Fig. 2H,I). The

triangular scales in the clear wing region have proceeded to generate two new branches,

which fork and elongate at the tips bidirectionally, while bristle-like scales have

elongated and curved (Fig. 2H). In the opaque region, scales were longer, wider, flatter,

and had developed serrations at the tips (Fig. 2I). F-actin bundles extended all the way to the distal tips of these serrations, which is necessary to produce finger-like projections at the tips of scales (*6*). Phalloidin staining also revealed that actin bundles were arranged in more symmetrical patterns around the periphery of the bristle-like scale morphologies, forked scales showed modified actin organization at the branching points, and actin bundle asymmetry was greatest in developing flat opaque scales, with larger bundles present on the adwing side (Fig. 2H,I).

Ultrastructure analysis of developing bristle, forked and opaque scales

To reveal ultrastructural detail of developing wing scale morphology, we performed transmission electron microscopy (TEM) on pupal wing tissue of *G. oto* at 48 hours APF (Fig. 3). In transverse sections, we could resolve distinct scale morphologies (bristle, forked and opaque) and their associated cytoskeletal elements.

Bristle-like scales in the clear wing regions were circular in cross sections (Fig. 3A-C). We could also distinguish between distal and basal regions of bristle-like scales, the latter of which had the presence of a surrounding socket cell in the cross section (Fig. 3B,C). TEM revealed that these bristle-like scales were ringed by peripheral bundles of actin filaments, which lay spaced just under the cell membrane (Fig. 3B-C'). On the adwing side of the scale, the actin bundles were larger and spaced closer to one another relative to the abwing side, and in more distal regions of the bristle-like scale, the actin bundles were more widely spaced and smaller in size. We also observed large populations of microtubules (MTs) distributed throughout the developing scales, which were internal relative to the actin bundles. Interestingly, we observed distinct patterns of microtubule distribution within different developing scale morphologies. The cross section of bristle-like scales revealed large populations of internal microtubules, which we identified due to their characteristic ring shape and diameter of ~25 nm (Fig. 3B',C'). The circular ring

shape of microtubules in cross sections of both the basal and distal parts of the bristle-like

scale suggested that microtubules are all longitudinally oriented, running in the same

direction as the actin filaments, parallel to growth. We also observed that populations of

MTs are localized primarily away from the surface of the scale in its interior, and MTs

were fewer distally than basally (Fig. 3B',C').

In our TEM cross sections we also observed scale types that appeared more

triangular in shape, suggesting that these corresponded to developing forked scales within

the clear wing region (Fig. 3D,E). We observed that these scales were ringed by

peripheral bundles of crosslinked actin filaments, with thicker actin bundles on the

adwing side of the scale. Interestingly, we observed two internal bundles of actin

filaments that were not observed in bristle-like scale morphologies (Fig. 3E'). We also

note that there was variability in MT orientation, rather than the ubiquitous longitudinal

orientations observed in bristle-like scales.

Finally, developing opaque scales were easily identified in cross sections due to

their large size and flattened morphology (Fig. 3F,G). We observed peripheral bundles of

crosslinked actin filaments that were widely spaced and smaller in size in distal parts of

the scale (Fig. 3G-G'). We observed a clear asymmetry in actin bundle size, which were

thicker on the adwing side of the scale relative to the abwing surface. In opaque wing

regions, TEM micrographs revealed what appeared to be concentrated parallel-running

populations of MTs near the narrow base of the scales, and then a more mesh-like

network of MTs in more distal flattened regions, indicating that MTs have varying

orientations within different regions of the scale (Fig. 3G,G', fig. S1). In contrast to the

bristle-like scales, large, flattened opaque scales appeared to contain populations of MTs

that were more widely distributed and less dense.

In all scale types we observed the presence of numerous internal organelles and

vesicles, including mitochondria, electron dense vesicles and free ribosomes (Fig. 3, fig.

S1). We also observed that the actin bundles contained dense, hexagonally packed F-actin

filaments, supporting previous patterns for actin bundle formation in elongating insect

scales (fig. S1). The neck regions of different scale morphologies were predominantly

filled with longitudinally oriented microtubules, actin bundles, and mitochondria.

Longitudinal views also supported that MTs are numerous in the outgrowing scale, and

their spatial arrangement differed with scale position and shape. More mature scales

around 120 hours APF exhibited developed ridge morphologies and thickened cuticle

layers (fig. S1).

Ontogeny of wing membrane nanostructures

The clear wing regions of *G. oto* contain nanopillars that cover the surface of the

membrane (Fig 1I, Fig 4A). These nanopillars were previously characterized in adult

wings, which feature an irregular height distribution and help to generate omnidirectional

anti-reflective properties (*10*). To gain insight into the development of these

nanostructures, we examined the surface of the wing membrane epithelial cells with TEM

(Fig. 4B-F). At 60 hours APF, a perpendicular section through the wing epithelia

showed a continuous epithelial lamina (Fig. 4B,C). We observed the epithelial cells

contained microvilli (MV), which appeared as slender linear extensions from the inner

margins of the developing cells that insert into electron-dense material (Fig. 4B,C). The

surface layer of the epithelia appeared as an extracellular lamellar system, and lamina

evaginations appeared in the section as domes distal to the microvillar extensions (Fig.

4C). By 72 hours APF, we observed a thin outer layer of the epicuticle that rose above

the epidermal cells and by 120 hours APF, we found that this upper layer above the

microvilli contained what appear to be dome-shaped protrusions and thickened cuticle,

possibly secreted from regularly spaced microvilli (Fig. 4D,E). Finally, in our TEM

cross section of a fully developed adult wing of *G. oto*, we observed that the membrane

surface harbors dome-shaped nanoprotrusions with similar morphologies to insect

corneal surface nipple arrays (e.g. *9, 17*), which we refer to throughout the text now as

"nipple nanostructures", and an upper layer containing pillar-like protrusions, which we

refer to as "nanopillars", that featured a more irregular height distribution (Fig. 4F).

These results show early subcellular processes of developing nanopillars within the

clear wing region, which arise distal to microvillar extension in epithelial cells.

Topographical organization and biochemical composition of wing surface

nanostructures

Based on our EM results of membrane nanostructures, we investigated the topographical

organization and biochemical composition of the adult wing surface. To do so, we treated

individual, disarticulated adult *G. oto* wings in two ways: by 1) physically removing wing

surface nanostructures by gently pressing and rubbing a wing in between paper and

Styrofoam (after *9*) and 2) testing the wing surface structures for solubility in organic

solvents, including hexane and chloroform to extract lipids (after *27*). We then performed

SEM to compare wing surface topography of untreated and treated wing samples (Fig.

5A-C'). SEM confirmed that the first treatment partially or completely removed

nanostructures across the wing membrane surface (Fig. 5B). In a region of partial

removal, we could identify smaller, dome-shaped nipple nanostructures underneath the

top layer of nanopillars (Fig. 5B'). SEM of the chemically treated wing surface revealed

that the upper layer of irregularly sized nanopillars were completely removed, revealing a

layer of regularly arranged dome-shaped nipple nanostructures that did not dissolve

through chloroform or hexane exposure (Fig. 5C,C'). Therefore, we hypothesized that the

upper layer of irregularly sized nanopillars consisted of a secreted wax-based material,

which sits above smaller chitin-based nipple nanostructures.

To test this hypothesis, we extracted the surface layer of *G. oto* clear wing regions

with either hexane or chloroform and analyzed the chemical composition by gas

chromatography–mass spectrometry (GC-MS). We found that the chemical profile

generated by both hexane and chloroform extracts yielded similar results (Fig. 5D). In all

extracts, we identified two straight-chain alkanes that made up approximately 2/3 of the

compounds detected: $41.64 \pm 5.75\%$ pentacosane ($C_{25}H_{52}$) and $23.32 \pm 5.35\%$

heptacosane ($C_{27}H_{56}$) (Table S1). The remaining compounds were primarily composed of

slightly larger methyl-branched alkanes (monomethyl and dimethyl C27, C29 and C31)

and esters. Therefore, our results suggest that in *G. oto* there are two components to wing

surface ultrastructure: procuticle-based nipple nanostructures, and an upper epicuticular

layer of irregularly sized nanopillars, composed mainly of straight chain alkanes (Fig.

5D,E).

Anti-reflective properties of wax-based nanopillars

To address whether the wax-based nanopillars play a role in wing reflection, we measured

the reflectance spectra of untreated and hexane-treated wings (Fig. 6). Additionally, we

measured nanostructure geometries and membrane thickness from wing SEM cross

sections (n = 6), and determined the average distance between two nanostructures as $d =$

174 nm, conical shaped cuticular nipple nanostructures height, $h_p = 77$ nm, wax-based

irregular nanopillars radius, $r_{np} = 53$ nm, mean height, $h_{np} = 224$ nm and variance $\sigma_{np} =$

49.3 nm, and membrane thickness, $h_m = 746$ nm and variance $\sigma_m = 43$ nm (Fig. 6B,D, fig.

S2). On the basis of SEM micrographs for treated and untreated samples, we modeled

three wing architectures consisting of 1) nanopillars with variable height together with

cuticle-based nipple nanostructures on the wing membrane, 2) cuticle-based nipple

nanostructures on wing membrane and 3) wing membrane without any nanostructures, to simulate the optical properties for different conditions (Fig. 6E). The simulated reflectance data of the untreated and treated conditions in Fig. 6F closely resembled the experimental ones. In untreated wings of *G. oto*, we found that transparent regions have a low total diffuse reflection of about 2%, which is in line with previous reflectance measurements of this species (Fig. 6F, *10*). By contrast, the hexane treated wings without the upper layer of wax nanopillars had about 2.5 times greater reflectance relative to the untreated wings, and generated an iridescent thin film spectra, even though they harbored dome-shaped nipple nanostructures (Fig. 6D,F).

For simulated data, the overall reflectance ratio of the hexane treated wing to that of the untreated was approximately three, similar to experimental reflectance data (Fig. 6F, Table S2). Most importantly, the simulated results for the untreated wing with wax-based irregular nanopillars make reflectance more uniform across wavelengths, which reduces the iridescent effect of the wing membrane. Finally, we simulated a thin film membrane without any nanostructures, which showed reflectance (averaged from all wavelengths) of the membrane itself to be 8.81 ± 3.46%, whereas the treated and untreated wing reflections were 5.78 ± 2.82% and 1.93 ± 0.77%, respectively (Fig. 6F). While treated wings harboring dome-shaped nipple nanostructures reduced the overall reflectance relative to the membrane only, their effect was not strong enough to reduce reflectance spectra oscillation. The wax-based irregular nanopillars on top introduced a more gradual transition between refractive indices to lessen the oscillation by approximately five-fold, in addition to reducing overall reflection (Fig, 6F). Additionally, we simulated the three wing architecture models considering different mean membrane thicknesses and variance in membrane thickness (fig. S3). We found that variance in wing membrane thickness reduced reflectance spectra oscillations, rather than mean membrane

342   thickness alone, and more peaks appear in the visible spectrum with increasing thickness

343   of the membrane. (fig. S3, Table S3). Overall, these results demonstrate that the non-

344   constant architecture of the wing membrane and wax-based irregular nanopillars on the

345   wing surface of *G. oto* function to dramatically enhance anti-reflective properties.

Discussion

Butterflies and moths have evolved sub-wavelength anti-reflective structural innovations

on their wings that enable them to be transparent. Here we report the details of pupal

wing development and cytoskeletal organization in the glasswing butterfly, *Greta oto*, as

well as insights into the ontogeny and biochemical basis of wing surface nanostructures

that reduce reflection.

The arrangement of unicellular projections in insect integument, such as bristles

and scales, has been a model for research on cellular pattern formation (*28*). Shortly after

pupation, sensory organ precursor (SOP) cells develop from a monolayer of epithelial

cells into orderly arrangements, then differentiate into scale and socket cells. In the

present study, we found that early SOP cell patterning impacts the final adult scale

density in *G. oto* and this feature of spacing scale cells farther apart, and therefore

reducing the overall density of scales, is an initial step to generate clear wings. During

early pupal development, the receptor molecule Notch is expressed in a grid-like pattern

in the wing epithelium (*29*). This may contribute to the parallel rows of uniformly spaced

SOP cells that express low levels of Notch, likely through a lateral inhibition mechanism.

The low-Notch SOP cells express a homolog of the *achaete-scute* proneural transcription

factors, which likely plays a role in scale precursor cell differentiation (*30*). Notch-

mediated lateral inhibition could establish a dense population of ordered SOP cells in the

developing wing, resulting in a characteristic ratio of scale-building and epithelial cells.

Future studies should investigate if modifications in Notch signaling play a role in scale

cell patterning in clearwing butterflies and moths, many of which contain reduced

densities of scale cells (*24, 25*).

The range of morphological diversity among scales and bristles within

Lepidoptera likely results developmentally from components or modifiers of the

cytoskeletal structures and cell membrane. One study surveyed a wide range of

developing butterfly and moth scales and identified that F-actin is required for several

aspects of scale development, including scale cell elongation and proper orientation (*6*).

In the present study, we found that *G. oto* serves as an excellent model to study

differences in bristle and scale morphogenesis, as the wing contains a wide range of

different scale types. In the developing bristle-like scales, we find symmetrical actin

bundles that outline the cell periphery and a large population of longitudinally running

interior microtubules. This is similar to what has been described for developing bristles in

Drosophila melanogaster pupae, which contain peripheral bundles of cross-linked actin

filaments and a large population of microtubules that run longitudinally along the bristle

(*31*). Recently, (*32*) showed that actin bundles play different roles in shaping scales and

bristles in the mosquito *Aedes aegypti*, in which developing bristles contained

symmetrically organized actin bundles, while actin bundle distribution in scales became

more asymmetrically organized. Given that actin dynamics play a variety of roles in

regulating the development of bristles and scales (*6, 7, 32, 33*), we hypothesize that

modifications in F-actin organization of scales in the transparent wing of *G. oto* are

responsible in part for their narrow bristle-like and forked morphologies. In *D.*

melanogaster, subunits of actin are rapidly added to the barbed ends of the actin filaments

of bristles, relying on actin polymerization and bundling for this purpose, and cross-

linking proteins are required early to bring filaments together (*31*). One cross-linking

protein, Fascin, connects filaments together into hexagonally packed bundles. Our TEM

of actin bundles, along with previous studies, support a similar mechanism of hexagonally

packed F-actin bundles in Lepidoptera (Fig. 3, fig. S1) (*6, 7*).

In animal cells, microtubules have been frequently observed in arrangements

parallel to the long axis of cellular extensions, such as axons, dendrites, and developing

lepidopteran scales (*33*). In an analysis of moth scale development, major shape changes

were found to be correlated with changes to the orientation of the cytoplasmic

microtubules (*33*). In the present study, we identified large populations of microtubules

organized throughout developing scales and showed that microtubules are more

concentrated at the base of the scale. We also found that microtubules exhibit different

distributions and orientations relative to distinct scale morphologies, namely between

bristle, forked, and flat, round scales. In *D. melanogaster*, it has been suggested that

bristle microtubules play a role in elongation, noting that they are highly stable, form at

the start of the elongation, and then extend along the shaft as the cell elongates (*30*). A

more recent reinvestigation of the role of MTs in *D. melanogaster* bristle elongation

suggests that two populations of microtubules help to guide bristle development: dynamic

microtubules (with mixed polarity) add bulk to the bristle cytoplasm and are thought to

contribute proper axial growth, while stable microtubules act to polarize the axis of bristle

elongation and are believed to aid organelle and protein distribution (*34, 35*). It would be

interesting for future studies to functionally characterize the role microtubules play in the

development of lepidopteran scales. Overall, we found conservation of developmental

processes in scale formation relative to other previously described Lepidoptera, with

notable differences in clear versus opaque wing regions. These findings lend further

support that general patterns of scale development, including patterns of F-actin

localization and microtubule distribution, seem to be well conserved in Lepidoptera, and

that modifications of scale morphology to achieve clearwing phenotypes, such as narrow bristle-like and forked scales, likely involve alteration of cytoskeletal organization during scale growth.

Chitinous wing membrane has a higher refractive index than air, so as a mechanism that reduces glare, some clearwing species have evolved sub-wavelength anti-reflective nanostructures (*9, 10*). In this study, we identified the early developmental processes of nanostructures that arise in the wing epithelium. We also note interesting parallels of our observations to previous descriptions of developing nanostructures on the surface of insect cornea. Early data on pupal development of corneal nanostructures were produced by detailed electron microscopy studies, showing that corneal nipples emerge during lens formation, and a chitinous layer may be subsequently secreted underlying the nanostructure (*36*). In these observations, development of initial laminar patches formed on top of underlying microvilli. Subsequently, nanostructures (termed nipple structure array) formed on the surface, with the tips of microvilli still attached to the inner surface. Another study subsequently investigated pupal eye development in *D. melanogaster* and identified features of corneal nipple array formation that matched observations previously made in moth eye nanostructure development (*37*). Gemne (*36*) proposed that the corneal nanostructures originate from secretion by the regularly spaced microvilli of the cone lens cells, although there is still debate about the exact nature of how microvilli pre-pattern nanostructure arrays (*38*). Our TEM results provide insight into the early developmental processes of anti-reflective nanostructure formation in the wings of *G. oto*, highlighting certain similarities to nipple array development in insect cornea. It would be interesting for future work to explore if features of nanostructure formation arose independently in insect cuticle as a mechanism to reduce surface reflection.

In contrast to previously described highly ordered nipple arrays on insect eyes (e.g. *18, 38*), the irregularly sized anti-reflective nanopillars in the clear regions of *G. oto* wings consist of an upper layer of wax-based epicuticle sitting above procuticle-based nipple nanostructures. Insect cuticle is an extracellular matrix formed by the epidermis and composed of three layers: the outermost envelope, the middle epicuticle and the inner procuticle (*39*). The envelope and the epicuticle are composed mainly of lipids and proteins, while the procuticle contains the polysaccharide chitin. Many terrestrial arthropods deposit a layer of wax lipids on the surface of their cuticle, which reduces evaporative water loss (*40*). In some species of dragonfly, epicuticular wax-based nanostructures have also been demonstrated to play a role in generating optical properties, such as an ultraviolet reflection (*27*). In mature males of the dragonflies, a dense wax secretion composed of long-chain methyl ketones, in particular 2-pentacosanone, was found to contribute to the UV reflection properties. The chemical composition of nanopillars on the wing surface of cicadas, which have been shown to contribute to wettability and antimicrobial properties, and found that the major epicuticular components are fatty acids and hydrocarbons ranging from C_{17} to C_{44} (*41*). Another study exploring the molecular organization of dragonfly wing epicuticle found that the major components identified were fatty acids and *n*-alkanes with even numbered carbon chains ranging from C_{14} to C_{30} (*42*). Here, we identified that the epicuticular layer of irregularly sized anti-reflective nanopillars in *G. oto* appear to be composed mainly of *n*-alkanes, including pentacosane (C_{25}) and heptacosane (C_{27}) and showed the importance of these structures to attain better transparency.

Due to thin film optics, the thin membranes of insect wings sometimes reflect distinct structural coloration and iridescence (*43*). However, variability and non-constant thickness render the wing membranes as non-ideal thin films, and additional surface

nanoprotrusions can introduce a gradient of refractive indices that reduces thin film

reflections (*44*). For instance, membrane thickness was found to vary over the transparent

wings of the damselfy *Hetaerina americana* from below 1 μm to up to 3 μm, yet

membrane nanoprotrusions acted as an effective impedance matching device to reduce

reflectance (*44*). In that study, average reflectance spectra for the Andromica clearwing

butterfly *Greta andromica* was also calculated, although the wing was treated as a thin

film, and did not address membrane surface nanostructures. By varying thickness in a

Gaussian way while maintaining average thickness, (*44*) found that an increasing width of

the Gaussian progressively reduced modulation of the reflectance spectrum. Similarly, in

the present study, measurements from SEM cross sections of *G. oto* transparent wings

indicate that the membrane thickness is non-constant, and in our optical simulations,

variance in membrane thickness was found to be an important parameter for reduced

reflectance spectra modulation (fig. S3). Overall, we found that variance in membrane

thickness and wax-based nanostructures with irregular height distributions in *G. oto*

reduce iridescence and maintain anti-reflection properties, which likely aid in crypsis

(*11*).

Turing reaction-diffusion mechanisms have been proposed as a model for the

formation of various corneal nanostructure morphologies (such as spacing, height, and

spatial organization) during insect eye development (reviewed in *38*). Although the

degree of height irregularity of nanopillars is important for achieving omnidirectional

anti-reflection in *G. oto*, we do not yet understand how the wax nanopillars are generated

to vary in height. Perhaps the pressure of the wax secretion varies across the microvillar

extensions' area, similar to how nozzle area plays a role in the propulsion force, and tunes

the height of the nanopillars in the process. In such a scenario, the degree of the height

variation could be synthetically engineered depending on the two-dimensional

nanopatterned mask design in the biomimetic processes, like molding or imprinting

techniques. Additionally, others have generated three-dimensional wax structures by

using *n*-alkanes, noting that wax-based crystals can generate different shapes, sizes and

densities depending on the chain length (*45*). Future work should investigate the possible

role of alkanes, and the two-dimensional surface growth geometry, in generating three-

dimensional anti-reflective nanostructures and potential applications for biomimetics. Our

exploration of *Greta oto* wing development can serve as a model for understanding how

transparent phenotypes evolved within Ithomiini, a diverse tribe of neotropical butterflies

that act as mimicry models for numerous species of Lepidoptera (*26*), as well as more

distantly related butterfly and moth species.

Materials and Methods

Samples

Glasswing butterfly (*Greta oto*) pre-pupae were purchased from Magic Wings Butterfly

House (Deerfield, Massachusetts, USA) and reared on *Cestrum nocturnum* (Solanaceae)

leaves at 27°C and 60% humidity on a 16:8 hour light:dark cycle at the Marine Biological

Laboratory (Woods Hole, MA) under the United States Department of Agriculture permit

number P526P-19-02269. At the appropriate time of development, pupal wings were

dissected and age was recorded as hours after pupal case formation (h APF) as in (*6*). The

average timeline from pupation to eclosion (adult emergence) for *G. oto* at 27°C is about

7 days, and we report our time series here which covers early aspects of wing scale

development.

Optical imaging and scale measurements

Images of whole mounted specimens were taken with a Canon EOS 70D digital camera

with an EF 100mm f/2.8L macro lens. High-magnification images of disarticulated wings

were taken with a Keyence VHX-5000 digital microscope. Scale density was determined

by counting the numbers of scales in a 1 mm^2 area. Scales were also removed from the

wings, laid flat onto a slide, and Keyence software was used to measure the surface area

of individual scales. Images of clear and opaque regions were processed with Keyence

software to measure the percentage of area covered by scales. Sample size was equal to

three individual butterflies reared in the same cohort, in which four measurements for

each individual were averaged. We performed Student's t-tests for scale density and

percent of exposed membrane, and one-way ANOVA test for scale surface area

comparisons.

Confocal microscopy

For confocal microscopy of fixed tissue, pupal wings were dissected and fixed in PIPES,

EGTA, MgSO$_4$ (PEM) buffer with 3.7% paraformaldehyde for 20-30 minutes at room

temperature, as described previously (*6*). Fixed wings were incubated in 1X PBS+0.1%

Triton-X 100 (PT) with 1:200 dilution of phalloidin, Alexa 555 conjugated (Invitrogen

A34055), and Wheat Germ Agglutinin, Alexa 647 conjugated (Invitrogen W32466) at a

dilution of 1:200 overnight at 4°C. Wings were washed in PT and then placed in 50%

glycerol:PBS with 1 μg/mL DAPI overnight at 4°C. Wing samples were placed on

microscope slides and mounted in 70% glycerol:PBS. A coverslip (#1.5 thickness) was

applied, and each preparation was sealed with nail polish. Slides of fixed tissue were

examined with an LSM 880 confocal microscope (Carl Zeiss, Germany) with 40x and

63x objectives. Confocal images and movies were generated using Imaris Image Analysis

Software (Bitplane, Oxford Instruments, UK).

Scanning electron microscopy

We cut 2mm square pieces from dry wings, coated them with a 10 nm layer of gold using

the BIO-RAD E5400 Sputter Coater, and imaged with a Hitachi TM-1000 SEM at 5 kV.

Top-view and cross section SEM images were analysed with ImageJ 1.52 to measure

membrane thickness and nanostructure dimensions (n = 6).

Transmission electron microscopy

For transmission electron microscopy, wings of *Greta oto* pupae were dissected and fixed

in 2% glutaraldehyde, 2% paraformaldehyde in 0.1 M sodium cacodylate buffer overnight

at 4°C (pH 7.4). Samples were then rinsed in 0.1 M cacodylate buffer (pH 7.4) and post-

fixed in1% aqueous osmium tetroxide in 0.1M cacodylic buffer overnight at 4°C, then

rinsed in water. Samples were en bloc stained with 1% uranyl acetate in water and then

rinsed in water. Samples were dehydrated through a graded ethanol series (50–100% in

10% steps), rinsed in propylene oxide, then infiltrated in 50% resin and propylene oxide

overnight. Samples were infiltrated with Epon/Alardite embedding medium (70%, 80%,

95% to 100% steps) and polymerized at 60°C for two days. Thin sections (~70nm) were

cut on an Ultramicrotome RMC PowerTome XL using a Diatome diamond knife. Digital

images were taken using a JEOL 200 transmission electron microscope (Jeol, USA).

Wing surface wax extraction and analysis

To identify the molecular composition of the transparent wing surface, we pooled wing

dissections from three individual adults and performed two replicates for chloroform-

based extractions and two replicates for hexane-based extractions (after *26*). First, the

samples were soaked with 100 μL of either hexane or chloroform and gently mixed for 15

minutes on a Thermolyne RotoMix 51300. The liquid solutions containing dissolved wing

surface compounds were then transferred to glass vials with fixed microvolume inserts

and the solvent was evaporated under a stream of high-purity nitrogen gas (99.99%).

Dried extracts were re-dissolved in fixed volumes of hexane (10 μL), and half of the

extract (5 μl) was injected by automatic liquid sampler into a gas chromatograph coupled

with a mass selective detector (GC: 7890A; MS: 5975C; Agilent Technologies, USA)

operating in electron impact mode. The injection was performed in a split/splitless

injector in the splitless mode. Separation of compounds was performed on a fused silica

capillary column (DB-5MS, 30 m $\times$ 0.32 mm $\times$ 0.25 μm, Agilent J&W GC columns,

USA) with a temperature program starting from 80°C for 5 min and increasing by 80°C

per min to 200 °C, followed by an increase of 5 °C per min to 325 °C which was held for

3 min, with helium used as the carrier gas, positive electron ionization (70 eV), Analog to

Digital (A/D) sampling rate was set at 4, and the scan range was m/z 40.0 to 650.0.

Chemical data processing was carried out using the software "Enhanced Chemstation"

(Agilent Technologies, USA). We retained peaks with abundances greater than 0.25% of

the total and compounds were identified according to their retention indices, diagnostic

ions, and mass spectra, which are provided in Table S1. For some peaks, it was not

possible to narrow the identity to a single specific compound because (1) some low

abundance substances produced poor quality mass spectra, (2) multiple compounds could

have produced the observed fragmentation patterns and/or (3) multiple compounds may

have co-eluted at the same retention time.

Optical measurements

The wing reflection measurements were performed on a Cary 5000 UV-Vis-NIR

spectrophotometer, equipped with a light source of tungsten halogen and an integrating

sphere diffuse reflectance accessory (Internal DRA 1800). Wing measurements from the

dorsal wing surface (n = 6) were recorded with unpolarized light with a spot size of 100

μm for an incident angle of 8° to avoid the loss of direct specular reflectance component

through the aperture. All measurements were taken in the dark to avoid possible stray

illumination from the surrounding environment. A reference measurement was done with

a calibrated commercial white spectralon standard to calculate the relative diffuse

reflectance. The reflectance measurements and mean data are presented in Table S2.

Optical simulations

The reflectance of the wing membrane before and after chemical treatment by hexane was

analytically modeled using effective medium theory and transfer matrix method (*10, 18*).

First, the effective volume fraction of the nanoprotuberances before and after the chemical

treatment were based on measurements taken from SEM micrographs of the wings. We

used the average distance between two hexagonally arranged nanostructures, *d*, conical

shaped nipple nanostructures with height, h_{p}, wax-based irregular nanopillars with radius,

r_{np}, mean height, h_{np} and variance σ_{np}, and membrane thickness, h_{m} and variance σ_{m} (fig.

S2). We considered a Gaussian distribution of irregular nanopillar height, as described

previously (*10*). We also modeled the membrane thickness with Gaussian distribution to

replicate the experimental membrane modulation in the calculation (*44, 46*). The total

volume fraction of the untreated wing along the height *h* can be given by:

$$\frac{\pi r_{np}^2}{\sqrt{3}d^2}\text{erfc}\left(\frac{h-h_{np}}{\sigma_{np}\sqrt{2}}\right);$$ zone: dorsal wax based nanopillar

$$\frac{\pi}{\sqrt{3}d^2}\left(r_{np}+\left(\frac{d}{2}-r_{np}\right)\left(1-\frac{h}{h_p}\right)\right)^2;$$ zone: dorsal chitin-based nipple array

$f_{untreated}(h)=$ 1;   zone: chitin membrane

$$\frac{\pi}{\sqrt{3}d^2}\left(r_{np}+\left(\frac{d}{2}-r_{np}\right)\left(1-\frac{h}{h_p}\right)\right)^2;$$ zone: ventral chitin-based nipple array

$$\frac{\pi r_{np}^2}{\sqrt{3}d^2}\text{erfc}\left(\frac{h-h_{np}}{\sigma_{np}\sqrt{2}}\right);$$ zone: ventral wax based nanopillar

611   where, $\text{erfc}(x)=\frac{2}{\sqrt{\pi}}\int_x^\infty e^{-t^2}dt$ is the complementary error function.

612   The volume fraction of the treated wing without the irregular nanopillars will be:

$$\frac{\pi}{2\sqrt{3}}\left(1-\frac{h}{h_p}\right);$$ zone: dorsal chitin-based nipple array

$f_{treated}(h)=$ 1;   zone: chitin membrane

$$\frac{\pi}{2\sqrt{3}}\left(1-\frac{h}{h_p}\right);$$ zone: ventral chitin-based nipple array

After determining the volume fraction, the corresponding refractive index changes along

the wing at any height h was calculated using the effective medium theory with the

Maxwell-Garnett approximation as shown in Fig. 6E, fig. S2. The refractive indices of the

different materials were considered as $n_{air}=1$, $n_{chitin}=1.56+i0.008$ (20, 21) and we

considered $n_{wax}=1.39$ (based on 47). Afterwards, the transfer matrix method computed

the reflectance from the stratified medium with calculated refractive index profiles as

shown in Fig. 6E for the unpolarized condition (taking the average of both s- and p-

polarization) at an incident angle of 8°. The membrane-only reflection at normal incident

light can be directly calculated from (46):

$$R\,(\lambda)=\int_0^\infty \left|\frac{r(1-e^{-2i\delta})}{1-r^2 e^{-2i\delta}}\right|^2 \frac{1}{\sigma_m\sqrt{2\pi}}e^{\frac{-(h-h_m)^2}{2\sigma_m^2}}\,dh.$$

623   Where, $\delta = (2\pi n_{chitin} h)/\lambda$ is the phase delay introduced by the membrane thickness of h,

624   and r is the reflection coefficient at the air-chitin boundary governed by Fresnel's

625   equation for a normal incident light, i.e., $r = (1 - n_{chitin}) / (1 + n_{chitin})$.

References and Notes

1. P. Beldade, P. M. Brakefield. The genetics and evo–devo of butterfly wing patterns. *Nature Reviews Genetics*. **3**, 442-52 (2002).

2. R. O. Prum, T. Quinn, R. H. Torres. Anatomically diverse butterfly scales all produce structural colours by coherent scattering. *Journal of Experimental Biology*. **209**, 748-65. (2006).

3. L. E. Gilbert, M. C. Singer. Butterfly ecology. *Ann. Rev. Ecol. Syst.* **6**, 365-95 (1975).

4. D. G. Stavenga, H. L. Leertouwer, B. D. Wilts. Coloration principles of nymphaline butterflies–thin films, melanin, ommochromes and wing scale stacking. *J. Exper. Biol.* **217**, 2171-80 (2014).

5. R. C. Thayer, F. I. Allen, N. H. Patel. Structural color in *Junonia* butterflies evolves by tuning scale lamina thickness. *Elife*. **9**, e52187 (2020).

6. A. Dinwiddie, R. Null, M. Pizzano, L. Chuong, A. L. Krup, H. E. Tan, N. H. Patel. Dynamics of F-actin prefigure the structure of butterfly wing scales. *Develop. biol.* **392**, 404-18 (2014).

7. A. R. Day, J. J. Hanly, A. Ren, A. Martin. Sub-micrometer insights into the cytoskeletal dynamics and ultrastructural diversity of butterfly wing scales. *Developmental Dynamics*. **248**, 657-70 (2019).

8. P. P. Goodwyn, Y. Maezono, N. Hosoda, K. Fujisaki. Waterproof and translucent wings at the same time: problems and solutions in butterflies. *Naturwissenschaften*. **96**, 781-7 (2009).

9. A. Yoshida, M. Motoyama, A. Kosaku, K. Miyamoto. Antireflective nanoprotuberance array in the transparent wing of a hawkmoth, *Cephonodes hylas*. *Zoological sci*. **14**, 737-41 (1997).

10. R. H. Siddique, G. Gomard, H. Hölscher. The role of random nanostructures for the omnidirectional anti-reflection properties of the glasswing butterfly. *Nature communications*. 6, 1-8 (2015).

11. M. Arias, J. Mappes, C. Desbois, S. Gordon, M. McClure, M. Elias, O. Nokelainen, D. Gomez. Transparency reduces predator detection in mimetic clearwing butterflies. *Func. Ecol.* **33**, 1110-9 (2019).

12. M. Arias, M. Elias, C. Andraud, S. Berthier, D. Gomez. Transparency improves concealment in cryptically coloured moths. *J. Evol. Biol.* **33**, 247-52 (2020).

13. M. Mcclure, C. Clerc, C. Desbois, A. Meichanetzoglou, M. Cau, L. Bastin-Héline, J. Bacigalupo, C. Houssin, C. Pinna, B. Nay, V. Llaurens. Why has transparency evolved in aposematic butterflies? Insights from the largest radiation of aposematic butterflies, the Ithomiini. *Proc. Royal Soc. B*. **286**, 20182769 (2019).

14. S. Johnsen. Hidden in plain sight: the ecology and physiology of organismal transparency. *The Biological Bulletin*. **201**, 301-18 (2001).

15. S. Johnsen. Hide and seek in the open sea: pelagic camouflage and visual countermeasures. *Annual review of marine science*. **6**, 369-92 (2014).

16. L. E. Bagge. Not as clear as it may appear: challenges associated with transparent camouflage in the ocean. *Integ. compar. biol*. **59**,1653-63 (2019).

17. C. G. Bernhard, W. H. Miller. A corneal nipple pattern in insect compound eyes. *Acta Physiologica Scandinavica*. **56**, 385-6 (1962).

18. D. G. Stavenga, S. Foletti, G. Palasantzas, K. Arikawa. Light on the moth-eye corneal nipple array of butterflies. *Proc. Royal Soc. B: Biological Sciences*. **273**, 661-7 (2006).

19. Y. F. Huang, Y. J. Jen, L. C. Chen, K. H. Chen, S. Chattopadhyay. Design for approaching cicada-wing reflectance in low-and high-index biomimetic nanostructures. *ACS nano*. **9**, 301-11 (2015).

20. P. Vukusic, J. R. Sambles, C. R. Lawrence, R. J. Wootton. Quantified interference and diffraction in single Morpho butterfly scales. *Proc. Royal Soc. London. Series B: Biol. Sci*. **266**, 1403-11 (1999).

21. V. Narasimhan, R. H. Siddique, J. O. Lee, S. Kumar, B. Ndjamen, J. Du, N. Hong, D. Sretavan, H. Choo. Multifunctional biophotonic nanostructures inspired by the longtail glasswing butterfly for medical devices. *Nature nanotechnology*. **13**, 512-9 (2018).

22. O. Deparis, S. R. Mouchet, L. Dellieu, J. F. Colomer, M. Sarrazin. Nanostructured surfaces: bioinspiration for transparency, coloration and wettability. *Mater. Today Proc. S*. **1**, 122-9 (2014).

23. V. R. Binetti, J. D. Schiffman, O. D. Leaffer, J. E. Spanier, C. L. Schauer. The natural transparency and piezoelectric response of the *Greta oto* butterfly wing. *Integr. Biol*. **1**, 324-9 (2009).

24. D. Gomez, C. Pinna, J. Pairraire, M. Arias, J. Barbut, A. Pomerantz, C. Nous, W. D. de Marcillac, S. Berthier, N. Patel, C. Andraud, M. Elias. Transparency in butterflies

and moths: structural diversity, optical properties and ecological relevance. *bioRxiv*. doi: https://doi.org/10.1101/2020.05.14.093450 (2020).

25. C. Pinna, M. Vilbert, S. Borenztajn, W. D. de Marcillac, F. Piron-Prunier, A. Pomerantz, N. Patel, S. Berthier, C. Andraud, D. Gomez, M. Elias. Convergence in light transmission properties of transparent wing areas in clearwing mimetic butterflies *bioRxiv*. doi: https://doi.org/10.1101/2020.06.30.180612 (2020).

26. M. Elias, Z. Gompert, C. Jiggins, K. Willmott. Mutualistic interactions drive ecological niche convergence in a diverse butterfly community. *PLoS biology*. **6**, e300 (2008).

27. R. Futahashi, Y. Yamahama, M. Kawaguchi, N. Mori, D. Ishii, G. Okude, Y. Hirai, R. Kawahara-Miki, K. Yoshitake, S. Yajima, T. Hariyama. Molecular basis of wax-based color change and UV reflection in dragonflies. *Elife*. **8**, e43045 (2019).

28. H. T. Ghiradella, M. W. Butler. Many variations on a few themes: a broader look at development of iridescent scales (and feathers). *J. Royal Soc. Interface*. Apr **6**, S243-51 (2009).

29. R. D. Reed. Evidence for Notch-mediated lateral inhibition in organizing butterfly wing scales. *Development Genes and Evolution*. **214**, 43-6 (2004).

30. R. Galant, J. B. Skeath, S. Paddock, D. L. Lewis, S. B. Carroll. Expression pattern of a butterfly achaete-scute homolog reveals the homology of butterfly wing scales and insect sensory bristles. *Current Biology*. **8**, 807-13 (1998).

31. I. G. Tilney, P. S. Connelly, K. A. Vranich, M. K. Shaw, G. M. Guild. Actin filaments and microtubules play different roles during bristle elongation in *Drosophila*. *J. Cell. Sci*. **113**, 1255-65 (2000).

32. S. Djokic, B. Anna, I. Zurim, N. Urakova, J. Ragson, U. Abdu. Actin bundles play different role in shaping scale as compare to bristle in mosquito *Aedes aegypti*. *bioRxiv*. doi: https://doi.org/10.1101/2020.04.06.027110 (2020).

33. I. Overton. Microtubules and microfibrils in morphogenesis of the scale cells of *Ephestia kuhniella*. *J. cell biology*. **29**, 293-305 (1966).

34. A. Bitan, G. M. Guild, D. Bar-Dubin, U. Abdu. Asymmetric microtubule function is an essential requirement for polarized organization of the Drosophila bristle. *Molec. cell. biol*. **30**, 496-507 (2010).

35. A. Bitan, I. Rosenbaum, U. Abdu. Stable and dynamic microtubules coordinately determine and maintain *Drosophila* bristle shape. *Development*. **139**, 1987-96 (2012).

36. G. Gemne. Ontogenesis of corneal surface ultrastructure in nocturnal Lepidoptera. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*. **262**, 343-63 (1971).

37. A. Fröhlich. A scanning electron-microscopic study of apical contacts in the eye during postembryonic development of *Drosophila melanogaster*. *Cell and tissue research*. **303**, 117-28 (2001).

38. M. Kryuchkov, A. Blagodatski, V. Cherepanov, V. L. Katanaev. "Arthropod corneal nanocoatings: diversity, mechanisms, and functions" in Functional Surfaces in Biology III. (Springer, Cham, Switzerland, 2017), pp. 29-52.

39. B. Moussian. Recent advances in understanding mechanisms of insect cuticle differentiation. *Insect biochem. molec. biol*. **40**, 363-75 (2010).

40. A. G. Gibbs. Water-proofing properties of cuticular lipids. American Zoologist. **38**, 471-82 (1998).

41. I. Román-Kustas, J. B. Hoffman, J. H. Reed, A. E. Gonsalves, J. Oh, L. Li, S. Hong, K. D. Jo, C. E. Dana, N. Miljkovic, D. M. Cropek. Molecular and Topographical Organization: Influence on Cicada Wing Wettability and Bactericidal Properties. *Advanced Materials Interfaces*. **7**, 2000112 (2020).

42. A. P. Ivanova, S. H. Nguyen, H. K. Webb, J. Hasan, V. K. Truong, R. N. Lamb, X. Duan, M. J. Tobin, P. J. Mahon, R. J. Crawford. Molecular organization of the nanoscale surface structures of the dragonfly *Hemianax papuensis* wing epicuticle. *PLoS One*. **8**, e67893. (2013).

43. A. Shevtsova, C. Hansson, D. H. Janzen, J. Kjærandsen. Stable structural color patterns displayed on transparent insect wings. *Proc. Natl. Acad. Sci*. **108**, 668-73 (2011).

44. D. G. Stavenga. Thin film and multilayer optics cause structural colors of many insects and birds. *Materials Today: Proceedings*. **1**, 109-21 (2014).

45. A. Gorb, S. Böhm, N. Jacky, L. P. Maier, K. Dening, S. Pechook, B. Pokroy, S. Gorb. Insect attachment on crystalline bioinspired wax surfaces formed by alkanes of varying chain lengths. *Beilstein J. Nanotech*. **5**, 1031-41 (2014).

46. R. H. Siddique, S. Vignolini, C. Bartels, I. Wacker, H. Hölscher. Colour formation on the wings of the butterfly *Hypolimnas salmacis* by scale stacking. *Scientific reports*. **6**, 36204 (2016).

47. I. R. Hooper, P. Vukusic, R. J. Wootton. Detailed optical study of the transparent wing membranes of the dragonfly *Aeshna cyanea*. *Optics express*. **14**, 4891-7 (2006).

Acknowledgments

We would like to thank the Angie Serrano, Paola Betucci, Idoia Quintana-Urzainqui and Helena Bilandzija from the MBL Embryology Course and Cao Lu Yan from the MBL Physiology Course for their preliminary work on scale morphology of clearwing Lepidoptera, and subsequent work by Jaap van Krugten, Raymundo Picos, and Johnny On at UC Berkeley. We also thank Rachel Thayer, Arnaud Martin, Damien Gailly, Melanie Mcclure, Luca Livraghi, Oscar Paneso, Rémi Mauxion and Owen McMillan for their assistance with fieldwork, rearing, and preliminary experiments. We thank Fred Gagnon of Magic Wings Butterfly Conservatory and Gardens for assistance with butterfly rearing and Neil Tsutsui for support with the GC-MS experiments. Members of the Patel Lab, Craig Miller, and Noah Whiteman provided helpful feedback on the manuscript. RHS acknowledges the support from the Beckman Institute of the California Institute of Technology to the Molecular Materials Research Center. **Funding:** This work was supported by a grant from the Human Frontier Science Program (RGP0014/2016), a France-Berkeley fund grant (FBF #2015--58) and an ANR grant (CLEARWING project, ANR-16-CE02-0012). **Author contributions:** Conceived and designed the experiments: AFP and NHP. Performed the experiments: AFP, RHS, EIC, KH. All authors contributed to analysis of the results, contributed reagents/materials/analysis tools and contributed to writing the paper. **Competing interests:** The authors declare they have no competing interests.

Fig. 1. Examples of clearwing butterflies and wing scale features in *Greta oto*

(**A**) Giant glasswing *Methona confusa* (Nymphalidae: Ithomiini). Scale bar = 1

cm. Wings under (**B**) reflected and (**B'**) transmitted light illustrating general

transparency, but strong light reflectance off the wing surface in this species. (**C**)

The surface of the wing membrane is smooth and devoid of nanostructures. Scale bar = 1 μm. **(D)** Longtail glasswing *Chorinea faunus* (Riodinidae). Scale bar = 1 cm. Wings under **(E)** reflected and **(E')** transmitted light illustrating minimal reflectance. **(F)** The membrane surface contains dome-shaped chitin nanoprotuberances that generate anti-reflective properties (*21*). Scale bar = 1 μm. **(G)** Glasswing *Greta oto* (Nymphalidae: Ithomiini). Red arrowhead indicates the representative clear and opaque wing region investigated, scale bar = 1 cm. Wings under **(H)** reflected and **(H')** transmitted light illustrating minimal reflectance. **(I)** The surface of the wing membrane contains irregularly sized nanopillars that enable omnidirectional anti-reflective properties (*10*). Scale bar = 1 μm. **(J)** High magnification of a transition boundary between a clear (left side) and opaque (upper right side) wing region in *G. oto*. Scale bar = 100 μm. **(K)** SEM of adult scales in a clear wing region of *G. oto*, revealing the alternating forked (green false coloring) and bristle-like (red false coloring) scale morphologies (socket false colored in blue). Scale bar = 20 μm. **(L)** SEM of scales in an opaque wing region in *G. oto*, highlighting typical large, flat scale morphologies. Scale bar = 20 μm. **(M)** Measurements of scale density in clear and opaque wing regions, **(N)** scale surface area for forked, bristle-like, and opaque scale morphologies, and **(O)** percent of wing membrane exposed in *G. oto* clear and opaque regions. Error bars indicate means + SD of four measurements taken from wings in three different individuals, P-values are based on Student's t-test for **(M)**, **(O)**, and ANOVA test for **(N)**, ***$P < 0.001$; **$P < 0.01$.

Fig. 2. Pupal wing development and cytoskeletal organization of scales in clear and opaque regions

(A) Representative image of a *Greta oto* pupa ~5 days after pupal formation (APF), **(A'-A'')** developing up to the melanic stage ~7 days APF, just prior to eclosion. **(B)** Early wing development 16 hours APF stained with DAPI (nuclei) in a clear wing region and **(C)** opaque wing region. The clear region contains a reduced number of sensory organ precursor (SOP) cells (which are the precursor cells to the scale and socket cells) relative to the opaque region. Scale bar = 20 μm. SOP cells are false colored magenta for better viewing. Simultaneous confocal imaging of fluorescently labeled scale cell membrane (wheat germ agglutinin; WGA, magenta), and F-actin (phalloidin, green), comparing clear

wing regions **(D, F, H)** to opaque wing regions **(E, G, I)**. At 30 h AFP **(D, E)**

WGA and phalloidin staining reveal early scale buds extending from the wing

epithelium. F-actin reveals loosely organized parallel actin filaments protruding

from the membrane. 48 h APF **(F, G)** scales have grown and changed in

morphology. Short actin filaments have reorganized and formed smaller numbers

of thick, regularly spaced parallel bundles under the surface of the cell membrane.

(F) In the clear wing region, scale cells alternate between triangular shapes and

bristles. 60 h APF **(H, I)**, developing scales have become more elongated. **(H)** The

triangular-shaped scales in the clear wing region have proceeded to generate two

new branches, which fork and elongate bidirectionally, while bristle-like scales

have rapidly elongated and curved. **(I)** In the opaque region, scales are longer,

wider, and have now developed serrations at the tips. Scale bar in **(D-I)** = 10 μm.

Fig. 3. Confocal and transmission electron microscopy transverse sections of

developing bristle (top), forked (middle) and flat (bottom) scales 48 hours APF

(A) Confocal projection of a bristle-like scale morphology (phalloidin) in a clear wing region. White arrowhead shows representative regions of transverse TEM sections. Scale bar = 5 μm. TEM of a bristle-like scale in a distal region (B-B') and a basal region near the socket cell (C-C'). Note the peripheral actin bundles (false colored green) and internal microtubule rings (false colored magenta). The more distal region of the scale (B) contains a lower density of microtubules relative to the base of the scale (C). Scale bars in (B,C) = 500 nm and scale bars in (B',C') = 100 nm. (D) Confocal projection of a developing forked scale (phalloidin) in a clear wing region. White arrowhead shows representative regions of transverse TEM sections. Scale bar = 5 μm. (E-E') TEM of a forked scale reveals peripheral bundles of actin (false colored green), with thicker actin bundles on the ventral side of the scale and internal microtubules (false colored magenta). Two internal bundles of actin filaments can be observed in the cytoplasm (E'). Scale bars in (E-E') = 500 nm. (F) Confocal projections of developing flat, round scale (phalloidin) in an opaque wing region. White arrowhead shows representative regions of transverse TEM sections. Scale bar = 5 μm. (G-G') TEM reveals asymmetry in the actin bundles (false colored green), which are thicker on the bottom side of the scale relative to the upper surface. Microtubules (false colored magenta) are found in various orientations. Scale bar in (G-G') = 500 nm.

Fig. 4. Ontogeny of wing membrane surface nanostructures

(A) SEM cross section (side view) of an adult *Greta oto* clear wing region. Scale bar = 10 µm. Bristle-like scale false colored in red, forked scale false colored in green, sockets false colored in blue. **(B)** TEM transverse section of epithelial tissue 60 h APF, showing lateral scale growth and wing membrane cells. Scale bar = 2 µm. **(C)** Higher magnification of developing wing epithelial cells at 60 h APF show microvilli (MV) projections, which appear as slender linear extensions from the inner margins of the developing cells that insert into a thin layer of electron-dense material. Lamina evaginations appear in the section as domes. **(D)** TEM of epithelial tissue 72 h APF and **(E)** 120 h APF shows wing surface nanostructures protruding from the surface, with tips of microvilli still attached to the inner surface of the wing membrane. **(F)** TEM of the adult wing membrane. The surface contains dome-shaped nipple nanostructures and an upper layer of nanopillars. Scale bar in **(C-E)** = 500 nm.

Fig. 5. Topographical organization and biochemical composition of wing surface

nanostructures

Scanning electron microscopy of the transparent wing membrane surface of *Greta oto*

under (A-A') untreated condition, highlighting the presence of irregularly arranged

nanopillar structures covering the surface, (B-B') physical treated condition, revealing

partial removal of surface nanopillars, and a lower layer of more regularly arranged

nipple-like nanostructures and (C) chloroform treated condition, revealing complete

removal of the upper layer of nanopillars, and remining lower layer of nipple-like

nanostructures. Scale bars in (A, B C) = 2 μm, scale bars in (A', B', C') = 1 μm. (D)

879   Chromatogram of hexane-treated (top; red line) and chloroform-treated (bottom; black

880   line) clearwing extracts. X-axis shows the retention time in minutes and Y-axis shows the

881   abundance of total ion current. **(E)** Schematic of proposed wing surface membrane

882   nanostructures in *Greta oto*, composed of chitin-based procuticle and wax-based

883   epicuticle.

Fig. 6. Structural elements, reflectance spectra and optical modeling of anti-reflective nanostructures

Optical images and cross section scanning electron microscopy of *Greta oto* **(A, B)**

untreated wings, illustrating low reflectance and the presence of nanopillars on the wing

membrane surface and **(C, D)** hexane-treated wings illustrating increased reflectance and

the loss of nanopillars on the wing membrane, but presence of nipple-like nanostructures

on the surface. Scale bars for **(B, D)** = 200 nm. **(E)** Optical modeling of effective

refractive index conditions for untreated (top), with nanopillars of variable height together

with cuticle-based nipple nanostructures on the wing membrane, treated (middle) with

cuticle-based nipple nanostructures on wing membrane and wing membrane without any

nanostructures (bottom). Y-axis represents height h and X-axis represents effective

refractive index condition of air (n_{air}), chitin (n_{chitin}), and wax (n_{wax}). **(F)** Representative

reflectance spectra of experimental (red) and simulation data (black) for untreated wings

with nanopillars on the membrane surface (solid line), hexane-treated wings with the

wax-based layer of nanopillars removed (dashed line) and membrane only (dotted line).

Supplementary Materials

Table S1. GC-MS relative proportions (mean ± standard deviation) of wing cuticular compounds isolated from *Greta oto*.

Table S2. Spectrometry of *Greta oto* untreated and hexane treated clear wing regions and simulated reflectance spectra.

Movie S1. 3D projection of developing scales in a clear wing region 48 hours after pupal formation. 3D projection and rotation of the same scales shown in Fig. 2F, 48 hours APF in a clear wing region. WGA (magenta) stains cell membranes and phalloidin (green) stains F-actin and DAPI (blue) stains nuclei. Short actin filaments have reorganized and formed smaller numbers of thick, regularly spaced parallel bundles just under the surface of the cell membrane. Scales alternate with future forked scales appearing as triangular shapes and longer future bristle-like shapes.

Fig. S1. TEM micrographs of scales 72 hours (top) and 120 hours (bottom) after

pupal formation

(**A**) TEM micrograph of a developing opaque scale 72 h APF, highlighting microtubule

arrangement (MT). (**B**) Thick actin bundles contain dense, hexagonally packed F-actin

filaments. (**C**) Basal region of a developing scale outgrowth and socket cell. Developing

scales 72 h APF contain dense populations of microtubules (MT) and numerous internal

organelles, including mitochondria (M), electron dense vesicles and free single

ribosomes. (**D**) Transverse section of developing scales around 120 h APF, highlighting

both flat and thin, bristle-like scale morphologies. Cross section near the (**E**) base and (**F**)

distal region of scales 120 h APF, showing thickened cuticle and ridge morphologies.

Fig. S2. Optical modeling parameters and effective refractive index condition for

untreated transparent wing of *Greta oto*

Schematic representation for the optical modeling parameters of wing membrane and

surface nanostructures. Average distance between two nanostructures represented as d,

conical shaped cuticular nipple nanostructures height as h_p, wax-based irregular

nanopillars radius as r_np, mean height as h_np and variance σ_np, and membrane thickness as

h_m and variance σ_m. Y-axis represents height h and X-axis represents effective refractive

index condition of air (n_air), chitin (n_chitin), and wax (n_wax).

Fig. S3. Optical simulations for mean membrane thickness and modulation of thickness under different wing architecture models

Simulation reflectance spectra of (**A**) Membrane only (lacking surface nanostructures) with varying mean membrane thickness. (**B**) Treated wings (containing cuticle-based nipple nanostructures but lacking wax-based irregular nanopillars) with varying mean membrane thickness. (**C**) Untreated wings (containing wax-based irregular nanopillars and nipple nanostructures) with varying mean membrane thickness and no modulation in thickness. (**D**) Untreated wings with variable mean membrane thickness and modulation of 43 nm variance in thickness.