The impact of allometry on vomer shape and its implications for the taxonomy and

cranial kinesis of crown-group birds

Olivia Plateau[1]*, Christian Foth[1]

[1]Department of Geosciences, University of Fribourg, Chemin du Musée 6, CH-1700

Fribourg, Switzerland

*e-mail: olivia.plateau@unifr.ch

O.P. (ORCID: 0000-0002-8321-2687)

C.F. (ORCID: 0000-0002-9410-4569)

Abstract

Crown birds are subdivided into two main groups, Palaeognathae and Neognathae, that can

be distinguished, among other means, by the organization of the bones in their pterygoid-

palatine complex (PPC). Shape variation of the vomer, which is the most anterior part of the

PPC, was recently analysed with help of geometric morphometrics to discover morphological

differences between palaeognath and neognath birds. Based on this study, the vomer was

identified as sufficient to distinguish the two main groups (and even some inclusive neognath

groups) and their cranial kinetic system. As there are notable size differences between the

skulls of Palaeognathae and Neognathae, we here investigate the impact of allometry on

vomeral shape and its implication for taxonomic classification by re-analysing the data of the

previous study. Different types of multivariate statistical analyses reveal that taxonomic

identification based on vomeral shape is strongly impaired by allometry, as the error of

correct identification is high when shape data is corrected for size. This finding is evidenced

by a great overlap between palaeognath and neognath subclades in morphospace. Correct

taxonomic identification is further impeded by the convergent presence of a flattened vomeral

morphotype in multiple neognath subclades. As the evolution of cranial kinesis has been

linked to vomeral shape in the original study, the correlation between shape and size of the

vomer across different bird groups found in the present study questions this conclusion. In

fact, cranial kinesis in crown birds results from the loss of the jugal-postorbital bar in the

temporal region and ectopterygoid in the PPC and the combination of a mobilized quadrate-

zygomatic arch complex and a flexible PPC. Therefore, we can conclude that vomer shape

itself is not a suitable proxy for exploring the evolution of cranial kinesis in crown birds and

their ancestors. In contrast, the evolution of cranial kinesis needs to be viewed in context of

the braincase, quadrate-zygomatic arch and the whole pterygoid-palatine complex.

Introduction

The pterygoid-palatine complex (PPC) of crown birds is mainly formed by five bones: the

unpaired vomer that results from the fusion of the originally paired vomer elements, and the

paired pterygoids and palatines. The general morphology of the PPC was first studied by

Huxley (1867), who distinguished the clade Palaeognathae from all other birds on the basis of

palatal morphology. Although the PPC of Palaeognathae is quite variable (McDowell 1948),

it is characterized by a large vomer that is only partly fused. The pterygoids and palatines are

highly connected, forming a rigid unit that articulates with the braincase via well-developed

basipterygoid processes, while a contact with the parasphenoid is not present (see Bellairs &

Jenkin 1960; Zusi 1993; Gussekloo et al. 2001, Mayr 2017; Fig. 1A). In contrast, neognath

birds possess a movable joint between pterygoid and palatine, which plays an important role

in the kinematic movement of the upper jaw. Here, the pterygoid articulates with the

parasphenoid, while the basipterygoid processes are often reduced. The vomer is highly

variable in size and shape and often has no connection with the upper jaw beyond an

association with the nasal septum and the palatine. In some neognaths, the vomer is greatly

reduced or even absent (see Bellairs & Jenkin 1960; Bock 1964; Zusi 1993; Mayr 2017, Fig.

1A).

In a recent paper, Hu et al. (2019) investigated palate evolution in crown birds and

their stem, focusing on the morphology of the vomer. Using 3D geometric morphometrics,

the study found that the vomeral shape of neognaths is clearly distinguishable from

Palaeognathae, in that the latter group has a stronger similarity with their non-avian

ancestors. Linking vomer shape with the kinetic abilities of the skull, the authors concluded

that cranial kinesis represents an innovation of Neognathae. Furthermore, the authors

concluded that vomeral morphology allows for a taxonomic differentiation between the major

groups of Neognathae, namely Aequorlitornithes, Galloanserae, Gruiformes, and Inopinaves.

However, according to their PCA results, all groups strongly overlap each other within PC1,

while a taxonomic differentiation is only noticeable within PC2 (other principal components

are not shown). Taking the great size variation of the vomer of neognath birds into account

(Zusi 1993), we wonder if the reported taxonomic differentiation between palaeognaths and

the neognath subclades could alternatively be related to allometry, i.e. the dependence of

shape on size (Klingenberg 1998), rather than pure shape variation. In order to test this

hypothesis, we re-analysed the dataset of Hu et al. (2019), comparing allometric shape data

with non-allometric residuals, and re-evaluating the role of the vomer in the evolution of

cranial kinesis in crown birds.

Materials and Methods

The published 3D models and landmarks data of 41 specimens including 36 species were

downloaded from Hu et al. 2019 (https://doi.org/10.6084/m9.figshare.7769279.v2). This

dataset contains five extinct species (two stem line representatives: the troodontid

Sinovenator changii, the Avialae *Sapeornis chaoyangensis*; and three fossil palaeognath

crown birds from the clade Dinornithiformes: *Pachyornis australis*, *Megalapteryx didinus*

and *Diornis robustus*), five extant Paleognathae and 27 extant Neognathae representing the

two major clades of crown birds.

The original landmark data (Dataset A) is composed of five anatomical landmarks and

43 semi-landmarks (see Hu et al. 2019). The landmark data were imported into the software

R v.3.5.2 (R Core Team, 2018). Using the *plotAllSpecimens* function of *Geomorph v.3.2.1*

(Adams *et al*. 2013) in *R*, we notice great variability for each anatomical landmark, resulting

from two main shapes of the vomer. The majority of bird possesses a fused vomer that is

bilaterally symmetric and roof-shaped in transection, with a horizontal orientation within the

pterygoid-palatine complex (Fig. 1B). In contrast, some members of Aequorlitornithes (e.g.,

Podiceps nigricollis, and *Podilymbus podiceps*), Galloanserae (e.g., *Anas crecca*, *Anseranas*

semipalmata, and *Cairina moschata*) and Inopinaves (e.g., *Aquila audax*, *Falco cenchroides*,

and *Haliastur sphenurus*) possess a fused vomer that is completely mediolaterally flattened in

transection and vertically orientated within the pterygoid-palatine complex (Fig. 1B).

Therefore, we created a second dataset (Dataset B), where species with this latter, flat vomer

morphology were excluded. Furthermore, the palaeognath birds *Struthio camelus* and

Dromaieus novaehollandiae of the original Dataset A were represented by both juvenile and

adult specimens. Because ontogenetic variation could, however, potentially affect size and

position of the palaeognath morphospace, we removed the juvenile and subadult specimens

of *Struthio camelus* and *Dromaieus novaehollandiae* in order to reran the analysis just with

adult semaphoronts (Dataset C). Finally, we created a fourth dataset, where both

juvenile/subadult specimens and species with flat vomers were removed from the sample

(Dataset D).

For superposition of the 3D landmark data, we followed Hu et al. (2019) by performing a *Generalized Procrustes Analysis* (*GPA*). The *GPA* was done with the help of the *gpagen* function in *Geomorph*. Afterward, we performed a *principal component analysis* (*PCA*) in order to visualize the shape variability of the vomer and the variance of morphospace for two groupings: (1) Paleognathae versus Neoganthae and (2) Paleognathae, Inopinaves, Galloanserae, Gruiformes and Aequorlitornithes. This was done with the *plotTangentSpace* function from *Geomorph*.

Because the vomer showed great variation in centroid size after superimposition, ranging from 14.60 (*Manorina melanocephala*) to 168.32 (*Dromaieus novehollandia*), we tested if there is a significant correlation between Procrustes coordinates and log-transformed centroid size (Goodall 1991) using the function *procD.lm* in *Geomorph*. This function performs a multivariate regression between the shape and size with a permutation of 10,000 iterations. A significant relationship between both parameters indicates that the superimposed shape still contains an allometric signal. Based on this correlation we estimated non-allometric residuals of the Procrustes coordinates and repeated the *PCA*. In addition, we tested each of the first eleven PCs that together describe more than 95 of total variation for allometric signals.

A set of 1,000 relaxed-clock molecular trees, which follow the topology of Hackett et al. (2008) and summarize the range of uncertainties in terms of time calibration of ancestral nodes, were downloaded from birdtree (http://birdtree.org) (Jetz et al. 2012, 2014) including all extant bird species in the dataset (Supplementary Data S1). Due to uncertainties in the taxonomic identification of *Aquila sp.*, this specimen was removed from the sample as we could not include it in the phylogeny. Because the specimen occupies almost the same position as *Aquilla audax*, we consider this deletion to have a negligible effect on the outcome of the analyses. Furthermore, the species *Sterna bergii* and *Grus rubincunda* used in

the analysis from Hu et al. (2019) are junior synonyms of *Thalasseus bergii* (Bridge et al. 2005) and *Antigone rubicunda* (Krajewski et al. 2010). Using the function *consensus.edges* in the *R* package *phytools v.0.7-20*, we computed a temporal consensus. The extinct dinornithiform species were placed as sister-group to Tinamidae following Mitchell et al. (2014). Because of their recent extinction (Holdaway & Jacomb 2000; Turvey & Holdaway 2005), the age was set to zero, similar to the other crown birds. The stem line representatives *Sinovenator changii* and *Sapeornis chaoyangensis* were added following the time-calibrated phylogeny of Rauhut et al. (2019). Because of the presence of juvenile specimens in dataset A and B, we added the juvenile specimens by splitting OTU (operational taxonomic unit) of *Struthio camelus* and *Dromaieus novehollandia* into a polytomy with each sub-OTU having a branch length of one year (this value had to be standardized, as *pFDA* requires an isometric tree). The impact of phylogeny on shape and centroid size (log-transformed) was tested using the function *physignal* in *Geomorph*. Based on *K*-statistics, this method evaluates the impact of phylogeny on a multivariate dataset relative to what is expected, if evolution is simulated under a Brownian motion model (Blomberg et al. 2003).

To explore the size of the single morphospaces of each taxon, we computed the morphological disparity using *morphol.disparity* in *Geomorph*. This analysis uses the Procrustes variance as disparity metric, which is the sum of all diagonal elements of a group covariance matrix divided by the number of observations within the group (Zelditch et al. 2012). Statistical comparisons of group sizes were executed with help of a permutation test with 999 iterations. To test for potential overlap in morphospace of vomer shapes in different clades of crown bird (see grouping 1 and 2) and their relation to the stem line representatives *Sinovenator changii* and *Sapeornis chaoyangensis*, we applied three different multivariate statistical methods, using the first eleven PCs as input data. We first applied a nonparametric multivariate analysis of variance (*perMANOVA*). This method evaluates the potential

overlapping of groups in morphospace by testing the significance of their distribution on the

basis of permutation (10,000 replications) and Euclidean distance (as one of several possible

distance measures), not requiring normal distribution of the data (Anderson, 2001; Hammer

& Harper, 2006). The spatial relationship of groups relative to each other is expressed by an

F value and *p* value. For the five-group comparison, the *p* values were Bonferroni-corrected

by multiplying the value with the number of comparisons. Next, we ran a discriminant

analysis (*DA*), which reduces a multivariate data set to a smaller set of dimensions by

maximizing the separation between two or more groups using Mahalanobis distances. This

distance measure is estimated from the pooled within-group covariance matrix, resulting in a

linear discriminant classifier and an estimated group assignment for each species. The results

were cross-validated using Jackknife resampling (Hammer & Harper, 2006; Hammer 2020).

Both multivariate tests were done with the program *PAST v.4.03* (Hammer et al. 2001).

Finally, we performed a phylogenetic flexible discriminant analysis (*pFDA*) (Schmitz &

Motani, 2011; Motani & Schmitz, 2011) in *R*. This method removes the phylogenetic bias

from the categorical variables before the actual discriminant analysis is unertaken by

estimating Pagel's lambda, which tests how the grouping correlates with phylogeny. This was

done for all allometric and non-allometric datasets.

The error of correct identification from the resulting confusion matrices was

compared between allometric and non-allometric data. For these comparisons, we used non-

parametric *Mann-Whitney U* and *Kruskal-Wallis* tests, which both estimates, whether or not

two univariate samples were taken from populations with equal medians, being more robust

against small sample sizes and data without normal distribution than parametric tests

(Hammer & Harper 2006). Both tests were carried out with *PAST*.

Finally, we applied an ordinary least square regression analysis to 19 species test the

correlation between log-transformed vomer size and the skull size using a log-transformed

box volume (Height x Width x Length). The measurements of the skull box volume were

taken from skullsite (https://skullsite.com).

Results

Based on the *PCA* of the original dataset, the first two PCs explain over 52% (Fig. 2A) of

total shape variation (PC1: 27.5%; PC2:25.1%). The morphospace of Palaeognathae and

Neognathae is almost equal in size. Taking the small sample size of Palaeognathae into

account, the size of their morphospace indicates relatively great shape variation. This is

supported by the Procrustes variances, which indicates a larger shape disparity in

Palaeognathae. Both Palaeognathae and Neognathae show a strong overlap along PC1 and a

partial overlap along PC2. When comparing neognath subclades, Aequorlitornithes show

strong overlap along both PCs with the palaeognath morphospace. Gruiformes lie in the

overlapping area of both groups. The morphospace of Inopinaves and Galloanserae overlap

with each other on both axes, but are separated from Palaeognathae, Aequorlitornithes and

Gruiformes along PC2. Within Neognathae, Galloanserae have the largest shape disparity,

becoming successively smaller in Inopinaves, Aequorlitornithes and Gruiformes

(Supplementary Data S2; S3: Fig. S1; S4: Table S1).

Allometry summarizes about 6.4% of total shape variation. Using non-allometric

residuals PC1 explains 29.3% and PC2 22.9% (Fig 2.B). While the general distribution of the

single bird clades does not change along PC1, the groups are less separated along PC2, which

contains the major allometric signal within the principal components (slope: -0.523; R^2:

0.185; *p*: 0.005; predicted variation: 19.5%), which is 4.9% of total shape variation in the

original dataset. Here, the palaeognath morphospace overlaps fully with Aequorlitornithes

and Gruiformes, partly with Inopinaves and marginally with Galloanserae. For the three other

datasets, we observe more or less similar general trends before and after size correction,

although the single morphospaces are partly shrunk. In all cases, the two stem line

representatives *Sinovenator changii* and *Sapeornis chaoyangensis* lie within the marginal

area of the palaeognaths/aequorlitornithines morphospace. Here, vomer morphology of the

troodontid *Sinovenator changii* is more bird-like than that of the avialian *Sapeornis*

chaoyangensis. The exclusion of the allometric shape variation has only a minor impact on

disparity. Thus, all disparity trends found in the original dataset persist in the non-allometric

datasets (Supplementary Data S2; S3: Fig. S1; S4: Table S1).

As previously detected by Hu et al. (2019), vomer shape is impacted by phylogeny.

Neither the modification of the sampling nor the exclusion of allometry changes this

relationship. In contrast, log centroid size does not contain a phylogenetic signal

(Supplementary Data S4: Table S2).

In all studied datasets, the *perMANOVA* found a significant separation between

palaeognath and neognath birds, showing no impact of allometry. For the five-group

comparison of the original dataset (A), the overall results still indicate significant separation

between clades for both allometric and non-allometric data. However, group-by-group

comparison of allometric data indicates an overlap in morphospace of Gruiformes with

Aequorlitornithes, Inopinaves, Galloanserae and Palaeognathae. These overlaps of

Gruiformes with other bird clades persist when allometry is removed from shape, but in

addition, Aequorlitornithes cannot be distinguished from Palaeognathae anymore, as

indicated by the *PCA* results (Fig. 2 A, B). The overlap between clades increases with the

exclusion of species with flat vomers and non-adult semaphoronts (Supplementary Data S4:

Table S3, S4).

For the original dataset (A) with allometry included, the *DA* identifies all species

correctly as Palaeognathae or Neognathae. The error of false identification increases to 2.6%

if the data are jackknifed. When allometry is removed, the error increases to 13.2% before

and 36.8% after jackknife resampling. In the former case, the misidentifications are restricted to neognath birds, which are wrongly classified as Palaeognathae, while jackknifing leads to identification errors in both groups. For the five-group comparison, all species of dataset (A) are correctly identified, when allometry is still present. The error is 18.4% after jackknife resampling, showing minor mismatches in all clades. Excluding allometry, the error increases to 10.5% before and 47.4% after jackknifing. While in the former case, a few Aequorlitornithes (2) and Inopinaves (1) species are wrongly identified as Palaeognathae (Fig. 2 C, D), Palaeognathae cannot be separated from the neognath subclades anymore after resampling. The exclusion of species with flat vomers and non-adult semaphoronts leads to an increase of error (Supplementary Data S4: Table S3-S6).

The *pFDA* found 15.8% of incorrect identifications when Palaeognathae are compared with neognaths in the original dataset (A). This error increases to 31.6% if shape is corrected for allometry. In both cases, error is primarily based on the incorrect identification of palaeognath specimens as neognaths. When Palaeognathae are compared with neognath subclades, the error of correct identification is 10.5% before and 26.3% after allometry is removed from the data. For the allometric data, the misidentifications result from the overlap between Paleognathae, Aequorlitornithes and Gruiformes. The misidentifications between these three groups are increased when shape is corrected for allometry, while Inopinaves are in part also wrongly identified as Palaeognathae. The exclusion of species with a flat vomer and/or non-adult semaphoronts usually causes a decrease of false identifications. However, the non-allometric dataset shows an increase in error for the two-group comparison, when species with flat vomers are excluded, and for the five-group comparison, when only adult semaphoronts are taken into account (Fig. 2 E, F). Nevertheless, for all four datasets, the error of correct identification is significantly higher for non-allometric vomer shapes (Fig 3A, Supplementary Data S4: Table S3-S6).

Discussion

The skull of crown birds possesses a complex kinetic system that includes a mobilized quadrate, the zygomatic arch (= jugal bar) and the pterygoid-palatine complex (PPC) that allows for the simultaneous, but restricted motion of both jaws (Bock 1964; Zusi 1984). According to Zusi (1984), the kinetic system can be differentiated into three main types: (1) prokinesis describes the rotation of the whole beak around the nasal-frontal hinge; (2) amphikinesis is derived from prokinesis, including the rotation of the beak around the nasal-frontal hinge plus an additional flexion of the anterior portion of the beak; and (3) rhynchokinesis, which in contrast includes a simple flexion of the beak around one or several bending zones rostral to the nasal-frontal suture, lacking a true hinge. Depending on the position of the bending zones, rhynchokinesis can be further differentiated into five subtypes (Zusi 1984). Most palaeognath birds possess central rhynchokinesis, while neognaths have realized all types of cranial kinesis (Zusi 1984), including some taxa with akinetic skulls (Reid 1835; Sims 1955; Degrange et al. 2010). In the past, several authors (Hofer 1954; Simonetta 1960; Bock 1963) suggested a close relationship between the morphology of the PPC and type of cranial kinesis. However, Gussekloo et al. (2001) demonstrated that all types of kinesis present in crown birds have similar movements of the quadrate, zygomatic arch and PPC. Palaeognathae and Neognathae only differ in the magnitude of kinetic movements in that Palaeognathae have slightly more restricted movement due to their rigid palate missing a movable joint between the pterygoid and palatine (Gussekloo et al. 2005).

Thus, although the results of geometric morphometric analysis of the vomer shape by Hu et al. (2019) imply at first glance a distinct separation between Palaeognathae and Neognathae, this separation does not necessarily reflect their conclusions regarding the evolution of cranial kinesis in crown birds, i.e., that cranial kinesis represents an innovation

of Neognathae. As indicated by the *PCA*, Palaeognathae occupy an enormous vomeral morphospace (Hu et al. 2019), which mirrors their generally large palatal disparity (see McDowell 1948) and partly overlaps with Gruiformes and Aequorlitornithes. In all cases tested, however, the exclusion of allometric shape variation generally increases the error of misidentification between all groups (Fig. 3A; Supplementary Data S4: Table S7), indicating that the taxonomic distinctions of shape found by Hu et al. (2019) are at least partly an artefact of size. This primarily concerns PC2, which according to Hu et al. (2019) separates Palaeognathae from Neognathae, but also contains the major part of allometric information. According to shape variation explained by PC2, larger birds tend to evolve vomers that are more dorsoventrally compressed. Only members of the Galloanserae could be still identified with a high amount of certainty when allometry is excluded.

Thus, our finding supports previous studies that demonstrated a relevant impact of allometry on skull shape evolution in birds (Klingenberg & Marugán-Lobón 2013; Bright et al. 2016; Linde-Medina 2016; Tokita et al. 2016, Bright et al. 2019). By modifying the dataset, it becomes further clear that both the homoplastic presence of flat vomers in Aequorlitornithes, Inopinaves, Galloanserae (Dataset B) and ontogenetic variation (Dataset C) affects the accuracy of taxonomic identification. In addition, Palaeognathae and Neognathae do not differ in vomer size when compared to the head size (Fig. 3B). Consequently, vomer shape is not practical for taxonomic identification and should not be used as a proxy to infer the presence or absence of cranial kinesis in crown birds or their stem. As the manifold shape diversity of crown bird's skulls is impaired by a tessellated evolution with multiple convergent events (e.g., Zusi 1993; Felice & Goswami 2018), the use of isolated elements for taxonomic identification and/or biomechanical implications should be treated generally with some caution.

In fact, *DA* and *pFDA* frequently identified the troodontid *Sinovenator changii* and

avialan *Sapeornis chaoyangensis* as neognaths or neognath subclades when allometry is

excluded, while the original dataset implied a referral to Palaeognathae (see also Hu et al.

2019). However, the skull anatomy of both species indicates no cranial kinesis (Xu et al.

2002; Wang et al. 2017; Yin et al. 2018; Hu et al. 2020).

The origin and evolution of cranial kinesis in the stem line of birds is still not well

understood due to the rarity of complete three-dimensional skulls. However, skull material

from the ornithurines *Ichthyornis dispars* and *Hesperornis regalis* indicates a certain degree

of rhynchokinesis (Bühler et al. 1988; Field et al. 2018) that might be comparable to that of

extant Palaeognathae or some Aequorlitornithes, but further shows that this functional

character was already present before the origin of the crown. Their kinesis is indicated by the

loss of the jugal-postorbital bar and the ectopterygoid (resulting in a loss of contact in the

jugal with the skull roof and the palate), the presence of a mobile bicondylar quadrate and a

mobile joint between quadrate and quadratojugal. Recently, Plateau & Foth (2020) speculated

that the peramorphic bone fusion in the braincase could be also related to cranial kinesis, in

which the fusion-induced immobility constrains a controlled kinetic dorsoventral flexion of

the avian beak during biting/picking. Based on these characters, most Mesozoic Avialae

(including *Sapeornis chaoyangensis*) still had akinetic skulls, although some Enantiornithes

possessing a reduced jugal-postorbital bar might have evolved primitive kinesis convergently

to Ornithurae (O'Connor & Chiappe 2011).

Allometry describes the relationship between size and shape, which is one of the key

concepts in biology to explain variation of shape in organisms (Klingenberg 1998), and is

crucial for studying taxonomy, ontogeny and functional morphology. Investigating the effect

of allometry on the vomer shape in crown birds with help of multiple multivariate statistical

methods, indicates that this bone is not a good proxy for taxonomy. Shape differences

between Palaeognathae and Neoganthae are clearly affected by size and can neither be used

to differentiate between different types of cranial kinesis nor to explain the evolution of

cranial kinesis within crown birds. In contrast, the evolution of cranial kinesis in birds needs

to be studied in context of the whole pterygoid-palatine complex and its contacts with the

braincase and quadrate-zygomatic arch.

Acknowledgements

We thank Walter Joyce Roland Sookias and Sergio Martínez Nebreda for their critical

comments on previous versions of the manuscript. The Swiss National Science Foundation is

thanked for its financial support (PZ00P2_174040 to C.F.).

Additional information

Funding

This study was funded by the Swiss National Science Foundation (PZ00P2_174040).

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the

content of this article. None of the authors is a *PCIPaleo* recommender.

Author Approvals

All authors have seen and approved the manuscript. The manuscript has not been accepted or

published elsewhere.

Author contributions

O.P. and C.F. designed the research project and analysed the data; and O.P. and C.F. wrote

the paper and prepared all figures.

Data availability

The 3D models and landmarks data of Hu et al. (2019) are available at Figshare (DOI:

https://doi.org/10.6084/m9.figshare.7769279.v2).

Supplementary information

• Data S1: Phylogenetic trees used for *pFDA*.

• Data S2: PCA results of all dataset before and after correction for allometry.

• Data S3: PCA plots of Dataset B-D before and after correction for allometry.

• Data S4: Results of npMANOVA, DA and pFDA.

• Data S5: R Code including all statistical analyses.

References

Adams DC, Otárola-Castillo E. 2013. *geomorph*: an *R* package for the collection and analysis

of geometricmorphometric shape data. *Methods in Ecology and Evolution* 4:393–399.

Anderson MJ. 2001. A new method for non-parametric multivariate analysis of variance.

Austral Ecology 26:32–46.

Bellairs ADA, Jenkin CR. 1960. The skeleton of birds. In: Marshall AJ ed. *Biology and*

Comparative Physiology of Birds, Vol. 1. New York: Academic Press, 241–300.

Blomberg SP, Garland TJ, Ives AR. 2003. Testing for phylogenetic signal in comparative

data: behavioral traits are more labile. *Evolution* 57:717–745.

Bock WJ. 1963. The cranial evidence for ratite affinities. *Proceedings of the 13th*

International Ornithological Congress 1:39–54.

Bock WJ. 1964. Kinetics of the avian skull. *Journal of Morphology* 114:1–42.

Bridge ES, Jones AW, Baker AJ. 2005. A phylogenetic framework for the terns (Sternini)

inferred from mtDNA sequences: implications for taxonomy and plumage evolution.

Molecular Phylogenetics and Evolution 35:459–469.

Bright JA, Marugán-Lobón J, Cobb SN, Rayfield J. 2016. The shapes of bird beaks are highly

controlled by nondietary factors. *Proceedings of the National Academy of Sciences,*

U.S.A. 113:5352–5357.

Bright JA, Marugán-Lobón J, Rayfield J, Cobb SN. 2019. The multifactorial nature of beak

and skull shape evolution in parrots and cockatoos (Psittaciformes). *BMC Evolutionary*

Biology 19:1–9.

Bühler P, Martin LD, Witmer LM. 1988. Cranial kinesis in the Late Cretaceous birds

Hesperornis and *Parahesperornis*. *The Auk* 105:111–122.

Degrange F, Tambussi C, Moreno K, Witmer L, Wroe S. 2010. Mechanical analysis of

feeding behavior in the extinct "Terror Bird" *Andalgalornis steulleti* (Gruiformes:

Phorusrhacidae). *PLoS ONE* 5:e11856.

Felice RN, Goswami A. 2018. Developmental origins of mosaic evolution in the avian

cranium. *Proceedings of the National Academy of Sciences, U.S.A.* 115:555–560.

Field DJ, Hanson M, Burnham DA, Wilson LE, Super K, Ehret D, Ebersole JA, Bhullar B-

AS. 2018. Complete *Ichthyornis* skull illuminates mosaic assembly of the avian head.

Nature 557:96–100.

Goodall C. 1991. Prorustes methods in the statistical analysis of shape. *Journal of the Royal*

Statistical Society: Series B (Methodological) 53:285–321.

Gussekloo SWS, Bout RG. 2005. The kinematics of feeding and drinking in palaeognathous

birds in relation to cranial morphology. *The Journal of Experimental Biology* 208:3395–

3407.

Gussekloo SWS, Vosselman MG, Bout RG. 2001. Three-dimensional kinematics of skeletal

elements in avian prokinetic and rhynchokinetic skull determined by roentgen

stereophotogrammetrics. *The Journal of Experimental Biology* 204:1735–1744.

Hackett SJ, Kimball RT, Reddy S, Browie RCK, Braun EL, Chojnowski JL, Cox A, Han K,

Harshman J, Huddleston CJ, Marks BD, Miglia KJ, Moore WS, Sheldon FH, Steadman

DW, Witt CC, Yuri T. 2008. A phylogenomic study of birds reveals their evolutionary

history. *Science* 320:1763–1768.

Hammer O, Harper DAT. 2006. *Paleontological Data Analysis*. Malden: Blackwell

Publishing, 351 pp.

Hammer O, Harper DAT, Ryan PD. 2001. PAST: paleontological statistics software package

for education and data analysis. *Palaeontologia Electronica* 4:1–9.

Hammer O. 2020. *PAST Paleontological Statistics v.4.03. Reference Manual.* Oslo:

University of Oslo, 283 pp.

Hofer H. 1954. Neuere Untersuchungen zur Kopfmorphologie der Vögel. *Acta XI Congressus*

Internationalis Ornithologici:104–137.

Holdaway RN, Jacomb C. 2000. Rapid extinction of the moas (Aves: Dinornithiformes):

model, test, and implications. *Science* 287:2250–2254.

Hu H, O'Connor J, McDonald P, Wroe S. 2020. Cranial osteology of the Early Cretaceous

Sapeornis chaoyangensis (Aves: Pygostylia). *Cretaceous Research* 113:104496.

Hu H, Sansalone G, Wroe S, McDonald PG, O'Connor JK, Li Z, Xu X, Zhou Z. 2019.

Evolution of the vomer and its implications for cranial kinesis in Paraves. *Proceedings*

of the National Academy of Sciences, U.S.A. 116:19571–19578.

Huxley TH. 1867. On the classification of birds; and on the taxonomic value of the

modifications of certain cranial bones observable in the class. *Proceedings of the*

Zoological Society of London 27:415–472.

Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. 2012. The global diversity of birds in

space and time. *Nature* 491:444–448.

Jetz W, Thomas GH, Joy JB, Redding DW, Hartmann K, Mooers AO. 2014. Global

distribution and conservation of evolutionary distinctness in birds. *Current Biology*

24:1–12.

Klingenberg CP. 1998. Heterochrony and allometry: the analysis of evolutionary change in

ontogeny. *Biological Reviews* 73:79–123.

Klingenberg CP, Marugán-Lobón J. 2013. Evolutionary covariation in geometric

morphometric data: analyzing integration, modularity, and allometry in phylogenetic

context. *Systematic Biology* 62:591–610.

Krajewski C, Sipiorski JT, Anderson FE. 2010. Complete mitochondrial genome sequences

and the phylogeny of cranes (Gruiformes: Gruidae). *The Auk* 127:440–452.

Linde-Medina M. 2016. Testing the cranial evolutionary allometric "rule" in Galliformes.

Journal of Evolutionary Biology 29:1873–1878.

Mayr G. 2017. *Avian evolution*. Chichester: John Wiley, 293 pp.

McDowell S. 1948. The bony palate of birds. Part I. The Palaeognathae. *The Auk* 65:520–

549.

Mitchell K, Llamas B, Soubrier J, Rawlence N, Worthy T, Wood J, Lee M, Cooper A. 2014.

Ancient DNA reveals elephant birds and kiwi are sister taxa and clarifies ratite bird

evolution. *Science* 344:898–900.

Motani R, Schmitz L. 2011. Phylogenetic versus functional signals in the evolution of form-

function relationships in terrestrial vision. *Evolution* 65:2245–2257.

O'Connor JK, Chiappe LM. 2011. A revision of enantiornithine (Aves: Ornithothoraces)

skull morphology. *Journal of Systematic Palaeontology* 9:135–157.

Plateau O, Foth C. 2020. Birds have peramorphic skulls, too: anatomical network analyses

reveal oppositional heterochronies in avian skull evolution. *Communications Biology* 3.

R Development Core Team. 2011. *R*: a language and environment for statistical computing.

http://www.r-project.org.

Rauhut OWM, Tischlinger H, Foth C. 2019. A non-archaeopterygid avialan theropod from

the Late Jurassic of southern Germany. *eLife* 8:e43789.

Reid J. 1835. Anatomical description of the Patagonian penguin. *Proceedings of the*

Zoological Society of London 3:132–148.

Revell LJ. 2012. *phytools*: an *R* package for phylogenetic comparative biology (and other

things). *Methods in Ecology and Evolution* 3:217–223.

Schmitz L, Motani R. 2011. Nocturnality in dinosaurs inferred from scleral ring and orbit

morphology. *Science* 332:705–708.

Simonetta A. 1960. On the mechanical implications of the avian skull and their bearing on

the evolution and classification of birds. *The Quaterly Review of Biology* 35:206–220.

Sims S. 1955. The morphology of the head of the hawfinch (*Coccothraustes coccothraustes*).

Bulletin of the British Museum 2:371–393.

Tokita M, Yano W, James H, Abzhanov A. 2016. Cranial shape evolution in adaptive

radiations of birds: comparative morphometrics of Darwin's finches and Hawaiian

honeycreepers. *Philosophical Transactions of the Royal Society B* 372:20150481.

Turvey ST, Holdaway RN. 2005. Postnatal ontogeny, population structure, and extinction of

the giant moa *Dinornis*. *Journal of Morphology* 265:70–86.

Wang Y, Hu H, O'Connor JK, Wang M, Xu X, Zhou Z, Wang X, Zheng X. 2017. A

previously undescribed specimen reveals new information on the dentition of *Sapeornis*

chaoyangensis. *Cretaceous Research* 74:1–10

Xu X, Norell MA, Wang X, Makovicky PJ, Wu X. 2002. A basal troodontid from the Early

Cretaceous of China. *Nature* 415:780–784.

Yin Y, Pei R, Zhou C. 2018. Cranial morphology of *Sinovenator changii* (Theropoda:

Troodontidae) on the new material from the Yixian Formation of western Liaoning,

China. *PeerJ* 6:e4977.

Zelditch ML, Swiderski DL, Sheets HD. 2012. *Geometric morphometrics for biologists: a*

*prime*r. Amsterdam: Elsevier Academic Press, 488 pp.

Zusi RL. 1984. A functional and evolutionary analysis of rhynchokinesis in birds.

Smithsonian Contributions to Zoology 395:1-40.

Zusi RL. 1993. Patterns of diversity in the avian skull. In: Hanken J, Hall BK eds. *The skull.*

Vol. 2. Patterns of structural and systematic diversity. Chicago: University of Chicago

Press, 391–437.

Figure legends

Fig. 1. Anatomical organization of the pterygoid-palatine complex (PPC) and shape

variability of the vomer in palaeognath and neognath birds. (A) Palates of *Dromaius*

novaehollandia (left), *Cygnus olor* (middle) and *Corvus corax* (right) in ventral view (all

specimens form the Natural History Museum of Fribourg/University of Fribourg). For

Dromaius novaehollandiae and *Corvus corax* the main organization of palate morphology is

highlighted in a coloured scheme. The blue arrow in *Dromaius novaehollandiae* indicates the

contact between the basipterygoid process and the pterygoid. The red arrow in *Corvus corax*

indicates the mobile joint between pterygoid and palatine. **(B)** 3D models of the vomer of

Dinornis robustus, *Anas crecca* and *Strepera graculina* in lateral view (left) and anterior

view (right) from (not at scale) (3D models from Hu et al. 2019). Abbreviations: Max,

Maxillary; Pal, Palatine; Premax, Premaxilary; Pter, Pterygoid; Vom, Vomer.

Fig. 2. Differences between allometric and non-allometric morphospaces of the vomer

(Dataset A) in palaeognath and neognath birds. (A) PCA *(Principal Component Analysis)*

results of allometric data. **(B)** *PCA* results of non-allometric data. **(C)** *DA (Discriminante*

Analysis) results of allometric data. **(D)** *DA* results of non-allometric data. **(E)** *pFDA*

(phylogenetic Flexible Discriminant Analysis) results of allometric data. **(F)** *pFDA* results of

non-allometric data. Centroid sizes (Csize) are indicated by the size of the symbols. All

silhouettes are taken from http://www.phylopic.org/.

Fig. 3. Errors of correct taxonomic identification for all comparisons of Dataset A-D.

(A) Two-group identification (Palaeognathae and Neognathae) before (red) and after (green)

correction for allometry. *DA*, Discriminant analysis; *DA*JK*, Discriminant Analysis with

jackknife resampling; *pFDA*, phylogenetic Flexible Discriminant Analysis. **(B)** Five-group

identification (Palaeognathae, Aequorlitornithes, Galloanserae, Gruiformes and Inopinaves).

(C) *OLS* regression (black line) between log-transformed skull box volume and log-

transformed centroid size of the vomer. Grey shadow marks the area of the 95% confidence

interval.

A

Premax
Max
Vom
Pal
Pter

Dromaius novaehollandiae

Premax
Max
Vom
Pal
Pter

Cygnus olor

Premax
Max
Vom
Pal
Pter

Corvus corax

1 cm

| | Premax | | Max | | Vom | | Pal | | Pter |

B

Dinornis robustus

Anas crecca

Strepera graculina

Dorsal

Anterior

Dorsal

Lateral

A Allometric

PC2[25.09%]
PC1[27.62%]

B Non Allometric

PC2[22.88%]
PC1[29.31%]

C Allometric

DA2
DA1

D Non Allometric

DA2
DA1

E Allometric

DA1
DA1

F Non Allometric

DA2
DA1

Aequorlitornithes Gruiformes Paleognathae
Galloanserae Inopinaves Csize
Sinovenator Sapeornis 40 160