1

The condensation of HP1-*α*/Swi6 imparts nuclear stiffness

Jessica F. Williams[1], Ivan V. Surovtsev[1,2], Sarah M. Schreiner[1], Ziyuan Chen[3], Gulzhan Raiymbek[4], Hang Nguyen[1], Yan Hu[1], Julie S. Biteen[3,5], Simon G.J. Mochrie[2], Kaushik Ragunathan[4,6]* and Megan C. King[1]*

[1] Department of Cell Biology, Yale School of Medicine, 333 Cedar Street, New Haven, Connecticut, USA, 06520

[2] Department of Physics, Yale University, 217 Prospect Street, New Haven, Connecticut, USA, 06511

[3] Department of Biophysics, University of Michigan, Ann Arbor, MI 48109 USA.

[4] Department of Biological Chemistry, University of Michigan, Ann Arbor, MI 48109 USA.

[5] Department of Chemistry, University of Michigan, Ann Arbor, MI 48109 USA.

[6] Current affiliation: Department of Biology, Brandeis University, Waltham, MA 02453 USA.

* To whom correspondence should be addressed: megan.king@yale.edu or kaushikr@brandeis.edu

SUMMARY

The condensation of proteins and nucleic acids underlies the formation of membraneless organelles, which have emerged as major drivers of cellular organization. It remains largely unexplored, however, whether these condensates can impart mechanical function(s) to the cell. The heterochromatin protein HP1-α (Swi6 in *S. pombe*) crosslinks histone H3K9 methylated nucleosomes and has been proposed to undergo condensation to drive the liquid-like clustering of heterochromatin domains. Here we leverage the genetically tractable *S. pombe* model and a separation-of-function Swi6 allele to elucidate a mechanical function imparted by its condensation. Using a combination of single-molecule imaging, force spectroscopy on individual nuclei, and high-resolution live-cell imaging, we show that Swi6 is critical for nuclear resistance to external force. Strikingly, it is this condensed yet dynamic pool of Swi6, rather than the chromatin-bound molecules, that is essential to imparting mechanical stiffness. Our findings suggest that Swi6 condensates embedded in the chromatin meshwork establish the emergent mechanical behavior of the nucleus as a whole, revealing that biomolecular condensation can influence organelle and cell mechanics.

KEYWORDS

HP1, Swi6, heterochromatin, biomolecular condensation, nuclear mechanics, *S. pombe*, microscopy, biophysics

INTRODUCTION

The cell nucleus provides a mechanically resilient compartment that encapsulates and protects the genome[1,2]. The nuclear compartment is defined by the nuclear envelope, an integrated network of the inner and outer nuclear membranes, nuclear pore complexes, integral inner nuclear membrane proteins, peripheral chromatin, and, in higher eukaryotes, a polymer mesh of intermediate filaments (lamins)[3–5]. Maintaining the integrity of the nuclear envelope is critical for cell survival, and mechanical defects that lead to nuclear blebbing and rupture are tied to human disease[1,2]. Nuclear ruptures can occur during cellular migration, when cells move through tissue and experience significant compressive force[6,7], but even intrinsic cytoskeletal forces in adherent or contractile cells are strong enough to induce nuclear rupture, indicating that the nucleus is under constant strain from its own cellular architecture[8,9].

In the effort to understand how the nucleus withstands these mechanical forces, much work has focused on the contribution of the nuclear lamina[10,11]. However, we previously demonstrated that the tethering of chromatin to the nuclear envelope is critical for maintaining nuclear stiffness in fission yeast, which lack nuclear lamins[12]. Other studies in mammalian cells have further demonstrated the unique mechanical role of chromatin in its contribution to the nuclear force response to small deformations (1-3 μm), while A-type lamins are critical for the force response to large deformations (> 3 μm)[13]. Additionally, when chromatin compaction is globally altered through histone deacetylase or demethylase inhibition, the relative proportion of euchromatin to heterochromatin (HC) affects the overall stiffness of the nucleus[14]. Thus, both the lamina and the underlying chromatin are important for maintaining mechanical stability, yet each play separate roles related to the physical—and perhaps temporal—scale of the external strain. Indeed, increasing the amount HC is sufficient to rescue the nuclear irregularity and nuclear blebbing phenotype seen in HeLa cells expressing the mutant form of lamin A (progerin) of the premature aging disease Hutchinson-Guilford Progeria Syndrome[14]. Together, these studies suggest that in addition to its function as a transcriptional insulator[15], HC acts as a dynamic mechanical scaffold for the nucleus[16]; however, the

biophysical mechanisms by which HC can influence nuclear mechanics remain largely

undefined.

An evolutionarily conserved feature of nuclear organization is the association of HC with

the nuclear periphery. This localization has classically been thought to reflect the ability

of HC to promote gene silencing. Several studies, however, suggest that the causality of

the periphery-silencing relationship may be more complex, implying that there may be

additional function(s) for peripherally-located HC. Indeed, in *C. elegans*, HC positioning

at the nuclear periphery does not dictate gene silencing[17]. Moreover, leveraging the

model of rod cells with "inverted" nuclei, re-localization of HC from the periphery to the

center of the nucleus was found to neither alter gene expression nor chromatin

organization at the 100s of kilobase to many megabase scale[18]. Our previous work

demonstrated that chromatin tethering to the nuclear periphery is important for resisting

nuclear deformations in response to microtubule forces in fission yeast[12,19], which is

echoed by studies in mammalian cells[14,20]. Here we further probe the hypothesis that a

primary function of peripherally-located HC is to mechanically support the nuclear

envelope.

The potential mechanism for this mechanical support stems from recent evidence

suggesting that a key protein responsible for HC compaction and gene silencing—

heterochromatin protein (HP1-α)—is important for maintaining normal nuclear

morphology and stiffness without affecting HC peripheral localization[21] and forms

phase-separated, HP1α-rich HC domains[22,23]. These (and other) biomolecular

condensates often exhibit liquid-like properties, including surface tension and

viscosity[24–29]. One model proposes that the surface tension of HC-associated

condensates, enhanced by strong electrostatic interactions as observed for HP1-α[30],

would resist deformation. Alternatively, or in addition, a distinct model of "elastic

ripening" is being developed[31], in which the formation of condensates embedded in a

polymer network gives rise to a composite system with emergent or synergistic

mechanical properties. Lastly, while it has been proposed that biomolecular

condensates can do mechanical work[32], for example, to drive local membrane

116 bending[33–35] or, in an artificial system, to bring two chromatin loci together[36], whether

117 biomolecular condensation can drive emergent mechanical behaviors at the scale much

118 larger than an individual condensate, i.e., at the level of organelles and cells, has yet to

119 be explored.

121 To biomechanically define the features of HP1-α/Swi6 HC domains, we employed both

122 *in vivo* fluorescence microscopy and *in vitro* force spectroscopy techniques. We

123 demonstrate that HC domains respond to mechanical force with viscoelastic properties,

124 and that proper coalescence of the domains—specifically telomeres—depends on HP1-

125 α/Swi6. Using single-molecule imaging we describe a separation-of-function allele of

126 HP1-α/Swi6—*swi6-sm1*—which displays disruption of the biomolecular condensation of

127 HC domains despite normal H3K9me2/3 binding and crosslinking *in vivo*. This defect in

128 Swi6 condensation leads to depletion of mobile Swi6 at HC domains and concomitantly

129 leads to substantially softer nuclei both *in vitro* and *in vivo*. Taken together, our findings

130 argue that it is the condensation of HP1-α/Swi6 at HC domains, rather than the binding

131 and/or crosslinking of H3K9me2/3 nucleosomes, that contributes to the overall stiffness

132 of the nucleus, showcasing a role for biomolecular condensates within a biological

133 meshwork (the chromatin) in driving the emergent mechanical stiffness of a membrane-

134 bound compartment.

RESULTS

Heterochromatin domains exhibit liquid-like properties *in vivo* that respond to cytoskeletal forces

The HC protein HP1-α/Swi6 is a key component of centromeric and telomeric HC, responsible for its condensation and transcriptional repression[37–39] and has been shown to oligomerize and form phase-separated domains that facilitate HC formation[22,23,40]. With this in mind, we sought to further characterize the liquid-like properties of Swi6 at two regions at which it is highly enriched: the centromeres and the telomeres[37,41,42].

When Swi6 is visualized as a fluorescent protein fusion expressed from its endogenous locus (Swi6-GFP), it resides in several nuclear foci/domains (Figure 1A, top). The largest domain is comprised of the centromeres of the three chromosomes, which are coupled to the spindle-pole body (SPB) interface (Figure 1A, bottom). The other, smaller heterochromatic foci contain the six telomeres, the mating type locus, and other silenced regions of the genome and are typically associated with the nuclear envelope far from the SPB. In *S. pombe*, microtubules continuously translocate the SPB, which transmits the MT-dependent force on to the centromeric HC. As the centromeric HC is driven across the nucleus by MT forces, we routinely observe a morphologic transition from a circular domain (as visualized from a two-dimensional image plane; Figure 1B) when "at rest" (at t = 0) to an elliptical domain when under maximum MT shearing force (at t = 30-80s), stretching along the SPB oscillation axis (Figure 1B; see also insets of reconstructed contours in green). At various time points during this translational motion (t > 60s), the centromere displays an apparent fission into smaller, individual domains that later merge (t > 100s). These observations suggest that mechanical forces from the cytoskeleton are directly transduced to HC, leading to changes in domain morphology consistent with liquid-like reorganization[26,43]. We therefore considered the possibility that HC domains can act as mechanical cushions, absorbing MT-driven forces exerted upon the nucleus through their resistance to deformation and/or their effects on the emergent properties of the chromatin itself.

Consistent with previous observations for HP1-α in mammalian cells[23], the addition of

5% of the aliphatic alcohol 1,6-hexanediol for 15 minutes, which disrupts weak

hydrophobic interactions, leads to both a rapid decrease in both the size and intensity of

nuclear GFP-Swi6 foci as well as the accumulation of a diffuse pool of GFP-Swi6 in the

cytoplasm (Figure S1). However, a pool of GFP-Swi6 is retained within the HC foci,

which are expected to be 1,6-hexanediol-resistant due to direct binding via the Swi6

chromodomain to nucleosomes, particularly those containing methylated histone H3K9.

Taken together, these observations support the existence of at least two populations of

Swi6 molecules within HC foci: one directly bound to chromatin through the Swi6

chromodomain and the other driven by biomolecular condensation that is sensitive to

1,6-hexanediol, consistent with our prior single-molecule experiments demonstrating

four distinct diffusive states of Swi6[44]. These observations suggest Swi6, as an

ensemble, is an attractive candidate to impart viscoelastic properties to HC and, in turn,

to the nucleus as a whole.

Heterochromatin domains exhibit liquid-like properties *in vivo* that are tied to

normal telomere clustering

As a role for biomolecular condensation in driving telomere clustering in cancer cells

has been described previously[45,46], we also suspected that liquid-like fusion could drive

the coalescence of the six (replicated) heterochromatic telomeres, which are clustered

together at the nuclear periphery in *S. pombe*[47]. When visualized by expressing a GFP

fusion of Taz1, a shelterin complex subunit that specifically binds telomeric DNA[48],

approximately half of nuclei display one or two foci containing the six telomeres, with the

other half displaying up to four foci (Figure 1C,D and Figure S2). This behavior is largely

cell cycle-independent, as assessed using cell length as a proxy for cell cycle

progression (Figure 1D). To further interrogate if telomere clustering relies on liquid-like

condensation mediated by weak hydrophobic interactions, we treated cells with 1,6-

hexanediol (5% for 30 minutes). This perturbation resulted in an increase in the number

of Taz1-GFP foci per nucleus as the telomeres de-clustered (Figure 1D).

Similar to the 1,6-hexanediol treatment, we found that *swi6Δ* cells have an increase in

the number of Taz1-GFP foci per nucleus (Figure 1D), indicating a clustering defect and

a reliance on Swi6 for proper telomere coalescence. We note that even in this condition

many cells display less than six discernible telomere foci, which we suspect occurs due

to stochastic coincidence of telomeres in the nuclear volume or an inability to fully

visualize the dimmer, de-clustered telomeres. Taken together, these observations

suggest that Swi6 HC foci have a component of their behavior that is liquid-like and

sensitive to treatment with 1,6-hexanediol. Furthermore, Swi6 contributes to the normal

coalescence of telomere foci in fission yeast.

It is important to note that in fission yeast (as in mammalian cells), Swi6 shares its

transcriptional silencing role in gene regulation with other chromodomain proteins,

including Chp1 and Chp2, which are necessary for the proper establishment and

balance of heterochromatic regions, respectively[49,50]. All three chromodomain proteins

rely on the H3K9 methyltransferase Clr4 to be recruited to methylated histones of

heterochromatic domains[49,51]. Their protein structures, however, differ from each other;

for example, Chp1 lacks a chromoshadow domain and contains a much smaller N-

terminal domain[49] (Figure S3A). Under wild-type conditions, Chp1—like Swi6—binds to

HC to form discrete foci (Figure S3B), and, also similar to Swi6, Chp1 is unable to bind

and localize to heterochromatic foci in the absence of Clr4[49] (Figure S3C,G). When

H3K9 HC was allowed to spread in the absence of the H3K9me2/3 demethylase

Epe1[52], we observed that Chp1-GFP intensity increased at HC foci (Figure S3D-G),

consistent with an enlargement of HC domains in Epe1-deficient cells. In the absence of

Swi6, Chp1-GFP foci intensity also significantly increased (Figure S3E-G), suggesting

competition of H3K9me2/3 binding between Swi6 and Chp1. With this, we can thus infer

that the other chromodomain proteins, even at increased levels, are unable to replace

Swi6's role in telomere clustering in *swi6Δ* cells (Figure 1D), highlighting a unique

feature of Swi6.

The *swi6-sm1* allele disrupts the ability of HP1-α/Swi6 to promote the coalescence

of heterochromatin domains

To determine the biochemical features of Swi6 tied to its ability to form and coalesce

liquid-like HC domains, we turned to prior genetic analysis of Swi6 function. In

particular, we interrogated the *swi6-sm1* allele, which carries two mutations in its N-

terminal domain, rendering it unable to silence HC while leaving its H3K9me2/3-binding

activity, dimerization, and cohesin interactions unaffected[53] (Figure 2A). In previous

studies of Swi6, defects in its N-terminal phosphorylation by casein kinase II also led to

loss of transcriptional silencing[54]. Indeed, the D21G mutation encoded in the *swi6-sm1*

allele disrupts the casein kinase II recognition site for phosphorylation of serine 18; this

site is phosphorylated *in vivo* and *in vitro* by casein kinase II, and its targeted mutation

disrupts Swi6's ability to promote gene silencing[54], thereby phenocopying the *swi6-sm1*

allele. As N-terminal phosphorylation enhances oligomerization leading to phase

separation for both HP1α and Swi6[22,55], we hypothesized that the *swi6-sm1* allele could

disrupt Swi6's ability to undergo biomolecular condensation.

Like WT Swi6-GFP, *swi6-sm1*-GFP accumulates in foci within the nucleus close to the

nuclear periphery (Figure 2). To begin to probe if and how the *swi6-sm1* mutation alters

the coalescence of HC domains, we compared the size, intensity, and number of Swi6-

GFP and *swi6-sm1*-GFP foci within individual nuclei (Figure 2B). To quantitatively

compare foci size and intensity, we employed an image analysis approach we

previously established that reconstructs HC foci from image z-stacks with sub-pixel

resolution[56] (Figure 2C). Comparing each foci's intensity to its cross-sectional area in an

ensemble scatter plot revealed divergent populations in Swi6-GFP versus *swi6-sm1*-

GFP-labeled HC (Figure 2D and Figure S4). In wild-type cells, Swi6-GFP HC domains

comprise a nearly uniform distribution of intensities ranging from bright (centromeric

HC) to dim (telomeric and other HC foci). Cells expressing *swi6-sm1*-GFP, however,

display a depletion of intermediately intense HC foci, indicative of non-centromeric HC

domain de-clustering (p < 0.0001). This depleted population is most likely to represent

intact, condensed telomeric domains, indicating that *swi6-sm1* affects proper telomere

coalescence. To support this, expression of the *swi6-sm1* allele with Taz1-GFP shows

disruption of normal telomere clustering similar to complete loss of Swi6 or treatment

with 1,6-hexanediol (Figure 2E and Figure 1D). To the extent that the condensation of

Swi6 facilitates proper telomere clustering, we conclude that the *swi6-sm1* allele could

specifically disrupt aspects of Swi6 function that underlie its ability to form condensates

(Figure 2F).

The *swi6-sm1* allele shows a depletion of mobile states despite normal chromatin

and H3K9me2/3 binding *in vivo*

Previously, we used a single-molecule approach to track the dynamics of Swi6 in the

fission yeast nucleus[44]. We fused Swi6 to the photoactivatable fusion protein

PAmCherry and captured the distinct biophysical states associated with its motion

within its native chromatin context (Figure 3A,B). Through mutational analysis, we

defined how each biophysical state of Swi6 is associated with a unique biochemical or

structural property that influences how Swi6 behaves at sites of H3K9 methylation[44].

Our analysis revealed four states of WT PAmCherry-Swi6 molecules with different

diffusion coefficients (Figure 3C, crosses) and transition probabilities between the states

(Figure 3D). The least mobile state (termed α, blue) comprises molecules bound to

H3K9me2/3 (weight fraction, π ~25%), the intermediate states indicate molecules

associated with unmethylated H3K9 nucleosomes (termed β, orange; π ~33%) or

nucleic acids (termed γ, yellow; π ~15%), and the remaining, most mobile state is

comprised of unbound molecules (termed δ, purple; π ~27%) (Figure 3C)[44]. Note that

this observed high-mobility state of Swi6 displays a ~40-fold slower diffusion coefficient

(D = 0.51 μm^2/s) than what has been reported for similar-size GFP-tagged HP1

monomers and dimers in euchromatic regions where it is expected to be relatively fully

unbound/free[57]. Thus, the δ state is slowed significantly compared to truly freely

diffusing molecules, consistent with it representing unbound but restrained movement in

the mobile phase of the condensate at HC domains. Indeed, the rapid diffusion of

completely free monomers and dimers would not detectable within the acquisition

parameters optimized for our PAmCherry-Swi6 experiments.

In light of the quantitative defects observed in the HC domains formed in the presence

of the *swi6-sm1* allele (Figure 2), we sought to characterize how *swi6-sm1* mutations

affect the distribution of mobility states and thus infer how its two N-terminal mutations

influence Swi6 dynamics. Overall, PAmCherry-*swi6-sm1* molecules move more slowly

compared to PAmCherry-Swi6 as noted by the ~30% increase in the fraction of

molecules in the H3K9me and chromatin bound states (α and β states) (Figure 3C,

scatter dots). Using Single-Molecule Analysis by Unsupervised Gibbs sampling

(SMAUG) analysis[58], we distinguished four mobility states associated with PAmCherry-

swi6-sm1; the same number as previously observed for wild-type PAmCherry-Swi6

(Figure 3C,E). The *swi6-sm1* mutations, however, led to significant changes in the

distributions of molecules within each state (Figure 3C) and the transitions between

them (Figure 3E and Table S1). Of note, the weight fraction, π, of the mobile (δ) Swi6

population, previously associated with unbound Swi6 molecules, decreased significantly

from $\pi = 27\%$ to only 3% of all molecules. Concomitant with this decrease, there is an

increase in the H3K9me2/3-bound (α) state weight fraction from $\pi = 25\%$ to 32% and a

substantial increase in the occupancy of the nucleosome-bound (β) state from $\pi = 33\%$

to 50%. We also analyzed the transition probabilities between the different mobility

states and observed that a major difference between WT Swi6 and *swi6-sm1* is a 4-fold

increase in the transition probability between the δ and β states, suggesting that the

small pool of free *swi6-sm1* molecules is likely to associate with chromatin (H3K9me0 or

H3K9me2/3). Taken together, these results confirm that PAmCherry-*swi6-sm1* is fully

competent to associate with chromatin and is, in fact, more likely to remain bound to

nucleosomes compared to wild-type Swi6. Importantly, the increase in the percentage

of chromatin-bound PAmCherry-*swi6-sm1* molecules leads to a profound depletion of

the mobile PAmCherry-*swi6-sm1* pool and a substantial reduction in the likelihood of

returning to the unbound state from the chromatin-bound state, a feature that potentially

correlates with the fluidity of liquid-like Swi6 condensates[59].

The *swi6-sm1* mutations disrupt the ability of HP1-α/Swi6 to form a condensed,

mobile phase around H3K9me2/3 chromatin

To further interrogate how the *swi6-sm1* mutations affect its global condensation at HC

domains, we imaged diploid cells expressing either (1) one copy of WT Swi6-GFP and

one copy of WT Swi6-mCherry (Figure 4A) or (2) one copy of WT Swi6-GFP and one

copy of *swi6-sm1*-mCherry (Figure 4B). We hypothesized that the depletion of the

condensed, mobile fraction of *swi6-sm1* molecules as revealed by our single molecule

experiments would manifest as a decrease in the total number of Swi6 molecules at HC

foci alongside an otherwise normal pool of relatively immobile *swi6-sm1* molecules.

Indeed, while both Swi6-GFP and *swi6-sm1*-mCherry co-localize within HC domains as

expected due to intact H3K9me2/3 and dimerization properties (Figure 3), WT Swi6

accumulates to a greater extent, forming brighter foci than the mutant *swi6-sm1* (Figure

4B) when compared to images of the WT/WT diploid, which we used to normalize the

scale between the co-localized GFP and mCherry foci (Figure 4C and Figure S5). When

averaged over thousands of HC foci, the mean WT Swi6-GFP foci intensity profiles

were similarly bright across the two diploid strains, while the mean *swi6-sm1*-mCherry

foci intensity was reduced by ~25% compared to the WT Swi6-mCherry foci (Figure

4D). Comparison of the brightest, intermediate, and dimmest foci within each diploid

nucleus revealed similar reductions in *swi6-sm1* concentration, suggesting the mutant

allele affects Swi6's condensation properties across all sizes of HC domains, with bright

foci corresponding to centromeric HC and intermediate to dim foci representing

telomeric and other smaller HC domains (Figure 4E). Our findings suggest that one pool

of Swi6 is bound directly to HC (intact in the *swi6-sm1* allele) while a second pool is

localized to and concentrated at HC domains via its condensation (disrupted in the

swi6-sm1 allele). Taken together, our data argue that the *swi6-sm1* allele represents a

separation-of-function mutant with a specific defect in Swi6 condensation.

H3K9me2/3 and the condensed, mobile pool of Swi6 are required to impart

nuclear stiffness *in vitro*

As we previously demonstrated that peripherally-tethered chromatin at the inner nuclear membrane is a source of mechanical support for the nucleus[12], we sought to evaluate the mechanical contribution of Swi6, and specifically its ability to undergo condensation, on nuclear mechanics. To directly interrogate how fission yeast nuclei respond to explicit magnitudes, time scales, and compression/tension force regimes, we employed an *in vitro* force spectroscopy approach on isolated nuclei that we developed previously[12] to directly measure how altering the extent of H3K9 methylation or loss of HP1-α/Swi6 condensation impacts nuclear stiffness.

Nuclei expressing a GFP fusion of the nucleoporin Cut11 were isolated and used to perform single-nuclei force spectroscopy measurements using optical tweezers (Figure 5). To directly measure its mechanics, a nucleus is sandwiched between two silica beads, one a larger bead adherent to a glass slide with the other, smaller bead trapped in the optical trap (Figure 5A,B). As the larger bead is moved by a piezo stage ($\Delta x_{piezostage}$), the displacement of the smaller bead in the trap is measured (Δx_{bead}); the difference in x-displacement is the amount by which the nucleus is either stretched or compressed ($\Delta x_{nucleus}$). Each oscillation yields force and displacement measurements that can be plotted with respect to each other (Figure 5C). The oscillation data display a linear relationship (black line); the slope (k) of which is equivalent to the spring constant, or effective stiffness, of the individual nucleus. Each nucleus is subjected to multiple rounds of tension and compression at varying time scales, with all oscillations performed over a range of biologically-relevant frequencies: 0.01 Hz – 2 Hz (Figure S6).

The response to compression and tension forces revealed a pronounced mechanical defect in nuclei purified from a strain in which the Suv39/Clr4 H3K9 methylase was deleted, which is necessary for H3K9me2/3 HC domain formation and therefore lies upstream of Swi6. These nuclei displayed a ~50% decrease in effective stiffness over the range of oscillation frequencies (Figure 5D,E). Interestingly, loss of the presumed

H3K9me2/3 demethylase KDM2/Epe1, which leads to a moderate expansion of HC

domains[52] (Figure S3D,F), leads to a ~30% increase in nuclear stiffness (Figure 5D,E),

suggesting that the extent of H3K9 methylation can dictate the mechanical response of

the nucleus to deformation.

We next tested whether Swi6 contributes to the mechanism by which the extent of

H3K9 methylation impacts nuclear stiffness. Strikingly, *swi6Δ* nuclei also had a ~50%

decrease in stiffness, phenocopying *clr4Δ* nuclei (Figure 5D) and suggesting that Swi6

is integral to the mechanism by which H3K9me imparts nuclear stiffness. To investigate

the aspect(s) of Swi6 function that underlie its mechanical contribution, we next

evaluated the mechanical behavior of *swi6-sm1* nuclei. To our surprise, although *swi6-*

sm1 appears to be entirely competent for its direct biochemical association with

H3K9me2/3 nucleosomes (Figure 3), these nuclei are mechanically deficient to the

same extent as *swi6Δ* nuclei (Figure 5D). This finding suggests that it is the mobile

molecules of Swi6 condensed at HC domains (not the molecules bound directly to

H3K9me or nucleosomes) that play a critical role in the contribution of Swi6 to nuclear

stiffness.

We previously observed that fission yeast nuclei display a viscoelastic response[12].

Repeating oscillations at different frequencies can reveal the time-dependent (viscous)

component of the nuclear response (see Figure S6). At longer oscillation frequencies

(100 s or 0.01 Hz), we observed a characteristic reduction in effective stiffness—a non-

linear mechanical response—in wild type, *clr4Δ*, *swi6Δ*, and *swi6-sm1* nuclei, indicating

that these systems are not perfect springs but instead combinations of an elastic spring

and a viscous dashpot, or dampener. To extract the individual mechanical properties of

nuclear stiffness and viscosity, we performed non-linear regression applying the

Maxwell viscoelastic model to our frequency series of effective stiffnesses (Figure 5D,

solid lines). From the viscoelastic fits in (D), the nuclear stiffness was extracted for each

genotype. As compared to the mean stiffness of wild-type nuclei (0.055 pN/nm), *epe1Δ*

nuclei were stiffer (0.073 pN/nm), and *clr4Δ*, *swi6Δ,* and *swi6-sm1* nuclei were softer

(0.041 pN/nm, 0.033 pN/nm, and 0.030 pN/nm respectively) (Figure 5E).

Relative to the viscosity of wild type nuclei (1.72 pN s/nm), *clr4Δ*, *swi6Δ,* and *swi6-sm1*

nuclei were also significantly less viscous (0.51 pN s/nm, 0.58 pN s/nm, and 0.41 pN

s/nm, respectively) (Figure 5F). In contrast, loss of Epe1 also led to an increase in

nuclear viscosity to the extent that we were unable to be determine its value from the

timescales sampled with our oscillation assay (note the complete time-independence of

the measured stiffness in Figure 5D). Thus, despite a rather subtle effect on the

recruitment of *swi6-sm1* to HC domains via its intact H3K9me2/3 binding and

dimerization activities (Figures 3 and 4), the loss of its ability to undergo condensation

and/or to drive coalescence of HC domains is critical to its mechanical contribution to

nuclear mechanics (Figure 5G,H).

Loss of HP1-α/Swi6 condensation increases microtubule-driven nuclear

deformations *in vivo*

To assess how loss of HP1-*α*/Swi6 or abrogation of its condensation affects the nuclear

response to MT dynamics, which primarily exert force through the SPB on the

centromeric HC domain, we monitored nuclear envelope (NE) deformations *in vivo*

(Figure 6A). To evaluate these nuclear deformations with precise spatial and temporal

resolution, the nuclear envelope was labeled by expression of the nucleoporin Cut11-

GFP and visualized in all four-dimensions (4D: x, y, z, and time) by live-cell

fluorescence microscopy (Figure 6B). We next employed a three-dimensional (3D)

surface reconstruction software we previously developed[12,56] applied to series of

fluorescent z-stacks with sub-pixel resolution to capture intrinsic MT-driven

deformations (Figure 6C). As expected, the largest NE deformations manifest at points

subjected to maximum shear forces as MTs drive SPB oscillations, which can be

visualized as local maxima on the nuclear surface (Figure 6B,C,D, left column); these

maxima are lost upon MT depolymerization with the drug carbendazim (MBC) (Figure

6D, right column, and E, dashed lines). When averaged over many nuclei, the largest

NE deformations occur at ~45 and 135 degrees with respect to the long axis of the cell

(Figure 6E, right panel, and Figure S7)—the points at which the SPB is maximally

translated due to the shearing force from the MTs (Figure 6A).

Similar to the mechanical defect seen in our *in vitro* experiments, cells lacking Swi6

displayed larger deformations at all angles as compared to wild-type cells, as reflected

in the root mean squared deformation size in both the presence and absence of MT

dynamics (Figure 6E, left and middle, p < 0.0001). Averaging over hundreds of nuclei

and thousands of individual NE deformations, the locations of the deformations in *swi6Δ*

cells mirrored the wild-type profile, but were shifted towards larger deformations (Figure

6E, right). This suggests that loss of Swi6 leaves the geometry of forces exerted on the

nucleus intact while reducing its ability to resist MT forces (in contrast to loss of

Suv39/Clr4, which cannot be interpreted because this genetic background disrupts the

characteristic MT-induced nuclear deformations, see Figure S8A,B).

Cells lacking Swi6 have a defect in cohesin loading at the centromeres[60], which are

subjected to tangential MT forces *in vivo* (Figure 1). In contrast, the *swi6-sm1* allele

retains the ability to support normal centromeric cohesion[41] and to associate with

H3K9me2/3 nucleosomes (Figure 3), but depletes the mobile phase of Swi6 molecules

and limits HC condensation of smaller domains (Figures 2-4). This allele therefore

provides an opportunity to investigate the Swi6 functions(s) that contribute most to the

response of the nucleus at the centromeric HC-MT interface. We observe that *swi6-sm1*

cells largely recapitulated the increase in nuclear deformability observed in *swi6Δ* cells,

albeit to a slightly lesser extent (Figure 6E, p < 0.0001 as compared to wild type). Taken

together, these observations suggest that defects in Swi6 condensation at HC domains

plays a more dominant role in the global, emergent viscoelasticity of the nucleus as a

system measured *in vitro*, while there is a clear but less quantitative effect on local

deformations in reponse to forces exerted on the centromeric HC-MT interface *in vivo*.

Expansion of HC or crosslinking of H3K9me2/3 nucleosomes by Swi6 are insufficient to impart nuclear stiffness in the absence of a mobile, condensed pool of Swi6

Cells harboring the *swi6-sm1* allele have HC domain clustering defects (Figure 2), reduced Swi6 concentration at HC sites (Figure 4), and ultimately loss of global nuclear stiffness (Figures 5 and 6), all features we couple to the depletion of the mobile pool of Swi6 molecules even as the enhanced association of *swi6-sm1* molecules bound to (and cross-linking of) nucleosomes (Figure 3) is insufficient to retain these features. To further probe this model in the context of WT Swi6, we sought to understand how expansion of HC domains affects Swi6 binding states, and, in turn, how this expansion affects nuclear mechanics. By single molecule tracking, we found that loss of H3K9me2/3 demethylation activity (by deletion of the H3K9me2/3 demethylase, KDM2/Epe1) does not significantly alter the relative populations of bound versus unbound Swi6 as compared to WT cells (Figure 7A, left). However, combined loss of Epe1 and Mst2, a histone acetylase that enhances HC spreading when deleted[52], leads to WT Swi6 being nearly entirely chromatin-bound, while the mobile pool normally found in the condensate is depleted[44] (Figure 7A, right), recapitulating the phenotype seen in *swi6-sm1* cells (Figure 3C). We then tested whether HC spreading affects the *in vivo* nuclear force response. In this *in vivo* assay, which measured the local response to MT dynamics, we observed no significant difference in NE deformations in Epe1-deficient nuclei (Figure 7B and Figure S8C,D). Strikingly, however, we found that *epe1Δmst2Δ* cells displayed much larger NE deformations than WT or *epe1Δ* cells (Figure 7B), showing a paradoxical increase in nuclear deformability when HC spreads. We conclude that enhanced levels of H3K9me2/3 substantial enough to drive depletion of the mobile pool of Swi6 from the condensates compromises rather than reinforces nuclear mechanics.

To further challenge this interpretation, we tested the sufficiency of an engineered H3K9me2/3-containing nucleosome crosslinker comprised of a GST moiety (which inherently drives dimerization[44,61]) fused to one or two units of the Swi6 chromodomain

(CD) to influence nuclear deformability in cells lacking WT Swi6 (Figure 7C). These

heterologous constructs, which lack the domains implicated in Swi6 condensation[44,62],

show a similar depletion of the mobile/unbound state and are found predominantly

bound to chromatin when examined by single molecule tracking (Figure 7D).

Concomitantly, we observed increased nuclear deformability upon expression of *swi6-*

2xCD-GST (Figure 7E), highlighting again that driving H3K9me2/3 nucleosomes into

dense arrays cannot support nuclear mechanics and indeed poisons mechanical

robustness of the nucleus in the absence of WT Swi6, which otherwise would be

condensed at HC domains. Thus, we conclude that Swi6 plays a key role in defining

nuclear stiffness not by its HC-binding and crosslinking abilities but rather by its ability to

undergo condensation into HC-associated droplets.

DISCUSSION

In an effort to understand the contribution of heterochromatin to nuclear mechanics, we

combined the use of specific H3K9me-associated genetic variants with a

complementary array of *in vivo* and *in vitro* assays aimed at elucidating its biophysical

mechanism. Our results support a model in which the mobile, condensed phase of the

HC protein HP1-α/Swi6 is a significant contributor to the viscoelastic behavior of the

nucleus. We propose that the ensemble mechanics of the nucleus arise from a

composite system (individual, HC condensates enmeshed within the chromatin

polymer) that has emergent properties, much like what is seen in aqueous foams (e.g.,

shaving cream)[63]. We suggest that it is this emergent property that provides a

mechanism by which these condensates can support nuclear resistance to potentially

damaging compression or tension forces. Moreover, this property likely influences the

response to forces that can activate changes in the relative amount of HC, specifically

of the H3K9me2/3-associated constitutive HC predominantly found at the nuclear

periphery[64,65,20] that may have large-scale consequences on gene silencing, prevention

of cellular differentiation[66], and genome-protective nuclear softening. We also suggest

that this composite system likely applies to other biomolecular condensates, particularly

those that are enmeshed in biological polymers such as the cytoskeleton.

As described previously for HP1-α and Swi6[22,23,67], our observations support a model in

which Swi6 exhibits liquid-like properties. As telomere clustering is compromised upon

loss of Swi6, expression of the *swi6-sm1* allele, or treatment with 1,6-hexanediol, we

suggest that the formation of Swi6-dependent condensates drives coalescence of

telomeric HC, as has been suggested recently for telomeres in other contexts[45,46,49].

The phenotypic mimicry of the *swi6-sm1* allele supports previous work showing that the

N-terminus of HP1-α/Swi6 serves an essential role in the formation of phase-separated

domains[22,55]. As the *swi6-sm1* allele also confers a silencing defect despite normal

recruitment to H3K9me regions[53], it also offers a potential link between HP1-α/Swi6

condensation and transcriptional silencing, which will require further study.

While recent work affirms that degradation of HP1-α also leads to softer nuclei in

mammalian cells suggested to arise from crosslinking of H3K9me2/3 nucleosomes[21],

our findings specifically suggest an alternate model for the underlying mechanism,

namely that the ability of the mobile phase of HP1-α/Swi6 to undergo condensation is

essential to its contribution to nuclear stiffness. Previously, we showed that the tethering

of chromatin to the nuclear membrane acts as a mechanical scaffold for the nucleus,

and that the nucleus behaves as a viscoelastic material, exhibiting both liquid and solid

characteristics that are time-dependent[12]. This dual nature allows the nucleus to resist

deformation (elasticity/stiffness) while allowing plasticity on longer time scales to enable

chromatin flow within the nucleus (viscosity). The increased nuclear viscosity associated

with expanding H3K9 HC domains in cells lacking Epe1 in our *in vitro* measurements

further implies reduced chromatin flow under these conditions. In this context,

expansion of small HC domains within the chromatin polymer mesh may promote a

more elastic, solid behavior on the minutes timescale (Figure 5D). *In vivo*, however, loss

of Epe1 has little effect on local nuclear deformations due to microtubule shearing force

while further expansion of HC with concomitant loss of Mst2 leads instead to increased

nuclear deformability, suggesting that the loss (and not the gain) of the mobile,

condensed Swi6 pool dominates the local nuclear behavior in response to intrinsic

cytoskeletal forces. This mechanical transformation has potential implications for

physiologic states where cells exhibit gains in repressive H3K9 methylation marks, for

example in response to critical stretching[20,65]. Here, our data suggest that increases in

H3K9 methylation not only act to repress transcription but could also increase nuclear

stiffness and decrease nuclear viscosity when at moderate levels. However, if

pronounced, high levels of H3K9 methylation could instead cause nuclear softening

should it outpace the available pool of unbound HP1-α/Swi6 in the HC-associated

condensate – suggesting a "Goldilocks" balance required to retain both bound and

mobile pools of HP1-α/Swi6 to impart optimal nuclear stiffness.

Loss of HP1-α/Swi6 in the condensate, either through complete deletion of Swi6 or by

expression of the *swi6-sm1* allele, destabilizes the nucleus, reducing both its ability to

resist deformation and impede chromatin flow. This effect is similar to the softer, less

viscous nuclei that lack the H3K9 methyltransferase, Clr4, which controls recruitment of

HC-condensing proteins (including Swi6). The similarity in the mechanical

consequences of disrupting Clr4 or Swi6 is consistent with a model in which the specific

properties of Swi6 associated with HC are necessary for the mechanical effects of

H3K9me2/3 HC to manifest rather than arising through a direct effect of H3K9

methylation levels. This finding also excludes a quantitative contribution of other

H3K9me2/3-binding HC proteins such as Chp1 and Chp2 to nuclear mechanics,

singling out Swi6 as both necessary and unique.

It is important to note that the biological importance of the ability of HP1-α to undergo

condensation *in vitro* on its *in vivo* function has been questioned, for example the failure

of HP1-α over-expression to increase the size of heterochromatic domains as predicted

for simple liquid-liquid phase-separation systems[68]. However, the complexity of

feedback within HC is rife with paradox, for example the requirement for siRNA

transcription to drive chromatin compaction and silencing in fission yeast[69]. Indeed,

Swi6 recruits the H3K9me2/3 demethylase Epe1 almost exclusively into HC domains[70],

which would be expected to decrease available H3K9me2/3 marks and thereby

delocalize H3K9me2/3-engaged Swi6 molecules, likely short-circuiting the ability of

elevated Swi6 levels to drive HC spreading. Moreover, the binding of HP1-α/Swi6 to

H3K9me2/3 and its subsequent compaction of HC involves an integrated set of activities driven by its (1) chromodomain avidity, (2) dimerization, (3) engagement of nucleosome arrays, and (4) its condensation properties, all of which make it challenging to arrive at mechanistic conclusions without clear separation-of-function mutants or through engineered systems[68]. We believe this study, and the use of the *swi6-sm1* allele, provides an additional context in which condensation of HP1-α/Swi6 has a critical function – namely in supporting nuclear mechanics.

Taken together, our results suggest that the mobile, condensed phase of HP1-α/Swi6 at HC domains contributes stiffness to the nucleus, similar to the ability of phase-separated droplets to provide structure to a polymer mesh in which they are embedded[71]. Thus, we have extended the roles for biomolecular condensation *in vivo* beyond cellular compartmentalization to include mechanical functions at the level of micron scale organelles, expanding our perspective on the multi-faceted functionality of condensates in cellular biology.

ACKNOWLEDGEMENTS

We thank members of the King, Mochrie, and Lusk laboratories for helpful discussions, commentary, and troubleshooting, in particular, Hao Yan, Elisa Rodriguez, Andrea Nguyen, and Patrick Lusk. We thank Saikat Biswas for his initial support in creating the PAmCherry-*swi6-sm1* strain for single molecule measurements. We thank the Yeast Genome Resource Center at Osaka University and the many investigators who have deposited strains at this resource. This work was supported by the NIH Pre-doctoral Program for Cellular and Molecular Biology (T32GM00722343), the NSF (CMMI-1334406), the Physical and Engineering Biology Program at Yale, a National Science Foundation Understanding Rules of Life Award (1921677) to JSB and KR, and an NIH award R35GM137832 to KR.

AUTHOR CONTRIBUTIONS

Conceptualization, J.W. and M.K.; Data curation, J.W., I.S., G.R., and H.N.; Formal analysis, J.W., I.S., H.N., Z.C., and M.K.; Funding acquisition, J.B., K.R., S.M. and M.K.; Investigation, J.W., I.S., S.S., Z.C., G.R., Y.H., and M.K.; Methodology, J.W., I.S., Z.C., J.B, K.R., S.M., and M.K.; Project administration, M.K.; Resources, J.B., K.R., S.M. and M.K.; Software, J.W., I.S., Z.C., H.N., and J.B.; Supervision, J.B., K.R., S.M. and M.K.; Validation, J.W., I.S., and M.K.; Visualization, J.W., I.S., G.R., K.R., and M.K.; Writing – original draft, J.W. and M.K.; Writing – review & editing, J.W., I.S., S.S., Z.C., G.R., H.N., Y.H., J.B., K.R., S.M., and M.K.

DECLARATION OF INTERESTS

The authors declare no competing interests.

645
646
Figure 1. Heterochromatin domains exhibit liquid-like properties *in vivo* that are tied to normal telomere clustering

(A) Schematic representation of the distribution of heterochromatin (HC) domains in *S. pombe* as visualized by expression of Swi6-GFP (light green) from its endogenous locus. HC domains are subject to shearing forces driven by microtubule dynamics.

(B) HC domains are deformed in response to MT-driven nuclear oscillations. Time-lapse montage of cells expressing Swi6-GFP shows a large HC focus (blue arrowhead) transitioning from a round to an oblong shape that aligns with the long axis of the cell as it is translated in response to MT dynamics (orange arrowheads, t = 30 to t = 80 s);

image insets display the contour of thresholded Swi6-GFP foci. At t = 60 s, the focus
begins to separate into two foci, and after MT dynamics cease (t = 100 s), these foci
begin to coalesce. Scale bar, 2 µm.
(C,D) Under normal conditions, the six fission yeast telomeres typically cluster into ~1-3
foci during interphase that de-cluster in response to either 1,6-hexanediol treatment or
deletion of Swi6.
(C) Fluorescence maximum intensity projections of representative *S. pombe* cells
expressing Taz1-GFP with 1-6 telomere clusters (see also Figure S2). Scale bar, 2 µm.
(D) Percentage of cells containing the indicated number of Taz1-GFP foci; foci per cell
were counted and grouped by cell length as a proxy for cell cycle progression. In wild-
type cells (n = 130), roughly half the population contained 1-2 (dark blue) or 3-4 (light
blue) Taz1-GFP foci across the cell cycle. Upon exposure to 1,6-hexanediol (n = 135),
Taz1-GFP foci number increased, particularly during G2, indicating telomere de-
clustering. In cells lacking Swi6 (n = 150), significant telomere de-clustering relative to
wild-type cells was observed. * $p < 0.05$; **** $p < 0.0001$ by Fisher's exact test.

700
701
Figure 2. The *swi6-sm1* allele disrupts the ability of HP1-α/Swi6 to promote the coalescence of heterochromatin domains

(A) The *swi6-sm1* allele encodes two mutations in the N-terminal domain of Swi6.

(B) Both wild-type Swi6-GFP and *swi6-sm1*-GFP localize to HC domains.

(C) HC domain size and intensity measured using 3D domain reconstruction. Maximum
intensity projections of transmitted light (upper left) and fluorescence images (upper right)
of *S. pombe* cells expressing Swi6-GFP. Foci are grouped by cell (lower left) and
individually reconstructed in three dimensions (lower right). Scale bars, 6 µm and 2 µm.
(D) The *swi6-sm1* allele leads to a depletion of intermediate scale and intense
heterochromatic domains relative to wild-type Swi6-GFP. Foci intensity versus cross-
sectional area density scatter plots for Swi6-GFP (n = 2500 foci) and *swi6-sm1*-GFP (n =
2402 foci) labeled HC domains. **** $p < 0.0001$ by Kolmogorov-Smirnov test after
calculation of cumulative distributions.
(E) The *swi6-sm1* allele leads to telomere de-clustering in a manner similar to 1,6-
hexanediol exposure and *swi6Δ*. Data expressed as (and wild-type data replotted from)
in Figure 1D. In cells expressing *swi6-sm1* (n = 145), the de-clustering effect recapitulates
that seen with 1,6-hexanediol-treated cells or cells lacking Swi6. *** $p < 0.001$; **** $p <$
0.0001 by Fisher's exact test.
(F) Summary of HC domain phenotypes. In wild-type cells, HC domains form discrete
clusters. Treatment with 1,6-hexanediol, deletion of Swi6, or expression of the N-terminal
mutant *swi6-sm1* causes HC domains to decluster.

Figure 3. The *swi6-sm1* allele shows a depletion of mobile states despite normal chromatin and H3K9me2/3 binding *in vivo*

(A) Schematic representation of Swi6. Each domain has a distinct biochemical property. NT: unstructured N-terminus subject to phosphorylation that regulates Swi6 condensation *in vitro* (*swi6-sm1* has two mutations as indicated); CD: H3K9me binding; H: nucleic acid binding; CSD: protein oligomerization.

(B) Single-molecule tracking photoactivation localization microscopy (spt-PALM) of wild-type Swi6 and *swi6-sm1* in the *S. pombe* nucleus. 1-3 PAmCherry-Swi6 molecules/nucleus are photoactivated (406 nm, 1.50-4.50 W/cm2, 200 ms), then imaged and tracked until photobleaching (561 nm, 71.00 W/cm2, 25 frames/sec). The cycle is repeated 10-20 times/cell.

(C) SMAUG identifies four distinct mobility states, (α blue, β orange, γ yellow, and δ purple) for PAmCherry-Swi6 and PAmCherry-*swi6-sm1*. Each point represents the

average single-molecule diffusion coefficient, D, versus weight fraction of Swi6/*swi6-sm1* molecules in that state at a saved iteration of the Bayesian algorithm after convergence. Dataset, PAmCherry-*swi6-sm1*: 29,657 steps from 3,744 trajectories and 97 cells. Crossbars: comparison to PAmCherry-Swi6 results from reference [44] with 10,095 steps from 1,491 trajectories and 36 cells.

(D) Average probabilities (arrows) of a PAmCherry-Swi6 molecule transitioning between the mobility states (circles) from reference[44]. Each circle area is proportional to the weight fraction, π; D is in μm^2/s. Low-significance transition probabilities below 0.04 are excluded.

(E) Average probabilities (arrows) of a PAmCherry-*swi6-sm1* molecule transitioning between the mobility states (circles) as in (D) (see also Table S1).

Figure 4. The *swi6-sm1* mutations disrupt the ability of HP1-*α*/Swi6 to form a condensed, mobile phase around H3K9me2/3 chromatin

(A) Maximum intensity projections of fluorescence images of diploid *S. pombe* cells expressing Swi6-GFP/Swi6-mCherry. Inset shows a representative nucleus with detected Swi6 domains marked by intensity as bright (red, the brightest domain in the nucleus), dim (blue, the dimmest domain), and intermediate (cyan, all the remaining domains). Right panel shows distribution of "bright", "intermediate", "dim" domains for combined domains across all cells with ≥ 3 foci. For display the intensity between GFP and mCherry was normalized (calibration from (C))**.** Scale bar, 2 µm, inset, 1 µm.

(B) Same as (A) but for cells expressing Swi6-GFP/*swi6-sm1*-mCherry and using the same calibration between channels as in (A).

(C) Scatter plot of Swi6 foci intensities in diploid cells from nuclei with ≥ 3 foci, shown as mCherry signal versus GFP signal for Swi6-GFP/Swi6-mCherry cells (black, n = 584 nuclei; 2,648 foci) and Swi6-GFP/*swi6-sm1*-mCherry cells (purple, n = 538 nuclei; 2,327 foci). Red lines are best fit of a linear model y(x) = a*x+b, with a = 0.4495, b = 1.142 x 10^5 and a = 0.3379, b = 8.655 x 10^4, respectively. Values in parentheses are 95% confidence intervals. Coefficient a = 0.4495 was used to rescale mCherry channel fluorescence in (A) and (B) (see also Figure S5).

(D) Average intensity profile for GFP and mCherry fluorescence for Swi6 domains after mCherry signal rescaling using calibration from (C).

(E) Pixel intensity as a function of distance from the center of Swi6 domain (determined in GFP channel) for GFP and mCherry signal for "bright", "intermediate", and "dim" domains. Upper panels: Swi6-GFP/Swi6-mCherry cells; bottom panels: Swi6-GFP/*swi6-sm1*-mCherry cells. * p < 0.05; ** p < 0.01; **** p < 0.0001 by two-sample Kolmogorov-Smirnov test.

Figure 5. H3K9me2/3 and the condensed, mobile pool of Swi6 are required to impart nuclear stiffness as assessed *in vitro*

(A) Transmitted light image showing the force spectroscopy configuration as viewed from above. Scale bar, 2 µm.

(B) Schematic representation of the force spectroscopy assay.

(C) Representative force versus extension curve of an isolated wild type nucleus. Each data point (gray) represents the force applied to the nucleus and the resulting displacement over one oscillation (extension and compression). The data display a linear relationship (black line); the slope (k) is equivalent to the spring constant, or effective stiffness, of the individual nucleus.

(D) Effective stiffnesses of wild type nuclei (black, n = 8 nuclei; 63 oscillations), KDM2/Epe1-deleted nuclei (pink, n = 18 nuclei; 172 oscillations), Suv39/Clr4-deleted nuclei (orange, n = 3 nuclei; 22 oscillations), HP1-α/Swi6-deleted nuclei (green, n = 5 nuclei; 40 oscillations), and *swi6-sm1*-expressing nuclei (purple, n = 3 nuclei; 29 oscillations) over a range of oscillation frequencies (0.01 – 2 Hz). Data are represented as mean +/- SD of the raw oscillation data at each oscillation frequency.

(E) Nuclear stiffnesses extracted from viscoelastic fit of data in (D). Data are represented as mean +/- SD of the fitted values.

(F) Nuclear viscosity extracted from viscoelastic fit of data in (D). Data are represented as mean +/- SD of the fitted values. N.D., not determined.

(G) Models of H3K9me HC with (left) and without (right) Swi6 condensation. Nucleosomes (gray) are methylated on histone H3K9 (yellow) by the methyltransferase Suv39/Clr4 (orange). Upon methylation, the HC protein Swi6 (green) binds H3K9me2/3, crosslinking adjacent nucleosomes to form compact HC. Condensation of additional, unbound Swi6 molecules form an extended phase nucleated at these H3K9me2/3 chromatin regions (represented by green cloud). The demethylase KDM2/Epe1 (pink) reverses this process, countering HC formation by promoting H3K9me2/3 demethylation. *swi6-sm1* (purple) binds to, crosslinks, and compacts H3K9me2/3 nucleosomes normally but disrupts the ability to condense and form this mobile phase.

(H) Schematic table representing the mechanical effects of altering H3K9me and binding proteins. $* p < 0.05$; $** p < 0.01$; $*** p < 0.001$; $*** p < 0.0001$ by ordinary one-way ANOVA with Dunnett's correction for multiple comparisons.

Figure 6. Loss of HP1-α/Swi6 or its condensation increases microtubule-driven nuclear deformations *in vivo*

(A) In *S. pombe*, shearing forces acting upon the nuclear envelope (red) arise from polymerizing microtubules (MTs, black) at the cell tips that drive oscillation of the associated spindle pole body (SPB, tan) and are transmitted to the centromeric HC (large green domain).

(B) Fluorescence images of a wild. type *S. pombe* cell expressing Cut11-GFP demonstrating a typical nuclear deformation (arrowhead). Scale bar, 2 μm.

(C) 3D reconstructions of the nucleus. At t = 0 (left), the nucleus is circular; ~55 seconds later (right), the same nucleus undergoes a MT-dependent deformation (arrowhead).

(D) Representative single nuclear surface deformations with (left column) and without (right column) MT dynamics shown as topological heat maps with respect to its latitudinal angle (points above and below the nuclear "equator")—a band between +45 and -45 degrees—and its longitudinal angle (points around the nucleus) between 0 and 360 degrees (see also Figure S7).

(E) Root mean squared deformation size across all nuclei observed (left and middle) and all nuclear radial angles (right): wild type (black, n = 760 nuclei; 12,556 deformations), *swi6Δ* (green, n = 218 nuclei; 4,421 deformations), *swi6-sm1* (purple, n = 371 nuclei; 6,583 deformations). MBC-treated nuclei shown as dashed lines: wild type (black, n = 58 nuclei; 561 deformations), *swi6Δ* (green; n = 69 nuclei; 976 deformations), *swi6-sm1* (purple, n = 36 nuclei; 534 deformations). Data are represented as mean (+/- SEM in right plots in (E)). **** p < 0.0001 by Mann-Whitney U test.

Figure 7. Expansion of HC or crosslinking of H3K9me2/3 nucleosomes by Swi6 are insufficient to impart nuclear stiffness in the absence of a mobile, condensed pool
(A) Depletion of both Epe1 and Mst2 (but not Epe1 alone), which drives heterochromatin spreading, leads to depletion of the mobile pool PA-mCherry-Swi6. SMAUG analysis and plots as in Figure 3C. Individual points represent the cells lacking Epe1 (left) or Epe1 and Mst2 (right) correspond to the indicated mobility states with the wild-type genotype shown by the cross-hatches (data from reference [44]).
(B) Expanding HC domains drives larger nuclear deformations. Root mean squared deformation size across all nuclei observed (left and middle) and all nuclear radial angles (right): wild type (black, n = 760 nuclei; 12,556 deformations, replotted from Figure 6E), *epe1Δ* (pink, n = 215 nuclei; 3,610 deformations), *epe1Δmst2Δ* (red, n = 361 nuclei; 8,419 deformations). Data are represented as mean (+/- SEM in right plots in (B)).
(C) Diagram of wild-type Swi6 that assembles by oligomerization mediated by the chromoshadow domains (top) and an engineered H3K9me nucleosome crosslinking construct, PA-mCherry-*swi6-2xCD-GST* (developed in reference [44]).
(D) The PA-mCherry-*swi6-2xCD-GST* construct (light colors, data from reference [44]; 42,382 steps from 5,182 trajectories and 140 cells), which lacks the domains mediating Swi6 condensation, largely phenocopies the diffusive states displayed by PAmCherry-*swi6-sm1* (dark colors). Wild-type PAmCherry-Swi6 (crosshatches) and PAmCherry-*swi6-sm1* data are re-plotted from Figure 3C (see also Table S1).
(E) Expression of the *swi6-2xCD*-GST construct leads to larger nuclear deformations. Root mean squared deformation size across all nuclei observed (left and middle) and all nuclear radial angles (right): wild type with PAmCherry-Swi6 (black, n = 1,502 nuclei; 27,059 deformations), *swi6Δ* (green, n = 218 nuclei; 4,421 deformations; replotted from Figure 6E), *swi6-sm1* (purple, n = 371 nuclei; 6,583 deformations; replotted from Figure 6E), and *swi6-2xCD-GST* (teal, n = 329 nuclei; 6,055 deformations). Data are represented as mean (+/- SEM in right plots in (E)). **** p < 0.0001 by Mann-Whitney U test.

STAR METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Megan King (megan.king@yale.edu).

Materials availability

All unique/stable reagents and strains generated in this study are available from the Lead Contact without restriction.

Data and code availability

The code generated during this study (3DMembraneReconstruction, Swi6/*swi6-sm1* diploid heterochromatin domain detection and analysis, TweezersAnalysis, and NuclearFluctuations) is available on GitHub at https://github.com/LusKingLab/.

METHOD DETAILS

Cell culture and strain generation

The strains used in this study are listed in the Key Resources Table (see also reference[72]). *S. pombe* cells were grown and maintained in standard cell culture conditions at 30°C[73]. Genetic knock-outs were made by gene replacement of the open reading frames with the kanMX6[74], hphMX6, or natMX6 cassettes[75]. C- and N-terminal GFP tagging was performed with pFa6a-GFP-kanMX6[74] or pFa6a-natMX6-Pnmt41[74] respectively; the pFa6a-mCherry-kanMX6 cassette was also used as a template for C-terminal mCherry tagging[76]. To edit PAmCherry-Swi6 to generate PAmCherry-Swi6-sm1, we used a CRISPR-Cas9 based approach. sgRNAs targeting the N-terminus of Swi6 were generated and transformed as previously described in Torres-Garcia *et al.*[77] along with a gene block (Twist Biosciences, CA) that contains the sm1 allele mutations

(D21G, ΔE74) and synonomous substitutions that destabilize the sgRNA binding site.

The resulting *S.pombe* colonies were screened using Sanger sequencing to confirm the

desired mutation[77]. All strains generated by cassette integration were confirmed by PCR

and/or DNA sequencing. Strains made through genetic crosses were confirmed by the

segregation of selection markers and/or by the presence of the designated

fluorescently-tagged protein.

1030

1031

Standard live-cell microscopy

S. pombe strains were grown in YE5S plus 250 mg/L adenine to log phase (OD$_{600}$ 0.5-

0.8). Cells were mounted on agarose pads (1.4% agarose in EMM5S) and sealed with

VALAP (1:1:1, vaseline:lanolin:paraffin). Cells treated with MBC were incubated on agar

pads with 50 mg/mL MBC for 10 min before imaging. Cells treated with 1,6-hexanediol

were incubated with final concentrations of 5% 1,6-hexanediol for 15 or 30 min as

indicated in the legend and imaged directly thereafter. Live-cell images were acquired

on a Deltavision Widefield Deconvolution Microscope (Applied Precision/GE

Healthcare) with a CoolSnap HQ2 CCD camera (Photometrics) or an Evolve 512

EMCCD camera (Photometrics). For Swi6-GFP time-lapse imaging, single mid-plane z-

slices with 0.15-sec exposure time were taken every 5 sec with the CCD camera. For

telomere/Taz1-GFP imaging, 15 z-slices with 50-ms exposure time and 320-nm spacing

were taken with the EMCCD camera. For Swi6/*swi6-sm1*-GFP imaging, 16 z-slices with

10-ms exposure time and 400-nm spacing with taken with the EMCCD camera. For

Swi6/ *swi6-sm1* diploid imaging, 21 z-slices with 500-ms exposure time and 200-nm

spacing with taken with the EMCCD camera. For nuclear contour imaging of Cut11-

GFP, 10 z-slices with 20-ms exposure time and 400-nm spacing were taken every 2.5 s

for 5 min with the CCD camera. For GFP-Swi6 imaging, 20 z-slices with 0.1-sec

exposure time and 200-nm spacing with taken with the EMCCD camera. For Chp1-GFP

imaging, 20 z-slices with 0.2-sec exposure time and 200-nm spacing were taken with

the CCD camera.

1053

1054

Heterochromatin shape and telomere analysis

To characterize the shape dynamics of Swi6-GFP foci, single mid-plane z-slices were read in MathWorks MATLAB R2016b (Natick, MA, USA), binarized with a pixel threshold value of 0.8, and focus boundary x and y values were found using the function bwboundaries. X and y values for each Swi6-GFP focus shape was plotted using the MATLAB fill function.

For Taz1-GFP and cell length analysis only live, interphase cells that had completed cytokinesis were considered. Maximum intensity projections were displayed as described in the Figure legends. Images were analyzed in ImageJ (Fiji;[78]). To calculate the number of Taz1-GFP foci per cell, maximum Z-projections were generated and total number of Taz1-GFP foci per cell were counted. For statistical comparison, percentage of cells treated with 1,6-hexanediol, *swi6Δ* cells, and *swi6-sm1* cells were individually compared to wild-type cells for each cell length (a proxy for cell cycle stage) using Fisher's exact test.

Three-dimensional heterochromatin foci analysis

GFP-Swi6, Swi6-GFP, and *swi6-sm1*-GFP cross-sectional foci areas and intensities were measured using three-dimensional reconstructions of GFP foci with a custom MATLAB software as previously described[56]. Foci were grouped by cell using a minimum separation distance of 40-50 pixels from the center of the grouping; this adequately grouped together foci of the same nucleus while excluding foci from nearby, neighboring cells. Z-stack images for each focus were individually fit with a three-dimensional Gaussian function to determine maximum focus width in both x and y-dimensions as well as foci integrated intensity. Foci intensity vs. cross-sectional area plots and histograms were generated using the MATLAB functions dscatter[79] and scatterhist. For statistical comparison, Swi6-GFP (wild type) and *swi6-sm1*-GFP foci areas and intensities were compared using the Kolmogorov-Smirnov test.

Single-molecule tracking photoactivation localization microscopy (spt-PALM)

Yeast strains containing a copy of *swi6-sm1*-PAmCherry under the control of the native Swi6 promoter were grown in standard complete YES media (US Biological, cat. Y2060) containing the full complement of yeast amino acids and incubated overnight at 32 °C. The seed culture was diluted and incubated at 25 °C with shaking to reach an OD_{600} ~0.5. To maintain cells in an exponential phase and eliminate extranuclear vacuole formation, the culture was maintained at OD_{600} ~0.5 for 2 days with dilutions performed at ~12-hour time intervals. To prepare agarose pads for imaging, cells were pipetted onto a pad of 2% agarose prepared in YES media, with 0.1 mM N-propyl gallate (Sigma, cat. P-3130) and 1% gelatin (Millipore, cat. 04055) as additives to reduce phototoxicity during imaging. *S. pombe* cells were imaged at room temperature with a 100× 1.40 NA oil-immersion objective in an Olympus IX-71 inverted microscope. First, the fluorescent background was decreased by exposure to 488-nm light (Coherent Sapphire, 200 W/cm^2 for 20 – 40 s). A 406-nm laser (Coherent Cube 405-100; 102 W/cm^2) was used for photo-activation (200-ms activation time) and a 561-nm laser (Coherent-Sapphire 561-50; 163 W/cm^2) was used for imaging. Images were acquired at 40-ms exposure time per frame. The fluorescence emission was filtered with Semrock LL02-561-12.5 filter and Chroma ZT488/561rpc 488/561 dichroic to eliminate the 561-nm excitation source and imaged using a 512 × 512-pixel Photometrics Evolve EMCCD camera.

Single-molecule trajectory analysis with SMAUG algorithm

Recorded *swi6-sm1*-PAmCherry single-molecule positions were detected and localized with 2D Gaussian fitting with home-built MATLAB software as previously described and connected into trajectories using the Hungarian algorithm[80,81]. These single-molecule trajectory datasets were analyzed by a non-parametric Bayesian framework to reveal heterogeneous dynamics[58]. This SMAUG algorithm uses non-parametric Bayesian statistics and Gibbs sampling to identify the number of distinct mobility states, *n*, in the single-molecule tracking dataset in an iterative manner. It also infers the weight fraction, π_i, and effective diffusion coefficient, D_i, for each mobility state ($i = 1,…,n$), assuming a

Brownian motion model. We ran the algorithm over >10,000 iterations to achieve a

thoroughly mixed state space. The state number and associated parameters were

updated in each iteration of the SMAUG algorithm and saved after convergence. The

final estimation shows the data after convergence for iterations with the most frequent

state number. Each mobility state, i, is assigned a distinct color, and for each saved

iteration, the value of D_i is plotted against the value of π_i. The distributions of estimates

over the iterations give the uncertainty in the determination of D_i. Furthermore, the

transition probabilities (e.g., Figure 3D,E) give the average probability of transitioning

between states from one step to another in any given trajectory.

1126

1127

Swi6/*swi6-sm1* diploid heterochromatin domain detection and analysis

Swi6 HC domains were detected in 3D images in both GFP and mCherry channels

independently using custom-written MATLAB code that finds irregular fluorescent blobs

regardless of their shape

(https://github.com/LusKingLab/Spot_Detection_n_Colocalization). First, the code

identifies all pixels that are brightest within sub-volumes of a given size. Next, it

calculates ratio of the mean intensities of "central" pixels (*i.e.*, pixels within a given

radius R_{cent}, from the brightest pixel) versus "edge" pixels (*i.e.*, pixels within a spherical

shell of a given thickness d_{shell} ith R_{edge} sphere radius). If the ratio is above a given

threshold, the code keeps the identified spot and calculates its 3D position as a

intensity-weighted centroid of pixels within R_{fit} radius, and its peak intensity as a mean

intensity of pixels within R_{fit} radius. After spot identification, individual GFP and mCherry

spots were paired if the peak positions were within 3 pixel distance. These paired spots

were used to compare intensities between GFP and mCherry signals and to calibrate

channels relative to each other (using data from cells bearing only wild-type Swi6 -

Swi6-GFP/Swi6-mCherry cells), and to compare recruitment of Swi6-mCherry versus

swi6-sm1-mCherry.

1145

To analyze data for centromeric HC domains versus other HC domains, domains

detected in the GFP channel were grouped into nuclei based on spatial proximity. Next,

for each nucleus, the nuclear background intensity was determined as a median

intensity value for all pixels within R_{nuc} radius. Then, for each nucleus with at least two

spots (n_{spots} > 2), the peak intensity of each spot was assigned to the "bright" (*i.e.*,

centromeric), "intermediate", or "dim" class. Note that in this "by nucleus" definition a

bright spot from one nucleus might be dimmer than an intermediate spot from another.

The peak intensities of the classified spots in the GFP channel were then compared to

their paired intensities in the mCherry channel. To determine the calibration coefficient

between GFP and mCherry channels, data from "centromeric" ("bright") domains below

an intensity threshold (to avoid the influence of few very bright domains) after correction

for different exposure time were fitted by *ImCherry* = a*$\underline{I_{GFP}}$ + b, resulting in calibration

coefficient a = 0.4412, that was then used in the subsequent analysis.

1159

1160

Nuclear isolation

Nuclei were isolated as previously described in [12]. Briefly, strains were grown to log

phase in rich media (YE5S) at 30°C and diluted into 1 liter YE5S for overnight growth.

Cells were harvested the following day at an OD_{600} ~0.8 and incubated in 100 mM Tris

pH 9.4 and 10 mM DTT for 10 min at 33°C to prepare for spheroplasting. Cells were

spheroplasted in 0.4 mg/mL zymolase (MP Biomedicals), 0.6 mg/mL lysing enzymes

(Sigma), 350 μL beta-glucuronidase (MP Biomedicals) and 5 mM DTT in 1.1 M sorbitol

at 33°C for 2-3 hours. Spheroplasts (nuclei) were isolated from cells by centrifugation

using a series of three density-step sucrose gradients, and aliquots were flash frozen in

liquid N_2 and stored at -80°C. To thaw for use, one aliquot was dialyzed overnight in 500

mL dialysis buffer (80 mM PIPES, 5% DMSO, 2 mM $MgCl_2$, 1 mM EGTA, 500 mM

sucrose).

1173

1174

Force spectroscopy oscillation assay

Force spectroscopy was performed on isolated nuclei as described previously in

Schreiner *et al.*[12]. In a flow cell, isolated *S. pombe* nuclei were trapped and fixed via

electrostatic interactions to poly-ornithine-coated silica beads by micromanipulation

using optical tweezers. Optical trapping and interferometry position detection was

achieved using a single-beam gradient focus from a 3-watt 1064 nm ND:YAG laser. The

focused laser was used to trap beads and nuclei within the flow cell, which was

mounted onto a piezo-controlled stage with nanometer precision. To visualize each

nucleus, an epifluorescence microscope was coupled with the optical tweezers,

equipped with a 470 nm excitation source (blue light emitting diode). Each nucleus

(using the optical tweezers) was affixed on its right side to a 5.2 μm bead adherent to a

glass coverslip, which elevates the nucleus off the coverslip and provides a stationary

structure upon which forces can be applied. Next, a smaller 1.2 μm bead (using the

optical tweezers) is affixed to the left (free) side of the nucleus, creating a bead-nucleus-

bead sandwich. While holding the smaller bead within the optical trap, the large bead

and coverslip—attached to a piezostage—is respectively translated away and towards

the optical trap by a pre-determined displacement ($\Delta x_{piezostage}$) to exert respective

tension or compression on the sandwiched nucleus. The resulting displacement of the

small bead in the optical trap (Δx_{bead}) is measured by its deflection within the trap using

back focal plane interferometry via an infrared-enhanced quadrant photodiode. The

difference in the two displacements is equal to the displacement of the nucleus

($\Delta x_{nucleus}$). To exert compression and tension forces, the stage is driven sinusoidally with

an amplitude range of 60-100 nm (physiologic length scales), at discrete frequencies of

0.01-2 Hz.

1199

1200

Force spectroscopy data analysis

To determine the effective stiffness of the nucleus ($k_{nucleus}$), the system was treated as a

spring that adheres to Hooke's law ($F = k_{nucleus}\Delta_{nucleus}$), where $\Delta_{nucleus}$ is the

displacement of the nucleus (either in extension or compression) and F is the force

imparted to the nucleus from the oscillating stage. $\Delta_{nucleus}$ was calculated as the

difference between the piezo stage displacement ($\Delta_{piezostage}$) and the displacement of

the small bead from the center of the trap (Δ_{bead}). The force F on the nucleus is equal

and opposite to the force on the bead in the trap, therefore determining the stiffness of

the trap allowed us to calculate the force acting upon the bead and thus the force acting

upon the nucleus. The inherent stiffness of the optical trap was determined for each

experiment by fitting the measured power spectral density of a trapped 1.2 μm bead

with a Lorentzian function to extract the corner frequency[82]. The trap stiffness is the

ratio of this corner frequency to the bead's friction coefficient. We subjected each

nucleus to multiple rounds of compression and extension at different frequencies (0.01

– 2 Hz), enabling us to observe any time-dependent mechanical properties. We then fit

the effective stiffness data with a Maxwell viscoelastic model to extract the elastic

(spring) and viscous (dashpot/dampener) components, K and η respectively, using the

corresponding fitting function for the effective stiffness:

1219

1220
$$k(f) = \frac{2\pi K\eta f}{\sqrt{(2\pi\eta f)^2 + K^2}}$$

1221

1222

Nuclear envelope deformation analysis

NE contour fitting and deformation time analysis were performed as previously

described in [12] and [56]. Briefly, our contour fitting algorithm reconstructed the three-

dimensional nuclear shape with sub-pixel resolution from a z-stack of 10 two-

dimensional images by optimizing NE shape to simultaneously maximize the total

intensity of the nuclear envelope across the z-stack and minimize the NE's total

curvature. To avoid the effects of aberration and difficulties with image noise at the

nuclear caps in the z-dimension, we limited our deformation analysis to a band 45

degrees above and below the center/"equator" of the nucleus (Figure 6D and Figure

S7a). After the 3D shape reconstruction at each time point (denoted by the radial

distance from the nuclear center r(θ, φ, t) at vertex angle (θ, φ) at frame t), a time-

average shape $R_{av}(\theta, \varphi) = <r(\theta, \varphi, t)>_t$ was calculated and shape deformations were

detected by deviation of the radial distance from a time-averaged radial distance at that

same (θ, φ) position. The average, or root mean squared, deformation size was

calculated according to:

1238

1239

$$RMSF(\theta, \varphi) = \frac{1}{T} \sum_{t=1}^{T} \sqrt{(r(\theta, \varphi, t) - \langle r(\theta, \varphi, t) \rangle_t)^2}$$

1240

Where T is the total number of frames, $r(\theta, \varphi, t)$ is the radial distance from the nuclear

center at vertex angle (θ, φ) at frame t and $< r(\theta, \varphi, t) >_t$ denotes the time-averaged

radial distance. Individual deformations were tracked over time using 2D particle

tracking, assigning longer deformations (> 10 frames/25 s) to those induced by

microtubules and shorter deformations arising from baseline thermal activity. Peak

height, formation (rise) time, and resolution (decay) time were determined by fitting each

deformation trajectory to a single, asymmetric triangle waveform.

1248

1249

1250   **QUANTIFICATION AND STATISTICAL ANALYSIS**

1251

Statistical details (statistical tests used, exact value for n, what n represents, definition

of center, and dispersion and precision of measures) for each experiment can be found

in the Figure legends. Microsoft Excel (Redmond, WA, USA) was used to tabulate

values, GraphPad Prism 9 (La Jolla, CA, USA) was used to create simple plots (Figure

1D, Figure 2E, Figure 5C, D, E, F, Figure S1C, D, Figure S3F, Figure S4, and Figure

S6) and perform statistical analysis, and MathWorks MATLAB R2016b (Natick, MA,

USA) was used for all other analysis and plotting. $* p < 0.05$; $** p < 0.01$; $*** p < 0.001$;

$**** p < 0.0001$ by Fisher's exact test, ordinary one-way ANOVA with Dunnett's multiple

comparisons test, unpaired t-test, Kolmogorov-Smirnov test, or Mann-Whitney U test.

1261

1262

1263   **SUPPLEMENTAL ITEM TITLES**

1264

1265   **Table S1. Diffusion coefficients and weights from Figures 3C and 7D**

1266

1267

1268

REFERENCES

1. Lusk, C.P., and King, M.C. (2017). The nucleus: keeping it together by keeping it apart. Current Opinion in Cell Biology *44*, 44–50. 10.1016/j.ceb.2017.02.001.

2. Shah, P., Wolf, K., and Lammerding, J. (2017). Bursting the Bubble - Nuclear Envelope Rupture as a Path to Genomic Instability? Trends Cell Biol *27*, 546–555. 10.1016/j.tcb.2017.02.008.

3. De Magistris, P., and Antonin, W. (2018). The Dynamic Nature of the Nuclear Envelope. Curr Biol *28*, R487–R497. 10.1016/j.cub.2018.01.073.

4. Beck, M., and Hurt, E. (2017). The nuclear pore complex: understanding its function through structural insight. Nat Rev Mol Cell Biol *18*, 73–89. 10.1038/nrm.2016.147.

5. de Leeuw, R., Gruenbaum, Y., and Medalia, O. (2018). Nuclear Lamins: Thin Filaments with Major Functions. Trends Cell Biol *28*, 34–45. 10.1016/j.tcb.2017.08.004.

6. Raab, M., Gentili, M., de Belly, H., Thiam, H.-R., Vargas, P., Jimenez, A.J., Lautenschlaeger, F., Voituriez, R., Lennon-Duménil, A.-M., Manel, N., et al. (2016). ESCRT III repairs nuclear envelope ruptures during cell migration to limit DNA damage and cell death. Science *352*, 359–362. 10.1126/science.aad7611.

7. Denais, C.M., Gilbert, R.M., Isermann, P., McGregor, A.L., te Lindert, M., Weigelin, B., Davidson, P.M., Friedl, P., Wolf, K., and Lammerding, J. (2016). Nuclear envelope rupture and repair during cancer cell migration. Science *352*, 353–358. 10.1126/science.aad7297.

8. Vargas, J.D., Hatch, E.M., Anderson, D.J., and Hetzer, M.W. (2012). Transient nuclear envelope rupturing during interphase in human cancer cells. Nucleus *3*, 88–100. 10.4161/nucl.18954.

9. Hatch, E.M., and Hetzer, M.W. (2016). Nuclear envelope rupture is induced by actin-based nucleus confinement. Journal of Cell Biology *215*, 27–36. 10.1083/jcb.201603053.

10. Harada, T., Swift, J., Irianto, J., Shin, J.-W., Spinler, K.R., Athirasala, A., Diegmiller, R., Dingal, P.C.D.P., Ivanovska, I.L., and Discher, D.E. (2014). Nuclear lamin stiffness is a barrier to 3D migration, but softness can limit survival. Journal of Cell Biology *204*, 669–682. 10.1083/jcb.201308029.

11. Lammerding, J., Fong, L.G., Ji, J.Y., Reue, K., Stewart, C.L., Young, S.G., and Lee, R.T. (2006). Lamins A and C but Not Lamin B1 Regulate Nuclear Mechanics. Journal of Biological Chemistry *281*, 25768–25780. 10.1074/jbc.M513511200.

12. Schreiner, S.M., Koo, P.K., Zhao, Y., Mochrie, S.G.J., and King, M.C. (2015). The tethering of chromatin to the nuclear envelope supports nuclear mechanics. Nat Commun *6*, 7159. 10.1038/ncomms8159.

13. Stephens, A.D., Banigan, E.J., Adam, S.A., Goldman, R.D., and Marko, J.F. (2017). Chromatin and lamin A determine two different mechanical response regimes of the cell nucleus. MBoC *28*, 1984–1996. 10.1091/mbc.e16-09-0653.

14. Stephens, A.D., Liu, P.Z., Banigan, E.J., Almassalha, L.M., Backman, V., Adam, S.A., Goldman, R.D., and Marko, J.F. (2018). Chromatin histone modifications and rigidity affect nuclear morphology independent of lamins. Mol Biol Cell *29*, 220–233. 10.1091/mbc.E17-06-0410.

15. Bickmore, W.A., and van Steensel, B. (2013). Genome architecture: domain organization of interphase chromosomes. Cell *152*, 1270–1284. 10.1016/j.cell.2013.02.001.

16. Bustin, M., and Misteli, T. (2016). Nongenetic functions of the genome. Science *352*, aad6933. 10.1126/science.aad6933.

17. Gonzalez-Sandoval, A., and Gasser, S.M. (2016). Mechanism of chromatin segregation to the nuclear periphery in *C. elegans* embryos. Worm *5*, e1190900. 10.1080/21624054.2016.1190900.

18. Falk, M., Feodorova, Y., Naumova, N., Imakaev, M., Lajoie, B.R., Leonhardt, H., Joffe, B., Dekker, J., Fudenberg, G., Solovei, I., et al. (2019). Heterochromatin drives compartmentalization of inverted and conventional nuclei. Nature *570*, 395–399. 10.1038/s41586-019-1275-3.

19. King, M.C., Drivas, T.G., and Blobel, G. (2008). A Network of Nuclear Envelope Membrane Proteins Linking Centromeres to Microtubules. Cell *134*, 427–438. 10.1016/j.cell.2008.06.022.

20. Nava, M.M., Miroshnikova, Y.A., Biggs, L.C., Whitefield, D.B., Metge, F., Boucas, J., Vihinen, H., Jokitalo, E., Li, X., García Arcos, J.M., et al. (2020). Heterochromatin-Driven Nuclear Softening Protects the Genome against Mechanical Stress-Induced Damage. Cell *181*, 800-817.e22. 10.1016/j.cell.2020.03.052.

21. Strom, A.R., Biggs, R.J., Banigan, E.J., Wang, X., Chiu, K., Herman, C., Collado, J., Yue, F., Ritland Politz, J.C., Tait, L.J., et al. (2021). HP1α is a chromatin crosslinker that controls nuclear and mitotic chromosome mechanics. Elife *10*, e63972. 10.7554/eLife.63972.

22. Larson, A.G., Elnatan, D., Keenen, M.M., Trnka, M.J., Johnston, J.B., Burlingame, A.L., Agard, D.A., Redding, S., and Narlikar, G.J. (2017). Liquid droplet formation by HP1α suggests a role for phase separation in heterochromatin. Nature *547*, 236–240. 10.1038/nature22822.

23. Strom, A.R., Emelyanov, A.V., Mir, M., Fyodorov, D.V., Darzacq, X., and Karpen,
G.H. (2017). Phase separation drives heterochromatin domain formation. Nature
547, 241–245. 10.1038/nature22989.

24. Berry, J., Weber, S.C., Vaidya, N., Haataja, M., and Brangwynne, C.P. (2015). RNA
transcription modulates phase transition-driven nuclear body assembly. Proc. Natl.
Acad. Sci. U.S.A. *112*. 10.1073/pnas.1509317112.

25. Brangwynne, C.P., Eckmann, C.R., Courson, D.S., Rybarska, A., Hoege, C.,
Gharakhani, J., Jülicher, F., and Hyman, A.A. (2009). Germline P Granules Are
Liquid Droplets That Localize by Controlled Dissolution/Condensation. Science *324*,
1729–1732. 10.1126/science.1172046.

26. Brangwynne, C.P., Mitchison, T.J., and Hyman, A.A. (2011). Active liquid-like
behavior of nucleoli determines their size and shape in *Xenopus laevis* oocytes.
Proc. Natl. Acad. Sci. U.S.A. *108*, 4334–4339. 10.1073/pnas.1017150108.

27. Elbaum-Garfinkle, S., Kim, Y., Szczepaniak, K., Chen, C.C.-H., Eckmann, C.R.,
Myong, S., and Brangwynne, C.P. (2015). The disordered P granule protein LAF-1
drives phase separation into droplets with tunable viscosity and dynamics. Proc.
Natl. Acad. Sci. U.S.A. *112*, 7189–7194. 10.1073/pnas.1504822112.

28. Molliex, A., Temirov, J., Lee, J., Coughlin, M., Kanagaraj, A.P., Kim, H.J., Mittag, T.,
and Taylor, J.P. (2015). Phase separation by low complexity domains promotes
stress granule assembly and drives pathological fibrillization. Cell *163*, 123–133.
10.1016/j.cell.2015.09.015.

29. Zhang, H., Elbaum-Garfinkle, S., Langdon, E.M., Taylor, N., Occhipinti, P., Bridges,
A.A., Brangwynne, C.P., and Gladfelter, A.S. (2015). RNA Controls PolyQ Protein
Phase Transitions. Mol Cell *60*, 220–230. 10.1016/j.molcel.2015.09.017.

30. Larson, A.G., and Narlikar, G.J. (2018). The Role of Phase Separation in
Heterochromatin Formation, Function, and Regulation. Biochemistry *57*, 2540–2548.
10.1021/acs.biochem.8b00401.

31. Rosowski, K.A., Sai, T., Vidal-Henriquez, E., Zwicker, D., Style, R.W., and Dufresne,
E.R. (2020). Elastic ripening and inhibition of liquid–liquid phase separation. Nat.
Phys. *16*, 422–425. 10.1038/s41567-019-0767-2.

32. Bergeron-Sandoval, L.-P., and Michnick, S.W. (2018). Mechanics, Structure and
Function of Biopolymer Condensates. J Mol Biol *430*, 4754–4761.
10.1016/j.jmb.2018.06.023.

33. Busch, D.J., Houser, J.R., Hayden, C.C., Sherman, M.B., Lafer, E.M., and
Stachowiak, J.C. (2015). Intrinsically disordered proteins drive membrane curvature.
Nat Commun *6*, 7875. 10.1038/ncomms8875.

34. Bergeron-Sandoval, L.-P., Kumar, S., Heris, H.K., Chang, C.L.A., Cornell, C.E., Keller, S.L., François, P., Hendricks, A.G., Ehrlicher, A.J., Pappu, R.V., et al. (2021). Endocytic proteins with prion-like domains form viscoelastic condensates that enable membrane remodeling. Proc Natl Acad Sci U S A *118*, e2113789118. 10.1073/pnas.2113789118.

35. Yuan, F., Alimohamadi, H., Bakka, B., Trementozzi, A.N., Day, K.J., Fawzi, N.L., Rangamani, P., and Stachowiak, J.C. (2021). Membrane bending by protein phase separation. Proc Natl Acad Sci U S A *118*, e2017435118. 10.1073/pnas.2017435118.

36. Shin, Y., Chang, Y.-C., Lee, D.S.W., Berry, J., Sanders, D.W., Ronceray, P., Wingreen, N.S., Haataja, M., and Brangwynne, C.P. (2018). Liquid Nuclear Condensates Mechanically Sense and Restructure the Genome. Cell *175*, 1481-1491.e13. 10.1016/j.cell.2018.10.057.

37. Ekwall, K., Javerzat, J.-P., Lorentz, A., Schmidt, H., Cranston, G., and Allshire, R. (1995). The Chromodomain Protein Swi6: A Key Component at Fission Yeast Centromeres. Science *269*, 1429–1431. 10.1126/science.7660126.

38. Lorentz, A., Ostermann, K., Fleck, O., and Schmidt, H. (1994). Switching gene swi6, involved in repression of silent mating-type loci in fission yeast, encodes a homologue of chromatin-associated proteins from Drosophila and mammals. Gene *143*, 139–143. 10.1016/0378-1119(94)90619-X.

39. Allshire, R.C., Nimmo, E.R., Ekwall, K., Javerzat, J.P., and Cranston, G. (1995). Mutations derepressing silent centromeric domains in fission yeast disrupt chromosome segregation. Genes Dev. *9*, 218–233. 10.1101/gad.9.2.218.

40. Canzio, D., Chang, E.Y., Shankar, S., Kuchenbecker, K.M., Simon, M.D., Madhani, H.D., Narlikar, G.J., and Al-Sady, B. (2011). Chromodomain-Mediated Oligomerization of HP1 Suggests a Nucleosome-Bridging Mechanism for Heterochromatin Assembly. Molecular Cell *41*, 67–81. 10.1016/j.molcel.2010.12.016.

41. Nakayama Ji, null, Allshire, R.C., Klar, A.J., and Grewal, S.I. (2001). A role for DNA polymerase alpha in epigenetic control of transcriptional silencing in fission yeast. EMBO J *20*, 2857–2866. 10.1093/emboj/20.11.2857.

42. Kanoh, J., Sadaie, M., Urano, T., and Ishikawa, F. (2005). Telomere Binding Protein Taz1 Establishes Swi6 Heterochromatin Independently of RNAi at Telomeres. Current Biology *15*, 1808–1819. 10.1016/j.cub.2005.09.041.

43. Alberti, S., Gladfelter, A., and Mittag, T. (2019). Considerations and Challenges in Studying Liquid-Liquid Phase Separation and Biomolecular Condensates. Cell *176*, 419–434. 10.1016/j.cell.2018.12.035.

44. Biswas, S., Chen, Z., Karslake, J.D., Farhat, A., Ames, A., Raiymbek, G.,
Freddolino, P.L., Biteen, J.S., and Ragunathan, K. (2022). HP1 oligomerization
compensates for low-affinity H3K9me recognition and provides a tunable
mechanism for heterochromatin-specific localization. Sci Adv *8*, eabk0793.
10.1126/sciadv.abk0793.

45. Zhang, H., Zhao, R., Tones, J., Liu, M., Dilley, R.L., Chenoweth, D.M., Greenberg,
R.A., and Lampson, M.A. (2020). Nuclear body phase separation drives telomere
clustering in ALT cancer cells. MBoC *31*, 2048–2056. 10.1091/mbc.E19-10-0589.

46. Min, J., Wright, W.E., and Shay, J.W. (2019). Clustered telomeres in phase-
separated nuclear condensates engage mitotic DNA synthesis through BLM and
RAD52. Genes Dev. *33*, 814–827. 10.1101/gad.324905.119.

47. Gasser, S.M., Hediger, F., Taddei, A., Neumann, F.R., and Gartenberg, M.R.
(2004). The Function of Telomere Clustering in Yeast: The Circe Effect. Cold Spring
Harbor Symposia on Quantitative Biology *69*, 327–338. 10.1101/sqb.2004.69.327.

48. Cooper, J.P., Watanabe, Y., and Nurse, P. (1998). Fission yeast Taz1 protein is
required for meiotic telomere clustering and recombination. Nature *392*, 828–831.
10.1038/33947.

49. Sadaie, M., Iida, T., Urano, T., and Nakayama, J. (2004). A chromodomain protein,
Chp1, is required for the establishment of heterochromatin in fission yeast. EMBO J
23, 3825–3835. 10.1038/sj.emboj.7600401.

50. Sadaie, M., Kawaguchi, R., Ohtani, Y., Arisaka, F., Tanaka, K., Shirahige, K., and
Nakayama, J. (2008). Balance between Distinct HP1 Family Proteins Controls
Heterochromatin Assembly in Fission Yeast. Mol Cell Biol *28*, 6973–6988.
10.1128/MCB.00791-08.

51. Haldar, S., Saini, A., Nanda, J.S., Saini, S., and Singh, J. (2011). Role of Swi6/HP1
Self-association-mediated Recruitment of Clr4/Suv39 in Establishment and
Maintenance of Heterochromatin in Fission Yeast. Journal of Biological Chemistry
286, 9308–9320. 10.1074/jbc.M110.143198.

52. Wang, J., Reddy, B.D., and Jia, S. (2015). Rapid epigenetic adaptation to
uncontrolled heterochromatin spreading. eLife *4*, e06179. 10.7554/eLife.06179.

53. Yamagishi, Y., Sakuno, T., Shimura, M., and Watanabe, Y. (2008). Heterochromatin
links to centromeric protection by recruiting shugoshin. Nature *455*, 251–255.
10.1038/nature07217.

54. Shimada, A., Dohke, K., Sadaie, M., Shinmyozu, K., Nakayama, J.-I., Urano, T., and
Murakami, Y. (2009). Phosphorylation of Swi6/HP1 regulates transcriptional gene
silencing at heterochromatin. Genes Dev. *23*, 18–23. 10.1101/gad.1708009.

55. Canzio, D., Liao, M., Naber, N., Pate, E., Larson, A., Wu, S., Marina, D.B., Garcia,
J.F., Madhani, H.D., Cooke, R., et al. (2013). A conformational switch in HP1
releases auto-inhibition to drive heterochromatin assembly. Nature *496*, 377–381.
10.1038/nature12032.

56. Zhao, Y., Schreiner, S.M., Koo, P.K., Colombi, P., King, M.C., and Mochrie, S.G.J.
(2016). Improved Determination of Subnuclear Position Enabled by Three-
Dimensional Membrane Reconstruction. Biophysical Journal *111*, 19–24.
10.1016/j.bpj.2016.05.036.

57. Müller, K.P., Erdel, F., Caudron-Herger, M., Marth, C., Fodor, B.D., Richter, M.,
Scaranaro, M., Beaudouin, J., Wachsmuth, M., and Rippe, K. (2009). Multiscale
Analysis of Dynamics and Interactions of Heterochromatin Protein 1 by
Fluorescence Fluctuation Microscopy. Biophys J *97*, 2876–2885.
10.1016/j.bpj.2009.08.057.

58. Karslake, J.D., Donarski, E.D., Shelby, S.A., Demey, L.M., DiRita, V.J., Veatch, S.L.,
and Biteen, J.S. (2021). SMAUG: Analyzing single-molecule tracks with
nonparametric Bayesian statistics. Methods *193*, 16–26.
10.1016/j.ymeth.2020.03.008.

59. Banani, S.F., Rice, A.M., Peeples, W.B., Lin, Y., Jain, S., Parker, R., and Rosen,
M.K. (2016). Compositional Control of Phase-Separated Cellular Bodies. Cell *166*,
651–663. 10.1016/j.cell.2016.06.010.

60. Nonaka, N., Kitajima, T., Yokobayashi, S., Xiao, G., Yamamoto, M., Grewal, S.I.S.,
and Watanabe, Y. (2002). Recruitment of cohesin to heterochromatic regions by
Swi6/HP1 in fission yeast. Nat Cell Biol *4*, 89–93. 10.1038/ncb739.

61. Maru, Y., Afar, D.E., Witte, O.N., and Shibuya, M. (1996). The dimerization property
of glutathione S-transferase partially reactivates Bcr-Abl lacking the oligomerization
domain. J Biol Chem *271*, 15353–15357. 10.1074/jbc.271.26.15353.

62. Hiragami-Hamada, K., Soeroes, S., Nikolov, M., Wilkins, B., Kreuz, S., Chen, C., De
La Rosa-Velázquez, I.A., Zenn, H.M., Kost, N., Pohl, W., et al. (2016). Dynamic and
flexible H3K9me3 bridging via HP1β dimerization establishes a plastic state of
condensed chromatin. Nat Commun *7*, 11310. 10.1038/ncomms11310.

63. Weaire, D.L., and Hutzler, S. (1999). The Physics of Foams (Clarendon Press).

64. Damodaran, K., Venkatachalapathy, S., Alisafaei, F., Radhakrishnan, A.V., Sharma
Jokhun, D., Shenoy, V.B., and Shivashankar, G.V. (2018). Compressive force
induces reversible chromatin condensation and cell geometry–dependent
transcriptional response. MBoC *29*, 3039–3051. 10.1091/mbc.E18-04-0256.

65. Le, H.Q., Ghatak, S., Yeung, C.-Y.C., Tellkamp, F., Günschmann, C., Dieterich, C.,
Yeroslaviz, A., Habermann, B., Pombo, A., Niessen, C.M., et al. (2016). Mechanical

regulation of transcription controls Polycomb-mediated gene silencing during
lineage commitment. Nat Cell Biol *18*, 864–875. 10.1038/ncb3387.

66. Maki, K., Nava, M.M., Villeneuve, C., Chang, M., Furukawa, K.S., Ushida, T., and
Wickström, S.A. (2021). Hydrostatic pressure prevents chondrocyte differentiation
through heterochromatin remodeling. J Cell Sci *134*, jcs247643.
10.1242/jcs.247643.

67. Sanulli, S., Trnka, M.J., Dharmarajan, V., Tibble, R.W., Pascal, B.D., Burlingame,
A.L., Griffin, P.R., Gross, J.D., and Narlikar, G.J. (2019). HP1 reshapes nucleosome
core to promote phase separation of heterochromatin. Nature *575*, 390–394.
10.1038/s41586-019-1669-2.

68. Erdel, F., Rademacher, A., Vlijm, R., Tünnermann, J., Frank, L., Weinmann, R.,
Schweigert, E., Yserentant, K., Hummert, J., Bauer, C., et al. (2020). Mouse
Heterochromatin Adopts Digital Compaction States without Showing Hallmarks of
HP1-Driven Liquid-Liquid Phase Separation. Molecular Cell *78*, 236-249.e7.
10.1016/j.molcel.2020.02.005.

69. Yamanaka, S., Mehta, S., Reyes-Turcu, F.E., Zhuang, F., Fuchs, R.T., Rong, Y.,
Robb, G.B., and Grewal, S.I.S. (2013). RNAi triggered by specialized machinery
silences developmental genes and retrotransposons. Nature *493*, 557–560.
10.1038/nature11716.

70. Zofall, M., and Grewal, S.I.S. (2006). Swi6/HP1 Recruits a JmjC Domain Protein to
Facilitate Transcription of Heterochromatic Repeats. Molecular Cell *22*, 681–692.
10.1016/j.molcel.2006.05.010.

71. Style, R.W., Sai, T., Fanelli, N., Ijavi, M., Smith-Mannschott, K., Xu, Q., Wilen, L.A.,
and Dufresne, E.R. (2018). Liquid-Liquid Phase Separation in an Elastic Network.
Phys. Rev. X *8*, 011028. 10.1103/PhysRevX.8.011028.

72. Hu, Y., Bennett, H.W., Liu, N., Moravec, M., Williams, J.F., Azzalin, C.M., and King,
M.C. (2019). RNA-DNA Hybrids Support Recombination-Based Telomere
Maintenance in Fission Yeast. Genetics *213*, 431–447.
10.1534/genetics.119.302606.

73. Moreno, S., Klar, A., and Nurse, P. (1991). Molecular genetic analysis of fission
yeast Schizosaccharomyces pombe. Methods Enzymol *194*, 795–823.
10.1016/0076-6879(91)94059-l.

74. Bähler, J., Wu, J.-Q., Longtine, M.S., Shah, N.G., Mckenzie III, A., Steever, A.B.,
Wach, A., Philippsen, P., and Pringle, J.R. (1998). Heterologous modules for
efficient and versatile PCR-based gene targeting inSchizosaccharomyces pombe.
Yeast *14*, 943–951. 10.1002/(SICI)1097-0061(199807)14:10<943::AID-
YEA292>3.0.CO;2-Y.

75. Hentges, P., Van Driessche, B., Tafforeau, L., Vandenhaute, J., and Carr, A.M.
(2005). Three novel antibiotic marker cassettes for gene disruption and marker
switching inSchizosaccharomyces pombe. Yeast *22*, 1013–1019. 10.1002/yea.1291.

76. Snaith, H.A., Samejima, I., and Sawin, K.E. (2005). Multistep and multimode cortical
anchoring of tea1p at cell tips in fission yeast. EMBO J *24*, 3690–3699.
10.1038/sj.emboj.7600838.

77. Torres-Garcia, S., Di Pompeo, L., Eivers, L., Gaborieau, B., White, S.A., Pidoux,
A.L., Kanigowska, P., Yaseen, I., Cai, Y., and Allshire, R.C. (2020). SpEDIT: A fast
and efficient CRISPR/Cas9 method for fission yeast. Wellcome Open Res *5*, 274.
10.12688/wellcomeopenres.16405.1.

78. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-
source platform for biological-image analysis. Nat Methods *9*, 676–682.
10.1038/nmeth.2019.

79. Eilers, P.H.C., and Goeman, J.J. (2004). Enhancing scatterplots with smoothed
densities. Bioinformatics *20*, 623–628. 10.1093/bioinformatics/btg454.

80. Munkres, J. (1957). Algorithms for the Assignment and Transportation Problems.
Journal of the Society for Industrial and Applied Mathematics *5*, 32–38.
10.1137/0105003.

81. Rowland, D.J., and Biteen, J.S. (2017). Measuring molecular motions inside single
cells with improved analysis of single-particle trajectories. Chemical Physics Letters
674, 173–178. 10.1016/j.cplett.2017.02.052.

82. Mack, A.H., Schlingman, D.J., Regan, L., and Mochrie, S.G.J. (2012). Practical axial
optical trapping. Review of Scientific Instruments *83*, 103106. 10.1063/1.4757862.

1547
1548