Analysis of a photosynthetic cyanobacterium rich in

internal membrane systems via gradient profiling by

sequencing (Grad-seq)

Matthias Riediger[1], Philipp Spät[2], Raphael Bilger[1], Boris Macek[2], Wolfgang R. Hess[1,*]

[1]Genetics and Experimental Bioinformatics, Faculty of Biology, University of Freiburg,

Schänzlestr. 1, 79104 Freiburg, Germany;

[2]Department of Quantitative Proteomics, Interfaculty Institute for Cell Biology,

University of Tübingen, Auf der Morgenstelle 15, 72076 Tübingen, Germany

Short title: Grad-seq in *Synechocystis* sp. PCC 6803

Category: Large-Scale Biology Article

*corresponding author: Wolfgang R. Hess: wolfgang.hess@biologie.uni-freiburg.de

Abstract

Regulatory sRNAs in photosynthetic cyanobacteria have been reported, but the lack of plausible RNA chaperones involved in this regulation appears enigmatic. Here, we analyzed the full ensemble of cellular RNAs and cofractionating proteins using gradient profiling by sequencing (Grad-seq) in *Synechocystis* 6803. Complexes with overlapping subunits such as the CpcG1-type versus the CpcL-type phycobilisomes or the PsaK1 versus PsaK2 photosystem I pre(complexes) could be distinguished supporting a high quality of the approach. Clustering of the in-gradient distribution profiles followed by several additional criteria yielded a short list of potential RNA chaperones that include a YlxR homolog and a cyanobacterial homolog of the KhpA/B complex. The data suggest previously undetected complexes between accessory proteins and CRISPR-Cas systems, such as a Csx1-Csm6 ribonucleolytic defense complex. Moreover, the exclusive association of either RpoZ or 6S RNA with the core RNA polymerase complex and the existence of a reservoir of inactive sigma-antisigma complexes is suggested. The *Synechocystis* Grad-seq resource is available online at https://trex90.shinyapps.io/GradSeqExplorer_SynPCC6803/, providing a comprehensive resource for the functional assignment of RNA-protein complexes and multisubunit protein complexes in a photosynthetic organism.

Keywords: cyanobacteria / post-transcriptional regulation / ribonucleoprotein complexes / RNA-binding proteins / small regulatory RNAs

Introduction

Non-coding (nc)RNAs constitute a major component of the transcriptional output in all classes of organisms (Cech and Steitz, 2014; Morris and Mattick, 2014). In photosynthetic cyanobacteria as well as in other bacteria, complex regulatory networks have been identified that include small regulatory ncRNAs (sRNAs) as major players in the posttranscriptional control of gene expression (Wagner and Romby, 2015; Kopf and Hess, 2015). While some sRNAs as well as cis-transcribed antisense RNAs (asRNAs) may function independent of proteins, the vast majority of ncRNAs require interactions with specific RNA binding proteins (RBPs).

Cyanobacteria form a single phylum of species with very different morphologies, with genome sizes varying by approximately one order of magnitude and highly diverse lifestyles. Nevertheless, cyanobacteria are monophyletic and the only bacteria that perform oxygenic photosynthesis. They are the evolutionary ancestors of chloroplasts, are of paramount ecological importance as primary producers and of high biotechnological interest (Hagemann and Hess, 2018; Vijay et al., 2019). In the unicellular model cyanobacterium *Synechocystis* sp. PCC 6803 (from here *Synechocystis*), three CRISPR systems (Scholz et al., 2013), a 6S RNA homolog (Heilmann et al., 2017) and hundreds of putative sRNAs have been identified (Kopf and Hess, 2015; Kopf et al., 2014). Some of these sRNAs are important posttranscriptional regulators. The sRNA PsrR1 caps the expression of several photosynthesis proteins and pigment biosynthesis proteins during the acclimation of the photosynthetic machinery to high light conditions (Georg et al., 2014). Likewise, the sRNA IsaR1 contributes to the adjustment of iron-sulfur cluster biosynthesis, photosynthetic electron transport and photosystem (PS) I gene expression in the acclimation response to low iron (Georg et al., 2017). Moreover, two different types of ncRNAs, the sRNA NsiR4 and the glutamine type I riboswitch, were found to control cyanobacterial nitrogen assimilation in ways that differ considerably from the archetypical model developed for *E. coli* (Klähn et al., 2015, 2018). Despite the observed abundance of riboregulatory processes, no RBPs functionally similar to ProQ, CsrA and Hfq could be identified in cyanobacteria. While there are no candidates for the former two, a structural homolog of Hfq is present in several cyanobacteria, including *Synechocystis* (Dienst et al., 2008; Bøggild et al., 2009). However, compared to the homologs in proteobacteria, cyanobacterial Hfq is

truncated, does not bind RNA and may function through protein:protein interactions
(Schuergers et al., 2014).

A recently developed method uses gradient profiling by sequencing (Grad-seq) to
directly detect groups of RNAs and comigrating proteins. In this approach, whole-cell
lysates are fractionated on a sucrose density gradient by ultracentrifugation. The
fractions are subject to mass spectrometry-based proteome measurements and
RNA-sequencing analyses, hence guiding the potential discovery of new globally
acting RBPs and polypeptides acting together in large multiprotein complexes
(Smirnov et al., 2016). This approach has been prolific in the discovery of ProQ as a
previously overlooked major small RNA-binding protein in the bacterial pathogen
Salmonella enterica (Smirnov et al., 2016; Westermann et al., 2019) and other
enteric bacteria (Melamed et al., 2020) and identified the unexpected involvement of
exoribonuclease activity in the stabilization and activation of sRNAs in the Gram-
positive pathogen *Streptococcus pneumonia* (Hör et al., 2020).

While there are reports on the composition of protein complexes (Xu et al., 2020) and
the association of these complexes with the thylakoid or cell membrane systems in
the model *Synechocystis* (Baers et al., 2019), no information on RNA/protein
complexes is currently available for any cyanobacterium. Here, we report the results
of Grad-seq analysis applied to *Synechocystis*, identifying the sedimentation profiles
of thousands of transcripts and proteins simultaneously.

We applied a hierarchical clustering approach and provide a database that supports
further analyses of this dataset based on multiple filter criteria. In contrast to previous
Grad-Seq studies, this strategy allows the delineation of RBPs or other proteins of
interest directly. The results provide the first RNA/protein complexome resource for a
photosynthetic cyanobacterium.

Results

Grad-seq analysis separates the full proteome and transcriptome of a photosynthetic bacterium

Triplicate cultures of *Synechocystis* were cultivated at moderate light intensities in BG11 medium. Whole cell lysates were prepared in the presence of beta-dodecylmaltoside, a mild detergent to allow the inclusion of membrane proteins, and fractionated by sucrose gradient ultracentrifugation as described (Smirnov et al., 2016). The gradients showed the colorful separation of the different pigment-containing complexes (**Figure 1A)**. After separation, the eluted fractions from each gradient were analyzed via mass spectrometry (MS)-based proteomics and RNA sequencing for the identification of their protein and RNA composition. The overall in-gradient distribution of RNA profiles (**Supplementary Figure S1A**) and protein profiles (**Supplementary Figure S1B**) showed a strong correlation (median RNA-seq r = 0.76, median MS r = 0.85) between the replicates and confirmed good reproducibility. The fractionated proteins and RNAs were also visualized using conventional denaturing polyacrylamide gels (**Figure 1A** and **B**).

Following the identification of proteins and RNAs by MS and RNA sequencing, we determined the sedimentation profile of 2,394 proteins and 4,251 different transcripts detected in all three replicates (**Figure 1C, Supplementary Table S1**). These transcripts consisted of 2,968 mRNAs, 530 separate 5'UTRs, 359 3'UTRs, 140 ncRNAs, 139 asRNAs, 65 transposase-associated RNAs, 42 tRNAs, 6 rRNAs and 3 types of crRNAs. A total of 3,559 annotated protein-coding genes were previously defined in this strain of *Synechocystis* (Trautmann et al., 2012); hence, we detected 67.3% of all annotated proteins. When including 122 more recently identified additional genes, we detected 65% of the 3,681 total proteins. The 4,251 different detected transcripts corresponded to 2,544 previously defined transcriptional units (TUs), out of a total of 4,091 (Kopf et al., 2014). Hence, 62.2% of all TUs were detected. In our previous definition of TUs, we assigned multiple genes and their UTRs to one and the same TU if the transcriptomic coverage did not indicate interrupting terminators or internal transcriptional start sites. However, aiming at maximum sensitivity, we kept all genes and UTRs separate here. The percentages of 67.3% detected proteins and of 62.2% detected TUs are quite exhaustive because the samples were from a single culture condition, while certain genes are expressed

143 only under specific conditions (Kopf et al., 2014). In our database available at

144 https://trex90.shinyapps.io/GradSeqExplorer_SynPCC6803/, we provide the

145 opportunity to scan for all detected proteins and RNAs of this dataset.

Hierarchical cluster analysis splits the Grad-seq dataset into protein- and RNA-
dominated main branches

Based on the distribution along the gradient, proteins and transcripts were assigned

to 17 different clusters using the dynamic tree cut algorithm from the WGCNA R

package (Langfelder and Horvath, 2008, 2012) (**Figure 2A**, **Supplementary Figure**

S2A). A striking result from this approach was the division of the entire dataset into a

protein-dominated branch whose members mainly sedimented in the lower molecular

fractions and an RNA-dominated branch whose members belonged predominantly to

the higher molecular fractions (**Supplementary Figure S2B** and **C**). The protein-

dominated branch, represented by clusters 1 to 9, consisted of soluble proteins, small

and medium protein complexes and tRNAs in lower molecular fractions 2 to 6 [~10 -

20% sucrose]. Approximately 80% of all detected proteins were classified into this

branch. The RNA-dominated branch, represented by clusters 10 to 17, included most

transcripts and large protein complexes, predominantly in higher molecular fractions

6 to 18 [~20 - 40% sucrose], and comprised ~90% of all detected transcripts (**Figure**

2B). As expected, the sedimentation properties of different proteins varied largely

and relied on the molecular weight or on the size of the associated protein complex.

According to the calibration by known complexes, proteins peaking in fractions above

5 were associated with larger complexes of more than 250 kDa in size

(**Supplementary Figure S3A**). In contrast, the majority of RNAs exhibited a largely

similar sedimentation pattern, accumulating mainly in clusters 10, 13 and 14. Notable

exceptions were RNase P RNA, tmRNA, 6S RNA as well as 5S, 16S and 23S rRNA,

which sedimented towards higher buoyancies. These results are in accordance with

previous observations (Smirnov et al., 2016) that RNA sedimentation profiles are

largely independent of the transcript length but rely more on their corresponding

binding protein[s] (**Supplementary Figure S3B**).

The proteins of our dataset were categorized into functional groups (**Figure 2C**),

defined by KEGG (Kanehisa, 2019; Kanehisa et al., 2019; Kanehisa and Goto,

2000). We then compared every single protein with the presence of likely orthologs in
selected other organisms (**Supplementary Table S2**). We chose a set of 57
cyanobacterial genomes to judge the proteins conserved in cyanobacteria only, in the
E. coli and *S. enterica* genomes combined as reference for bacteria with
characterized sRNA chaperones and in the *A. thaliana* genome to detect proteins
more widely conserved in the green lineage.

This led to classifying the proteins into five categories: "*Synechocystis* only",
"cyanobacteria only", "bacteria", "cyanobacteria and *Arabidopsis*" and "all". The
dominance of proteins of unknown function in the "*Synechocystis* only" category and
their proportion of more than 50% in the "cyanobacteria only" category illustrate that
these organisms are still largely unexplored. Only 20% of the *Synechocystis*
proteome was assigned to clusters of the RNA-dominated branch. These clusters
included proteins that are part of large, well-known complexes, such as the majority
of PS subunits, RubisCO, RNAP, most ribosomal proteins and a small number of
largely undescribed proteins, depicting promising candidates for interaction with such
complexes or potential RBPs, especially if not conserved in *E. coli*, *S. enterica* and *A.*
thaliana (**Figure 2C**).

Many transcripts show bimodal sedimentation profiles

Most transcripts were assigned to one of the three clusters 10, 13 or 14. The
transcripts of clusters 13 and 14 had largely similar sedimentation profiles and were
analyzed together, while the sedimentation profiles of cluster 10 transcripts clearly
differed. The transcripts of cluster 10 mainly occurred in fractions 5 to 7 but with a 2^{nd}
occurrence above fraction 8, often peaking around fraction 9. Therefore, the
transcripts of cluster 10 were grouped according to their main peaks, yielding three
subgroups (**Figure 3A**). In these subgroups, the main peaks of mRNAs, their
associated asRNAs, and, as far as we know, their respective regulatory ncRNAs
largely overlapped. Hence, these fractions likely contained mRNA:asRNA and
mRNA:sRNA hybrids. However, the secondary peaks for mRNA occurrences did not
overlap with asRNA or ncRNA peaks. Hence, the respective mRNAs cofractionated
with their putative RNA regulators in the primary peak fractions and fractionated
separately in the secondary peak fractions. The transcripts of clusters 13 and 14

were grouped accordingly. Transposase mRNAs are a major constituent of these clusters. Notably, only two annotated ncRNAs, Ncr0560 and Ncr1310, are included in cluster 14, exhibiting a single peak in fraction 9. Ncr0560 originates from a transcriptional start site internal to gene *slr2062* encoding an IS200/IS605 TnpB protein and therefore is closely connected to transposases. However, most transcripts of clusters 13 and 14 occurred mainly in a narrow range of fractions 9 to 10, with secondary peaks below fraction 8, often in fraction 4 (**Figure 3B**). Therefore, the vast majority of *Synechocystis* transcripts seem to occur in at least two states, either bound to a large complex or as unbound RNA, probably protected from degradation by one or multiple RNA chaperones.

Grad-seq analysis allows the detection of ribonucleoprotein complexes

Fifty-three ribosomal or ribosome-associated proteins and the ribosomal RNAs (rRNAs) were present in the fractions containing the complexes with the highest sedimentation coefficients. A preference for the entire set of ribosomal components was restricted to fraction 18 (**Figure 4A**). Notable exceptions were Rpl11/Sll1743, Rpl12/sll1746, Rps2/sll1260, Rps10/sll1101 and Rpl16/sll1805, pointing to their association with other complexes or transcripts. Moreover, Ycf65/slr0923, previously reported as a plastid-specific ribosomal protein in some eukaryotic algae (Turmel et al., 2009) and higher plants (Yamaguchi and Subramanian, 2003), was not found to be associated with other r-proteins, nor were the Rps1a/slr1356 and Rps1b/slr1984 proteins (**Figure 4A**). Rps1a and Rps1b rather may play a role in the Shine-Dalgarno-independent initiation of translation in cyanobacteria (Nakagawa et al., 2010; Mutsuda and Sugiura, 2006).

CRISPR-Cas systems are particularly interesting macromolecular complexes that are known to consist of both protein and RNA components. CRISPR-Cas systems confer a form of acquired immunity against invading nucleic acids in many different bacteria and archaea and can be classified into greatly varying subtypes that belong to two major classes (Brouns et al., 2008; Makarova et al., 2015). Despite the high level of divergence among different systems, CRISPR-Cas systems usually consist of repeat-spacer arrays and the associated protein-coding *cas* genes. *Synechocystis* encodes three separate and complete CRISPR-Cas systems that are highly expressed and

active in interference assays (Scholz et al., 2013; Reimann et al., 2017; Kieper et al., 2018; Behler et al., 2018; Scholz et al., 2019). According to the presence of signature genes, these three systems were classified as subtype I-D, subtype III-D and subtype III-Bv systems (Scholz et al., 2013; Behler et al., 2018). The gradient profiles of the crRNAs of the three CRISPR systems of *Synechocystis* varied vastly from one another. While CRISPR1 RNAs (cr1RNAs) were mainly detected in fraction 9 (cluster 14), CRISPR3 RNAs (cr3RNAs) peaked in fraction 10 (**Figure 4B**). However, the detected CRISPR2-associated proteins colocalized together with the cr2RNAs in low molecular fractions 4 to 6 at the same sedimentation level as tRNAs. The detected CRISPR1-associated proteins were scattered throughout the gradient, with only the Cas7/Csc2 protein Slr7012 cofractionating with cr1RNAs in higher molecular fractions 9 to 11. This tight association with cr1RNAs is consistent with the known function of this protein in binding the variable crRNA spacer sequence (Hrle et al., 2014). All structural proteins associated with the CRISPR3 system (proteins Cmr1, Cmr5, Cmr4, Cmr3 and Cmr2/Cas10 encoded by the genes *sll7085* to *sll7087, sll7089* and *sll7090*) fractionated together and with the CRISPR3-associated crRNAs, with a peak in fraction 10. This co-fractionation indicates that the interference complex remained largely intact during the preparation. However, those proteins exhibited a second peak in the lower fractions without associated cr3RNA or the Cas10 protein Sll7090, hence constituting a possibly nonfunctional subcomplex (**Figure 4B**). Interestingly, Slr7080, a Csx3 accessory protein, cofractionated with these two CRISPR3 complexes. Cas accessory genes encode diverse groups of proteins that cooccur only sporadically with CRISPR-Cas systems and are not essential for their functionality (Shmakov et al., 2018; Shah et al., 2019). Slr7080 has been characterized as a Csx3-AAA protein (Shah et al., 2019), a distant member of the Cas Rossmann fold (CARF) protein family (Makarova et al., 2014). CARF proteins sense ligands such as cyclic oligoadenylate (cOA) signaling molecules that can be synthesized by the polymerase domain of Cas10 upon recognition of its target (Kazlauskiene et al., 2017; Niewoehner et al., 2017; Han et al., 2018). The *Archaeoglobus fulgidus* Csx3 protein AfCsx3 is a ring nuclease that degrades the CRISPR signaling molecule cOA4 (Athukoralage et al., 2020). However, Slr7080/Csx3 is substantially larger than AfCsx3 (310 compared to 110 amino acids) and possesses an additional AAA ATPase domain in the C-terminal region (Shah et al., 2019). Our data suggest that Slr7080/Csx3 is expressed relatively abundantly

and likely is physically linked to both the cr3RNA-free subcomplex and the complete
functional interference complex. Compelling evidence for an even more intriguing
complex of Cas accessory proteins was found for the CRISPR2 system in which the
four proteins Slr7061/Csx1, Sll7062/Csm6, Sll7070 and Slr7073 overlapped with the
fractionation of cr2RNAs (**Figure 4B**). Slr7061/Csx1 was bioinformatically classified
as a CARF-HEPN domain protein and Sll7062/Csm6 as a CARF-RelE domain-
containing protein (Shah et al., 2019). The protein most closely structurally related to
Slr7061/Csx1 is the *Sulfolobus islandicus* Csx1 protein, an RNase that becomes
allosterically activated upon cOA4 binding (Molina et al., 2019). Csm6 is a cOA-
dependent CRISPR ribonuclease as well (Garcia-Doval et al., 2020); hence, our data
suggest the presence of a ribonucleolytic defense complex associated with cr2RNAs.
The other two proteins, Sll7070 and Slr7073, are entirely uncharacterized, but the
localization of their genes that frame the CRISPR2 *cas1cas2* adaptation module
suggests a functionality closely associated with the CRISPR2 system. Prior to this
work, neither the expression of Slr7061/Csx1, Sll7062/Csm6, Sll7070 and Slr7073 at
the protein level nor their likely physical association with cr2RNAs was known.
Hence, these proteins are suggested as highly interesting candidates for future
experimental analyses.

The cyanobacterial RNAP is unique due to splitting of the β' subunit into an N-
terminal γ (RpoC1) and a C-terminal β' (RpoC2) subunit (Schneider and Hasekorn,
1988) and a >600 amino acid insertion into RpoC2 (Iyer et al., 2004). Moreover, the
interplay between the core subunits, multiple sigma factors (9 in *Synechocystis*) and
inhibitory 6S RNA is only partially understood (Heilmann et al., 2017). We observed
that all RNAP core subunits colocalized with the sigma factors SigB and SigC, as well
as 6S RNA, SigE and SigA in higher molecular fractions, indicating the parallel
presence of multiple forms of the RNAP holoenzyme. In contrast, the bulk of the
RNAP ω subunit (RpoZ), as well as SigF and SigG, together with three detected anti-
sigma factors was distributed only into the low molecular fractions, suggesting the
presence of a reservoir of inactive binary complexes (**Figure 4C**). Furthermore, we
noticed the mutually exclusive occurrence of either RpoZ or the 6S RNA together
with the RNAP core subunits (see F8 in **Figure 4C**). This observation may explain
the observation that cyanobacterial RpoZ facilitates the association of the primary
sigma factor SigA with the RNAP core stimulating transcription of highly expressed
genes (Gunnelius et al., 2014). In *E. coli*, 6S binds the β/β' subunits (Wassarman and

Storz, 2000). Hence, RpoZ may affect this binding by interacting with the γ and β'
subunits in *Synechocystis* (Gunnelius et al., 2014).

RNase P and the tmRNA showed similar sedimentation patterns as the RNAP
holoenzyme and 6S RNA, while the signal recognition particle mostly clustered
together with the bulk of mRNAs (**Figure 4C**). Most other factors, which were
annotated to different functionalities in RNA metabolism, seemed to not be
associated with a larger complex. Notable exceptions were the DEAD-box RNA
helicase CrhR, Sll1366 (a putative SNF2 helicase), polynucleotide phosphorylase
(PNPase), enolase, RNase J and RNase E. The latter enzyme is central for RNA
metabolism in bacteria because it is involved in numerous functions, such as the
posttranscriptional regulation of gene expression, maturation of transcripts or
initiation of mRNA degradation (Bandyra and Luisi, 2018). Examples in
Synechocystis include the posttranscriptional regulation of gene expression with the
sRNA PsrR1 as specificity factor (Georg et al., 2014), operon discoordination of the
rimO-crhR dicistron (Rosana et al., 2020) or the maturation of functionally active
crRNAs from the CRISPR3 system (Behler et al., 2018). Indeed, RNase E was found
in several fractions, peaking together with ribosomal proteins, cooccurring with the
bulk of mRNAs as well as cr3RNAs (**Figure 4C**). Cyanobacterial RNase E forms a
complex with PNPase through an oligoarginine repeat (Zhang et al., 2014b),
consistent with their cooccurrence in fraction 10 while their separate occurrences in
other fractions (**Figure 4C**) indicates that the PNPase-RNase E complex is likely only
one of several different combinations in which RNase E exists in the cell.

These results illustrate the sedimentation profiles of native ribonucleoprotein
complexes in a sucrose density gradient, supplying valuable information about the
compositional structure of such complexes in a cyanobacterial model organism. In
contrast to the Grad-seq analysis performed with *Salmonella* (Smirnov et al., 2016),
Hfq was in the density gradients not even close to any RNAs or known
ribonucleoprotein complexes (**Figure 4C**), confirming the different functionality of this
protein in cyanobacteria.

Sedimentation profiles of mature and intermediate protein complexes associated with the thylakoid membrane provide insights into the complexity of the photosynthetic lifestyle

Focusing on protein complexes of the cyanobacterial thylakoid membrane helps to understand the factors that maintain its photosynthetic machinery. Most components of these complexes were cosedimented, indicating the presence of largely intact PSI, PSII and cytochrome-b_6f complexes (**Figure 5A**), NADH-dehydrogenase (NDH-1) and cytochrome-c-oxidase complexes as well as F0F1ATP synthase (**Supplementary Figure S4A-C**). The detection of several small proteins, such as NdhP/sml0013 (Schwarz et al., 2013; Zhang et al., 2014a), NdhN, NdhH, NdhJ, and NdhO (He and Mi, 2016), cosedimenting with other NDH-1 proteins, supports the validity of the data. The soluble electron carriers plastocyanin (PetE), cytochrome c_6 (PetJ) and ferredoxins (PetF, Fdx) were mainly located between the lowest molecular fractions 2 and 3 and were not associated with any complex except for Fdx7 (**Supplementary Figure S4D**).

The occurrence of the PSII and PSI core complexes is in accordance with the measured absorption spectra for fully assembled complexes in fractions 6 and 7 (**Figure 5B**). The three membrane-extrinsic PSII subunits PsbO, PsbV and PsbU, located at the lumenal side of the thylakoid membrane, were disconnected and cosedimented in fraction 3 (**Figure 5A**). Several other PSII accessory proteins were identified in lower fractions as well, including Psb27 (**Figure 5A**) and most PSII assembly factors (reviewed by (Nickelsen and Rengstl, 2013)) (**Supplementary Figure S4E**). A PSII subcomplex lacking PSII core reaction center polypeptides but containing Psb27 was described as a PSII repair cycle chlorophyll-protein complex (Weisz et al., 2019); therefore, corresponding complexes in transition between disassembled and assembled PSII were likely present in fractions 4 to 5. This hypothesis is supported by the measured absorption spectra, which mainly showed the presence of carotenoids but no chlorophylls (**Figure 5B**). Detected PSII assembly factors cosedimenting with and potentially associated with PSII subcomplexes include Slr1471, PratA/Slr2048, CtpA/Slr0008 and Ycf48/Slr2034, which are involved in D1 integration, processing and stabilization in the early stage of reaction center assembly; Psb27/Slr1645 and Sll0606, which are involved in CP43 assembly and integration in the later stage of reaction center assembly and PSII repair cycle; and

Slr1768, Sll1390 and Sll1414, which are involved in D1 turnover as part of the PSII repair cycle (Nickelsen and Rengstl, 2013). Interestingly, few PSII assembly factors showed a more complex sedimentation pattern. Slr1768 exhibited a second and Slr2013 two more peaks in higher molecular fractions, while Sll0933, the homolog of the PSII assembly factor PAM68 in *Arabidopsis* (Rast et al., 2016; Rengstl et al., 2013), solely sedimented in higher molecular fractions, leaving their exact role in PSII assembly in *Synechocystis* open for discussion. Furthermore, our data illustrate the roles of PsaL and PsaK in PSI trimer formation. PsaL and PsaK are the last subunits to be incorporated into PSI in the trimer assembly process, with the incorporation of PsaK2 mainly involved in state transition and PsaK1 in the constitutive state (Dühring et al., 2007; Fujimori et al., 2005), while PsaL is required for PSI trimer formation (Chitnis and Chitnis, 1993). The intact PSI monomer could be easily detected, but the PSI trimer peak was only weakly detectable, since the majority of PSI seemed to occur monomerically, possibly caused by the preparation. However, the peak appeared more clearly when the proteomics data were normalized to equal protein amounts per fraction (**Supplementary Figure S4G** and **H**). Matching the findings of previous research, our data show the cosedimentation of PsaL and PsaK1 with the fully assembled PSI monomer as well as the trimer, while the majority of PsaK2 was only partially incorporated into the PSI monomer together with most assembly factors. Similar to previous observations of multiple peaks of some PSII assembly factors, Ycf4/Sll0226 exhibited a second and VIPP1/Sll0617 exhibited two more peaks in higher molecular fractions (**Supplementary Figure S4F**).

The sedimentation profiles of the phycobiliproteins pointed to another aspect of the complex dynamics of the photosynthetic apparatus. Two types of phycobilisomes (PBS) were previously described. The main phycobilisome (CpcG1-PBS type) is characterized by CpcG1 as a rod-core linker and the smaller CpcL-PBS type is characterized by CpcG2 as a rod core linker (Kondo et al., 2005; Liu et al., 2019). The distinction between those two types according to their protein composition illustrates the different sedimentation profiles of CpcG2-PBS and CpcG1-PBS within the pigmented sedimentation range (**Figure 5**). The association of CpcG2 with PsaK2 was suggested under high light conditions to function as a PSI antenna involved in state transitions (Kondo et al., 2007, 2009). As part of an NDH1L–PSI–CpcG2-PBS supercomplex, CpcG2 is involved in facilitating efficient cyclic electron transport (Gao et al., 2016). Interestingly, ferredoxin-NADP$^+$ reductase (FNR, PetH)

404 and PsaK1 colocalize with CpcG1-PBS. The association of FNR with phycobilisomes

405 has previously been shown (Kondo et al., 2005; Liu et al., 2019), while the interaction

406 of PsaK1 and CpcG1-PBS, analogous to the interaction of PsaK2 and CpcG2-PBS,

407 has not yet been shown.

Smaller complexes and protein-protein interactions in toxin-antitoxin systems

409
and regulation

In addition to the analysis of ribonucleoprotein complexes and membrane-bound

protein complexes, the dataset also provides interesting clues about smaller

complexes consisting of just some interacting proteins.

One class of such complexes with frequently just two proteins are the toxin–antitoxin

(TA) systems. TA systems consist of a stable toxic protein and its unstable cognate

antitoxin. In the case of Type II TA systems, both of these components are small

proteins (Schuster and Bertram, 2013). Previously, 69 Type II TA systems were

predicted for *Synechocystis* (Kopfmann et al., 2016). While the majority of these

proteins are plasmid-localized, consistent with their function in plasmid maintenance,

22 predicted TA systems are encoded in the main chromosome. These must have

functions distinct from postsegregational killing but are severely understudied. We

detected at least four such TA systems: Slr0770/Slr0771 and Ssl0258/Ssl0259 in

cluster 2; Ssl2138/Sll1092 in cluster 5 and Ssl2245/Sll1130 in cluster 8 (**Figure 6A**).

From these, the first two have not been studied so far, while the functionality of

Ssl2138/Sll1092 as a TA system was experimentally verified (Ning et al., 2013) and

for Ssl2245/Sll1130, some evidence exists that it is involved in a wider regulatory

context (Srikumar et al., 2017). Interestingly, Ssl2245/Sll1130 were assigned to

cluster 8, suggesting that they were part of a larger complex. Hence, our data not

only support the existence of these TA systems but also show their active expression

and support their supposed mode of action due to their co-fractionation.

Other smaller protein complexes have regulatory functions. Here, we chose proteins

interacting with components of the nitrogen regulatory system for illustration. The

GntR-like transcriptional regulator PlmA is in an inactive state when bound to P_{II}-PipX

(Labella et al., 2016) but is involved in phycobilisome degradation when activated

under nitrogen starvation (Sato et al., 2008). The formation of the P_{II}-PipX complex

also controls the activity of the transcriptional master regulator for nitrogen starvation,

NtcA (for review see (Labella et al., 2020a)). During high nitrogen abundance, P_{II}

forms a complex with either N-acetylglutamate kinase (NAGK, ArgB), which stabilizes

it and therefore enhances arginine synthesis. Furthermore, it binds approximately

85% of PipX to sequester it from interacting with NtcA and thus inhibits its regulatory

properties. The other 15% of PipX is bound by PlmA to keep it inactive. During low

nitrogen, P_{II} is sequestered by 2OG, and NtcA-dependent gene expression is

activated upon binding to PipX (Llacer et al., 2010; Forcada-Nadal et al., 2018). Our

data illustrate the cooccurrence of the P_{II}-PipX complex and all mentioned proteins

interacting with its single components (**Figure 6B**). Another protein interacting with P_{II}

during high nitrogen conditions is the membrane protein PamA, implicated in the

control of genes of nitrogen and sugar metabolism (Osanai et al., 2005). Notably, this

is the only one of the previously mentioned proteins that exhibited peaks outside the

range of the P_{II}-PipX complex, thus potentially belonging to a larger complex (**Figure**

6B). Interestingly, the single occurrence of PipX in fraction 9 (**Figure 6B**) might be

linked to a recently suggested new functionality as an interactor with the translation

machinery (Cantos et al., 2019).

Classification of sedimentation zones linked to various steps of protein

biosynthesis assisting in the identification of cyanobacterial RNA chaperones

Most transcripts accumulated in a narrow gradient range, equivalent to large protein

complexes of ~300 kDa in size, similar to the PS monomers. Intriguingly, this section

was largely free of known RNA interacting complexes, indicating the presence of one

or more RNA chaperones responsible for the characteristic sedimentation pattern.

We classified these transcripts of cluster 10 as group 1 RNAs. Other transcripts

assigned to clusters 13 and 14 were classified as group 2 RNAs (**Figure 7**).

Most RNA-interacting complexes involved in protein biosynthesis sedimented deeper,

accumulating in higher molecular fractions. These proteins included the RNAP

holoenzyme, 6S RNA, RNase P, tmRNA, several other proteins functionally linked to

RNA degradation and a subset of ribosomal proteins. A large set of transcripts,

including 16S rRNA, peaked in the same range, indicating the location of a large

transcription-translation complex still binding RNAs (**Figure 7**). Proteins that

sedimented in these fractions are another group of interesting candidates with potential involvement in RNA metabolism. Interestingly, although CRISPR1, CRISPR3 and most transposase mRNAs mainly cosedimented with those complexes, their involvement in specific regulatory aspects is not known.

Based on these findings, further analyses were performed to characterize the proteins cosedimenting with transcripts and to determine interesting candidates for future research (**Supplementary Figure S5**). We primarily focused on 19 % (451 / 2394) of all detected proteins, which were assigned to the RNA-dominated clusters in the higher molecular fractions (cluster ≥10), to which the essential complexes of protein biosynthesis also belonged. Specifically, 27 % (124 / 451) of those proteins that were annotated as unknown function were of high interest. Of those, a total of .41% (51 / 124) were conserved in at least 50 % of selected cyanobacterial genomes but not in *E. coli*, *S. enterica* or *A. thaliana* (**Supplementary Table S2**). Hence, these 51 proteins are conserved, largely undescribed and exhibit sedimentation profiles linking them to RNA metabolism. To aid the selection further, an SVM score was computed for the prediction of RNA-binding proteins from their amino acid sequence using the RNApred web server (Kumar et al., 2011). We performed this prediction for all proteins and included it in the database associated with this paper. Approximately 27 % of the resulting 51 proteins fulfilled the SVM score criteria, which were set according to the RNApred performance for *Synechocystis* proteins (**Supplementary Figure S6**), narrowing the list of RNA chaperone candidates to only 14 proteins (0.67%) of the initially 2,394 detected proteins (**Table 1**).

Manual curation of the shortlist of 14 proteins that had passed the criteria of appropriate cluster assignment, unknown functional categorization, minimum phylogenetic conservation and minimum SVM score identified three proteins that are unlikely to be involved with nucleic acids due to known functions. These are Sll1515, the glutamine synthetase inactivating factor IF17 (Garcia-Dominguez et al., 1999), Slr0169, a CsoD1 ortholog that is part of the β-carboxysome, containing two bacterial microcompartment (BMC) domains (Kinney et al., 2011) or Sll1939, a FtsZ interacting Ftn6 homolog involved in cell division (Marbouty et al., 2009). Nevertheless, we kept them in the list because a small number of bifunctional proteins exist in bacteria that

are active in metabolism and in controlling gene expression, including mechanisms in
which RNA is bound (Commichau and Stülke, 2008).

Another four proteins might interact with nucleic acids but less so with RNA. These
proteins are Ssl2874, a homolog of RemA, an essential regulator of biofilm formation
in *Bacillus subtilis* (Winkelman et al., 2009, 2013), Sll0639, a DHH phosphoesterase
domain-containing protein with similarities to a RecJ domain protein,
nucleotidyltransferase/poly(A) polymerase or exopolyphosphatase-like enzyme,
Slr1081 (DUF820) featuring a restriction endonuclease domain and Sll0284 (DUF87)
containing an N-terminal HAS-barrel domain similar to archaeal proteins, such as
DNA double-strand break repair helicase HerA. However, bacterial proteins
containing similar domains seem to be largely undescribed.

Another four poorly characterized proteins were Slr0211, an acyl-CoA N-
acyltransferase domain containing protein, Sll0319 (DUF3747), a periplasmic protein,
Slr1660 (DUF3172) containing an N-terminal transmembrane domain and Ssl2064, a
75 amino acid DUF4160-containing protein.

Ultimately, three candidates appeared most promising. The first was Sll1424, an
undescribed protein with very conserved synteny to the transcriptional regulator NtcA
(**Supplementary Figure S7A**). Furthermore, we noticed Ssr1238/Slr0743a, an 84
amino acid YlxR homologue with a DUF448 domain and a pronounced synteny in
cyanobacteria, expressed from a gene between *nusA* encoding the transcription
termination factor NusA and *infB* encoding the translation initiation factor InfB
(**Supplementary Figure S7B**). While the order of adjacent genes *rimP-nusA-infB* is
conserved even in *Salmonella* and *E. coli*, a gene corresponding to *ssr1238* is
lacking in their genomes. The relationship of *ylxR* with *rimP-nusA-infB* has also been
observed in recent analyses of cyanobacterial gene linkage networks (Labella et al.,
2020b).

Finally, we identified Slr0287, a KH domain-containing 118 amino acid protein that
shares 30% identical and 56% similar residues with the corresponding protein KhpA
from *S. pneumonia*. KhpA forms together with the JAG/EloR-domain protein KhpB
and RNA-binding complex in *S. pneumonia* (Zheng et al., 2017). Indeed, we noticed
that with Slr1472, a KhpB homolog also exists in *Synechocystis*, and both proteins
are widely conserved throughout the cyanobacterial phylum and even cluster

together in the gradient (**Supplementary Figure S5**). The genes of both proteins are syntenic with genes encoding proteins involved in RNA metabolism and translation, such as *khpA/slr0287* with *rps16* (**Supplementary Figure S7C**) or *khpB/slr1472* with *rpl34, rnpA and slr1470* encoding a membrane protein and *slr1471* encoding a YidC/Oxa1 homolog (**Supplementary Figure S7D**). Slr1471/Oxa1 is essential for thylakoid biogenesis and the membrane integration of the reaction center precursor protein pD1 (Spence et al., 2004; Ossenbühl et al., 2006). The synteny of *rnpA, yidC/oxa1* and *khpB* as well as *khpA* and *rps16* is strikingly wide, extending even to gram-positive bacteria, such as *S. pneumonia* (not shown).

Data accessibility/Visualizations

To facilitate maximum usability of this resource, we have made the entire dataset available at https://trex90.shinyapps.io/GradSeqExplorer_SynPCC6803/, providing multiple tools to focus on different fractions of the proteome or transcriptome, from groups of functionally related proteins and transcripts to the level of the individual protein or sRNA. Multiple filter options, a customizable graphical output, as well as a tabular output options provide extensive background information to the findings presented in this analysis.

Discussion

Based on the interactions with different types of RBPs, five major classes of ncRNAs were differentiated that are common to many bacteria (Hör et al., 2018). These include the 6S RNA as a global modulator of RNA polymerase (RNAP) specificity and activity (Barrick et al., 2005; Wassarman and Storz, 2000), the Hfq-binding sRNAs (Møller et al., 2002; Zhang et al., 1998; Moll et al., 2003), the ProQ-binding sRNAs (Smirnov et al., 2016), the CsrB-like sRNAs that antagonize the translational repressor CsrA (Liu et al., 1997), and CRISPR RNAs (crRNAs), which form together with the CRISPR-associated (Cas) proteins ribonucleoprotein particles for defense (Hale et al., 2009; Brouns et al., 2008). While 6S RNA and crRNAs exist in cyanobacteria, the affiliation of certain sRNAs to certain RBPs has remained unresolved.

Our Grad-seq analysis of *Synechocystis* provides the first global analysis of major
stable RNA-protein complexes in a photosynthetic cyanobacterium. We anticipate
that our data not only provide crucial knowledge for this particular model
cyanobacterium but also constitute a valuable resource for other cyanobacteria,
including the models *Anabaena* (*Nostoc*) sp. PCC 7120 or *Synechococcus*
elongatus.

Synechocystis as a cyanobacterial model deviates substantially from other bacteria
investigated in previous Grad-seq studies (Smirnov et al., 2016; Hör et al., 2020).
One aspect is the magnitude of internal membrane systems and membrane-bound
protein complexes. We showed that even macromolecular membrane-bound protein
complexes remained largely intact during the process, and their in-gradient
localization matched the pigment composition of the gradient in a meaningful way.
Hence, the analysis provides a spatially resolved proteome dataset of high quality for
the macromolecular complexes of a photosynthetic bacterium. In addition, the
occurrence of subunits specific for certain subcomplexes or specialized functions
provides information about the *in vivo* state and illustrates the dynamics in complex
formation. Examples for such acclimation processes include the incorporation of
PsaK2 into PSI as the high light-adapted form instead of the constitutive form, which
incorporates mainly PsaK1. In our data, the occurrence of PsaK2 was restricted to
disassembled and monomeric PSI, while PsaK1 occurred mainly in fully assembled
monomeric and trimeric PSI, capturing the *in vivo* state of *Synechocystis* PSI and a
multitude of other complexes under standard growth conditions.

Our data not only provide information about known protein-protein interactions but
also provide clues about currently unknown interactions, such as the given examples
of the PSII assembly factors Sll0933, Slr1768 and Slr2013, the PSI assembly factor
VIPP1 homolog, the P_{II} interacting protein PipX or the membrane protein PamA with
unexplained occurrences in the heavy fractions and outside the range of their known
interaction partners, which illustrate the potential value of this dataset.

However, some subcomplexes likely became unstable during the procedure and
dissociated from the respective main complex, such as the oxygen evolving complex
of PSII. The possibility that some complexes are only fully stable when membrane
bound cannot be excluded and cannot be avoided during the solubilization of the
membranes. Nevertheless, the chosen clustering approach to divide the content of

the dataset into groups of similarly sedimenting proteins and RNAs proved quite
helpful in that matter. To our surprise, the gradient profiles of the majority of RNAs
diverged vastly from most detected proteins, locating generally in the heavier
fractions and with largely similar sedimentation patterns, independent of transcript
lengths. This fact is a strong indication of association with one or more major RNA
chaperones or to large intact protein complexes involved in transcription or
translation, such as RNAP, ribosomes and associated factors.

The colocalization of many known intact RNA-protein complexes in fractions 8 - 11
(cluster 11-15) involved in transcription, RNA metabolism and translation initiation
shows the stability of such complexes, while the cooccurrence of a large number of
RNAs illustrates that those complexes were still capturing the *in vivo* state of
Synechocystis regulatory processes by binding RNAs. Furthermore, the fact that
these fractions constituted a poor protein complexity but still harbored most of the
well-studied factors involved in protein biosynthesis makes them interesting for the
identification of novel candidates functionally linked to posttranscriptional regulation.

Another surprising observation was that the majority of detected transcripts were
largely not associated with the previously mentioned complexes but mainly
sedimented in fractions 6-7 (cluster 10), indicating the involvement of one or more
RNA chaperones stably binding their target RNAs. The data permit for the first time
the identification of candidate RNA-binding proteins involved in the formation of major
RNA–protein complexes in this group of bacteria. Although *Synechocystis* expresses
multiple sRNAs of crucial relevance for oxygenic photosynthesis and other important
physiological processes (Klähn et al., 2015; Georg et al., 2017, 2014; de Porcellinis
et al., 2016), homologs of CsrA, ProQ or Hfq do not exist in cyanobacteria or are not
RNA binding (Schuergers et al., 2014). Rather, a prediction from our Grad-seq
cluster analysis is that cyanobacteria may not utilize a general RBP analogous to Hfq
or ProQ as characterized in enterobacteria, in which these two RBPs together are
associated with more than 80% of all sRNAs (Holmqvist and Vogel, 2018). On the
one hand, sRNAs in cyanobacteria may frequently function independent of a
common RNA chaperone, or they may rely more upon specialized RBPs. In this
respect, our data resemble the conclusions recently drawn for gram-positive bacteria
based on the Grad-seq analysis of the human pathogen *S. pneumonia* (Hör et al.,
2020). We filtered out a short list of previously uncharacterized protein candidates for

the interaction with sRNAs. Among them, we discovered a cyanobacterial homolog of
the KhpA/B complex (Zheng et al., 2017). The nearly congruent peaks of the
homologs Slr0287 and Slr1472 in gradient fractions 8–10 indicate that these proteins
are part of a higher molecular weight complex, actually much larger than the complex
reported for *S. pneumonia,* which sedimented in fractions 2 and 3.

Materials and Methods

Cell cultivation, cell lysis, gradient preparation and fractionation

Triplicates of 400 mL of *Synechocystis* liquid cultures were cultivated in BG11
medium under constant shaking in standard growth conditions (30 °C, 50 µE) until
reaching exponential phase (OD_{750} = 0.8). The cultures were harvested by
centrifugation at 4000 g for 15 min at RT. After centrifugation, the cell pellets were
resuspended in ice cold lysis buffer (20 mM Tris-HCl, pH 7.5; 150 mM KCl; 1 mM
MgCl2; 1 mM DTT; 0.2 % [v/v] Triton X-100; RiboLock RNase Inhibitor; DNase I;
cOmplete Protease Inhibitor cocktail) and mechanically lysed with glass beads at
4 °C (Precellys). From here on, the samples were kept at 4 °C during the whole
procedure. Unbroken cells were removed by a centrifugation step for 5 min at 4000 g,
and the supernatant was transferred to a fresh tube. The membrane proteins were
solubilized by incubating the samples under shaking for 60 min in the dark with n-
dodecyl β-D-maltoside in a 3:1 ratio according to the protein content. Thylakoid
membranes were removed by centrifugation at 30,000 g for 30 min, and the
supernatant was applied on top of linear sucrose gradients. The 10 - 40 % sucrose
gradients were made using a gradient mixer with two solutions of lysis buffer,
supplemented with either 10 % or 40 % sucrose [w / v]. Ultracentrifugation of the
lysates was performed in a swing out rotor (Beckman SW40 Ti) for 16 h at 285,000 g.
The fractionation of equal volumes (600 µL) was performed manually, and the
fractions were split into samples for MS analysis (100 µL) and library preparation for
RNA sequencing (360 µL).

Spectroscopy

Absorption spectra were measured using a Specord® 250 Plus (Analytik Jena)
spectrophotometer at room temperature and were normalized to equal fraction
volumes.

RNA sample preparation, generation of cDNA libraries and sequencing

Equal volumes of each fraction were spiked with a barcoded 197 bp long spike-in
RNA mix ranging from 1 - 100 fmol and subjected to proteinase K treatment (1 %
SDS, 10 mM Tris-HCl pH 7.5, proteinase K) for 30 min at 50 °C, followed by RNA
extraction with PGTX (Pinto et al., 2009). After the first phase separation step, the
aqueous phase was transferred to a fresh microcentrifuge tube and mixed 1:1 [v / v]
with 100 % EtOH, and RNA was further extracted using the "RNA clean &
concentrator" kit from Zymo according to the manufacturer's instructions. The spike-
in RNA mix was created by in vitro transcription (Mega-shortscript, Thermo Fisher) of
PCR-amplified parts from the genomic DNA of a phiX174 bacteriophage genome
(**Supplementary Table S3**), purified from a 10% urea-PAA gel (RNA PAGE
Recovery, (Zymo). Determination of RNA concentration was performed using the
Qubit RNA HS Assay kit (Thermo Fisher). For the control, equal volumes of purified
RNA were separated on a 10% denaturing urea-PAA gel.

Due to low protein yields in fractions 14 - 17, fractions 14 and 15, as well as fractions
16 and 17, were pooled prior to RNA sample preparation, resulting in a total of 16
samples per replicate for transcriptome analysis. The library preparation and Illumina
paired end sequencing (75 bp read length) were performed by vertis Biotechnologie
AG. In short, RNA samples were fragmented using ultrasound (4 pulses of 30 s each
at 4 °C), followed by 3' end adapter ligation and first strand cDNA synthesis using M-
MLV reverse transcriptase. The 5' Illumina TrueSeq sequencing adapter was ligated
to the 3' end of the antisense from the purified first-strand cDNA, and the resulting
cDNA was PCR amplified (11 cycles) and purified using the Agencourt AMPure XP
kit (Beckman Coulter Genomics). For Illumina NextSeq sequencing, the samples
were pooled, and the cDNA pool was eluted in the size range of 160 - 500 bp from a
preparative agarose gel. The primers used for PCR amplification were designed for
TruSeq sequencing according to the instructions of Illumina (**Supplementary Table
S3**).

Read mapping and normalization

The analysis of the raw reads from the RNA-sequencing was performed using the galaxy web platform. The workflow can be accessed and reproduced with the following link: https://usegalaxy.eu/u/mr559/w/-grad-seq-pipeline. In short, the read quality was checked at the beginning and after each step with FastQC. Adapter trimming was performed using Cutadapt (Martin, 2011), and the trimmed reads were mapped to the *Synechocystis* genome using segemehl (Otto et al., 2014) with default parameters and were assigned to annotated regions using FeatureCounts. Prior to normalization, all features with a maximum read count ≤ 5 across all fractions were filtered prior to downstream analysis. Fraction-wise normalization was performed by normalizing according to the mapped reads of the barcodes from the spike-in RNAs to compensate for the huge differences in RNA amounts between the fractions. The reproducibility between the replicates was checked by comparing the distribution of the Spearman correlation coefficient from the relative distribution of each RNA within a replicate with the same RNAs in the other replicates (**Supplementary Figure S1A**).

Sample preparation for mass spectrometry-based proteome measurements

As an initial step, proteins were purified by acetone/methanol precipitation. Therefore, 100 µL of each of the initial 18 fractions were mixed with an ice-cold mixture of 800 µL acetone and 100 µL methanol. After incubation o.n. at -20 °C, precipitated proteins were subsequently collected by centrifugation at 1000 × g and 4 °C. Subsequently, protein pellets were washed twice with 1 mL cold 80 % (v / v) acetone (aq.). During the washing procedure, precipitated sucrose was removed almost completely, also in fractions with high sucrose content. Due to low protein yields, fractions were partially pooled (F14+15 and F16+17), resulting in a total of 16 samples per replicate for proteome analysis.

Dried protein pellets were consequently redissolved in 50 µL denaturation buffer (6 M urea, 2 M thiourea in 100 mM Tris/HCl, pH 7.5). Intrachain disulfide bonds were reduced with dithiothreitol and resulting thiol groups were alkylated with iodoacetamide as described before (Spät et al., 2015). Subsequently, proteins were predigested with endoprotease Lys-C for 3 h, further diluted with 200 µL 20 mM

ammonium bicarbonate buffer, pH 8.0, and digested with trypsin o.n., applying a
protease:protein ratio of 1:100 for both enzymes. The resulting peptide solutions
were acidified to pH 2.5 with 10 % (v / v) trifluoroacetic acid (aq.) and purified
following the stage tip protocol (Rappsilber et al., 2007).

For MS-based proteome measurements, a constant sample volume was analyzed for
all fractions of one replicate. Consequently, the measured protein yields ranged
between a maximum of 1000 ng and <100 ng in fractions with the lowest protein
concentrations (**Supplementary Table S4**). As a control, equal volumes of the initial
protein fractions were separated in 15% SDS-PAA gels and visualized by Coomassie
staining.

MS measurements

For nanoLC-MS/MS-based proteome measurements, purified samples were loaded
onto an in-house produced C18 nanoHPLC column (20 cm PicoTip fused silica
emitter with 75 µm inner diameter (New Objective, Woburn, USA), packed with
1.9 µm ReproSil-Pur C18-AQ resin (Dr Maisch, Ammerbuch, Germany)) on an
EASY-nLC 1200 system (Thermo Scientific, Bremen, Germany). Peptides were
separated using a 49 min segmented linear gradient (**Supplementary Table S5**),
and eluting peptides were directly ionized through an on-line coupled electrospray
ionization (ESI) source and analyzed on a Q Exactive HF mass spectrometer
(Thermo Scientific, Bremen, Germany), operated in the positive-ion mode and data
dependent acquisition. MS full scans were acquired with a mass-to-charge (m/z)
range of 300 - 1650 at a resolution of 60,000. The 12 most intense multiple charged
ions were selected for fragmentation by higher-energy collisional dissociation (HCD)
and recorded in MS/MS scans with a resolution of 30,000. Dynamic exclusion of
sequenced precursor ions was set to 30 s. Wash runs (**Supplementary Table S5**)
were performed after each of the three samples.

Proteomics data processing

Acquired raw MS data of all three replicates were processed using the MaxQuant
software suite (version 1.5.2.8) (Cox and Mann, 2008). Each file was defined as an

individual sample. The data processing was performed at default settings, applying

intensity based absolute quantification (iBAQ) and the following search criteria:

trypsin was defined as cleaving enzyme and a maximum of two missed cleavages

was allowed. Carbamidomethylation of cysteine residues was set as a fixed

modification and methionine oxidation and protein N-terminal acetylation were

defined as variable modifications. MS/MS spectra were searched against a combined

target-decoy database of *Synechocystis* downloaded from Cyanobase

(http://genome.microbedb.jp/cyanobase; version 06/2014), and the sequences of

newly discovered small proteins (Mitschke et al., 2011; Kopf et al., 2014;

Baumgartner et al., 2016), representing a total of 3,681 protein sequences plus 245

common contaminants. To enable the identification of low abundant proteins which

were not selected for MS/MS sequencing in any fraction, ancillary measurements

were performed for the first replicate with a constant protein yield of 1000 ng per

fraction. These supporting measurements were included in the data processing as a

resource to promote overall protein identifications via the matching between runs

option. Reported protein and peptide false discovery rates were each limited to 1%.

The reproducibility between the replicates was checked by comparing the distribution

of the Spearman correlation coefficient from the relative distribution of each protein

within a replicate with the same protein in the other replicates (**Supplementary**

Figure S1B).

Hierarchical clustering and module detection

Clustering and module detection were performed using the dynamic tree cut

algorithm implemented in the WGCNA R package (Langfelder and Horvath, 2008,

2012) from the standardized relative abundances of the normalized RNA-seq reads

and IBAQ intensities.

RNApred benchmarking for *Synechocystis*

The performance of RNApred was tested for *Synechocystis* proteins categorized into

Gene Ontology groups "Nucleotide binding", "DNA binding", "RNA binding" or "No

binding" if no other category was assigned. The majority of the "no binding" group did

not reach an SVM score >0.49, while the majority of the "RNA binding" group

reached an SVM score of at least 0.26 (**Supplementary Figure S6A**). The distribution of those groups barely overlapped, showing the validity of RNApred for *Synechocystis* proteins. Furthermore, it was tested against all ribosomal proteins (**Supplementary Figure S6B**) and selected known RNA-binding proteins (**Supplementary Figure S6C**). Notably, the cyanobacterial Hfq failed to fulfill the set criteria by far, matching previous observations of *Synechocystis* Hfq as not binding RNA (Schuergers et al., 2014), thus confirming this approach. The groups "Nucleotide binding" and "DNA binding" were distributed between those two other groups, with the majority of the "DNA binding" group below an SVM score of 1.07. Therefore, a conservative minimum SVM score of at least ≥ 0.5 should serve sufficient to remove most proteins without nucleic acid binding properties. The median SVM score of all orthologs in the genomes of **Supplementary Table S2** was included to enhance the robustness of the approach, applying a minimum score of ≥ -0.2, which is the default threshold for RBPs in the webserver.

Data availability

The datasets produced in this study are available in the following databases:

- FastQ files from RNA sequencing in SRA (BioProject accession number PRJNA608723): https://www.ncbi.nlm.nih.gov/sra/PRJNA608723
- Galaxy workflow used for raw read processing: https://usegalaxy.eu/u/mr559/w/-grad-seq-pipeline
- Mass spectrometry raw data deposited at the ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via the PRIDE partner repository (Vizcaíno et al., 2013) under the identifier: PXD020025
- Data accessibility and visualization: https://trex90.shinyapps.io/GradSeqExplorer_SynPCC6803/

Acknowledgments

Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - 322977937/GRK2344 to WRH and through the "SCyCode" research group 397695561/FOR 2816 to WRH and BM.

Author contributions

WRH and MR designed the research and performed the experiments and bioinformatics analyses. PS and BM carried out the MS and the related analyses. MR developed the bioinformatics analysis workflow. RB performed validation experiments. WRH and MR wrote the manuscript with input from all authors. WRH and BM supervised and funded the project.

Conflict of interest

The authors declare that they have no conflicts of interest.

References

Athukoralage, J.S., McQuarrie, S., Grüschow, S., Graham, S., Gloster, T.M., and White, M.F. (2020). Tetramerisation of the CRISPR ring nuclease Crn3/Csx3 facilitates cyclic oligoadenylate cleavage. eLife 9.

Baers, L.L., Breckels, L.M., Mills, L.A., Gatto, L., Deery, M.J., Stevens, T.J., Howe, C.J., Lilley, K.S., and Lea-Smith, D.J. (2019). Proteome mapping of a cyanobacterium reveals distinct compartment organization and cell-dispersed metabolism. Plant Physiol. 181: 1721–1738.

Bandyra, K.J. and Luisi, B.F. (2018). RNase E and the high-fidelity orchestration of RNA metabolism. Microbiol. Spectr. 6.

Barrick, J.E., Sudarsan, N., Weinberg, Z., Ruzzo, W.L., and Breaker, R.R. (2005). 6S RNA is a widespread regulator of eubacterial RNA polymerase that resembles an open promoter. RNA 11: 774–784.

Baumgartner, D., Kopf, M., Klähn, S., Steglich, C., and Hess, W.R. (2016). Small proteins in cyanobacteria provide a paradigm for the functional analysis of the bacterial micro-proteome. BMC Microbiol. 16: 285.

Behler, J., Sharma, K., Reimann, V., Wilde, A., Urlaub, H., and Hess, W.R. (2018). The host-encoded RNase E endonuclease as the crRNA maturation enzyme in a CRISPR–Cas subtype III-Bv system. Nat. Microbiol. 3: 367–377.

Bøggild, A., Overgaard, M., Valentin-Hansen, P., and Brodersen, D.E. (2009). Cyanobacteria contain a structural homologue of the Hfq protein with altered RNA-binding properties. FEBS J. 276: 3904–3915.

Brouns, S.J.J., Jore, M.M., Lundgren, M., Westra, E.R., Slijkhuis, R.J.H., Snijders, A.P.L., Dickman, M.J., Makarova, K.S., Koonin, E.V., and van der Oost, J. (2008). Small CRISPR RNAs guide antiviral defense in prokaryotes. Science 321: 960–964.

Cantos, R., Labella, J.I., Espinosa, J., and Contreras, A. (2019). The nitrogen regulator PipX acts in cis to prevent operon polarity. Environ. Microbiol. Rep. 11: 495–507.

Cech, T.R. and Steitz, J.A. (2014). The noncoding RNA revolution-trashing old rules to forge new ones. Cell 157: 77–94.

Chitnis, V.P. and Chitnis, P.R. (1993). PsaL subunit is required for the formation of photosystem I trimers in the cyanobacterium Synechocystis sp. PCC 6803. FEBS Lett. 336: 330–334.

Commichau, F.M. and Stülke, J. (2008). Trigger enzymes: bifunctional proteins active in metabolism and in controlling gene expression. Mol. Microbiol. 67: 692–702.

Cox, J. and Mann, M. (2008). MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat. Biotechnol. 26: 1367–1372.

Dienst, D., Dühring, U., Möllenkopf, H.-J., Vogel, J., Golecki, J., Hess, W.R., and Wilde, A. (2008). The cyanobacterial homologue of the RNA chaperone Hfq is essential for motility of Synechocystis sp. PCC 6803. Microbiology 154: 3134–3143.

Dühring, U., Ossenbühl, F., and Wilde, A. (2007). Late assembly steps and dynamics of the cyanobacterial ühotosystem I. J. Biol. Chem. 282: 10915–10921.

Forcada-Nadal, A., Llácer, J.L., Contreras, A., Marco-Marín, C., and Rubio, V.
(2018). The PII-NAGK-PipX-NtcA regulatory axis of Cyanobacteria: A tale of
changing partners, allosteric effectors and non-covalent interactions. Front.
Mol. Biosci. 5: 91.

Fujimori, T., Hihara, Y., and Sonoike, K. (2005). PsaK2 subunit in photosystem I is
involved in state transition under high light condition in the cyanobacterium
Synechocystis sp. PCC 6803. J. Biol. Chem. 280: 22191–22197.

Gao, F., Zhao, J., Chen, L., Battchikova, N., Ran, Z., Aro, E.-M., Ogawa, T., and Ma,
W. (2016). The NDH-1L-PSI supercomplex is important for efficient cyclic
electron transport in cyanobacteria. Plant Physiol. 172: 1451–1464.

Garcia-Dominguez, M., Reyes, J.C., and Florencio, F.J. (1999). Glutamine
synthetase inactivation by protein-protein interaction. Proc. Natl. Acad. Sci. 96:
7161–7166.

Garcia-Doval, C., Schwede, F., Berk, C., Rostøl, J.T., Niewoehner, O., Tejero, O.,
Hall, J., Marraffini, L.A., and Jinek, M. (2020). Activation and self-inactivation
mechanisms of the cyclic oligoadenylate-dependent CRISPR ribonuclease
Csm6. Nat. Commun. 11: 1596.

Georg, J. et al. (2017). Acclimation of oxygenic photosynthesis to iron starvation is
controlled by the sRNA IsaR1. Curr. Biol. 27: 1425-1436.e7.

Georg, J., Dienst, D., Schurgers, N., Wallner, T., Kopp, D., Stazic, D., Kuchmina, E.,
Klahn, S., Lokstein, H., Hess, W.R., and Wilde, A. (2014). The small regulatory
RNA SyR1/PsrR1 controls photosynthetic functions in cyanobacteria. Plant
Cell 26: 3661–3679.

Gunnelius, L., Hakkila, K., Kurkela, J., Wada, H., Tyystjärvi, E., and Tyystjärvi, T.
(2014). The omega subunit of the RNA polymerase core directs transcription
efficiency in cyanobacteria. Nucleic Acids Res. 42: 4606–4614.

Hagemann, M. and Hess, W.R. (2018). Systems and synthetic biology for the
biotechnological application of cyanobacteria. Curr Opin Biotechnol 49: 94–99.

Hale, C.R., Zhao, P., Olson, S., Duff, M.O., Graveley, B.R., Wells, L., Terns, R.M.,
and Terns, M.P. (2009). RNA-guided RNA cleavage by a CRISPR RNA-Cas
protein complex. Cell 139: 945–956.

Han, W., Stella, S., Zhang, Y., Guo, T., Sulek, K., Peng-Lundgren, L., Montoya, G.,
and She, Q. (2018). A Type III-B Cmr effector complex catalyzes the synthesis
of cyclic oligoadenylate second messengers by cooperative substrate binding.
Nucleic Acids Res. 46: 10319–10330.

He, Z. and Mi, H. (2016). Functional characterization of the subunits N, H, J, and O of
the NAD(P)H dehydrogenase complexes in *Synechocystis* sp. strain PCC
6803. Plant Physiol. 171: 1320–1332.

Heilmann, B., Hakkila, K., Georg, J., Tyystjärvi, T., Hess, W.R., Axmann, I.M., and
Dienst, D. (2017). 6S RNA plays a role in recovery from nitrogen depletion in
Synechocystis sp. PCC 6803. BMC Microbiol. 17: 229.

Holmqvist, E. and Vogel, J. (2018). RNA-binding proteins in bacteria. Nat. Rev.
Microbiol. 16: 601–615.

Hör, J., Garriss, G., Di Giorgio, S., Hack, L.-M., Vanselow, J.T., Förstner, K.U.,
Schlosser, A., Henriques-Normark, B., and Vogel, J. (2020). Grad-seq in a
Gram-positive bacterium reveals exonucleolytic sRNA activation in
competence control. EMBO J.: e103852.

Hör, J., Gorski, S.A., and Vogel, J. (2018). Bacterial RNA biology on a genome scale.
Mol. Cell 70: 785–799.

Hrle, A., Maier, L.-K., Sharma, K., Ebert, J., Basquin, C., Urlaub, H., Marchfelder, A.,
and Conti, E. (2014). Structural analyses of the CRISPR protein Csc2 reveal
the RNA-binding interface of the type I-D Cas7 family. RNA Biol. 11: 1072–
1082.

Iyer, L.M., Koonin, E.V., and Aravind, L. (2004). Evolution of bacterial RNA
polymerase: implications for large-scale bacterial phylogeny, domain
accretion, and horizontal gene transfer. Gene 335: 73–88.

Kanehisa, M. (2019). Toward understanding the origin and evolution of cellular
organisms. Protein Sci. 28: 1947–1951.

Kanehisa, M. and Goto, S. (2000). KEGG: Kyoto encyclopedia of genes and
genomes. Nucleic Acids Res. 28: 27–30.

Kanehisa, M., Sato, Y., Furumichi, M., Morishima, K., and Tanabe, M. (2019). New
approach for understanding genome variations in KEGG. Nucleic Acids Res.
47: D590–D595.

Kazlauskiene, M., Kostiuk, G., Venclovas, Č., Tamulaitis, G., and Siksnys, V. (2017).
A cyclic oligonucleotide signaling pathway in type III CRISPR-Cas systems.
Science 357: 605–609.

Kieper, S.N., Almendros, C., Behler, J., McKenzie, R.E., Nóbrega, F.L., Haagsma,
A.C., Vink, J.N.A., Hess, W.R., and Brouns, S.J.J. (2018). Cas4 facilitates
PAM-compatible spacer selection during CRISPR adaptation. Cell Rep. 22:
3377–338.

Kinney, J.N., Axen, S.D., and Kerfeld, C.A. (2011). Comparative analysis of
carboxysome shell proteins. Photosynth. Res. 109: 21–32.

Klähn, S., Bolay, P., Wright, P.R., Atilho, R.M., Brewer, K.I., Hagemann, M., Breaker,
R.R., and Hess, W.R. (2018). A glutamine riboswitch is a key element for the
regulation of glutamine synthetase in cyanobacteria. Nucleic Acids Res. 46:
10082–10094.

Klähn, S., Schaal, C., Georg, J., Baumgartner, D., Knippen, G., Hagemann, M.,
Muro-Pastor, A.M., and Hess, W.R. (2015). The sRNA NsiR4 is involved in
nitrogen assimilation control in cyanobacteria by targeting glutamine
synthetase inactivating factor IF7. Proc. Natl. Acad. Sci. 112: E6243–E6252.

Kondo, K., Geng, X.X., Katayama, M., and Ikeuchi, M. (2005). Distinct roles of
CpcG1 and CpcG2 in phycobilisome assembly in the cyanobacterium
Synechocystis sp. PCC 6803. Photosynth. Res. 84: 269–273.

Kondo, K., Mullineaux, C.W., and Ikeuchi, M. (2009). Distinct roles of CpcG1-
phycobilisome and CpcG2-phycobilisome in state transitions in a
cyanobacterium Synechocystis sp. PCC 6803. Photosynth. Res. 99: 217–225.

Kondo, K., Ochiai, Y., Katayama, M., and Ikeuchi, M. (2007). The membrane-
associated CpcG2-phycobilisome in *Synechocystis*: A new photosystem I
antenna. Plant Physiol. 144: 1200–1210.

Kopf, M. and Hess, W.R. (2015). Regulatory RNAs in photosynthetic cyanobacteria.
FEMS Microbiol. Rev. 39: 301–315.

Kopf, M., Klähn, S., Scholz, I., Matthiessen, J.K.F., Hess, W.R., and Voß, B. (2014).
Comparative analysis of the primary transcriptome of *Synechocystis* sp. PCC
6803. DNA Res. 21: 527–539.

Kopfmann, S., Roesch, S.K., and Hess, W.R. (2016). Type II toxin-antitoxin systems
in the unicellular cyanobacterium *Synechocystis* sp. PCC 6803. Toxins 8:
228.1-228.23.

Kumar, M., Gromiha, M.M., and Raghava, G.P.S. (2011). SVM based prediction of
RNA-binding proteins using binding residues and evolutionary information. J.
Mol. Recognit. 24: 303–313.

Labella, J.I., Cantos, R., Salinas, P., Espinosa, J., and Contreras, A. (2020a).
Distinctive features of PipX, a unique signaling protein of Cyanobacteria. Life
10: 79.

Labella, J.I., Llop, A., and Contreras, A. (2020b). The default cyanobacterial linked
genome: an interactive platform based on cyanobacterial linkage networks to
assist functional genomics. FEBS Lett. 594: 1661–1674.

Labella, J.I., Obrebska, A., Espinosa, J., Salinas, P., Forcada-Nadal, A., Tremiño, L.,
Rubio, V., and Contreras, A. (2016). Expanding the cyanobacterial nitrogen
regulatory network: The GntR-like regulator PlmA interacts with the PII-PipX
complex. Front. Microbiol. 7: 1677.

Langfelder, P. and Horvath, S. (2012). Fast *R* functions for robust correlations and
hierarchical clustering. J. Stat. Softw. 46: i11.

Langfelder, P. and Horvath, S. (2008). WGCNA: an R package for weighted
correlation network analysis. BMC Bioinformatics 9: 559.

Liu, H., Weisz, D.A., Zhang, M.M., Cheng, M., Zhang, B., Zhang, H., Gerstenecker,
G.S., Pakrasi, H.B., Gross, M.L., and Blankenship, R.E. (2019).
Phycobilisomes harbor FNR $_L$ in Cyanobacteria. mBio 10: e00669-19,
/mbio/10/2/mBio.00669-19.atom.

Liu, M.Y., Gui, G., Wei, B., Preston, J.F., Oakford, L., Yüksel, U., Giedroc, D.P., and
Romeo, T. (1997). The RNA molecule CsrB binds to the global regulatory
protein CsrA and antagonizes its activity in *Escherichia coli*. J. Biol. Chem.
272: 17502–17510.

Llacer, J.L., Espinosa, J., Castells, M.A., Contreras, A., Forchhammer, K., and Rubio,
V. (2010). Structural basis for the regulation of NtcA-dependent transcription
by proteins PipX and PII. Proc. Natl. Acad. Sci. 107: 15397–15402.

Makarova, K.S. et al. (2015). An updated evolutionary classification of CRISPR-Cas
systems. Nat. Rev. Microbiol. 13: 722–736.

Makarova, K.S., Anantharaman, V., Grishin, N.V., Koonin, E.V., and Aravind, L.
(2014). CARF and WYL domains: ligand-binding regulators of prokaryotic
defense systems. Front. Genet. 5: 102.

Marbouty, M., Saguez, C., Cassier-Chauvat, C., and Chauvat, F. (2009).
Characterization of the FtsZ-interacting septal proteins SepF and Ftn6 in the
spherical-celled cyanobacterium *Synechocystis* strain PCC 6803. J. Bacteriol.
191: 6178–6185.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet.journal 17: 10.

Melamed, S., Adams, P.P., Zhang, A., Zhang, H., and Storz, G. (2020). RNA-RNA
interactomes of ProQ and Hfq reveal overlapping and competing roles. Mol.
Cell 77: 411-425.e7.

Mitschke, J., Georg, J., Scholz, I., Sharma, C.M., Dienst, D., Bantscheff, J., Voß, B.,
Steglich, C., Wilde, A., Vogel, J., and Hess, W.R. (2011). An experimentally

anchored map of transcriptional start sites in the model cyanobacterium *Synechocystis* sp. PCC6803. Proc. Natl. Acad. Sci. 108: 2124–2129.

Molina, R., Stella, S., Feng, M., Sofos, N., Jauniskis, V., Pozdnyakova, I., López-Méndez, B., She, Q., and Montoya, G. (2019). Structure of Csx1-cOA4 complex reveals the basis of RNA decay in Type III-B CRISPR-Cas. Nat. Commun. 10: 4302.

Moll, I., Leitsch, D., Steinhauser, T., and Bläsi, U. (2003). RNA chaperone activity of the Sm-like Hfq protein. EMBO Rep. 4: 284–289.

Møller, T., Franch, T., Højrup, P., Keene, D.R., Bächinger, H.P., Brennan, R.G., and Valentin-Hansen, P. (2002). Hfq: a bacterial Sm-like protein that mediates RNA-RNA interaction. Mol. Cell 9: 23–30.

Morris, K.V. and Mattick, J.S. (2014). The rise of regulatory RNA. Nat. Rev. Genet. 15: 423–437.

Mutsuda, M. and Sugiura, M. (2006). Translation initiation of cyanobacterial rbcS mRNAs requires the 38-kDa ribosomal protein S1 but not the Shine-Dalgarno sequence: development of a cyanobacterial in vitro translation system. J. Biol. Chem. 281: 38314–38321.

Nakagawa, S., Niimura, Y., Miura, K., and Gojobori, T. (2010). Dynamic evolution of translation initiation mechanisms in prokaryotes. Proc. Natl. Acad. Sci. U. S. A. 107: 6382–6387.

Nickelsen, J. and Rengstl, B. (2013). Photosystem II assembly: from cyanobacteria to plants. Annu. Rev. Plant Biol. 64: 609–635.

Niewoehner, O., Garcia-Doval, C., Rostøl, J.T., Berk, C., Schwede, F., Bigler, L., Hall, J., Marraffini, L.A., and Jinek, M. (2017). Type III CRISPR-Cas systems produce cyclic oligoadenylate second messengers. Nature 548: 543–548.

Ning, D., Zhao, W., and Qian, Y. (2013). [A hypothetical gene pair, ssl2138 and sll11092, constitutes a functional TA system on the chromosome of *Synechocystis* PCC 6803]. Wei Sheng Wu Xue Bao 53: 1043–1049.

Osanai, T., Sato, S., Tabata, S., and Tanaka, K. (2005). Identification of PamA as a PII-binding membrane protein important in nitrogen-related and sugar-catabolic gene expression in *Synechocystis* sp. PCC 6803. J. Biol. Chem. 280: 34684–34690.

Ossenbühl, F., Inaba-Sulpice, M., Meurer, J., Soll, J., and Eichacker, L.A. (2006). The *Synechocystis* sp PCC 6803 Oxa1 homolog is essential for membrane integration of reaction center precursor protein pD1. Plant Cell 18: 2236–2246.

Otto, C., Stadler, P.F., and Hoffmann, S. (2014). Lacking alignments? The next-generation sequencing mapper segemehl revisited. Bioinforma. Oxf. Engl. 30: 1837–1843.

Pinto, F., Thapper, A., Sontheim, W., and Lindblad, P. (2009). Analysis of current and alternative phenol based RNA extraction methodologies for cyanobacteria. BMC Mol. Biol. 10: 79.

de Porcellinis, A.J., Klähn, S., Rosgaard, L., Kirsch, R., Gutekunst, K., Georg, J., Hess, W.R., and Sakuragi, Y. (2016). The non-coding RNA Ncr0700/PmgR1 is required for photomixotrophic growth and the regulation of glycogen accumulation in the cyanobacterium *Synechocystis* sp. PCC 6803. Plant Cell Physiol. 57: 2091–2103.

Rappsilber, J., Mann, M., and Ishihama, Y. (2007). Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. Nat. Protoc. 2: 1896–1906.

Rast, A., Rengstl, B., Heinz, S., Klingl, A., and Nickelsen, J. (2016). The role of Slr0151, a tetratricopeptide repeat protein from *Synechocystis* sp. PCC 6803, during photosystem II assembly and repair. Front. Plant Sci. 7: 605.

Reimann, V., Alkhnbashi, O.S., Saunders, S.J., Scholz, I., Hein, S., Backofen, R., and Hess, W.R. (2017). Structural constraints and enzymatic promiscuity in the Cas6-dependent generation of crRNAs. Nucleic Acids Res. 45: 915–925.

Rengstl, B., Knoppová, J., Komenda, J., and Nickelsen, J. (2013). Characterization of a *Synechocystis* double mutant lacking the photosystem II assembly factors YCF48 and Sll0933. Planta 237: 471–480.

Rosana, A.R.R., Denise S. Whitford, Migur, A., Steglich, C., Kujat-Choy, S.L., Hess Wolfgang R., and Owttrim George W. (2020). RNA helicase-regulated processing of the *Synechocystis rimO-crhR* operon results in differential cistron expression and accumulation of two sRNAs. J Biol Chem 295: 6372–6386.

Sato, H., Fujimori, T., and Sonoike, K. (2008). sll1961 is a novel regulator of phycobilisome degradation during nitrogen starvation in the cyanobacterium *Synechocystis* sp. PCC 6803. FEBS Lett. 582: 1093–1096.

Schneider, G.J. and Hasekorn, R. (1988). RNA polymerase subunit homology among cyanobacteria, other eubacteria and archaebacteria. J. Bacteriol. 170: 4136–4140.

Scholz, I., Lange, S.J., Hein, S., Hess, W.R., and Backofen, R. (2013). CRISPR-Cas systems in the cyanobacterium *Synechocystis* sp. PCC6803 exhibit distinct processing pathways involving at least two Cas6 and a Cmr2 protein. PloS One 8: e56470.

Scholz, I., Lott, S.C., Behler, J., Gärtner, K., Hagemann, M., and Hess, W.R. (2019). Divergent methylation of CRISPR repeats and cas genes in a subtype I-D CRISPR-Cas-system. BMC Microbiol. 19: 147.1-147.11.

Schuergers, N., Ruppert, U., Watanabe, S., Nürnberg, D.J., Lochnit, G., Dienst, D., Mullineaux, C.W., and Wilde, A. (2014). Binding of the RNA chaperone Hfq to the type IV pilus base is crucial for its function in Synechocystis sp. PCC 6803. Mol. Microbiol. 92: 840–852.

Schuster, C.F. and Bertram, R. (2013). Toxin-antitoxin systems are ubiquitous and versatile modulators of prokaryotic cell fate. FEMS Microbiol. Lett. 340: 73–85.

Schwarz, D., Schubert, H., Georg, J., Hess, W.R., and Hagemann, M. (2013). The gene sml0013 of *Synechocystis* sp. strain PCC 6803 encodes for a novel subunit of the NAD(P)H oxidoreductase or complex I that is ubiquitously distributed among Cyanobacteria. Plant Physiol. 163: 1191–1202.

Shah, S.A., Alkhnbashi, O.S., Behler, J., Han, W., She, Q., Hess, W.R., Garrett, R.A., and Backofen, R. (2019). Comprehensive search for accessory proteins encoded with archaeal and bacterial type III CRISPR-cas gene cassettes reveals 39 new cas gene families. RNA Biol. 16: 530–542.

Shmakov, S.A., Makarova, K.S., Wolf, Y.I., Severinov, K.V., and Koonin, E.V. (2018). Systematic prediction of genes functionally linked to CRISPR-Cas systems by gene neighborhood analysis. Proc. Natl. Acad. Sci. 115: E5307–E5316.

Smirnov, A., Förstner, K.U., Holmqvist, E., Otto, A., Günster, R., Becher, D.,
Reinhardt, R., and Vogel, J. (2016). Grad-seq guides the discovery of ProQ as
a major small RNA-binding protein. Proc. Natl. Acad. Sci. 113: 11591–11596.

Spät, P., Macek, B., and Forchhammer, K. (2015). Phosphoproteome of the
cyanobacterium *Synechocystis* sp. PCC 6803 and its dynamics during
nitrogen starvation. Front. Microbiol. 6.

Spence, E., Bailey, S., Nenninger, A., Møller, S.G., and Robinson, C. (2004). A
homolog of Albino3/OxaI is essential for thylakoid biogenesis in the
cyanobacterium *Synechocystis* sp. PCC6803. J. Biol. Chem. 279: 55792–
55800.

Srikumar, A., Krishna, P.S., Sivaramakrishna, D., Kopfmann, S., Hess, W.R., Swamy,
M.J., Lin-Chao, S., and Prakash, J.S.S. (2017). The Ssl2245-Sll1130 Toxin-
Antitoxin system mediates heat-induced programmed cell death in
Synechocystis sp. PCC6803. J. Biol. Chem. 292: 4222–4234.

Trautmann, D., Voss, B., Wilde, A., Al-Babili, S., and Hess, W.R. (2012).
Microevolution in cyanobacteria: re-sequencing a motile substrain of
Synechocystis sp. PCC 6803. DNA Res. 19: 435–448.

Turmel, M., Gagnon, M.-C., O'Kelly, C.J., Otis, C., and Lemieux, C. (2009). The
chloroplast genomes of the green algae Pyramimonas, Monomastix, and
Pycnococcus shed new light on the evolutionary history of prasinophytes and
the origin of the secondary chloroplasts of euglenids. Mol. Biol. Evol. 26: 631–
648.

Uchiyama, I., Mihara, M., Nishide, H., Chiba, H., and Kato, M. (2019). MBGD update
2018: microbial genome database based on hierarchical orthology relations
covering closely related and distantly related comparisons. Nucleic Acids Res.
47: D382–D389.

Vijay, D., Akhtar, M.K., and Hess, W.R. (2019). Genetic and metabolic advances in
the engineering of cyanobacteria. Curr. Opin. Biotechnol. 59: 150–156.

Vizcaíno, J.A. et al. (2013). The PRoteomics IDEntifications (PRIDE) database and
associated tools: status in 2013. Nucleic Acids Res. 41: D1063-1069.

Wagner, E.G.H. and Romby, P. (2015). Small RNAs in bacteria and archaea: who
they are, what they do, and how they do it. Adv. Genet. 90: 133–208.

Wassarman, K.M. and Storz, G. (2000). 6S RNA regulates E. coli RNA polymerase
activity. Cell 101: 613–623.

Weisz, D.A., Johnson, V.M., Niedzwiedzki, D.M., Shinn, M.K., Liu, H., Klitzke, C.F.,
Gross, M.L., Blankenship, R.E., Lohman, T.M., and Pakrasi, H.B. (2019). A
novel chlorophyll protein complex in the repair cycle of photosystem II. Proc.
Natl. Acad. Sci. U. S. A. 116: 21907–21913.

Westermann, A.J., Venturini, E., Sellin, M.E., Förstner, K.U., Hardt, W.-D., and Vogel,
J. (2019). The major RNA-binding protein ProQ impacts virulence gene
expression in *Salmonella enterica* serovar Typhimurium. mBio 10: e02504-18.

Winkelman, J.T., Blair, K.M., and Kearns, D.B. (2009). RemA (YlzA) and RemB
(YaaB) regulate extracellular matrix operon expression and biofilm formation in
Bacillus subtilis. J. Bacteriol. 191: 3981–3991.

Winkelman, J.T., Bree, A.C., Bate, A.R., Eichenberger, P., Gourse, R.L., and Kearns,
D.B. (2013). RemA is a DNA-binding protein that activates biofilm matrix gene
expression in *Bacillus subtilis*: RemA activates matrix genes directly. Mol.
Microbiol. 88: 984–997.

Xu, C., Wang, B., Yang, L., Hu, L.Z., Yi, L., Wang, Y., Chen, S., Emili, A., and Wan,
C. (2020). Global landscape of native protein complexes in *Synechocystis* sp.
PCC 6803. bioRxiv: 2020.03.07.980128.

Yamaguchi, K. and Subramanian, A.R. (2003). Proteomic identification of all plastid-
specific ribosomal proteins in higher plant chloroplast 30S ribosomal subunit.
Eur. J. Biochem. 270: 190–205.

Zhang, A., Altuvia, S., Tiwari, A., Argaman, L., Hengge-Aronis, R., and Storz, G.
(1998). The OxyS regulatory RNA represses *rpoS* translation and binds the
Hfq (HF-I) protein. EMBO J. 17: 6061–6068.

Zhang, J., Gao, F., Zhao, J., Ogawa, T., Wang, Q., and Ma, W. (2014a). NdhP is an
exclusive subunit of large complex of NADPH dehydrogenase essential to
stabilize the complex in *Synechocystis* sp. strain PCC 6803. J. Biol. Chem.
289: 18770–18781.

Zhang, J.-Y., Deng, X.-M., Li, F.-P., Wang, L., Huang, Q.-Y., Zhang, C.-C., and Chen,
W.-L. (2014b). RNase E forms a complex with polynucleotide phosphorylase
in cyanobacteria via a cyanobacterial-specific nonapeptide in the noncatalytic
region. RNA N. Y. N 20: 568–579.

Zheng, J.J., Perez, A.J., Tsui, H.-C.T., Massidda, O., and Winkler, M.E. (2017).
Absence of the KhpA and KhpB (JAG/EloR) RNA-binding proteins suppresses
the requirement for PBP2b by overproduction of FtsA in *Streptococcus*
pneumoniae D39. Mol. Microbiol. 106: 793–814.

1176

1177

1178

Tables

Table 1: Shortlist of RNA chaperone candidates based on multiple filter criteria (shown in Figure 7). The locus tag is given,

followed by cluster no. to which the protein was assigned, the fractions in which the abundance peaked (Max) and the fractions in

which peaks occurred. The following columns give the predicted number of amino acids (AA), the relative conservation and SVM

scores, and finally comments and annotation. All listed proteins were assigned to the phylogenetic group "Cyanoabacteria".

Locus tag	Cluster	Max	Fractions	AA	Relative conservation	SVM score	Comments and annotation
sll1424	10	6	6, 10	491	1	0.78	Putative SMC (structural maintenance of chromosomes) domain
sll1515/gifB	10	6	6	149	0.71	1.97	glutamine synthetase inactivating factor IF17
slr0169	10	7	7	213	0.94	0.99	CsoD1 ortholog (part of β-Carboxysome
slr0211	10	7	4, 7	403	0.99	0.83	acyl-CoA N-acyltransferase domain
slr1081	10	7	4, 7	210	0.75	0.56	DUF820 featuring a restriction endonuclease domain
ssl2064	10	7	4, 7	75	0.64	2.22	DUF4160
ssl2874	10	6	4, 6, 11	89	0.9	2.42	RemA homolog; KEGG K09777
sll0319	11	8	3, 8	297	0.57	1.55	DUF3747, periplasmic protein
slr0287	12	9	5, 9	118	0.9	1.02	KH domain-containing protein; KEGG K06960
sll1939	14	9	3, 9, 11, 13	214	0.89	0.53	FtsZ interacting Ftn6 homolog
sll0284	15	11	11	579	0.93	0.64	DUF87 containing an N-terminal HAS-barrel domain; KEGG K06915
sll0639	15	12	7, 12	417	0.74	0.63	DHH phosphoesterase domain-
ssr1238, slr0743a	15	11	11		0.71	1.04	YlxR homologue
slr1660	17	18	18	192	0.91	1.41	DUF3172 with N-terminal transmembrane domain

Figures

1185

Figure 1. Fractionation of cyanobacterial proteins and RNAs by Grad-Seq
analysis. (A) Top part: Gradient tube following centrifugation. The different colors
match known pigment-protein complexes, containing yellow-orange carotenoids, light
blue phycobilins, green chlorophyll and white material at the bottom of the tube. The
different sucrose concentrations are given in % (m/v). One representative tube out of
three is shown. Lower part: Separation of RNA in a 10% urea-PAA gel. The position
of abundant RNA species is given for orientation. The low-range ssRNA ladder (NEB)
served as a size marker (numbers to the left in nt). **(B)** Top part: Fractions after
elution into collection tubes. Based on the low sample complexity of fractions 14 to 17
determined in pre-experiments, fractions 14 and 15 as well as 16 and 17 were pooled
before preparing samples for mass spectrometry and RNA sequencing. Lower part:
Separation of proteins in a 15% SDS-PAA gel. The positions of abundant
phycobiliproteins (phycocyanin and allophycocyanin), small and large RuBisCO
subunits (RbcS and RbcL) and ribosomal proteins are given for orientation. The
PageRuler protein ladder (Thermo Fisher) served as a molecular mass marker
(numbers to the left in kDa). **(C)** Characteristic sedimentation profiles of the entire set
of proteins and of major RNA classes.

Figure 2. Composition of cluster profiles, functional and phylogenetic analysis.
(A) Profiles of clusters 1 to 17 (numbers to the right) along the different fractions 1 to 18 (numbers at the X axis). The grayscaled clusters 1-9 represent the protein-dominated branch, while the colored clusters 10-17 represent the RNA-dominated branch (see **Supplementary Figure S2**). The red cluster 10 represents the main RNA cluster, the yellow clusters 11-12 represent the small clusters between the RNA clusters, the green clusters 13-15 represent the other RNA clusters containing transposase-associated RNAs and housekeeping ncRNAs, such as 6S RNA, tmRNA or RNase P, while the purple clusters 16-17 represent the main ribosomal RNA clusters. **(B)** Pie charts illustrating the relative content in proteins and different RNA species of each cluster. The circle sizes correlate with the respective number of

different elements in them (scale in lower left corner). **(C)** Pie charts illustrating the association of proteins in each cluster with a functional category (Kanehisa and Goto, 2000; Kanehisa, 2019; Kanehisa et al., 2019). Proteins not included in the KEGG database or annotated as "Function unknown" or "General function prediction only" were manually annotated as "Function unknown". The phylogenetic distribution of likely orthologs was determined based on the domclust algorithm of the Microbial Genome Database (Uchiyama et al., 2019) and 58 selected genomes (**Supplementary Table S2**). As some examples, the RNAP subunits, most ribosomal and photosystem subunits or the RubisCO cluster together based on their sedimentation profiles and group together based on their phylogenetic occurrence. Largely uncharacterized proteins assigned to the RNA-dominated branch and not conserved in *Arabidopsis*, *Salmonella* and *E. coli* seem promising candidates as potential cyanobacterial RNA chaperones (highlighted by the dashed rectangle). The circle sizes correlate with the respective number of different proteins in them (scale to the right lower end). The distribution of homologs in the selected reference organisms is indicated by red (absence) or green (presence) rectangles at the bottom of the panel.

1233

Figure 3. RNA sedimentation profiles. (A) Detailed sedimentation profiles of asRNAs (blue), mRNAs (pink) and ncRNAs (red) of cluster 10 by their peaks yield three subgroups. Transcripts with main peaks in (1) fractions 5 and 7, (2) fraction 6 or (3) fraction 7. **(B)** Detailed sedimentation profiles of transposase mRNAs (green) mRNAs (pink) and ncRNAs (red) of clusters 13 and 14 by their peaks yield five subgroups. Transcripts with secondary peaks in (1) fraction 4, (2) fraction 5, (3) fraction 4 and 6, (4) fraction 4 and 7 or (5) no secondary peaks in those fractions. The peaks relevant for the subgrouping are indicated with a red arrow, and the other peaks with a black arrow.

1240

Figure 4. In-gradient distribution of ribonucleoprotein complexes of *Synechocystis*. The heatmap illustrates the standardized relative abundances (z-score) of each detected protein and RNA in different shades of gray (proteins) or red (RNA) alongside the fractions of the sucrose density gradient. Cluster affiliations are shown on the right (cluster color code as in **Figure 2** and **Supplementary Figure S2**). **(A)** Ribosomal RNAs and proteins. The fractions of cooccurring 16S rRNA and 30S ribosomal proteins as well as 5S, 23S rRNA and 50S ribosomal proteins are edged by a red box. The names of ribosome-associated proteins with diverging gradient profiles are indicated in red, such as Ycf65, Rps1b, Rps1A, Rps2, Rps10 (30S subunit), Rpl12, Rpl11 and Ssl0438 (50S subunit). **(B)** CRISPR RNAs and associated proteins. The fractions of respective cRNAs and cooccurring CRISPR

proteins are edged by red boxes. **(C)** RNAP subunits, sigma factors and 6S RNA, essential housekeeping RNAs and their protein binding partners, as well as selected proteins involved in RNA metabolism. The fractions of cooccurring RNAP subunits, sigma factors and 6S RNA, as well as the tmRNA, RNase P RNA and signal recognition particle (SRP) RNA cooccurring with their respective cognate binding proteins, are edged by red boxes. Sigma factors and proteins involved in RNA metabolism with peaks above fraction 5 are marked in red.

Figure 5. Heatmap showing the in-gradient distribution of the subunits of the major protein complexes involved in photosynthesis. (A) The sucrose density gradient [w / v 10 % - 40 %] showed a characteristic color pattern (left) corresponding to the pigments bound to distinct macromolecular complexes detected in the same sedimentation range (grayscale heatmap). The in gradient distribution of detected photosystem subunits and phycobiliproteins is given in the heatmap with standardized relative abundances (z-score). The profiles of CpcG1 and CpcG2 are shown separately to illustrate the difference between the G-type PBS (CpcG1 as rod-core linker) and L-type PBS (CpcG2 as rod-core linker) types (Kondo et al., 2005; Liu et al., 2019). **(B)** Absorption spectra taken from the indicated pigmented fractions. The dashed lines indicate the average absorption spectrum of all pigmented fractions. Complex subunits are abbreviated with the last letter of the gene name. Subunits of PSII, e.g., psb**A**, of the cytochrome-b$_6$f complex, e.g., pet**A**, subunits of PSI, e.g., psa**A**, subunits of phycobilisomes, e.g., phycocyanins cpc**A** or allophycocyanins apc**A**. Green coloring indicates membership to the core complex, while green-yellow coloring refers to subcomplexes, accessorial parts of the complex or assembly factors, showing gradient profiles differing from the corresponding core complex.

1273

1274

1275

Figure 6. In-gradient distribution of protein-protein complexes consisting of few interacting proteins. The heatmap illustrates the standardized relative abundances (z-score), giving the in gradient distribution of **(A)** all detected TA systems and **(B)** proteins interacting with the nitrogen regulatory proteins P_{II} and PipX.

1281

1282

Figure 7. Sedimentation zones functionally linked to gene expression. The scheme illustrates the sedimentation profiles of intact macromolecular protein complexes (left) alongside the sucrose density gradient, matching the observed pigment pattern. The overall protein and RNA distribution along the gradient is depicted on the right of the gradient scheme. The majority of proteins (~80 %) occur in the lower molecular fractions (below F6, clusters 1-9), indicating their association with small complexes or occurrence as soluble proteins. Proteins that fall into that group and are involved in gene expression are the majority of transcription factors and two component systems, anti-sigma factors, the sigma factors SigG and SigF, several proteins involved in RNA degradation and tRNA

metabolism. The minority of proteins (~20 %) occur in the higher molecular fractions (above F5, cluster 10-17), indicating their association with large complexes. Proteins that fall into that group and are involved in protein biosynthesis are the RNA polymerase (RNAP), the sigma factors SigC, SigB, SigE and SigA, several proteins and RNA-protein complexes involved in RNA metabolism, such as RNase D, RNase J, RNase E, RNase P, PNPase, Enolase, CrhR, 6S RNA, tmRNA, the signal recognition particle (SRP) and the ribosomes. A large portion of these proteins and RNA-protein complexes cooccur in fractions 8-11. Most detectable RNAs occur in fractions 8-11 as well, indicating their likely direct association with complexes composed of proteins involved in transcription, RNA degradation, RNA processing or translation initiation. Notably, most detected RNAs fall into two groups. Group one comprises those that sediment in high abundance in fractions 6-7 (cluster 10), potentially forming stable interactions with RNA chaperones. Group two includes those RNAs that generally sediment in lower abundances in fraction 4 (cluster 13-14), potentially not bound to any RNA chaperone and thus are not as stable when not directly associated in a larger complex. tRNAs are the exception, mainly sedimenting in high abundance in fraction 5 (cluster 8) together with the majority of tRNA synthetases and likely not bound to a larger complex, while rRNAs seem to be solely attached to the ribosomes.