Original Article

Landscape and climatic features drive genetic differentiation processes in a South American coastal plant

GUSTAVO A. SILVA-ARIAS[1,2], LINA CABALLERO-VILLALOBOS[1], GIOVANNA C. GIUDICELLI[1], LORETA B. FREITAS[1*]

[1]Laboratory of Molecular Evolution, Department of Genetics, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil.

[2] Section of Population Genetics, Center of Life and Food Sciences Weihenstephan, Technische Universität München, Liesel-Beckmann-Straße 2 85354 Freising, Germany.

[*]For correspondence. E-mail: loreta.freitas@ufrgs.br

Genetic differentiation in South America´s coastline

ABSTRACT

Background and aims Historical and ecological processes shaped the patterns of genetic

diversity in plant species; among these, colonization to new environments such as coastal

regions generate multiple signals of interest to understand the influence of landscape features

on the population differentiation.

Methods We analysed the genetic diversity and population structure of *Calibrachoa*

heterophylla to infer the influence of abiotic landscape features on this coastal species' gene

flow in the South Atlantic Coastal Plain (SACP). We used ten microsatellite loci to genotype

253 individuals from 15 populations, covering the species' entire geographical range. We

applied population genetics analyses to determine population diversity and structure along the

SACP, migration model inference and correlative analyses to disentangle the most likely

drivers of gene flow in the SACP.

Key Results The *C. heterophylla* populations located more distantly from the seashore

showed higher genetic diversity than those closer to the sea. The genetic differentiation had a

consistent signal of isolation-by-distance. Landscape features, such as water bodies and wind

corridors, and raw geographical distances equally explained the genetic differentiation,

whereas the precipitation seasonality showed a strong signal for isolation-by-environment in

marginal populations. The estimated gene flow suggested that marginal populations had

restricted immigration rates, which could enhance the genetic differentiation.

Conclusions The influence of topographical features in population differentiation in *C.*

heterophylla is related with the history of the coastal plain deposition. Gene flow is mainly

restricted to nearby populations and facilitated by wind fields but with no apparent influence

of large water bodies. Furthermore, differential rainfall regimes in marginal populations can

promote local genetic differentiation.

Key words: *Calibrachoa heterophylla; c*oastal species; colonization; gene flow; landscape

genetics; Solanaceae; South Atlantic Coastal Plain.

INTRODUCTION

Coastal areas in South America constitute distinct landscapes with unique biotic composition.

Many different geomorphological, climate, oceanographic features, and colonization events

from the surrounding biomes shaped these areas (Hulton *et al.,* 2002; Scarano, 2002; Behling,

2003; Carnaval and Moritz, 2008; Saillard *et al.,* 2009; Miloslavich *et al.,* 2011). Therefore,

South American coastal flora shows a peculiar diversity with a range of biogeographical

processes involving different population demographic processes (Silva *et al.,* 2018; Massante

and Gerhold, 2020). Although the studies on plant diversification in South America have

received increased attention in the last years, analyses focusing the post-glacial re-

colonization, speciation, migration, and colonization of coastal areas are still scarce (Sérsic *et*

al., 2011; Turchetto-Zolet *et al.,* 2013; Leal *et al.,* 2016).

The species' geographical distribution and genetic diversity result from historical and

contemporary processes acting together with ecological factors (Loveless and Hamrick, 1984;

Huang *et al.,* 2016; Schierenbeck, 2017). The multiple environmental particularities in the

coastal areas constitute exciting models for studying genetic differentiation in response to

climate changes, physical barriers, and ecological features (Kadereit and Westberg, 2007;

Escudero *et al.,* 2010; Sork, 2016). Coastal regions have common characteristics, such as

intrinsic linear distributions, high salinity, wind strength, and tidal influence, that investigate

convergent demographic patterns among species from different areas (Escudero *et al.,* 2010).

The colonization and the genetic isolation are critical events in the evolutionary dynamics of

coastal plant populations (Thompson, 1999), mainly because the spread to new environments

generates signals on the genetic diversity and structure of species (Excoffier *et al.,* 2009).

The geographical distance and environmental differences influence the genetic

differentiation across species range because they affect the genetic variation and structure

between populations (Manel *et al.,* 2003; Nosil *et al.,* 2009; Lee and Mitchell-Olds, 2011).

Analyses such as the isolation-by-distance (IBD) and isolation-by-environment (IBE) can

identify the causes of the genetic differentiation resulting from geographic distance and

interactions between organisms and their environments, respectively (Orsini *et al.,* 2013;

Sexton *et al.,* 2014; Wang and Bradburd, 2014).

Some studies have indicated that the differences in micro-environments or resulting from

abiotic factors (temperature and superficial marine currents) may promote differential

selection that limits the establishment of species (e.g., Tellier *et al.,* 2009, 2011; Mori *et al.,*

2015; Francisco *et al.,* 2018). Despite that, the evaluation of the relative influence of space

and environment on the genetic differentiation in South American coastal plains is still scarce

(Baranzelli *et al.,* 2014; Silva-Arias *et al.,* 2017; Meireles and Manos, 2018).

The South Atlantic coastal Plain (SACP) is a flat, continuous, and open region

constituting the most extensive coastal region in South America. The SACP extends NE-SW

for approximately 600 km, occupied mostly by large coastal lakes, and crossed by two

perennial water channels (Tomazelli *et al.,* 2000; Weschenfelder *et al.,* 2010). The SACP

gradually arose during sea-level transgressions and regressions processes caused by glacial-

interglacial cycles during the last 400 thousand years. The most substantial transgression and

regression cycles let the formation of four main sand barriers to be positioned parallel to the

coastline (barrier-lagoon systems I to IV; Tomazelli *et al.*, 2000; Tomazelli and Dillenburg,

2007). The harsh environment, such as strong spring-summer sea breezes from the northeast

and high insolation, strongly influences this region such as strong spring-summer sea breezes

from the northeast and high insolation (Dillenburg *et al.,* 2009), which was responsible for the

current topography and, consequently, influenced the distribution and variability of genetic

lineages in contemporary plants (e.g., Mäder *et al.,* 2013; Ramos-Fregonezi *et al.,* 2015;

Silva-Arias *et al.,* 2017).

In this study, we aimed to investigate the historical and contemporary processes involved

in the diversification of coastal plants in South America. We examined potential

topographical and climatic predictors for population structure and gene flow during the

colonization of the SACP based on a small shrub, perennial, and coastal nightshade species,

Calibrachoa heterophylla. Our results provided information that can support the

establishment of general scenarios describing evolutionary processes for plants from the

coastal regions in South America and addressing conservation gaps in the face of climate

changes.

86   **MATERIALS AND METHODS**

Study system

The species of *Calibrachoa* (Solanaceae) occur in subtropical and temperate grasslands in

southern Brazil, northeast Argentina, and Uruguay. The genus encompasses ca. 30 species,

among which *C. heterophylla* is the only species that colonized coastal environments (Mäder

and Freitas, 2019). This species is diploid (2n = 18), semi-prostrated, and displays purplish

bee-pollinated flowers; the fruits are capsules and produce dozens of tiny seeds (< 1.4 mm)

with no dispersal mechanisms. The species occupies open sandy grasslands, dunes, or rocky

outcrops in lakeside or marine environments from ~ 28 Lat S to 32 Lat S in the SACP (Mäder

et al., 2013). Longitudinally, populations of *C. heterophylla* occur from the seashore to less

than 90 km from the coast, with the populations separated from the sea by big lagoons. Just

one disjointed and small group of populations (Fig. 1A) can be found outside from SACP,

restricted to the sandbanks alongside the Santa Maria River basin, ~ 55 Long W.

The phylogeographic structure of *C. heterophylla* reveals one inland and three coastal

intra-specific plastid DNA (cpDNA) lineages that likely resulted from divergence before the

SACP formation. Two river channels acted as paleo-barriers, splitting the coastal lineages

(Mäder *et al.,* 2013). The current distribution of *C. heterophylla* in SACP could be shaped

through a population expansion following the last marine regression (Mäder and Freitas,

2019), ca. 7-8 thousand years ago (kya; Tomazzeli *et al.,* 2000).

Sample collections

We sampled a total of 253 individuals from 15 locations (hereafter called populations; Fig.

1A) that covered the entire *C. heterophylla* known distribution. We collected leaves of all

individuals found in each locality and preserved them in silica gel. The number of individuals

per population varied from three to 41 (Table 1).

Laboratory procedures

The total DNA was extracted following a CTAB-based protocol (Roy *et al.,*1992) and

amplified for ten anonymous microsatellite loci (Che18, Che59, Che119, Che26, Che34,

Che81, Che82, Che85, Che72, and Che126) developed for *C. heterophylla*, following

standard protocols for PCR and genotyping procedures (Silva-Arias *et al.,* 2015).

Characterization of the genetic diversity

We performed tests for linkage disequilibrium and deviations from Hardy-Weinberg

equilibrium (HWE) within each population for each locus. We assessed the significance of

HWE deviations using 10^6 Markov chain steps and Fisher's exact probability tests in

ARLEQUIN 3.5 (Excoffier and Lischer, 2010). We estimated the genetic diversity based on

average rarefied allelic richness, private alleles, observed heterozygosity (H_O), expected

heterozygosity (H_E), the Garza-Williamson (G-W) index, and inbreeding coefficient (F_{IS}; with

confidence limits from 1000 bootstrap resampling over loci) using the POPPR 2.8.5 (Kamvar

et al., 2014, 2015) and HIERFSTAT 0.04-22 (Goudet, 2005, 2014) in R 3.6.3 package (R Core

Team, 2019), and ARLEQUIN.

Genetic structure

We assessed the genetic structure employing two model-based clustering methods and two

exploratory data analyses (François and Waits, 2016). The model-based clustering methods

used were STRUCTURE 2.3.4 (Pritchard *et al.,* 2000) and the spatial Bayesian clustering

program TESS 2.3. These analyses provide estimates for the number of genetic clusters (K) in

HWE equilibrium, individual assignment probabilities, and compute the proportion of the

genome of each individual that can be assigned to the inferred clusters.

For STRUCTURE analysis, the number of clusters evaluated ranged from 1 to the total

number of populations (15), with ten independent runs per K-value. We performed each run

using 2.5×10^5 burn-in periods and 1.0×10^6 Markov chain Monte Carlo repetitions after the

burn-in, under an admixture model, assuming correlated allele frequencies (Falush *et al.,*

2003), and including a priori sampling locations as prior (*locprior*) to detect weak population

structure. The *locprior* option is not biased toward detecting structure when it is not present

and can improve the STRUCTURE results when implemented with few loci (Hubisz *et al.,*

2009). To obtain the K value that better explains the structure based on the genetic dataset, we

assessed the measures of the ΔK method (Evanno *et al.,* 2005) that is useful to recover the

highest level of genetic structure.

TESS implements a spatial assignment approach to group individuals into clusters

accounting for samples' geographical locations, giving them higher probabilities of belonging

to the same genetic cluster to those that are spatially closer in the connection network. For

TESS, we ran100 000 generations, with 50 000 generations as the burn-in, using the

conditional autoregressive (CAR) admixture model, and starting from a neighbour-joining

tree. We ran 20 iterations for each value of maxK ranging from 2 to 15. We added small

perturbations to the original spatial with a standard deviation equal to 0.2 to obtain single

different coordinates for each individual, because of the spatial proximity among individuals

from each collection site allowed to obtain only one coordinate. We assessed the convergence

inspecting the post-run log-likelihood plots and obtained the support for alternative K values

inspecting the statistical measure of the model prediction capability from deviance

information criterion (DIC; Spiegelhalter *et al.*, 2002). We computed and plotted the average

of DIC values to detect maxK value at the beginning of a plateau. Replicated runs of best K

results for both STRUCTURE and TESS were summarized and plotted with the POPHELPER

(Francis, 2017) R package.

　　Additionally, to detect genetic structure, we implemented two exploratory data analyses

multivariate method (François and Waits, 2016), the Discriminant Analysis of Principal

Components (DAPC; Jombart *et al.*, 2010) and the spatial Principal Component Analysis

(sPCA; Jombart *et al.,* 2008), both using the ADEGENET 2.1.3 (Jombart, 2008) R package. For

the DAPC analysis, the SSR data were first transformed using Principal Component Analysis

and keeping all principal components (PCs). After that, we implemented the function

find.clusters to obtain the optimal number of clusters that maximizes the between-group

variability using the lowest Bayesian Information Criterion (BIC) score. To avoid overfitting,

we set an optimal reduced number of PCs to retain given the best number of clusters using the

function *optim.a.score*. Finally, the DAPC was implemented with the number of clusters and

PCs to the optimal values and plotted a scatter plot of the two first components with the

function *s.class*.

　　sPCA of genetic structure incorporates spatial information to maximize the product of

spatial autocorrelation (Moran's I) and the variance for each eigenvector, which produces

orthogonal axes that describe spatial patterns of genetic variation. The spatial information is

included in the analysis using a spatial weighting matrix derived from a connection network.

To test the effect of the neighbouring definition on the results, we ran the sPCA using six

different connection networks available in the function *chooseCN*. For these analyses, we

used the same perturbed coordinates used in TESS analysis. Monte Carlo simulations (global

and local tests) were used with 10 000 permutations to test for non-random spatial association

of population allele frequencies for all sPCA implemented.

Historical and contemporary gene flow estimations

Contemporary asymmetric migration rates were estimated using a Bayesian approach

implemented in BAYESASS 3.0 (Wilson and Rannala, 2003). We ran 10^8 iterations and a

burn-in of 10^7. We adjusted the mixing of allele frequencies, inbreeding coefficients, and

migration rates parameters to 0.6, 0.6, and 0.3, respectively, to obtain acceptance rates around

40%. We assessed convergence examining the log-probability plots and the effective sample

sizes for each run using TRACER 1.6 (Rambaut *et al.,* 2014) and looking for consistency of the

migration estimates among three independent runs with different initial seed numbers.

We assessed historical gene flow testing the support of four alternative scenarios given

our genetic dataset using Bayes factors calculated from the marginal likelihood

approximations (Beerli and Palczewski, 2010). We used the coalescent-based MIGRATE-N

3.2.6 (Beerli and Felsenstein, 2001) software to estimate the mutation-scaled population size

(Θ) and a mutation-scaled migration (M) parameter. To assess the support of each migration

model, we used the Bézier log-marginal likelihood approximations.

For all models, we pooled the populations into four groups according to the geographical

distribution and genetic structure (see Results Figs. 1A and 2). The 'Inland' group included

the São Francisco de Assis, Cacequi 1, and Cacequi 2 populations; 'West' group

encompassed the Arambaré, Barra do Ribeiro, and Pelotas populations; 'North' included the

Santo Antônio da Patrulha, Torres, and Laguna populations; and 'South' group clustered the

Mostardas 1, Mostardas 2, São José do Norte 1, São José do Norte 2, Rio Grande, and Santa

Vitória do Palmar populations.

We evaluated four migration models: (1) source-sink from inland with unidirectional

migration from 'Inland' group to the remaining groups; (2) source-sink from the west with

unidirectional migration from 'West' group to the remaining groups; (3) step-stone from

inland with unidirectional migration from Inland to West and from West to North and South;

and (4) step-stone from the coast with unidirectional migration from North to West, from

South to West, and from West to Inland (Supplementary Data Fig. S1).

We ran the MIGRATE-N Bayesian inference in the Cipres Science Gateway 3.3 (Miller *et*

al., 2010), with one long chain of 5×10^6 steps, sampling at every 100[th] increment, and a

burn-in of 3×10^4 steps. We used uniform priors and slice sampling for both Θ and M ranging

from 0 to 20 (mean = 10, delta = 0.5). We used a heating scheme (Metropolis-coupled

Markov Chain Monte Carlo) with four parallel chains and temperatures of 1, 1.5, 3, and 10^6.

Space, topography, environment, and genetic differentiation

Spatial correlation patterns under isolation-by-distance (IBD) arouse bias in several genetic

structure tests (Frantz *et al.,* 2009; Meirmans, 2012). Therefore, we assessed the IBD through

linear regression of linearized pairwise F_{ST} genetic distances and log-transformed

geographical distances (Rousset, 1997) using a Mantel test, assessing the significance with 10

000 randomizations in VEGAN 2.5-6 (Oksanen *et al.,* 2015) R package.

We calculated the pairwise F_{ST} (Weir and Cockerham, 1984) matrix with the HIERFSTAT

package. We obtained the geographical inter-population distance matrix calculating the linear

Euclidean distance between X and Y UTM 22S (reference EPSG: 32722) populations'

coordinates transformed from Long/Lat coordinates with RGDAL 1.0-4 (Bivand *et al.,* 2015) R

package.

We tested isolation-by-environment (IBE) models to examine whether differences in

climatic conditions explain the observed inter-population genetic differentiation in *C.*

heterophylla. We calculated the climatic dissimilarity matrices between each populations' pair

for the following bioclimatic variables: total annual precipitation, total annual days with rain,

precipitation seasonality, mean annual temperature, mean summer maximum temperature,

mean winter minimum temperature, mean temperature range, and temperature seasonality.

We took the values of those climatic variables from raster layers specifically developed for

the SACP derived from a high-density sampling of climate stations throughout the region,

geostatistical modelling, and spatial interpolation (Silva-Arias *et al.,* 2017).

Considering that strong summer winds in the SACP can be an important vector for seed

or vegetative propagules movement, we also included a wind connectivity matrix in the IBE

tests. We calculated surface wind direction and speed data for southern Hemisphere's spring

months (September to November) 2011–2016 sampled every three hours. We downloaded the

data from the Global Forecasting System using the RWIND 1.1.5 (Fernández-López and

Schliep, 2019) R package. For each sampled time, we transformed direction and speed values

into raster layers using the *wind2raster* function to obtain transition layers using the function

flow.dispersion. Finally, we calculated pairwise cost distance matrices with the function

costDistance in GDISTANCE 1.3-1 (van Etten, 2017) R package. We then averaged the matrices

for all-time series. We plotted the final matrix with the QGRAPH 1.6.5 (Epskamp *et al.,* 2012)

in the R package.

To test for possible models of inter-population differentiation linked to landscape

discontinuities alongside the SACP, we extended the IBD and IBE analyses using raster grids.

We outlined two coast distance models (Supplementary Data Fig. S2): (1) the *continuous* (or

null) model wherein no landscape discontinuity affects the inter-population connectivity. We

created a raster grid with all cells values equal to 1, including all cells on freshwater surfaces.

This model is expected to resemble a Euclidean geographical distance, but it is more proper

for comparisons with models based on circuit theory; and (2) the *water bodies* model, which

proposes that the widespread freshwater bodies in the SACP restrict the connectivity between

populations. For that, we created a raster grid with all land cells values equal to 1, and cells

within freshwater surfaces as complete barriers (*no data*). We generated pairwise cost

distance matrices using the function *transition* in GDISTANCE package considering an eight-

neighbour cell connection scheme, Long/Lat coordinates per population as nodes, and raster

resolution of 0.09 degrees (~ 10 km).

We examined the relationships between genetic differentiation (F_{ST}) and geographical or

topographical distances (IBD) and environmental dissimilarity (IBE) using multiple matrix

regressions with randomization (MMRR; Wang, 2013) implemented in R.

264   **RESULTS**

Genetic diversity

We found a total of 140 alleles across the ten microsatellite loci. The mean number of alleles

per locus was 14, ranging from seven (Che59) to 17 (Che81). All loci showed higher H_e

(Supplementary Data Fig. S3) with 25% of the locus-population combinations showing a

departure of HWE (P < 0.05). We detected a significant linkage disequilibrium signal (P <

0.01) for several loci pairs, but as the linkage pattern was not consistent across populations for

any loci pair, we assumed linkage equilibrium and maintained all loci in the analyses.

MICROCHECKER analysis did not show null alleles, scoring errors, or stutter peaks for any

locus.

Populations located outside of SACP and those collected at the west side of the Patos

Lagoon showed higher genetic diversity overall loci (Fig. 1B). In contrast, the coastal

populations located at the northern and southern edges of species distribution in SACP

(Laguna and Santa Vitória do Palmar populations, respectively) had lower genetic diversity

values. Average values across loci for H_o ranged from 0.72 (São Francisco de Assis) to 0.31

(Laguna) and for H_e from 0.74 (Cacequi 2) to 0.48 (São José do Norte 1). We found 22% of

the alleles being exclusive to one population, with Barra do Ribeiro showing the highest

number of private alleles (eight), whereas Pelotas, Mostardas 1, São José do Norte 1, and São

José do Norte 2 populations had no private alleles. Garza-Williamson values ranged from

0.39 (Cacequi 1) to 0.83 (Torres). We found positive and significant inbreeding coefficients

(F_{IS}) for five populations (Fig. 1B), all of them located at the borders (northern and southern)

of species' distribution in the SACP (Fig. 1A; Table 1).

Genetic structure

The best K value inferred from the ΔK method in STRUCTURE analysis was K = 4

(Supplementary Data Fig. S4A), whereas the curve of average DIC values obtained with

TESS analyses showed a plateau after maxK = 8 and maxK = 2, 3, and 4 runs had the lowest

standard deviations (Supplementary Data Fig. S4B). Accounting for these results, we

analysed the bar plots of membership probability from K = 2 to K = 8.

The recovered population structure had a strong geographic signal. Genetic clustering

obtained with STRUCTURE (Fig. 2) and TESS (Supplementary Data Fig. S5) showed similar

results. For both approaches, assignment probabilities obtained for K = 2 delimited one

cluster composed of populations located at northern SACP and a second group composed by

remain populations. When K = 3, one group encompassed the northern coastal populations,

the second clustered the southern coastal populations, and the third brought together all

populations located on the west side of the Patos Lagoon and the three inland populations.

Higher values of K showed differences between the two Bayesian clustering methods in the

sequence that groups were incorporated; however, both recovered the same clustering pattern

at K = 8. With this K, the first group was composed by the three inland populations; the

second and third groups encompassed solely a single population each (Barra do Ribeiro and

Pelotas, respectively, both located on the west of Patos Lagoon); the forth cluster grouped two

populations from the northern SACP (Santo Antonio da Patrulha and Torres); the fifth cluster

encompassed the individuals from Laguna population (the northernmost distributed

population); the sixth cluster grouped São José do Norte 1 and São José do Norte 2

populations; the seventh cluster enclosed the individuals from the southernmost located

population (Santa Vitória do Palmar); and the eighth group was not preferentially linked to

any population, instead showed low membership probabilities for individuals from several

populations.

We found higher admixed membership for individuals from populations located in

geographical transitional regions (Fig. 2 and Supplementary Data Fig. S5). The most notable

cases were Mostardas 1 and Mostardas 2 populations located between the north and south

SACP portions, with all individuals from these populations showing membership probability

assigned into all clusters except the inland group. Individuals from the Rio Grande population

also showed high mixed membership and achieved some discrepancy between the two

Bayesian assignment tests. For the individuals from the Rio Grande population, STRUCTURE

assigned around 60% of the membership to the Torres genetic component, while TESS

supported a higher membership probability to the Santa Vitória do Palmar component.

Exploratory analyses revealed similar results to the Bayesian clustering analyses.

Although the lowest BIC score was for K = 8 (Supplementary Data Fig. S6), DAPC

scatterplot of the two main discriminant components (Fig. 1C) revealed three main groups,

delimiting northern and southern coastal populations in two different groups, and a third

group encompassing individuals from inland, Patos Lagoon western side, and the two

Mostardas populations. sPCA (Fig. 1D) recovered the two main groups formed by the

northern and southern edge coastal populations on the first sPC axis. The Torres and Santo

Antônio de Patrulha from the northern and Santa Vitória do Palmar from the southern

occupied opposed spaces related to their respective groups on the first sPC axis, while the

remaining populations showed a gradient of differentiation on the second sPC axis.

Migration rates

The mean migration rate, as estimated with BAYESASS, was 0.015. However, the vast

majority of the pairwise population estimations showed relatively wide confidence intervals

ranging between 0 and 0.1, with only four population pairs showing higher posterior effective

migration rates and confidence intervals above zero. Among these four cases, the most

outstanding was Mostardas2 to Arambaré ($N_m \approx 0.08$; 95% CI 0.01 - 0.14) that supports

migration between populations separated by the Patos Lagoon. The other three cases involved

neighbour populations, Cacequi1 to Cacequi2 ($N_m \approx 0.12$; 95% CI 0.05-0.19), Mostardas2 to

Mostardas1 ($N_m \approx 0.07$; 95% CI 0.01-0.13), and SJoséNorte2 to SJoséNorte1 ($N_m \approx 0.16$;

95% CI 0.09-0.22). Migration estimates obtained from independent runs of BAYESASS

showed similar values (Supplementary Data Table. S1).

The model-based coalescent approach implemented in MIGRATE-N supported the *step-*

stone from the coast as the most likely historical migration model between population groups

(Table 2, Supplementary Data Fig. S1C). Parameter estimation taken from the best-supported

model showed that 'Inland' group had the highest mean mutation scaled population size (Θ),

which was around eight times higher than the Θ estimated for 'West' and 'North' groups, and

around of 20 times higher than the Θ estimated for 'South' group that showed the lowest

value (Table 2). Migration from 'West' to 'Inland' had the highest mean mutation scaled

migration rate that was twice higher than the 'North' to 'West' and five times when compared

to the 'South' to 'West' values (Table 2).

Isolation-by-distance, isolation-by-environment, and resistance tests

Measures of population differentiation (Fig. 3A) F_{ST} ranged from 0.01 between Mostardas1

and Mostardas2 populations to 0.54 between Laguna and São José do Norte1 populations.

Linear regressions showed a significant positive relationship between genetic (Fig. 3A) and

linear geographical (Fig. 3B) distances ($R^2 = 0.14$; $P < 0.001$) supporting a IBD pattern for the

data set (Fig. 4A). Mantel test also supported a correlation between the genetic and

geographical distance matrices (Mantel's r = 0.38, P < 0.001).

Analyses testing the relationship among geographical distance, climate variables, and

genetic differentiation based on MMRR approach showed significant association between the

genetic dissimilarity and Euclidean distance ($R^2 = 0.21$, $\beta = 1.7 \times 10^{-7}$, $P < 0.001$) when

regressed with F_{ST} as the unique explanatory variable. MMRR models implemented with each

of the assessed climate variables and Euclidean distance as explanatory variables showed

significant relationship only for precipitation seasonality (Fig. 3C; $R^2 = 0.35$, $P = 0.003$;

$\beta_{precseason} = 7.5 \times 10^{-3}$, $P = 0.05$; $\beta_{Euc} = 1.1 \times 10^{-7}$, $P = 0.02$, respectively). IBD tests based on

topographic models based on coast distances showed that the *continuous* model (i.e.,

landscape matrix with no topographic discontinuities; Supplementary Data Fig. S2A)

explained slightly better the genetic differentiation (F_{ST} ; $R^2 = 0.16$, $\beta = 0.022$, $P = 0.006$) than

the *water bodies* ($R^2 = 0.14$, $\beta = 0.02$, $P = 0.011$) model (i.e., landscape matrix with water

bodies as full barriers to population connectivity; Supplementary Data Fig. S2B). The wind

connectivity matrix (Figs. 4B and 4C) showed a consistent north-to-south asymmetric step-

stone pattern where marginal populations appeared strongly isolated. Moreover, populations

located at the west side of the Patos Lagoon became receptors from coastline populations. The

coast distance wind matrix (Fig. 4A) showed significant correlation with the F_{ST} genetic

distance matrix ($R^2 = 0.19$, $\beta = 0.001$, $P = 0.0037$).

378   **DISCUSSION**

In this study, we analysed patterns of genetic diversity and structure of *Calibrachoa*

heterophylla to infer the influence of topographical and environmental features on the gene

flow during the recent colonization of a coastal plain in South America.

Our results provided consistent evidence for limited and asymmetric gene flow, mainly

limited by the geographical distance. The populations from northern and southern edges of

the species distribution showed negligible historical and contemporary immigration rates

probably related with an isolation-by-environment through the precipitation conditions. We

also found that the more outstanding topographical feature in the South Atlantic Coastal Plain

(the big water bodies) does not seem to constrain *C. heterophylla* populations' gene flow.

Historical and contemporary drivers of genetic structure in Calibrachoa heterophylla

There is a hierarchical pattern of genetic structure related to both historical and contemporary

landscape features. The highest level of population structure showed three well-supported

groups, northern, central, and southern groups. The central group also included the inland

populations. This main clustering pattern mirrors the phylogeographical structure of *C.*

heterophylla based on cpDNA (Mäder *et al.,* 2013). The retention of historical signals of

genetic structure in highly variable markers, such as microsatellites, is expected for studies

involving the entire geographic range of species, which reinforces the importance of

considering the historical patterns for landscape genetic approaches (Anderson *et al.,* 2010).

The populations from the intersection between northern and southern regions, such as

Mostardas 1 and Mostardas 2 populations, or those between west Patos Lagoon and southern

regions, as Pelotas population, had higher admixture values to a secondary gene flow between

previously differentiated intraspecific lineages (Fig 1).

The influence of geographical distances on the genetic structure was evident in several

levels of *C. heterophylla* genetic structure. The northern, southern, and central groups showed

a main latitudinal pattern of structure. At a fine scale, we found that individuals from

peripheral populations conformed independent groups with individuals' membership higher

than 80%, as seen in Santa Vitória do Palmar, inland populations São Francisco de Assis,

Cacequi 1, and Cacequi 2, and Laguna locations (Figs. 1 and 2; Supplementary Data Fig. S3).

Genetic drift due to strong geographical isolation (Whigham *et al.,* 2008) can explain

individuals' assignment to a completely separated group.

We observed few departures from the geographical frame in the genetic structure in

some distant populations for which the assignment tests suggested as sharing partial genetic

identities, such as Laguna and São José do Norte and Torres and Rio Grande population pairs.

In both cases, populations from northern and southern SACP regions were involved. Either

persistent ancestral variation after population divergence or randomly driven processes, such

as the fixation of the same allele during an expansion wave (Excoffier and Ray, 2008) or

homoplasy (Grimaldi and Crouau-Roy, 1997; van Oppen *et al.,* 2000; Schaal and Olsen,

2000) could equally explain that partial genetic affinity between spatially distant populations.

A long-distance gene flow seems an unlikely scenario based on the high F_{ST} between these

populations and the observed low migration rates.

The Pelotas population showed several different results according to the methodology

and structure level inferred. Exploratory and STRUCTURE analyses grouped Pelotas with the

inland populations (Figs. 1 and 2), while TESS suggested that Pelotas would be more related

to the southern group (Supplementary Data Fig. S3). Similarly, a population fine structure

patterns (i.e., K = 5-8) for both STRUCTURE and TESS suggested a higher affinity of Pelotas

population with São José do Norte or forming an independent group with all individuals

assigned at ~ 100% of membership probability (Fig. 2, Supplementary Data Fig. S3). The

geographic position of Pelotas population and environmental features could have drawn this

scenario. Regarding the geographic position, the Pelotas population is located close to the

southern coastal populations, such as the Rio Grande and São José do Norte. However, the

Patos Lagoon separates the Pelotas population from the seashore, which can lead Pelotas to

receive migrants from coastal populations in a more continental environment. Nevertheless,

inter-annual rainfall conditions, as well as the continental-scale periodic climatic fluctuations,

such as El Niño, can also affect the fluvial discharge and the wind responsible for the

salinization and desalination processes in the Patos Lagoon (Möller *et al.,* 1996, 2001; Möller

and Castaing, 1999). This environmental dynamic could continuously change the individuals'

establishment or survival rates, with the coastal or continental gene pools probably leading to

a differential genetic profile.

Patos Lagoon development let to a secondary contact between previously diverged lineages

The populations located close to the Patos Lagoon, Arambaré, Mostardas 1, Mostardas 2, São

José do Norte 1, São José do Norte 2, and Rio Grande (Fig. 1A) showed the highest levels of

genetic admixture (Fig. 2; Supplementary Data Fig. S3) and the lowest F_{ST} values (Fig. 3A)

among the coastal populations. These results could be related to the recent geomorphological

history of the SAPC region. During most of the Quaternary Period, two rivers (Jacuí and

Camaquã rivers), including several channels corresponding to their temporal dynamic delta

systems, crossed the Patos Lagoon (Weschenfelder *et al.,* 2014). After the formation of the

barrier systems III and IV, the two most recent ocean regressive-transgressive events, the

Patos Lagoon was completely close by sands from the inlets (Santos-Fischer *et al.,* 2016),

which could have constituted an obstacle for the establishment of the population in the east

side of the lagoon. In contrast, the northern and southern regions, which correspond to the

well-established barrier systems I and II (cf. Fig. 1 in Tomazelli and Dillenburg, 2007), could

let to the establishment and differentiation of main coastal lineages that, later, spread and

experienced a secondary contact on the east side of the Patos Lagoon generating the current

patterns of genetic admixture in this region. This recent admixture processes can also explain

the lack of private alleles in Pelotas, São José do Norte 1, São José do Norte 2, and Mostardas

1 populations (Fig. 1B).

The east side of Patos Lagoon (seashore side) has the strongest wind influence within

the SACP (Fig. 4B; Martinho *et al.,* 2010) that can increase the seed dispersal alongside the

region generating, in consequence, higher admixture rates. The gene flow estimations among

C. heterophylla populations support an asymmetric migration from coastal to inland locations,

even at great distances crossing the coastal lakes (Figs. 1 and 4). Despite the differences in the

intraspecific divergence history between *C. heterophylla* (Mäder *et al.,* 2013) and the co-

distributed *P. integrifolia* subsp. *depauperata* (Ramos-Fregonezi *et al.,* 2015), both taxa

shared the same pattern of high genetic admixture based on microsatellite data in populations

located at the east of Patos Lagoon (Silva-Arias *et al.,* 2017). This congruent pattern between

species indicates that the current and historical dynamics in the topographical and

environmental conditions in SACP are responsible for the admixture patterns and the genetic

structure in this region.

The precipitation seasonality is different between the northern and southern extremes

of SACP (Fig. 3C), and there is a significant positive relationship between this environment

feature and genetic differentiation. Moreover, the immigration for the SAPC edges falls

within the lowest estimates (Supplementary Data Fig. S5). Altogether, these results point to a

genetic divergence process enhanced by differential rainfall regimes alongside SACP. The

strong correlation between precipitation seasonality and genetic differentiation was also found

for the coastal populations of *Petunia integrifolia* subsp. *depauperata* (Silva-Arias *et al.,*

2017). Ecological differentiation can promote selection against immigrants that leads, in a

genome-wide context, to a reduction of gene flow, reproductive isolation, and enhances the

stochastic effects of genetic drift (Hendry, 2004; Nosil *et al.,* 2005, 2009). For *C.*

heterophylla, the limited migration rates from the central to marginal distribution could

enhance the fixation of locally adaptive alleles in peripheral populations by preventing the

gene swamping (Alleaume-Benharira *et al.,* 2006).

We found that both geomorphological and environmental conditions influenced population

demographic processes in coastal plants, such as differentiation, connectivity, and local

adaptation. The main current population differentiation in *Calibrachoa heterophylla* is

determined by historical processes and the age of the deposition of the coastal plain. In

contrast, population connectivity is mainly determined by geographic distance and wind fields

but seems to be not affected by significant barriers like water bodies. Besides, marginal

populations appear to present local differentiation related to rainfall conditions.

490   **ACKNOWLEDGEMENTS**

The authors thank to G. Mäder for help in fieldwork and J.R Stehmann for plant

identification. G.A.S.-A. and L.B.F. designed the study; G.A.S.-A. performed fieldwork,

laboratory experiments, analysed data, and interpreted the results; G.A.S.-A. and L.C.-V.

discussed the results and wrote the manuscript. G.C.G. performed laboratory genotyping;

L.B.F. supervised the project and provided resources for the research. All authors reviewed

the manuscript. All authors declare no conflict of interest. All data underlying the study are

included in the manuscript or its supplementary materials.

FUNDING

This work was supported by the Conselho Nacional de Desenvolvimento Científico e

Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior

(CAPES), and Programa de Pós-Graduação em Genética e Biologia Molecular da

Universidade Federal do Rio Grande do Sul (PPGBM-UFRGS). G.A.S.-A. was supported by

a scholarship from the Francisco José Caldas Institute for the Development of Science and

Technology (COLCIENCIAS).

LITERATURE CITED

Alleaume-Benharira M, Pen IR, Ronce O. 2006. Geographical patterns of adaptation within

a species' range: interactions between drift and gene flow. *Journal of Evolutionary*

Biology **19**: 203–215.

Anderson CD, Epperson BK, Fortin M-J, *et al*. **2010.** Considering spatial and temporal

scale in landscape-genetic studies of gene flow. *Molecular Ecology* **19**: 3565–3575.

Baranzelli MC, Johnson LA, Cosacov A, Sérsic AN. 2014. Historical and ecological

divergence among populations of *Monttea chilensis* (Plantaginaceae), an endemic

endangered shrub bordering the Atacama Desert, Chile. *Evolutionary Ecology* **28**: 751–

774.

Beerli P, Felsenstein J. 2001. Maximum likelihood estimation of a migration matrix and

effective population sizes in n subpopulations by using a coalescent approach.

Proceedings of the National Academy of Sciences USA **98**: 4563–4568.

Beerli P, Palczewski M. 2010. Unified framework to evaluate panmixia and migration

direction among multiple sampling locations. *Genetics* **185**: 313–326.

Behling H. 2003. Late glacial and Holocene vegetation, climate and fire history inferred from

Lagoa Nova in the southeastern Brazilian lowland. *Vegetation History and*

Archaeobotany **12**: 263–270.

Bivand R, Keitt T, Rowlingson B. 2015. rgdal: Bindings for the 'Geospatial' data

abstraction library. https://cran.r-project.org/package=rgdal.

Carnaval AC, Moritz C. 2008. Historical climate modelling predicts patterns of current

biodiversity in the Brazilian Atlantic forest. *Journal of Biogeography* **35**: 1187–1201.

Dillenburg SR, Barboza EG, Tomazelli LJ, Ayup-Zouain RN, Hesp PA, Clerot LCP.

2009. The Holocene coastal barriers of Rio Grande do Sul. In: *Geology and*

Geomorphology of Holocene Coastal Barriers of Brazil. Lecture Notes in Earth Sciences,

vol. 107. Berlin:Springer, 53-91.

Epskamp S, Cramer AOJ, Waldorp LJ, Schmittmann VD, Borsboom D. 2012. *qgraph*:

network visualization of relationships in psychometric data. *Journal of Statistical*

Software **48**: 1-18.

Escudero M, Vargas P, Arens P, Ouborg NJ, Luceño M. 2010. The east-west-north

colonization history of the Mediterranean and Europe by the coastal plant *Carex extensa*

(Cyperaceae). *Molecular Ecology* **19**: 352–370.

van Etten J. 2017. *R* Package gdistance: distances and routes on geographical grids. *Journal*

of Statistical Software **76**: 1-21.

Evanno G, Regnaut S, Goudet J. 2005. Detecting the number of clusters of individuals

using the software structure: a simulation study. *Molecular Ecology* **14**: 2611–2620.

Excoffier L, Lischer HEL. 2010. *Arlequin* suite ver 3.5: a new series of programs to perform

population genetics analyses under Linux and Windows. *Molecular Ecology Resources*

10: 564–567.

Excoffier L, Ray N. 2008. Surfing during population expansions promotes genetic

revolutions and structuration. *Trends in Ecology & Evolution* **23**: 347–351.

Excoffier L, Foll M, Petit RJ. 2009. Genetic consequences of range expansions. *Annual*

Review of Ecology, Evolution, and Systematics **40**: 481–501.

Falush D, Stephens M, Pritchard JK. 2003. Inference of population structure using

multilocus genotype data: linked loci and correlated allele frequencies. *Genetics* **164**:

1567–1587.

Fernández-López J, Schliep K. 2019. *rWind*: download, edit and include wind data in

ecological and evolutionary analysis. *Ecography* **42**: 804–810.

Francis RM. 2017. POPHELPER: an R package and web app to analyse and visualize

population structure. *Molecular Ecology Resources* **17**: 27–32.

Francisco PM, Mori GM, Alves FM, Tambarussi EV, Souza AP. 2018. Population genetic

structure, introgression, and hybridization in the genus *Rhizophora* along the Brazilian

coast. *Ecology and Evolution* **8**: 3491–3504.

François O, Waits LP. 2016. Clustering and assignment methods in landscape genetics. In

Balkenhol N, Cushman SA, Storfer AT, Waits LP, eds. *Landscape genetics: concepts,*

methods, applications. Chichester: Wiley, 114-128.

Frantz AC, Cellina S, Krier A, Schley L, Burke T. 2009. Using spatial Bayesian methods

to determine the genetic structure of a continuously distributed population: clusters or

isolation by distance? *Journal of Applied Ecology* **46**: 493–505.

Goudet J. 2005. *hierfstat*, a package for R to compute and test hierarchical F-statistics.

Molecular Ecology Notes **5**: 184–186.

Goudet J. 2014. *hierfstat*: estimation and tests of hierarchical F-statistics. R package version

0.04-14. http://CRAN.R-project.org/package=hierfstat.

Grimaldi M-C, Crouau-Roy B. 1997. Microsatellite allelic homoplasy due to variable

flanking sequences. *Journal of Molecular Evolution* **44**: 336–340.

Hendry AP. 2004. Selection against migrants contributes to the rapid evolution of

ecologically dependent reproductive isolation. *Evolutionary Ecology Research* **6**: 1219–

1236.

Hickerson MJ, Carstens BC, Cavender-Bares J, *et al.* **2010.** Phylogeography's past,

present, and future: 10 years after Avise, 2000. *Molecular Phylogenetics and Evolution*

54: 291–301.

Huang W, Zhao H, Zhao X, Li Y, Lian J. 2016. Effects of environmental factors on genetic

diversity of *Caragana microphylla* in Horqin sandy land, northeast China. *Ecology and*

Evolution **6**: 8256–8266.

Hubisz,MJ, Falush D, Stephens M, Pritchard JK. 2009. Inferring weak population

structure with the assistance of sample group information. *Molecular Ecology Resources*

9: 1322–1332.

Hulton NRJ, Purves RS, McCulloch RD, Sugden DE, Bentley MJ. 2002. The Last Glacial

Maximum and deglaciation in southern South America. *Quaternary Science Reviews* **21**:

233–241.

Jombart T. 2008. *adegenet*: a R package for the multivariate analysis of genetic markers.

Bioinformatics **24**: 1403–1405.

Jombart T, Devillard S, Balloux F. 2010. Discriminant analysis of principal components: a

new method for the analysis of genetically structured populations. *BMC Genetics* **11**: 94.

Jombart T, Devillard S, Dufour A-B, Pontier D. 2008. Revealing cryptic spatial patterns in

genetic variability by a new multivariate method. *Heredity* **101**: 92–103.

Kadereit J. Westberg E. 2007. Determinants of phylogeographic structure: a comparative

study of seven coastal flowering plant species across their European range. *Watsonia* **26**:

229–238.

Kamvar ZN, Brooks JC, Grünwald NJ. 2015. Novel R tools for analysis of genome-wide

population genetic data with emphasis on clonality. *Frontiers in Genetics* **6**: 208.

Kamvar ZN, Tabima JF, Grünwald NJ. 2014. *Poppr*: an R package for genetic analysis of

populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* **2**: e281.

Leal BSS, Palma da Silva C, Pinheiro F. 2016. Phylogeographic studies depict the role of

space and time scales of plant speciation in a highly diverse Neotropical region. *Critical*

Reviews in Plant Sciences **35**: 215–230.

Lee C-R, Mitchell-Olds T. 2011. Quantifying effects of environmental and geographical

factors on patterns of genetic differentiation. *Molecular Ecology* **20**: 4631–4642.

Loveless MD, Hamrick JL. 1984. Ecological determinants of genetic structure in plant

populations. *Annual Review of Ecology and Systematics* **15**: 65–95.

Mäder G, Fregonezi JN, Lorenz-Lemke AP, Bonatto SL, Freitas LB. 2013. Geological

and climatic changes in Quaternary shaped the evolutionary history of *Calibrachoa*

heterophylla, an endemic South-Atlantic species of petunia. *BMC Evolutionary Biology*

13: 178.

Mäder G, Freitas LB. 2019. Biogeographical, ecological, and phylogenetic analyses

clarifying the evolutionary history of *Calibrachoa* in South American grasslands.

Molecular Phylogenetics and Evolution **141**: 106614.

Manel S, Schwartz MK, Luikart G, Taberlet P. 2003. Landscape genetics: combining

landscape ecology and population genetics. *Trends in Ecology & Evolution* **18**: 189–197.

Martinho CT, Hesp PA, Dillenburg SR. 2010. Morphological and temporal variations of

transgressive dune fields of the northern and mid-littoral Rio Grande do Sul coast,

Southern Brazil. *Geomorphology* **117**: 14–32.

Massante JC, Gerhold P. 2020. Environment and evolutionary history depict phylogenetic

alpha and beta diversity in the Atlantic coastal white-sand woodlands. *Journal of*

Vegetation Science **31**: 634-645.

Meireles JE, Manos PS. 2018. Pervasive migration across rainforest and sandy coastal plain

Aechmea nudicaulis (Bromeliaceae) populations despite contrasting environmental

conditions. *Molecular Ecology* **27**: 1261–1272.

Meirmans PG. 2012. The trouble with isolation by distance. *Molecular Ecology* **21**: 2839–

2846.

Miller MA, Pfeiffer W, Schwartz T. 2010. Creating the CIPRES Science Gateway for

inference of large phylogenetic trees. In: *2010 Gateway Computing Environments*

Workshop (GCE). New Orleans: IEEE, 1–8.

Miloslavich P, Klein E, Díaz JM, *et al.* 2011. Marine biodiversity in the Atlantic and Pacific

coasts of South America: knowledge and gaps. *PLoS ONE* **6**: e14631.

Möller OO, Castaing P. 1999. Hydrographical characteristics of the estuarine area of Patos

Lagoon (30°S, Brazil). *In* Perillo GME, Piccolo MC, Pino-Quivira M, eds, *Estuaries of*

*South America.*Berlin: Springer, 83–100.

Möller OO, Castaing P, Salomon JC, Lazure P. 2001. The influence of local and non-local

forcing effects on the sub-tidal circulation of Patos Lagoon. *Estuaries* **24**: 297–311.

Möller OO, Lorenzzentti JA, Stech JL,Math MM. 1996. The Patos Lagoon summertime

circulation and dynamics. *Continental Shelf Research* **16**: 335–351.

Mori GM, Zucchi MI, Sampaio I, Souza AP. 2015. Species distribution and introgressive

hybridization of two *Avicennia* species from the Western Hemisphere unveiled by

phylogeographic patterns. *BMC Evolutionary Biology* **15**: 61.

Nosil P, Funk PD, Ortiz-Barrientos D. 2009. Divergent selection and heterogeneous

genomic divergence. *Molecular Ecology* **18**: 375–402.

Nosil P, Vines TH, Funk DJ. 2005. Reproductive isolation caused by natural selection

against immigrants from divergent habitats. *Evolution* **59**: 705–719.

Oksanen J, Blanchet FG, Kindt R, *et al.* 2015. *vegan*: community ecology package. R

package version 2.3-0. http://CRAN.R-project.org/package=vegan.

Orsini L, Vanoverbeke J, Swillen I, Mergeay J, Meester L. 2013. Drivers of population

genetic differentiation in the wild: isolation by dispersal limitation, isolation by

adaptation and isolation by colonization. *Molecular Ecology* **22**: 5983–5999.

Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure using

multilocus genotype data. *Genetics* **155**: 945–959.

R Core Team. 2019. *R*: a language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. http://www.R-project.org/.

Rambaut A, Suchard MA, Xie D, Drummond A. 2014. *Tracer* v1.6.

http://beast.bio.ed.ac.uk/Tracer.

Ramos-Fregonezi AMC, Fregonezi JN, Cybis GB, Fagundes NJR, Bonatto SL, Freitas

LB. 2015. Were sea level changes during the Pleistocene in the South Atlantic Coastal

Plain a driver of speciation in *Petunia* (Solanaceae)? *BMC Evolutionary Biology* **15**: 92.

Rissler LJ. 2016. Union of phylogeography and landscape genetics. *Proceedings of the*

National Academy of Sciences USA **113**: 8079–8086.

Rousset F. 1997. Genetic differentiation and estimation of gene flow from F-statistics under

isolation by distance. *Genetics* **145**: 1219–1228.

Roy A, Frascaria N, MacKay J, Bousquet J. 1992. Segregating random amplified

polymorphic DNAs (RAPDs) in *Betula alleghaniensis*. *Theoretical and Applied Genetics*

85: 173–180.

Saillard M, Hall SR, Audin L, *et al.* 2009. Non-steady long-term uplift rates and Pleistocene

marine terrace development along the Andean margin of Chile (31°S) inferred from 10Be

dating. *Earth and Planetary Science Letters* **277**: 50–63.

Santos-Fischer CB, Corrêa ICS, Weschenfelder J, Torgan LC, Stone JR. 2016.

Paleoenvironmental insights into the Quaternary evolution of the southern Brazilian coast

based on fossil and modern diatom assemblages. *Palaeogeography, Palaeoclimatology,*

Palaeoecology **446**: 108–124.

Scarano FR. 2002. Structure, function and floristic relationships of plant communities in

stressful habitats marginal to the Brazilian Atlantic Rainforest. *Annals of Botany* **90**: 517–

524.

Schaal BA, Olsen KM. 2000. Gene genealogies and population variation in plants.

Proceedings of the National Academy of Sciences USA **97**: 7024–7029.

Schierenbeck KA. 2017. Population-level genetic variation and climate change in a

biodiversity hotspot. *Annals of Botany* **119**: 215–228.

Sérsic, AN, Cosacov A, Cocucci AA, *et al*. 2011. Emerging phylogeographical patterns of

plants and terrestrial vertebrates from Patagonia. *Biological Journal of the Linnean*

Society **103**: 475–494.

Sexton JP, Hangartner SB, Hoffmann AA. 2014. Genetic isolation by environment or

distance: which pattern of gene flow is most common? *Evolution* **68**: 1–15.

Silva GAR, Antonelli A, Lendel A, Moraes EM, Manfrin MH. 2018. The impact of early

Quaternary climate change on the diversification and population dynamics of a South

American cactus species. *Journal of Biogeography* **45**: 76–88.

Silva-Arias GA, Mäder G, Bonatto SL, Freitas LB. 2015. Novel Microsatellites for

Calibrachoa heterophylla (Solanaceae) endemic to the South Atlantic Coastal Plain of

South America. *Applications in Plant Sciences* **3**: 1500021.

Silva-Arias GA, Reck-Kortmann M, Carstens BC, Hasenack H, Bonatto SL, Freitas LB.

2017. From inland to the coast: spatial and environmental signatures on the genetic

diversity in the colonization of the South Atlantic Coastal Plain. *Perspectives in Plant*

Ecology, Evolution and Systematics **28**: 47–57.

Slatkin M. 1987. Gene flow and the geographic structure of natural populations. *Science* **236**:

787–792.

Sork VL. 2016. Gene flow and natural selection shape spatial patterns of genes in tree

populations: implications for evolutionary processes and applications. *Evolutionary*

Applications **9**: 291–310.

Spiegelhalter DJ, Best NG, Carlin BP, van der Linde A. 2002. Bayesian measures of

model complexity and fit. *Journal of the Royal Statistical Society: Series B (Statistical*

Methodology) **64**: 583–639.

Tellier F, Meynard AP, Correa JA, Faugeron S, Valero M. 2009. Phylogeographic

analyses of the 30° S south-east Pacific biogeographic transition zone establish the

occurrence of a sharp genetic discontinuity in the kelp *Lessonia nigrescens*: vicariance or

parapatry? *Molecular Phylogenetics and Evolution* **53**: 679–693.

Tellier F, Tapia J, Faugeron S, Destombe C, Valero M. 2011. The *Lessonia nigrescens*

species complex (Laminariales, Phaeophyceae) shows strict parapatry and complete

reproductive isolation in a secondary contact zone: reproductive isolation between kelp

species. *Journal of Phycology* **47**: 894–903.

Thompson JD. 1999. Population differentiation in Mediterranean plants: insights into

colonization history and the evolution and conservation of endemic species. *Heredity* **82**:

229–236.

Tomazelli LJ, Dillenburg SR. 2007. Sedimentary facies and stratigraphy of a last

interglacial coastal barrier in south Brazil. *Marine Geology* **244**: 33–45.

Tomazelli LJ, Dillenburg SR, Villwock JA. 2000. Late Quaternary geological history of

Rio Grande do Sul coastal plain, southern Brazil. *Revista Brasileira de Geociências* **30**:

474–476.

Turchetto-Zolet AC, Pinheiro F, Salgueiro F, Palma-Silva C. 2013. Phylogeographical

patterns shed light on evolutionary process in South America. *Molecular Ecology* **22**:

1193–1213.

van Oppen MJH, Rico C, Turner GF, Hewitt GM. 2000. Extensive homoplasy, non-

stepwise mutations, and shared ancestral polymorphism at a complex microsatellite locus

in lake Malawi cichlids. *Molecular Biology and Evolution* **17**: 489–498.

Wang IJ. 2013. Examining the full effects of landscape heterogeneity on spatial genetic

variation: a multiple matrix regression approach for quantifying geographic and

ecological isolation. *Evolution* **67**: 3403–3411.

Wang IJ, Bradburd GS. 2014. Isolation by environment. *Molecular Ecology* **23**: 5649–

5662.

Weir BS, Cockerham CC. 1984. Estimating F-statistics for the analysis of population

structure. *Evolution* **38**: 1358–1370.

Weschenfelder J, Baitelli R, Corrêa ICS, Bortolin EC, Santos CB. 2014. Quaternary

incised valleys in southern Brazil coastal zone. *Journal of South American Earth Sciences*

55: 83–93.

Weschenfelder J, Corrêa ICS, Aliotta S, Baitelli R. 2010. Paleochannels related to Late

Quaternary sea-level changes in Southern Brazil. *Brazilian Journal of Oceanography* **58**:

35–44.

Whigham PA, Dick GC, Spencer HG. 2008. Genetic drift on networks: ploidy and the time

to fixation. *Theoretical Population Biology* **74**: 283–290.

Wilson G, Rannala B. 2003. Bayesian inference of recent migration rates using multilocus

genotypes. *Genetics* **163**: 1177-1191.

 TABLES

745 **Table 1.** Sampling information and inbreeding coefficients for *Calibrachoa heterophylla*

746 populations. Population ID codes follow Fig. 1A.

Pop ID	N	Location	Long	Lat	F_{IS}
1	4	São Francisco de Assis	-55.10077	-29.58307	-0.16
2	10	Cacequi 1	-54.85375	-29.8947	-0.01
3	9	Cacequi 2	-54.90852	-29.85478	0
4	27	Barra do Ribeiro	-51.20255	-30.40754	0.08
5	4	Arambaré	-51.49195	-30.90082	0.12
6	23	Pelotas	-52.16478	-31.70757	0.07
7	10	Laguna	-48.76501	-28.45991	0.34**
8	23	Torres	-49.79809	-29.43227	0.17**
9	37	S Antônio da Patrulha	-50.42936	-29.89291	0.10*
10	3	Mostardas 1	-50.73934	-30.93746	0.35
11	13	Mostardas 2	-50.90112	-31.10909	0.08
12	12	S José do Norte 1	-51.42576	-31.66673	-0.01
13	11	S José do Norte 2	-52.03612	-32.02393	-0.05
14	26	Rio Grande	-52.54661	-32.52396	0.27***
15	41	S Vitória do Palmar	-52.7323	-32.98765	0.25***

748   *P-value < 0.05; ** P-value < 0.01; *** P-value < 0.001

750 **Table 2.** Model support statistics (upper panel) and parameter estimations taken from the

751 best-supported model (lower panel).

Model name	Ln marginal Likelihood	Log Bayes Factor	Model probability
Step stone from coast	-5900.7	0	1.0
Source-sink from west	-5940.5	-79.6	5.1E-18
Source-sink from inland	-5941.5	-81.6	1.9E-18
Step stone from inland	-5944.9	-88.4	6.5E-20

Parameter	Median	Mean	95 % CI
Θ 'Inland'	4.11	4.25	2.55 - 6.28
Θ 'West'	0.55	0.56	0.12 - 0.96
Θ 'North'	0.51	0.52	0.05 - 0.95
Θ 'South'	0.27	0.22	0 - 0.55
M 'West' -> 'Inland'	0.41	0.45	0 - 1.16
M 'North' -> 'West'	0.3	0.26	0 - 0.6
M 'South' -> 'West'	0.19	0.08	0 - 0.41

753 Bayes factor values < 2 indicate a strong preference for the model with the highest probability. Θ = mutation

754 scaled population size; M = mutation scaled migration rate; CI = confidence interval. For graphical model

755 descriptions see Fig. S1.

FIGURES

Figure 1. (A) Geographic locations of *Calibrachoa heterophylla* populations. (B) Graphical

representation of the mean genetic diversity statistics estimated for each population across all

microsatellite loci. (C) Scatterplot of the DAPC analysis. (D) Scatterplot of the sPCA.

Populations' numbers and colours follow each panel.

Figure 2. Bar plots of the individual membership for K = 2-8 genetic clusters as estimated

with STRUCTURE. Populations are separated by white dashed lines and named on the figure

top side of.

Figure 3. Interpopulation genetic, geographic, and environmental dissimilarity matrices. (A)

Genetic differentiation and F_{ST}; (B) Linear geographic distance; (C) Precipitation seasonality

dissimilarity. The values are represented by a colour on a continuous scale. Populations are

numbered as in Fig. 1A and Table 1.

Figure 4. Spring season wind fields in the South Atlantic Coastal Plain and inference of wind

in the population connectivity and gene flow. (A) Correlation between geographical distance

(red) and wind connectivity (blue) coast distance matrices with the F_{ST} genetic distance

matrix; (B) Mean values 2011-2016 of wind speed (blue scale m/s) and direction (arrows) for

the spring season (Sept-Nov); (C) Inter-population wind connectivity network.

Figure S1. Graphical representation of the four coalescent migration models tested in

MIGRATE-N for *Calibrachoa heterophylla*. (A) Source-sink from inland; (B) Source-sink

from the west; (C) Step-stone from inland; (D) Step-stone from the coast.

793

794 **Figure S2**. Graphical representation of the raster layers used to calculate the connectivity

795 values in topographic tests (A) Continuous model; (B) Water bodies model.

Figure S3. The plot of observed *vs*. expected heterozygosity for each locus.

802 **Figure S4**. (A) Plots of the best K estimates for STRUCTURE results. (B) Plots of the mean and

803 standard deviation of the deviance information criterion (DIC) obtained for each maxK

804 assessed with TESS.

Figure S5. Bar plots of the individual membership for each genetic cluster obtained with

TESS. Populations are separated by white dashed lines and names are indicated on the figure

top side.

Figure S6. The plot of Bayesian information criterion (BIC) values obtained for each K

number as assessed with the multivariate method Discriminant Analyses of Principal

Components.

Table S1. Migration estimates obtained with three independent runs of BAYESASS. Populations are numbered as in Fig. 1. The values indicate the estimated posterior mean effective migration rate per generation [the fraction of individuals in population i (rows) that are migrants derived from population j (columns)] and the numbers in parenthesis show the standard deviation. Bold values indicate the diagonal (intra-population estimates) and red values indicate the highest migration estimates (those with above zero 95% confidence intervals).

	1_j	2_j	3_j	4_j	5_j	6_j	7_j	8_j	9_j	10_j	11_j	12_j	13_j	14_j	15_j
1_i	**0.6850** (**0.0180**)	0.0454 (0.0389)	0.0179 (0.0171)	0.0183 (0.0185)	0.0177 (0.0168)	0.0330 (0.0280)	0.0179 (0.0169)	0.0405 (0.0306)	0.0175 (0.0164)	0.0179 (0.0168)	0.0186 (0.0183)	0.0181 (0.0172)	0.0176 (0.0169)	0.0173 (0.0164)	0.0174 (0.0164)
2_i	0.0136 (0.0132)	**0.7980** (**0.0339**)	0.0136 (0.0129)	0.0134 (0.0131)	0.0136 (0.0131)	0.0168 (0.0153)	0.0136 (0.0130)	0.0160 (0.0155)	0.0135 (0.0130)	0.0173 (0.0159)	0.0168 (0.0159)	0.0135 (0.0130)	0.0135 (0.0131)	0.0135 (0.0130)	0.0133 (0.0126)
3_i	0.0144 (0.0139)	0.1189 (0.0356)	**0.6806** (**0.0134**)	0.0140 (0.0134)	0.0211 (0.0185)	0.0140 (0.0133)	0.0140 (0.0133)	0.0232 (0.0199)	0.0144 (0.0137)	0.0139 (0.0131)	0.0146 (0.0142)	0.0153 (0.0145)	0.0138 (0.0132)	0.0140 (0.0132)	0.0138 (0.0132)
4_i	0.0175 (0.0169)	0.0203 (0.0192)	0.0174 (0.0164)	**0.6842** (**0.0165**)	0.0179 (0.0171)	0.0189 (0.0178)	0.0174 (0.0165)	0.0793 (0.0330)	0.0177 (0.0167)	0.0175 (0.0167)	0.0177 (0.0169)	0.0174 (0.0168)	0.0179 (0.0170)	0.0212 (0.0195)	0.0176 (0.0167)
5_i	0.0080 (0.0079)	0.0101 (0.0094)	0.0079 (0.0077)	0.0081 (0.0080)	**0.8792** (**0.0257**)	0.0080 (0.0079)	0.0080 (0.0079)	0.0105 (0.0100)	0.0080 (0.0079)	0.0087 (0.0085)	0.0086 (0.0083)	0.0081 (0.0078)	0.0081 (0.0078)	0.0105 (0.0099)	0.0082 (0.0081)
6_i	0.0088 (0.0087)	0.0088 (0.0086)	0.0088 (0.0087)	0.0087 (0.0085)	0.0089 (0.0087)	**0.8694** (**0.0270**)	0.0089 (0.0086)	0.0098 (0.0095)	0.0087 (0.0085)	0.0088 (0.0086)	0.0087 (0.0087)	0.0092 (0.0089)	0.0088 (0.0085)	0.0149 (0.0124)	0.0088 (0.0085)
7_i	0.0185 (0.0175)	0.0181 (0.0175)	0.0187 (0.0176)	0.0186 (0.0173)	0.0191 (0.0177)	0.0193 (0.0181)	**0.6849** (**0.0171**)	0.0711 (0.0320)	0.0187 (0.0177)	0.0187 (0.0179)	0.0185 (0.0178)	0.0203 (0.0194)	0.0185 (0.0175)	0.0186 (0.0178)	0.0183 (0.0175)
8_i	0.0119 (0.0114)	0.0119 (0.0113)	0.0122 (0.0117)	0.0120 (0.0116)	0.0138 (0.0134)	0.0123 (0.0117)	0.0121 (0.0117)	**0.8201** (**0.0329**)	0.0119 (0.0115)	0.0129 (0.0126)	0.0138 (0.0131)	0.0143 (0.0134)	0.0119 (0.0116)	0.0156 (0.0144)	0.0134 (0.0131)
9_i	0.0063 (0.0063)	0.0065 (0.0064)	0.0064 (0.0064)	0.0064 (0.0063)	0.0064 (0.0064)	0.0065 (0.0063)	0.0065 (0.0065)	0.0066 (0.0064)	**0.9055** (**0.0216**)	0.0064 (0.0061)	0.0086 (0.0086)	0.0086 (0.0078)	0.0064 (0.0063)	0.0064 (0.0063)	0.0065 (0.0064)
10_i	0.0132 (0.0127)	0.0134 (0.0131)	0.0133 (0.0127)	0.0136 (0.0134)	0.0134 (0.0127)	0.0135 (0.0126)	0.0133 (0.0129)	0.0136 (0.0131)	0.0135 (0.0127)	**0.7483** (**0.0703**)	0.0768 (0.0705)	0.0136 (0.0130)	0.0133 (0.0129)	0.0135 (0.0129)	0.0138 (0.0133)
11_i	0.0088 (0.0085)	0.0087 (0.0084)	0.0088 (0.0086)	0.0088 (0.0086)	0.0088 (0.0085)	0.0088 (0.0086)	0.0088 (0.0084)	0.0088 (0.0085)	0.0973 (0.0967)	0.0094 (0.0090)	**0.7873** (**0.0966**)	0.0095 (0.0092)	0.0088 (0.0086)	0.0087 (0.0085)	0.0086 (0.0086)
12_i	0.0081 (0.0080)	0.0081 (0.0080)	0.0081 (0.0079)	0.0081 (0.0080)	0.0081 (0.0079)	0.0087 (0.0083)	0.0080 (0.0079)	0.0088 (0.0085)	0.0098 (0.0094)	0.0082 (0.0078)	0.0088 (0.0086)	**0.8814** (**0.0253**)	0.0081 (0.0080)	0.0085 (0.0083)	0.0092 (0.0089)
13_i	0.0122 (0.0117)	0.0124 (0.0120)	0.0125 (0.0120)	0.0125 (0.0123)	0.0124 (0.0119)	0.0124 (0.0120)	0.0122 (0.0118)	0.0131 (0.0126)	0.0136 (0.0133)	0.0125 (0.0119)	0.0125 (0.0120)	0.0143 (0.0136)	**0.6792** (**0.0121**)	0.1557 (0.0323)	0.0125 (0.0120)
14_i	0.0127 (0.0121)	0.0125 (0.0122)	0.0129 (0.0124)	0.0129 (0.0125)	0.0127 (0.0121)	0.0150 (0.0144)	0.0130 (0.0123)	0.0131 (0.0126)	0.0129 (0.0123)	0.0127 (0.0124)	0.0129 (0.0124)	0.0129 (0.0124)	0.0127 (0.0122)	**0.8180** (**0.0322**)	0.0130 (0.0126)
15_i	0.0060 (0.0059)	0.0059 (0.0058)	0.0059 (0.0058)	0.0060 (0.0058)	0.0059 (0.0059)	0.0061 (0.0060)	0.0060 (0.0058)	0.0060 (0.0060)	0.0059 (0.0058)	0.0059 (0.0058)	0.0059 (0.0058)	0.0061 (0.0060)	0.0060 (0.0059)	0.0059 (0.0058)	**0.9164** (**0.0193**)

	1_j	2_j	3_j	4_j	5_j	6_j	7_j	8_j	9_j	10_j	11_j	12_j	13_j	14_j	15_j
1_i	**0.6848 (0.0172)**	0.0418 (0.0393)	0.0399 (0.0400)	0.0175 (0.0163)	0.0175 (0.0165)	0.0262 (0.0246)	0.0178 (0.0169)	0.0306 (0.0289)	0.0176 (0.0164)	0.0192 (0.0188)	0.0174 (0.0166)	0.0176 (0.0167)	0.0174 (0.0164)	0.0174 (0.0165)	0.0173 (0.0166)
2_i	0.0135 (0.0132)	**0.7976 (0.0340)**	0.0145 (0.0139)	0.0138 (0.0135)	0.0138 (0.0131)	0.0166 (0.0153)	0.0132 (0.0126)	0.0154 (0.0146)	0.0132 (0.0126)	0.0203 (0.0179)	0.0142 (0.0134)	0.0134 (0.0129)	0.0135 (0.0131)	0.0134 (0.0127)	0.0136 (0.0129)
3_i	0.0138 (0.0132)	0.1111 (0.0383)	**0.6858 (0.0186)**	0.0139 (0.0133)	0.0235 (0.0197)	0.0138 (0.0131)	0.0140 (0.0138)	0.0236 (0.0202)	0.0143 (0.0137)	0.0143 (0.0137)	0.0148 (0.0140)	0.0151 (0.0144)	0.0141 (0.0136)	0.0140 (0.0134)	0.0140 (0.0134)
4_i	0.0175 (0.0167)	0.0204 (0.0189)	0.0179 (0.0172)	**0.6839 (0.0163)**	0.0173 (0.0167)	0.0192 (0.0180)	0.0176 (0.0168)	0.0805 (0.0335)	0.0175 (0.0170)	0.0170 (0.0163)	0.0182 (0.0175)	0.0175 (0.0166)	0.0177 (0.0169)	0.0201 (0.0185)	0.0176 (0.0167)
5_i	0.0080 (0.0077)	0.0097 (0.0093)	0.0091 (0.0087)	0.0080 (0.0079)	**0.8789 (0.0260)**	0.0082 (0.0081)	0.0080 (0.0079)	0.0101 (0.0097)	0.0081 (0.0080)	0.0093 (0.0091)	0.0082 (0.0080)	0.0081 (0.0079)	0.0080 (0.0078)	0.0104 (0.0096)	0.0079 (0.0078)
6_i	0.0087 (0.0086)	0.0089 (0.0086)	0.0087 (0.0085)	0.0087 (0.0084)	0.0089 (0.0086)	**0.8697 (0.0270)**	0.0088 (0.0085)	0.0099 (0.0096)	0.0088 (0.0085)	0.0086 (0.0085)	0.0088 (0.0085)	0.0091 (0.0088)	0.0087 (0.0085)	0.0149 (0.0124)	0.0088 (0.0085)
7_i	0.0186 (0.0174)	0.0186 (0.0175)	0.0187 (0.0176)	0.0181 (0.0174)	0.0186 (0.0176)	0.0189 (0.0181)	**0.6854 (0.0178)**	0.0713 (0.0319)	0.0182 (0.0171)	0.0188 (0.0176)	0.0184 (0.0175)	0.0202 (0.0190)	0.0185 (0.0177)	0.0188 (0.0180)	0.0188 (0.0178)
8_i	0.0118 (0.0114)	0.0120 (0.0118)	0.0119 (0.0116)	0.0118 (0.0113)	0.0138 (0.0132)	0.0124 (0.0119)	0.0116 (0.0114)	**0.8204 (0.0323)**	0.0121 (0.0115)	0.0135 (0.0128)	0.0137 (0.0131)	0.0143 (0.0137)	0.0119 (0.0115)	0.0154 (0.0144)	0.0134 (0.0128)
9_i	0.0065 (0.0064)	0.0064 (0.0062)	0.0065 (0.0065)	0.0064 (0.0063)	0.0066 (0.0063)	0.0064 (0.0063)	0.0064 (0.0063)	0.0066 (0.0064)	**0.9034 (0.0216)**	0.0064 (0.0064)	0.0102 (0.0097)	0.0090 (0.0080)	0.0065 (0.0064)	0.0064 (0.0063)	0.0064 (0.0062)
10_i	0.0134 (0.0128)	0.0133 (0.0126)	0.0132 (0.0126)	0.0132 (0.0127)	0.0133 (0.0127)	0.0135 (0.0132)	0.0135 (0.0130)	0.0138 (0.0130)	0.0132 (0.0127)	**0.8033 (0.0452)**	0.0219 (0.0358)	0.0135 (0.0130)	0.0135 (0.0129)	0.0133 (0.0129)	0.0140 (0.0133)
11_i	0.0087 (0.0085)	0.0090 (0.0087)	0.0088 (0.0086)	0.0088 (0.0087)	0.0088 (0.0084)	0.0089 (0.0087)	0.0086 (0.0084)	0.0088 (0.0086)	0.0244 (0.0399)	0.0097 (0.0093)	**0.8593 (0.0457)**	0.0096 (0.0093)	0.0088 (0.0085)	0.0088 (0.0087)	0.0090 (0.0087)
12_i	0.0082 (0.0079)	0.0081 (0.0079)	0.0082 (0.0080)	0.0083 (0.0080)	0.0081 (0.0079)	0.0088 (0.0084)	0.0082 (0.0079)	0.0089 (0.0086)	0.0097 (0.0092)	0.0082 (0.0081)	0.0096 (0.0092)	**0.8802 (0.0255)**	0.0081 (0.0079)	0.0086 (0.0083)	0.0090 (0.0086)
13_i	0.0123 (0.0119)	0.0123 (0.0119)	0.0122 (0.0118)	0.0123 (0.0117)	0.0123 (0.0119)	0.0122 (0.0116)	0.0123 (0.0116)	0.0133 (0.0126)	0.0135 (0.0131)	0.0123 (0.0118)	0.0129 (0.0125)	0.0144 (0.0137)	**0.6790 (0.0117)**	0.1564 (0.0318)	0.0124 (0.0120)
14_i	0.0130 (0.0125)	0.0128 (0.0125)	0.0128 (0.0121)	0.0128 (0.0124)	0.0130 (0.0125)	0.0149 (0.0141)	0.0126 (0.0122)	0.0129 (0.0123)	0.0128 (0.0122)	0.0129 (0.0125)	0.0132 (0.0128)	0.0128 (0.0124)	0.0126 (0.0120)	**0.8182 (0.0322)**	0.0128 (0.0123)
15_i	0.0059 (0.0059)	0.0060 (0.0058)	0.0060 (0.0059)	0.0059 (0.0058)	0.0059 (0.0058)	0.0061 (0.0060)	0.0060 (0.0060)	0.0061 (0.0060)	0.0061 (0.0060)	0.0060 (0.0058)	0.0060 (0.0058)	0.0062 (0.0062)	0.0060 (0.0057)	0.0061 (0.0061)	**0.9157 (0.0194)**

	1_j	2_j	3_j	4_j	5_j	6_j	7_j	8_j	9_j	10_j	11_j	12_j	13_j	14_j	15_j
1_i	**0.6852** **(0.0185)**	0.0557 (0.0411)	0.0219 (0.0246)	0.0177 (0.0170)	0.0175 (0.0167)	0.0259 (0.0244)	0.0178 (0.0169)	0.0338 (0.0299)	0.0174 (0.0164)	0.0187 (0.0180)	0.0180 (0.0177)	0.0175 (0.0167)	0.0177 (0.0167)	0.0178 (0.0171)	0.0175 (0.0164)
2_i	0.0134 (0.0130)	**0.7989** **(0.0338)**	0.0139 (0.0132)	0.0135 (0.0128)	0.0136 (0.0131)	0.0171 (0.0158)	0.0134 (0.0129)	0.0156 (0.0150)	0.0133 (0.0130)	0.0179 (0.0165)	0.0156 (0.0149)	0.0133 (0.0127)	0.0135 (0.0128)	0.0135 (0.0130)	0.0135 (0.0131)
3_i	0.0143 (0.0136)	0.1194 (0.0364)	**0.6813** **(0.0143)**	0.0139 (0.0133)	0.0211 (0.0188)	0.0139 (0.0132)	0.0137 (0.0133)	0.0222 (0.0196)	0.0143 (0.0138)	0.0142 (0.0137)	0.0146 (0.0141)	0.0150 (0.0144)	0.0139 (0.0133)	0.0141 (0.0135)	0.0140 (0.0135)
4_i	0.0176 (0.0167)	0.0207 (0.0194)	0.0174 (0.0165)	**0.6845** **(0.0168)**	0.0176 (0.0170)	0.0185 (0.0175)	0.0175 (0.0168)	0.0801 (0.0331)	0.0177 (0.0167)	0.0176 (0.0168)	0.0177 (0.0169)	0.0175 (0.0165)	0.0174 (0.0162)	0.0209 (0.0192)	0.0172 (0.0164)
5_i	0.0081 (0.0079)	0.0099 (0.0093)	0.0081 (0.0079)	0.0082 (0.0078)	**0.8791** **(0.0253)**	0.0081 (0.0079)	0.0080 (0.0079)	0.0104 (0.0099)	0.0080 (0.0079)	0.0088 (0.0085)	0.0083 (0.0081)	0.0083 (0.0081)	0.0080 (0.0078)	0.0104 (0.0097)	0.0082 (0.0079)
6_i	0.0088 (0.0086)	0.0088 (0.0085)	0.0087 (0.0086)	0.0088 (0.0087)	0.0087 (0.0086)	**0.8698** **(0.0270)**	0.0089 (0.0084)	0.0097 (0.0092)	0.0088 (0.0086)	0.0087 (0.0084)	0.0088 (0.0086)	0.0091 (0.0086)	0.0087 (0.0083)	0.0148 (0.0122)	0.0088 (0.0086)
7_i	0.0185 (0.0178)	0.0186 (0.0178)	0.0183 (0.0172)	0.0184 (0.0175)	0.0188 (0.0178)	0.0194 (0.0185)	**0.6848** **(0.0173)**	0.0720 (0.0326)	0.0183 (0.0172)	0.0183 (0.0172)	0.0188 (0.0179)	0.0202 (0.0187)	0.0185 (0.0177)	0.0187 (0.0177)	0.0183 (0.0175)
8_i	0.0120 (0.0115)	0.0120 (0.0115)	0.0117 (0.0114)	0.0119 (0.0117)	0.0138 (0.0134)	0.0123 (0.0118)	0.0119 (0.0116)	**0.8204** **(0.0323)**	0.0121 (0.0117)	0.0132 (0.0128)	0.0139 (0.0133)	0.0141 (0.0132)	0.0119 (0.0116)	0.0154 (0.0145)	0.0133 (0.0128)
9_i	0.0065 (0.0065)	0.0064 (0.0063)	0.0064 (0.0063)	0.0065 (0.0064)	0.0065 (0.0064)	0.0064 (0.0062)	0.0065 (0.0063)	0.0065 (0.0065)	**0.9046** **(0.0216)**	0.0065 (0.0063)	0.0091 (0.0089)	0.0086 (0.0079)	0.0064 (0.0063)	0.0065 (0.0063)	0.0066 (0.0064)
10_i	0.0132 (0.0126)	0.0133 (0.0127)	0.0133 (0.0127)	0.0132 (0.0127)	0.0136 (0.0129)	0.0134 (0.0129)	0.0134 (0.0126)	0.0135 (0.0132)	0.0133 (0.0127)	**0.7636** **(0.0693)**	0.0623 (0.0674)	0.0136 (0.0130)	0.0132 (0.0126)	0.0133 (0.0130)	0.0139 (0.0133)
11_i	0.0087 (0.0085)	0.0088 (0.0086)	0.0088 (0.0087)	0.0087 (0.0085)	0.0086 (0.0084)	0.0090 (0.0087)	0.0087 (0.0084)	0.0089 (0.0087)	0.0836 (0.0938)	0.0092 (0.0089)	**0.8010** **(0.0933)**	0.0092 (0.0089)	0.0087 (0.0084)	0.0089 (0.0090)	0.0090 (0.0087)
12_i	0.0081 (0.0078)	0.0083 (0.0081)	0.0081 (0.0079)	0.0083 (0.0080)	0.0082 (0.0079)	0.0088 (0.0086)	0.0083 (0.0081)	0.0088 (0.0085)	0.0098 (0.0095)	0.0080 (0.0080)	0.0090 (0.0089)	**0.8810** **(0.0253)**	0.0081 (0.0079)	0.0084 (0.0083)	0.0089 (0.0086)
13_i	0.0123 (0.0116)	0.0125 (0.0120)	0.0123 (0.0118)	0.0125 (0.0120)	0.0124 (0.0118)	0.0124 (0.0117)	0.0121 (0.0117)	0.0128 (0.0123)	0.0132 (0.0125)	0.0125 (0.0121)	0.0129 (0.0125)	0.0145 (0.0136)	**0.6792** **(0.0121)**	0.1560 (0.0322)	0.0124 (0.0122)
14_i	0.0129 (0.0126)	0.0127 (0.0123)	0.0128 (0.0123)	0.0129 (0.0124)	0.0129 (0.0124)	0.0150 (0.0142)	0.0129 (0.0124)	0.0130 (0.0127)	0.0129 (0.0123)	0.0127 (0.0122)	0.0130 (0.0124)	0.0130 (0.0126)	0.0128 (0.0121)	**0.8176** **(0.0324)**	0.0129 (0.0124)
15_i	0.0059 (0.0058)	0.0060 (0.0059)	0.0059 (0.0058)	0.0061 (0.0060)	0.0059 (0.0060)	0.0060 (0.0058)	0.0060 (0.0059)	0.0060 (0.0060)	0.0059 (0.0058)	0.0058 (0.0058)	0.0061 (0.0059)	0.0060 (0.0059)	0.0059 (0.0059)	0.0059 (0.0057)	**0.9165** **(0.0192)**