The gut microbiome influences host diet selection behavior

Brian K. Trevelline[#,1,2,3] and Kevin D. Kohl[1]

[1] Department of Biological Sciences, University of Pittsburgh, Pittsburgh, Pennsylvania,

15260, USA

[2] Cornell Lab of Ornithology, Cornell University, Ithaca, New York, 14850 USA

[3] Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, New York

14850, USA

[#]Address correspondence to Dr. Brian K. Trevelline – brian.trevelline@gmail.com

SIGNIFICANCE: The behavior of diet choice or diet selection can have wide-reaching

implications, scaling from individual animals to ecological and evolutionary processes.

Previous work in this area has largely ignored the potential for intestinal microbiota to

modulate these signals. This notion has been highly speculated for years but has not

yet been explicitly tested. Here we show that germ-free mice colonized by differential

microbiomes (from wild rodents with varying natural feeding strategies) exhibited

significant differences in their voluntary dietary selection. Specifically, differences in

voluntary carbohydrate selection were associated with plasma amino acid levels and

bacterial genes involved in the metabolism of tryptophan. Together, these results

demonstrate a role for the microbiome in host nutritional physiology and behavior.

ABSTRACT

Diet selection is a fundamental aspect of animal behavior with numerous ecological and evolutionary implications. While the underlying mechanisms are complex, the availability of essential dietary nutrients can strongly influence diet selection behavior. The gut microbiome has been shown to metabolize many of these same nutrients, leading to the untested hypothesis that intestinal microbiota may influence diet selection. Here we show that germ-free mice colonized by gut microbiota from three rodent species with distinct foraging strategies differentially selected diets that varied in macronutrient composition. Specifically, we found that herbivore-conventionalized mice voluntarily selected a higher protein:carbohydrate ratio diet, while omnivore- and carnivore-conventionalized mice selected a lower P:C ratio diet. In support of the long-standing hypothesis that tryptophan – the essential amino acid precursor of serotonin – serves as a peripheral signal regulating diet selection, bacterial genes involved in tryptophan metabolism and plasma tryptophan availability prior to the selection trial were significantly correlated with subsequent voluntary carbohydrate intake. Finally, herbivore-conventionalized mice exhibited larger intestinal compartments associated with microbial fermentation, broadly reflecting the intestinal morphology of their donor species. Together, these results demonstrate that gut microbiome can influence host diet selection behavior, perhaps by mediating the availability of essential amino acids, thereby revealing a novel mechanism by which the gut microbiota can influence host foraging behavior.

INTRODUCTION

Proper nutrition is essential to life, and thus animals have evolved complex internal sensory systems that help maintain nutritional homeostasis by regulating macronutrient intake[1]. The intestinal tract plays a critical role in this process by liberating dietary nutrients (*e.g.,* essential amino acids) that communicate meal quality to the central nervous system by direct stimulation of enteric nerves or through post-absorptive peripheral signals[2–4]. The intestinal tract also harbors trillions of microorganisms (collectively known as the gut microbiome), which have been shown to influence numerous aspects of host behavior, most likely through metabolites that interact with host sensory systems[5]. Given the importance of dietary nutrients in the regulation of food intake and diet selection[6], the gut microbiome may influence host foraging behavior through metabolic processes that affect the availability of nutrients (or their derivatives) recognized by the central nervous system[2,7,8]. For example, a recent study showed that experimental colonization of *Providencia* bacteria in the gut of the model organism *C. elegans* resulted in divergent foraging preferences through the bacterial synthesis of the neurotransmitter tyramine from the essential amino acid tyrosine[9]. While studies in model systems provide powerful opportunities to dissect host-microbe interactions[10], the microbiome field recognizes the need to address and study the complexity of these interactions in ecologically-realistic scenarios in which animals can harbor thousands of microbial taxa[11,12]. It has been suggested that these complex microbial communities could elicit host foraging behaviors that enrich the intestinal environment in nutrients on which they depend (*i.e.,* promoting their own fitness)[7], while

others have posited that a positive-feedback relationship between dietary nutrients and

microbial community composition eventually results in stable microbial communities and

host foraging behaviors[8]. However, these potential mechanisms operate under the

assumption that the gut microbiome influences diet selection behavior – a hypothesis

that has existed for years[7,8], but has never been tested using complex microbial

communities, or within an ecological or evolutionary context.

The transplantation of intestinal microbiota into germ-free mice is a powerful

approach for disentangling the effects of the gut microbiome on host phenotypes from

other potentially confounding factors (*e.g.,* host genetics)[13]. This approach has been

successfully applied using a wide range of donor species (*e.g.,* termites, zebrafish)[14],

demonstrating that germ-free mice are a tractable model system for understanding the

function of gut microbiota in evolutionarily-distant organisms. In one notable example,

Sommer *et al.* used fecal microbiome transplants from brown bears into germ-free mice

(two species separated by ~94 million years of evolution) to show that seasonal

changes in gut microbiota influence host energy metabolism[15]. In our study, we used

this approach to determine whether the gut microbiome influences diet selection

behavior. We chose three rodent species with distinct foraging strategies as microbial

donors for germ-free mice: a carnivore/insectivore (southern grasshopper mouse,

Onychomys torridus), an omnivore (white-footed mouse, *Peromyscus leucopus*), and an

herbivore (montane vole, *Microtus montanus*). These three species are in the same

taxonomic family (Cricetidae) and are all equally distantly related to lab mice (~27 MYA;

Mus musculus, family Muridae)[16]. Under sterile laboratory conditions, we randomly

divided 30 adult male germ-free mice into Carn-CONV, Omni-CONV, and Herb-CONV

treatment groups (n = 10 mice per group), where each mouse in a given group was

"conventionalized" (*i.e.*, inoculated) with the cecal contents of a unique, wild-caught

donor individual (to better reflect natural interindividual variation) (Fig. 1a). One recipient

mouse from the Herb-CONV group was excluded from our dataset due to aberrant

behaviors   that   indicated   possible   injury   during   microbiome   transplants.

Conventionalized mice were acclimated to their microbiota for 7 days, during which they

were offered only sterile water and a low protein:carbohydrate ratio diet (LPC; Table

S1). There were no differences in daily or cumulative macronutrient and food intake

across treatment groups during the acclimation period (Fig. S1; Dataset S1). After

acclimation, conventionalized mice were given a choice between the LPC diet and one

with a higher P:C ratio (HPC; Table S1) for a period of 11 days (Fig. 1a). Importantly,

these diets had identical energy densities (caloric content per gram).

To determine whether treatment groups differed in foraging behavior, we

employed a state-space approach known as the Geometric Framework in which

foraging decisions are analyzed within a multi-dimensional nutritional space where each

functionally relevant nutrient forms a single dimension[17,18]. In this study, we defined

these nutritionally-explicit dimensions as protein and carbohydrate intake, thereby

allowing us to measure the effect of the gut microbiome on host diet selection.

Supporting the hypothesis that the gut microbiome influences diet selection behavior,

this approach revealed statistically significant differences in macronutrient intake across

groups of conventionalized mice (Fig. 1b). Treatment groups differed significantly in

daily (Fig. S1) and cumulative carbohydrate intake (Fig. 1b) during the diet selection

trial. Specifically, Herb-CONV mice voluntarily consumed fewer carbohydrates than

Carn-CONV and Omni-CONV mice. This trend was most apparent after approximately 1

week of diet choice (Fig. S1), suggesting that it may take time for internal nutritional

signals to stabilize[19] and for associative learning[20] to affect host feeding behavior. In

contrast, treatment groups did not differ in either daily (Fig. S1) or cumulative protein

intake (Fig. 1b). Lower cumulative carbohydrate intake among Herb-CONV mice led to

their selection of a significantly higher P:C ratio diet compared to Omni-CONV and

Carn-CONV mice (Fig. S2). Interestingly, we also observed a significant difference in

total food intake among Herb-CONV mice compared to the other treatment groups (Fig.

S1), suggesting that Herb-CONV mice's preference for the higher P:C ratio diet may

have permitted them to reduce total energy intake without affecting nutritional

homeostasis (*i.e.*, protein-leveraging)[19]. Under natural scenarios, such differences in

selected P:C ratios could be accomplished by animals incorporating different levels of

insects, seeds, or foliage into their diets. The ratio of macronutrients an animal

consumes, rather than the total amount of any individual nutrient, has significant effects

on animal physiology, life history, and reproductive fitness[21–23]. The preference of Herb-

CONV mice for the HPC diet are also consistent with previous studies showing that

Microtus voles prefer high-protein foods when available[24,25], though a follow-up study on

the foraging preferences of *M. montanus* with respect to specific dietary nutrients would

more robustly support the ecological significance of our findings. More generally, these

results are also consistent with the "nitrogen limitation hypothesis", which posits that the

relative scarcity of nitrogen in plant materials may drive the opportunistic consumption

of higher protein foods among herbivores[26–28]. Interestingly, the hindgut microbiota of

herbivorous mammals are also nitrogen-limited[29], and so our findings offer support to

the hypothesis that microbes may alter host foraging behaviors to enrich the intestinal

environment in necessary nutrients[7].

Next, we characterized day 0 (7 days post-inoculation and just prior to diet

selection trial) gut microbial community structure, microbiome function, and plasma

metabolites of conventionalized mice to determine how these aspects were associated

with differential diet selection across treatment groups. 16S rRNA inventories confirmed

that both donors and recipients harbored distinct bacterial communities that differed

significantly from blank extraction controls (Fig. 1c; Fig. S3; Fig. S4). We observed

significant differences in colonization efficiency across treatment groups. Specifically,

microbial communities of Carn- and Omni-CONV recipients were significantly most

similar to those of their donors, while Herb-CONV recipients were not significantly

similar to any donor group (Fig. S4). It is expected that recipient communities would not

match donors identically, as the *Mus* host physiology reshapes donor communities[30],

and our donor communities were collected from individuals in the wild, and thus our

design does not account for the well-documented effects of captivity on the

microbiome[31]. The comparatively lower colonization efficiency among Herb-CONV mice

may have been driven by the low content of indigestible plant fibers that are primarily

fermented by microbes. Even in established microbiomes, differences in the content or

composition of dietary fiber can result in the extirpation of some fermentive

microbes[32,33]. However, Herb-CONV mice were successfully colonized by donor

microbiota in the phylum Firmicutes (classes Bacilli and Clostridia), notably those in the

family Lachnospiracae, which are strict anaerobes known for their ability to transform

plant fibers into volatile fatty acids in the mammalian digestive tract[34]. Additionally,

microbiomes from herbivorous mammals colonize germ-free mice at a lower absolute

density than microbiomes from omnivorous or carnivorous mammals[35]. More work is

required to understand differential transfer of microbiomes across species, and we

discuss this limitation in more detail below.

Bacterial ASV richness and phylogenetic diversity were similar across donor

groups, but significantly lower in Herb-CONV mice compared to the other treatment

groups (Fig. S4). In general, the bacterial communities of conventionalized mice were

dominated by the phyla Bacteroidetes and Firmicutes (Fig. S4). Importantly, all recipient

fecal samples tested negative for the presence of pathogenic microorganisms.

Metagenomic analysis of recipient fecal samples revealed a statistically significant effect

of donor species on the relative abundances of 183 (51%) KEGG functional modules

(Fig. 1d; Dataset S2). These differences in microbiome community structure and

function were accompanied by concomitant differences in plasma metabolites (Fig. 1e),

with 27 identified metabolites (16%) differing significantly across treatment groups

(Dataset S3). Together, these results demonstrate that interspecific differences in gut

microbial communities across rodents with divergent foraging strategies translate to

distinct microbial functions and metabolite profiles independent of host diet.

There is substantial evidence that the availability of circulating essential amino

acids (EAAs) provide peripheral signals that act to regulate macronutrient intake and

diet selection[4,6]. Despite consuming identical diets prior to the selection trial, treatment

groups differed in circulating levels of several amino acids, with Herb-CONV mice

exhibiting significantly higher amounts of the EAAs lysine, isoleucine, methionine,

phenylalanine, and tryptophan (Fig. 2a). While EAAs are primarily derived from the diet,

bacteria can also produce these peptides through their own metabolic processes[36], and

thus the gut microbiome may act as a source of EAAs for their hosts. In support of this

hypothesis, treatment groups exhibited broad differences in the microbial synthesis and

degradation of EAAs (Fig. 2b). Notably, the microbiome of Herb-CONV mice had a

higher abundance of genes involved in the synthesis of aromatic amino acids

(phenylalanine, tryptophan, and tyrosine) (Fig. 2b), all of which are synthesized from

chorismate (product of the Shikimate pathway)[37]. The ratios of bacterial genes involved

in tryptophan biosynthesis (M00023) to those involved in tryptophan degradation via the

kynurenine pathway (M00038) were significantly correlated with plasma tryptophan (Fig.

2c). Given that conventionalized mice consumed identical diets prior to blood

collections, these results demonstrate that bacterial metabolism can alter the availability

of circulating levels of plasma EAAs, consistent with recent studies conducted in

Drosophila[38].

There is emerging evidence that bacterial tryptophan metabolism is a key

mechanism by which the gut microbiome can influence host behavior[39,40]. This

relationship is a consequence of tryptophan's role as the primary regulatory molecule

for the synthesis of central serotonin (5-hydroxytryptamine, 5-HT)[41], which has been

shown to drive foraging behavior and diet selection in several experimental studies[42,43].

For example, when given a choice between low- or high-carbohydrate meals, rats

receiving hypothalamic injections of 5-HT significantly reduced their carbohydrate

intake[44]. Importantly, serotonin synthesis is extraordinarily sensitive to plasma

tryptophan availability, and thus plasma tryptophan is generally considered a reliable

proxy for central serotonin[45]. Therefore, we predicted that plasma tryptophan would be

associated with differences in diet selection among conventionalized mice. Indeed, we

found a statistically significant correlation between day 0 plasma tryptophan and

subsequent voluntary carbohydrate intake (Fig. 2c). More recent work has argued that

serotonin synthesis is affected by the availability of tryptophan relative to the large

neutral amino acids (LNAA: Leu, Ile, Phe, Tyr, and Val) that compete for transport

across the blood brain barrier[46]. Consistent with these studies, we found a statistically

significant correlation between day 0 Trp:LNAA ratios and cumulative carbohydrate

intake (Fig. 2c). Further, the ratio of tryptophan biosynthesis and degradation KEGG

modules were also statistically significant predictors of carbohydrate and P:C intake

(Fig. 2c). Overall, these results support the hypothesis that bacterial tryptophan

metabolism influences host diet selection behavior.

Interspecific differences in foraging behavior are generally associated with diet-

specific adaptations to intestinal physiology. For example, herbivores generally maintain

an enlarged cecum (fermentation chamber) that enhances the digestibility of low-quality,

carbohydrate-rich foods[47]. Given that the gut microbiome can profoundly alter host

intestinal gene expression and physiology[48–50], divergent microbial communities may

drive differences in intestinal morphology across feeding strategies. At the conclusion of

the diet selection trial (day 11), we quantified intestinal morphology with the prediction

that conventionalized mice would exhibit differences that broadly reflected that of their

donor species. While there was no change in body mass over the duration of the

experiment ($F = 1.01$, $P = 0.377$), treatment groups differed significantly in empty colon

mass (Fig. 3b), with Herb-CONV mice exhibiting comparatively larger colons than those

in other treatment groups. There were no significant differences in cecum mass (Fig.

3a) or colon length (Fig. 3c). In general, the comparatively larger colons observed in

Herb-CONV mice are consistent with evolutionary adaptations observed in herbivorous

animals, which generally maintain larger hindguts to promote digestion[47]. The gut is a

highly dynamic organ that can rapidly change in mass and length in response to

environmental conditions, often through altered rates of cellular proliferation in intestinal

crypts and cell loss through sloughing or apoptosis at the ends of intestinal villi, but also

through the change of the size of individual enterocytes[51]. In the future, histological

analyses could be conducted to investigate whether these changes in gut size are

driven by hyperplasia (increase in cell number) and/or hypertrophy (increase in cell

size), and to rule out the possibility for these differences to be driven by intestinal

inflammation.

While the observed differences in gut size are consistent with adaptations

observed in herbivores, our study only tested the microbiome of a single species from

each feeding strategy. A robust test of whether the microbiome recapitulates the

differences in gut size observed across feeding strategies would require several donor

species from each dietary strategy. Another question is whether the gut microbiome

affected intestinal morphology directly or via differential diet selection. While our

experimental design makes it difficult to disentangle the effects of differential diet

selection from those of microbiome, it is worth noting that previous work has

demonstrated that lab mice fed low P:C ratio diets had larger intestinal compartments

(*e.g.,* colon) compared to those fed higher P:C diets[50]. In our study, we observed the

opposite – Herb-CONV mice, which consumed a higher P:C ratio diet (Fig. 1b),

exhibited larger colon masses (Fig. 3). These results contradict the generally accepted

model of adaptive physiological responses to dietary carbohydrates, suggesting that the

gut microbiome may drive interspecific differences in host intestinal physiology to some

extent, independent from the effects of diet and genetics.

Here, we present evidence for an effect of the gut microbiome on host diet

selection behavior, however, it is important to recognize that our approach has several

substantial limitations. For example, the relative differences in nutrient composition

between diets have been shown to greatly influence animals' ability to distinguish and

differentially feed[19], suggesting that our differential diet selection results may have been

more pronounced if we had used diets with greater differences in macronutrient content.

Further, previous work has shown that the evolutionary distance between donor species

and germ-free mice can affect the efficacy of microbiome transplants[14]. While our

selected donor species were similarly distant to *Mus musculus*, there were significant

differences in colonization success across donor species, suggesting that cecal

microbiota may be specifically adapted to their hosts. While differences in colonization

efficiency may limit our ability to robustly connect our study to the ecology of donor

species, this limitation should not diminish our major finding that conventionalized germ-

free mice harboring compositionally and functionally distinct microbiotas differing in

microbial diversity exhibited different feeding preferences. Overall, our approach is

stronger than comparing conventional mice with the highly artificial state of germ-free

mice, and the complex microbial communities that we used better reflect reality, which

is recognized as pressing need in the field of host-microbe interactions[11,12].

In this study, we found that conventionalized germ-free mice harboring distinct

gut microbiota exhibited significant differences in diet selection behavior, providing

support for our core hypothesis that microbiota can influence foraging decisions.

Specifically, our study provides evidence that variation in the gut microbiota alters host

nutrient availability and can yield significant differences in the diet selection of

conventionalized mice in just 11 days, likely through differential bacterial metabolism

and downstream availability of EAAs, especially tryptophan. These findings are largely

consistent with recent mechanistic work in model systems[9,38], but address the natural

variation in microbial communities that exist among individuals and across species[11,12].

Therefore, this study not only represents a novel contribution to a large body of work

showing that the gut microbiome is a key player in host physiology and performance[52],

but also more broadly supports the hypothesis that the gut microbiota can influence

ecological and evolutionary processes shaping animal behavior. Foraging strategies

and feeding behaviors can influence many aspects of an animal's ecology (*e.g.,* the

need to obtaining specific nutrients while also avoiding predators[53]), and animal feeding

can also shape the structures of entire plant and animal communities[54]. Thus, there may

be an underexplored role for gut microbes to influence far-reaching aspects of animal

and ecosystem ecology through influencing the feeding behavior of their hosts.

Acknowledgements. This work was supported by the National Science Foundation

(IOS-1942587 to K. Kohl) and a grant from the University of Pittsburgh Central

Research Development Fund. We thank A. Darracq, M. D. Dearing, T. Derting, R.

Martínez-Mota, and S. Secor for help with animal collections and dissections.

Author Contributions. K.D.K conceived the project. B.K.T designed the experiments,

performed the experiments, collected the data, interpreted the results, and wrote the

manuscript with guidance from K.D.K.

Additional Information. Supplementary information is available for this paper.

Sequencing data are deposited in the NCBI SRA database under PRJNA629007. The

authors declare no competing financial interests. Correspondence and requests for

materials should be addressed to B.K.T. (brian.trevelline@gmail.com).

MATERIALS AND METHODS

Wild rodents. Wild *Onychomys torridus* were collected in August 2018 from field

sites in near Green Valley, Pima Co., AZ (31.802834, -110.891172), *Peromyscus*

leucopus in May 2018 near Murray, Calloway Co., KY (36.686582, -88.221204), and

Microtus montanus in July 2018 at Timpie Springs Waterfowl Management Area,

Dugway, Tooele Co., UT (40.753708, -112.639903). Ten individuals from each species

were collected using baited Sherman live traps under the following state permits: *O.*

torridus (AZ Game and Fish Dept., SP627958), *P. leucopus* (KY Dept. of Fish and

Wildlife, SC1911097), and *M. montanus* (UT Division of Wildlife Resources,

1COLL5194-2). Animals were euthanized within 12 hours and immediately dissected

under IACUC protocols registered at the University of Utah (16-02011 to D. Dearing),

Murray State University (2018-026 to T. Derting), and University of Alabama (18-04-

1159 to S. Secor). Cecum contents for microbiome transplants were transferred to 1.7

mL Eppendorf tubes using sterile instruments and temporarily frozen at -20℃ in the

field before long-term lab storage at -80℃.

Microbiome transplants. Donor cecum contents were diluted at 100mg/mL in

sterile phosphate-buffered saline containing 0.2 g/L Na_2S and 0.5 g/L cysteine as

reducing agents[55,56]. Under sterile laboratory conditions, 30 adult (aged 6-8 weeks)

male germ-free C57BL/6 mice (Taconic Biosciences, Inc., Rensselaer, NY) were

randomly divided into Carn-CONV, Omni-CONV, and Herb-CONV groups (n = 10 mice

per group), where each mouse in a given group was colonized by oral gavage of 200 μL

of fecal slurry from a unique, wild-caught donor individual. Conventionalized mice were

then singly-housed in sterile static cages (Innovive, Inc., San Diego, CA; MSX2-AD)

modified by the addition of two metabolic feeder hoods (Laboratory Products, Inc.,

Seaford, DE; 2110S) that prevent mice from caching powdered diets, and thus enable

the tracking daily macronutrient intake (see below). Due to a lack of similar studies on

this topic, we were unable to conduct an *a priori* power analysis to justify the number of

donor/recipient mice per group. Instead, we decided on n = 10 per group based on the

number of animals typically used in studies involving germ-free mice, the vast majority

of which used 5-10 individuals per group[13]. One recipient mouse from the Herb-CONV

group (V57) was excluded from our dataset due to aberrant behaviors that indicated

possible injury during microbiome transplants. All recipient fecal samples were screened

for 21 of the most common rodent pathogenic microorganisms using PCR tests

conducted by a third-party diagnostic company (Charles River Research Animal

Diagnostic Services, Wilmington, MA).

Diet selection experiment. After colonization, conventionalized mice were acclimated for 7 days (to allow the gut microbiome to stabilize[55]), during which they were offered only sterile water and a low protein:carbohydrate ratio diet (LPC [0.27]; Table S1), as this diet is rather similar to standard mouse chow. After acclimation (day 0), mice were briefly removed from their cages for a 200 µL blood draw for metabolomics analysis (see details below). Mice were weighed (rounded to nearest hundredth) and returned to empty cages to facilitate the collection of fresh fecal samples for 16S rRNA microbial inventories and shotgun metagenomics (see details below). Conventionalized mice were then presented with a choice between two isocaloric diets (Table S1): (1) the LPC (0.27) diet offered during acclimation and (2) a diet with a higher P:C ratio (HPC [0.71]). The positions of these two diets were rotated daily to avoid learned preferences. Diets were designed by Teklad/Envigo (Indianapolis, IN), and were powdered prior sterilization to be visually indistinguishable from each other and to prevent food caching. Daily food consumption was calculated as the difference between the mass (rounded to nearest thousandth) of each diet presented (~8 g) and the mass of each diet remaining after a 24-hour period. After tracking diet preferences for 11 consecutive days, animals were euthanized and dissected to investigate differences in the empty masses (rounded to nearest thousandth) of intestinal compartments. Conventionalized mice were maintained on a 12:12-h light:dark cycle, with 21°C ambient temperature and 40% humidity for the duration of the experiment. Animal experiments were conducted at the University of Pittsburgh Plum Borough Primate Facility under IACUC protocol 19074445.

Metabolomics. Blood plasma was analyzed for primary metabolites (amino

acids, hydroxyl acids, carbohydrates, sugar acids, sterols, aromatics, nucleosides,

amines, and miscellaneous compounds) by the West Coast Metabolomics Center at the

University of California – Davis, which performed all sample preparation, data

acquisition, and data processing as previously described[57]. Briefly, metabolites were

extracted using a mixture of acetonitrile:isopropanol:water (3:3:2, v/v/v) as well as 1:1

acetonitrile:water for removal of protein from serum. Dried metabolite extracts were

resuspended in methoxyamine hydrochloride in pyridine for derivatization before being

analyzed using gas chromatography-time-of-flight (GC-TOF) using a LECO Pegasus IV

mass spectrometer equipped with automated liner exchange (ALEX; Gerstel

corporation) and cold injection system (CIS; Gerstel corporation) for data acquisition.

The CIS temperature was set at 50 °C to 250 °C final temperature at a rate of 12 °C

s−1. Raw GC-TOF MS data were preprocessed with ChromaTOF (version 2.32) and

apex masses were used to identify metabolites using the BinBase database. Values

were reported as peak height for the quantification ion (*m/z* value) at the specific

retention index, which is more precise than peak area for low abundant metabolites. All

database entries that were positively detected in more than 10% of the samples of a

study design class for unidentified metabolites were reported. Raw peak heights were

vector normalized to reduce the impact of between-series drifts of instrument sensitivity,

caused by machine maintenance status and tuning parameters.

DNA extractions. DNA was extracted from donor cecal contents and day 0

conventionalized mouse feces using the Qiagen PowerFecal DNA Kit (Qiagen, Hilden,

Germany; 12830) following the manufacturer's instructions.

16S rRNA microbial inventories. Extracted DNA from conventionalized mice and donor cecum contents was amplified and sequenced by the Genome Research Core of the University of Illinois at Chicago as previously described[58]. Briefly, polymerase chain reaction (PCR) was used to amplify a portion of the bacterial 16S rRNA gene for Illumina sequencing using the Earth Microbiome Project primers 515F (GTGCCAGCMGCCGCGGTAA) and 806R (GGACTACNVGGGTWTCTAAT) targeting the V4 region of microbial small subunit ribosomal RNA gene[59]. Amplicon libraries were sequenced using a 2x251 paired-end run on an Illumina MiSeq. In addition to donor and recipient fecal samples, we sequenced five 'blank' extractions to control for the possibility of microbial contamination during the extraction procedure and microbial DNA present in commercial extraction kits[60]. A total of 1,398,994 raw Illumina sequencing reads (mean of 22,206 per sample (n = 63) ± 1111 SE) were paired and quality filtered via the DADA2 pipeline[61] in QIIME2 (version 2020.4)[62] using default parameters. Sequences that passed the quality filter were clustered into amplicon sequence variants (ASVs), which were identified using the SILVA reference database (release 138)[63]. Identified ASVs were filtered to exclude non-bacterial sequences (archaea, chloroplast, eukaryote, and mitochondria), reducing our total number of reads to 1,396,450 (mean of 22,166 per sample ± 1,112 SE) and 4,359 ASVs. We detected a total of 4,118 ASVs in donor and recipient fecal samples, 19 (0.46%) of which were also detected in blank extractions (total of 260 ASVs from 27,807 reads with mean of 5,561 per sample ± 1,419 SE). As recommended by McMurdie and Holmes (2014)[64], we used un-rarefied ASV tables for comparisons of colonization efficiency (Bray-Curtis distances), alpha

diversity (ASV richness and Faith's phylogenetic diversity), and beta diversity (Bray-

Curtis and unweighted/weighted UniFrac distances[65]).

Shotgun metagenomics. Extracted DNA from conventionalized mice was sent

to CoreBiome, Inc. (St. Paul, MN) for shotgun metagenomic analysis using

BoosterShot™□. Briefly, sequencing libraries were prepared using a procedure adapted

from the Illumina Nextera Library Prep Kit (Illumina, 20018705) and sequenced on an

Illumina NovaSeq using single-end 1x100 reads with the Illumina NovaSeq SP reagent

kit (Illumina, 20027464). A total of 122,190,150 raw sequence reads (mean of 4,213,453

per sample (n = 29) ± 151,158 SE) were filtered for low quality (Q-Score < 30) and

length (< 50), trimmed of adapter sequences, and converted into a single fasta using

SHI7 (version 0.99)[66]. Sequences were then trimmed to a maximum length of 100 bp

and aligned using BURST (version 0.99.8)[67] at 97% identity against CoreBiome's Venti

database consisting of all RefSeq bacterial genomes with additional manually curated

strains as well as a bacterial KEGG[68] annotated database created from dereplicating

the bacterial genes within the Venti database. KEGG orthology counts were converted

to relative abundance within a sample and collapsed into KEGG modules for statistical

analysis.

Statistics. Differences in macronutrient and total diet intake across treatment

groups were tested using a multivariate analysis of variance (MANOVA) while

controlling for the effects of body mass. A *post hoc* power analysis for MANOVA was

conducted using G*Power[69] (version 3.1) to confirm that statistical power was

sufficiently greater than the widely-accepted minimum threshold of 0.80[70]. Microbial

community structure (from 16S rRNA inventories) was visualized using principal

coordinates analysis (PCoA) on ASV relative abundances, which were then assessed

for differences (controlling for multiple comparisons using false discovery rate corrected

P-values) across treatment groups using non-parametric permutational multivariate

analysis of variance (PERMANOVA), analysis of similarity (ANOSIM), and

permutational analysis of dispersion (PERMDISP) in QIIME2[62]. Microbiome function

was visualized using PCoA on KEGG module relative abundances and analyzed for

differences across treatment groups with PERMANOVA in QIIME2. Differences in the

relative abundance of functional KEGG modules across conventionalized mice were

tested using the non-parametric Krustal-Wallis test and linear discriminant analysis in

LEfSe using the "one-against-all" strategy for multi-class analysis[71]. Identified plasma

metabolites were filtered (based on mean intensity and IQR) and auto-scaled before

using non-parametric median tests to identify metabolites that varied significantly across

treatment groups and visualized using supervised partial least square discriminant

analysis (PLS-DA) in MetaboAnalyst (version 4.0)[72]. Non-parametric Spearman rank

correlations between plasma Trp availability, Trp KEGG modules, and macronutrient

intake were conducted using non-parametric Spearman's test (controlling for the effect

of donor species) in the R package *ppcor* (version 1.1)[73] and visualized using *corrplot*

(version 0.85)[74]. Differences in empty cecum mass, empty colon mass, and colon length

across treatment groups were tested using ANOVA with body mass as a covariate and

corrected for multiple comparisons using Tukey's HSD. Unless otherwise noted, all

statistical tests were two-sided and conducted in JMP Pro version 14.1.0 (SAS Institute

Inc., Cary, NC). For all statistical analyses, P-values ≤ 0.05 were defined as 'significant'.

REFERENCES

1. Kyriazakis, I., Tolkamp, B. J. & Emmans, G. Diet selection and animal state: an integrative framework. *Proc. Nutr. Soc.* **58**, 765–772 (1999).

2. Blanco, A. M., Calo, J. & Soengas, J. L. The gut–brain axis in vertebrates: Implications for food intake regulation. *J Exp Biol* **224**, jeb231571 (2021).

3. Furness, J. B. The enteric nervous system and neurogastroenterology. *Nature Reviews Gastroenterology & Hepatology* **9**, 286–294 (2012).

4. Stubbs, R. J. Peripheral signals affecting food intake. *Nutrition* **15**, 614–625 (1999).

5. Cryan, J. F. *et al.* The microbiota-gut-brain axis. *Physiological Reviews* **99**, 1877–2013 (2019).

6. Berthoud, H.-R., Münzberg, H., Richards, B. K. & Morrison, C. D. Neural and metabolic regulation of macronutrient intake and selection. *Proceedings of the Nutrition Society* **71**, 390–400 (2012).

7. Alcock, J., Maley, C. C. & Aktipis, C. A. Is eating behavior manipulated by the gastrointestinal microbiota? Evolutionary pressures and potential mechanisms. *Bioessays* **36**, 940–949 (2014).

8. Norris, V., Molina, F. & Gewirtz, A. T. Hypothesis: Bacteria control host appetites. *Journal of Bacteriology* **195**, 411–416 (2013).

9. O'Donnell, M. P., Fox, B. W., Chao, P.-H., Schroeder, F. C. & Sengupta, P. A neurotransmitter produced by gut bacteria modulates host sensory behaviour. *Nature* **583**, 415–420 (2020).

10.   Douglas, A. E. Simple animal models for microbiome research. *Nature Reviews*
Microbiology **17**, 764–775 (2019).

11.   Hanage, W. P. Microbiology: Microbiome science needs a healthy dose of
scepticism. *Nature News* **512**, 247 (2014).

12.   Greyson-Gaito, C. J. *et al.* Into the wild: Microbiome transplant studies need
broader ecological reality. *Proceedings of the Royal Society B* **287**, 20192834
(2020).

13.   Luczynski, P. *et al.* Growing up in a bubble: Using germ-free animals to assess
the influence of the gut microbiota on brain and behavior. *International Journal of*
Neuropsychopharmacology **19**, (2016).

14.   Seedorf, H. *et al.* Bacteria from diverse habitats colonize and compete in the
mouse gut. *Cell* **159**, 253–266 (2014).

15.   Sommer, F. *et al.* The gut microbiota modulates energy metabolism in the
hibernating brown bear *Ursus arctos. Cell Reports* **14**, 1655–1661 (2016).

16.   Steppan, S. J., Adkins, R. M. & Anderson, J. Phylogeny and divergence-date
estimates of rapid radiations in muroid rodents based on multiple nuclear genes.
Systematic Biology **53**, 533–553 (2004).

17.   Raubenheimer, D., Simpson, S. J. & Mayntz, D. Nutrition, ecology and nutritional
ecology: toward an integrated framework. *Functional Ecology* **23**, 4–16 (2009).

18.   Simpson, S. J., Raubenheimer, D. & Bone, Q. A multi-level analysis of feeding
behaviour: the geometry of nutritional decisions. *Philosophical Transactions of*
the Royal Society of London. Series B: Biological Sciences **342**, 381–402 (1993).

19. Sørensen, A., Mayntz, D., Raubenheimer, D. & Simpson, S. J. Protein-leverage

in mice: The geometry of macronutrient balancing and consequences for fat

deposition. *Obesity* **16**, 566–571 (2008).

20. Integration of learning and metabolic signals into a theory of dietary choice and

food intake. in *Ruminant physiology: digestion, metabolism, growth, and*

reproduction (eds. Cronjé, P. & Boomker, E. A.) (CABI Pub, 2000).

21. Reddiex, A. J., Gosden, T. P., Bonduriansky, R. & Chenoweth, S. F. Sex-specific

fitness consequences of nutrient intake and the evolvability of diet preferences.

The American Naturalist **182**, 91–102 (2013).

22. Solon-Biet, S. M. *et al.* The ratio of macronutrients, not caloric intake, dictates

cardiometabolic health, aging, and longevity in ad libitum-fed mice. *Cell Metab*

19, 418–430 (2014).

23. Solon-Biet, S. M. *et al.* Macronutrient balance, reproductive function, and lifespan

in aging mice. *Proc Natl Acad Sci USA* **112**, 3481–3486 (2015).

24. Bergeron, J. M. & Jodoin, L. Defining 'high quality' food resources of herbivores:

the case for meadow voles (*Microtus pennsylvanicus*). *Oecologia* **71**, 510–517

(1987).

25. Harju, A. & Hakkarainen, O. Effect of protein and birch-bark powder on selection

of food by root voles (*Microtus oeconomus*). *Journal of Mammalogy* **78**, 563–568

(1997).

26. White, T. C. R. *The inadequate environment: Nitrogen and the abundance of*

animals. (Springer Science & Business Media, 2012).

27.   White, T. C. R. The importance of a relative shortage of food in animal ecology.

523   *Oecologia* **33**, 71–86 (1978).

524   28.   Mattson, W. J. Herbivory in relation to plant nitrogen content. *Annual Review of*

525   *Ecology and Systematics* **11**, 119–161 (1980).

526   29.   Reese, A. T. *et al.* Microbial nitrogen limitation in the mammalian large intestine.

527   *Nature Microbiology* **3**, 1441–1450 (2018).

528   30.   Rawls, J. F., Mahowald, M. A., Ley, R. E. & Gordon, J. I. Reciprocal gut

529   microbiota transplants from zebrafish and mice to germ-free recipients reveal

530   host habitat selection. *Cell* **127**, 423–433 (2006).

531   31.   Alberdi, A., Martin Bideguren, G. & Aizpurua, O. Diversity and compositional

532   changes in the gut microbiota of wild and captive vertebrates: a meta-analysis.

533   *Sci Rep* **11**, 22660 (2021).

534   32.   Martínez-Mota, R., Kohl, K. D., Orr, T. J. & Denise Dearing, M. Natural diets

535   promote retention of the native gut microbiota in captive rodents. *ISME J* **14**, 67–

536   78 (2020).

537   33.   Sonnenburg, E. D. *et al.* Diet-induced extinctions in the gut microbiota compound

538   over generations. *Nature* **529**, 212–215 (2016).

539   34.   Meehan, C. J. & Beiko, R. G. A phylogenomic view of ecological specialization in

540   the Lachnospiraceae, a family of digestive tract-associated bacteria. *Genome*

541   *Biol Evol* **6**, 703–713 (2014).

542   35.   Contijoch, E. J. *et al.* Gut microbiota density influences host physiology and is

543   shaped by host and microbial factors. *eLife* **8**, e40553 (2019).

36.   Dai, Z.-L., Wu, G. & Zhu, W.-Y. Amino acid metabolism in intestinal bacteria:

Links between gut ecology and host health. *Frontiers in bioscience: a journal and*

virtual library **16**, 1768–86 (2011).

37.   Hermann, K. M. & Weaver, L. M. The shikimate pathway. *Annual Review of Plant*

Biology 473–503.

38.   Lesperance, D. N. A. & Broderick, N. A. Gut bacteria mediate nutrient availability

in *Drosophila* diets. *Appl. Environ. Microbiol.* **87**, (2020).

39.   Gao, K., Mu, C., Farzi, A. & Zhu, W. Tryptophan metabolism: A link between the

gut microbiota and brain. *Advances in Nutrition* **11**, 709–723 (2020).

40.   O'Mahony, S. M., Clarke, G., Borre, Y. E., Dinan, T. G. & Cryan, J. F. Serotonin,

tryptophan metabolism and the brain-gut-microbiome axis. *Behavioural Brain*

Research **277**, 32–48 (2015).

41.   Fernstrom, J. D. Role of precursor availability in control of monoamine

biosynthesis in brain. *Physiological Reviews* **63**, 484–546 (1983).

42.   Harrold, J. A., Dovey, T. M., Blundell, J. E. & Halford, J. C. G. CNS regulation of

appetite. *Neuropharmacology* **63**, 3–17 (2012).

43.   Blundell, J. E. & Halford, J. CG. Serotonin and appetite regulation. *Mol Diag Ther*

9, 473–495 (1998).

44.   Shor-Posner, G., Grinker, J. A., Marinescu, C., Brown, O. & Leibowitz, S. F.

Hypothalamic serotonin in the control of meal patterns and macronutrient

selection. *Brain Research Bulletin* **17**, 663–671 (1986).

45.   Fernstrom, J. D. & Wurtman, R. J. Brain serotonin content: Physiological

dependence on plasma tryptophan levels. *Science* **173**, 149–152 (1971).

46. Fernstrom, J. D. Large neutral amino acids: dietary effects on brain
neurochemistry and function. *Amino Acids* **45**, 419–430 (2013).

47. Stevens, C. E. & Hume, I. D. *Comparative physiology of the vertebrate digestive*
system. (Cambridge University Press, 2004).

48. Chevalier, C. *et al.* Gut microbiota orchestrates energy homeostasis during cold.
Cell **163**, 1360–1374 (2015).

49. Broderick, N. A., Buchon, N. & Lemaitre, B. Microbiota-induced changes in
Drosophila melanogaster host gene expression and gut morphology. *mBio* **5**,
e01117-14 (2014).

50. Sørensen, A., Mayntz, D., Simpson, S. J. & Raubenheimer, D. Dietary ratio of
protein to carbohydrate induces plastic responses in the gastrointestinal tract of
mice. *J Comp Physiol B* **180**, 259–266 (2010).

51. Starck, J. M. Shaping up: How vertebrates adjust their digestive system to
changing environmental conditions. *Animal Biology* **53**, 245–257 (2003).

52. McFall-Ngai, M. *et al.* Animals in a bacterial world, a new imperative for the life
sciences. *Proc Natl Acad Sci USA* **110**, 3229–3236 (2013).

53. McArthur, C., Banks, P. B., Boonstra, R. & Forbey, J. S. The dilemma of foraging
herbivores: Dealing with food and fear. *Oecologia* **176**, 677–689 (2014).

54. Martin, T. E. & Maron, J. L. Climate impacts on bird and plant communities from
altered animal–plant interactions. *Nature Climate Change* **2**, 195–200 (2012).

55. Molinaro, A. *et al.* Host–microbiota interaction induces bi-phasic inflammation
and glucose intolerance in mice. *Molecular Metabolism* **6**, 1371–1380 (2017).

56. Tremaroli, V. *et al.* Roux-en-Y gastric bypass and vertical banded gastroplasty
induce long-term changes on the human gut microbiome contributing to fat mass
regulation. *Cell Metabolism* **22**, 228–238 (2015).

57. Fiehn, O. *et al.* Quality control for plant metabolomics: Reporting MSI-compliant
studies. *The Plant Journal* **53**, 691–704 (2008).

58. Trevelline, B. K., MacLeod, K. J., Langkilde, T. & Kohl, K. D. Gestation alters the
gut microbiota of an oviparous lizard. *FEMS Microbiology Ecology* **95**, (2019).

59. Caporaso, J. G. *et al.* Global patterns of 16S rRNA diversity at a depth of millions
of sequences per sample. *Proceedings of the National Academy of Sciences*
108, 4516–4522 (2011).

60. Salter, S. J. *et al.* Reagent and laboratory contamination can critically impact
sequence-based microbiome analyses. *BMC Biol* **12**, 87 (2014).

61. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina
amplicon data. *Nature Methods* **13**, 581–583 (2016).

62. Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome
data science using QIIME 2. *Nature Biotechnology* **37**, 852–857 (2019).

63. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Research* **41**, D590–D596
(2013).

64. McMurdie, P. J. & Holmes, S. Waste not, want not: Why rarefying microbiome
data is inadmissible. *PLoS Comput Biol* **10**, e1003531 (2014).

65. Lozupone, C. & Knight, R. UniFrac: A new phylogenetic method for comparing
microbial communities. *AEM* **71**, 8228–8235 (2005).

66. Al-Ghalith, G. A., Hillmann, B., Ang, K., Shields-Cutler, R. & Knights, D. SHI7 is a self-learning pipeline for multipurpose short-read DNA quality control. *mSystems* **3**, e00202-17 (2018).

67. Al-Ghalith, G. & Knights, D. BURST enables optimal exhaustive DNA alignment for big data. (2017).

68. Kanehisa, M. & Goto, S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Research* **28**, 27–30 (2000).

69. Faul, F., Erdfelder, E., Lang, A.-G. & Buchner, A. G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods* **39**, 175–191 (2007).

70. Cohen, J. *Statistical Power Analysis for the Behavioral Sciences*. (Academic Press, 2013).

71. Segata, N. *et al.* Metagenomic biomarker discovery and explanation. *Genome Biology* **12**, R60 (2011).

72. Chong, J., Wishart, D. S. & Xia, J. Using MetaboAnalyst 4.0 for comprehensive and integrative metabolomics data analysis. *Current Protocols in Bioinformatics* **68**, e86 (2019).

73. Kim, S. ppcor: An R package for a fast calculation to semi-partial correlation coefficients. *Communications for Statistical Applications and Methods* **22**, 665 (2015).

74. Wei, T. & Simko. R package 'corrplot': Visualization of a correlation matrix. (2017).

 FIGURE LEGENDS

Fig. 1. The gut microbiome influences host diet selection behavior. a, Overview of

experimental design. Germ-free mice were colonized with the gut microbiome of three

species of wild rodents with distinct foraging strategies: carnivorous *Onychomys torridus*

(Carn-CONV), omnivorous *Peromyscus leucopus* (Omni-CONV), and herbivorous

Microtus montanus (Herb-CONV). Conventionalized mice were acclimated on LPC diet

for 7 days before day 0 blood and fecal sampling. After acclimation, conventionalized

mice were then given a choice between LPC and HPC diets for 11 days. Daily diet

intakes were tracked via two feeder hoods, which were rotated daily to avoid learned

preferences. **b,** Treatment groups differed significantly in macronutrient intake (Wilks' λ

= 0.455, Cohen's f^2 = 0.41, power = 0.98, P = 0.0007), with Herb-CONV mice voluntarily

consuming fewer carbohydrates than the Omni- and Carn-CONV groups (F = 9.22, P =

0.001). There was no difference in cumulative protein intake across treatment groups (F

= 1.362, P = 0.275). Dashed rails and associated P:C ratios indicate the expected result

if mice consumed only a single diet. Error bars represent the standard error of the mean

(SEM). **c,** Principal coordinate analysis (PCoA) of 16S rRNA inventories of wild donors

(squares) and conventionalized recipients at day 0 (circles) using Bray-Curtis

dissimilarity. Microbial community structure differed significantly among wild donors

(Pseudo-F = 7.41, P = 0.001) and recipients (Pseudo-F = 3.24, P = 0.001). All groups

differed significantly from blank extraction controls (gray diamonds; Pseudo-F = 4.78, P

= 0.001). **d,** PCoA analysis showing a statistically significant difference in the relative

abundances of microbial KEGG modules using Bray-Curtis dissimilarity (Pseudo-F =

5.96, P = 0.001). **e,** PLS-DA analysis illustrating broad differences in identified plasma

metabolites across conventionalized mice at day 0. * denotes P ≤ 0.05.

Fig. 2. Day 0 plasma tryptophan availability and bacterial tryptophan metabolism

are associated with differential macronutrient intake across treatment groups. a,

Heatmap illustrating broad differences in plasma levels of essential amino acids across

treatment groups, with Herb-CONV mice exhibiting significantly greater levels of lysine

(X^2 = 6.13, P = 0.047), isoleucine (X^2 = 11.42, P = 0.003), methionine (X^2 = 6.13, P =

0.047), phenylalanine (X^2 = 6.13, P = 0.047), and tryptophan (X^2 = 9.10, P = 0.011)

compared Carn-CONV and Omni-CONV mice. Columns represent individual

conventionalized mice for each treatment group. * denotes P ≤ 0.05 and color indicates

the treatment group with greatest circulating plasma levels (red = Carn-CONV, blue =

Omni-CONV, and yellow = Herb-CONV). **b,** Heatmap illustrating broad differences in

the abundances of microbial genes associated with metabolism of essential amino acids

(Dataset S2). * denotes P ≤ 0.05 and color indicates the treatment group with greatest

relative abundance. **c,** Correlation plot summarizing relationships between plasma

tryptophan availability, bacterial tryptophan metabolism, and host diet selection among

conventionalized mice. The direction and color of the ellipses indicate whether

correlations were positive or negative, and asterisks indicate whether Spearman's

correlations were statistically significant (* denotes P ≤ 0.05, ** denotes P < 0.01, and

*** denotes P < 0.001).

Fig. 3. Treatment groups exhibit differences in intestinal morphology. a, Empty

cecum mass did not differ significantly across treatment groups (F = 2.18, P = 0.133). **b,**

Empty colon mass differed significantly across treatment groups (F = 6.91, P = 0.004),

with Herb-CONV mice exhibiting a greater colon mass than Carn-CONV mice (FDR-adj.

P = 0.003). **c,** Colon length did not differ significantly across treatment groups (F = 2.03,

P = 0.151)**.** ** denotes P ≤ 0.01.

a

O. torridus *P. leucopus* *M. montanus*

Donors (n=10)

Colonization (n=10)

Acclimation (7 days)

LPC
LPC

Sampling (day 0)

Selection trial (11 days)

HPC
LPC

Carn-CONV **Omni-CONV** **Herb-CONV**

b Diet choice

P:C = 0.27 (LPC only)
P:C = 0.71 (HPC only)

Cumulative carbohydrate intake (g)

Cumulative protein intake (g)

● Carn-CONV
● Omni-CONV
● Herb-CONV

c Microbiome structure

PCoA 2 (7.43%)

PCoA 1 (9.47%)

d Microbiome function

PCoA 2 (19.4%)

PCoA 1 (32.3%)

e Plasma metabolites

PLS-DA 2 (8.8%)

PLS-DA 1 (16.1%)

◇ Blank extraction □ Donor ○ Recipient ● Blank extraction ● Carn-CONV ● Omni-CONV ● Herb-CONV