KDM6B promotes activation of the oncogenic CDK4/6-pRB-E2F pathway by maintaining

enhancer activity in MYCN-amplified neuroblastoma

Alexandra D'Oto[1*], Jie Fang[1*], Hongjian Jin[2], Beisi Xu[2], Shivendra Singh[1], Anoushka Mullasseril[1],

Victoria Jones[1], Ahmed Abu-Zaid[1], Xinyu von Buttlar[1], Bailey Cooke[1], Dongli Hu[1], Jason Shohet[3],

Andrew J Murphy[1], Andrew M Davidoff[1], Jun Yang[1]

[1]Department of Surgery, [2]Center for Applied Bioinformatics, St. Jude Children's Research Hospital, 262

Danny Thomas Place, Memphis, TN 38105

[3] Department of Pediatrics, University of Massachusetts Medical School, 55 Lake Avenue North,

Worcester, MA 01655

*equal contribution

Correspondence: Jun.Yang2@stjude.org or Andrew.Davidoff@stjude.org

ABSTRACT

The H3K27me2/me3 histone demethylase KDM6B is over-expressed in neuroblastoma and is essential to neuroblastoma cell survival. While the KDM6B inhibitor, GSK-J4, has shown activity in *in vitro* and *in vivo* preclinical models, the mechanism of action remains poorly defined. We demonstrate that genetic and pharmacologic inhibition of KDM6B downregulates the pRB-E2F transcriptome and *MYCN* expression. Chemical genetic analyses show that high expression of the E2F transcriptome is positively correlated with sensitivity of cancer cells to GSK-J4. Mechanistically, inhibition of KDM6B activity 1) reduces the chromatin accessibility of E2F target genes and *MYCN, 2)* selectively leads to an increase of H3K27me3 but a decrease of the enhancer mark H3K4me1 at the CTCF and BORIS binding sites, which may, consequently, disrupt the long-range chromatin interaction of *MYCN* and E2F target genes, and 3) phenocopies the transcriptome induced by the specific CDK4/6 inhibitor palbociclib. Overexpression of CDK4/6 or *Rb1* knockout confers neuroblastoma cell resistance to both palbociclib and GSK-J4. A gene signature targeted by KDM6B inhibition is associated with poor survival of patients with neuroblastoma regardless of the *MYCN* status. These data indicate that KDM6B activity promotes an oncogenic CDK4/6-pRB-E2F pathway in neuroblastoma cells via H3K27me3-dependent enhancer-promoter interactions, providing a rationale to target KDM6B for high-risk neuroblastoma.

Key words: KDM6B, MYC, E2F, Neuroblastoma, CDK4/6, Palbociclib

INTRODUCTION

KDM6 is a Jumonji domain-containing H3K27me3/me2 demethylase subfamily[1-4] that antagonizes the activity of polycomb repressive complex 2 (PRC2)[5], the methyltransferase of H3K27. KDM6 consists of 3 members, KDM6A, KDM6B and UTY, although UTY has low lysine demethylase activity[6]. While KDM6A is believed to be a tumor suppressor and is mutated in many different types of cancer[7,8], the function of KDM6B in cancer remains poorly defined. Early studies have shown that KDM6B contributes to activation of the Ink4a/Arf locus in fibroblasts in response to oncogenic stress[9,10], and nuclear p53 protein stabilization in glioblastoma cells[11]. Loss of KDM6B enhances aggressiveness of pancreatic cancer cells[12]. These data suggest KDM6B might have a tumor suppressive function. However, KDM6B is overexpressed in various cancers[13], indicating that KDM6B may play different roles depending on the cellular context. KDM6B appears to be essential for the initiation and maintenance of NOTCH-driven acute T-cell lymphoblastic leukemia[14], regulates multiple myeloma cell growth and survival by modulating the MAPK pathway[15], drives glioblastoma stem cell plasticity and drug tolerance by chromatin remodeling[16], and promotes migration, invasion, and stem cell-like behaviors in hepatocellular carcinoma[17]. KDM6B is also involved in chemotherapy resistance[18].

Neuroblastoma, the most common extra-cranial solid tumor in children, arises as a result of blocked differentiation of neural crest precursors (NCCs) during development of the sympathetic nervous system[19,20]. This aggressive malignancy accounts for 15% of cancer-related deaths in children[21]. Although outcomes for children with low- or intermediate-risk disease are excellent, fewer than 50% of children with high-risk disease survive despite aggressive multimodal therapy. In patients with high-risk disease, one key biological feature associated with poor prognosis is amplification of the *MYCN* oncogene (MNA)[22]. As a master lineage transcription factor that drives tumorigenesis, the functions of MYCN have been associated with targetable epigenetic modifiers[23-25]. Our previous study demonstrated that the histone demethylase KDM4B regulates MYC activity and promotes tumor growth and maintenance of neuroblastoma[26]. More recently, we identified a novel KDM4B inhibitor ciclopirox[27], that inhibits tumor growth and promotes differentiation[27]. EZH2, the essential catalytic unit of the PRC2 complex, is a MYCN target that represses neuronal differentiation in a H3K27me3-dependent manner, leading to inactivation of a tumor suppressive program in neuroblastoma[28].

In this study, we show that KDM6B is highly expressed in neuroblastoma and its genomic locus is broadly marked with transcriptionally active histone modifications and transcription factor binding. Using chemical genetic and epigenetic approaches, we show that KDM6B is involved in regulation of the key oncogenic CDK4/6-pRB-E2F pathway and expression of both MYCN and C-MYC proto-oncogenes.

Furthermore, the E2F transcriptome serves as a biological marker of KDM6 inhibitor sensitivity.

Pharmacological inhibition of KDM6B activity alters the chromatin accessibility of E2F target genes and

MYCN, induces the redistribution of H3K27me3 and the enhancer mark H3K4me1. We propose that this

may disrupt the long-range chromatin interaction of the E2F transcriptome within the same topologically

associated domains (TAD). At the level of transcriptome, KDM6B inhibition mimics the CDK4/6

inhibitor palbociclib. Cancer cells resistant to CDK4/6 inhibitor also are resistant to KDM6B inhibition.

The gene signature targeted by KDM6 inhibition was associated with poor patient survival. Thus, our

studies reveal that KDM6B regulates the oncogenic CDK4/6-pRB-E2F pathway in MYCN-amplified

neuroblastoma, revealing a new mechanism of regulation of the E2F transcriptome by an epigenetic

modulator.

RESULTS

***KDM6B* is highly expressed in human neuroblastoma and is associated with a transcriptionally**

active epigenetic landscape

We have previously shown that KDM6B, but not KDM6A or UTY, is essential for neuroblastoma cell

survival[27]. To more clearly define the role of KDM6B in neuroblastoma, we first compared expression of

KDM6 family genes in normal trunk neural crest-derived tissues, neuroblastoma being a neural crest-

derived cancer of the peripheral nervous system[29], with expression in tumors in 4 different neuroblastoma

patient cohorts[30-32]. While expression of *KDM6A* and *UTY* did not show consistent differences in tumors

as compared to normal tissue of neural crest origin, *KDM6B* expression was significantly elevated across

all 4 neuroblastoma cohorts (p<0.001) (Figure 1A-1C). We also found that levels of *KDM6B* expression,

but not *KDM6A* or *UTY,* in neuroblastoma cell lines were among the highest, across 40 different cancer

lineages (Supplementary Figure 1A). Analysis of the data from 15 subtypes of pediatric cancers (PCGP

project performed at St Jude) also showed *KDM6B* expression was among the highest in neuroblastoma

(Supplementary Figure 1B). We then compared the RNA-seq transcript counts of *KDM6B*, *KDM6A* and

UTY in three large neuroblastoma cohorts and found that expression of *KDM6B* was significantly higher

than that of *KDM6A* or *UTY* (Figure 1D-1F).

Gene transcription status is usually associated with a distinct epigenetic landscape. For example,

high levels of H3K4me1, H3K4me3 and H3K27Ac are marks of active gene transcription while high

levels of H3K27me3 and H3K9me3 are marks of repressive gene transcription[33]. To determine whether

high expression of KDM6B in neuroblastoma is associated with active epigenetic modifications, we

investigated the epigenetic landscapes at the genomic loci of *KDM6B*, *KDM6A* and *UTY* in 7 primary

human neuroblastoma tissues sequenced at St Jude, 4 with MNA and 3 without MNA (Figure 1G,

Supplementary Figure 2). Regardless of the *MYCN* status, the genomic locus of *KDM6B* across 22-kb in

length in all 7 human neuroblastoma cell lines is occupied by RNA polymerase II, which drives gene

transcription, in accordance with enrichment of active gene transcription marks H3K4me1, H3K4me2,

H3K4me3, H3K27Ac, H3K9-K14Ac, and H3K36me3. However, no transcriptionally repressive

H3K27me3 or H3K9me3 marks were occupied at the *KDM6B* locus. The broad occupation of H3K4me1

and H3K27Ac across the whole locus of *KDM6B* indicates that a super-enhancer[34,35] may regulate the

expression of *KDM6B* in neuroblastoma. In contrast, the epigenetic landscapes at the *KDM6A* and *UTY*

loci appeared to lack transcriptionally active gene transcription marks (Figure 1G, Supplementary Figure

2). The promoter region and gene body of *KDM6A* were occupied by the transcriptionally repressive marks

H3K27me3 and H3K9me3, respectively. In summary, *KDM6B* is highly expressed in human

neuroblastoma and its genomic locus is epigenetically modified for active gene transcription, which is

distinct from its paralogs *KDM6A* and *UTY*.

KDM6B regulates MYC expression

We further validated the importance of KDM6B in neuroblastoma by assessing colony formation after

knocking down *KDM6B* with 4 different siRNAs in BE2C cells (with MNA) and SK-N-AS cells (with

high C-MYC activity). KDM6B depletion greatly reduced colony formation in both neuroblastoma cell

lines, and was correlated with the KDM6B knockdown efficiency (Figure 1H,1I), consistent with our

previous study[27]. In support of our data, neuroblastoma is the third most sensitive among 25 different

cancer lineages that are sensitive to knockdown of KDM6B but not KDM6A (Supplementary Figure 3A,

3B), just below prostate cancer and myeloma, in a genome-wide shRNA screen (20 shRNAs per gene) in

709 cancer cell lines (DepMap dataset). Notably, depletion of KDM6B led to MYCN and C-MYC

reduction in neuroblastoma cells (Figure 1I, Supplementary Figure 3C-F). Importantly, introduction of

retroviral based KDM6B cDNA without its endogenous 3'UTR rescued the colony formation and MYCN

expression in BE2C cells when endogenous KDM6B was depleted using a siRNA designed to target its

3'UTR (Figure 1J-L), demonstrating the phenotype induced by KDM6B depletion is not due to an off-

target effect. Conversely, overexpression of KDM6B in BE2C cells reduced H3K27me3, increased

MYCN expression and promoted cell proliferation (Supplementary Figure 3G, H). Additionally, we found

that KDM6B also regulated MYC expression in other cancer lineages such as lung cancer, osteosarcoma,

and colorectal cancer cells and that this was independent of p53 status (Supplementary Figure 3I-L). These

data indicate that KDM6B is needed for MYC expression in cancer cells.

KDM6B regulates a pRB-E2F transcription program in neuroblastoma

To investigate the molecular function of KDM6B in neuroblastoma, we performed RNA-seq followed by gene set enrichment analysis (GSEA) after knockdown of KDM6B in MYCN-amplified (BE2C and KELLY) and MYCN-non-amplified cells (SK-N-AS, SK-N-FI). Strikingly, GSEA results showed that the top hits of KDM6B downstream targets were enriched with E2F gene sets, *Rb1* knockout signatures (*Rb1* encodes pRB, is a negative regulator of E2F activity[36]) and other cell cycle gene signatures (Figure 2A-2C, Supplementary Figure 4A, 4B), in the two MYCN amplified cell lines. Motif analysis of the KDM6B target genes also revealed that they were mainly enriched with E2F binding motifs (Table 1). These data indicate that KDM6B regulates the transcriptome of the E2F pathway in MYCN-amplified neuroblastoma cells. The E2F transcription factors are critical regulators of cell cycle progression by directly regulating more than 100 genes including those encoding cyclins[37]. Early studies showed that E2F also binds to promoters of *C-MYC* and *MYCN* to regulate their expression[38,39]. GSEA data revealed that MYC targets were also significantly downregulated by KDM6B depletion (Figure 2A). Interestingly, the top signatures upregulated by KDM6B knockdown in SK-N-FI were enriched with E2F gene sets (Supplementary Figure 4C, 4D), although C-MYC expression was reduced (Supplementary Figure 3F). However, the E2F gene signatures were less affected in SK-N-AS cells than BE2C, KELLY and SK-N-FI although one E2F3 gene set was downregulated by KDM6B knockdown (Supplementary Figure 3E, 3F).

We further investigated KDM6B function using pharmacologic inhibition. GSK-J1 is the only selective KDM6 inhibitor available[40], but, unfortunately, it is unable to penetrate cells. To make GSK-J1 cell-permeable, GSK-J1 was modified by adding an ester (new reagent was named GSK-J4)[40]. Thus, GSK-J4 is not itself a chemical tool for direct KDM inhibition, but was designed specifically to enable efficient intracellular delivery of GSK-J1 into cells. The intracellular conversion of the ester pro-drug is complete within 15 min after which levels of intracellular GSK-J4 are negligible[41]. Once entering cells, GSK-J4 converts back to GSK-J1, consequently inhibiting KDM6 (Supplementary Figure 5A)[40]. To validate this and profile the selectivity of GSK-J1/GSK-J4, we used an AlphaLISA approach, a method that tests the *in vitro* inhibitory activity of GSK-J1 and GSK-J4 against purified KDMs (KDM2A, KDM3A, KDM4A, KDM5A, KDM6A, KDM6B). We confirmed that GSK-J1 was selectively effective against KDM6A and KDM6B, had some effect on KDM5A but showed no activity against KDM2A, KDM3A and KDM4A (Supplementary Figure 5B), a result consistent with published GSK data[40]. However, GSK-J4 showed no *in vitro* activity against all tested KDMs (Supplementary Figure 5C). While GSK-J4 could have off-target

effects when administered at high concentrations, the concentration we used in cells (2.5 μM) only induced

the upregulation of H3K27me3, but not H3K9me3, H3K36me3 and H3K4me3 (Supplementary Figure

5D), three other major histone methylation sites of H3, indicating that GSK-J4 induced its cellular effect

by selectively targeting KDM6 but not other KDMs. Additionally, a selective KDM5 inhibitor, KDM5-

C70[42], showed no effect on neuroblastoma cell proliferation even when administered at very high

concentrations (40μM) (Supplementary Figure 5E, 5F).

Treatment of neuroblastoma cells and normal human fibroblasts (HS68) with GSK-J4, in a colony

formation assay, showed that GSK-J4 inhibited colony formation in a concentration-dependent manner,

with a significant effect observed in the majority of cell lines at concentrations of 1.0-2.0μM

(Supplementary Figure 6A). However, HS68 cells were unaffected by GSK-J4. Consistent with these

results, another recent study showed that GSK-J4 has a potent effect on neuroblastoma cell survival[43]. To

define the molecular mechanism of GSK-J4 action on neuroblastoma cells, we performed global gene

expression profiling for GSEA analysis after GSK-J4 treatment of BE2C cells. GSK-J4 induced a very

similar transcriptome profile as depletion of KDM6B and the gene sets most significantly downregulated

by this inhibitor were enriched with cell cycle, *Rb1* knockout and E2F pathways (Figure 2D-2F).

Transcription factor motif binding analysis showed that genes downregulated by GSK-J4 were greatly

enriched with E2F binding (Table 2). Again, MYCN and C-MYC expression were also downregulated by

GSK-J4 treatment (Supplementary Figure 6B-6D), consistent with KDM6B depletion (Supplementary

Figure 3). Re-analysis of an independent RNA-seq data set[43] also showed that KDM6 inhibition led to

downregulation of gene sets enriched with *Rb1* knockout and DREAM complex (dimerization partner,

RB-like, E2F and multi-vulval class B) in neuroblastoma cells (Supplementary Figure 7). Thus, GSK-J4

phenocopied KDM6B depletion in neuroblastoma cells and, therefore, its therapeutic effect may be

mediated through inhibition of KDM6B. Taken together, genetic and pharmacologic inhibition of

KDM6B predominantly leads to downregulation of the pRB-E2F transcriptional program, particularly in

MYCN-amplified neuroblastoma cells.

Pharmacogenetics reveals that the E2F transcriptome is associated with sensitivity to KDM6B

inhibition

While the presence of KDM6B is a prerequisite for GSK-J4 activity in neuroblastoma cells[43], expression

levels of KDM6B do not correlate with GSK-J4 sensitivity[43]. A biomarker that predicts therapy response

is critical for patient stratification. Since KDM6B inhibition appears to effect its anti-tumor activity

through downregulation of the E2F transcriptome, we hypothesized that high levels of the E2F

transcriptome may correlate with the response of cancer cells to KDM6B inhibition. The gene signatures
induced by chemical (i.e., small-molecule) or genetic perturbations (i.e., siRNA knockdown) can be used
to connect unknown mechanisms of action (MoA) of chemical probes. Differential gene expression has
been correlated with patterns of small-molecule sensitivity across many cell lines to illuminate the actions
of compounds whose MoA are unknown[44,45]. The Cancer Therapeutic Response Portal correlated the
sensitivity patterns of 481 compounds including GSK-J4 with 19,000 basal transcript levels across 823
different human cancer cell lines and identified selective outlier transcripts[44,45], which allowed us to
interrogate whether the KDM6B inhibitor activity was correlated with specific transcripts. The E2F genes
such as *E2F8* were the ones most significantly affected by KDM6B inhibition (Figure 2). E2F8 is an
atypical E2F and downstream target of E2F1-3[46]. We chose *E2F*8 as a representative gene to correlate
E2F pathway activity with drug effect. Strikingly, GSK-J4 ranked highest among the top compounds
whose cellular activity was correlated with *E2F8* expression (Pearson correlation R=-0.528, z-score = -
4.56, Figure 3A). We arbitrarily chose 13 cell lines with high *E2F8* expression as group A, and 8 cell lines
with low *E2F8* expression as group B, which had high sensitivity and low sensitivity to GSK-J4,
respectively (Figure 3B, Supplementary Table 1). We then used GSEA to infer the pathways that differed
in the two groups. We found that the pRB-E2F gene sets were the most significant hits in group A cell
lines (Figure 3C, 3D). The genes highly expressed in group A cells were profoundly enriched with E2F
transcription factor binding motifs (Supplementary Table S2), and most of the top 50 genes such as *E2F1*
and *E2F8* are involved in regulation of the cell cycle (Supplementary Figure 8A). MYC signatures were
also enriched in group A cells (Supplementary Figure 8B-8D). We also selectively investigated several
other E2F genes including *ASF1B*, *CDK2* and *E2F1*, all of which were positively correlated with GSK-J4
sensitivity (Supplementary Figure 8E-8G). Therefore, cancer cells with high E2F activity appear to be
more sensitive to KDM6B inhibition.

To specifically investigate the conclusion that high E2F activity serves as a biomarker for sensitivity
to KDM6B inhibition in neuroblastoma, we categorized neuroblastoma cell lines into two groups
according to their IC_{50} to GSK-J4[43]. GSEA analysis of RNA-seq data of these cell lines showed a very
similar transcriptome pattern to KDM6B knockdown or GSK-J4 inhibition for those with lower IC_{50} to
GSK-J4 (more sensitive), which were enriched with E2F and MYC gene signatures (Figure 3E). For
example, SK-N-FI cells with high E2F transcriptome expression were more sensitive to GSK-J4 than SK-
N-SH cells (Figure 3F), which had lower expression of the E2F transcriptome. The genes highly expressed
in neuroblastoma cells that were more sensitive to GSK-J4 were predominantly enriched with E2F
transcription factor binding motifs (Table 3). Taken together, these data demonstrate that a high E2F gene

signature is correlated with GSK-J4 sensitivity, and thus it may serve as a biomarker for effectiveness of

KDM6B inhibitors in cancer treatment in the future. These data also further support that KDM6B regulates

the E2F transcriptome.

Inhibition of KDM6B activity reduces the chromatin accessibility of E2F target genes and *MYCN*

To understand how KDM6B regulates the E2F pathway, we performed Assay for Transposase-Accessible

Chromatin with high-throughput sequencing (ATAC-seq) after 24h and 48h of GSK-J4 treatment, for

mapping genome-wide chromatin accessibility[47]. The 24h GSK-J4 treatment resulted in upregulation of

8386 nucleosome-free, open chromatin regions (log2 fold change >=0.5, p<0.05, Figure 4A), 46.9% of

which were located at the annotated promoter regions (within 2-kb from transcription start site, TSS) and

39% of which were located at the annotated enhancer regions (2-50kb distance from TSS). The rest

(14.1%) were located at more distal regions. However, 9574 nucleosome free, open chromatin regions

were downregulated by GSK-J4 treatment (Figure 4A), only 18.5% of which were located at the promoter

regions and 61.8% of which were located at enhancer regions, while 19.7% were located at more distal

regions. The 48h GSK-J4 treatment further enhanced its effect on DNA accessibility, resulting in an

increase of 13546 and decrease of 11027 peaks, respectively (Figure 4A). The proportions of the annotated

regions tended to be similar to that of 24h treatment. Among the regions with upregulated chromatin

accessibility, 57.6% were located at promoter regions, 29.5% were located at enhancer regions, and 12.9%

were located at more distal regions (Figure 4A). Among the regions with downregulated chromatin

accessibility, 17.9% were located at promoter regions and 57.8% were located at enhancer regions, while

24.3% were located at more distal regions (Figure 4A). We further assessed whether altered chromatin

accessibility was associated with changes in gene expression. Indeed, the genes annotated with differential

accessibility regions (DARs) at promoters or enhancers were significantly enriched in differentially

expressed genes by GSK-J4 treatment (Supplementary Figure 9).   For example, DNA accessibility at the

promoter region of the oncogene *AKT1* was reduced by GSK-J4, and one additional peak at a potential

enhancer region near the long non-coding RNA *LINC00638* was also decreased (Figure 4B). The

reduction of chromatin accessibility was consistent with the reduced expression of *AKT1* transcripts.

AKT1 is a key molecule in the PI3K-mTOR pathway[48], and E2F upregulates AKT activity through a

transcription-dependent mechanism[49]. *E2F8*, whose expression was downregulated by KDM6B

knockdown and GSK-J4 treatment, did not show significant changes in DNA accessibility within its

promoter region. However, several peaks at the distal region of E2F8 which extended to the adjacent

NAV2 locus were altered by GSK-J4 (Figure 4C). Notably, our RNA-seq analysis showed that the *NAV2*

locus is not expressed in BE2C cells (Figure 4C). These data suggest that *E2F8* expression might be

regulated by a distal enhancer that is silenced by KDM6B inhibition. Similarly, we did not observe

significant changes at the promoter regions of *MYCN*, and cell cycle genes (*CCNE1*, *BUB1* and *BIRC5*);

however, the DNA accessibility at the distal regions of these genes were reduced (Supplementary Figure

10A-10D). The downregulation of gene expression by GSK-J4 was not always correlated with reduction

of DNA accessibility. For example, chromatin accessibility at the *AURKB* promoter was increased by

GSK-J4 although its transcript expression was reduced (Supplementary Figure 10E), suggesting that the

open chromatin may recruit transcription suppressors to repress its gene transcription. For *CDKN1A*,

which encodes p21 to inhibit the CDK2/Cyclin E complex during S phase of the cell cycle, the DNA

accessibility at its promoter and enhancer regions was enhanced, consistent with its transcript induction

by KDM6B inhibition (Supplementary Figure 10F).

Transcription factor motif analysis of the ATAC-seq data using genomic footprinting

demonstrated that E2F binding motifs had reduced DNA accessibility (Figure 4D, 4E). Downregulated

open chromatin regions were also observed for MYCN and MYC/MAX binding motifs (Figure 4E). These

data indicate that KDM6B inhibition impacts the chromatin accessibility and transcription of E2F and

MYC pathway genes.

KDM6B inhibition induces an increase of H3K27me3 but a decrease of the enhancer mark H3K4me1 at CTCF and BORIS binding sites

To examine the impact of KDM6B inhibition on epigenetic marks, we performed Cleavage Under Targets

& Tagmentation (CUT&Tag) to assess the alterations of H3K27me3, the KDM6B substrate, and

H3K4me1, the enhancer mark, after KDM6B knockdown and GSK-J4 treatment of BE2C cells.

CUT&Tag is an epigenomic profiling strategy that overcomes shortcomings of ChIP-seq[50]. In CUT&Tag,

the H3K27me3 and H3K4me1 were bound in situ by specific antibodies, which were then tethered to a

protein A-Tn5 transposase fusion protein. Activation of the transposase efficiently generates fragment

libraries with high resolution and exceptionally low background[50]. The CUT&Tag library size for each

sample was adjusted by E. Coli residual genomic DNA as a spike-in scaling factor, which was used to

normalize the read counts for the downstream differential analysis. KDM6B inhibition induced a modest

increase of global H3K27me3 peaks (Supplemental Table 3, Supplemental Figure 11A). We specifically

assessed the H3K27me3 in the differentially expressed genes induced by KDM6B inhibition. Among the

genes differentially regulated by KDM6B knockdown and GSK-J4 treatment, the patterns of H3K27me3

changes tended to be similar in both groups (Figure 5A), suggesting that GSK-J4 mainly acts through

KDM6B to modulate the H3K27me3 levels. In particular, the H3K27me3 signals at the super-enhancer region of *MYCN* (the genomic region between *MYCN* and *FAM149A*, with a noncoding RNA *GACAT3* between both of them[23,51]) were most significantly upregulated by KDM6B knockdown or GSK-J4 treatment (Figure 5C, Supplemental Figure 11A), which is consistent with the downregulation of MYCN by KDM6B inhibition. We further assessed H3K4me1, the enhancer mark, using CUT&Tag after KDM6B knockdown and GSK-J4 treatment (Supplemental Table 4). Similar patterns of H3K4me1 changes were observed in both groups (Figure 5B). Interestingly, while more peaks of H3K4me1 were increased in genes downregulated by KDM6B inhibition, the super-enhancer region of *MYCN* showed most significant downregulation of H3K4me1 although GSK-J4 also induced an increase of several peaks at this region (Figure 5C, Supplemental Figure 11B), suggesting that MYCN super-enhancer region is particularly susceptible to perturbation of KDM6B inhibition.

The alterations of H3K27me3 and H3K4me1 by KDM6B inhibition mainly occurred at non-promoter regions (Supplemental Figure 11C), suggesting that KDM6B has a major impact on distal regulatory elements of transcription. Transcriptional factor motif analyses of H3K27me3 and H3K4me1 peaks revealed that KDM6B inhibition dominantly led to an increase of H3K27me3 but a decrease of H3K4me1 at the CTCF and BORIS binding sites (Figure 5D, E), and no enrichment with MYCN or E2F binding sites. CTCF and BORIS (also named CTCFL) play an important role in chromatin looping and long-range chromatin interactions[52-54]. H3K4me1 is an essential feature of enhancers and is bound by multiple chromatin-associated factors such as BAF complex and Cohesion[55], which facilitate enhancer-promoter looping[56,57]. Since the majority of upregulated H3K27me3 peaks were located at distal regulatory regions, we hypothesized that the H3K27me3 changes at these distal regions repressed transcription of *MYCN* and E2F target genes via alteration of chromatin structure and enhancer/promoter interactions. To investigate this possibility, we cross-referenced the Hi-C chromatin conformation of BE2C cells. Interestingly, *MYCN* is located at the same topologically associated domain (TAD) as the *GACAT3 and FAM149A* loci, marked as a super-enhancer by broad H3K27Ac occupancy, showing dysregulated H3K27me3 and H3K4me1 by KDM6B inhibition (Figure 5C). A TAD domain is a self-interacting genomic region that regulates gene expression by limiting the enhancer-promoter interaction to each TAD[58,59]. Chromatin interaction was observed across this region (Figure 5F), supporting the hypothesis that KDM6B inhibition may disrupt the chromatin interaction, leading to reduced transcription of *MYCN*. Similarly, we found the *E2F8* gene locus showed multiple chromatin interactions with adjacent genes within the TAD, particularly to the *NAV2* locus (Supplemental Figure 11D, E). Taken together,

these data suggest that KDM6B inhibition might alter the long-range interactions of *MYCN* and E2F target

genes in TAD, repressing the MYCN and E2F regulated transcriptome.

GSK-J4 reduces the expression of the PRC2 complex

The homeostasis of H3K27me3 is balanced by PRC2 and KDM6[5]. More H3K27me3 peaks were expected

by KDM6 inhibition. However, GSK-J4 treatment did not lead to expected high number loci of

H3K27me3 (Supplemental Figure 11A). To explain this discrepancy, we examined the expression of the

PRC2 complex. The key PRC2 components (EZH2, SUZ12 and EED) have been shown to be pRB-E2F

targets[60,61]. Consistent with the data that KDM6B inhibition mainly downregulated an pRB-E2F

transcriptome, GSK-J4 treatment indeed reduced the expression of the PRC2 complex (Supplementary

Figure 12A), However, the expression of global H3K27me3 was minimally affected, indicating that the

expected H3K27me3 upregulation by KDM6 inhibition is counterbalanced by reduced expression of

PRC2. A recent study identified a 37-gene signature of EZH2[28], which is repressed by EZH2 and silenced

in *MYCN*-amplified high-risk neuroblastoma. GSEA analysis revealed that this 37-gene signature was

significantly associated with the GSK-J4 treatment (Supplementary Figure 12B, 12C), suggesting that

KDM6B inhibition consequently downregulates the functions of PRC2, leading to the induction of tumor-

suppressive program repressed by EZH2.

Inhibition of KDM6B mimics CDK4/6 inhibitor Palbociclib

To further investigate the biological functions of KDM6B, we integrated the KDM6B downstream target

genes with information from the Library of Integrated Network-Based Cellular Signatures (LINCS) data.

LINCS data is composed of tens of thousands of gene sets, indicative of the transcriptional responses to a

large library of chemical compounds[62]. With this approach, we found that genes downregulated by

KDM6B knockdown in neuroblastoma cells (Supplementary Table S5) overlapped significantly with the

transcriptomes downregulated by palbociclib (Figure 6A), a specific CDK4/6 inhibitor that acts upstream

of the E2F pathway in cell cycle regulation. Similarly, genes downregulated by GSK-J4 (Supplementary

Table S6) also significantly overlapped the transcriptomes downregulated by palbociclib (Figure 6B).

CDK4/6 inhibitor signatures being identified among the most correlated signatures with KDM6B

knockdown or GSK-J4 treatment further demonstrated that KDM6B is mainly involved in regulation of

the E2F pathway. To corroborate these findings, we treated 4 different neuroblastoma cell lines (2 with

MNA and 2 without MNA) with palbociclib and performed RNA-seq for differential gene expression

analyses. By comparing the genes most significantly downregulated by palbociclib among these cell lines

via Venn program (Supplementary Figure 13), we extracted 89 common genes (Supplementary Table S7).

GSEA results showed that the 89-gene signature was significantly enriched within gene sets

downregulated by KDM6B knockdown (Figure 6C) or GSK-J4 treatment (Figure 6D). Although single

drug treatment effectively inhibited colony formation in a dose dependent manner, the combination of

GSK-J4 and palbociclib did not enhance the inhibitory effect on colony formation of BE2C cells, as shown

by colony formation and Bliss combination index (Figure 6E, 6F), likely because both drugs target the

same pathway. However, GSK-J4 greatly enhanced the effect of 17-DMAG, an HSP90 inhibitor

(Supplementary Figure 14A, 14B), and JQ-1, a bromodomain inhibitor that targets BRD4[63]

(Supplementary Figure 14C, 14D). Interestingly, we observed some degree of synergy of GSK-J4 and

palbociclib, as well as 17-DMAG and JQ-1 in SK-N-AS cells (Supplementary Figure 14E-G), the non-

MYCN amplified cells with less impact on E2F pathways by KDM6B inhibition.

396 Cancer cells with acquired CDK4/6 inhibitor resistance are less responsive to GSK-J4

Upon mitogen stimulation, CDK4 and CDK6 phosphorylate the pRB protein, leading to release of E2F

transcription factors from pRB for gene transcription[64]. CDK4/6 inhibitors including palbociclib showed

promising efficacy in ER$^+$/HER2$^-$ breast cancer and have been approved for clinical use[65,66]. Over 100

clinical trials of CDK4/6 inhibitors for a variety of cancers are currently in progress[65]. However, acquired

resistance to CDK4/6 inhibitors, as observed during breast cancer treatment, is expected and the

mechanism involves amplification of *CDK6* or loss of function of *Rb1*[65,66]. To test if these mechanisms

also diminish the effect of GSK-J4, we transduced the BE2C cells with lentiviral based CDK4 and CDK6

(Figure 6G). Overexpression of CDK4 or CDK6 conferred expected resistance in neuroblastoma cells to

palbociclib, as shown by colony formation and EC50 shift (Figure 6H, I). Cells which overexpressed

CDK4 or CDK6 also gained resistance to GSK-J4, although a higher concentration of GSK-J4 at 2.5μM

was still able to kill them (Figure 6H, 6J). We then generated an *Rb1* knockout cell line for testing the

response to GSK-J4 and palbociclib (Figure 6K). As expected, loss of *Rb1* conferred resistance to

palbociclib (Figure 6L). Similarly, the *Rb1* knockout cells were also resistant to GSK-J4 treatment in

comparison with the *Rb1* wildtype cells (Figure 6M). Similar to the CDK4 or CDK6 overexpressing cells,

the combination of palbociclib and GSK-J4 was not synergistic (Figure 6M). These data further

demonstrate that KDM6B acts on the same pathway as pRB-E2F in neuroblastoma.

414 A gene signature targeted by KDM6B inhibition is associated with poor outcome

To correlate the clinical relevance of target genes of KDM6B inhibition, we identified a 149-gene

signature (Supplementary Table 8) that was commonly downregulated by GSK-J4 in BE2C and three

other neuroblastoma cell lines (IMR5, LAN5 and SK-N-FI)[43]. Among these 149 genes, 85 were highly

expressed in the high-risk neuroblastomas while 20 were highly expressed in the low-risk neuroblastomas

(Supplementary Table 9). Based on the expression levels of these differentially expressed genes in

neuroblastomas, they were categorized into 4 clusters (Figure 7A). Cluster 3 and 4 neuroblastomas

expressed higher levels of the GSK-J4 signature, and were enriched with high-risk, high-stage and

MYCN-amplified tumors. While patients in clusters 1 and 2 showed excellent event-free and overall

survival, clusters 3 and 4 patients had a significantly worse outcome (Figure 7B, 7C). No difference was

observed between clusters 1 and 2. Cluster 4 showed a poorer survival than that of cluster 3, in line with

the higher expression of GSK-J4 target genes in cluster 4. *MYCN* amplification (MNA) is a well-known

poor prognostic risk factor in neuroblastoma. To investigate whether the correlation of the GSK-J4

signature with patient outcome was affected by MNA status, we compared the event-free survival of

clusters 3 and 4 with MNA. The results showed that cluster 4, which had higher levels of GSK-J4

signature, was still correlated with a poorer event-free survival of patients with MNA although the overall

survival was not statistically significant (Figure 7D, 7F). However, in stage 4 patients, cluster 4 showed

significantly worse outcome in both event-free and overall survival than that of cluster 3 (Supplementary

Figure 15). In high-risk patients, cluster 4 had a significantly worse outcome in event-free survival

compared to cluster 3, but there was no difference in overall survival (Figure 7E, G). These data indicate

that, in stage 4 or MNA patients, higher expression levels of the GSK-J4 signature is a high-risk factor.

437 DISCUSSION

Despite one recent study suggesting that KDM6B might be tumor suppressive in neuroblastoma[67], we and

others have found that KDM6B plays an important role in MYC-driven tumorigenesis and that

pharmacologically targeting KDM6B by the small molecule GSK-J4 is therapeutically efficacious in

multiple tumor models[14,27,68-71], including high-risk neuroblastoma[43]. However, the anticancer

mechanisms of KDM6B inhibition are poorly understood. Here we show that *KDM6B* is significantly

over-expressed in human primary neuroblastoma in comparison with its paralogs *KDM6A* and *UTY*.

Strikingly, the 22-kb genomic locus of *KDM6B* is heavily occupied by active gene transcription marks,

including H3K4me1, H3K27Ac, BRD4 and RNA polymerase II binding while the *KDM6A* and *UTY* loci

lack such histone modifications and transcription factor binding. These data explain why *KDM6B* is

expressed at much higher levels than *KDM6A* and *UTY* in neuroblastoma. The long-distance broad marks

at the KDM6B locus with H3K4me1 and H3K27Ac suggest *KDM6B* expression is under the control of a

super-enhancer, a large cluster of transcriptional enhancers that drive expression of genes that define cell

identity[72]. We further found that KDM6B is predominantly involved in regulation of MYC expression and

downstream target genes of the pRB-E2F pathway. Genetic and pharmacologic inhibition of KDM6B

consistently repressed the expression of MYCN and C-MYC, and the transcriptome of E2F target genes.

These results support previous studies demonstrating that E2F and MYC are functionally associated[38,39],

and that E2F drives MYC expression. MYC also regulates E2F expression and requires distinct E2F

activities to induce S phase and apoptosis[73]. Moreover, a close association between E2F and MYC binding

sites and their target genes has been observed[74]. Nevertheless, the impact of KDM6B inhibition on E2F

pathways was more dramatic in MYCN-amplified neuroblastoma cells.

E2F deregulation in cancer often occurs through loss-of-function of the pRB tumor suppressor[75].

The E2F transcription factors (E2F1, E2F2, and E2F3a) bind at target gene promoters in complexes with

their dimerization partner (DP1 or DP2) and pRB[36], which suppresses target gene transcription through

the recruitment of chromatin modifiers and remodeling factors such as HDAC and EZH2[36,76]. During the

G1 phase of the cell cycle, oncogenes such as Ras induce D-type cyclins to activate cyclin dependent

serine/threonine kinases CDK4 and CDK6, which phosphorylate pRB, consequently leading to release of

E2F transcription factors from pRB, allowing for cell proliferation[64]. While *Rb1* loss is very rare in

neuroblastoma, several studies have shown that the cyclin D/CDK4/CDK6 pathway is hyperactive in

neuroblastoma[77-82]. Genome-wide CRISPR and shRNA screening demonstrated that knockout of *CDK4*

or CDK6 inhibited neuroblastoma cell proliferation/survival while *Rb1* knockout promoted cell

proliferation (https://depmap.org/portal/). A majority of neuroblastoma models with MNA are sensitive

to CDK4/6 inhibition[83]. These data indicate that the deregulated E2F pathway is essential to

neuroblastoma. The genetic and pharmacologic KDM6B inhibition results in a predominant reduction of

the E2F transcriptome and MYC, which may account for the therapeutic effect of KDM6B blockade in

neuroblastoma. In addition, the transcriptome of KDM6B inhibition mimics the CDK4/6 inhibitor,

palbociclib. When cells gained resistance to palbociclib by overexpressing CDK4/6 or by *Rb1* knockout,

they also gained resistance to GSK-J4. Thus, these data further support that KDM6B regulates the

CDK4/6-pRB-E2F pathway in neuroblastoma.

Intriguingly, while transcriptome, chemical genetics and ATAC-seq analyses show that

KDM6B inhibition impacts MYCN and the E2F transcriptome, the loci of H3K27me3 alterations were

not enriched with MYCN and E2F genes. As the majority of enhancer mark H3K4me1 and H3K27me3

peaks altered by KDM6B inhibition were located at distal non-promoter regions, one possible explanation

is that KDM6B inhibition disrupted the long-range chromatin interaction between its targets and the

regions with altered H3K27me3 and H3K4me1 because H3K4me1 functions to mediate the enhancer-

promoter looping[56,57]. H3K4me1 is basically catalyzed by KMT2 family methyltransferases. Previous

studies have shown that KMT2 members can form complexes with KDM6[84-86], suggesting a concerted

mechanism for transcriptional activation in which cycles of H3K4 methylation by KMT2 are linked with

the demethylation of H3K27[84-86]. The transcriptional factor motif analysis showed that CTCF and BORIS

binding motifs are the major sites impacted by altered H3K27me3 and H3K4me1. KDM6B inhibition

specifically leads to increased H3K27me3 at CTCF and BORIS binding sites while H3K4me1 at CTCF

and BORIS binding sites are selectively reduced. These data suggest that increased H3K27me3 displaces

H3K4me1 modifiers (highly likely the H3K4me1 methyltransferase KMT2) from CTCF and BORIS sites,

consequently disrupting the enhancer activity. Previous studies have shown that enhancers, CTCF and

H3K4me1 peaks overlap[87]. CTCF regulates the long-range chromatin interactions at enhancers. Thus,

KDM6B inhibition may disrupt the chromatin interaction at its target genes. *MYCN* and many E2F target

genes reside in the same topologically associated domains (TAD) as the regions with elevated H3K27me3

by KDM6B inhibition. TADs include chromatin loops that mediate promoter–enhancer contacts that

regulate gene expression[58,59]. Our CUT&Tag data showed that the super-enhancer region of *MYCN*, which

exhibits long-range chromatin interactions within a TAD domain, has the most significant changes of

H3K27me3 and H3K4me1 by KDM6B knockdown and GSK-J4 treatment, in line with the reduction of

MYCN expression by KDM6B inhibition. In erythroid cells, long-range control of epigenetic regulation

has been observed in that KDM6B is recruited to the enhancer regions to erase H3K27me3, consequently

evicting the gene silencing PcG protein for a high rate of transcription[88]. During early differentiation steps,

the embryonic stem cell factor Tbx3 associates with KDM6B at the enhancer element of the *Eomes* locus

to allow enhancer-promoter interactions[89]. This spatial reorganization of the chromatin primes the cells to

respond to Activin signaling. In neural stem cells, SMAD3 recruits KDM6B at the enhancers in response

to TGF-β, for enhancer transcription[90]. Here we propose a model that, once stimulated by mitogens due

to oncogenic activity, KDM6B is recruited to chromatin to maintain the low levels of H3K27me3 at the

distal regulatory enhancer regions bound by CTCF and BORIS, which loops and physically interacts with

E2F transcription factors that bind at the promoters of target genes, together with associated transcriptional

machinery including RNA polymerase II and transcription mediators, driving the expression of *MYCN*

and the E2F transcriptome (Figure 8A). When KDM6B is inhibited, the H3K27me3 accumulates at the

distal regions and disrupts the assembly of transcriptionally active interaction of promoter-enhancers,

leading to the downregulation of *MYCN* and the E2F transcriptome (Figure 8B).

Interestingly, expression of the PRC2 complex, which is a downstream target of pRB-E2F[60,61]

and MYCN[28], was downregulated by KDM6B inhibition, in line with of the decrease of a number of

H3K27me3 peaks and de-repression of a tumor suppressive program governed by EZH2[28]. The connection

between KDM6B and EZH2 is interesting. Since EZH2 is a downstream target of E2F and MYCN, we

assumed the reduction of EZH2 by KDM6B inhibition is through this mechanism. However, considering

both proteins are epigenetic modifiers that confer cellular plasticity, we also speculate that cells strive for

adaptation to the KDM6B inhibition by suppressing EZH2 in order to counterbalance the net increase of

H3K27me3.

Neuroblastoma is a disease that arises as a result of blocked differentiation of neural crest

precursors (NCCs) during development[19,20]. EZH2 is required to maintain the undifferentiatied state of

neuroblastoma[28,91]. Thus, our data and that from other studies indicate that a network composed of MYCN,

E2F, PRC2 and KDM6B regulates cell proliferation and differentiation of neuroblastoma (Figure 8C),

highlighting the importance of KDM6B in coupling the two essential features of cancer cells. In summary,

we have defined a novel chromatin-dependent mechanism of action of KDM6B inhibition that modulates

the CDK4/6-pRB-E2F pathway in neuroblastoma. We also demonstrate that E2F target genes can act as

biomarkers for sensitivity to KDM6B inhibitors in neuroblastoma and that this may well be predictive for

other MYC-driven tumors.

MATERIALS AND METHODS

Cell Culture and Reagents

Neuroblastoma cell lines BE2(C) (ATCC), SIMA (DSMZ, Germany), SKNDZ (ATCC), SK-N-AS

(ATCC), SK-N-SH (ATCC), IMR32 (ATCC), CHLA20 (COG), NB-1691 (Peter Houghton), HS68

(ATCC) were cultured in standard RPMI media supplemented with 10% FBS (Sigma), 1% L-glutamine

(MediaTech), at 37 °C in 5% CO_2. Colorectal cancer cell lines HCT116, and isogenic $p53^{-/-}$ HCT116 cells

were kindly provided by Bert Vogelstein (Baltimore, MD), osteosarcoma cell line U2OS, lung cancer cell

lines A549 purchased from ATCC, and they were cultured in standard DMEM media supplemented with

10% FCBS, 1% L-glutamine (MediaTech), at 37 °C in 5% CO_2. Cell lines were validated by short tandem

repeat (STR) using Promega PowerPlex 16 HS System once per month. PCR-based method was used for

detection of Mycoplasma with LookOut Mycoplasma PCR Detection Kit (Sigma) and JumpStart Taq

DNA Polymerase (Sigma) once per month to ensure cells were mycoplasma negative. GSK-J4,

Palbociclib, 17-DMAG, JQ-1 were purchased from Selleckchem. KDM5-C70 was purchased from Xcess

Biosciences Inc.

AlphaLISA

Materials: GSK J1 is purchased from Tocris (Bristol, United Kingdom, Catalog number 4593). 2,4-

Pyridine Dicarboxylic Acid (2,4-PCA) is purchased from Acros Organics (New Jersey, Catalog number

101860010). JIB-04 is purchased from Sigma-Aldrich (St. Louis, MO, Catalog number SML0808).

Daminozide is purchased from Tocris (Bristol, United Kingdom, MO, Catalog number 4684). AlphaLISA

anti-mIgG acceptor beads from PerkinElmer (Santa Clara, CA, Catalog number AL105C). AlphaLISA

anti-rIgG acceptor beads from PerkinElmer (Santa Clara, CA, Catalog number AL104C). AlphaScreen

Streptavidin-conjaugated donor beads from PerkinElmer (Santa Clara, CA, Catalog number 6760002).

Primary antibody 5 from BPS (Catalog number 52140E). Primary antibody 6 from BPS (Catalog number

52140F). Primary antibody 13-4 from BPS (Catalog number 52140M4). Primary antibody 16-2 from BPS

(Catalog number 52140P-2). Primary antibody 17-4 from BPS (Catalog number 52140Q4). Biotinylated

histone H3 peptide substrate (KDM4A) BPS (Catalog number 79841). Biotinylated histone H3 peptide

substrate (KDM5A) BPS (Catalog number 79840). Biotinylated histone H3 peptide substrate (KDM2A)

BPS (Catalog number 79843). Biotinylated histone H3 peptide substrate (KDM6A and KDM6B) BPS

(Catalog number 79841). Biotinylated histone H3 peptide substrate (KDM3A) BPS (Catalog number

79841). 4x HDM Assay Buffer 2 from BPS (Catalog number 52407). 4x HDM Assay Buffer 3 from BPS

(Catalog number 52408). 4x HDM Assay Buffer 4 from BPS (Catalog number 52409). 4x HDM Assay

Buffer 5 from BPS (Catalog number 79847). 4x Detection buffer from BPS (Catalog number 52301).

Assay Conditions: All of the enzymatic reactions were conducted in duplicate at room temperature for 60

minutes in a 10 µl mixture containing assay buffer, histone H3 peptide substrate, demethylase enzyme,

and the test compound.   These 10 µl reactions were carried out in wells of 384-well Optiplate

(PerkinElmer). The dilution of the control compounds was first performed in 100 % DMSO. Each

intermediate compound dilution (in 100 % DMSO) will then get directly diluted 30x fold into assay buffer

for 3.3x conc (DMSO). Enzyme only and blank only wells have a final DMSO concentration of 1 %.

From this intermediate step, 3 µl of compound is added to 4 µl of demethylase enzyme dilution is

incubated for 30 minutes at room temperature. After this incubation, 3 µl of peptide substrate is added.

The final DMSO concentration is 1 %.   After enzymatic reactions, 5 µl of anti-Mouse Acceptor beads

(PerkinElmer, diluted 1:500 with 1x detection buffer) or 5 µl of anti-Rabbit Acceptor beads (PerkinElmer,

diluted 1:500 with 1x detection buffer) and 5 µl of Primary antibody (BPS, diluted 1:200 with 1x detection

buffer) were added to the reaction mix. After brief shaking, plate was incubated for 30 minutes. Finally,

10 µl of AlphaScreen Streptavidin-conjugated donor beads (Perkin, diluted 1:125 with 1x detection buffer)

were added. In 30 minutes, the samples were measured in AlphaScreen microplate reader (EnSpire Alpha

2390 Multilabel Reader, PerkinElmer).

Data Analysis: Enzyme activity assays were performed in duplicates at each concentration. The A-screen

intensity data were analyzed and compared. In the absence of the compound, the intensity (Ce) in each

data set was defined as 100 % activity. In the absence of enzyme, the intensity (C0) in each data set was

defined as 0 % activity. The percent activity in the presence of each compound was calculated according

to the following equation: % activity = (C-C0)/(Ce-C0), where C = the A-screen intensity in the presence

of the compound. The values of % activity were plotted on a bar graph.

SDS-PAGE and Western blotting

For western blotting, samples were mixed with calculated volume of 2X sample buffer (1M TRIS/HCl,

10% SDS, 0.1% bromophenol-blue, 10% b−mercaptoethanol, 10% glycerol), and heated for 25 minutes

at 96°C. Proteins were resolved on SDS-polyacrylamide gels (SDS-PAGE) and transferred onto PVDF

membrane (Immobilon-P, Millipore). After being incubated with the primary antibody, horseradish

peroxidase-(HRP) conjugated secondary antibody (Novex, Life technologies) at 1: 5000 was used for 1

hour incubation. The signals were detected by chemiluminescence (ECL, Thermo scientific). Antibodies

including H3K4me1(Abcam Cat# ab8895, RRID:AB_306847), H3K4me3 (RevMAb Biosciences Cat#

31-1226-00, RRID:AB_2783580), H3K27ac (Active Motif Cat# 39133, RRID:AB_2561016),

H3K27me3 (Cell Signaling Technology Cat# 9733, RRID:AB_2616029), CDK4 (Cell Signaling

Technology Cat# 12790, RRID:AB_2631166), CDK6 (Cell Signaling Technology Cat# 13331,

RRID:AB_2721897), β-ACTIN (Sigma-Aldrich Cat# A1978, RRID:AB_476692), GAPDH (Cell

Signaling Technology Cat# 5174, RRID:AB_10622025), pRB (4H1) (Cell Signaling Technology Cat#

9309, RRID:AB_823629), MYCN (Cell Signaling Technology Cat# 9405, RRID:AB_10692664) or

(Santa Cruz Biotechnology Cat# sc-53993, RRID:AB_831602), C-MYC (Cell Signaling Technology Cat#

13987, RRID:AB_2631168), KDM6B (Abclonal, A12763), H3K9me3 (Active Motif, 39161),

H3K36me3 (Abcam, ab9050), were used for western blot.

Crystal Violet Staining

After removing media, cells were washed with Dulbecco's phosphate buffered saline without calcium or

magnesium (DPBS, Lonza) and treated with 4% Formaldehyde in PBS (PFA) for 20 minutes. Once PFA

was removed, cells were stained with 0.1% crystal violet stain for 1 hour.

Synergy Assay

BE2C and SK-N-AS cells were seeded in 96-well plates (3,000 cells per well). After 24 hours, cells were

treated with GSK-J4 (0, 0.25, 0.5, 1 µM, for BE2C, 0, 0.5, 1, 2.5 µM for SK-N-AS) and palbociclib (0,

0.25, 0.5, 1.0, 2.0, 4.0 µM) or JQ1 (0, 0.0625, 0.125, 0.25, 0.5, 1.0 µM) or 17-DMAG (0, 0.0625, 0.125,

0.25, 0.5, 1.0 µM) in a 4x6 matrix. Every combination treatment was performed in quadruplicate. Cells

were treated for 5 days, and cell viability was determined using the Prestoblue assay (Invitrogen, A-

13262). Cell viability for each treatment was normalized against the control group. A Bliss independence

model was used to evaluate combination effects. Percentage over the Bliss score index was calculated

with the equation (A+B)-AxB, in which A and B are the percentage of growth inhibitions induced by

agents A and B at a given dose, respectively. The difference between the Bliss expectation and the

observed growth inhibition induced by the combination of agent A and B at the same dose is the Bliss

excess.

Cell viability assay

BE2C, and CDK4 or CDK6 overexpressing BE2C (BE2C-CDK4-OE, BE2C-CDK6-OE) cells were

seeded in 96-well plates at 3,000 cells per well. After 24 hours, cells were treated with GSK-J4 (0, 0.02,

0.04, 0.08, 0.16, 0.3125, 0.625, 1.25, 2.5, 5.0,10 µM) and palbociclib (0, 0.02, 0.04, 0.08, 0.16, 0.3125,

0.625, 1.25, 2.5, 5.0, 10 µM). Cells were treated for 5 days. Cell viability was determined using the

Prestoblue assay (Invitrogen, A-13262). Cell viability for each treatment was normalized against no

treatment well. EC50 was determined with GraphPad Prism equation Y=100/(1+10^((LogEC50-

X)*HillSlope)), EC50 is the concentration that gives a 50% response. HillSlope represents the steepness

of the curve.

Small interfering RNA Transfection

Small interfering RNAs (siRNA) were transfected into subconfluent cells using Lipofectamine RNAiMax

(Invitrogen) according to manufacturer's instructions. Non-Targeting siRNA#1 (Dharmacon, D-001810-

0105) used as siRNA control. The siRNA oligos for KDM6B have sequences as follows: KDM6B#1, 5-

GGAAUGAGGUGAAGAACGU-3, KDM6B#2, 5-GGAGACCUCGUGUGGAUUA-3, KDM6B#31, 5-

GCAUCUAUCUGGAGAGCAA-3, KDM6B#33, 5-GGAAGAGGAACAGCAACA-3. KDM6B-

3'UTR, 5-AGAAAGAACUAUGAGGAAAUU-3.

CRISPR knockout of Rb1

The Rb1 CRISPR plasmid with gRNA sequence 5-GCTCTGGGTCCTCCTCAGGA-3 (TLCV2-RB1,

Addgene#87836) was purchased from Addgene. Plasmids were maxiprepped by using NucleoBond Xtra

EF kits (Takara Bio USA, 740424-50) according to manufacturer's protocol. Lentivirus was produced by

transient transfection of PEI-pro DNA complex (6µg of TLCV2-RB1, 3 µg of 1-1r, 1 µg RTR, 1 µg of

VSVg with 22 ul of PEI pro in 400 µl of DMEM medium) with 5×10^6 HEK293T cells in 10 ml complete

medium (DMEM, 100 U/mL penicillin/streptomycin, 1 x L-glutamine and 10% FBS) in a 10 cm dish.

Virus supernatant was collected every 8-12 hours for 3 days, which were passed through a 0.45 µm filter

and concentrated by ultracentrifuge at 28,500 rpm for 1.5 hours at 4°C. The TLCV2-RB1 virus particles

were added to BE2C cells with polybrene to final concentration of 8 µg/ml. Puromycin (2.5 µg/ml in

complete medium) selection were performed in the next day after virus transduction. To generate Rb1

knockout in BE2C, BE2C-Rb1 cells were treated with 1µM of doxycycline for 72 h, then sorted for green

fluorescent protein (GFP) positive cells to enrich for RB1 knockout. The sorted cells were expanded

without doxycycline in RPMI 1640 media (Corning, 10-040-CM) supplemented with 100 U/mL

penicillin/streptomycin (Gibco, 15140122), and 10% FBS (Sigma-Aldrich, F2442).

Generation of KDM6B, CDK4 and CDK6 overexpressing cell lines

The CDK4 plasmid (pHAGE-CDK4, Addgene#116724) and CDK6 plasmid (pHAGE-CDK6,

Addgene#116725) were purchased from Addgene. The plasmid maxiprep and lentiviral packaging were

followed by the same protocol as described in Rb1 knockout. To overexpress CDK4 and CDK6 in BE2C

cells, the pHAGE-CDK4 or pHAGE-CDK6 viral particles were transduced to BE2C cells and sorted for

green fluorescent protein (GFP) positive ones. To overexpression KDM6B, MSCV-JMJD3

(Addgene#21212) were packaged into retroviral particles, which were transduced into BE2C cells and

selected with puromycin for stable expression.

RNA-seq and microarray

Total RNA was extracted from cells by using RNeasy Mini Kit (cat. # 74104) from QIAGEN. Paired-end

sequencing was performed using the High-Seq platform with 100bp read length. Reads were aligned to

the human GRCh37-lite using SJCRH's Strongarm pipeline. Counts per gene were obtained using htseq-

count version 0.6.1 with Gencode vM5 level 1 and 2 gene annotations. Counts were normalized with

VOOM and analyzed with LIMMA within the R statistical environment. Significance was defined as

having a false discovery rate (FDR) <0.05. VOOM normalized counts were analyzed with Gene Set

Enrichment Analysis (GSEA)[92]. For Affymetrix microarray, after quality control with Agilent RNA

analyzer, RNA was subjected to hybridization using an Affymetrix Clariom S human array. Differential

gene expression was analyzed by t-test using the Differential Expression Analysis

module at GenePattern server (http://genepattern.broadinstitute.org/gp/pages/protocols/DiffExp.html).

Assay for Transposase-Accessible Chromatin using sequencing (ATAC-seq)

Library preparations for ATAC-seq were based on the protocol with minor modifications[47,93]. Briefly,

fresh cultured BE2C cells (100,000 per sample) with or without 2.5μM of GSK-J4 treatment were

harvested and washed with 150μl cold Dulbecco's Phosphate-Buffered Saline (DPBS) containing protease

inhibitor (PI). Nuclei were collected by centrifuging at 500g for 10 minutes at 4°C after cell pellets were

resuspended in lysis buffer (10 mM Tris-Cl pH 7.4, 10 mM NaCl, and 3 mM MgCl2 containing 0.1% NP-

40 and PI). Nuclei were incubated with Tn5 transposon enzyme in transposase reaction mix buffer

(Illumina) for 30 min at 37°C. DNAs were purified from transposition sample by using Min-Elute PCR

purification kit (Qiagen, Valencia, CA) and measured by Qubit. Polymerase chain reaction (PCR) was

performed to amplify with High-Fidelity 2X PCR Master Mix [72°C/5mins+ 98 °C /30 s +12 × (98 °C /10

s + 63 °C /30 s + 72 °C /60 s) + 72 °C /5 min]. The libraries were purified using Min-Elute PCR

purification kit (Qiagen, Valencia, CA). ATAC-seq libraries followed by pair-end sequencing on

HiSeq4000 (Illumina) in the Hartwell Center at St Jude Children's Research Hospital.

ATAC-seq data analysis

2x100bp paired-end reads obtained from all samples were trimming for Nextera adapter by

cutadapt(version 1.9, paired-end mode, default parameter with " -m 25 -O 6 ")[94] and aligned to human

genome hg19(GRCh37-lite) by BWA (version 0.7.12-r1039, default parameter)[95] , duplicated reads were

then marked with Picard(version 2.6.0-SNAPSHOT)[96] and only non-duplicated proper paired reads have

been kept by samtools (parameter "-q 1 -F 1804" version 1.2)[97]. After adjustment of Tn5 shift (reads were

offset were offset by +4 bp for the sense strand and -5 bp for the antisense strand) we separated reads into

nucleosome free, mononucleosome, dinucleosome, trinucleosome as described in Buenrostro et.al[98] by

fragment size and generated bigwig files by using the center 80bp of fragments and scale to 20M

nucleosome free reads. We observed reasonable nucleosome free peaks and pattern of mono-, di-, tri-

nucleosome on IGV (version 2.4.13)[99]. Next we merged each 2 replicates to enhance peak calling on nucleosome free reads by MACS2(version 2.1.1.20160309 default parameters with "--extsize 200 --nomodel ")[100], all cell types have more than 20M nucleosome free reads after merge so we consider all important nucleosome free regions were called. To assure the replicability, we first merge peaks from different treatmeans to create a set of reference chromatin accessible regions. We then counted nucleosome free reads from each of samples overlap the reference regions by bedtools (v2.24.0)[101]. We conclude the reproducibility is good since spearman correlation coefficient between replicates are larger than between samples from different groups. To find the differential accessible regions, we first normalized raw nucleosome free reads counts used trimmed mean of M-values normalization method(TMM) and applied Empirical Bayes Statistics test after linear fitting from voom packge(R 3.23, edgeR 3.12.1, limma 3.26.9)[102]. FDR-correct p-value 0.05 and fold change > 2 were used as cutoff for differential accessible regions (DARs) while p-value > 0.5 and fold change < 1.05 were used for control regions. To find the Transcription factor enriched for DARs, we scanned the TRANSFAC [103] motif database using FIMO(parameter "--motif-pseudo 0.0001 --thresh 1e-4") from MEME suite(v4.11.3)[104], then for each motif, we counted how many DARs or control regions have the motif matches and using Fisher exact test to estimate their enrichment over the background(DAR or control regions don't have the motif matches).

CUT&Tag and analysis

CUT&Tag for KDM6B knockdown and GSK-J4 treatment was prepared by following the protocol as described previously (Kaya-Okur et al. 2019 and https://www.protocols.io/view/bench-top-cut-amp-tag-bcuhiwt6?step=1) with minor modifications. Approximately 500,000 BE2C cells were treated with siRNAs (siCtrl or siKDM6B) or compounds (DMSO 0.1% or GSK-J4 2.5µM) for 72 hours, followed by washing with wash buffer (20 mM HEPES pH 7.5; 150 mM NaCl; 0.5 mM Spermidine; 1× Protease inhibitor cocktail). Nuclei were isolated with cold NE1 buffer (20 mM HEPES–KOH, pH 7.9; 10 mM KCl 0.1%; Triton X-100; 20% Glycerol, 0.5 mM Spermidine; 1x Protease Inhibitor) for 10 min on ice. Nuclei were collected by 600 x g centrifuge and resuspended in 1ml washing buffer containing with 10 µL of activated concanavalin A-coated beads (Bangs laboratories, BP531) at RT for 10 min. Bead-bound nuclei were collected with placing tube on magnet stand and removing clear liquid. The nuclei bound with bead were resuspended in 50 µL Dig-150 buffer (20 mM HEPES pH 7.5; 150 mM NaCl; 0.5 mM Spermidine; 1× Protease inhibitor cocktail; 0.05% Digitonin; 2 mM EDTA) and incubated with a 1:50 dilution of H3K4me1 (Abcam, ab8895) and H3K27me3 (CST, 9733S) antibodies overnight at 4 °C. The

unbound primary antibody was removed by placing the tube on the magnet stand and withdrawing the
liquid. The primary antibody bound nuclei bead was mixed with 100uL of Dig-150 buffer containing
guinea pig anti-Rabbit IgG antibody (Antibodies, ABIN101961) in 1:100 dilution for 1 hour at RT. Beads
bound nuclei were washed using the magnet stand 3× for 5 min in 1 mL Dig-150 buffer to remove unbound
antibodies. A 1:100 dilution of pA-Tn5 adapter complex was prepared in Dig-300 buffer (20 mM HEPES,
pH 7.5, 300 mM NaCl, 0.5 mM Spermidine, 0.05% Digitonin, 1× Protease inhibitor cocktail). After
removing the liquid on the magnet stand, 100 µL mixture of pA-Tn5 and Dig- 300 buffer was added to
the nuclei bound beads with gentle vortex and incubated at RT for 1 h. After 3× 5 min in 1 mL Dig-300
buffer to remove unbound pA-Tn5 protein, nuclei were resuspended in 250 µL Tagmentation buffer
(10 mM MgCl2 in Dig-300 buffer) and incubated at 37 °C for 1 h. 10 µL of 0.5 M EDTA, 3 µL of 10%
SDS and 2.5 µL of 20 mg/mL Proteinase K were added to stop tagmentation and incubated at 55 °C for 1
hour. DNA was then precipitated by phenol/chloroform/isoamylalcohol followed by ethanol precipitation
with glycogen and then dissolved in water. Sequencing libraries were prepared using NEBNext HiFi 2×
PCR Master Mix (NEB, M0541L) according to the manufacturer's instructions. The PCR products were
cleaned up with SPRIselect beads and quantified using Qubit dsDNA HS assay kit (Agilent Technologies).
The libraries were sequenced on a HiSeq2500 with paired-end 50-bp reads (Illumina).
Mapping reads and peak calling. The Cut&Tag raw reads were aligned to the human reference genome
(hg19) using BWA (version 0.7.12; BWA aln+sampe). Duplicate reads were marked and removed by
Picard (version 1.65). Only properly paired uniquely mapped reads with a fragment size of 150bp-2000bp
were extracted by samtools (version 1.3.1 parameters used were -q 1 -f 2 -F 1804) for calling peaks and
generating bigwig file. Narrow peaks were called by MACS2 (version 2.2.7.1) with parameters of " -t
cut_tag_file -q 0.05 -f BED --keep-dup all" for CUT&Tag data).We used SICER (version 1.1, with
parameters of redundancy threshold 1, window size 200bp, effective genome fraction 0.86, gap size
600bp, FDR 0.00001 with fragment size defined above) for calling broad enriched regions. Unmapped
reads were further aligned to E.Coli genome as a spike-in control to calculated ChIP-Rx scaling factor as
described[105,106]. The Cut&Tag library size of each sample was adjusted by E.Coli spike-in scaling factor
and then used to normalize the read count for the downstream differential analysis, IGV bigwig and
heatmap visualization.
Visualization. We used genomeCoverageBed (BEDtools 2.25.0) to produce genome-wide coverage in
BEDGRAPH file and then converted it to a bigwig file by bedGraphToBigWig. The bigwig signals were
adjusted by E.Coli spike-in scaling factor and scaled to 15 million reads to allow comparison across
samples. To show average of several replicates as a single track in the browser, the bigwig files were

merged to a single average bigwig file using UCSC tools bigWigtoBedGraph, bigWigMerge and bedGraphToBigWig. The Integrated Genomics Viewer (IGV 2.3.82) was used for visual exploration of data. DeepTools (version 2.3)[107] was used to plot heatmap.

Differential analysis. Cut&Tag raw read counts were reported for each region/each sample using BEDtools 2.25.0. Raw read counts were Voom normalized and statistically contrasted using the R (version 3.5.1) packages limma and edgeR (version 3.16.5) for CPM calculation and differential analysis. An empirical Bayes fit was applied to contrast treated samples to control samples and to generate log fold changes, p values and false discovery rates for each peak region.

Peak overlap and annotation. Peak regions were defined to be the union of peak intervals in replicates from control or treated cells respectively. For peak overlap analysis, mergeBed (BEDtools version 2.25.0) was used to combine overlapping regions from multiple peak sets into a new region and then a custom script was used to summarize common or distinct peaks and visualize in a venn diagram. Promoter regions was defined as the regions 1.0 kb upstream and 1.0 kb downstream of the transcription start sites based on the human RefSeq annotation (hg19). Genomic feature annotation of peaks was done by annotatePeaks.pl, a program from the HOMER suite (v4.8.3, http://homer.salk.edu/homer/).

Motif analysis. Adjacent histone mark peaks (within a distance of 100bp) from both control and treated samples were merged first and then separated to 3 subgroups including control-only, treatment-only and shared peaks (at least 1bp overlap) using Bedtools (version 2.25.0). The regions of peak center +/- 100 bp were used for HOMER known motif search. Top ten most significant motifs from each sub-group were visualized as a heatmap.

Hi-C data mining

To view the 3D chromatin conformation of *MYCN* and *E2F8* gene in BE2C, we mined the Hi-C data, of BE2C, KELLY and SK-N-AS cells, whic were downloaded from St Jude Cloud (https://www.stjude.cloud/)[108]. The Arc TRAC data showing the chromatin interactions, which were generated from Jurkat ChIA-PET SMC1 (Mango), were downloaded from St Jude Protein Paint program under St Jude Cloud (https://proteinpaint.stjude.org/).

Library of Integrated Network-based Cellular Signatures (LINCS) analysis

We had two sets of gene expression data: microarray and RNA-seq for siKDM6B and GSK-J4 treatment. While the pathway analysis showed very similar results for each, we used them in different purposes. Because Lincs used a limited number of genes in charactering chemical compounds signatures, we chose

our microarray data to compare since RNA-seq data gave rise to a large amount of gene expression data
that were not covered by Lincs chemical signatures. We first identify differentially expressed genes in
microarray data using GenePattern program
(https://cloud.genepattern.org/gp/pages/index.jsf). Then we chose the downregulated genes (with logFC
>=0.7) by KDM6B knockdown or GSK-J4 treatment to compare with the chemical signatures in LINCS
database (http://www.lincsproject.org/LINCS/tools). The datasets from LINCS L1000 Chem Pert down
were downloaded and analyzed using PRISM program.
803

The Cancer Therapeutics Response Portal (CTRP) analysis
Cancer Therapeutic Response Portal (https://portals.broadinstitute.org/ctrp.v2.1/) correlated the
sensitivity patterns of 481 compounds including GSK-J4 with 19,000 basal transcript levels across 823
different human cancer cell lines and identified selective outlier transcripts[44,45], which allowed to
interrogate whether KDM6B inhibitor was correlated with specific transcripts. We chose the features that
show the correlation of drug sensitivity with gene expression or copy number of quired genes such as
E2F8.
811

Kaplan-Meier analysis
A 149-gene signature commonly downregulated by GSK-J4 in BE2C and three other neuroblastoma cell
lines (IMR5, LAN5 and SK-N-FI)[43] was uploaded into R2 genomics analysis and visualization program
(https://hgserver1.amc.nl/cgi-bin/r2/main.cgi) to find the differentially expressed genes in high-risk
neuroblastoma in dataset (GSE49710) that has 498 cases, followed by k-means cluster analysis. The 4
clusters were stored as track for Kaplan-Meier curve analysis using Log-Rank method for event-free
survival and overall survival.
819

820 **Statistical analysis**
821 To determine statistical significance, the unpaired, two-tailed Student t test was calculated using the t test
822 calculator available on GraphPad Prism 8.0 software. A p value of less than 0.05 was considered
823 statistically significant.
824

Data accessibility
Super series GSE149539 includes 3 datasets.
https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE149539&token=ehozmooupbajlaf

GSE182884 [CUT&TAG]

KDM6B regulates an CDK4/6-pRB-E2F transcriptome that is correlated with GSK-J4 sensitivity

GSE149519 [ATAC-seq]

KDM6B regulates an CDK4/6-pRB-E2F transcriptome that is correlated with GSK-J4 sensitivity

GSE149537 [RNA-seq]

KDM6B regulates an CDK4/6-pRB-E2F transcriptome that is correlated with GSK-J4 sensitivity

Acknowledgments

This work was partly supported by American Cancer Society-Research Scholar (130421-RSG-17-071-01-

TBG, Jun Yang) and National Cancer Institute (1R01CA229739-01, Jun Yang). The content is solely the

responsibility of the authors and does not necessarily represent the official views of the National Institutes

of Health.

Contributions

Alexandra D'Oto, Jie Fang, Shivendra Singh, Anoushka Mullasseril, Victoria Jones, Xinyu von Buttlar,

Bailey Cooke, Dongli Hu, Ahmed Abu-Zaid performed experiments. Beisi Xu and Hongjian Jin analyzed

sequencing data. Jun Yang and Andrew M Davidoff conceived the project. Jun Yang wrote the

manuscript with input from Jason Shohet, Andrew J Murphy and Andrew M Davidoff.

References

1   Agger, K. *et al.* UTX and JMJD3 are histone H3K27 demethylases involved in
849   HOX gene regulation and development. *Nature* **449**, 731-734,
850   doi:10.1038/nature06145 (2007).
851   2   De Santa, F. *et al.* The histone H3 lysine-27 demethylase Jmjd3 links
852   inflammation to inhibition of polycomb-mediated gene silencing. *Cell* **130**,
853   1083-1094, doi:10.1016/j.cell.2007.08.019 (2007).
854   3   Hong, S. *et al.* Identification of JmjC domain-containing UTX and JMJD3 as
855   histone H3 lysine 27 demethylases. *Proc Natl Acad Sci U S A* **104**, 18439-
856   18444, doi:10.1073/pnas.0707292104 (2007).
857   4   Xiang, Y. *et al.* JMJD3 is a histone H3K27 demethylase. *Cell Res* **17**, 850-857,
858   doi:10.1038/cr.2007.83 (2007).
859   5   Margueron, R. *et al.* Ezh1 and Ezh2 maintain repressive chromatin through
860   different mechanisms. *Mol Cell* **32**, 503-518,
861   doi:10.1016/j.molcel.2008.11.004 (2008).
862   6   Walport, L. J. *et al.* Human UTY(KDM6C) is a male-specific N-methyl lysyl
863   demethylase. *J Biol Chem* **289**, 18302-18313, doi:10.1074/jbc.M114.555052
864   (2014).

865 7 Kandoth, C. *et al.* Mutational landscape and significance across 12 major
866  cancer types. *Nature* **502**, 333-339, doi:10.1038/nature12634 (2013).
867 8 Huether, R. *et al.* The landscape of somatic mutations in epigenetic regulators
868  across 1,000 paediatric cancer genomes. *Nat Commun* **5**, 3630,
869  doi:10.1038/ncomms4630 (2014).
870 9 Agger, K. *et al.* The H3K27me3 demethylase JMJD3 contributes to the
871  activation of the INK4A-ARF locus in response to oncogene- and stress-
872  induced senescence. *Genes Dev* **23**, 1171-1176, doi:10.1101/gad.510809
873  (2009).
874 10 Barradas, M. *et al.* Histone demethylase JMJD3 contributes to epigenetic
875  control of INK4a/ARF by oncogenic RAS. *Genes Dev* **23**, 1177-1182,
876  doi:10.1101/gad.511109 (2009).
877 11 Ene, C. I. *et al.* Histone demethylase Jumonji D3 (JMJD3) as a tumor
878  suppressor by regulating p53 protein nuclear stabilization. *PLoS One* **7**,
879  e51407, doi:10.1371/journal.pone.0051407 (2012).
880 12 Yamamoto, K. *et al.* Loss of histone demethylase KDM6B enhances
881  aggressiveness of pancreatic cancer through downregulation of C/EBPalpha.
882  *Carcinogenesis* **35**, 2404-2414, doi:10.1093/carcin/bgu136 (2014).
883 13 Arcipowski, K. M., Martinez, C. A. & Ntziachristos, P. Histone demethylases in
884  physiology and cancer: a tale of two enzymes, JMJD3 and UTX. *Curr Opin*
885  *Genet Dev* **36**, 59-67, doi:10.1016/j.gde.2016.03.010 (2016).
886 14 Ntziachristos, P. *et al.* Contrasting roles of histone 3 lysine 27 demethylases
887  in acute lymphoblastic leukaemia. *Nature* **514**, 513-517,
888  doi:10.1038/nature13605 (2014).
889 15 Ohguchi, H. *et al.* KDM6B modulates MAPK pathway mediating multiple
890  myeloma cell growth and survival. *Leukemia*, doi:10.1038/leu.2017.141
891  (2017).
892 16 Sherry-Lynes, M. M., Sengupta, S., Kulkarni, S. & Cochran, B. H. Regulation of
893  the JMJD3 (KDM6B) histone demethylase in glioblastoma stem cells by
894  STAT3. *PLoS One* **12**, e0174775, doi:10.1371/journal.pone.0174775 (2017).
895 17 Tang, B. *et al.* Aberrant JMJD3 Expression Upregulates Slug to Promote
896  Migration, Invasion, and Stem Cell-Like Behaviors in Hepatocellular
897  Carcinoma. *Cancer Res* **76**, 6520-6532, doi:10.1158/0008-5472.CAN-15-
898  3029 (2016).
899 18 Dalvi, M. P. *et al.* Taxane-Platin-Resistant Lung Cancers Co-develop
900  Hypersensitivity to JumonjiC Demethylase Inhibitors. *Cell Rep* **19**, 1669-
901  1684, doi:10.1016/j.celrep.2017.04.077 (2017).
902 19 Maris, J. M. Recent advances in neuroblastoma. *N Engl J Med* **362**, 2202-2211,
903  doi:10.1056/NEJMra0804577 (2010).
904 20 Cheung, N. K. & Dyer, M. A. Neuroblastoma: developmental biology, cancer
905  genomics and immunotherapy. *Nat Rev Cancer* **13**, 397-411,
906  doi:10.1038/nrc3526 (2013).
907 21 Bosse, K. R. & Maris, J. M. Advances in the translational genomics of
908  neuroblastoma: From improving risk stratification and revealing novel
909  biology to identifying actionable genomic alterations. *Cancer*,
910  doi:10.1002/cncr.29706 (2015).

22   Gustafson, W. C. & Weiss, W. A. Myc proteins as therapeutic targets. *Oncogene*
29, 1249-1259, doi:10.1038/onc.2009.512 (2010).
23   Chipumuro, E. *et al.* CDK7 inhibition suppresses super-enhancer-linked
oncogenic transcription in MYCN-driven cancer. *Cell* **159**, 1126-1139,
doi:10.1016/j.cell.2014.10.024 (2014).
24   Puissant, A. *et al.* Targeting MYCN in neuroblastoma by BET bromodomain
inhibition. *Cancer Discov* **3**, 308-323, doi:10.1158/2159-8290.CD-12-0418
(2013).
25   He, S., Liu, Z., Oh, D. Y. & Thiele, C. J. MYCN and the epigenome. *Front Oncol* **3**,
1, doi:10.3389/fonc.2013.00001 (2013).
26   Yang, J. *et al.* The role of histone demethylase KDM4B in Myc signaling in
neuroblastoma. *J Natl Cancer Inst* **107**, djv080, doi:10.1093/jnci/djv080
(2015).
27   Yang, J. *et al.* Targeting Histone Demethylases in MYC-Driven
Neuroblastomas with Ciclopirox. *Cancer Res* **77**, 4626-4638,
doi:10.1158/0008-5472.CAN-16-0826 (2017).
28   Chen, L. *et al.* CRISPR-Cas9 screen reveals a MYCN-amplified neuroblastoma
dependency on EZH2. *J Clin Invest* **128**, 446-462, doi:10.1172/JCI90793
(2018).
29   Matthay, K. K. *et al.* Neuroblastoma. *Nat Rev Dis Primers* **2**, 16078,
doi:10.1038/nrdp.2016.78 (2016).
30   Molenaar, J. J. *et al.* Sequencing of neuroblastoma identifies chromothripsis
and defects in neuritogenesis genes. *Nature* **483**, 589-593,
doi:10.1038/nature10910 (2012).
31   Fix, A. *et al.* Characterization of amplicons in neuroblastoma: high-resolution
mapping using DNA microarrays, relationship with outcome, and
identification of overexpressed genes. *Genes Chromosomes Cancer* **47**, 819-
834, doi:10.1002/gcc.20583 (2008).
32   Lastowska, M. *et al.* Identification of candidate genes involved in
neuroblastoma progression by combining genomic and expression
microarrays with survival data. *Oncogene* **26**, 7432-7444,
doi:10.1038/sj.onc.1210552 (2007).
33   Bannister, A. J. & Kouzarides, T. Regulation of chromatin by histone
modifications. *Cell Res* **21**, 381-395, doi:10.1038/cr.2011.22 (2011).
34   Whyte, W. A. *et al.* Master transcription factors and mediator establish super-
enhancers at key cell identity genes. *Cell* **153**, 307-319,
doi:10.1016/j.cell.2013.03.035 (2013).
35   Pott, S. & Lieb, J. D. What are super-enhancers? *Nat Genet* **47**, 8-12,
doi:10.1038/ng.3167 (2015).
36   Dyson, N. J. RB1: a prototype tumor suppressor and an enigma. *Genes Dev* **30**,
1492-1502, doi:10.1101/gad.282145.116 (2016).
37   Thurlings, I. & de Bruin, A. E2F Transcription Factors Control the Roller
Coaster Ride of Cell Cycle Gene Expression. *Methods Mol Biol* **1342**, 71-88,
doi:10.1007/978-1-4939-2957-3_4 (2016).
38   Strieder, V. & Lutz, W. E2F proteins regulate MYCN expression in
neuroblastomas. *J Biol Chem* **278**, 2983-2989, doi:10.1074/jbc.M207596200
(2003).

39   Ishida, S., Shudo, K., Takada, S. & Koike, K. A direct role of transcription factor
E2F in c-myc gene expression during granulocytic and macrophage-like
differentiation of HL60 cells. *Cell Growth Differ* **6**, 229-237 (1995).
40   Kruidenier, L. *et al.* A selective jumonji H3K27 demethylase inhibitor
modulates the proinflammatory macrophage response. *Nature* **488**, 404-408,
doi:10.1038/nature11262 (2012).
41   Heinemann, B. *et al.* Inhibition of demethylases by GSK-J1/J4. *Nature* **514**,
E1-2, doi:10.1038/nature13688 (2014).
42   Johansson, C. *et al.* Structural analysis of human KDM5B guides histone
demethylase inhibitor development. *Nat Chem Biol* **12**, 539-545,
doi:10.1038/nchembio.2087 (2016).
43   Lochmann, T. L. *et al.* Targeted inhibition of histone H3K27 demethylation is
effective in high-risk neuroblastoma. *Sci Transl Med* **10**,
doi:10.1126/scitranslmed.aao4680 (2018).
44   Rees, M. G. *et al.* Correlating chemical sensitivity and basal gene expression
reveals mechanism of action. *Nat Chem Biol* **12**, 109-116,
doi:10.1038/nchembio.1986 (2016).
45   Seashore-Ludlow, B. *et al.* Harnessing Connectivity in a Large-Scale Small-
Molecule Sensitivity Dataset. *Cancer Discov* **5**, 1210-1223, doi:10.1158/2159-
8290.CD-15-0235 (2015).
46   Bertoli, C., Skotheim, J. M. & de Bruin, R. A. Control of cell cycle transcription
during G1 and S phases. *Nat Rev Mol Cell Biol* **14**, 518-528,
doi:10.1038/nrm3629 (2013).
47   Buenrostro, J. D., Wu, B., Chang, H. Y. & Greenleaf, W. J. ATAC-seq: A Method
for Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol*
109, 21 29 21-21 29 29, doi:10.1002/0471142727.mb2129s109 (2015).
48   Hoxhaj, G. & Manning, B. D. The PI3K-AKT network at the interface of
oncogenic signalling and cancer metabolism. *Nat Rev Cancer* **20**, 74-88,
doi:10.1038/s41568-019-0216-7 (2020).
49   Chaussepied, M. & Ginsberg, D. Transcriptional regulation of AKT activation
by E2F. *Mol Cell* **16**, 831-837, doi:10.1016/j.molcel.2004.11.003 (2004).
50   Kaya-Okur, H. S., Janssens, D. H., Henikoff, J. G., Ahmad, K. & Henikoff, S.
Efficient low-cost chromatin profiling with CUT&Tag. *Nat Protoc* **15**, 3264-
3283, doi:10.1038/s41596-020-0373-x (2020).
51   Helmsauer, K. *et al.* Enhancer hijacking determines extrachromosomal
circular MYCN amplicon architecture in neuroblastoma. *Nat Commun* **11**,
5823, doi:10.1038/s41467-020-19452-y (2020).
52   Phillips, J. E. & Corces, V. G. CTCF: master weaver of the genome. *Cell* **137**,
1194-1211, doi:10.1016/j.cell.2009.06.001 (2009).
53   Wendt, K. S. *et al.* Cohesin mediates transcriptional insulation by CCCTC-
binding factor. *Nature* **451**, 796-801, doi:10.1038/nature06634 (2008).
54   Debruyne, D. N. *et al.* BORIS promotes chromatin regulatory interactions in
treatment-resistant cancer cells. *Nature* **572**, 676-680, doi:10.1038/s41586-
019-1472-0 (2019).
55   Local, A. *et al.* Identification of H3K4me1-associated proteins at mammalian
enhancers. *Nat Genet* **50**, 73-82, doi:10.1038/s41588-017-0015-6 (2018).

56 Kagey, M. H. *et al.* Mediator and cohesin connect gene expression and
chromatin architecture. *Nature* **467**, 430-435, doi:10.1038/nature09380
(2010).
57 Kim, S. I., Bultman, S. J., Kiefer, C. M., Dean, A. & Bresnick, E. H. BRG1
requirement for long-range interaction of a locus control region with a
downstream promoter. *Proc Natl Acad Sci U S A* **106**, 2259-2264,
doi:10.1073/pnas.0806420106 (2009).
58 Beagrie, R. A. *et al.* Complex multi-enhancer contacts captured by genome
architecture mapping. *Nature* **543**, 519-524, doi:10.1038/nature21411
(2017).
59 Rao, S. S. *et al.* A 3D map of the human genome at kilobase resolution reveals
principles of chromatin looping. *Cell* **159**, 1665-1680,
doi:10.1016/j.cell.2014.11.021 (2014).
60 Bracken, A. P. *et al.* EZH2 is downstream of the pRB-E2F pathway, essential
for proliferation and amplified in cancer. *EMBO J* **22**, 5323-5335,
doi:10.1093/emboj/cdg542 (2003).
61 Lee, S. R. *et al.* Activation of EZH2 and SUZ12 Regulated by E2F1 Predicts the
Disease Progression and Aggressive Characteristics of Bladder Cancer. *Clin*
Cancer Res **21**, 5391-5403, doi:10.1158/1078-0432.CCR-14-2680 (2015).
62 Keenan, A. B. *et al.* The Library of Integrated Network-Based Cellular
Signatures NIH Program: System-Level Cataloging of Human Cells Response
to Perturbations. *Cell Syst* **6**, 13-24, doi:10.1016/j.cels.2017.11.001 (2018).
63 Filippakopoulos, P. *et al.* Selective inhibition of BET bromodomains. *Nature*
468, 1067-1073, doi:10.1038/nature09504 (2010).
64 Sherr, C. J., Beach, D. & Shapiro, G. I. Targeting CDK4 and CDK6: From
Discovery to Therapy. *Cancer Discov* **6**, 353-367, doi:10.1158/2159-8290.CD-
15-0894 (2016).
65 Alvarez-Fernandez, M. & Malumbres, M. Mechanisms of Sensitivity and
Resistance to CDK4/6 Inhibition. *Cancer Cell* **37**, 514-529,
doi:10.1016/j.ccell.2020.03.010 (2020).
66 Klein, M. E., Kovatcheva, M., Davis, L. E., Tap, W. D. & Koff, A. CDK4/6
Inhibitors: The Mechanism of Action May Not Be as Simple as Once Thought.
Cancer Cell **34**, 9-20, doi:10.1016/j.ccell.2018.03.023 (2018).
67 Yang, L. *et al.* Histone demethylase KDM6B has an anti-tumorigenic function
in neuroblastoma by promoting differentiation. *Oncogenesis* **8**, 3,
doi:10.1038/s41389-018-0112-0 (2019).
68 Hashizume, R. *et al.* Pharmacologic inhibition of histone demethylation as a
therapy for pediatric brainstem glioma. *Nat Med* **20**, 1394-1396,
doi:10.1038/nm.3716 (2014).
69 Morozov, V. M., Li, Y., Clowers, M. M. & Ishov, A. M. Inhibitor of H3K27
demethylase JMJD3/UTX GSK-J4 is a potential therapeutic option for
castration resistant prostate cancer. *Oncotarget* **8**, 62131-62142,
doi:10.18632/oncotarget.19100 (2017).
70 Mathur, R. *et al.* Inhibition of demethylase KDM6B sensitizes diffuse large B-
cell lymphoma to chemotherapeutic drugs. *Haematologica* **102**, 373-380,
doi:10.3324/haematol.2016.144964 (2017).

71   Hong, B. J. *et al.* Oncogenic KRAS Sensitizes Lung Adenocarcinoma to GSK-J4-
Induced Metabolic and Oxidative Stress. *Cancer Res* **79**, 5849-5859,
doi:10.1158/0008-5472.CAN-18-3511 (2019).
72   Hnisz, D. *et al.* Super-enhancers in the control of cell identity and disease. *Cell*
155, 934-947, doi:10.1016/j.cell.2013.09.053 (2013).
73   Leone, G. *et al.* Myc requires distinct E2F activities to induce S phase and
apoptosis. *Mol Cell* **8**, 105-113, doi:10.1016/s1097-2765(01)00275-1 (2001).
74   Chen, X. *et al.* Integration of external signaling pathways with the core
transcriptional network in embryonic stem cells. *Cell* **133**, 1106-1117,
doi:10.1016/j.cell.2008.04.043 (2008).
75   Sherr, C. J. & McCormick, F. The RB and p53 pathways in cancer. *Cancer Cell* **2**,
103-112, doi:10.1016/s1535-6108(02)00102-2 (2002).
76   Dick, F. A., Goodrich, D. W., Sage, J. & Dyson, N. J. Non-canonical functions of
the RB protein in cancer. *Nat Rev Cancer* **18**, 442-451, doi:10.1038/s41568-
018-0008-5 (2018).
77   Easton, J., Wei, T., Lahti, J. M. & Kidd, V. J. Disruption of the cyclin D/cyclin-
dependent kinase/INK4/retinoblastoma protein regulatory pathway in
human neuroblastoma. *Cancer Res* **58**, 2624-2632 (1998).
78   Krasnoselsky, A. L. *et al.* Altered expression of cell cycle genes distinguishes
aggressive neuroblastoma. *Oncogene* **24**, 1533-1541,
doi:10.1038/sj.onc.1208341 (2005).
79   Molenaar, J. J. *et al.* Copy number defects of G1-cell cycle genes in
neuroblastoma are frequent and correlate with high expression of E2F target
genes and a poor prognosis. *Genes Chromosomes Cancer* **51**, 10-19,
doi:10.1002/gcc.20926 (2012).
80   Mosse, Y. P. *et al.* Neuroblastomas have distinct genomic DNA profiles that
predict clinical phenotype and regional gene expression. *Genes Chromosomes
Cancer* **46**, 936-949, doi:10.1002/gcc.20477 (2007).
81   Mosse, Y. P. *et al.* High-resolution detection and mapping of genomic DNA
alterations in neuroblastoma. *Genes Chromosomes Cancer* **43**, 390-403,
doi:10.1002/gcc.20198 (2005).
82   Molenaar, J. J. *et al.* Cyclin D1 and CDK4 activity contribute to the
undifferentiated phenotype in neuroblastoma. *Cancer Res* **68**, 2599-2609,
doi:10.1158/0008-5472.CAN-07-5032 (2008).
83   Rader, J. *et al.* Dual CDK4/CDK6 inhibition induces cell-cycle arrest and
senescence in neuroblastoma. *Clin Cancer Res* **19**, 6173-6182,
doi:10.1158/1078-0432.CCR-13-1675 (2013).
84   Lee, M. G. *et al.* Demethylation of H3K27 regulates polycomb recruitment and
H2A ubiquitination. *Science* **318**, 447-450, doi:10.1126/science.1149042
(2007).
85   Kim, J. H. *et al.* UTX and MLL4 coordinately regulate transcriptional programs
for cell proliferation and invasiveness in breast cancer cells. *Cancer Res* **74**,
1705-1717, doi:10.1158/0008-5472.CAN-13-1896 (2014).
86   Wong, A. S. *et al.* Multiplexed barcoded CRISPR-Cas9 screening enabled by
CombiGEM. *Proc Natl Acad Sci U S A* **113**, 2544-2549,
doi:10.1073/pnas.1517883113 (2016).

87   Yan, J. *et al.* Histone H3 lysine 4 monomethylation modulates long-range
chromatin interactions at enhancers. *Cell Res* **28**, 204-220,
doi:10.1038/cr.2018.1 (2018).
88   Vernimmen, D. *et al.* Polycomb eviction as a new distant enhancer function.
Genes Dev **25**, 1583-1588, doi:10.1101/gad.16985411 (2011).
89   Kartikasari, A. E. *et al.* The histone demethylase Jmjd3 sequentially associates
with the transcription factors Tbx3 and Eomes to drive endoderm
differentiation. *EMBO J* **32**, 1393-1408, doi:10.1038/emboj.2013.78 (2013).
90   Fueyo, R. *et al.* Lineage specific transcription factors and epigenetic
regulators mediate TGFbeta-dependent enhancer activation. *Nucleic Acids*
Res **46**, 3351-3365, doi:10.1093/nar/gky093 (2018).
91   Li, Z. *et al.* EZH2 regulates neuroblastoma cell differentiation via NTRK1
promoter epigenetic modifications. *Oncogene* **37**, 2714-2727,
doi:10.1038/s41388-018-0133-3 (2018).
92   Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based
approach for interpreting genome-wide expression profiles. *Proc Natl Acad*
Sci U S A **102**, 15545-15550, doi:10.1073/pnas.0506580102 (2005).
93   Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J.
Transposition of native chromatin for fast and sensitive epigenomic profiling
of open chromatin, DNA-binding proteins and nucleosome position. *Nat*
Methods **10**, 1213-1218, doi:10.1038/nmeth.2688 (2013).
94   Martin, M. Cutadapt removes adapter sequences from high-throughput
sequencing reads. *EMBnet.journal* **17**, pp.-10-12 (2011).
95   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-
Wheeler transform. *Bioinformatics* **25**, 1754-1760,
doi:10.1093/bioinformatics/btp324 (2009).
96   Tischler, G. & Leonard, S. biobambam: tools for read pair collation based
algorithms on BAM files. *Source Code for Biology and Medicine* **9**, 13,
doi:10.1186/1751-0473-9-13 (2014).
97   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools.
Bioinformatics **25**, 2078-2079, doi:10.1093/bioinformatics/btp352 (2009).
98   Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J.
Transposition of native chromatin for fast and sensitive epigenomic profiling
of open chromatin, DNA-binding proteins and nucleosome position. *Nature*
Methods **10**, 1213-1218, doi:10.1038/nmeth.2688 (2013).
99   Robinson, J. T. *et al.* Integrative genomics viewer. *Nature Biotechnology* **29**,
24-26, doi:10.1038/nbt.1754 (2011).
100   Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biology* **9**,
R137, doi:10.1186/gb-2008-9-9-r137 (2008).
101   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**, 841-842,
doi:10.1093/bioinformatics/btq033 (2010).
102   Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: precision weights unlock
linear model analysis tools for RNA-seq read counts. *Genome Biology* **15**, R29,
doi:10.1186/gb-2014-15-2-r29 (2014).
103   Matys, V. *et al.* TRANSFAC and its module TRANSCompel: transcriptional
gene regulation in eukaryotes. *Nucleic Acids Res* **34**, D108 - 110 (2006).

104   Bailey, T. L. *et al.* MEME SUITE: tools for motif discovery and searching.
1144   *Nucleic Acids Res* **37**, W202-208, doi:10.1093/nar/gkp335 (2009).
1145   105   Skene, P. J. & Henikoff, S. An efficient targeted nuclease strategy for high-
1146   resolution mapping of DNA binding sites. *Elife* **6**, doi:10.7554/eLife.21856
1147   (2017).
1148   106   Orlando, D. A. *et al.* Quantitative ChIP-Seq normalization reveals global
1149   modulation of the epigenome. *Cell Rep* **9**, 1163-1170,
1150   doi:10.1016/j.celrep.2014.10.018 (2014).
1151   107   Ramirez, F. *et al.* deepTools2: a next generation web server for deep-
1152   sequencing data analysis. *Nucleic Acids Res* **44**, W160-165,
1153   doi:10.1093/nar/gkw257 (2016).
1154   108   McLeod, C. *et al.* St. Jude Cloud: A Pediatric Cancer Genomic Data-Sharing
1155   Ecosystem. *Cancer Discov* **11**, 1082-1099, doi:10.1158/2159-8290.CD-20-
1156   1230 (2021).
1157   109   Zeid, R. *et al.* Enhancer invasion shapes MYCN-dependent transcriptional
1158   amplification in neuroblastoma. *Nat Genet* **50**, 515-523,
1159   doi:10.1038/s41588-018-0044-9 (2018).
1160
1161

Figure 1. KDM6B is marked with a highly active epigenetic landscape and highly expressed in neuroblastoma and regulates MYC expression.

(A-C) The expression of KDM6B, KDM6A and UTY in normal human trunk neural crest (GSE14340) and 4 different neuroblastoma cohorts (Versteeg GSE16476, Delattre GSE14880, Hiyama GSE16237, Lastowska GSE13136). Y-axis represents the normalized log2 expression value. ***p<0.001, student t test

(D-F) The expression of KDM6B, KDM6A and UTY in 3 different neuroblastoma RNA-seq cohorts. The RNA-seq data of St Jude was downloaded from https://pecan.stjude.cloud. The RNA-seq data of TARGET and SEQC datasets were downloaded from R2 (https://hgserver1.amc.nl/cgi-bin/r2/main.cgi). Y-axis represents the Fragment Per Kilobase of transcript per Million (FPKM) mapped reads.

(G) The epigenetic landscapes consisting of histone marks and transcription factor binding distinguish KDM6B from KDM6A in primary neuroblastoma tissues with MYCN amplification or without MYCN amplification (https://pecan.stjude.cloud/proteinpaint/study/mycn_nbl_2018).

(H) Crystal violet staining of colonies after BE2C and SKNAS cells were transfected with four different siRNA oligos to knockdown KDM6B for 7 days. siCtrl = siRNA control oligo, NT= no treatment.

(I) Western blot analysis with indicated antibodies to assess MYCN or C-MYC expression after 3-day transfection of 4 different siRNA to knockdown KDM6B in BE2C and SKNAS.

(J) BE2C and MSCV-KDM6B overexpressing (KDM6B-OE) BE2C cells were transfected with siRNA control (siCtrl) and siRNA targeting the endogenous 3' untranslated region (3'UTR) of KDM6B (siKDM6B-3'UTR). 4 days later, cells were stained with crystal violet.

(K) Quantification of cell density of each group (n=3) using imageJ. P-value calculated using student t test.

(L) BE2C and MSCV-KDM6B overexpressing (KDM6B-OE) BE2C cells were transfected with siRNA control (siCtrl) and siRNA siKDM6B-3'UTR. 2 days later, cells were subject to immunoblotting with indicated antibodies.

Figure 2. KDM6B predominantly regulates the pRB-E2F pathway.

(A) Quantitative comparison of all chemical and genetic perturbation gene sets (n = 3403) from the MSigDB by gene set enrichment analysis (GSEA) for increased (left) and reduced (right) expression of global genes caused by KDM6B knockdown. Data are presented as a scatterplot of normalized p value/false discovery q value vs normalized enrichment score (NES) for each evaluated gene set. The gene sets circled in red color indicate cell cycle, pRB-E2F and MYC pathway gene sets.

(B) Two examples of GSEA show that genes downregulated by depletion of KDM6B are enriched with Rb1 knockout and E2F targets.

(C) Heatmap shows the gene list from the E2F targets (B).

(D-F) Similar analysis for GSK-J4 treatment as shown in (A-C).

Figure 3. Chemogenetic data show E2F gene signature is correlated with sensitivity of GSK-J4.

(A) The Pearson correlation Z-score for each tested compound with E2F8 expression, which was extracted from The Cancer Therapeutics Response Portal (CTRP). The correlation of GSK-J4 and E2F8 ranked on the top with R=-0.626, Z-score =-4.56.

(B) The plot of E2F8 expression and GSK-J4 sensitivity for each tested cell line. The y-axis represents the transcript expression levels of E2F8 in each cell line. The x-axis represents the drug response metrics (area under curve, AUC) to GSK-J4. The lower the AUC, the more sensitive to GSK-J4. The circled group A and B populations are arbitrarily chosen as GSK-J4 sensitive vs resistant.

(C) Quantitative comparison of all chemical and genetic perturbation gene sets (n = 3403) from the MSigDB by gene set enrichment analysis (GSEA) for GSK-J4 sensitive (group A) and resistant (group B). Data are presented as a scatterplot of normalized p value/false discovery q value vs normalized enrichment score (NES) for each evaluated gene set. The red dots highlight cell cycle and Rb-E2F pathway gene sets.

(D) GSEA show that genes highly expressed in GSK-J4 sensitive (group A) cells are enriched with Rb1 knockout.

(E) Quantitative comparison of all chemical and genetic perturbation gene sets (n = 3403) from the MSigDB by gene set enrichment analysis (GSEA) for GSK-J4 sensitive and resistant neuroblastoma cells, as shown in the right panel of cell lines. Data are presented as a scatterplot of normalized p value/false discovery q value vs normalized enrichment score (NES) for each evaluated gene set. The red and green dots highlight cell cycle and Rb-E2F pathway gene sets and MYC gene sets, respectively.

(F) Crystal violet staining of colonies after SK-N-SH and SK-N-FI neuroblastoma cell lines were treated with different concentrations of GSK-J4 for 7 days.

Figure 4. KDM6B inhibition represses chromatin accessibility of E2F genes.

(A) ATAC-seq was performed after BE2C cells were treated with 2.5μM of GSK-J4 for 48 hours. Summary of peak calling number of ATAC-seq (p<0.05, log2 fold change >=0.5) including the upregulated and downregulated nucleosome free regions, and the annotated locations of the peaks at defined promoter and enhancer regions.

(B) Snapshot of *AKT1* locus using Integrative Genomic Viewer (IGV) for ATAC-seq, H3K27Ac ChIP-seq, and RNA-seq. The ATAC-seq analysis shows the downregulation of two peaks at the 5' promoter region of *AKT1* and one peak at the non-coding RNA *LINC00638*. The RNA-seq results showed that the AKT1 transcript was downregulated by 48h of GSK-J4 treatment and KDM6B knockdown. H3K27Ac in BE2C cells was referenced to GSM2113518[109].

(C) Snapshot of *E2F8* locus using Integrative Genomic Viewer (IGV) for ATAC-seq, H3K27Ac ChIP-seq, and RNA-seq. The ATAC-seq shows the downregulation of three peaks at the enhancer region of *E2F8*, next to the *NAV2* gene locus. The RNA-seq results showed that *E2F8* transcript was downregulated by KDM6B inhibition while the adjacent *NAV2* expression is barely detectable.

(D) Cartoon indicates the rationale of footprinting analysis. The DNA motifs bound by transcription factors such as E2F1 protect the cut from transposase Tn5, while the adjacent open chromatin gives rise to a high signal of nucleosome free region after ATAC-seq analysis. After GSK-J4 treatment, the open chromatin was repressed and consequently reducing the reads of free DNA.

 (E) Footprinting plot shows the reduction of open chromatin at predicted binding motifs of E2Fs and MYC after BE2C cells were treated with GSK-J4 for 24 h and 48h.

Figure 5. KDM6B inhibition induces an increase of H3K27me3 but a decrease of the enhancer mark H3K4me1 at CTCF and BORIS binding sites

(**A, B**) Heatmap indicating the CUT&Tag signal intensity of H3K27me3 (A) and H3K4me1(B) around differentially expressed genes (from TSS-5kb to TES+5kb). CUT&Tag experiments were done with two biological replicates in KDM6B knockdown and GSK-J4 treatment respectively. Differentially expressed genes were identified by cutoffs (log2FC>0.5) from RNA-seq. TSS= transcription start site, TES=transcription end site

(**C**) The signals of H3K27me3 and H3K4me1 (CUT&Tag), H3K27Ac (ChIP-seq)[109], and ATAC-seq between the genomic locus of *MYCN, GACAT3,* and *FAM149A*, snapshot using IGV program.

(**D, E**) Motif analysis of H3K27me3 (D) and H3K4me1 (E) peaks by using HOMER known motif search. Top ten most significant motifs from each sub-group were visualized as a heatmap. The heatmap scale indicates -$\log_{10}$(p value).

(**F**) The Hi-C data of BE2C neuroblastoma cells (data extracted from the St Jude Cloud) showing that *MYCN* and its adjacent *FAM149A* locus reside in the same topologically associated domain (TAD). The Arc indicates the chromatin interactions, which were generated from Jurkat ChIA-PET SMC1 (Mango) (https://proteinpaint.stjude.org/).

Figure 6. GSK-J4 induces a transcriptome that mimics CDK4/6 inhibition, and vice versa

(**A**) Genes downregulated by KDM6B knockdown were compared with gene profiles induced with

chemical compounds from Library of Integrated Network-Based Cellular Signatures (LINCS). The top

chemical signature hits, which are basically overrepresented by palbociclib, are highlighted in pink dots.

(**B**) Same analysis performed as in (A) for GSK-J4 treatment.

(**C**) The 89-gene signature derived from palbociclib treatment of 4 neuroblastoma cell lines was included

in gene sets for GSEA analysis of KDM6B knockdown.

(**D**) The 89-gene signature derived from palbociclib treatment was included in gene sets for GSEA analysis

of GSK-J4.

(**E**) BE2C cells were seeded with low numbers in 6-well plate and treated with different concentrations of

GSK-J4 or/and palbociclib for 10 days. The cell colonies were stained with crystal violet.

(**F**) Bliss index for the combination of GSK-J4 and palbociclib. Positive scores indicate synergy, negative

scores indicate antagonism.

(**G**) Western blot to assess the expression of CDK4 and CDK6 that were transduced into BE2C cells.

(**H**) The BE2C parental and CDK4/6 overexpressing cells were seeded with low numbers in 6-well plate

and treated with different concentrations of GSK-J4 or palbociclib for 8 days. The cell colonies were

stained with crystal violet.

(**I, J**) BE2C and CDK4 or CDK6 overexpressing BE2C cells (BE2C-CDK4-OE, BE2C-CDK6-OE) were

treated with different concentrations of palbociclib (I) or GSK-J4 (J). EC50 was calculated using

Prestoblue assay.

(**K**) Western blot to assess the expression of pRB after inducible knockout in BE2C cells.

(**L**) Photos taken under microscope (10x) show Rb1 knockout leads to resistance to palbociclib.

(**M**) The BE2C wildtype and Rb1 knockout cells were seeded with low numbers in 6-well plate and treated

with different concentrations of GSK-J4 or/and palbociclib for 8 days. The cell colonies were stained with

crystal violet.

1314

1315

1316

1317

1318

1319

1320

Figure 7. A gene signature targeted by KDM6B inhibition is associated with poor outcome

(**A**) 149-gene signature was uploaded into R2 genomics analysis program (https://hgserver1.amc.nl/cgi-bin/r2/main.cgi) to find the differentially expressed genes in high-risk neuroblastoma in a cohort dataset (GSE49710) that has 498 neuroblastoma cases, followed by k-means cluster analysis. Risk (Red =high risk; green =low risk); MYCN (red = Amplification; green = non-Amplification); Stage (red = stage 4; blue= stage 4S; brown =stage 3; dark green = stage 2, green= stage 1).

(**B-C**) Kaplan-Meier curve using Log-Rank method shows the event-free survival and overall survival of 4 clusters that have differential expression levels of GSK-J4 signature.

(**D, F**) Kaplan-Meier curve using Log-Rank method shows the event-free survival and overall survival of clusters 3 and 4 with MNA.

(**E, G**) Kaplan-Meier curve using Log-Rank method shows the event-free survival and overall survival of clusters 3 and 4 with high-risk disease.

1347

Figure 8. Working model of KDM6B inhibition in neuroblastoma.

(**A**) Upon stimulation by mitogens, KDM6B is recruited to chromatin to maintain the low levels of
H3K27me3 at the distal regulatory enhancer regions marked by H3K4me1 overlapping with the
CTCF/BORIS binding sites, which loops and physically interacts with E2F that binds at the promoter of
target genes, together with associated transcriptional machinery including RNA polymerase II and
mediators, driving the MYCN and E2F transcriptome.

(**B**) When inhibited by CDK4/6 inhibitor, the inhibitory pRB complexes with HDAC to suppress gene
transcription. When KDM6B is inhibited by GSK-J4, the H3K27me3 will accumulate at the distal regions
to displace the H3K4me1 modifier KMT2 and evicts the transcription activators of promoter-enhancer,
leading to the downregulation of MYCN and E2F transcriptome (Figure 7B).

(**C**) A network composed by MYCN, E2F, EZH2 and KDM6B regulates the cell proliferation and
differentiation of neuroblastoma.

1360

1361

1362

1363

1364

1365

1366

1367

Table 1. Transcription factor binding motif enrichment for KDM6B target genes (top15)

	Gene Sets	NES	NOM p-val	FDR q-val
1	KRCTCNNNNMANAGC_UNKNOWN	2.76	0	0
2	SGCGSSAAA_E2F1DP2_01	2.17	0	0
3	TTTNNANAGCYR_UNKNOWN	2.14	0	0
4	E2F4DP1_01	2.1	0	0
5	E2F_03	2.06	0	0
6	E2F1_Q6	2.03	0	0
7	E2F_02	2.03	0	0
8	E2F_Q6	2.02	0	0
9	E2F_Q3	2.01	0	0
10	E2F4DP2_01	2	0	0
11	E2F1DP2_01	2	0	0
12	E2F1DP1_01	1.99	0	0
13	E2F_Q4	1.99	0	0
14	E2F_Q6_01	1.92	0	0
15	E2F1_Q6_01	1.9	0	0

Table 2. Transcription factor binding motif enrichment for GSK-J4 target genes (top 15)

	Gene sets	NES	NOM p-val	FDR q-val
1	KRCTCNNNNMANAGC_UNKNOWN	2.43	0	0
2	SGCGSSAAA_E2F1DP2_01	1.48	0.003	0.253
3	E2F_01	1.42	0.037	0.31
4	E2F_Q3	1.39	0.012	0.325
5	TTTNNANAGCYR_UNKNOWN	1.39	0.023	0.262
6	E2F_Q6	1.36	0.012	0.274
7	E2F1_Q3	1.33	0.015	0.308
8	E2F4DP1_01	1.32	0.018	0.301
9	E2F1_Q6	1.32	0.02	0.269
10	E2F_Q4	1.31	0.013	0.273
11	RGTTAMWNATT_HNF1_01	1.31	0.071	0.25
12	E2F1DP2_01	1.29	0.027	0.268
13	E2F_02	1.28	0.028	0.266
14	E2F1_Q6_01	1.28	0.042	0.255
15	E2F1DP1_01	1.28	0.037	0.246

Table 3. Transcription factor binding motif enrichment for neuroblastoma cells sensitive to GSK-J4

	Gene Sets	NES	NOM p-val	FDR q-val
1	E2F_Q6	1.89019	0.00000	0.00000
2	E2F_Q4	1.86916	0.00000	0.00112
3	E2F4DP2_01	1.84061	0.00000	0.00075
4	E2F_02	1.83652	0.00000	0.00112
5	E2F1DP1_01	1.83527	0.00000	0.00090
6	E2F1_Q6	1.83142	0.00000	0.00075
7	E2F1DP2_01	1.83124	0.00000	0.00064
8	E2F1_Q3	1.82484	0.00000	0.00056
9	E2F1DP1RB_01	1.80926	0.00000	0.00050
10	E2F4DP1_01	1.80680	0.00000	0.00045
11	E2F_Q3_01	1.80364	0.00000	0.00041
12	E2F_03	1.79745	0.00000	0.00064
13	SGCGSSAAA_E2F1DP2_01	1.79145	0.00000	0.00076
14	E2F1_Q4_01	1.77454	0.00000	0.00125
15	E2F_Q4_01	1.77275	0.00000	0.00117
16	E2F_Q6_01	1.71608	0.00000	0.00258
17	E2F1_Q6_01	1.67011	0.00000	0.00468
18	E2F_Q3	1.66943	0.00000	0.00448
19	KRCTCNNNNMANAGC_UNKNOWN	1.63733	0.00184	0.00603
20	NRF1_Q6	1.55445	0.00000	0.01859
21	NRSF_01	1.54447	0.00000	0.01968
22	TMTCGCGANR_UNKNOWN	1.52952	0.00000	0.02146
23	CAGNWMCNNNGAC_UNKNOWN	1.52403	0.00000	0.02186
24	ACTAYRNNNCCCR_UNKNOWN	1.51553	0.00000	0.02396
25	MCAATNNNNGCG_UNKNOWN	1.51439	0.00372	0.02336