Relationships between macro-fungal dark diversity and habitat parameters using LiDAR

Jose W. Valdez[1], Ane Kirstine Brunbjerg[1], Camilla Fløjgaard[1], Lars Dalby[1], Kevin K. Clausen[1], Meelis Pärtel[2], Norbert Pfeifer[3], Markus Hollaus[3], Michael H. Wimmer[3], Rasmus Ejrnæs[1], Jesper Erenskjold Moeslund[1]*

[1]Department of Bioscience, Aarhus University, Grenåvej 14, 8410, Rønde, Denmark

[2] Department of Botany, Institute of Ecology and Earth Sciences, University of Tartu, Lai 40, Tartu, 51005, Estonia

3 Department of Geodesy and Geoinformation, Technische Universität Wien, Wiedner Hauptstraße 8/E120, 1040 Vienna, Austria

*Corresponding author. Email: jesper.moeslund@bios.au.dk

Abstract

Despite the important role of fungi for ecosystems, relatively little is known about the factors underlying the dynamics of their diversity. Moreover, studies do not typically consider their dark diversity: the species absent from an otherwise suitable site. Here, we examined potential drivers of local fungal dark diversity in temperate woodland and open habitats using LiDAR and in-situ field measurements, combined with a systematically collected and geographically comprehensive macro-fungi and plant data set. For the first time, we also estimated species pools of fungi by considering both plant and fungi co-occurrences. The most important LiDAR variables for modelling fungal dark diversity were amplitude and echo ratio, which are both thought to represent vegetation structure. These results suggest that the local fungal dark diversity is highest in production forests like plantations and lowest in more open forests and in open habitats with little woody vegetation. Plant species richness was the strongest explanatory factor overall and negatively correlated with local fungal dark diversity. Soil fertility showed a positive relationship with dark diversity in open habitats. These findings may indicate that the local dark diversity of macro-fungi is highest in areas with a relatively high human impact (typically areas with low plant species richness and high soil fertility). Overall, this study brings novel insights into local macro-fungi dark diversity patterns, suggesting that a multitude of drivers related to both soil and vegetation act in concert to determine fungal dark diversity.

Key Words: airborne laser scanning, bog, fens, forests, fungal diversity, grasslands,

mycorrhiza, regional species pool, remote sensing, shrublands, wetlands

Introduction

Understanding the underlying drivers shaping biodiversity patterns is a central goal in

ecology and conservation biology. This is also true for fungi which play a vital role in

ecosystem functioning as decomposers, mutualists, and pathogens. However, fungi and the

underlying environmental factors influencing fungal diversity is less studied than animals and

plants, and quantifying fungal diversity is far from trivial. The most commonly used

biodiversity metric is observed species richness (Mueller 2011). However, this measure is not

always suitable for comparisons across habitats and conveys no information on the part of the

diversity that is potentially missing in a given site (Pärtel *et al.* 2011). In addition, monitoring

fungal diversity is often severely hampered by detectability issues and the life history of the

involved species (Yahr *et al.* 2016, Blackwell and Vega 2018). Several alternative

approaches have been developed to more effectively monitor and compare biodiversity across

landscapes (Solow and Polasky 1994, Sarkar and Margules 2002, Ricotta 2005, Ricotta

2007). Although these methods can provide valuable insights, they do not consider the dark

diversity, the absent part of the species pool which can potentially inhabit an environmentally

suitable site (Pärtel *et al.* 2011). This often-ignored aspect of diversity provides a novel and

ecologically meaningful metric for estimating how much of the potential species diversity –

the site-specific species pool – is lacking (Pärtel *et al.* 2011). This information is important to

understand the underlying mechanisms and dynamics of community assembly (e.g.,

community saturation) (Mateo *et al.* 2017). Dark diversity may also become an important

conservation tool to measure biodiversity potential, such as aiding managers or policy-makers

to prioritize certain habitats, estimate restoration potential of degraded habitats, or forecast

potential impacts of invasions (Lewis *et al.* 2017). Here, we use fungal data from 130

thoroughly inventoried sites covering all terrestrial habitats, from open to forest, and wet to

arid, to investigate important drivers of fungal dark diversity.

Dark diversity aims to reconcile the role of simultaneous, and potentially

confounding, regional and local processes underlying biodiversity patterns and biological

communities (Pärtel *et al.* 2011, Pärtel 2014). In any given landscape, the biodiversity

potential is ultimately determined by large-scale biogeographic and evolutionary processes

(i.e., species diversification and historic migration patterns) determining the set of species

which can theoretically inhabit a site, defined as the regional pool (Pärtel *et al.* 1996, Cornell and Harrison 2014, Zobel 2016). This regional pool is further filtered by local processes such as environmental gradients, species interactions, population dynamics, dispersal, disturbance, and stochastic events and referred to as the site-specific species pool, i.e., species that could possibly live in a given site (Pärtel *et al.* 2013, Cornell and Harrison 2014, Ronk *et al.* 2015, Zobel 2016). While many studies have investigated the drivers of fungal diversity, only a few studies have focused on the determinants of fungal dark diversity. These studies demonstrate that higher temperatures increases arbuscular mycorrhizal dark diversity (Pärtel *et al.* 2017a) and annual precipitation decreases the dark diversity of ectomycorrhizal fungi at the global scale (Pärtel *et al.* 2017b). These results concur with previous research suggesting that large scale climatic factors are strong drivers of fungal richness and community composition, attributed to the direct and indirect effects which alter soil and floristic conditions (Staddon *et al.* 2003, Kivlin *et al.* 2011, Tedersoo *et al.* 2014). Local edaphic conditions such as soil moisture, pH, and calcium concentration are also known to influence fungal diversity (Geml *et al.* 2014, Tedersoo *et al.* 2014, Tonn and Ibáñez 2017, Frøslev *et al.* 2019), but it is not known how these environmental factors affect fungal dark diversity. In fact, the general mechanisms determining dark diversity in fungal communities remain largely unknown. Clearly, species can be absent from an area just by chance (Hubbell 2011). Species can also be absent from a site because of some kind of unexpected disturbance – for example human, but it could also be natural – altering species' dispersal, establishment, or persistence possibilities. In principle, these disturbances can be both biological and chemical and act at various spatial scales. An example could be extreme drought. Often habitats have a constant level of relatively low or usual disturbances that the habitat's species are adapted to (e.g., grazing), and these do not count towards disturbances that can cause dark diversity. It is important not to confuse dark diversity with hidden diversity; i.e., species that are actually present in a given site but just not recorded (Milberg *et al.* 2008, Abrego *et al.* 2016).

Besides the influence of environmental gradients, other factors particularly important for fungi are vegetation and habitat structure, such as vegetation height, shrub layer, vegetation cover, dead wood, and other woody features (Humphrey *et al.* 2000, Nordén and Paltto 2001, Nordén *et al.* 2004, Gómez-Hernández and Williams-Linera 2011, Zuo *et al.* 2016). As the dominant primary producer in terrestrial ecosystems, plants also form the living and dead organic carbon pools and biotic surfaces that are the niche space for not only fungi but other taxonomic groups as well (DeAngelis 2012, Brunbjerg *et al.* 2017). These structural

elements are an important element for biodiversity, and can influence not only fungal

diversity, but the diversity of plants, animals, and bacteria as well (Penone *et al.* 2019).

However, despite the obvious contribution of these variables, such factors are rarely covered

extensively since they are difficult to measure and require large amounts of resources to

obtain sufficient and high quality data. However, emerging technologies such as LiDAR

(light detection and ranging) could potentially remedy this situation.

Airborne LiDAR records a three-dimensional set of points using laser ranging from an

aircraft or a drone (Lefsky *et al.* 2002). It captures data suitable to represent many of the

vegetation and landscape structural measures important to fungi (Vehmas *et al.* 2009, Lopatin

et al. 2016, Peura *et al.* 2016, Thers *et al.* 2017, Mao *et al.* 2018). As a relatively new

methodology, biodiversity studies that employ LiDAR have been limited in scope, typically

addressing only one taxonomic group or habitat type at the local scale, and strongly biased

towards forest ecosystems. However, studies using LiDAR-based indicators have already

been shown to explain up to 66% and 82% of local plant and fungi richness, respectively

(Lopatin *et al.* 2016, Peura *et al.* 2016, Thers *et al.* 2017). A recent study has demonstrated its

potential to provide spatially accurate and comprehensive measures by predicting the local

biodiversity of different taxonomic groups (plants, fungi, lichens, and bryophytes) across

multiple habitat types and large geographic extent (Moeslund *et al.* 2019). LiDAR may also

be a useful tool in studying dark diversity by incorporating potentially important

spatiotemporal dynamics such as succession and disturbance (Mokany and Shine 2003, Scott

et al. 2011, Pärtel *et al.* 2013). For example, recent studies have found that human impact

increases dark diversity in arbuscular mycorrhizal fungi (Pärtel *et al.* 2017a), that ruderal

plants are more likely to be in dark diversity (Moeslund *et al.* 2017), and that human density

and agricultural land-use influence dark diversity of vascular plants (Riibak *et al.* 2017).

Alongside these structural and environmental factors, fungal diversity depends on

biotic interactions, with a large proportion of fungi deriving their nutrients and carbon from

host plants (Tedersoo *et al.* 2014, Nguyen *et al.* 2016). Recent evidence has hinted on the

influence of these interactions on dark diversity, as plant species dependent on mycorrhiza

have been found to have greater dark diversity than those without these mutualist

relationships (Moeslund *et al.* 2017). Moreover, ectomycorrhizal fungal diversity seems to

increase exponentially with an increasing proportion of their host plants, suggesting that

competitive interactions among fungi might also drive their dark diversity (Pärtel *et al.*

2017b). Typical and strong species interactions are indeed typically considered in dark

diversity, as the estimation hereof is usually based on species co-occurrences (Beals 1984,

McCune 1994, Münzbergová and Herben 2004, de Bello *et al.* 2012, Lewis *et al.* 2016).

However, this is usually done only within the species group being studied. For example, in

studies of plant dark diversity, only co-occurrences with other plants, and not fungi or other

species groups, are typically considered. However, recognizing the close and interconnected

relationship between plants and fungi allows for stronger and more realistic estimations of the

fungal dark diversity. Incorporating other taxonomic groups when determining species pools

and estimating dark diversity is not a new insight, and the importance of biotic interactions

across trophic groups has been discussed since the concept of dark diversity was first

introduced (Pärtel *et al.* 2011). Yet, such cross-species group data has never been included in

dark diversity estimates meaning that they may not sufficiently account for cross-species

group interactions, and this may be part of the explanation for why dark diversity is

sometimes over-estimated (Boussarie *et al.* 2018).

In this study, we examined a number of environmental factors influencing the local

dark diversity of fungi across habitat-types nationwide within Denmark. We used a

comprehensive biodiversity datasets covering major environmental gradients (Brunbjerg *et*

al. 2019) and combined it with LiDAR-based measurements. We also included fungi-plant-

co-occurrence information to estimate local fungal dark diversity and thereby acknowledge

the importance of their biotic interactions. More specifically, we addressed the following

questions: 1. To what degree can we explain local fungal dark diversity by abiotic and biotic

environmental factors? 2. Can vegetation and terrain structural factors that are important to

local fungal dark diversity be derived from LiDAR and if so, 3. how important are they

compared to field-measured factors?

Methods

Study area and site selection

The dataset was collected from a national biodiversity inventory in Denmark as part

of the "Biowide" research project (Brunbjerg *et al.* 2019). A total of 130 study sites (40 × 40

m) were selected with a minimum distance of 500 m between each to reduce spatial

covariance with 30 sites allocated to cultivated habitats and 100 sites to natural habitats

(Figure 1). The cultivated subset was stratified according to the type of land use and the

natural subset was selected amongst uncultivated habitats and stratified according to

gradients in soil fertility, soil moisture, and successional stage. The "Biowide" project

deliberately excluded saline and aquatic habitats but included temporarily inundated
depressions along with mires, bogs, and fens. The final set of 24 habitat strata consisted of
three types of fields (rotational, grass leys, set aside) and three types of plantations (beech,
oak, spruce), and the remaining 18 strata were natural or semi-natural habitats, constituting
all possible combinations of positions along three major natural environmental gradients: soil
fertility (rich, poor), soil moisture (dry, moist, wet), and successional stage (early, mid, late).
These 24 strata were replicated in five geographical regions in Denmark. The "Biowide"
dataset also includes a subset of 10 sites (two in each region) of hotspots for different
taxonomic groups in Denmark, which were selected by voting amongst active naturalists in
the Danish conservation and management societies. For further details on the design and data
collection procedures see Brunbjerg *et al.* (2019).

172
Figure 1. The (A) 130 selected study sites from a national biodiversity inventory and (B) the

four quadrants and 5□m circle plots within each site. Reprinted from Ejrnæs *et al.* (2018),

with permission from Elsevier.

Field-measured variables

We used fungi observational data from the "Biowide" field inventories (summarized

data and additional details in Brunbjerg *et al.* (2019)). Macro-fungal species were surveyed in

2014–2015 by an expert field-mycologist and volunteers during three inventories (up to one

hour per site) in the main fruiting season (August - November) by actively searching
microhabitats and substrates (soil, herbaceous vegetation and debris, dead wood, litter, and
bark of trees up to 2 m) within the 40 × 40 m sites. Since truffles are difficult to find, we did
not consider these in this study. Subspecies and varieties were lumped to the species level.
After pre-processing, the dataset consisted of 6,269 observations of 1,017 species.

Vascular plant species observations were also taken from the "Biowide" database and
were originally inventoried by trained botanists during the summer 2014 and spring 2015 to
account for variations in phenology. We removed hybrids and neophytes (i.e. species that are
not considered a natural part of the vegetation given their history and dispersal ability, see
appendix tables 6–8 in Buchwald *et al.* (2013)) and lumped subspecies and varieties to the
species level. Species nomenclature for both plants and fungi follow the species checklist of
Denmark (allearter.dk).

Apart from the LiDAR-based measures (detailed below), we also considered field-
measured variables representing both abiotic conditions and available biotic resources known
to influence fungal diversity and communities (Table 1). For further details on data collection
and how the environmental field measurements were made see Brunbjerg *et al.* (2019).

Table 1. Overview of the explanatory variables for fungal dark diversity models along with our hypothesized relationship with dark

diversity. If the standard deviation of a variable was calculated, in addition to its mean, the variable is denoted with an asterisk.

	Explanatory variables	Hypothesis	References
Field-based measures	Plant richness	High plant richness could mean higher community stability or reflect low human impact. This would result in a lower dark diversity	(Kuiters 2013, Pellkofer *et al.* 2016, Yang *et al.* 2018)
	Litter (open habitats) and dead wood (forests)	The more substrate represented by more organic matter gives more resources for fungi and hence a lower dark diversity. Alternatively, organic matter increases competition between fungi and soil bacteria or causes many unfilled microhabitats which would increase dark diversity	(Leigh *et al.* 2011, Averill *et al.* 2014)
	Soil pH	Despite increasing importance of bacteria and invertebrates in decomposition with increasing soil pH, we also expect less dominance and better opportunities for fungal colonization as the base-rich soil environment is less restrictive, where only few species can cope with low pH values, possibly resulting in a lower dark diversity at high pH	(Clark 1997, Rousk *et al.* 2009, Rousk *et al.* 2010)
	Soil fertility index (SFI)	Asymmetric competition among vascular plants	(Buckland and Grime 2000, Liu *et*

		increases with increasing soil fertility leading to loss of plant richness at high fertility levels and therefore potentially also increasing dark diversity	*al.* 2015, Nadeau and Sullivan 2015, Luo *et al.* 2017)
	Soil moisture index (SMI)	Most fungi thrive at intermediate soil moisture levels, but have a rapid and opportunistic growth response to high moisture. Assuming establishment is easier in intermediate moist environments dark diversity could be higher at high moisture. Alternatively, dark diversity could be high at intermediate moisture levels as this could yield many unfilled microhabitats	(Jacobson 1997, Kennedy and Peay 2007)
LiDAR-based measures	Vegetation height*	Taller vegetation could reflect encroachment by shrubs and trees in open habitats and forests resulting in more available microhabitats for potential fungal species increasing their dark diversity	(Zuo *et al.* 2016, Gómez *et al.* 2019)
	Succession (Amplitude)	Amplitude could reflect successional processes with later successional stages allowing fungi to become more established, resulting in lower dark diversity	(Fernández-Toirán *et al.* 2006, Twieg *et al.* 2007, Hui *et al.* 2017)

Microtopography -Terrain roughness (SigmaZ) -Terrain openness*	Microtopography increases availability of microhabitats and hence could increase dark diversity	(Cantelmo Jr and Ehrenfeld 1999)
Light/heat - Canopy openness (forests)* -Heat load index* -Solar irradiation* -Vegetation cover*	Light availability increases fungal establishment success which would decrease dark diversity	(Graham *et al.* 1982, Turner *et al.* 2009)
Canopy complexity -Echo ratio*	Canopy complexity provides more niches directly in the canopy and could also be expected to be associated with a higher variation in the forest floor with small openings in the canopy and variation in tree ages. This could increase the number of unfilled microhabitats and hence the dark diversity	(Unterseher and Tal 2006, Gómez-Hernández and Williams-Linera 2011, Dove and Keeton 2015)

198 ***LiDAR-based measures***

199  To enable the calculation of measures representing vegetation and terrain environmental

and structural aspects, we used the latest nationally covering LiDAR-based point cloud for

Denmark from the Danish Ministry of Environment. This dataset is freely available from

www.kortforsyningen.dk and has a point density of 4-5 points/m^2. Originally, this dataset was

recorded from fixed-wing airplanes at an altitude of approximately 680 m above ground level

and a speed over ground of approximately 240 km/h. The data was recorded by Riegl LMS-680i

scanners operating in the near-infrared wavelength (1550 nm) in a parallel line scan pattern

during the springs and autumns of 2014 and 2015. For all calculations, we relied on the

classification of points into ground, building and vegetation classes already present in the data

set upon download.

209  To represent vegetation and terrain environmental and structural aspects, we calculated

observed measures based on the point cloud data set. We calculated all measures at 1.5 m

resolution (except for terrain roughness which was at 0.5 m resolution) and their means and

standard deviations within 30 m radius circles centered in each study site. For all LiDAR

processing and calculation, we used the OPALS tools (Pfeifer *et al.* 2014) version 2.3.1 in a

Python 2.7 environment.

Vegetation-related measures

216  To represent *succession* and to some degree moisture balance in both vegetation and soil,

we used the amplitude of each echo representing a point in the LiDAR point cloud. This

amplitude is high if the reflecting surface is flat (i.e., smooth) and with high reflectivity. It is low

when the light energy is distributed between several returns for example in tree canopies, or

when surfaces have low reflectivity, are complex, or translucent (e.g., leaves). The wavelength

used to record the point cloud data is sensitive to leaf water content (Junttila *et al.* 2018) and soil

moisture (Zlinszky *et al.* 2014). Since the amplitude depends on reflectivity, which varies across

months and aircraft types (slightly different flying heights) used for data recording, the amplitude

was corrected to account for these biases. We constructed a Generalized Linear Model (GLM)

with Gaussian link having the raw amplitude as response and flight month as well as aircraft type

as explanatory factors and used only the residuals of this model for input in our statistical

modelling. We also tried using flight year as an explanatory factor, but this did not improve the

model (ΔAIC < 2). These residuals will be referred to as the *corrected amplitude* in the

following. Unfortunately, we did not have reference data enabling a full calibration of this

measure (Höfle and Pfeifer 2007).

To represent *vegetation height*, we estimated this measure by subtracting the terrain

model from the surface model (two raster files, detailed in the following). The terrain model

(DTM) calculation details are given in the section on "Terrain-structure measure". The surface

model was calculated using the DSM module in OPALS using all vegetation and ground points.

To reflect the penetrability and succession of the vegetation we calculated the *echo ratio*

(Höfle *et al.* 2012). Echo ratio is high where the surface is impenetrable and relatively smooth

and lower where the surface is uneven or penetrable. In order to calculate the echo ratio,

estimating normals for each point is required. We did this using the Normals module in OPALS

with a robust plane fit based on the 12 nearest neighboring points. Subsequently, we calculated

the echo ratio for each terrain and vegetation point using a search radius of 1.5 m along with the

slope adaptive method implemented in the EchoRatio module of OPALS.

To estimate light conditions, we calculated the *canopy openness* for all points categorized

as "ground", but contrary to terrain openness (see below), we calculated this considering

vegetation points as well. Therefore, canopy openness represents the actual blocking of the sky

view by the canopy around each ground point. Canopy openness is high for ground points inside

canopy gaps and low for ground points beneath a closed canopy.

Lastly, as an estimate of *vegetation cover*, we calculated the fraction of vegetation points

to all points (excluding unclassified points and those representing buildings and noise). This

measure will be high if the vegetation is dense or the cover of vegetation is relatively high, and

low for areas with no vegetation.

Terrain-structure measures

To enable the calculation of several terrain-related measures, we calculated a digital

terrain model (DTM) for each study site representing the elevation above sea level. To do this we

used the DTM module of OPALS based on only ground points. We set the module to use 8

neighboring points and a search radius of 6 m. To represent key features of the local terrain (e.g.,

soil moisture or heat balance (Moeslund *et al.* 2013)), we calculated *terrain slope* and *terrain*

aspect (used for heat load index calculation, see below). For this task, we used the GridFeature

module of OPALS using the DTM as input, a kernel size of 1 and requesting the terrain slope

and aspect (slope direction) in radians.

To reflect local heat input, we calculated the *heat load index* based on the terrain aspect

following the heat load index formula in McCune and Keon (2002). This index reaches

maximum values on southwest-facing slopes and zero on northeast-facing slopes. We also

calculated the potential *solar irradiation* based on terrain slope, aspect, and latitude following

equation 3 in McCune and Keon (2002).

To estimate micro-scale terrain heterogeneity, we calculated the *terrain roughness*

(SigmaZ) using only ground points as input. This measure represents the standard deviation of

the interpolated grid height. The OPALS DTM module outputs this measure as a by-product

when constructing a DTM. However, unlike the rest of the LiDAR measures in this study, the

terrain roughness was calculated at 0.5×0.5 m resolution mirroring micro-scale terrain

variations.

To represent site-scale terrain heterogeneity, we calculated the *terrain openness* (Doneus

2013). Terrain openness is defined as the opening angle of a cone (having the radius of the

kernel) turned upside down – with its tip restrained to the point of interest – that touches the

terrain surface. To calculate this, we used the PointStats module of OPALS requesting "positive

openness" based on only ground points and a search radius of 5 m. This measure is high in flat

(relative to the scale at which it is calculated) areas and low in heterogeneous terrains.

Finally, to test the importance of variability in the LiDAR measures we calculated the

standard deviation for LiDAR measures for which we believed it made ecological sense (Table

1).

Data analysis

Data preparation

Prior to statistical analysis, we removed the six intensively managed fields from the study

sites, as these were ploughed fields. We also removed two study sites because they were flooded

during the LiDAR data recording period. Finally, we removed one site due to an extreme outlier

285 in the LiDAR amplitude values (300 vs. a range of values between 10 and 130). Our final dataset
286 therefore comprised a total of 121 study sites.

287   Our initial visual inspection of the data revealed that many of the LiDAR measures were
288 relevant only for woodlands and therefore strongly zero-inflated in the open landscapes. The
289 analyses in this study were therefore separately run for open habitats and woodlands. Open
290 habitats included grasslands, fens, bogs, and other habitats with only few sporadic occurrences of
291 trees. The woodlands dataset consisted of forests, thickets and shrubland (e.g. willow).

292   In the following, we detail the steps we took to prepare the LiDAR and measured
293 variables for statistical modelling as explanatory factors. Obviously, a number of these variables
294 were strongly inter-correlated (Appendix 1). For example, echo ratio was strongly related to
295 canopy and light measures (Appendix 1). Therefore, we selected only those variables that we
296 hypothesized to affect local fungal dark diversity (See Table 1). Subsequently, to avoid issues
297 with multi-collinearity we calculated Variance Inflation Factors (VIFs) causing us to remove
298 vegetation height as an explanatory factor for the open landscapes to ensure VIF values below 10
299 following Kutner *et al.* (2005). Subsequently, the maximum VIF value of explanatory factors
300 used together in the same models was 4.8 and 5.7 for woodlands and open habitats respectively.
301 We scaled all explanatory variables to a mean of zero and a standard deviation of one. To strive
302 for normal distribution of explanatory variables we log- or square-root transformed those where
303 this made obvious distributional improvements based on visual examination of the histograms.

305　**Appendix 1.** Correlation plot between environmental and LiDAR variables. All significant
306　interactions are colored red for negative relationships and blue for positive relationships, with the
307　size and darkness of the color representing the strength of the relationship. Non-significant
308　correlations are blank.

Dark diversity

All statistical analyses mentioned in the following were performed in R version 3.5.3 (R
Core Team 2019). To calculate the dark diversity for each site, we first calculated the site-
specific species pool using the Beals' index (Beals 1984), as recommended by Lewis *et al.*

(2016), using the 'beals' function in the 'vegan' package (Oksanen *et al.* 2017). The Beals' index

represents the probability that a particular species will occur within a given site based on the

assemblage of co-occurring species (Beals 1984, McCune 1994, Münzbergová and Herben

2004). The threshold used for including a particular species in the site-specific species pool was

the 5^{th} percentile of the Beals' index value for each species (Gijbels *et al.* 2012, Ronk *et al.*

2015). Preceding the calculation of each threshold, the lowest Beals' index value among plots

with the occurrence of the species in question was identified, and all plots having values below

the minimum were not considered. We calculated two measures of the site-specific species pool

for each site: (1) using only fungi co-occurrence and (2) co-occurrences of both observed fungi

and vascular plants at each site to acknowledge the fungal-plant linkages (i.e., both fungi and

plant species were in the presence/absence matrix used to calculate Beals' index). Dark diversity

was calculated by subtracting observed fungal species richness from the site-specific species

pool. Since site-specific species pools differ between sites, we calculated the *relative dark*

diversity for each site as dark diversity (species predicted from the site-specific species pool but

not observed) divided by the regional pool to enable comparison of results across habitats.

Statistical analysis

To investigate what characterizes sites with a high fungal dark diversity we constructed

GLMs with a Gaussian link having the estimated relative dark diversity (described above) as the

response variable. We constructed models for both open habitats and woodlands, and for both

dark diversity estimates (see the section on dark diversity). Initially, we fitted models using only

the LiDAR measures as explanatory factors, to test the degree to which fungal dark diversity

patterns could be explained using LiDAR data alone. Subsequently, we fitted a similar model

with both measured and LiDAR variables as explanatory factors (Table 1), giving insight into

how much more explanatory power one gains by using measured variables in addition to LiDAR.

To allow for non-linear relationships for variables corresponding to the intermediate disturbance

hypothesis (Connell 1978, Townsend *et al.* 1997) and intermediate productivity hypothesis

(Fraser *et al.* 2015), we used Akaike's Information Criterion (AIC) (Burnham and Anderson

2002) to evaluate if inclusion of squared terms for the variables SMI, SFI, light, soil pH, and

bare soil (see Table 1) improved the model fit. If so, we kept the squared term of the variable in

question instead of the linear effect. After the initial fit and checking for non-linearity as

described above, we ran a backward model selection procedure for each model based on AIC.

The procedure stopped when AIC did not drop anymore and ΔAIC was above 2 (Burnham and

Anderson 2002). In each iteration, we dropped the variable causing the smallest change in AIC

value. As a final step, we checked model residuals to ensure that these were normally distributed.

We did not conduct spatial modelling in this study, since the original field-work design lying

behind the data was designed to avoid spatial autocorrelation, and tests for all the species groups

originally inventoried concluded that spatial signals were of minor to little importance (see

Brunbjerg *et al.* (2019), and also "Study area and site selection").

Results

The two relative fungal dark diversity estimates (based on fungi-only and both fungi- and

plant co-occurrences) were between 0.17 – 0.93 (open habitats, median 0.51) and 0.20 – 0.63

(woodland, median: 0.39); and 0.21 – 0.95 (open habitats, median: 0.56) and 0.24 – 0.7

(woodland, median: 0.44) respectively. In most cases, our models explained between 20-30 % of

the variation in fungal dark diversity and more than 40 % for the woodlands models when

including both LiDAR and measured variables (Table 2). In the "fungi-only dark diversity"

model for woodland habitats (Table 2) the squared term of soil pH did improve model fit, but

this term was left out during the subsequent model selection procedure.

The only LiDAR variable significant in both open habitats and woodlands was amplitude,

which was significant in all models for woodlands and in LiDAR-only models for open habitats

(Table 2). This variable had a positive effect on dark diversity in woodlands (Table 2) but a

negative influence in open habitats (Table 2). Echo ratio was the only other significant LiDAR

variable in our analyses and positively influenced dark diversity in open habitats (Table 2). Plant

richness was negatively related to local fungal dark diversity and had the strongest impact of all

the field-measured factors included in our analyses (Table 2). Also, soil fertility and moisture

were positively correlated with fungal dark diversity in open habitats (Table 2).

In all cases, models considering only the structural environment (LiDAR only) were

outperformed by models considering plant richness and the abiotic environment in addition to

the structural, notably in the open habitats (Table 2). Appendix 1 shows all pair-wise Spearman

correlations between the explanatory variables used and Appendix 2 shows observed fungi

richness against the two dark diversity measures for both open and woodland habitats.

Table 2. Modelling coefficients for the best models (i.e. after model selection) regressing dark diversity estimates based on only fungi

co-occurrences (*Fungi-only dark diversity*) or based on both fungi and plant co-occurrences (*Fungi-plant dark diversity*) against the

selected explanatory variables. Significant variables were from either a LiDAR-only or a full model with both LiDAR-based and field-

measured predictors.

| | | | R^2 | LiDAR variables | | Field-measured variables | | |
				Amplitude	Echo ratio	Plant richness	Soil fertility	Soil moisture
Woodland habitats	Fungi-only dark diversity	Lidar-only model	0.24	0.03	-	-	-	-
		Full model	0.40	0.04	-	-0.05	-	-
	Fungi-plant dark diversity	Lidar-only model	0.23	0.03	-	-	-	-
		Full model	0.44	0.04	-	-0.05	-	-
Open habitats	Fungi-only dark diversity	Lidar-only model	0.10	-0.05	0.01	-	-	-
		Full model	0.30	-	0.01	-0.08	0.06	0.04
	Fungi-plant dark diversity	Lidar-only model	0.07	-0.04	0.01	-	-	-
		Full model	0.34	-	0.01	-0.10	0.06	-

380 **Appendix 2.** Relationship between observed fungal diversity and dark diversity estimates in
381 woodland based on (A) only fungi co-occurrences (*Fungi-only dark diversity*) or (B) fungi and
382 plant co-occurrences (*Fungi-plant dark diversity*), and in open habitats based on (C) only fungi
383 co-occurrences (*Fungi-only dark diversity*) or (D) fungi and plant co-occurrences (*Fungi-plant*
384 *dark* *diversity*).

Discussion

In this study, we demonstrate for the first time that LiDAR derived variables, alone and
in combination with field-measured variables, can explain a significant amount of the variation
in local dark diversity of temperate macro-fungal communities. Our findings indicate that the
dark diversity of fungi is influenced by the local vegetation structure, plant associations, and the
abiotic environment. This is not surprising since local observed fungal diversity is also
determined by these factors to a large degree (Thers *et al.* 2017, Yang *et al.* 2017, Moeslund *et*
al. 2019). Indeed, the relative dark diversity analyzed in this study correlated with the observed
fungal species richness (Appendix 2), which was expected given that the dark diversity
calculations are based on the species present (Noreika *et al.* 2020). On the other hand, there are

differences between these two measures, so the results presented here are not necessarily

applicable to the observed fungal diversity. We also find that models including field-based

variables explained the dark diversity of fungi far better than models relying solely on LiDAR,

notably in open landscapes. While LiDAR has the advantage that one can record data from huge

areas in very fine detail for relatively low cost, our results indicates that to get the best

explanation of local fungal diversity patterns fieldwork is still needed.

LiDAR-based measures

This study shows that LiDAR captures habitat characteristics potentially important for

fungal dark diversity which are not represented by traditional field-measured variables. Notably,

the relationship between fungal dark diversity and LiDAR-derived vegetation structure in

woodlands was relatively strong. Although LiDAR can successfully quantify biophysical

characteristics in all types of habitats, it is known to be more effective in forested habitats (Su

and Bork 2007), supporting these findings. The most important LiDAR variables for modelling

fungal dark diversity were amplitude and echo ratio which gives us important insights into what

environmental aspects, which are not typically recorded in field-surveys, can be captured by a

LiDAR approach.

The LiDAR measure of amplitude is sensitive both to surface reflectivity and to the

number of targets hit by the laser pulse (Moeslund *et al.* 2019). The lower the reflectivity and the

more targets between which the light energy is distributed, the lower the amplitude associated

with a given point. This would result in high amplitudes in flatter surfaces while yielding low

amplitudes in tall and more complex canopies, or translucent surfaces such as leaves. Hence, this

variable can be a proxy for succession stage, surface evenness, or vegetation density; since both

flat and sparsely vegetated as well as densely vegetated canopies preventing light penetration

will yield high amplitude. Supporting this, amplitude was positively correlated with vegetation

height and vegetation cover (denser vegetation resulted in higher amplitude) and negatively

correlated with echo ratio (vegetation complexity, see below) and canopy openness. In

woodlands, the dark diversity of fungi was positively related to amplitude, suggesting that more

species are missing in the relatively tall and dense forests compared to more complex woodlands

(e.g., with canopy openings). The positive association between LiDAR amplitude and dark

diversity could therefore be a consequence of communities in older well-developed shrubland or

old-growth forests with windthrows or other openings, having allowed fungi more time to
become established with their associated plants (Fernández-Toirán *et al.* 2006, Twieg *et al.*
2007). Indeed, among the top half of woodland plots with regards to amplitude were plantations
(which often have dense canopy and equal-aged trees) and most of them contained high relative
fungal dark diversity, while the bottom half of the plots, those having the lowest amplitude and
dark diversity, were mostly old forests or shrublands with a more well-developed vegetation
structure (e.g., dead or fallen wood, or complex sub-canopy layer).

In open habitats, amplitude and echo-ratio were negatively and positively related to
fungal dark diversity, respectively. These results indicate that fewer species are missing from the
more even early-successional grasslands without trees and shrubs. We suggest this could be the
result of encroachment due to the widespread abandonment of ancient grassland management
practices resulting in a loss of small-statured typical grassland plant species without a
corresponding gain in species associated with scrub and woodland. It could also reflect that
fewer species are missing from calcareous or sandy grasslands since open limestone and white
sand have a relatively high reflectivity.

Plant richness

The most important field-measured variable for modelling fungal dark diversity was plant
species richness which was negatively related to dark diversity in both open habitats and
woodlands. Plant richness and composition are well-known to correlate with fungal richness and
composition (Zak *et al.* 2003, Chen *et al.* 2017, Yang *et al.* 2017, Brunbjerg *et al.* 2018, Wang *et*
al. 2018), and sites with lower plant species richness have previously been found to have a
relatively higher proportion of plants in the dark diversity (Fløjgaard *et al.* 2020). Here, the
negative relation between plant richness and fungal dark diversity may be attributed to greater
plant richness frequently associated with more stable communities and ecosystems (Kuiters
2013, Pellkofer *et al.* 2016, Yang *et al.* 2018), which could indicate longer continuity and hence
time for fungi to establish. Alternatively, host specific fungi species could be missing due to
absence of their symbiotic plant species (Dickie 2007). However, in our study, plant richness had
almost the same effect on both fungal dark diversity accounting for present plant species and
where these presences where unaccounted for. This points to plant hosts playing a minor role for
fungal dark diversity, which is unsurprising as it likely indicates that the calculation of fungal

dark diversity using the Beals' index does not allow fungi having non-present hosts into the site-

specific species pool. Another possible explanation is that plant richness mirrors human impact.

Generally plant species richness have declined over several decades and continue to as a

consequence of agricultural intensification and abandonment of extensive land-use (Hülber *et al.*

2017). Other studies have found human disturbance to be strongly related to fungal richness and

dark diversity patterns (Epp Schmidt *et al.* 2017, Pärtel *et al.* 2017a), and future studies may help

to tease apart these effects.

Abiotic environment

Soil fertility is often associated with fungal diversity (Balser *et al.* 2005, Kalliokoski *et*

al. 2010, Sterkenburg *et al.* 2015) and was found to have a positive relationship with dark

diversity in open habitats. In general, soil fertility influences plant species richness negatively

through asymmetric competition (Buckland and Grime 2000, Dybzinski *et al.* 2008, Nadeau and

Sullivan 2015, Luo *et al.* 2017). This could explain the negative relationship between the local

dark diversity of fungi and soil fertility: lower plant species richness yields a higher dark

diversity (see discussion above on "Plant richness"). However, the effect might also be

uncoupled from plants and simply due to changes in the soil decomposition microbiota from

fungal to bacterial dominance along a gradient of soil fertility and pH (Blagodatskaya and

Anderson 1998). Another alternative explanation is that soil fertility affects the density of soil

mycophagous and microarthropod species (Cole *et al.* 2005) which also may affect fungal dark

diversity (Crowther *et al.* 2013). However, while this explanation might be plausible, the

underlying mechanisms are largely unknown, calling for further research to dissect the

interactions between soil fertility, soil microarthropods and fungal diversity.

We also found soil moisture had a positive relationship with fungal dark diversity in open

habitats. Moisture is known to influence fungal communities (Gómez-Hernández and Williams-

Linera 2011, Gupta *et al.* 2018, Frøslev *et al.* 2019) as it affects the growth, colonization rate,

and spore and fruit body production (Salusso and Moraña 1995, Jacobson 1997, Kennedy and

Peay 2007). The relationship between fungi and plants is probably important in this regard

because soil moisture causes a high turnover in plant species composition (e.g., Moeslund *et al.*

(2013), and in turn, affects the quality and availability of resources for below-ground fungal

communities (Chen *et al.* 2017). Additionally, high soil moisture is a strong environmental filter

excluding most macro-fungi species from the wet habitats (Heilmann-Clausen *et al.* 2019). This
filter may also be the main reason for the higher fungal dark diversity found in the wet habitats
when plant co-occurrences was not included in determining the site-specific species pool. On the
other hand, the interdependencies between fungi and plants, and the strong link between plant
communities and soil moisture gradients (Xiong *et al.* 2003, Silvertown *et al.* 2015, Valdez *et al.*
2019), may explain why soil moisture was not significant in models where plant co-occurrences
was included in determining the site-specific species pool, as this approach perhaps accounts for
these interactions.

Uncertainties

One drawback of basing a study on an organism group like fungi, which live a mostly
hidden life, is the unavoidable issues concerning overlooked species and hidden diversity in
general (Milberg *et al.* 2008, Abrego *et al.* 2016). If such errors are biased towards specific plots
our results could be affected. However, the field-work behind the dataset used here was planned
to avoid this exact issue by including several visits at each site at different times of the year, and
we do not believe this confounds our findings. Nevertheless, there is always uncertainty when
estimating unknowns, such as which species are actually absent despite the fact that they could
indeed be in a given site. In this study we used on of the most comprehensive and best possible
data set along with state-of-the-art methods to calculate these estimates (see e.g. (Lewis *et al.*
2016, Moeslund *et al.* 2017), and we therefore believe our results are indeed sound and realistic
despite this uncertainty.

Conclusion

This is the first study to investigate potential drivers of the local dark diversity of fungi
using both LiDAR derived vegetation and terrain structure as well as field-measured variables.
We showed that local fungal dark diversity is strongly dependent on the environment with
vegetation structure, plant diversity, and abiotic factors playing important roles. Also, to our
knowledge, this is the first study using cross-species group co-occurrence data to determine
species pools. This may be a more ecologically sound methodology than using only one taxon
group, especially for interdependent taxonomic groups. Future studies and novel approaches will
be required to unravel the causal links between fungal communities and habitat and vegetation
characteristics; and to gain a better understanding of how LIDAR-based measures can be

interpreted as measures of vegetation and terrain structure. Using LiDAR as a tool to determine dark diversity, in conjunction with ecological field measurements, may provide a valuable tool to better guide conservation and restoration planning by identifying sites with high restoration potential (high dark diversity) and high priority for conservation (low dark diversity, or sites where the fungal communities are more "complete").

Acknowledgements

We thank Thomas Læssøe for collecting and identifying macro-fungi. We sincerely thank Aage V. Jensen Nature Fund for financial support to CF, AKB, JM, LD, KC and JV through the project "Dark Diversity in Nature Management". The Biowide project and REJ was supported by a grant from the Villum Foundation (VKR-023343). MP has been supported by the Estonian Ministry of Education and Research (IUT20–29), and the European Regional Development Fund (Centre of Excellence EcolChange). The authors declare no conflict of interest.

Literature Cited

Abrego, N., Halme, P., Purhonen, J. and Ovaskainen, O. (2016) 'Fruit body based inventories in wood-inhabiting fungi: Should we replicate in space or time?', *Fungal Ecology*, 20, 225-232.

Averill, C., Turner, B. L. and Finzi, A. C. (2014) 'Mycorrhiza-mediated competition between plants and decomposers drives soil carbon storage', *Nature*, 505(7484), 543-545.

Balser, T. C., Treseder, K. K. and Ekenler, M. (2005) 'Using lipid analysis and hyphal length to quantify AM and saprotrophic fungal abundance along a soil chronosequence', *Soil Biology and Biochemistry*, 37(3), 601-604.

Beals, E. W. (1984) 'Bray-Curtis ordination: An effective strategy for analysis of multivariate ecological data' in MacFadyen, A. and Ford, E. D., eds., *Advances in Ecological Research* Academic Press, 1-55.

Blackwell, M. and Vega, F. E. (2018) 'Lives within lives: Hidden fungal biodiversity and the importance of conservation', *Fungal Ecology*, 35, 127-134.

Blagodatskaya, E. V. and Anderson, T.-H. (1998) 'Interactive effects of pH and substrate quality on the fungal-to-bacterial ratio and qCO2 of microbial communities in forest soils', *Soil Biology and Biochemistry*, 30(10-11), 1269-1274.

551 Boussarie, G., Bakker, J., Wangensteen, O. S., Mariani, S., Bonnin, L., Juhel, J.-B., Kiszka, J. J.,
552   Kulbicki, M., Manel, S. and Robbins, W. D. (2018) 'Environmental DNA illuminates the
553   dark diversity of sharks', *Science advances*, 4(5), eaap9661.

555 Brunbjerg, A. K., Bruun, H. H., Broendum, L., Classen, A. T., Fog, K., Froeslev, T. G.,
556   Goldberg, I., Hansen, M. D. D., Hoeye, T. T., Laessoee, T., Newman, G., Skipper, L.,
557   Soechting, U. and Ejrnaes, R. (2019) 'A systematic survey of regional multitaxon
558   biodiversity: evaluating strategies and coverage', *BMC Ecology*, 19(43), 158030.

560 Brunbjerg, A. K., Bruun, H. H., Dalby, L., Fløjgaard, C., Frøslev, T. G., Høye, T. T., Goldberg,
561   I., Læssøe, T., Hansen, M. D. and Brøndum, L. (2018) 'Vascular plant species richness
562   and bioindication predict multi☐taxon species richness', *Methods in Ecology and*
563   *Evolution*, 9(12), 2372-2382.

565 Brunbjerg, A. K., Bruun, H. H., Moeslund, J. E., Sadler, J. P., Svenning, J.-C. and Ejrnæs, R.
566   (2017) 'Ecospace: A unified framework for understanding variation in terrestrial
567   biodiversity', *Basic and Applied Ecology*, 18, 86-94.

569 Buchwald, E., Wind, P., Bruun, H. H., Møller, P. F., Ejrnæs, R. and Svart, H. E. (2013) 'Hvilke
570   planter er hjemmehørende i Danmark?', *Flora & Fauna*, 118, 73-96.

572 Buckland, S. M. and Grime, J. P. (2000) 'The effects of trophic structure and soil fertility on the
573   assembly of plant communities: a microcosm experiment', *Oikos*, 91(2), 336-352.

575 Burnham, K. P. and Anderson, D. R. (2002) *Model Selection and Multi-model Inference: A*
576   *Practical Information-Theoretic Approach*, New York: Springer.

578 Cantelmo Jr, A. J. and Ehrenfeld, J. G. (1999) 'Effects of microtopography on mycorrhizal
579   infection in Atlantic white cedar (Chamaecyparis thyoides (L.) Mills.)', *Mycorrhiza*, 8(4),
580   175-180.

582 Chen, Y.-L., Xu, T.-L., Veresoglou, S. D., Hu, H.-W., Hao, Z.-P., Hu, Y.-J., Liu, L., Deng, Y.,
583   Rillig, M. C. and Chen, B.-D. (2017) 'Plant diversity represents the prevalent determinant
584   of soil fungal community structure across temperate grasslands in northern China', *Soil*
585   *Biology and Biochemistry*, 110, 12-21.

587 Clark, R. B. (1997) 'Arbuscular mycorrhizal adaptation, spore germination, root colonization,
588   and host plant growth and mineral acquisition at low pH', *Plant and Soil*, 192(1), 15-22.

Cole, L., Buckland, S. M. and Bardgett, R. D. (2005) 'Relating microarthropod community structure and diversity to soil fertility manipulations in temperate grassland', *Soil Biology and Biochemistry*, 37(9), 1707-1717.

Connell, J. H. (1978) 'Diversity in tropical rain forests and coral reefs', *Science*, 199(4335), 1302-1310.

Cornell, H. V. and Harrison, S. P. (2014) 'What are species pools and when are they important?', *Annual Review of Ecology, Evolution, and Systematics*, 45(1), 45-67.

Crowther, T. W., Stanton, D. W., Thomas, S. M., A'Bear, A. D., Hiscox, J., Jones, T. H., Voříšková, J., Baldrian, P. and Boddy, L. (2013) 'Top☐down control of soil fungal community composition by a globally distributed keystone consumer', *Ecology*, 94(11), 2518-2528.

de Bello, F., Price, J. N., Münkemüller, T., Liira, J., Zobel, M., Thuiller, W., Gerhold, P., Götzenberger, L., Lavergne, S. and Lepš, J. (2012) 'Functional species pool framework to test for biotic effects on community assembly', *Ecology*, 93(10), 2263-2273.

DeAngelis, D. L. (2012) *Dynamics of nutrient cycling and food webs*, Springer Science & Business Media.

Dickie, I. A. (2007) 'Host preference, niches and fungal diversity', *The New Phytologist*, 174(2), 230-233.

Doneus, M. (2013) 'Openness as visualization technique for interpretative mapping of airborne lidar derived digital terrain models', *Remote Sensing*, 5(12), 6427-6442.

Dove, N. C. and Keeton, W. S. (2015) 'Structural Complexity Enhancement increases fungal species richness in northern hardwood forests', *Fungal Ecology*, 13, 181-192.

Dybzinski, R., Fargione, J. E., Zak, D. R., Fornara, D. and Tilman, D. (2008) 'Soil fertility increases with plant species diversity in a long-term biodiversity experiment', *Oecologia*, 158(1), 85-93.

Ejrnæs, R., Frøslev, T. G., Høye, T. T., Kjøller, R., Oddershede, A., Brunbjerg, A. K., Hansen, A. J. and Bruun, H. H. (2018) 'Uniquity: A general metric for biotic uniqueness of sites', *Biological Conservation*, 225, 98-105.

629 Epp Schmidt, D. J., Pouyat, R., Szlavecz, K., Setälä, H., Kotze, D. J., Yesilonis, I., Cilliers, S.,
630   Hornung, E., Dombos, M. and Yarwood, S. A. (2017) 'Urbanization erodes
631   ectomycorrhizal fungal diversity and may cause microbial communities to converge',
632   *Nature ecology & evolution*, 1, 0123.

633
634 Fernández-Toirán, L. M., Ágreda, T. and Olano, J. M. (2006) 'Stand age and sampling year
635   effect on the fungal fruit body community in Pinus pinaster forests in central Spain',
636   *Canadian Journal of Botany*, 84(8), 1249-1258.

637
638 Fløjgaard, C., Valdez, J. W., Dalby, L., Moeslund, J. E., Clausen, K. K., Ejrnæs, R., Pärtel, M.
639   and Brunbjerg, A. K. (2020) 'Dark diversity reveals importance of biotic resources and
640   competition for plant diversity across broad environmental gradients', *Ecology and*
641   *Evolution*, 00, 1– 11.

642
643 Fraser, L. H., Pither, J., Jentsch, A., Sternberg, M., Zobel, M., Askarizadeh, D., Bartha, S.,
644   Beierkuhnlein, C., Bennett, J. A. and Bittel, A. (2015) 'Worldwide evidence of a
645   unimodal relationship between productivity and plant species richness', *Science*,
646   349(6245), 302-305.

647
648 Frøslev, T. G., Kjøller, R., Bruun, H. H., Ejrnæs, R., Hansen, A. J., Læssøe, T. and Heilmann-
649   Clausen, J. (2019) 'Man against machine: Do fungal fruitbodies and eDNA give similar
650   biodiversity assessments across broad environmental gradients?', *Biological*
651   *Conservation*, 233, 201-212.

652
653 Geml, J., Gravendeel, B., van der Gaag, K. J., Neilen, M., Lammers, Y., Raes, N., Semenova, T.
654   A., de Knijff, P. and Noordeloos, M. E. (2014) 'The Contribution of DNA Metabarcoding
655   to Fungal Conservation: Diversity Assessment, Habitat Partitioning and Mapping Red-
656   Listed Fungi in Protected Coastal Salix repens Communities in the Netherlands', *PLOS*
657   *ONE*, 9(6), e99852.

658
659 Gijbels, P., Adriaens, D. and Honnay, O. (2012) 'An orchid colonization credit in restored
660   calcareous grasslands', *Écoscience*, 19(1), 21-28.

661
662 Gómez-Hernández, M. and Williams-Linera, G. (2011) 'Diversity of macromycetes determined
663   by tree species, vegetation structure, and microenvironment in tropical cloud forests in
664   Veracruz, Mexico', *Botany*, 89(3), 203-216.

665
Gómez, F. J. R., Navarro-Cerrillo, R. M., Pérez-de-Luque, A., Oβwald, W., Vannini, A. and
Morales-Rodríguez, C. (2019) 'Assessment of functional and structural changes of soil
fungal and oomycete communities in holm oak declined dehesas through metabarcoding
analysis', *Scientific reports*, 9(1), 5315.

Graham, J. H., Leonard, R. T. and Menge, J. A. (1982) 'Interaction of light intensity and soil
temperature with phosphorus inhibition of vesicular-arbuscular mycorrhiza formation',
New Phytologist, 91(4), 683-690.

Gupta, M. M., Gupta, A. and Kumar, P. (2018) 'Urbanization and biodiversity of arbuscular
mycorrhizal fungi: The case study of Delhi, India', *Revista de Biología Tropical*, 66(4),
1547-1558.

Heilmann-Clausen, J., Frøslev, T. G., Læssøe, T. and Petersen, J. H. (2019) *Danmarks*
svampeatlas 2009-2013, Svampetryk.

Höfle, B., Hollaus, M. and Hagenauer, J. (2012) 'Urban vegetation detection using
radiometrically calibrated small-footprint full-waveform airborne LiDAR data', *ISPRS*
Journal of photogrammetry and remote sensing, 67, 134-147.

Höfle, B. and Pfeifer, N. (2007) 'Correction of laser scanning intensity data: Data and model-
driven approaches', *ISPRS Journal of photogrammetry and remote sensing*, 62(6), 415-
433.

Hubbell, S. P. (2011) *The unified neutral theory of biodiversity and biogeography (MPB-32)*,
Princeton, New Jersey, USA: Princeton University Press.

Hui, N., Liu, X., Kotze, D. J., Jumpponen, A., Francini, G. and Setälä, H. (2017)
'Ectomycorrhizal Fungal Communities in Urban Parks Are Similar to Those in Natural
Forests but Shaped by Vegetation and Park Age', *Applied and Environmental*
Microbiology, 83(23), e01797-17.

Hülber, K., Moser, D., Sauberer, N., Maas, B., Staudinger, M., Grass, V., Wrbka, T. and Willner,
W. (2017) 'Plant species richness decreased in semi-natural grasslands in the Biosphere
Reserve Wienerwald, Austria, over the past two decades, despite agri-environmental
measures', *Agriculture, ecosystems & environment*, 243, 10-18.

Humphrey, J. W., Newton, A. C., Peace, A. J. and Holden, E. (2000) 'The importance of conifer
plantations in northern Britain as a habitat for native fungi', *Biological Conservation*,
96(2), 241-252.

Jacobson, K. M. (1997) 'Moisture and substrate stability determine VA-mycorrhizal fungal
community distribution and structure in an arid grassland', *Journal of Arid Environments*,
35(1), 59-75.

Junttila, S., Sugano, J., Vastaranta, M., Linnakoski, R., Kaartinen, H., Kukko, A., Holopainen,
M., Hyyppä, H. and Hyyppä, J. (2018) 'Can leaf water content be estimated using
multispectral terrestrial laser scanning? A case study with Norway spruce seedlings',
Frontiers in plant science, 9, 299.

715
716 Kalliokoski, T., Pennanen, T., Nygren, P., Sievänen, R. and Helmisaari, H.-S. (2010)
717   'Belowground interspecific competition in mixed boreal forests: fine root and
718   ectomycorrhiza characteristics along stand developmental stage and soil fertility
719   gradients', *Plant and Soil*, 330(1-2), 73-89.

720
721 Kennedy, P. G. and Peay, K. G. (2007) 'Different soil moisture conditions change the outcome of
722   the ectomycorrhizal symbiosis between Rhizopogon species and Pinus muricata', *Plant
723   and Soil*, 291(1), 155.

724
725 Kivlin, S. N., Hawkes, C. V. and Treseder, K. K. (2011) 'Global diversity and distribution of
726   arbuscular mycorrhizal fungi', *Soil Biology and Biochemistry*, 43(11), 2294-2303.

727
728 Kuiters, A. T. (2013) 'Diversity–stability relationships in plant communities of contrasting
729   habitats', *Journal of Vegetation Science*, 24(3), 453-462.

730
731 Kutner, M. H., Nachtsheim, C. J., Neter, J. and Li, W. (2005) *Applied linear statistical models*,
732   McGraw-Hill Irwin Boston.

733
734 Lefsky, M. A., Cohen, W. B., Parker, G. G. and Harding, D. J. (2002) 'Lidar Remote Sensing for
735   Ecosystem Studies: Lidar, an emerging remote sensing technology that directly measures
736   the three-dimensional distribution of plant canopies, can accurately estimate vegetation
737   structural attributes and should be of particular interest to forest, landscape, and global
738   ecologists', *BioScience*, 52(1), 19-30.

739
740 Leigh, J., Fitter, A. H. and Hodge, A. (2011) 'Growth and symbiotic effectiveness of an
741   arbuscular mycorrhizal fungus in organic matter in competition with soil bacteria', *FEMS
742   Microbiology Ecology*, 76(3), 428-438.

743
744 Lewis, R. J., Bello, F., Bennett, J. A., Fibich, P., Finerty, G. E., Götzenberger, L., Hiiesalu, I.,
745   Kasari, L., Lepš, J., Májeková, M., Mudrák, O., Riibak, K., Ronk, A., Rychtecká, T.,
746   Vitová, A. and Pärtel, M. (2017) 'Applying the dark diversity concept to nature
747   conservation', *Conservation Biology*, 31(1), 40-47.

748
Lewis, R. J., Szava☐Kovats, R., Pärtel, M. and Evolution (2016) 'Estimating dark diversity and
species pools: an empirical assessment of two methods', *Methods in Ecology*, 7(1), 104-
113.

Liu, Y., Johnson, N. C., Mao, L., Shi, G., Jiang, S., Ma, X., Du, G., An, L. and Feng, H. (2015)
'Phylogenetic structure of arbuscular mycorrhizal community shifts in response to
increasing soil fertility', *Soil Biology and Biochemistry*, 89, 196-205.

Lopatin, J., Dolos, K., Hernández, H., Galleguillos, M. and Fassnacht, F. (2016) 'Comparing
generalized linear models and random forest to model vascular plant species richness
using LiDAR data in a natural forest in central Chile', *Remote Sensing of Environment*,
173, 200-210.

Luo, S., De Deyn, G. B., Jiang, B. and Yu, S. (2017) 'Soil biota suppress positive plant diversity
effects on productivity at high but not low soil fertility', *Journal of Ecology*, 105(6),
1766-1774.

Mao, L., Dennett, J., Bater, C. W., Tompalski, P., Coops, N. C., Farr, D., Kohler, M., White, B.,
Stadt, J. J. and Nielsen, S. E. (2018) 'Using airborne laser scanning to predict plant
species richness and assess conservation threats in the oil sands region of Alberta's boreal
forest', *Forest Ecology and Management*, 409, 29-37.

Mateo, R. G., Mokany, K. and Guisan, A. (2017) 'Biodiversity Models: What If Unsaturation Is
the Rule?', *Trends in Ecology & Evolution*, 32(8), 556-566.

McCune, B. (1994) 'Improving community analysis with the Beals smoothing function',
Écoscience, 1(1), 82-86.

McCune, B. and Keon, D. (2002) 'Equations for potential annual direct incident radiation and
heat load', *Journal of Vegetation Science*, 13(4), 603-606.

Milberg, P., Bergstedt, J., Fridman, J., Odell, G. and Westerberg, L. (2008) 'Observer bias and
random variation in vegetation monitoring data', *Journal of Vegetation Science*, 19(5),
633-644.

Moeslund, J., Arge, L. and Bøcher, P. (2013) 'Topographically controlled soil moisture is the
primary driver of local vegetation patterns across a lowland region', *Ecosphere*, 4(7), 1-
26.

Moeslund, J. E., Brunbjerg, A. K., Clausen, K. K., Dalby, L., Fløjgaard, C., Juel, A. and Lenoir,
J. (2017) 'Using dark diversity and plant characteristics to guide conservation and
restoration', *Journal of Applied Ecology*, 54(6), 1730-1741.

Moeslund, J. E., Zlinszky, A., Ejrnæs, R., Brunbjerg, A. K., Bøcher, P. K., Svenning, J.-C. and
Normand, S. (2019) 'Light detection and ranging explains diversity of plants, fungi,
lichens, and bryophytes across multiple habitats and large geographic extent', *Ecological*
Applications, 29(5), e01907.

796
797 Mokany, A. and Shine, R. (2003) 'Competition between tadpoles and mosquito larvae',
798   *Oecologia*, 135(4), 615-620.

799
800 Mueller, G. M. (2011) *Biodiversity of fungi: inventory and monitoring methods*, Elsevier.

801
802 Münzbergová, Z. and Herben, T. (2004) 'Identification of suitable unoccupied habitats in
803   metapopulation studies using co‑occurrence of species', *Oikos*, 105(2), 408-414.

804
805 Nadeau, M. B. and Sullivan, T. P. (2015) 'Relationships between plant biodiversity and soil
806   fertility in a mature tropical forest, Costa Rica', *International Journal of Forestry*
807   *Research*, 2015, 1-13.

808
809 Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., Schilling, J. S. and
810   Kennedy, P. G. (2016) 'FUNGuild: an open annotation tool for parsing fungal community
811   datasets by ecological guild', *Fungal Ecology*, 20, 241-248.

812
813 Nordén, B. and Paltto, H. (2001) 'Wood-decay fungi in hazel wood: species richness correlated
814   to stand age and dead wood features', *Biological Conservation*, 101(1), 1-8.

815
816 Nordén, B., Ryberg, M., Götmark, F. and Olausson, B. (2004) 'Relative importance of coarse and
817   fine woody debris for the diversity of wood-inhabiting fungi in temperate broadleaf
818   forests', *Biological Conservation*, 117(1), 1-10.

819
820 Noreika, N., Pärtel, M. and Öckinger, E. (2020) 'Community completeness as a measure of
821   restoration success: multiple-study comparisons across ecosystems and ecological
822   groups', *Biodiversity and Conservation*, 29(13), 3807-3827.

823
824 Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R., Simpson, G.
825   L., Solymos, P., Stevens, M. and Wagner, H. (2017) 'vegan: Community Ecology
826   Package. R package version 2.4-3'.

827
Pärtel, M. (2014) 'Community ecology of absent species: hidden and dark diversity', *Journal of*
Vegetation Science, 25(5), 1154-1159.

Pärtel, M., Öpik, M., Moora, M., Tedersoo, L., Szava-Kovats, R., Rosendahl, S., Rillig, M. C.,
Lekberg, Y., Kreft, H., Helgason, T., Eriksson, O., Davison, J., Bello, F., Caruso, T. and
Zobel, M. (2017a) 'Historical biome distribution and recent human disturbance shape the
diversity of arbuscular mycorrhizal fungi', *New Phytologist*, 216(1), 227-238.

835
836 Pärtel, M., Szava-Kovats, R. and Zobel, M. (2011) 'Dark diversity: shedding light on absent
837   species', *Trends in Ecology & Evolution*, 26(3), 124-128.

838
839 Pärtel, M., Szava-Kovats, R. and Zobel, M. (2013) 'Community completeness: linking local and
840   dark diversity within the species pool concept', *Folia Geobotanica*, 48(3), 307-317.

841
842 Pärtel, M., Zobel, M., Öpik, M. and Tedersoo, L. (2017b) 'Global patterns in local and dark
843   diversity, species pool size and community completeness in ectomycorrhizal fungi' in
844   Tedersoo, L., ed., *Biogeography of Mycorrhizal Symbiosis*, Cham, Switzerland: Springer
845   International Publishing, 395-406.

846
847 Pärtel, M., Zobel, M., Zobel, K. and van der Maarel, E. (1996) 'The species pool and its relation
848   to species richness: Evidence from estonian plant communities', *Oikos*, 75(1), 111-117.

849
850 Pellkofer, S., van der Heijden, M. G. A., Schmid, B. and Wagg, C. (2016) 'Soil Communities
851   Promote Temporal Stability and Species Asynchrony in Experimental Grassland
852   Communities', *PLOS ONE*, 11(2), e0148015.

853
Penone, C., Allan, E., Soliveres, S., Felipe-Lucia, M. R., Gossner, M. M., Seibold, S., Simons,
N. K., Schall, P., van der Plas, F., Manning, P., Manzanedo, R. D., Boch, S., Prati, D.,
Ammer, C., Bauhus, J., Buscot, F., Ehbrecht, M., Goldmann, K., Jung, K., Müller, J.,
Müller, J. C., Pena, R., Polle, A., Renner, S. C., Ruess, L., Schönig, I., Schrumpf, M.,
Solly, E. F., Tschapka, M., Weisser, W. W., Wubet, T. and Fischer, M. (2019)
'Specialisation and diversity of multiple trophic groups are promoted by different forest
features', *Ecology letters*, 22(1), 170-180.

861
862 Peura, M., Gonzalez, R. S., Müller, J., Heurich, M., Vierling, L. A., Mönkkönen, M. and Bässler,
863   C. (2016) 'Mapping a 'cryptic kingdom': Performance of lidar derived environmental
864   variables in modelling the occurrence of forest fungi', *Remote Sensing of Environment*,
865   186, 428-438.

866
867 Pfeifer, N., Mandlburger, G., Otepka, J. and Karel, W. (2014) 'OPALS–A framework for
868   Airborne Laser Scanning data analysis', *Computers, Environment and Urban Systems*, 45,
869   125-136.

870
871 R Core Team (2019) 'R: A language and environment for statistical computing ', 3.6.0

872
873 Ricotta, C. (2005) 'Through the Jungle of Biological Diversity', *Acta Biotheoretica*, 53(1), 29-38.

874
875 Ricotta, C. (2007) 'A semantic taxonomy for diversity measures', *Acta Biotheoretica*, 55(1), 23-
876   33.

877
878 Riibak, K., Ronk, A., Kattge, J. and Pärtel, M. (2017) 'Dispersal limitation determines
879   large□scale dark diversity in Central and Northern Europe', *Journal of Biogeography*,
880   44(8), 1770-1780.

881
882 Ronk, A., Szava□Kovats, R. and Pärtel, M. (2015) 'Applying the dark diversity concept to plants
883   at the European scale', *Ecography*, 38(10), 1015-1025.

884
885 Rousk, J., Bååth, E., Brookes, P. C., Lauber, C. L., Lozupone, C., Caporaso, J. G., Knight, R. and
886   Fierer, N. (2010) 'Soil bacterial and fungal communities across a pH gradient in an arable
887   soil', *The ISME journal*, 4(10), 1340.

888
889 Rousk, J., Brookes, P. C. and Bååth, E. (2009) 'Contrasting soil pH effects on fungal and
890   bacterial growth suggest functional redundancy in carbon mineralization', *Applied and
891   Environmental Microbiology*, 75(6), 1589-1596.

892
893 Salusso, M. M. and Moraña, L. B. (1995) 'Estructura de la comunidad de hongos
894   ectomicorrízicos en bosques de Pinus spp. de Altos La Sierra, Argentina', *Revista Chilena
895   de Historia Natural*, 68, 509-513.

896
897 Sarkar, S. and Margules, C. (2002) 'Operationalizing biodiversity for conservation planning',
898   *Journal of Biosciences*, 27(4), 299-308.

899
900 Scott, C. E., Alofs, K. M. and Edwards, B. A. (2011) 'Putting dark diversity in the spotlight',
901   *Trends in Ecology & Evolution*, 26(6), 263-264.

902
903 Silvertown, J., Araya, Y. and Gowing, D. (2015) 'Hydrological niches in terrestrial plant
904   communities: a review', *Journal of Ecology*, 103(1), 93-108.

905
906 Solow, A. R. and Polasky, S. (1994) 'Measuring biological diversity', *Environmental and
907   Ecological Statistics*, 1(2), 95-103.

908
909 Staddon, P. L., Thompson, K., Jakobsen, I., Grime, J. P., Askew, A. P. and Fitter, A. H. (2003)
910   'Mycorrhizal fungal abundance is affected by long-term climatic manipulations in the
911   field', *Global Change Biology*, 9(2), 186-194.

Sterkenburg, E., Bahr, A., Brandström Durling, M., Clemmensen, K. E. and Lindahl, B. D. (2015) 'Changes in fungal communities along a boreal forest soil fertility gradient', *New Phytologist*, 207(4), 1145-1158.

Su, J. G. and Bork, E. W. (2007) 'Characterization of diverse plant communities in Aspen Parkland rangeland using LiDAR data', *Applied Vegetation Science*, 10(3), 407-416.

Tedersoo, L., Bahram, M., Põlme, S., Kõljalg, U., Yorou, N. S., Wijesundera, R., Ruiz, L. V., Vasco-Palacios, A. M., Thu, P. Q., Suija, A., Smith, M. E., Sharp, C., Saluveer, E., Saitta, A., Rosas, M., Riit, T., Ratkowsky, D., Pritsch, K., Põldmaa, K., Piepenbring, M., Phosri, C., Peterson, M., Parts, K., Pärtel, K., Otsing, E., Nouhra, E., Njouonkou, A. L., Nilsson, R. H., Morgado, L. N., Mayor, J., May, T. W., Majuakim, L., Lodge, D. J., Lee, S. S., Larsson, K.-H., Kohout, P., Hosaka, K., Hiiesalu, I., Henkel, T. W., Harend, H., Guo, L.-d., Greslebin, A., Grelet, G., Geml, J., Gates, G., Dunstan, W., Dunk, C., Drenkhan, R., Dearnaley, J., De Kesel, A., Dang, T., Chen, X., Buegger, F., Brearley, F. Q., Bonito, G., Anslan, S., Abell, S. and Abarenkov, K. (2014) 'Global diversity and geography of soil fungi', *Science*, 346(6213), 1256688.

Thers, H., Brunbjerg, A. K., Læssøe, T., Ejrnæs, R., Bøcher, P. K. and Svenning, J.-C. (2017) 'Lidar-derived variables as a proxy for fungal species richness and composition in temperate Northern Europe', *Remote Sensing of Environment*, 200, 102-113.

Tonn, N. and Ibáñez, I. (2017) 'Plant-mycorrhizal fungi associations along an urbanization gradient: implications for tree seedling survival', *Urban Ecosystems*, 20(4), 823-837.

Townsend, C. R., Scarsbrook, M. R. and Dolédec, S. (1997) 'The intermediate disturbance hypothesis, refugia, and biodiversity in streams', *Limnology and oceanography*, 42(5), 938-949.

Turner, G. D., Lewis, J. D., Mates-Muchin, J. T., Schuster, W. F. and Watt, L. (2009) 'Light availability and soil source influence ectomycorrhizal fungal communities on oak seedlings grown in oak- and hemlock-associated soilsContribution No. 225 of the Louis Calder Center and Biological Station, Fordham University, Armonk, New York', *Canadian Journal of Forest Research*, 39(7), 1247-1258.

Twieg, B. D., Durall, D. M. and Simard, S. W. (2007) 'Ectomycorrhizal fungal succession in mixed temperate forests', *New Phytologist*, 176(2), 437-447.

Unterseher, M. and Tal, O. (2006) 'Influence of small scale conditions on the diversity of wood decay fungi in a temperate, mixed deciduous forest canopy', *Mycological Research*, 110(2), 169-178.

Valdez, J. W., Hartig, F., Fennel, S. and Poschlod, P. (2019) 'The recruitment niche predicts
plant community assembly across a hydrological gradient along plowed and undisturbed
transects in a former agricultural wetland', *Frontiers in plant science*, 10, 88.

Vehmas, M., Eerikäinen, K., Peuhkurinen, J., Packalén, P. and Maltamo, M. (2009)
'Identification of boreal forest stands with high herbaceous plant diversity using airborne
laser scanning', *Forest Ecology and Management*, 257(1), 46-53.

Wang, J., Chen, C., Ye, Z., Li, J., Feng, Y. and Lu, Q. (2018) 'Relationships Between Fungal and
Plant Communities Differ Between Desert and Grassland in a Typical Dryland Region of
Northwest China', *Frontiers in Microbiology*, 9(2327).

Xiong, S. J., Johansson, M. E., Hughes, F. M. R., Hayes, A., Richards, K. S. and Nilsson, C.
(2003) 'Interactive effects of soil moisture, vegetation canopy, plant litter and seed
addition on plant diversity in a wetland community', *Journal of Ecology*, 91(6), 976-986.

Yahr, R., Schoch, C. L. and Dentinger, B. T. (2016) 'Scaling up discovery of hidden diversity in
fungi: impacts of barcoding approaches', *Philosophical Transactions of the Royal Society
B: Biological Sciences*, 371(1702), 20150336.

Yang, G., Wagg, C., Veresoglou, S. D., Hempel, S. and Rillig, M. C. (2018) 'How Soil Biota
Drive Ecosystem Stability', *Trends in Plant Science*, 23(12), 1057-1067.

Yang, T., Adams, J. M., Shi, Y., He, J.-s., Jing, X., Chen, L., Tedersoo, L. and Chu, H. (2017)
'Soil fungal diversity in natural grasslands of the Tibetan Plateau: associations with plant
diversity and productivity', *New Phytologist*, 215(2), 756-765.

Zak, D. R., Holmes, W. E., White, D. C., Peacock, A. D. and Tilman, D. (2003) 'Plant diversity,
soil microbial communities, and ecosystem function: Are there any links?', *Ecology*,
84(8), 2042-2050.

Zlinszky, A., Schroiff, A., Kania, A., Deák, B., Mücke, W., Vári, Á., Székely, B. and Pfeifer, N.
(2014) 'Categorizing grassland vegetation with full-waveform airborne laser scanning: A
feasibility study for detecting Natura 2000 habitat types', *Remote Sensing*, 6(9), 8056-
8087.

Zobel, M. (2016) 'The species pool concept as a framework for studying patterns of plant
diversity', *Journal of Vegetation Science*, 27(1), 8-18.

Zuo, X., Wang, S., Lv, P., Zhou, X., Zhao, X., Zhang, T. and Zhang, J. (2016) 'Plant functional
995   diversity enhances associations of soil fungal diversity with vegetation and soil in the
996   restoration of semiarid sandy grassland', *Ecology and Evolution*, 6(1), 318-328.

998