Title: Development of a free radical scavenging probiotic to mitigate coral bleaching

Running title: Making a probiotic to mitigate coral bleaching

Ashley M. Dungan[a#], Dieter Bulach[b], Heyu Lin[c], Madeleine J. H. van Oppen[a,d], Linda L. Blackall[a]

[a]School of Biosciences, The University of Melbourne, Melbourne, VIC, Australia

[b]Melbourne Bioinformatics, The University of Melbourne, Melbourne, VIC, Australia

[c] School of Earth Sciences, The University of Melbourne, Melbourne, VIC, Australia

[d]Australian Institute of Marine Science, Townsville, QLD, Australia

[#]Address correspondence to Ashley M. Dungan, adungan31@gmail.com

Abstract word count: 216

Text word count:

Keywords: symbiosis, *Exaiptasia diaphana*, *Exaiptasia pallida*, probiotic, antioxidant, ROS,

Symbiodiniaceae, bacteria

ABSTRACT

Corals are colonized by symbiotic microorganisms that exert a profound influence on the

animal's health. One noted symbiont is a single-celled alga (from the family *Symbiodiniaceae*),

which provides the coral with most of its carbon. During thermal stress, the algae's

photosystems are impaired, resulting in a toxic accumulation of reactive oxygen species (ROS)

that cause cellular damage to both the host and symbiont. As a protective mechanism the coral

host and algal symbiont disassociate; this process is known as bleaching. Our goal was to

construct a probiotic comprised of host-associated bacteria able to neutralize free radicals such

as ROS. Using the coral model, the anemone *Exaiptasia diaphana*, and pure bacterial cultures

isolated from the model animal, we identified six strains with high free radical scavenging

ability belonging to the families *Alteromonadaceae*, *Rhodobacteraceae*, *Flavobacteriaceae*, and

Micrococcaceae. In parallel, we established a "negative" probiotic consisting of genetically

related strains with poor free radical scavenging capacities. From their whole genome

sequences, we explore genes of interest that may contribute to the potential beneficial roles of

these putative probiotic members, which may help facilitate the therapeutic application of a

bacterial probiotic. Probiotics is one of several interventions currently being developed with the

aim of augmenting climate resilience in corals and increasing the likelihood of coral reef

persistence into the future.

IMPORTANCE

Coral reefs are experiencing severe declines due to environmental impacts linked to climate

change. Consequently, interventions are required to ensure their persistence until climate

change is addressed. Interventions targeting toxic oxygen production for coral's algal partner

have shown promise. Here we develop a potential probiotic to neutralize toxic oxygen with the

aim of improving coral resistance to environmental stress. The results presented here will help

direct future work that investigates the ability of probiotics to mitigate coral bleaching during

periods of thermal stress.

INTRODUCTION

Coral reefs are among the most biologically and economically productive ecosystems on Earth [1, 2]. Though they make up less than 0.1% of the ocean floor [3], coral reefs support fisheries, tourism, pharmaceuticals and coastal development with a global value of $8.9 trillion "international $"/year [4]. Corals and other reef organisms have been dying, largely due to anthropogenic influences such as climate change[5, 6], which has led to an increased frequency, intensity and duration of summer heat waves that cause coral bleaching [7, 8].

The coral holobiont [the sum of the coral animal and its symbiotic partners, including intracellular algae, endolithic algae, fungi, protozoans, bacteria, archaea and viruses; 9] is an ecosystem engineer. By secreting a calcium carbonate skeleton, the reef structure rises from the ocean floor, forming the literal foundation of the coral reef ecosystem. The success of corals to survive and build up reefs over thousands of years [10] is tightly linked to their obligate yet fragile symbioses with endosymbiotic dinoflagellates of the family *Symbiodiniaceae* [11].

Intracellular *Symbiodiniaceae* translocate photosynthetically fixed carbon to the coral host [12] in exchange for inorganic nitrogen, phosphorus and carbon and location in a high light environment with protection from herbivory [13, 14]. During periods of intense thermal stress, the relationship between the coral host and their *Symbiodiniaceae* breaks down, resulting in a separation of the partners and a state of dysbiosis. This phenomenon, 'coral bleaching', is devastating to the host and detrimental to the reef system. Debilitating effects of bleaching on the coral include reduced skeletal growth and reproductive activity, a lowered capacity to shed sediments, and an inability to resist invasion of competing species and diseases. Severe and prolonged bleaching can cause partial to total colony death, resulting in diminished reef growth, the transformation of reef-building communities to alternate, non-reef building community types, bioerosion and ultimately the disappearance of reef structures [11].

Although there are several hypotheses detailing the mechanisms driving bleaching [see 15, 16-18], a common theme is the overproduction and toxic accumulation of reactive oxygen species

(ROS) from the algal symbiont. Excess ROS are generated by a number pathways including heat

damage to both photosynthetic and mitochondrial membranes [19, 20], and are shown to play

a central role in injury to both symbiotic partners and to inter-partner communication of a

stress response [15]. Once generated, ROS causes damage to many cell components including

photosystem II (PSII) reaction centers in the *Symbiodiniaceae*, specifically at the D1 and D2

proteins , [see review in 21]. Exposure to elevated temperatures [22] can result in

photoinhibition of photosynthesis in *Symbiodiniaceae*. Once damaged, *Symbiodiniaceae* are no

longer able to maintain their role in the symbiotic relationship with corals and separate from

the host tissue via *in situ* degradation, exocytosis, host cell detachment, host cell apoptosis or

host cell necrosis [15].

Probiotics are preparations of viable microorganisms that are introduced to alter a microbial

community in a way that is beneficial to the host. Microbiome engineering through the addition

of probiotics has been postulated as a key strategy to manipulate host phenotypes and

ecosystem functioning for coral reefs [23-28]. The differences in the bacterial species

composition of healthy and thermally stressed corals [29-34] and the coral model *Exaiptasia*

diaphana [35-37] suggest a role for microbiome engineering in cnidarian health. A disruption to

the bacterial community of *Pocillopora damicornis* with antibiotic treatment diminished the

resilience of the holobiont during thermal stress, whereas intact microbial communities

conferred resilience to thermal stress and increased the rate of recovery after bleaching events

to the coral holobiont [38]. The relative stability of coral-associated bacterial communities have

also been linked to coral heat tolerance; the bacterial community of heat sensitive *Acropora*

hyacinthus corals shifted when transplanted to thermal stress conditions, whereas heat-

tolerant *A. hyacinthus* corals harbored a stable bacterial community [39].

In recent years, researchers have begun to explore the use of probiotics in corals and the model

organism for corals, *E. diaphana*. To inhibit the progression of white pox disease in *E. diaphana*,

caused by the pathogen *Serratia marcescens*, an *Alphaproteobacteria* cocktail containing

several *Marinobacter* spp. isolates was applied [40]. These strains were able to inhibit both

biofilm formation and swarming in *S. marcescens,* which halted disease progression in

E. diaphana. The probiotic was deemed effective as anemones exposed to both the cocktail and

pathogen survived after seven days, while anemones in the *S. marcescens* control experiment

died. A bacterial consortium native to the coral *Mussismilia harttii* was selected to degrade

water-soluble oil fractions[41]. This bioremediation strategy reduced the negative impacts of oil

on *M. harttii* health and accelerated the degradation of petroleum hydrocarbons [41]. Coral

microbiomes have also been manipulated to mitigate the effects of thermal stress. This

manipulation of the coral-associated microbiome was facilitated through addition of a

consortium of native or seawater derived bacteria to the surface of *P. damicornis* [42]. The

results from this study suggest the consortium was able to partially mitigate coral bleaching and

provides promising initial results in the field of coral probiotics.

The goal of this research was to identify bacterial strains suitable for use in a probiotic to

mitigate the effects of thermal stress in *E. diaphana*. Given the potential role of ROS in the

bleaching process, our focus was to select diverse *E. diaphana*-sourced bacterial strains with

antioxidant properties while avoiding potential pathogens. Antioxidant properties were

measured using the stable free radical 2,2-diphenyl-1-picrylhydrazyl (DPPH), which is reduced in

the presence of an antioxidant molecule, undergoing a color change from a violet to a colorless

solution.

RESULTS

Diversity of culturable bacteria associated with *E. diaphana*. A total of 842 isolates were obtained from four genotypes of GBR-sourced *E. diaphana*, with no significant differences in bacterial colony forming units (CFUs) between the four genotypes, regardless of growth medium, with (mean±SE) 5.9-10.3 x 10^3 cells per anemone on Reasoner's 2A agar (R2A) and 6.3-10.4 x10^3 cells per anemone on marine agar (MA) ($p > 0.05$). Partial 16S rRNA gene sequences (~1000 bp) were used to identify the closest matches from the NCBI database using BLASTn. In total there were 109 species in 64 genera, 27 families and six phyla (Fig. 1). The most abundant genera were *Alteromonas, Labrenzia*, and *Ruegeria* (Table 1). Gram-positive bacteria comprised 23 species, including *Microbacterium* (31 isolates) and *Micrococcus* (28 isolates). Eight genera were found to be associated with all four genotypes (Table 1); these eight genera made up 59.4% of all *E. diaphana*-associated bacterial isolates.

130   **Table 1:** Details of cultured bacteria associated with all four genotypes of GBR-sourced

131   *E. diaphana* (AIMS1-4).

Genus	Class	No. of species	AIMS1	AIMS2	AIMS3	AIMS4	Total Isolates
Alteromonas	*Gamma-proteobacteria*	6	10	52	24	30	116
Labrenzia	*Alpha-proteobacteria*	4	11	10	26	38	85
Marinobacter	*Gamma-proteobacteria*	4	7	25	10	13	55
Muricauda	*Flavobacteriia*	1	16	11	10	5	42
Roseovarius	*Alpha-proteobacteria*	3	9	12	5	6	32
Ruegeria	*Alpha-proteobacteria*	3	29	8	5	39	81
Shimia	*Alpha-proteobacteria*	2	2	3	9	40	54
Vibrio	*Gamma-proteobacteria*	3	3	7	31	15	56

FIG 1: Neighbor-Joining phylogenetic tree showing the diversity and relatedness of 842 *E.*
diaphana-associated bacterial isolates inferred using partial 16S rRNA sequences. These isolates
covered six phyla indicated by shading over the tree with *Proteobacteria* split into the classes
Gamma and *Alphaproteobacteria*. The positions of selected probiotic strains are highlighted by
arrows with blue arrows indicating the high FRS strains and orange arrows indicating low FRS
strains.

Bacterial probiotic selection. Of the original 842 isolates, 709 were screened for their ability to scavenge free radicals. Those isolates were divided into three categories, positive (144), weakly positive (121), and negative (444). There was no clear pattern of free radical scavenging (FRS) capacity at the family level (Fig. 2A) with strain specific responses by species. Ninety-eight strains representing eight families and 18 genera were quantitatively assessed for FRS (Fig. 2B). From these isolates, probiotic members were selected by choosing *E. diaphana*-associated bacteria species with separate strains displaying a high ("positive") or low ("negative") FRS ability (Fig. 2C-G; Table 2). Of the 12 selected probiotic members, seven were catalase positive and five were catalase negative (Table 2). In each probiotic set (i.e. high or low FRS strains), none of the selected isolates showed antagonistic activity against one other as evidenced by the absence of any zone of inhibition and growth from each combination of isolates on a plate.

FIG 2: Quantitative FRS ability of *E. diaphana*-associated bacteria isolates, separated by Family. Families with high relative

abundance among all cultured bacteria (*Rhodobacteraceae* – A, *Alteromondaceae* – B, *Pseudoalteromonadaceae* – C,

Flavobacteraceae – D, and *Micrococcaceae* – E) were separately analyzed to identify strains with a high FRS ability (blue arrows) and

a corresponding strain of the same species* with a low FRS ability (red arrows). *The two strains selected from *Micrococcaceae*

belonged to the Genus *Micrococcus* but were not the same species. In each panel, the light dashed vertical line on the left

represents the mean FRS of a 0.025% (w/v) ascorbic acid standard, the middle dark dashed vertical line is the mean FRS for 0.05%

(w/v) ascorbic acid standard, and the far right dashed line is the mean FRS of the 0.075% (w/v) ascorbic acid standard.

158 **Table 2:** Details on probiotic strains. Mean media FRS (% ± SE) was 27.2 ± 2.3% (n=12) and was catalase negative. All sequence data

159 can be found under BioProject# PRJNA574193. References to each probiotic candidate at the Genus level are identified in the last

160 three columns.

Strain	Bacteria species	FRS (% ± SE)	*Positive VS Negative	*Strain VS Growth Medium	Catalase	*Exaiptasia* Literature	Coral Literature	Probiotic Literature
MMSF01163 (n=3)	*Alteromonas oceani* (Gamma-proteobacteria)	61.7 ± 5.2	0.065	**0.004**	Negative	[43-45]	[25, 46-48]	[49-52]
MMSF00404 (n=4)		35.5 ± 13.7		0.571	Negative			
MMSF00958 (n=3)	*Alteromonas macleodii* (Gamma-proteobacteria)	62.0 ± 4.2	0.175	**0.017**	Negative			
MMSF00257 (n=3)		30.3 ± 0.9		0.431	Positive			
MMSF00132 (n=8)	*Labrenzia aggregata [53]* (Alpha-proteobacteria)	53.6 ± 8.7	**0.016**	0.101	Negative	[44]	[33, 48]	[54]
MMSF00249 (n=3)		14.0 ± 3.6		0.103	Positive			
MMSF01190 (n=3)	*Marinobacter salsuginis* (Gamma-proteobacteria)	62.0 ± 4.5	**0.041**	**<0.0001**	Negative	[43, 44, 55]	[47, 56, 57]	[40]
MMSF00964 (n=3)		43.7 ± 5.8		**0.020**	Positive			
MMSF00068 (n=6)	*Micrococcus luteus* (Actinobacteria)	56.3 ± 7.3	0.210	**0.002**	Positive	[44]	[58]	[49, 59, 60]
MMSF00107 (n=3)	*Micrococcus yunnanensis* (Actinobacteria)	38.0 ± 8.0		0.306	Positive			

| MMSF00046 (n=3) | _Winogradskyella poriferorum [61]_ (Bacteroidetes) | 73.3 ± 1.9 | 0.284 | **0.010** | Negative | ^ | [41, 62] | NA |
| MMSF00910 (n=3) | | 36.0 ± 3.5 | | 0.174 | Negative | | | |

^While there are no instances of _Winogradskyella_ specifically identified in current _E. diaphana_ literature, there are several

Flavobacteriaceae that are not resolved to the genus level from metabarcoding data _[36, 37, 43, 45]_.

*Indicates p values for pairwise comparisons from respective one-way analysis of variance (Tukey HSD) or Kruskal-Wallis rank sum

test (Dunn test) with **bold** values representing significant differences.

Comparative genomics. As part of the characterization of the 12 isolates (six positive and six

conspecific negative FRS isolates), a draft genome sequence was determined for each isolate. A

summary of the data and metrics for the draft genome sequences is presented in Table S1. The

diversity of the six pairs of isolates is indicated by the %G+C range (35% to 72%) and genome

size (2.4 Mb to 6.8 Mb). Each isolate pair was classified as the same species except the

Micrococcus stains. Isolates MMSF00068 (high FRS strain) and MMSF00107 (low FRS strain) are

classified as *Micrococcus luteus* by both 16S rRNA gene and genome-based methods, but

MMSF00107 is classified as *Micrococcus yunnanensis* by NCBI. These isolates have the smallest

genomes among the isolates (2.43 Mb 2.48 Mb, respectively) and the highest G+C content of

72.8% and 72.4%, respectively. The classification of isolate MMSF00107 is uncertain,

particularly given that the genome sequence for the type strain for *M. yunnanensis* is not yet

available.

Direct pairwise comparison of the genome sequences between the pairs of isolates revealed a

wide range of genome variation, ranging from ~190,000 single nucleotide polymorphisms (SNP)

differences between the *Alteromonas oceani* strains, to fewer than five SNPs between the

genomes of *Labrenzia aggregata* (Table S2). The genome sequences of *Winogradskyella*

poriferorum isolates MMSF00910 and MMSF00046 are nearly identical with fewer than ten

pairwise core SNP differences; but there were some accessory genome differences.

Genes of interest. The annotated genome sequences of each selected probiotic member were

searched for key genes of interest (Table S3-S4). Dimethylsulfoniopropionate (DMSP) cleavage

to dimethylsulfide (DMS) was identified by presence of one or more of the DMSP lyase genes;

dddP, *dddD*, *dddL*, *dddW*, and *dddQ*. Only *L. aggregata* strains contained DMSP lyase genes

(*dddP* and *dddL*) in their whole genome sequences (WGS). DMSP biosynthesis was identified by

the presence of *dsyB*, which is the only described gene for an enzyme in the DMSP biosynthesis

pathway. The presence of *dsyB* in both *L. aggregata* isolates suggests that they have the

capacity to produce DMSP. *cobP* was used as an indicator gene for the presence of the dynamic

vitamin B_{12} pathway, which contains 27 genes (Table S4). Again, only the *L. aggregata* isolates

contained *cobP*. Catalase positive strains were identified by the presence of *katG*; all positive

and negative FRS strains contained *katG* except the *Micrococcus* spp. strains, in which *katA* and

katE were detected.

16S rRNA gene copy number.

196   The 16S rRNA gene copy numbers of the 12 draft genomes were estimated using a read depth

197   approach (Table S1). The copy numbers were close between each other within pairs of isolates,

198   in which the pair of isolates MMSF00257 and MMSF00958 contained the most copies (5.15 and

199   4.79, respectively) and the isolates MMSF00046 and MMSF00910 contained the fewest copies

200   (1.03 and 0.77, respectively).

DISCUSSION

The 842 *E. diaphana* bacterial isolates reported here and based on 16S rRNA gene-based

taxonomic classification comprise 109 species from 64 genera and six phyla. Using

metabarcoding, studies of microbiomes associated with *E. diaphana* have revealed a similar

diversity at the phylum level for *E. diaphana* sourced from the GBR [35], Hawaii [strain H2; 43],

Pacific and Caribbean [55], Atlantic [strain CC7; 45] and Red Sea [36], as well as stony corals

(see a review by Blackall, Wilson [63]). Thus, our culture collection of *E. diaphana* bacterial

isolates suitably represents the diversity of the *E. diaphana*-associated microbiome. Though

previous studies have used only a narrow bacterial species diversity to develop a probiotic [40,

42], our culture collection is sufficiently diverse to better select potential probiotics.

Inoculated bacteria can be acquired by developing coral larvae [25] and probiotic mixes applied

to cnidarians have been shown to be effective in inhibiting disease progression [40] and

increasing resistance to the negative effects of oil [41] and heat exposure [42]. The *E. diaphana*

probiotics generated in this study were assembled from 12 isolated bacterial strains selected

based on their FRS ability since free radical production, specifically ROS, is relevant in coral

bleaching. The broader culture collection contains bacteria with a wide range of FRS capacity.

The consistent and frequent reporting of our probiotic bacteria genera in *E. diaphana* and coral

studies (see Table 2) suggests these bacteria may have key functions in cnidarian holobionts.

Among these potential functions are the production of antioxidants such as DMSP, the

breakdown of DMSP to other antioxidants (dimethyl sulfide (DMS), acrylate, dimethyl sulfoxide

(DMSO), and methane sulfinic acid (MSA) [64]. *L. aggregata* has been reported to produce

DMSP in the absence of any methylated sulfur compounds with *dsyB* identified as the first

DMSP biosynthesis gene in any organism [65]. *dsyB* was found in the whole genome sequences

of both high and low FRS *L. aggregata* strains (Table S3). Many *E. diaphana*-sourced bacterial

species, specifically relatives of our selected probiotic members, are implicated in the

degradation of DMSP to DMS (*Alteromonas* spp., [66]; *Labrenzia* spp., [67]). *dddP* codes for the

enzyme responsible for cleaving DMSP to DMS and acrylate and was used (from the Prokka

annotation) as an indicator of a DMSP degradation genotype. Only the *L. aggregata* isolates

were found to be able to degrade DMSP (Table S3).

Carotenoids are among the strongest antioxidants and are highly reactive against both reactive

oxygen species and free radicals [68-72]. Carotenoids are lipid-soluble pigments, and in bacteria

they give an orange-yellow hue to colonies. Two of the five selected probiotic genera produce

orange/yellow colonies (*Winogradskyella*, *Micrococcus*), and there is evidence of carotenoid

production by marine *Flavobacteriaceae* [72, 73] and *Micrococcus* strains [74]. A marine

Flavobacteriaceae (strain GF1) was found to produce the potent antioxidant carotenoid

zeaxanthin that protected *Symbiodiniaceae* from thermal and light stress [75].

Vitamin B_{12} is a cofactor involved in the production of the amino acid methionine, which is

needed to synthesize every protein and in diverse metabolic pathways including generation of

the antioxidants glutathione and DMSP [76]. Vitamin B_{12} is synthesized by many heterotrophic

bacteria [77]. Genomic evidence suggests that *Symbiodiniaceae* may have lost the capacity to

synthesize vitamin B_{12} due to changes in their metabolic enzymes during their evolution [78],

agreeing with other work that free-living *Symbiodiniaceae* depend on bacterial symbionts to

gain access to this important cofactor [79]. The genes involved in the biosynthesis of vitamin B_{12}

have been found in coral-associated bacteria, specifically *L. aggregata* cultured from the

Caribbean coral, *Orbicella faveolata* [80].From the annotated draft genome sequence, using the

cobP gene [77] as the genotypic indicator, both *L. aggregata* isolates are capable of Vitamin B_{12}

biosynthesis. The *Marinobacter salsuginis* isolates lack this gene but have other detectable

vitamin B_{12} synthesis genes (Table S4).

Bacteria have developed highly specific mechanisms to protect themselves against oxidative

stress with enzymes such as catalase/peroxidase and superoxide dismutase (SOD), small

proteins like thioredoxin and glutaredoxin, and molecules such as glutathione (GSH) in

combination with glutathione peroxidase and glutathione reductase [see review 81]. It has

been suggested that increasing the *in hospite* concentration of catalase in the coral holobiont

by the application of a probiotic with catalase-positive organisms, could possibly minimize the

impact of thermal stress by neutralizing hydrogen peroxide [27]. While this is a justifiable

hypothesis worth investigating further, to our knowledge, no studies have shown that catalase

producing bacteria can reduce the concentration of ROS *in hospite*. Here we tested all probiotic

candidates for catalase production using a standard hydrogen peroxide assay [82]. Catalase

participates in cellular antioxidant defense by decomposing hydrogen peroxide; this

decomposition is enzymatically driven by two catalases (yielding H_2O and O_2): hydroperoxidase

I (HPI),which is present during aerobic growth and transcriptionally controlled at different

levels, and hydroperoxidase II (HPII), which is induced during stationary phase [83]. Most

phenotypically determined catalase positive strains (MMSF00249, *L. aggregata*; MMSF00257,

A. macleodii; MMSF00964, *M. salsuginis*) had homologs for the catalase-peroxidase gene, *katG*;

the *Micrococcus* spp. isolates did not (Table S3). However, *katG* homologs were found in all

strains that were phenotypically catalase negative, indicating that these genes may not be

active during culture. Given the inconsistency between the catalase and DPPH results (in many

cases high FRS isolates were catalase negative while the low FRS strain was catalase positive)

the catalase results were not used as a primary factor in selecting members of the positive and

negative probiotic.

Members of the selected probiotic consortium (e.g., *Marinobacter* spp. and *Winogradskyella*

spp.) have some described roles in coral/*E. diaphana* health. For example, coral-bleaching

events are often followed by disease outbreaks [84-86] such as white pox [87]. When

inoculated in the presence of the white pox pathogen, *S. marcescens*, *Marinobacter* strains

were able to inhibit the progression of the infection in *E. diaphana* [40]. In a bioremediation

study on coral and oil, *Winogradskyella* sp. showed a significant decrease in abundance during

oil treatment, which was correlated with a decrease in coral holobiont health measured via

maximum quantum yield [41]. The presence of this microbiome member could play a role in

protecting the *Symbiodiniaceae* photochemical ability during periods of stress.

A critical characteristic in the selection of probiotic members is the maintenance and

proliferation of the inoculated bacteria in the host over time [49] and their potential

transmission to the next generation. There is evidence that corals release bacteria with their

offspring such as *Alteromonas* [46], *Flavobacteriaceae* [46], *Rhodobacteraceae* [88], and

Marinobacter [56]. While the majority of broadcast spawning corals do not transfer their

bacterial symbionts with their gametes (vertical transfer) [89], the brooding coral *Porites*

astreoides transmits bacteria vertically to planulae with two bacterial taxa (*Roseobacter* clade-

associated bacteria and *Marinobacter* spp.) consistently and stably associating with juvenile *P.*

astreoides [56]. In addition to the potential antioxidant properties of *Marinobacter* [90], others

like *Roseobacter* spp. might be beneficial in facilitating larval settlement. If adult corals stably

associate with inoculated probiotic candidates like *Marinobacter, Alteromonas,* and

Winogradskyella, they may be passed on to offspring and thus have a long-term positive impact

on these individuals.

Interactions within the microbiota associated with marine holobionts are undoubtedly complex.

Results presented in this manuscript show that pure cultured bacteria from *E. diaphana* can

scavenge free radicals, albeit at a strain-specific rate. This suggests that the selection and

inoculation of these high FRS strains could be beneficial to the host under high oxidative stress

conditions, such as those that contribute to coral bleaching. Conspecific pairs of six bacteria

provide an opportunity to determine the genetic basis for measured phenotypic differences

between the pairs. An essential element of this future work will be to investigate the stability of

the phenotypic differences observed and this stability may be reflected in the nature of the

genetic differences between the pairs of strains. Where isolate pairs are distantly related, it is

unlikely that the genetic basis for the phenotypic difference will be identified using comparative

genomics.

The probiotic members were chosen from a highly diverse pool of *E. diaphana*-sourced

bacterial isolates. While the selected probiotic members are phylogenetically diverse,

potentially promising probiotic bacteria in the culture collection were omitted based on our

selection criteria. For example, *Ruegeria* spp., which were excluded based on the absence of a

high FRS strain, have the ability to breakdown DMSP and can participate in denitrification [80].

Muricauda isolates had high FRS abilities, but they did not grow consistently in the selected

medium and therefore were excluded from the consortium. Like *Ruegeria*, *Muricauda* also has

genes for denitrification [80], can oxidize DMS to produce DMSO [67], and produces potent

carotenoids [91] that can mitigate thermal and light stresses in *Symbiodiniaceae* cultures [75].

Muricauda will be involved in future probiotic evaluations.

At present there is no biological treatment that can minimize coral bleaching in the field.

Management priorities for coral reefs must move beyond documenting their declines and

toward investigating potential approaches for mitigating coral bleaching, such as the

application of coral probiotics. We believe that the application of a coral probiotic specifically

tailored to address coral bleaching by neutralizing ROS could provide hope for the future of

coral reefs. We also understand that this form of intervention may not work alone, but could

benefit by pairing with other strategies such as enhancing coral resistance and resilience using

other assisted evolution approaches such as assisted gene flow, hybridization and experimental

evolution of the algal symbionts [24, 26, 92-98].Climate warming will continue even with the

most drastic reductions in greenhouse gas emissions, thus, additional interventions such as

coral probiotics present an alternative that could lead to relief from coral bleaching in real time.

MATERIALS AND METHODS

Isolation of bacterial isolates. Great Barrier Reef (GBR) origin *E. diaphana* were maintained in

the laboratory at 26°C [99] and used to isolate probiotic candidates. Sixteen individuals from

each of four *E. diaphana* genotypes (AIMS1-4) were collected using sterile disposable pipets

and gently transferred to filter-sterilized (0.2 μm) reverse osmosis (RO) water reconstituted Red

Sea Salt™ (Red Sea; RSS) at ~34 parts per thousand (ppt) salinity (fRSS) and placed in the dark

for 30 min. This was done to remove some of the influence of the external seawater on the

bacterial community. After 30 min, each anemone was transferred to a sterile glass

homogenizer with 1 mL of fRSS. Each homogenate was used to prepare serial dilutions from 10^{-1}

to 10^{-4}. From each dilution, 50 μL was spread plated onto three replicate plates each of MA

(Difco™ Marine Agar 2216) and R2A (CM0906, Oxoid) supplemented with 40 g L^{-1} RSS and

incubated at 26˚C. After one week, CFU counts were completed. Individual bacterial isolates

were sub-cultured to purification from plates with <100 CFUs onto the initial isolation medium.

All purified bacterial isolates were resuspended in 40% glycerol, aliquoted into 1.2 mL

cryotubes and stored at -80°C.

Identification of *Exaiptasia*-sourced isolates. Colony PCR with the universal bacterial primers

27f (5' – AGA GTT TGA TCM TGG CTC AG – 3') and 1492r (5' – TAC GGY TAC CTT GTT ACG ACT T

– 3') [100] was used to generate 16S rRNA gene amplicons from each isolate. Briefly, cells from

each pure culture were suspended in 20 μL Milli-Q water and denatured at 95°C for 10 min. The

suspension was then centrifuged at 2,000 x *g* at 4°C for two minutes and the supernatant was

used as the DNA template for PCR amplification. The PCR was performed with 20 μL Mango

Mix™ (Bioline), 0.25 μM of each primer and 2 μL of DNA template in a final volume of 40 μL.

The thermal cycling protocol was as follows: 95°C for 5 min; 35 cycles of 95°C for 1 min, 50 °C

for 1 min, and 72°C for 1 min; and a final extension of 10 min at 72°C. Amplicons were purified

and Sanger sequenced on an ABI sequencing instrument by Macrogen Inc. (Seoul, South Korea)

or by the Australian Genome Research Facility (AGRF) using the 1492r primer. Trimmed high

quality read data from each isolate was used for presumptive identification by querying the 16S

rRNA gene sequences via the Basic Local Alignment Search Tool (BLASTn). For some isolates the

near-complete 16S rRNA gene sequence was determined by sequencing with additional primers

(27f, 357f (5'-CCT ACG GGA GGC AGC AG-3', [101]), 926f (CCG TCA ATT CMT TTR AGT TT, [102]),

519r (5'-GWA TTA CCG CGG CKG CTG-3', [101]), 926r (5'–AAA CTR AAA MGA ATT GAC GG–3',

[102]), and 1492r). The six reads for each isolate were aligned using Geneious Prime 2019.1.2

(https://www.geneious.com) via the Geneious global alignment default settings with automatic

determination of read direction. From this alignment, a consensus sequence for the 16S rRNA

gene was constructed based on the frequency of a base and its quality (from chromatogram

data) in each alignment column. The consensus sequence length for each of the six isolate pairs

varied from 1352 to 1495 nucleotides. GenBank accession numbers for sequences are shown in

Table 1.

Qualitative free radical scavenging assay. DPPH is a stable free radical that is purple in its

oxidized state but becomes white-yellow when reduced by antioxidants, and has been used to

identify FRS marine bacteria [103, 104]. To qualitatively assess *E. diaphana*-associated bacteria

isolates for FRS ability, a sterile Whatman #1 filter paper was gently pressed against fresh (2-4

days old) colonies from a streak plate. Plates (with filter paper) were then incubated overnight

at 26°C. The following day, filter papers were removed with forceps, allowed to dry in a fume

hood for 30 min, and 500 µL of a 0.2 mM DPPH (Cat# D9132, Sigma-Aldrich) solution in

methanol was applied with a pipette over individual colonies. As a positive control, a few drops

of 0.1% (w/v) L-ascorbic acid (Cat# A7631, Sigma-Aldrich) were placed on a separate filter. The

response of each isolate to DPPH was recorded within 3 min of DPPH application; a positive

response was recorded when a white-yellow halo appeared around individual colonies within 1

min, a weak positive response was assigned to strains that had a halo form between 1 and 3

min after DPPH application, and a negative response was listed for strains that failed to form a

halo (Fig. 3). Approximately 700 isolates were screened using the qualitative DPPH assay.

A.

 B.

FIG 3: DPPH is a stable free radical that is purple in its oxidized state. When reduced by an

antioxidant, a white-yellow halo will appear around individual bacteria colonies (A), this was

qualitatively deemed a positive response. Isolates that did not have a halo around colonies

within 3 min of DPPH application were deemed negative (B).

Quantitative free radical scavenging assay. To quantitatively assess the FRS ability, select

isolates were grown in R2A broth (see Table S5 for composition) made by suspending 43.12 g in

1 L of MilliQ water, dissolving the medium completely, and sterilization by autoclaving at 121˚C

for 15 min. Fifty mL aliquots of autoclaved medium were distributed into sterile 250 mL

Erlenmeyer flasks and each flask was inoculated with an isolate colony grown on agar (R2A or

MA). Cultures were grown with shaking (150 rpm; Ratek orbital incubator) at 37˚C for 48 h. A

minimum of three replicate cultures were grown per isolate. After 48 h, the optical density of

each culture was measured at 600 nm (OD_{600}, CLARIOstar PLUS, BMG Labtech), and the cultures

(including negative medium controls) were centrifuged at 3000 x g at 4˚C for 30 min (Allegra X-

12R) to pellet the bacterial cells. The cell-free supernatants (CFSs) were collected, frozen at -

80˚C, freeze dried (Alpha 1-4 LDplus, Martin Christ), and stored under inert gas in a dark, dry

environment until analysis. Antioxidants were extracted from the CFSs by resuspending at 50

mg mL^{-1} in 100% methanol, sonicating (Branson 2510) for 5 min, then centrifuging at 3000 x g

for 5 min at 4˚C. Quantitative DPPH assays were run by creating a 1:1 solution of 0.2 mM DPPH

in methanol and CFS extract to a final volume of 1 mL, vortexing, and reaction in the dark for 30

min at room temperature. Samples were then vortexed briefly, and three 300 µL replicates of

each sample were transferred into a well of a 96 well plate. FRS was measured by determining

absorbance at 517 nm (Enspire 2300 plate reader, Perkin Elmer). Decolourization of DPPH was

determined by measuring the decrease in absorbance at 517 nm, and the FRS activity was

calculated according to the formula, % DPPH scavenging activity = (Control – Sample) / Control

×100, where, Control is the absorbance of the DPPH control (1:1 0.2 mM DPPH:methanol), and

Sample is the absorbance of CFS extract in DPPH. All samples were measured against a 100%

methanol blank. Positive controls consisting of 0.01 - 0.001% (w/v) L-ascorbic acid were run on

each 96-well plate. FRS activity ranged from 0-90%.

Catalase assay. The pelleted cells from above were resuspended in 2 mL fRSS and 500 µL

hydrogen peroxide giving a final concentration of 16 mM. If bubbles appeared, the organism

was considered catalase positive. If there were no bubbles, the organism was classified as

catalase negative.

Inhibition testing. Each paired set of high and low FRS strains were inoculated crosswise along

the middle of MA plates to test for antagonism. Plates were kept at 26 °C and monitored daily

for up to 7 days for antagonistic activity by documenting the presence or absence of both

inoculated isolates and if there was a zone of inhibition between them.

Phylogenetic analysis. All partial 16S rRNA gene sequences (842) were aligned with reference

sequences (72) of closely related organisms using Geneious Prime 2019.1.2

(https://www.geneious.com). This alignment was used to construct a neighbor-joining

phylogenetic tree using the Jukes-Cantor method. Maximum-likelihood dendrograms were

generated with bootstrap values of 1000.

Whole genome sequence analysis. Positive FRS strains along with conspecific negative FRS

strains were selected for genome sequencing; in total, six pairs of isolates were sequenced.

Genomic DNA was isolated from a single colony using a JANUS Chemagic Workstation and

Chemagic Viral DNA/RNA kit (PerkinElmer). Libraries were prepared with the Nextera XT DNA

sample preparation kit (Illumina). Readsets were produced using the Illumina sequencing

platform (Instrument: Illumina NextSeq 500, 150 base, paired-end) and the whole genome

shotgun (WGS) method. Read depth coverage was approximately 100 times assuming a

genome size of 4 M bases.

Illumina readsets for each isolate were assembled using Skesa [105] and the draft genome

sequence annotated using Prokka [106]. No evidence of mixed colonies or sequence

contamination was detected. A genome sequence based taxonomic classification for each

isolate was determined using Kraken2 [107] with the Genome Taxonomy Database [GTDB; 108]

as the curated genomic data source. Classification was primarily based on the genome

sequence of related isolates (within the relevant species where possible), which were obtained

from GenBank. In situations where genomes of taxonomically relevant individuals were

available, a species level classification was possible. Where available, closed genome sequences

from GenBank were used for comparative genomics analysis. Core genome comparisons were

performed, as implemented in Nullarbor (https://github.com/tseemann/nullarbor), for each of

the six pairs of isolates, with phylogenies inferred using core SNP differences. Genes of interest

for DMSP synthesis and degradation, vitamin B_{12} synthesis, and catalase were identified from

the annotated genome sequence (GFF format) produced by Prokka; specific genes were

identified by both name and Refseq accession number.

16S rRNA gene copy number estimation.

The 16S rRNA gene copy number of the 12 draft genomes was predicted by the 16Stimator

pipeline [109]. Briefly, all the 12 genomic assemblies were submitted to the RAST server [110],

and the positions of 16S rRNA and a set of single-copied housekeeping genes (Table S6) were

extracted from the RAST annotations. The clean readsets were mapped back to the

corresponding genomic assemblies by Bowtie 2 [111] to determine the read depth of each

position. Finally, the 16S copy number of each isolate was calculated by dividing the median

depth of 16S gene by the median depth of the single-copied housekeeping genes after the read

depths were calibrated by the model parameters provided by 16Stimator.

Statistical analysis. CFU counts were analyzed in R [v3.6.2, 112] by first checking the

assumptions of equal variance and homogeneity. An analysis of variance test was used to

detect differences in the mean number of bacterial colonies from each anemone genotype by

solid growth media (R2A or MA). A one-way analysis of variance [one-way ANOVA; 113] was

used to determine if there were significant differences between FRS abilities of selected

positive (high FRS), negative (low FRS), and media controls, and pairwise comparisons were

performed using Tukey's HSD [114, 115]. Each probiotic pair and media control was tested to

determine if data met the assumptions of normality and homoscedasticity. If either assumption

was violated, the non-parametic Kruskal-Wallis rank sum test [116] was used with a Dunn test

[117] for multiple comparisons (p-values adjusted with the Benjamini-Hochberg method [118])

with the R package "FSA" [119].

Data availability. WGS raw reads are freely available in the Sequence Read Archive under

BioProject PRJNA574193; the complete data set is listed in Table S1.

ACKNOWLEDGEMENTS

This research was supported by the Australian Research Council Discovery Project grant DP160101468 (to MJHvO and LLB). MJHvO acknowledges Australian Research Council Laureate Fellowship FL180100036. We are grateful to Leon Hartman, Giulia Holland, and Shona Elliot-Kerr for their contributions in the preliminary culturing and screening of anemone-associated bacteria. Dr. Gayle Philip contributed with bacterial whole genome sequence analysis and Leon Hartman assisted with figure designs and reviewed the manuscript. Xavier Smith assisted with bacterial inhibition tests. Whole genome sequencing was organized by Dr. Glen Carter at the Peter Doherty Institute, Melbourne, Australia.

AMD, MvO and LLB conceived and designed the study. AMD performed the sampling and sample processing. AMD, DB, and HL completed bioinformatic analyses. AMD wrote the first draft. All authors edited and approved the final manuscript.

1.   Cesar, H., L. Burke, and L. Pet-Soede, *The economics of worldwide coral reef degradation*. 2003.

2.   Alder, J., et al., *Marine and coastal ecosystems and human well-being: A synthesis report based
on the findings of the Millennium Ecosystem Assessment*, C. Brown, et al., Editors. 2006.

3.   Spalding, M. and A. M. Grenfell, *New estimates of global and regional coral reef areas*. Vol. 16.
1997. 225-230.

4.   de Groot, R., et al., *Global estimates of the value of ecosystems and their services in monetary
units.* Ecosystem Services, 2012. **1**(1): p. 50-61.

5.   Stuart-Smith, R.D., et al., *Ecosystem restructuring along the Great Barrier Reef following mass
coral bleaching.* Nature, 2018. **560**(7716): p. 92-96.

6.   Hughes, T.P., et al., *Global warming and recurrent mass bleaching of corals.* Nature, 2017.
543(7645): p. 373-377.

7.   Hoegh-Guldberg, O., et al., *The human imperative of stabilizing global climate change at 1.5° C.*
Science, 2019. **365**(6459): p. eaaw6974.

8.   Hughes, T.P., et al., *Spatial and temporal patterns of mass bleaching of corals in the
Anthropocene.* Science, 2018. **359**: p. 80-83.

9.   Rohwer, F., et al., *Diversity and distribution of coral-associated bacteria.* Mar Ecol Prog Ser,
2002. **243**: p. 10.

10.   Devlin-Durante, M.K., et al., *How old are you? Genet age estimates in a clonal animal.* Molecular
Ecology, 2016. **25**(22): p. 5628-5646.

11.   Glynn, P.W., *Coral reef bleaching: facts, hypotheses and implications.* Global Change Biology,
1996. **2**(6): p. 495-509.

12.   Muscatine, L. and J.W. Porter, *Reef corals: mutualistic symbioses adapted to nutrient-poor
environments.* Bioscience, 1977. **27**(7): p. 454-460.

13.   Venn, A.A., J.E. Loram, and A.E. Douglas, *Photosynthetic symbioses in animals.* J Exp Bot, 2008.
59(5): p. 1069-80.

14.   Yellowlees, D., T.A.V. Rees, and W. Leggat, *Metabolic interactions between algal symbionts and
invertebrate hosts.* Plant, Cell & Environment, 2008. **31**(5): p. 679-694.

15.   Weis, V.M., *Cellular mechanisms of Cnidarian bleaching: stress causes the collapse of symbiosis.*
Journal of Experimental Biology, 2008. **211**(19): p. 3059-3066.

16.   Cunning, R. and A.C. Baker, *Excess algal symbionts increase the susceptibility of reef corals to
bleaching.* Nature Climate Change, 2012. **3**(3): p. 259-262.

17.   Wiedenmann, J., et al., *Nutrient enrichment can increase the susceptibility of reef corals to
bleaching.* Nature Climate Change, 2012. **3**: p. 160.

18.   Wooldridge, S.A., *Breakdown of the coral-algae symbiosis: towards formalising a linkage
between warm-water bleaching thresholds and the growth rate of the intracellular
zooxanthellae.* Biogeosciences, 2013. **10**(3): p. 1647-1658.

19.   Dykens, J.A., et al., *Oxygen radical production in the sea anemone Anthopleura elegantissima
and its endosymbiotic algae.* Journal of Experimental Biology, 1992. **168**(1): p. 219-241.

20.   Nii, C.M. and L. Muscatine, *Oxidative Stress in the Symbiotic Sea Anemone Aiptasia pulchella
(Carlgren, 1943): Contribution of the Animal to Superoxide Ion Production at Elevated
Temperature.* Biol Bull, 1997. **192**(3): p. 444-456.

21.   Lesser, M.P., *Experimental biology of coral reef ecosystems.* Journal of Experimental Marine
Biology and Ecology, 2004. **300**(1): p. 217-252.

22.   Iglesias-Prieto, R., et al., *Photosynthetic response to elevated temperature in the symbiotic
dinoflagellate Symbiodinium microadriaticum in culture.* Proc Natl Acad Sci U S A, 1992. **89**(21):
p. 10302-5.

23.   Damjanovic, K., et al., *The contribution of microbial biotechnology to mitigating coral reef
degradation.* Microb Biotechnol, 2017. **10**(5): p. 1236-1243.

24. van Oppen, M.J.H., et al., *Shifting paradigms in restoration of the world's coral reefs.* Glob Chang
Biol, 2017. **23**(9): p. 3437-3448.
25. Damjanovic, K., et al., *Experimental Inoculation of Coral Recruits With Marine Bacteria Indicates*
Scope for Microbiome Manipulation in Acropora tenuis and Platygyra daedalea. Frontiers in
Microbiology, 2019. **10**(1702).
26. van Oppen, M.J., et al., *Building coral reef resilience through assisted evolution.* Proc Natl Acad
Sci U S A, 2015. **112**(8): p. 2307-13.
27. Peixoto, R.S., et al., *Beneficial Microorganisms for Corals (BMC): Proposed Mechanisms for Coral*
Health and Resilience. Front Microbiol, 2017. **8**: p. 341.
28. Epstein, H.E., et al., *Microbiome engineering: enhancing climate resilience in corals.* Frontiers in
Ecology and the Environment, 2019. **17**(2): p. 100-108.
29. Vega Thurber, R., et al., *Metagenomic analysis of stressed coral holobionts.* Environ Microbiol,
2009. **11**(8): p. 2148-63.
30. Epstein, H.E., G. Torda, and M.J.H. van Oppen, *Relative stability of the Pocillopora acuta*
microbiome throughout a thermal stress event. Coral Reefs, 2019. **38**(2): p. 373-386.
31. Sunagawa, S., C.M. Woodley, and M. Medina, *Threatened corals provide underexplored*
microbial habitats. PLoS One, 2010. **5**(3): p. e9554.
32. Mouchka, M.E., I. Hewson, and C.D. Harvell, *Coral-associated bacterial assemblages: current*
knowledge and the potential for climate-driven impacts. Integr Comp Biol, 2010. **50**(4): p. 662-
74.
33. Littman, R., B.L. Willis, and D.G. Bourne, *Metagenomic analysis of the coral holobiont during a*
natural bleaching event on the Great Barrier Reef. Environ Microbiol Rep, 2011. **3**(6): p. 651-60.
34. Pootakham, W., et al., *Heat-induced shift in coral microbiome reveals several members of the*
Rhodobacteraceae family as indicator species for thermal stress in Porites lutea.
MicrobiologyOpen, 2019. **8**(12): p. e935.
35. Hartman, L.M., M.J. van Oppen, and L.L. Blackall, *The Effect of Thermal Stress on the Bacterial*
Microbiome of Exaiptasia diaphana. Microorganisms, 2020. **8**(1): p. 20.
36. Ahmed, H.I., et al., *Long-term temperature stress in the coral model Aiptasia supports the "Anna*
Karenina principle" for bacterial microbiomes. Frontiers in microbiology, 2019. **10**: p. 975.
37. Plovie, A., *Comparison of bacterial communities associated with healthy and bleached Aiptasia*
pallida, a novel model organism for coral studies: Implications and variation during bleaching.
Biochemistry and Biotechnology. 2010, Liège University.
38. Gilbert, J.A., et al., *Microbial consortia increase thermal tolerance of corals.* Marine Biology,
2012. **159**(8): p. 1763-1771.
39. Ziegler, M., et al., *Bacterial community dynamics are linked to patterns of coral heat tolerance.*
Nat Commun, 2017. **8**: p. 14213.
40. Alagely, A., et al., *Signaling-mediated cross-talk modulates swarming and biofilm formation in a*
coral pathogen Serratia marcescens. ISME J, 2011. **5**(10): p. 1609-20.
41. dos Santos, F.A.H., et al., *Impact of oil spills on coral reefs can be reduced by bioremediation*
using probiotic microbiota. Sci Rep, 2015. **5**: p. 18268.
42. Rosado, P.M., et al., *Marine probiotics: increasing coral resistance to bleaching through*
microbiome manipulation. The ISME Journal, 2018.
43. Herrera, M., et al., *Laboratory-Cultured Strains of the Sea Anemone Exaiptasia Reveal Distinct*
Bacterial Communities. Frontiers in Marine Science, 2017. **4**.
44. Binsarhan, M., *Comparative analysis and culturing of the microbial community of Aiptasia*
pallida, A Sea Anemone Model for Coral Biology. 2016.
45. Röthig, T., et al., *Distinct Bacterial Communities Associated with the Coral Model Aiptasia in*
Aposymbiotic and Symbiotic States with Symbiodinium. Frontiers in Marine Science, 2016. **3**.

46.   Ceh, J., M. van Keulen, and D.G. Bourne, *Intergenerational transfer of specific bacteria in corals*
and possible implications for offspring fitness. Microb Ecol, 2013. **65**(1): p. 227-31.
47.   Rothig, T., et al., *Microbial community composition of deep-sea corals from the Red Sea provides*
insight into functional adaption to a unique environment. Sci Rep, 2017. **7**: p. 44714.
48.   Chiu, J.M., et al., *Bacteria associated with skeletal tissue growth anomalies in the coral Platygyra*
carnosus. FEMS Microbiol Ecol, 2012. **79**(2): p. 380-91.
49.   Hai, N.V., *The use of probiotics in aquaculture.* Journal of Applied Microbiology, 2015. **119**(4): p.
917-935.
50.   Kesarcodi-Watson, A., et al., *Alteromonas macleodii 0444 and Neptunomonas sp. 0536, two*
novel probiotics for hatchery-reared Greenshell™ mussel larvae, Perna canaliculus. Aquaculture,
2010. **309**(1-4): p. 49-55.
51.   Riquelme, C., et al., *Potential probiotic strains in the culture of the Chilean scallop Argopecten*
purpuratus (Lamarck, 1819). Aquaculture, 1997. **154**(1): p. 17-26.
52.   Haryanti, H., et al., *ENHANCE PRODUCTION OF BLACK TIGER SHRIMP Penaeus monodon*
POSTLARVAE BY PROBIOTIC BACTERIUM Alteromonas sp. Indonesian Fisheries Research Journal,
2017. **7**(1): p. 1-6.
53.   Biebl, H., et al., *Description of Labrenzia alexandrii gen. nov., sp. nov., a novel*
alphaproteobacterium containing bacteriochlorophyll a, and a proposal for reclassification of
Stappia aggregata as Labrenzia aggregata comb. nov., of Stappia marina as Labrenzia marina
comb. nov. and of Stappia alba as Labrenzia alba comb. nov., and emended descriptions of the
genera Pannonibacter, Stappia and Roseibium, and of the species Roseibium denhamense and
Roseibium hamelinense. 2007.
54.   Mancuso, M., et al., *In vitro selection of bacteria and isolation of Probionts from farmed Sparus*
aurata with potential for use as probiotics. International Journal of Animal Biology, 2015. **1**(4): p.
93-98.
55.   Brown, T., et al., *Worldwide exploration of the microbiome harbored by the cnidarian model,*
Exaiptasia pallida (Agassiz in Verrill, 1864) indicates a lack of bacterial association specificity at a
lower taxonomic rank. PeerJ, 2017. **5**: p. e3235.
56.   Sharp, K.H., D. Distel, and V.J. Paul, *Diversity and dynamics of bacterial communities in early life*
stages of the Caribbean coral Porites astreoides. ISME J, 2012. **6**(4): p. 790-801.
57.   Littman, R.A., et al., *Diversities of coral-associated bacteria differ with location, but not species,*
for three acroporid corals on the Great Barrier Reef. FEMS Microbiol Ecol, 2009. **68**(2): p. 152-63.
58.   Kellogg, C.A., et al., *Comparing bacterial community composition between healthy and white*
plague-like disease states in Orbicella annularis using PhyloChip G3 microarrays. PLoS One, 2013.
8(11): p. e79801.
59.   El-Rhman, A.M.A., Y.A. Khattab, and A.M. Shalaby, *Micrococcus luteus and Pseudomonas species*
as probiotics for promoting the growth performance and health of Nile tilapia, Oreochromis
niloticus. Fish & Shellfish Immunology, 2009. **27**(2): p. 175-180.
60.   Osman, H., et al., *Improvement growth and immune status using a potential probiotic bacteria*
Micrococcus species among cultured Oreochromis niloticus. New York Science Journal, 2010.
3(10): p. 5-11.
61.   Lau, S.C.K., et al., *Winogradskyella poriferorum sp. nov., a novel member of the family*
Flavobacteriaceae isolated from a sponge in the Bahamas. International Journal of Systematic
and Evolutionary Microbiology, 2005. **55**(4): p. 1589-1592.
62.   Franco, A., et al., *Winogradskyella pocilloporae sp. Nov. Isolated from healthy tissue of the coral*
Pocillopora damicornis. Vol. 68. 2018.
63.   Blackall, L.L., B. Wilson, and M.J. van Oppen, *Coral-the world's most diverse symbiotic*
ecosystem. Mol Ecol, 2015. **24**(21): p. 5330-47.

64.   Sunda, W., et al., *An antioxidant function for DMSP and DMS in marine algae.* Nature, 2002.
418(6895): p. 317-20.

65.   Curson, A.R., et al., *Dimethylsulfoniopropionate biosynthesis in marine bacteria and*
identification of the key gene in this process. Nat Microbiol, 2017. **2**: p. 17009.

66.   Raina, J.-B., et al., *Coral-Associated Bacteria and Their Role in the Biogeochemical Cycling of*
Sulfur. Applied and Environmental Microbiology, 2009. **75**(11): p. 3492-3501.

67.   Hatton, A.D., et al., *Metabolism of DMSP, DMS and DMSO by the cultivable bacterial community*
associated with the DMSP-producing dinoflagellate Scrippsiella trochoidea. Biogeochemistry,
2012. **110**(1): p. 131-146.

68.   Fiedor, J. and K. Burda, *Potential role of carotenoids as antioxidants in human health and*
disease. Nutrients, 2014. **6**(2): p. 466-488.

69.   Fiedor, J., et al., *Cyclic endoperoxides of β-carotene, potential pro-oxidants, as products of*
chemical quenching of singlet oxygen. Biochimica et Biophysica Acta (BBA) - Bioenergetics, 2005.
1709(1): p. 1-4.

70.   Flórez, L.V., et al., *Defensive symbioses of animals with prokaryotic and eukaryotic*
microorganisms. Natural Product Reports, 2015. **32**(7): p. 904-936.

71.   Asker, D., T. Beppu, and K. Ueda, *Mesoflavibacter zeaxanthinifaciens gen. nov., sp. nov., a novel*
zeaxanthin-producing marine bacterium of the family Flavobacteriaceae. Systematic and applied
microbiology, 2007. **30**(4): p. 291-296.

72.   Shindo, K., et al., *Rare carotenoids, (3R)-saproxanthin and (3R,2'S)-myxol, isolated from novel*
marine bacteria (Flavobacteriaceae) and their antioxidative activities. Applied Microbiology and
Biotechnology, 2007. **74**(6): p. 1350.

73.   Gammone, M., et al., *Marine Carotenoids against Oxidative Stress: Effects on Human Health.*
Marine Drugs, 2015. **13**(10): p. 6226.

74.   Mohana, D., S. Thippeswamy, and R. Abhishe, *Antioxidant, antibacterial, and ultraviolet-*
protective properties of carotenoids isolated from Micrococcus spp. Radiation Protection and
Environment, 2013. **36**(4): p. 168-174.

75.   Motone, K., et al., *A Zeaxanthin-Producing Bacterium Isolated from the Algal Phycosphere*
Protects Coral Endosymbionts from Environmental Stress. mBio, 2020. **11**(1): p. e01019-19.

76.   Croft, M.T., et al., *Algae acquire vitamin B 12 through a symbiotic relationship with bacteria.*
Nature, 2005. **438**(7064): p. 90.

77.   Raux, E., H. Schubert, and M. Warren, *Biosynthesis of cobalamin (vitamin B12): a bacterial*
conundrum. Cellular and Molecular Life Sciences CMLS, 2000. **57**(13-14): p. 1880-1893.

78.   Matthews, J.L., et al., *Symbiodiniaceae-bacteria interactions: rethinking metabolite exchange in*
reef-building corals as multi-partner metabolic networks. Environmental Microbiology.

79.   Agostini, S., et al., *Coral symbiotic complex: Hypothesis through vitamin B12 for a new*
evaluation. Galaxea, Journal of Coral Reef Studies, 2009. **11**(1): p. 1-11.

80.   Smith, S.M., *Complementarity in the Coral Holobiont: A Genomic Analysis of Bacterial Isolates of*
Orbicella faveolata and Symbiodinium spp. 2018.

81.   Imlay, J.A., *Where in the world do bacteria experience oxidative stress?* Environ Microbiol, 2018.

82.   Taylor, W.I. and D. Achanzar, *Catalase Test as an Aid to the Identification of*
Enterobacteriaceae. Applied Microbiology, 1972. **24**(1): p. 58-61.

83.   Cabiscol, E., J. Tamarit, and J. Ros, *Oxidative stress in bacteria and protein damage by reactive*
oxygen species. Int Microbiol, 2000. **3**(1): p. 3-8.

84.   Bruno, J.F., et al., *Thermal stress and coral cover as drivers of coral disease outbreaks.* PLoS
biology, 2007. **5**(6).

85.   Selig, E.R., et al., *Analyzing the relationship between ocean temperature anomalies and coral*
disease outbreaks at broad spatial scales. Coral reefs and climate change: Science and
management, 2006. **61**: p. 111-128.
86.   Muller, E., et al., *Bleaching increases likelihood of disease on Acropora palmata (Lamarck) in*
Hawksnest Bay, St John, US virgin islands. Coral Reefs, 2008. **27**(1): p. 191-195.
87.   Patterson, K.L., et al., *The etiology of white pox, a lethal disease of the Caribbean elkhorn coral,*
Acropora palmata. Proceedings of the National Academy of Sciences, 2002. **99**(13): p. 8725-
8730.
88.   Damjanovic, K., et al., *Mixed-mode bacterial transmission in the common brooding coral*
Pocillopora acuta. Environmental Microbiology, 2020. **22**(1): p. 397-412.
89.   Sharp, K.H., et al., *Bacterial acquisition in juveniles of several broadcast spawning coral species.*
PLoS One, 2010. **5**(5): p. e10898.
90.   Sharp, K.H., et al., *Induction of Larval Settlement in the Reef Coral Porites astreoides by a*
Cultivated Marine Roseobacter Strain. Biol. Bull., 2015. **228**: p. 98-107.
91.   Prabhu, S., P. Rekha, and A. Arun, *Zeaxanthin biosynthesis by members of the genus Muricauda.*
Pol. J. Microbiol, 2014. **63**: p. 115-119.
92.   Omori, M., *Degradation and restoration of coral reefs: Experience in Okinawa, Japan.* Marine
Biology Research, 2011. **7**(1): p. 3-12.
93.   Young, C., S. Schopmeyer, and D. Lirman, *A review of reef restoration and coral propagation*
using the threatened genus Acropora in the Caribbean and Western Atlantic. Bulletin of Marine
Science, 2012. **88**(4): p. 1075-1098.
94.   Rinkevich, B., *Rebuilding coral reefs: does active reef restoration lead to sustainable reefs?*
Current Opinion in Environmental Sustainability, 2014. **7**: p. 28-36.
95.   Buerger, P., et al., *Heat-evolved microalgal symbionts increase coral bleaching tolerance.* Science
Advances, 2020. **in press**.
96.   Chan, W.Y., et al., *Interspecific hybridization may provide novel opportunities for coral reef*
restoration. Frontiers in Marine Science, 2018. **5**: p. 160.
97.   Morikawa, M.K. and S.R. Palumbi, *Using naturally occurring climate resilient corals to construct*
bleaching-resistant nurseries. Proceedings of the National Academy of Sciences, 2019. **116**(21):
p. 10586-10591.
98.   Quigley, K., et al., *Assessing the role of historical temperature regime and algal symbionts on the*
heat tolerance of coral juveniles. Biology Open, 2020.
99.   Dungan, A.M., et al., *Exaiptasia diaphana from the great barrier reef: a valuable resource for*
coral symbiosis research. Symbiosis, 2020: p. 1-12.
100.   Lane, D., *16S/23S rRNA sequencing.* Nucleic acid techniques in bacterial systematics, 1991: p.
115-175.
101.   Muyzer, G., E.C. De Waal, and A.G. Uitterlinden, *Profiling of complex microbial populations by*
denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes
coding for 16S rRNA. Appl. Environ. Microbiol., 1993. **59**(3): p. 695-700.
102.   Lane, D.J., et al., *Rapid determination of 16S ribosomal RNA sequences for phylogenetic analyses.*
Proceedings of the National Academy of Sciences, 1985. **82**(20): p. 6955-6959.
103.   Takao, T., et al., *A simple screening method for antioxidants and isolation of several antioxidants*
produced by marine bacteria from fish and shellfish. Bioscience, Biotechnology, and
Biochemistry, 1994. **58**(10): p. 1780-1783.
104.   Velho-Pereira, S., P. Parvatkar, and I.J. Furtado, *Evaluation of antioxidant producing potential of*
halophilic bacterial bionts from marine invertebrates. Indian Journal of Pharmaceutical Sciences,
2015. **77**(2): p. 183-189.

105.   Souvorov, A., R. Agarwala, and D.J. Lipman, *SKESA: strategic k-mer extension for scrupulous*
713   *assemblies.* Genome Biol, 2018. **19**(1): p. 153.
714   106.   Seemann, T., *Prokka: rapid prokaryotic genome annotation.* Bioinformatics, 2014. **30**(14): p.
715   2068-9.
716   107.   Wood, D.E. and S.L. Salzberg, *Kraken: ultrafast metagenomic sequence classification using exact*
717   *alignments.* Genome Biol, 2014. **15**(3): p. R46.
718   108.   Parks, D.H., et al., *A complete domain-to-species taxonomy for Bacteria and Archaea.* Nature
719   Biotechnology, 2020.
720   109.   Perisin, M., et al., *16Stimator: statistical estimation of ribosomal gene copy numbers from draft*
721   *genome assemblies.* The ISME journal, 2016. **10**(4): p. 1020-1024.
722   110.   Brettin, T., et al., *RASTtk: a modular and extensible implementation of the RAST algorithm for*
723   *building custom annotation pipelines and annotating batches of genomes.* Scientific reports,
724   2015. **5**: p. 8365.
725   111.   Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nature methods,
726   2012. **9**(4): p. 357.
727   112.   R Core Team, *A language and environment for statistical computing. Vienna, Austria: R*
728   *Foundation for Statistical Computing; 2012.* URL https://www. R-project. org, 2019.
729   113.   Chambers, J.M. and T.J. Hastie, *Statistical models in S*. Vol. 251. 1992: Wadsworth & Brooks/Cole
730   Advanced Books & Software Pacific Grove, CA.
731   114.   Miller, R.G., *Normal univariate techniques*, in *Simultaneous statistical inference*. 1981, Springer.
732   p. 37-108.
733   115.   Yandell, B., *Practical data analysis for designed experiments*. 2017: Routledge.
734   116.   Holland, M. and D. Wolfe, *Nonparametric statistical methods*. 1973, John Wiley & Sons, New
735   York.
736   117.   Dunn, O.J., *Multiple comparisons using rank sums.* Technometrics, 1964. **6**(3): p. 241-252.
737   118.   Ferreira, J. and A. Zwinderman, *On the Benjamini–Hochberg method.* The Annals of Statistics,
738   2006. **34**(4): p. 1827-1849.
739   119.   Ogle, D.H., *FSA: fisheries stock analysis.* R package version 0.8, 2017. **17**: p. 636.

741   **Table S1**: Isolate Genome Sequence Data Summary. Strains are presented as high FRS (grey) followed by low FRS (white). 16S rRNA
742   gene presumptive identity is derived from the NCBI classification of near-complete 16S rRNA gene sequences. *We were unable to
743   determine the 16S rRNA copy number of isolate MMSF00068 due to the presence of contaminant sequences in the WGS.

Isolate	Genus Level Classification	Genus % Confidence	Family	Family % Confidence	SRA Accession	Sample Accession
MMSF00257	*Alteromonas*	96.65	*Alteromonadaceae*	96.74	SRR10186803	SAMN12851724
MMSF00958	*Alteromonas*	92.93	*Alteromonadaceae*	92.98	SRR10186806	SAMN12851731
MMSF01163	*Alteromonas*	9.28	*Alteromonadaceae*	12.01	SRR10186808	SAMN12851721
MMSF00404	*Alteromonas*	8.15	*Alteromonadaceae*	10.49	SRR10186805	SAMN12851732
MMSF00132	*Labrenzia*	90.9	*Rhodobacteraceae*	90.91	SRR10186800	SAMN12851727
MMSF00249	*Labrenzia*	90.83	*Rhodobacteraceae*	90.83	SRR10186799	SAMN12851728
MMSF00964	*Marinobacter*	59.99	*Alteromonadaceae*	60.1	SRR10186802	SAMN12851725
MMSF01190	*Marinobacter*	65.77	*Alteromonadaceae*	65.85	SRR10186797	SAMN12851730
MMSF00068	*Micrococcus*	71.81	*Micrococcaceae*	72.88	SRR10186807	SAMN12851722
MMSF00107	*Micrococcus*	83.18	*Micrococcaceae*	84.87	SRR10186798	SAMN12851729
MMSF00046	*Winogradskyella*	49.63	*Flavobacteriaceae*	49.72	SRR10186801	SAMN12851726
MMSF00910	*Winogradskyella*	54.44	*Flavobacteriaceae*	54.53	SRR10186804	SAMN12851723

Isolate	16S rRNA gene presumptive identity	Confidence	Reads	Total Bases	G+C%	Avg. Read Length	Max Read Length	Avg Quality
MMSF00257	*Alteromonas macleodii*	1.00	2233080	334079592	44.7	149	151	30.9
MMSF00958	*Alteromonas macleodii*	1.00	2971718	443870758	44.6	149	151	31.9
MMSF01163	*Alteromonas oceani*	0.83	2757340	412741366	48.7	149	151	33.7
MMSF00404	*Alteromonas oceani*	0.83	2067800	306005114	48.7	147	151	33.3
MMSF00132	*Labrenzia aggregata*	0.98	3758944	557529936	59.2	148	151	31.1
MMSF00249	*Labrenzia aggregata*	0.98	3333098	497032699	59.3	149	151	30.7
MMSF00964	*Marinobacter salsuginis*	1.00	1713198	256635737	57.1	149	151	31
MMSF01190	*Marinobacter salsuginis*	1.00	3239568	483874048	57.1	149	151	33.1
MMSF00068	*Micrococcus luteus*	0.99	2658898	398496031	72.4	149	151	30.3

MMSF00107	*Micrococcus yunnanensis*	0.99	2323654	348617190	72.8	150	151	31.9
MMSF00046	*Winogradskyella poriferorum*	0.59	1930584	285978916	35	148	151	31.8
MMSF00910	*Winogradskyella poriferorum*	0.59	3939880	579510587	35.6	147	151	33.9

Isolate	Est. Read Coverage	Contigs in Draft Genome	Bases in Draft Genome	Min Contig	Avg Contig	Max Contig	N50	16S rRNA gene copy number
MMSF00257	69	136	4831263	515	35523	210048	72973	5.15
MMSF00958	94	42	4732026	620	112667	580903	303003	4.79
MMSF01163	75	100	5507488	526	55074	394030	155957	4.18
MMSF00404	51	111	6014142	502	54181	256548	93725	3.97
MMSF00132	82	34	6792087	855	199767	1019681	302139	3.57
MMSF00249	73	44	6791310	851	154347	1286212	294458	3.02
MMSF00964	56	41	4588310	512	111910	932206	291012	3.67
MMSF01190	110	40	4404819	529	110120	769441	404351	3.21
MMSF00068	160	316	2484978	509	7863	78799	12670	*
MMSF00107	143	501	2435904	515	4862	42365	7717	1.75
MMSF00046	83	104	3446386	553	33138	299826	78549	1.03
MMSF00910	168	50	3456468	553	69129	355406	162623	0.77

747   **Table S2**: Pairwise comparison of the genome sequences between the pairs of isolates

NCBI Classification	Type strain	Core Genome	SNP difference
Alteromonas macleodii	ATCC 27126	85%	~60000
Alteromonas oceani	S35	60%	~190000
Labrenzia aggregata	IAM12614	55%	5
Marinobacter salsuginis	SD14B	15%	~120000
Micrococcus spp.	NCTC2665	70%	~35000
Winogradskyella poriferorum	NA	NA	10

749　**Table S3:** Search outcomes for genes of interest.

Isolate	NCBI Classification	Gene	Gene Classification	Contig	Start	End	Strand
MMSF00132	*Labrenzia aggregata*	*dddL*	positive	Contig_16_56.34	104067	104756	-
MMSF00249	*Labrenzia aggregata*	*dddL*	positive	Contig_18_45.1167	104045	104734	-
MMSF00132	*Labrenzia aggregata*	*dddP*	positive x2 (Copy 1)	Contig_6_57.2063	50787	52046	+
MMSF00132	*Labrenzia aggregata*	*dddP*	positive x2 (Copy 2)	Contig_6_57.2063	650583	651881	-
MMSF00249	*Labrenzia aggregata*	*dddP*	positive x2 (Copy 1)	Contig_17_46.7174	317316	318575	+
MMSF00249	*Labrenzia aggregata*	*dddP*	positive x2 (Copy 2)	Contig_17_46.7174	917112	918410	-
MMSF00132	*Labrenzia aggregata*	*dsyB*	positive	Contig_32_56.2994	210397	211419	+
MMSF00249	*Labrenzia aggregata*	*dsyB*	positive	Contig_4_46.0396	460215	461237	+
MMSF00046	*Winogradskyella poriferorum*	*katG*	positive	Contig_15_55.0851	16430	18661	+
MMSF00068	*Micrococcus luteus*	*katG*	ND; katA & katE detected	NA	NA	NA	NA
MMSF00107	*Micrococcus yunnanensis*	*katG*	ND; katA & katE detected	NA	NA	NA	NA
MMSF00132	*Labrenzia aggregata*	*katG*	positive	Contig_6_57.2063	425873	428044	-
MMSF00249	*Labrenzia aggregata*	*katG*	positive	Contig_17_46.7174	692402	694573	-
MMSF00257	*Alteromonas macleodii*	*katG*	positive	Contig_101_46.6864	22130	24502	+
MMSF00404	*Alteromonas oceani*	*katG*	positive x2 (Copy 1)	Contig_26_39.8992	70873	73035	-
MMSF00404	*Alteromonas oceani*	*katG*	positive x2 (Copy 2)	Contig_52_41.1626	55994	58225	-
MMSF00910	*Winogradskyella poriferorum*	*katG*	positive	Contig_39_101.483	25621	27852	+
MMSF00958	*Alteromonas macleodii*	*katG*	positive	Contig_21_66.072	22086	24458	+
MMSF00964	*Marinobacter salsuginis*	*katG*	positive	Contig_25_36.8412	50391	52565	-
MMSF01163	*Alteromonas oceani*	*katG*	positive x2 (Copy 1)	Contig_46_63.2119	54248	56479	-
MMSF01163	*Alteromonas oceani*	*katG*	positive x2 (Copy 2)	Contig_86_59.9301	369720	371882	-
MMSF01190	*Marinobacter salsuginis*	*katG*	positive	Contig_22_84.8909	7780	9954	+
MMSF00132	*Labrenzia aggregata*	*cobA*	positive	Contig_6_57.2063	552920	553741	-

MMSF00249	*Labrenzia aggregata*	*cobA*	positive	Contig_17_46.7174	819449	820270	-
MMSF00132	*Labrenzia aggregata*	*cobB*	positive	Contig_6_57.2063	551611	552930	-
MMSF00249	*Labrenzia aggregata*	*cobB*	positive	Contig_17_46.7174	818140	819459	-
MMSF00132	*Labrenzia aggregata*	*cobC*	positive	Contig_6_57.2063	842748	843392	-
MMSF00249	*Labrenzia aggregata*	*cobC*	positive	Contig_17_46.7174	1109277	1E+06	-
MMSF00132	*Labrenzia aggregata*	*cobD*	positive x2	Contig_26_56.6932	208207	209184	-
MMSF00132	*Labrenzia aggregata*	*cobD*	positive x2	Contig_26_56.6932	209194	210195	-
MMSF00249	*Labrenzia aggregata*	*cobD*	positive x2	Contig_15_47.504	84229	85230	+
MMSF00249	*Labrenzia aggregata*	*cobD*	positive x2	Contig_15_47.504	85240	86217	+
MMSF00132	*Labrenzia aggregata*	*cobH*	positive	Contig_6_57.2063	545294	545950	+
MMSF00249	*Labrenzia aggregata*	*cobH*	positive	Contig_17_46.7174	811823	812479	+
MMSF00132	*Labrenzia aggregata*	*cobI*	positive	Contig_6_57.2063	547369	548139	+
MMSF00249	*Labrenzia aggregata*	*cobI*	positive	Contig_17_46.7174	813898	814668	+
MMSF00132	*Labrenzia aggregata*	*cobK*	positive	Contig_6_57.2063	555025	555786	+
MMSF00249	*Labrenzia aggregata*	*cobK*	positive	Contig_17_46.7174	821554	822315	+
MMSF00132	*Labrenzia aggregata*	*cobL*	positive	Contig_6_57.2063	546203	547417	+
MMSF00249	*Labrenzia aggregata*	*cobL*	positive	Contig_17_46.7174	812732	813946	+
MMSF00132	*Labrenzia aggregata*	*cobM*	positive	Contig_6_57.2063	550028	550801	+
MMSF00249	*Labrenzia aggregata*	*cobM*	positive	Contig_17_46.7174	816557	817330	+
MMSF00132	*Labrenzia aggregata*	*cobN*	positive	Contig_26_56.6932	201965	205723	+
MMSF00249	*Labrenzia aggregata*	*cobN*	positive	Contig_15_47.504	88701	92459	-
MMSF00132	*Labrenzia aggregata*	*cobO*	positive	Contig_26_56.6932	205720	206358	+
MMSF00249	*Labrenzia aggregata*	*cobO*	positive	Contig_15_47.504	88066	88704	-
MMSF00964	*Marinobacter salsuginis*	*cobO*	positive	Contig_25_36.8412	168373	168930	+
MMSF01190	*Marinobacter salsuginis*	*cobO*	positive	Contig_21_84.3861	85977	86534	+
MMSF00132	*Labrenzia aggregata*	*cobP*	positive	Contig_26_56.6932	200255	200806	+
MMSF00249	*Labrenzia aggregata*	*cobP*	positive	Contig_15_47.504	93618	94169	-
MMSF00132	*Labrenzia aggregata*	*cobQ*	positive	Contig_26_56.6932	206715	208205	+
MMSF00249	*Labrenzia aggregata*	*cobQ*	positive	Contig_15_47.504	86219	87709	-

Supplemental Tables

MMSF00132	*Labrenzia aggregata*	*cobU*	positive	Contig_24_56.1446	78969	79994	+
MMSF00249	*Labrenzia aggregata*	*cobU*	positive	Contig_13_45.8339	40345	41370	+
MMSF00132	*Labrenzia aggregata*	*cobV*	positive	Contig_24_56.1446	77990	78862	-
MMSF00249	*Labrenzia aggregata*	*cobV*	positive	Contig_13_45.8339	39366	40238	-
MMSF00132	*Labrenzia aggregata*	*cbiD*	positive	Contig_6_57.2063	553941	555035	+
MMSF00249	*Labrenzia aggregata*	*cbiD*	positive	Contig_17_46.7174	820470	821564	+
MMSF00964	*Marinobacter salsuginis*	*cbiO*	positive	Contig_36_38.8121	420124	420720	+

752 **Table S4:** Summary of vitamin B12 biosynthesis pathway genes. Genes in red were not found in any isolate.

Isolate	Genus Level Classification	CobP as indicator	Gene Count	*CobA*	*CobB*	*CobC*	*CobD*	*CobE*	*CobF*	*CobG*
MMSF01163	*Alteromonas*	-	0	-	-	-	-	-	-	-
MMSF00257	*Alteromonas*	-	0	-	-	-	-	-	-	-
MMSF00958	*Alteromonas*	-	0	-	-	-	-	-	-	-
MMSF00404	*Alteromonas*	-	0	-	-	-	-	-	-	-
MMSF00132	*Labrenzia*	+	16	+	+	+	+ x2	-	-	-
MMSF00249	*Labrenzia*	+	16	+	+	+	+ x2	-	-	-
MMSF00964	*Marinobacter*	-	2	-	-	-	-	-	-	-
MMSF01190	*Marinobacter*	-	1	-	-	-	-	-	-	-
MMSF00068	*Micrococcus*	-	0	-	-	-	-	-	-	-
MMSF00107	*Micrococcus*	-	0	-	-	-	-	-	-	-
MMSF00910	*Winogradskyella*	-	0	-	-	-	-	-	-	-
MMSF00046	*Winogradskyella*	-	0	-	-	-	-	-	-	-

Isolate	Genus Level Classification	*CobH*	*CobI*	*CobJ*	*CobK*	*CobL*	*CobM*	*CobN*	*CobO*	*CobP*	*CobQ*
MMSF01163	*Alteromonas*	-	-	-	-	-	-	-	-	-	-
MMSF00257	*Alteromonas*	-	-	-	-	-	-	-	-	-	-
MMSF00958	*Alteromonas*	-	-	-	-	-	-	-	-	-	-
MMSF00404	*Alteromonas*	-	-	-	-	-	-	-	-	-	-
MMSF00132	*Labrenzia*	+	+	-	+	+	+	+	+	+	+
MMSF00249	*Labrenzia*	+	+	-	+	+	+	+	+	+	+
MMSF00964	*Marinobacter*	-	-	-	-	-	-	-	+	-	-
MMSF01190	*Marinobacter*	-	-	-	-	-	-	-	+	-	-
MMSF00068	*Micrococcus*	-	-	-	-	-	-	-	-	-	-
MMSF00107	*Micrococcus*	-	-	-	-	-	-	-	-	-	-

MMSF00910	*Winogradskyella*	-	-	-	-	-	-	-	-	-	-	
MMSF00046	*Winogradskyella*	-	-	-	-	-	-	-	-	-	-	

Isolate	Genus Level Classification	CobP	CobQ	CobR	CobS	CobT	CobU	CobV	CobW	CobX	CobY	CobZ	Cbi
MMSF01163	*Alteromonas*	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00257	*Alteromonas*	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00958	*Alteromonas*	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00404	*Alteromonas*	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00132	*Labrenzia*	+	+	-	-	-	+	+	-	-	-	-	cbiD
MMSF00249	*Labrenzia*	+	+	-	-	-	+	+	-	-	-	-	cbiD
MMSF00964	*Marinobacter*	-	-	-	-	-	-	-	-	-	-	-	cbiO
MMSF01190	*Marinobacter*	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00068	*Micrococcus*	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00107	*Micrococcus*	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00910	*Winogradskyella*	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00046	*Winogradskyella*	-	-	-	-	-	-	-	-	-	-	-	-

756   **Table S5:** Composition of R2A broth adjusted to suit marine bacteria. Final pH = 7.2 +/- 0.2 at

757   26 °C.

Component	grams L^{-1}	Supplier
Casein acid hydrolysate	0.500	Cat#C0501, Sigma-Aldrich
Yeast extract	0.500	Cat#LP0021, Oxoid
Proteose peptone	0.500	Cat#211684, ThermoFisher
Dextrose	0.500	Cat#G360, PhytoTech Laboratories
Starch, soluble	0.500	Cat#AJA526, Univar
Dipotassium phosphate	0.300	Cat#P3786, Sigma-Aldrich
Magnesium sulfate	0.024	Cat#M2643, Sigma-Aldrich
Sodium pyruvate	0.300	Cat#P2256, Sigma-Aldrich
Red Sea Salt™	40.00	Cat#R11065, Red Sea

759 **Table S6:** Single-copy housekeeping genes extracted from the RAST annotations

Gene	Function	NCBI
16S	SSU rRNA ## 16S rRNA, small subunit ribosomal RNA	16S ribosomal RNA
23S	LSU rRNA ## 23S rRNA, large subunit ribosomal RNA	23S ribosomal RNA
fusA	Translation elongation factor G	elongation factor G
gyrB	DNA gyrase subunit B (EC 5.99.1.3)	DNA gyrase subunit B
gyrA	DNA gyrase subunit A (EC 5.99.1.3)	DNA gyrase subunit A
pyrG	CTP synthase (EC 6.3.4.2)	CTP synthetase
lepA	Translation elongation factor LepA	GTP-binding protein LepA
recA	RecA protein	recA protein
recG	ATP-dependent DNA helicase RecG (EC 3.6.4.12)	ATP-dependent DNA helicase RecG
rpoB	DNA-directed RNA polymerase beta subunit (EC 2.7.7.6)	DNA-directed RNA polymerase subunit beta
rpoD	RNA polymerase sigma factor RpoD	RNA polymerase sigma factor RpoD
atpD	ATP synthase beta chain (EC 3.6.3.14)	F0F1 ATP synthase subunit beta
ppK	Polyphosphate kinase (EC 2.7.4.1)	polyphosphate kinase
polA	DNA polymerase I (EC 2.7.7.7)	DNA polymerase I
pheSa	Phenylalanyl-tRNA synthetase alpha chain (EC 6.1.1.20)	phenylalanyl-tRNA synthetase subunit alpha
pheSb	Phenylalanyl-tRNA synthetase beta chain (EC 6.1.1.20)	phenylalanyl-tRNA synthetase subunit beta
ileS	Isoleucyl-tRNA synthetase (EC 6.1.1.5)	isoleucyl-tRNA synthetase
leuS	Leucyl-tRNA synthetase (EC 6.1.1.4)	leucyl-tRNA synthetase
aspS	Aspartyl-tRNA synthetase (EC 6.1.1.12)	aspartyl-tRNA synthetase
alaS	Alanyl-tRNA synthetase (EC 6.1.1.7)	alanyl-tRNA synthetase
argS	Arginyl-tRNA synthetase (EC 6.1.1.19)	arginyl-tRNA synthetase
hisS	Histidyl-tRNA synthetase (EC 6.1.1.21)	histidyl-tRNA synthetase
valS	Valyl-tRNA synthetase (EC 6.1.1.9)	valyl-tRNA synthetase
srp1	SSU ribosomal protein S1p	30S ribosomal protein S1
srp2	SSU ribosomal protein S2p (SAe)	30S ribosomal protein S2
srp3	SSU ribosomal protein S3p (S3e)	30S ribosomal protein S3
srp4	SSU ribosomal protein S4p (S9e)	30S ribosomal protein S4
srp5	SSU ribosomal protein S5p (S2e)	30S ribosomal protein S5
srp6	SSU ribosomal protein S6p	30S ribosomal protein S6
srp7	SSU ribosomal protein S7p (S5e)	30S ribosomal protein S7
srp8	SSU ribosomal protein S8p (S15Ae)	30S ribosomal protein S8
srp9	SSU ribosomal protein S9p (S16e)	30S ribosomal protein S9
srp10	SSU ribosomal protein S10p (S20e)	30S ribosomal protein S10

srp11	SSU ribosomal protein S11p (S14e)	30S ribosomal protein S11
srp12	SSU ribosomal protein S12p (S23e)	30S ribosomal protein S12
srp13	SSU ribosomal protein S13p (S18e)	30S ribosomal protein S13
srp14	SSU ribosomal protein S14p (S29e) @ SSU ribosomal protein S14p (S29e), zinc-independent	30S ribosomal protein S14
srp15	SSU ribosomal protein S15p (S13e)	30S ribosomal protein S15
srp16	SSU ribosomal protein S16p	30S ribosomal protein S16
srp17	SSU ribosomal protein S17p (S11e)	30S ribosomal protein S17
srp18	SSU ribosomal protein S18p @ SSU ribosomal protein S18p, zinc-independent	30S ribosomal protein S18
srp19	SSU ribosomal protein S19p (S15e)	30S ribosomal protein S19
srp20	SSU ribosomal protein S20p	30S ribosomal protein S20
srp21	SSU ribosomal protein S21p	30S ribosomal protein S21
lrp1	LSU ribosomal protein L1p (L10Ae)	50S ribosomal protein L1
lrp2	LSU ribosomal protein L2p (L8e)	50S ribosomal protein L2
lrp3	LSU ribosomal protein L3p (L3e)	50S ribosomal protein L3
lrp4	LSU ribosomal protein L4p (L1e)	50S ribosomal protein L4
lrp5	LSU ribosomal protein L5p (L11e)	50S ribosomal protein L5
lrp6	LSU ribosomal protein L6p (L9e)	50S ribosomal protein L6
lrp7	LSU ribosomal protein L7/L12 (P1/P2)	50S ribosomal protein L7
lrp8	LSU ribosomal protein L8p	50S ribosomal protein L8
lrp9	LSU ribosomal protein L9p	50S ribosomal protein L9
lrp10	LSU ribosomal protein L10p (P0)	50S ribosomal protein L10
lrp11	LSU ribosomal protein L11p (L12e)	50S ribosomal protein L11
lrp12	LSU ribosomal protein L12p	50S ribosomal protein L12
lrp13	LSU ribosomal protein L13p (L13Ae)	50S ribosomal protein L13
lrp14	LSU ribosomal protein L14p (L23e)	50S ribosomal protein L14
lrp15	LSU ribosomal protein L15p (L27Ae)	50S ribosomal protein L15
lrp16	LSU ribosomal protein L16p (L10e)	50S ribosomal protein L16
lrp17	LSU ribosomal protein L17p	50S ribosomal protein L17
lrp18	LSU ribosomal protein L18p (L5e)	50S ribosomal protein L18
lrp19	LSU ribosomal protein L19p	50S ribosomal protein L19
lrp20	LSU ribosomal protein L20p	50S ribosomal protein L20
lrp21	LSU ribosomal protein L21p	50S ribosomal protein L21
lrp22	LSU ribosomal protein L22p (L17e)	50S ribosomal protein L22
lrp23	LSU ribosomal protein L23p (L23Ae)	50S ribosomal protein L23
lrp24	LSU ribosomal protein L24p (L26e)	50S ribosomal protein L24
lrp25	LSU ribosomal protein L25p	50S ribosomal protein L25
lrp26	LSU ribosomal protein L26p	50S ribosomal protein L26
lrp27	LSU ribosomal protein L27p	50S ribosomal protein L27
lrp28	LSU ribosomal protein L28p	50S ribosomal protein L28

lrp29	LSU ribosomal protein L29p (L35e)	50S ribosomal protein L29
lrp30	LSU ribosomal protein L30p (L7e)	50S ribosomal protein L30
lrp31	LSU ribosomal protein L31p @ LSU ribosomal protein L31p, zinc-independent	50S ribosomal protein L31