Genomic analysis of the diversity, antimicrobial resistance and virulence potential

of *Campylobacter jejuni* and *Campylobacter coli* strains from a private health care

center in Central Chile

Bravo V[1,2], Katz A[3], Porte L[4], Weitzel T[4,5], Varela, C[4], Gonzalez-Escalona N[6], Blondel

CJ[2,7]*

[1] Unidad de Microbiología, Escuela de Medicina, Facultad de Ciencias Médicas,

Universidad de Santiago de Chile, Santiago, Chile.

[2] Instituto de Ciencias Biomédicas, Facultad de Medicina y Facultad de Ciencias de la

Vida, Universidad Andrés Bello, 8320000, Santiago, Chile.

[3] Programa de Biología Celular y Molecular, ICBM, Facultad de Medicina, Universidad

de Chile, Santiago, 8380453, Chile.

[4] Laboratorio Clínico, Clínica Alemana de Santiago, Facultad de Medicina Clínica

Alemana, Universidad del Desarrollo, Santiago, Chile.

[5] Instituto de Ciencias e Innovación en Medicina (ICIM), Facultad de Medicina Clínica

Alemana, Universidad del Desarrollo, Santiago, Chile.

[6] Center for Food Safety and Applied Nutrition, Office of Regulatory Science, Division of

Microbiology, U.S. Food and Drug Administration, College Park, Maryland, USA.

[7] Instituto de Ciencias Biomédicas, Facultad de Ciencias de la Salud, Universidad

Autónoma de Chile, Santiago, Chile.

Key words: *Campylobacter jejuni*, *Campylobacter coli*, cgMLST, comparative genomics,

foodborne pathogen, Chile.

*Corresponding author. carlos.blondel@unab.cl

Mailing address, Instituto de Ciencias Biomédicas, Universidad Andrés Bello. Echaurren

183, Santiago, 8320000, Chile.

ABSTRACT

Campylobacter jejuni and *Campylobacter coli* are the leading causes of human gastroenteritis in the developed world and an emerging threat in developing countries. There is increasing evidence that the incidence of campylobacteriosis in Chile and other South American countries is greatly underestimated, mostly due to the lack of adequate diagnostic methods. Accordingly, there is limited genomic and epidemiological data from these countries, leaving an important knowledge gap in our understanding of the population structure, virulence potential, and antimicrobial resistance profiles of clinical *Campylobacter* strains. We recently published the whole-genome sequences of a collection of 68 *C. jejuni* and 12 *C. coli* strains isolated from patients with enteric infections during a 2-year period (2017-2019) attending Clinica Alemana in Santiago, Chile. In the present study, we sequenced an additional *C. jejuni* genome and performed a genome-wide analysis of the genetic diversity, virulence, and antimicrobial resistance potential of these 81 strains. cgMLST analysis identified a high genetic diversity of *C. jejuni* as well as 13 novel STs and alleles in both *C. jejuni* and *C. coli*. Pangenome and virulome analyses showed a differential distribution of virulence factors, including both plasmid and chromosomally encoded T6SSs and T4SSs. Resistome analysis predicted widespread resistance to fluoroquinolones, but a low rate of erythromycin resistance. This study provides valuable genomic and epidemiological data and highlights the need for larger genomic epidemiology studies to better understand the genetic diversity, virulence potential and antimicrobial resistance of human clinical *Campylobacter* strains in South America.

DATA SUMMARY

1. Genome sequences and assemblies of all *C. jejuni* and *C. coli* used in this study have been deposited in public databases (**Table S1**) and most strains have also been published as genome resource announcements [1,2].

2. Bacterial draft genome sequence reannotations and nucleotide sequence fasta files used for virulome and plasmid screening analysis are available as a supplementary dataset on Zenodo (DOI: 10.5281/zenodo.3925206) https://doi.org/10.5281/zenodo.3925206.

3. The reconstructed phylogenetic trees and associated metadata of clinical strains from Chile in the global context of *C. jejuni* and *C. coli* sequenced strains can be interactively visualized on Microreact at https://microreact.org/project/VbEQsZtQD.

INTRODUCTION

Campylobacteriosis caused by *Campylobacter jejuni* and *Campylobacter coli* has emerged as an important public health problem and is the most common bacterial cause of human gastroenteritis worldwide [3–5]. Although the mortality rates associated with campylobacteriosis are low (<0.01% to 8.8%), this foodborne illness has emerged as an important public health threat worldwide due to its high morbidity rates and the emergence of antimicrobial resistance [6,7]. In this context, the European Food Safety Authority (EFSA) has estimated a disease burden and annual cost of 2.4 billion euros [8].

Campylobacteriosis is usually associated with a self-limiting illness characterized by diarrhea, nausea, abdominal cramps and bloody stools [9]. Only a minor percentage of cases are severe or fatal. Nevertheless, extraintestinal manifestations might cause long-term complications such as reactive arthritis, and Miller-Fisher or Guillain-Barré syndrome [10,11]. Despite the self-limiting course, certain populations such as children under 5 years of age, elderly patients, and immunocompromised patients might suffer severe infections and require antimicrobial treatment [9]. In this context, the World Health Organization (WHO) has included *Campylobacter* as a high priority pathogen [7] due to the worldwide emergence of strains with high level fluoroquinolone resistance [12,13].

In South America, there is limited data available on the prevalence of *C. jejuni* and *C. coli* in comparison to other enteric bacterial pathogens [3,14]. Furthermore, the incidence and number of cases reported varies substantially between countries, most probably due to differences in sensitivity of the diagnostic methods used in each surveillance program.

In Chile, the National Laboratory Surveillance Program (DTO. N° 7/2020) of the Public

Health Institute of Chile (ISP) has reported a national incidence rate of

campylobacteriosis of 0.1 to 0.6/100.000 [15]. This is quite low compared to other high-

income countries, where incidence rates are two orders of magnitude higher [3].

Nevertheless, recent reports suggest that the burden of campylobacteriosis in Chile has

been greatly underestimated due to suboptimal diagnostic protocols in both public and

private health centers [15]. A recent study showed a 10-fold increase of the detection

rate of *Campylobacter* in human stool samples after changing from microscopic stool

examination by Gram staining (the routine protocol in Chilean health centers) to culture-

dependent techniques [16]. In another study from the same group, *Campylobacter* spp.

emerged as the second most prevalent bacterial agents after diarrheagenic *Escherichia*

coli when a multiplex PCR based assay (FilmArray® Gastrointestinal Panel) was used

[17].

To date, only few reports have analyzed the genetic diversity and antimicrobial

resistance profiles of clinical human isolates of *Campylobacter* in Chile [18–21].

Furthermore, these studies have either used a limited number of strains or lacked the

resolution provided by sequence-based typing methods such as multilocus sequence

typing (MLST). There is only one report that used whole-genome sequence data

(wgMLST) to determine the genetic diversity of *C. jejuni* strains from the country, which

was based on the first 3 genomes of Chilean *C. jejuni* strains. Therefore larger genomic

epidemiology studies are needed [22].

We recently published the whole-genome sequences of 69 *C. jejuni* and 12 *C. coli*

human clinical strains isolated from patients with enteric infection attending Clínica

Alemana in Santiago, Chile, during a 2-year period (2017-2019) [1]. For 17 of the *C.*

jejuni strains, we obtained complete genome assemblies [2]. This represents the largest

collection of genomes from Chilean human clinical *Campylobacter* strains and includes

the first *C. coli* genome in the country [1].

In the present study, we performed a genome-wide analysis of the genetic diversity,

virulence potential, and antimicrobial resistance profiles of these 81 clinical strains.

Importantly, we identified a high genetic diversity of both *C. jejuni* and *C. coli* strains,

including 13 novel sequence types (STs) and alleles. Resistome analysis predicted a

widespread resistance to fluoroquinolones, but a decreased resistance to tetracycline in

comparison to the reported resistance from other regions of the world, in addition to the

differential distribution of virulence factors, including plasmids and chromosomally

encoded T6SS and T4SS.

METHODS

Genome data set, sequencing and reannotation

The genome sequence dataset analyzed in this study is listed in **Table S1** and was

compiled from our recently published collection of human clinical strains of *C. jejuni* and

C. coli from Central Chile [1,2] and available whole-genome sequences of *C. jejuni* and

C. coli strains from public databases. We additionally included in this dataset the draft

whole genome sequence of *C. jejuni* strain CFSAN096303, from the same collection of

Clinica Alemana isolates which was sequenced for this study. The final genome

composed of 2 contigs was obtained using a combination of long (nanopore) and short

(Illumina, NextSeq) sequencing reads. Briefly, for the short reads the sequencing library

was prepared using 1 ng DNA for the Nextera XT kit according to the manufacturer's

instructions. The strain was sequenced using the NextSeq sequencer (Illumina). We

used a NextSeq 500/550 high-output kit v2.5 (300 cycles) with 2 × 150-bp paired-end

chemistry, according to the manufacturer's instructions, with >200X average coverage.

The long reads were generated in a MinION sequencer (Nanopore Technologies,

Oxford, UK). The sequencing library was prepared using the rapid barcoding sequencing

kit (SQK-RBK004). The library for this strain was pooled with other 9 libraries and run in

a FLO-MIN106 (R9.4.1) flow cell, according to the manufacturer's instructions, for 48 h.

The run was base called live using default settings (Minknow v 19.06.8, Guppy v3.0.7).

Default parameters were used for all software unless otherwise specified. The

sequencing outputs for that run was 7.2 Gb (quality score, 11.46; N50, 6,000 bp; total

reads, 569,349) for an estimated genome average coverage of 60 to 90 X. Reads less

than 5,000 bp and quality scores of <7 were discarded for downstream analysis. To

assemble the genome of *C. jejuni* strain CFSAN096303, the Illumina reads were

managed with the CLC Genomics Workbench v9.5.2 (Qiagen), assessed for quality

(Q,□>30) with the quality control tool, and trimmed (adapter trimming, quality trimming,

and length trimming) with the trim sequences tool. The final complete genome sequence

for this strain was obtained using a previously described pipeline [1], except that Flye

was used instead of CANU for long read de novo assembling. The genomes were

annotated using the NCBI Prokaryotic Genomes Automatic Annotation Pipeline (PGAP,

http://www.ncbi.nlm.nih.gov/genome/annotation_prok). Default parameters were used for

all software unless otherwise specified. The NextSeq data was deposited in NCBI under

accession number SRR10859573. The nanopore raw data was deposited in NCBI under

accession number SRR11786978. The final draft genome was deposited under

accession number WXZQ01000000. Fasta files of all draft genomes from the collection

of strains from Chile were reordered to the reference genomes of the *C. coli* aerotolerant

strain OR12 (accID GCF_002024185.1) and *C. jejuni* strain NCTC 11168 (accID

GCF_000009085.1) using the Mauve Contig Mover (MCM) from the Mauve package

[23]. The ordered genomes were reannotated using Prokka [24] with the same reference

files mentioned above and by forcing GenBank compliance.

MLST and cgMLST analysis:

For MLST and cgMLST, genome assemblies ere mapped against an MLST scheme

based on seven housekeeping genes [25] and the 1,343-locus cgMLST scheme [26],

using Ridom SeqSphere software (Münster, Germany) [27]. *C. hepaticus HV10* (accID

GCF-001687475.2)*, C. upsaliensis DSM 5365* (accID GCF-000620965.1), *C. lari*

RM2100 (accID GCF-000019205.1), *C. jejuni NCTC 11168* (accID GCF-000009085.1*),*

*C. jejuni doylei 269.97 (*accID GCA-000017485.1*),* aerotolerant *C. coli OR12 (*accID

*GCF-002024185.1), C. coli BIGS0008 (*accID *GCA-000314165.1), C. coli 76339 (*accID

GCA-000470055.1*),* and *C. coli H055260513* (accID GCA-001491555.1) were used as

reference genomes. Phylogenetic comparison of the 81 *Campylobacter* genome

sequences was performed using a neighbor-joining (NJ) tree based on a distance matrix

of the core genomes of all the strains. STs and clonal complexes (CCs) were determined

after automated allele submission to the PubMLST server for *Campylobacter* MLST

(https://pubmlst.org/campylobacter/). Additionally, to compare the isolates with

previously described *C. jejuni* and *C. coli* isolates worldwide, we performed a similar

phylogenetic analysis using the NCBI database including the isolates reported for the

whole Latin American region. Further annotations and visualizations were performed

using iTOL v.4 [28].

Virulome, resistome and comparative genomic analysis

A local BLAST database was constructed using the fasta files of all sequenced genomes

and the makeblastdb program from BLAST (NCBI) [29]. Subsequently, the database

was screened for the nucleotide sequence of known pathogenicity genes using BLASTn

(version 2.8.1) [29]. A threshold of 90% of the original sequence length and 90% identity

was used to select positive matches. A fasta file containing the nucleotide sequence of

Campylobacter pathogenicity genes screened in this work can be found at

https://doi.org/10.5281/zenodo.3925206. Mutations in *gyrA* and 23S rRNA were

searched as indicative of resistance to fluoroquinolones and macrolides, respectively.

The sequences of both genes were screened by BLAST in the local BLAST database

and sequences were aligned to the reference gene from strain Campylobacter jejuni

ATCC 33560 using ClustalX [30]. We then manually searched for mutation C257T in

gyrA and mutations A2058C and A2059G in the genes for 23S RNA. Antimicrobial

resistance genes were additionally screened in each *Campylobacter* genome by means

of the ABRicate pipeline [31], using the included Resfinder [32], CARD [33], ARG-

ANNOT [34] and NCBI ARRGD [35] databases. BLAST-based genome comparison

analysis and visualization of genetic clusters were performed using EasyFig [36].

Pangenome and plasmid screening analysis

Annotated genome sequences from all sequenced genomes were compared using Roary [37], setting a minimum percentage identity of 90% for BLASTp. Visualization of genes shared between genomes was performed using roary_plots.py. Additionally, the file gene_presence_absence.csv [https://github.com/sanger-pathogens/Roary/tree/master/contrib/roary_plots] produced by Roary was further analyzed using a spreadsheet program to find genes that are shared by subgroups of isolates. Using this information Venn diagrams were constructed using the tool available at http://bioinformatics.psb.ugent.be/webtools/Venn/. The screen for plasmids was performed on the FASTQ files of each sequencing reaction using PlasmidSeeker [38] based on the default bacterial plasmid database. Additional searches were performed using an inhouse database containing known plasmids from *C. coli* or *C. jejuni*. The fasta file for in-house databases can be found in https://doi.org/10.5281/zenodo.3925206. Pangenome BLAST analysis was performed using the GView server (https://server.gview.ca/) [39].

Results and Discussion

Phylogenetic analysis of clinical Campylobacter strains

cgMLST analysis showed high genetic diversity among the 69 *C. jejuni* strains (**Figure**

1A), as shown by the 14 CCs and 31 distinct STs identified. Almost 93% (n=64) of the *C.*

jejuni strains belonged to one of the CCs previously described for this species. Among

our strains, CC-21 was the most common (37.6%) and diverse (5 distinct STs) (**Figure**

1A, Table S1). This is consistent with other reports, which describe CC-21 as the largest

and most widely distributed clonal complex representing 18.9% of all *C. jejuni* isolates

submitted to the PubMLST database. ST-1359 was the most prevalent ST of CC-21

(48%) of all strains in this study (17.4%) (**Figure 1A and Table S1**). The high prevalence

of ST-1359 was unexpected, since it is infrequent within the PubMLST database (0.06%

by February 2020). However, ST-1359 has been recognized as a prominent ST in Israel

[40]. Other STs within CC-21 included ST-50 (8.7%) and ST-21 (1.45%), which were

predominant in reports from Israel, Belgium, and the United Kingdom [40–42] as well as

worldwide in PubMLST (4.1% and 4.3% respectively). ST-45, the second most prevalent

sequence type worldwide worldwide (4.2%), was not identified among our strains. In

addition to CC-21, we identified high prevalence rates of CC-48 (13%) and CC-52

(7.3%). ST-475 from CC-48 was the second most prevalent ST (11.6%) identified

(**Figure 1A**).

There are only two previous reports of MLST data of clinical *C. jejuni* strains from Chile

[18,22]. Both reports identified four major CCs and STs, including CC-21 (ST-50), CC-48

(ST-475), CC-257 (ST-257), and CC-353 (ST-353). Our study is consistent with these

reports, except for the high prevalence of ST-1359 (CC-21), which has not been

previously reported in Chile. Further epidemiological information is needed to determine

if the high prevalence of these four CCs is specific to Chile and to determine if the high

prevalence of ST-1359 strains in our report is an isolated finding or if it represents an

emergent ST in Chile.

C. coli is divided into three genetic clades [43,44]. Clade 1 is subdivided into two major

clonal complexes, CC-828 and CC-1150, and often isolated from farm animals and

human cases of gastroenteritis [44]. Clades 2 and 3 lack clonal complex substructures

and are mainly isolated from environmental sources [43,44]. cgMLST analysis of the 12

C. coli strains from our collection revealed that 83% (10/12) of the strains belonged to

clade 1, while 17% (2/12) belonged to clade 3 and none to clade 2. Most clade 1 strains

(8/10) belonged to CC-828 and included four previously described STs (828, 829, 1173,

and 1556) and one novel ST (10201). Only one strain from clade 1 belonged to the

usually common CC-1150 (strain CFSAN093253 from ST-10203), and another strain

could not be assigned to any clonal complex (strain CFSAN096322 from ST-10202).

Both clade 3 strains belonged to ST-10204. Overall, clade 1 strains from CC-828 were

most prevalent, which is consistent with the worldwide distribution of this species.

Interestingly, despite the low incidence of cases of gastroenteritis associated with *C. coli*,

in the 2-year period of our study (only 12 strains in total), we were able to identify strains

from the uncommon clade 3. Considering that these are the first strains of *C. coli*

described from Chile and considering the size of our study, further genomic

epidemiological studies are needed to fully grasp the diversity of the strains from this

country.

A global cgMLST analysis comparing our 81 *Campylobacter* genomes with other *C.*

jejuni and *C. coli* genomes available from the NCBI database was performed. The

strains of our collections from Central Chile showed great diversty and did not group with

any existing clusters (**Figure 1B)** (the interactive tree and metadata are available in

Microreact, see data summary).

Pangenome analysis of Campylobacter strains

To identify the genetic elements that are shared and distinct among the Chilean *C. jejuni*

and *C. coli* strains, we performed pangenome analysis of the 81 strains from both

species (**Figure 2**). The analysis identified a pangenome of 6609 genes for both species

and identified the core genes shared between *C. jejuni* and *C. coli* (532 genes) and the

core genome of each species (1300 genes for *C. jejuni* and 1207 genes for *C. coli*). In

addition, the analysis was able to identify the differences among the core and accessory

genomes of *C. coli* strains from clade 1 and clade 3. *C. coli* strains from clade 1

harbored 340 core genes absent from clade 3 core genes, which in turn, harbored 159

core genes not present in clade 1 core genes (**Figure 2**). The estimated core genome

for *C. jejuni* in our study (1300 genes) was close to the 1343 core genes identified in the

development of the cgMLST scheme of *C. jejuni* [25] and in agreement with recent

reports [45–47] (**Table S2**). In addition, a significant variation in the accessory genome

between *C. jejuni* and *C. coli* strains was identified, confirming the high genetic variability

of *Campylobacter*. Accessory genes that are common to a particular cluster seem to

contribute to modifications of carbohydrates and flagella as well as antibiotic resistance.

There is also a proportion of genomes harboring phage- and type IV and VI secretion

system-related genes (see below).

Genomic analysis of the resistome of Chilean clinical Campylobacter strains.

Although most *Campylobacter* infections are self-limiting and do not require antibiotic

treatment, the use of antibiotics is indicated for patients with persistent and severe

gastroenteritis, patients with extraintestinal infections, and patients who are

immunocompromised [3].

To determine the resistome of the *C. jejuni* and *C. coli* strains, we performed an *in silico*

analysis to identify genes associated with antimicrobial resistance (**Figure 3**). We

screened each *Campylobacter* genome by means of the ABRicate pipeline [31], using

the Resfinder, CARD, ARG-ANNOT and NCBI ARRGD databases, to identify

antimicrobial resistance genes. We additionally performed sequence alignments to

identify point mutations in *gyrA* and 16S rRNA genes that have been previously

described to confer antimicrobial resistance.

The *cmeABC* operon was present in all *C. jejuni* and *C. coli* strains analyzed. This

operon encodes a common multidrug efflux pump characterized in different

Campylobacter species, which mainly confers resistance to fluoroquinolones and

macrolides [48,49]. Additionally, the point mutation in the quinolone resistance-

determining region (QRDR) of the *gyrA* gene, leading to the T86I substitution in GyrA,

was identified in 54.3% of the strains. This mutation is the most frequently reported

mutation worldwide and confers high levels of resistance to fluoroquinolones [48,49].

The high percentage of *C. jejuni* strains with this mutation is consistent with recent

studies from Chile, reporting ciprofloxacin resistance in 30-60% of *C. jejuni* strains

[18,20–22]. While Porte *et al.* reported that most resistant strains were associated with

ST-48 and ST-353, in our study, there was no particular association of this mutation

specific STs. However, it is important to note that 100% of strains belonging to the most

frequent ST described in this work (ST-1359) harbored the mutation in the QRDR region.

, while most ST-475 isolates (the second most prevalent ST) not have this mutation.

Although fluoroquinolones are not first choice antimicrobial for human

campylobacteriosis, these high levels of resistance might be a consequence of their

broad use in animal husbandry. Fernandez and Perez-Perez concluded that the primary

source of transmission of resistant *Campylobacter* strains to humans in Latin America is

related to the consumption of animal-origin food and interactions with environmental

animal sources [50]. However, further studies are needed to obtain a better

understanding of the relevance of animal production practices in the dissemination of

antibiotic resistance genes among Chilean *Campylobacter* strains.

Regarding macrolide resistance, neither mutation of the 23S rRNA gene nor presence of

the *ermB* gene was detected, which is consistent with previous reports on Chilean *C.*

jejuni strains [18,22]. The low incidence of resistance against this group of antibiotics

(especially azithromycin and erythromycin) makes them the first choice for treatment of

Campylobacter infections [51]. However, further studies are necessary to analyze a

higher number of *C. coli* strains. This is particularly important given that according to

EFSA, macrolide resistance has been mostly detected in this species [52].

In terms of tetracycline resistance, 22.2% of the strains harbored the *tetO* gene, which is

the main genetic factor mediating tetracycline resistance in *Campylobacter* spp [53]. This

gene can be harbored on a plasmid or chromosomally encoded and is widespread

among *Campylobacter* strains worldwide [53]. Recently, the European Union (EU)

reported very high levels of tetracycline resistance in *C. jejuni* strains isolated from

humans (45.4%) and even higher levels of resistance in *C. coli* (68.3%). This antibiotic is

widely used, especially in the poultry industry worldwide [54], and might serve as an

important reservoir for resistant strains [55]. In fact, Lapierre *et al.* reported that 32.3% of

C. jejuni isolates from poultry in Chile were resistant to tetracycline [21]. Our data are

consistent with a recent report from Chile [21] that found that 22% of *C. jejuni* strains of

clinical origin were tetracycline resistant, which suggests that Chilean strains of

Campylobacter have lower levels of tetracycline resistance than strains from other

countries. This highlights the need for a permanent surveillance program to implement

control measures for tetracycline usage in animal production to avoid an increase in

tetracycline resistance levels in the human population. Finally, we also found a high

proportion of strains (79%) harboring the blaOXA-61 gene among *C. jejuni* and *C. coli*

strains. This genetic marker is associated with β-lactam resistance and has high

prevalence in *Campylobacter* isolates worldwide [56,57].

Presence and diversity of virulence gene content among Chilean Campylobacter

strains

The ability of *Campylobacter* to cause disease in humans is thought to be multifactorial,

as it lacks several of the key virulence factors used by other enteric bacterial pathogens

to cause disease (e.g., type III secretion systems). Nevertheless, there are several

genes and loci that are associated with the virulence and pathogenic potential of

Campylobacter, even though their role in the development of campylobacteriosis is not

fully understood [58]. Furthermore, most of our knowledge regarding the virulence

potential of *Campylobacter* strains comes from studies performed in *C. jejuni*; therefore,

information regarding the genetic determinants that contribute to the virulence of *C. coli*

strains is still lacking.

To determine the virulence gene content of the *C. jejuni* and *C. coli* strains, we first

assembled a list of 220 potential virulence genes, including genes described in the

Virulence Factor Database [59] and genes reported to contribute to the virulence of

Campylobacter in the literature [38,60–63]. These genes were grouped into five distinct

categories (adhesion and colonization, invasion, motility, secretion systems and toxins)

and were used to screen each of the 81 genomes using BLASTn**.** The distribution of the

major virulence factors described for *C. jejuni* and *C. coli* is shown in **Figure 4** (see

Figure S1 and **Table S3** for the full list).

Most of the described factors involved in adhesion and colonization (*cadF*, *racR*, *jlpA,*

pdlA, dnaJ, aphC) were prevalent among the *C. jejuni* strains and were not associated

with any particular STs (**Figure 4**, **Table S3**). This finding is consistent with other reports

indicating that the prevalence of these factors in clinical strains is very high [63] and

shows the relevance of adherence in the pathogenicity of *C. jejuni*. The genes

responsible for the production of the CDT toxin (*cdtA*, *cdtB* and *cdtC*) were also found in

most strains of *C. jejuni* and in every *C. coli* strain tested. The *cdtA* and *cdtB* genes were

not detected in the *C. jejuni* strain CFSAN093229. In contrast, the *ciaB* gene was

identified only in *C. jejuni* (in 98.7% of the strains) (**Figure 4**). The product of this gene is

involved in cecal colonization in chickens and in translocation into host cells in *C. jejuni*

[64,65]. The absence of this gene in *C. coli* might suggest that it does not play a crucial

role in the pathogenesis of this *Campylobacter* species. The major differences among *C.*

jejuni strains were observed for genes located at the *cps* locus (Cj1413c -Cj1442)

involved in capsule polysaccharide synthesis, the LOS locus (C8J1074-C8J1095)

involved in lipooligosaccharide synthesis, a small set of genes in the O-linked flagellin

glycosylation island (Cj1321-Cj1326) and the T6SS and T4SS gene clusters (**Figure 4**,

Figure S1 and Table S3).

Most virulence genes described for *C. jejuni* were absent in *C. coli* clades 1 and 3

strains, highlighting the need for a better understanding of how *C. coli* strains cause the

disease. However, *C. coli* strains carried most of the genes known to be associated with

invasiveness in *C. jejuni*, including the *iamb, flaC* and *ciaB* genes. Both *C. coli* clades

also harbored some genes associated with other steps of infection such as the *cdt* toxin

gene and most of the flagellar biosynthesis and chemiotaxis-related genes described for

C. jejuni (**Figure S1**). The high prevalence of the *iamb* gene in *C. coli* has also been

observed in other studies, but its relevance in pathogenicity remains unclear [66].

Distribution of the T4SS and T6SS gene clusters.

To date, there is only one T6SS cluster described in *Campylobacter* [60], and it has

been linked to a wide variety of phenotypes, including host cell adherence, cellular

invasion and resistance to bile salt, and oxidative stress [67–69]. In addition,

Campylobacter T6SS mutant strains exhibit colonization defects in both mouse [67] and

chicken [69] *in vivo* infection models. The *Campylobacter* T6SS gene cluster is often

encoded within the CjIE3 conjugative element in the chromosome of *C. jejuni* [61], but it

can also be found in plasmids [70,71]. Our analysis identified the T6SS gene cluster in 9

C. jejuni strains and in 1 *C. coli* strain from clade 1 (**Figure 4**). As shown in **Figure 5A**,

whenever present the genetic structure and the sequence identity of this cluster was

highly conserved among *C. jejuni* and *C. coli* strains. For strains with genomes that were

not closed, we were not able to confirm whether the *tssI* gene is located within the

genetic context of the of the T6SS cluster, since it was not assembled in the same contig

as the rest of the cluster [70].

Since most of our sequenced strains correspond to draft genome assemblies, we could

not determine whether the T6SSs were plasmid or chromosomally-encoded in most of

them. However, in two T6SS-positive strains with closed genomes (CFSAN093238 and

CFSAN093227), T6SS was inserted in the chromosomal CjIE3 element, while in strain

CFSAN093246, it was plasmid-encoded. Sequence-based analysis of this plasmid

showed a high degree of identity with plasmid pMTVDDCj13-2 (**Figure 6A**), which

encodes a complete T6SS gene cluster and was recently sequenced from the

MTVDSCj13 strain of *C. jejuni* [71].

No correlation was found between the T6SS gene cluster and a particular ST or clonal

complex. This is different from what has been described in other countries, such as

Israel, where most T6SS-positive strains belong to ST-1359 [40]. Although ST-1359 was

the most common ST described in this study, none of the strains carried genes encoding

T6SS.

To date, two distinct T4SSs have been described in *Campylobacter*. One of them is

encoded in the pVir plasmid of *C. jejuni* and is homologous to the T4SSs from

Helicobacter pylori [72]. The pVir T4SS has been linked to the virulence of *C. jejuni*.

T4SS mutants show reduced invasion capacity of INT407 cells and decreased diarrhea-

inducing ability in a ferret model [62,73]. The second T4SS is encoded in the pTet and

pCC31 family of plasmids of *C. jejuni* and *C. coli*. This distinct T4SS is involved in

bacterial conjugation and is not directly linked to the virulence of *C. jejuni* and *C. coli*

strains [74].

We identified a T4SS gene cluster homologous to this second T4SS in 12 strains of *C.*

jejuni and 1 strain of *C. coli* clade 1 (**Figure 4**). It harbored the 16 core genes and was

correctly assembled in individual contigs in 11 strains, which allowed the analysis of the

genetic context of the cluster. As shown in **Figure 5B**, in each strain, the T4SS cluster

had a conserved genetic structure. One of the main differences between the T4SS gene

clusters encoded in plasmids pTet and pCC31 is the sequence divergence of the *cpp33*

gene [74]. Only the *cp33* present in strain CFSAN096321 showed high identity with the

gene harbored in the pTet plasmid. T4SS gene clusters from every other *C. jejuni* and *C.*

coli strain possessed high identity to the *cp33* encoded in pCC31 plasmid.

As shown in **Figure 6B**, plasmids pCFSAN093226 and pCFSAN096297 share

significant similarity with both the pCC31 and pTET plasmids beyond the T4SS gene

cluster. For all other strains, BLASTn analysis of each of our draft genome assemblies

identified most of the plasmid genes found in pCFSAN093226, pCFSAN096297 and

pTet-like plasmids, suggesting that all the T4SS gene clusters identified in these strains

are encoded in plasmids as well.

Conclusions:

The genomes reported in the current study represent 96.4% of the sequenced Chilean

Campylobacter strains and 36.8% of the strains from the South American region. Taken

together, the results of this study provide valuable epidemiological data and relevant

information about the population structure, virulence-associated genes and AMR profiles

of an important number of clinical *Campylobacter* isolates. Further studies are needed to

determine the dynamics of transmission of pathogenic *Campylobacter* from animals to

humans and to monitor the emergence of new virulence and antimicrobial resistance

markers that might represent a threat to the human population. Finally, we expect that

these actionable public health data will support the design of strengthened surveillance

programs, which will be essential for the development of an effective integrated health

system in Chile.

AUTHOR CONTRIBUTIONS

Veronica Bravo: Conceptualization, Methodology, Formal analysis, Investigation,
Visualization, Data curation, Writing-Original Draft and review and editing.

Assaf Katz: Methodology, Formal analysis, Investigation, Visualization, Data Curation,
Resources, Writing-Original Draft and review and editing.

Carmen Varela: Resources, Writing - Review and editing

Lorena Porte: Resources, Writing - Review and editing.

Thomas Weitzel: Resources, Writing - Review and editing.

Narjol   Gonzalez-Escanola: Conceptualization,   Methodology,   Formal   analysis,
Resources, review and editing final manuscript

Carlos J Blondel: Conceptualization, Methodology, Formal analysis, Investigation,
Visualization, Resources, Data curation, Writing - Review and editing, Funding
acquisition, Supervision and Project administration.

CONFLICT OF INTERESTS

The authors declare that there are no conflicts of interest.

FUNDING INFORMATION

This work was funded by the Howard Hughes Medical Institute (HHMI)-Gulbenkian
International Research Scholar Grant #55008749, FONDECYT Grant 1201805 (ANID)
and REDI170269 (ANID) from CJB. VB is supported by DICYT grant 022001BZ
(USACH). AK is supported by FONDECYT Grant 1191074 (ANID). NJG is supported by
MCMi Challenge Grants program proposal number 2018-646 and the FDA Foods
Program Intramural Funds (FDA employees).

REFERENCES

[1] Bravo V, Varela C, Porte L, Weitzel T, Kastanis GJ, Balkey M, et al. Draft Whole-Genome Sequences of 51 Campylobacter jejuni and 12 Campylobacter coli Clinical Isolates from Chile. Microbiol Resour Announc 2020;9:MRA.00072-20, e00072-20. https://doi.org/10.1128/MRA.00072-20.

[2] Bravo V, Porte L, Weitzel T, Varela C, Blondel C, Gonzalez-Escalona N. Complete genome sequences of 17 clinical Campylobacter jejuni strains from Chile. Microbiology Resource Announcements n.d.:9:e00535-20. https://doi.org/10.1128/MRA.00535-20.

[3] Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, Man SM. Global Epidemiology of Campylobacter Infection. Clin Microbiol Rev 2015;28:687–720. https://doi.org/10.1128/CMR.00006-15.

[4] World Health Organization, Food and Agriculture Organization of the United Nations, World Organisation for Animal Health. The global view of campylobacteriosis: report of an expert consultation, Utrecht, Netherlands, 9-11 July 2012. Geneva: World Health Organization; 2013.

[5] EFSA. The European Union One Health 2018 Zoonoses Report. EFSA Journal 2019;17:e05926. https://doi.org/10.2903/j.efsa.2019.5926.

[6] Eurosurveillance editorial team. European Union Summary Report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food 2012 published. Euro Surveill 2014;19:20748. https://doi.org/10.2807/ese.19.12.20748-en.

[7] World Health Organization E. Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics 2017:7.

[8] EFSA. Scientific Opinion on Campylobacter in broiler meat production: control options and performance objectives and/or targets at different stages of the food chain. EFSA Journal 2011;9:2105. https://doi.org/10.2903/j.efsa.2011.2105.

[9] Acheson D, Allos BM. Campylobacter jejuni Infections: Update on Emerging Issues and Trends. Clinical Infectious Diseases 2001;32:1201–6. https://doi.org/10.1086/319760.

[10] Wijdicks EFM, Klein CJ. Guillain-Barré Syndrome. Mayo Clinic Proceedings 2017;92:467–79. https://doi.org/10.1016/j.mayocp.2016.12.002.

[11] Scallan Walter EJ, Crim SM, Bruce BB, Griffin PM. Incidence of *Campylobacter* -Associated Guillain-Barré Syndrome Estimated from Health Insurance Data. Foodborne Pathogens and Disease 2020;17:23–8. https://doi.org/10.1089/fpd.2019.2652.

[12] Yang Y, Feye KM, Shi Z, Pavlidis HO, Kogut M, J. Ashworth A, et al. A Historical Review on Antibiotic Resistance of Foodborne Campylobacter. Front Microbiol 2019;10:1509. https://doi.org/10.3389/fmicb.2019.01509.

[13] Sproston EL, Wimalarathna HML, Sheppard SK. Trends in fluoroquinolone resistance in Campylobacter. Microbial Genomics 2018;4. https://doi.org/10.1099/mgen.0.000198.

[14] Fernández H. [Campylobacter and campylobacteriosis: a view from South America]. Rev Peru Med Exp Salud Publica 2011;28:121–7. https://doi.org/10.1590/s1726-46342011000100019.

[15] Instituto de Salud Pública de Chile. Vigilancia de laboratorio de Campylobacter spp. Chile, 2005–2013. Boletín ISP 2014;4:1–17.

[16] Porte L, Varela C, Haecker T, Morales S, Weitzel T. Impact of changing from staining to culture techniques on detection rates of Campylobacter spp. in routine

stool samples in Chile. BMC Infect Dis 2016;16:196.
https://doi.org/10.1186/s12879-016-1546-7.
[17] Lorena Porte, Weitzel T, Barbe M, Armijo Y, Vollrath V. Routine molecular diagnosis
of enteropathogens by the FilmArray gastrointestinal panel in a microbiological
laboratory in Santiago, Chile. 28th European Congress of Clinical Microbiology and
Infectious Diseases 2018.
[18] Collado L, Muñoz N, Porte L, Ochoa S, Varela C, Muñoz I. Genetic diversity and
clonal characteristics of ciprofloxacin-resistant Campylobacter jejuni isolated from
Chilean patients with gastroenteritis. Infection, Genetics and Evolution
2018;58:290–3. https://doi.org/10.1016/j.meegid.2017.12.026.
[19] Rivera F N, Bustos B R, Montenegro H S, Sandoval M M, Castillo N J, Fernández J
H, et al. [Genotyping and antibacterial resistance of Campylobacter spp strains
isolated in children and in free range poultry]. Rev Chilena Infectol 2011;28:555–62.
https://doi.org//S0716-10182011000700008.
[20] González-Hein G, Cordero N, García P, Figueroa G. [Molecular analysis of
fluoroquinolones and macrolides resistance in Campylobacter jejuni isolates from
humans, bovine and chicken meat]. Rev Chilena Infectol 2013;30:135–9.
https://doi.org/10.4067/S0716-10182013000200003.
[21] Lapierre L, Gatica MA, Riquelme V, Vergara C, Yañez JM, San Martín B, et al.
Characterization of Antimicrobial Susceptibility and Its Association with Virulence
Genes Related to Adherence, Invasion, and Cytotoxicity in *Campylobacter jejuni*
and *Campylobacter coli* Isolates from Animals, Meat, and Humans. Microbial Drug
Resistance 2016;22:432–44. https://doi.org/10.1089/mdr.2015.0055.
[22] Levican A, Ramos-Tapia I, Briceño I, Guerra F, Mena B, Varela C, et al. Genomic
Analysis of Chilean Strains of *Campylobacter jejuni* from Human Faeces. BioMed
Research International 2019;2019:1–12. https://doi.org/10.1155/2019/1902732.
[23] Darling AE, Mau B, Perna NT. progressiveMauve: Multiple Genome Alignment with
Gene Gain, Loss and Rearrangement. PLoS ONE 2010;5:e11147.
https://doi.org/10.1371/journal.pone.0011147.
[24] Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics
2014;30:2068–9. https://doi.org/10.1093/bioinformatics/btu153.
[25] Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics:
BIGSdb software, the PubMLST.org website and their applications. Wellcome
Open Res 2018;3:124. https://doi.org/10.12688/wellcomeopenres.14826.1.
[26] Cody AJ, Bray JE, Jolley KA, McCarthy ND, Maiden MCJ. Core Genome Multilocus
Sequence Typing Scheme for Stable, Comparative Analyses of Campylobacter
jejuni and C. coli Human Disease Isolates. J Clin Microbiol 2017;55:2086–97.
https://doi.org/10.1128/JCM.00080-17.
[27] Jünemann S, Sedlazeck FJ, Prior K, Albersmeier A, John U, Kalinowski J, et al.
Updating benchtop sequencing performance comparison. Nat Biotechnol
2013;31:294–6. https://doi.org/10.1038/nbt.2522.
[28] Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. Nucleic Acids Research 2019;47:W256–9.
https://doi.org/10.1093/nar/gkz239.
[29] Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search
tool. J Mol Biol 1990;215:403–10. https://doi.org/10.1016/S0022-2836(05)80360-2.
[30] Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, et
al. Clustal W and Clustal X version 2.0. Bioinformatics 2007;23:2947–8.
https://doi.org/10.1093/bioinformatics/btm404.
[31] Seemann T. tseemann/abricate. 2020.

[32] Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al.
Identification of acquired antimicrobial resistance genes. Journal of Antimicrobial
Chemotherapy 2012;67:2640–4. https://doi.org/10.1093/jac/dks261.
[33] Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, et al. CARD 2017:
expansion and model-centric curation of the comprehensive antibiotic resistance
database. Nucleic Acids Res 2017;45:D566–73.
https://doi.org/10.1093/nar/gkw1004.
[34] Gupta SK, Padmanabhan BR, Diene SM, Lopez-Rojas R, Kempf M, Landraud L, et
al. ARG-ANNOT, a New Bioinformatic Tool To Discover Antibiotic Resistance
Genes in Bacterial Genomes. Antimicrob Agents Chemother 2014;58:212–20.
https://doi.org/10.1128/AAC.01310-13.
[35] Feldgarden M, Brover V, Haft DH, Prasad AB, Slotta DJ, Tolstoy I, et al. Validating
the AMRFinder Tool and Resistance Gene Database by Using Antimicrobial
Resistance Genotype-Phenotype Correlations in a Collection of Isolates.
Antimicrob Agents Chemother 2019;63:e00483-19, /aac/63/11/AAC.00483-
19.atom. https://doi.org/10.1128/AAC.00483-19.
[36] Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer.
Bioinformatics 2011;27:1009–10. https://doi.org/10.1093/bioinformatics/btr039.
[37] Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, et al. Roary:
rapid large-scale prokaryote pan genome analysis. Bioinformatics 2015;31:3691–3.
https://doi.org/10.1093/bioinformatics/btv421.
[38] Roosaare M, Puustusmaa M, Möls M, Vaher M, Remm M. PlasmidSeeker:
identification of known plasmids from bacterial whole genome sequencing reads.
PeerJ 2018;6:e4588. https://doi.org/10.7717/peerj.4588.
[39] Stothard P, Grant JR, Van Domselaar G. Visualizing and comparing circular
genomes using the CGView family of tools. Briefings in Bioinformatics
2019;20:1576–82. https://doi.org/10.1093/bib/bbx081.
[40] Rokney A, Valinsky L, Moran-Gilad J, Vranckx K, Agmon V, Weinberger M. Genomic
Epidemiology of Campylobacter jejuni Transmission in Israel. Front Microbiol
2018;9:2432. https://doi.org/10.3389/fmicb.2018.02432.
[41] Elhadidy M, Arguello H, Álvarez-Ordóñez A, Miller WG, Duarte A, Martiny D, et al.
Orthogonal typing methods identify genetic diversity among Belgian Campylobacter
jejuni strains isolated over a decade from poultry and cases of sporadic human
illness. Int J Food Microbiol 2018;275:66–75.
https://doi.org/10.1016/j.ijfoodmicro.2018.04.004.
[42] Cody AJ, McCarthy NM, Wimalarathna HL, Colles FM, Clark L, Bowler ICJW, et al. A
Longitudinal 6-Year Study of the Molecular Epidemiology of Clinical Campylobacter
Isolates in Oxfordshire, United Kingdom. Journal of Clinical Microbiology
2012;50:3193–201. https://doi.org/10.1128/JCM.01086-12.
[43] Sheppard SK, McCarthy ND, Falush D, Maiden MCJ. Convergence of
Campylobacter Species: Implications for Bacterial Evolution. Science
2008;320:237–9. https://doi.org/10.1126/science.1155532.
[44] Sheppard SK, Didelot X, Jolley KA, Darling AE, Pascoe B, Meric G, et al.
Progressive genome wide introgression in agricultural *Campylobacter coli*. Mol
Ecol 2013;22:1051–64. https://doi.org/10.1111/mec.12162.
[45] Méric G, Yahara K, Mageiros L, Pascoe B, Maiden MCJ, Jolley KA, et al. A
Reference Pan-Genome Approach to Comparative Bacterial Genomics:
Identification of Novel Epidemiological Markers in Pathogenic Campylobacter.
PLoS ONE 2014;9:e92798. https://doi.org/10.1371/journal.pone.0092798.
[46] Biggs PJ, Fearnhead P, Hotter G, Mohan V, Collins-Emerson J, Kwan E, et al.
Whole-Genome Comparison of Two Campylobacter jejuni Isolates of the Same

Sequence Type Reveals Multiple Loci of Different Ancestral Lineage. PLoS ONE 2011;6:e27121. https://doi.org/10.1371/journal.pone.0027121.

[47] Friis C, Wassenaar TM, Javed MA, Snipen L, Lagesen K, Hallin PF, et al. Genomic Characterization of Campylobacter jejuni Strain M1. PLoS ONE 2010;5:e12253. https://doi.org/10.1371/journal.pone.0012253.

[48] Luangtongkum T, Jeon B, Han J, Plummer P, Logue CM, Zhang Q. Antibiotic resistance in *Campylobacter*□: emergence, transmission and persistence. Future Microbiology 2009;4:189–200. https://doi.org/10.2217/17460913.4.2.189.

[49] Shen Z, Wang Y, Zhang Q, Shen J. Antimicrobial Resistance in Campylobacter spp. In: Schwarz, Cavaco, Shen, editors. Antimicrobial Resistance in Bacteria from Livestock and Companion Animals, American Society of Microbiology; 2018, p. 317–30. https://doi.org/10.1128/microbiolspec.ARBA-0013-2017.

[50] Fernández H, Pérez-Pérez G. Campylobacter: fluoroquinolone resistance in Latin-American countries. Archivos de Medicina Veterinaria 2016;48:255–9. https://doi.org/10.4067/S0301-732X2016000300002.

[51] Bolinger H, Kathariou S. The Current State of Macrolide Resistance in Campylobacter spp.: Trends and Impacts of Resistance Mechanisms. Appl Environ Microbiol 2017;83:e00416-17, e00416-17. https://doi.org/10.1128/AEM.00416-17.

[52] Engberg J, Aarestrup FM, Taylor DE, Gerner-Smidt P, Nachamkin I. Quinolone and macrolide resistance in Campylobacter jejuni and C. coli: resistance mechanisms and trends in human isolates. Emerging Infect Dis 2001;7:24–34. https://doi.org/10.3201/eid0701.010104.

[53] Connell SR, Trieber CA, Dinos GP, Einfeldt E, Taylor DE, Nierhaus KH. Mechanism of Tet(O)-mediated tetracycline resistance. EMBO J 2003;22:945–53. https://doi.org/10.1093/emboj/cdg093.

[54] Van Boeckel TP, Brower C, Gilbert M, Grenfell BT, Levin SA, Robinson TP, et al. Global trends in antimicrobial use in food animals. Proc Natl Acad Sci USA 2015;112:5649–54. https://doi.org/10.1073/pnas.1503141112.

[55] Maron D, Smith TJ, Nachman KE. Restrictions on antimicrobial use in food animal production: an international regulatory and economic survey. Global Health 2013;9:48. https://doi.org/10.1186/1744-8603-9-48.

[56] Alfredson DA, Korolik V. Isolation and expression of a novel molecular class D beta-lactamase, OXA-61, from Campylobacter jejuni. Antimicrob Agents Chemother 2005;49:2515–8. https://doi.org/10.1128/AAC.49.6.2515-2518.2005.

[57] Marotta F, Garofolo G, di Marcantonio L, Di Serafino G, Neri D, Romantini R, et al. Antimicrobial resistance genotypes and phenotypes of Campylobacter jejuni isolated in Italy from humans, birds from wild and urban habitats, and poultry. PLoS ONE 2019;14:e0223804. https://doi.org/10.1371/journal.pone.0223804.

[58] Hansson I, Sandberg M, Habib I, Lowman R, Engvall EO. Knowledge gaps in control of *Campylobacter* for prevention of campylobacteriosis. Transbound Emerg Dis 2018;65:30–48. https://doi.org/10.1111/tbed.12870.

[59] Liu B, Zheng D, Jin Q, Chen L, Yang J. VFDB 2019: a comparative pathogenomic platform with an interactive web interface. Nucleic Acids Research 2019;47:D687–92. https://doi.org/10.1093/nar/gky1080.

[60] Redondo N, Carroll A, McNamara E. Molecular characterization of Campylobacter causing human clinical infection using whole-genome sequencing: Virulence, antimicrobial resistance and phylogeny in Ireland. PLoS ONE 2019;14:e0219088. https://doi.org/10.1371/journal.pone.0219088.

[61] Agnetti J, Seth-Smith HMB, Ursich S, Reist J, Basler M, Nickel C, et al. Clinical impact of the type VI secretion system on virulence of Campylobacter species

during infection. BMC Infect Dis 2019;19:237. https://doi.org/10.1186/s12879-019-
3858-x.
[62] Bacon DJ, Alm RA, Burr DH, Hu L, Kopecko DJ, Ewing CP, et al. Involvement of a
Plasmid in Virulence of Campylobacter jejuni 81-176. Infect Immun 2000;68:4384–
90. https://doi.org/10.1128/IAI.68.8.4384-4390.2000.
[63] Dasti JI, Tareen AM, Lugert R, Zautner AE, Groß U. Campylobacter jejuni: A brief
overview on pathogenicity-associated factors and disease-mediating mechanisms.
International Journal of Medical Microbiology 2010;300:205–11.
https://doi.org/10.1016/j.ijmm.2009.07.002.
[64] Konkel ME, Kim BJ, Rivera-Amill V, Garvis SG. Bacterial secreted proteins are
required for the internaliztion of Campylobacter jejuni into cultured mammalian
cells. Mol Microbiol 1999;32:691–701. https://doi.org/10.1046/j.1365-
2958.1999.01376.x.
[65] Novik V, Hofreuter D, Galán JE. Identification of Campylobacter jejuni genes
involved in its interaction with epithelial cells. Infect Immun 2010;78:3540–53.
https://doi.org/10.1128/IAI.00109-10.
[66] Ghorbanalizadgan M, Bakhshi B, Kazemnejad Lili A, Najar-Peerayeh S, Nikmanesh
B. A molecular survey of Campylobacter jejuni and Campylobacter coli virulence
and diversity. Iran Biomed J 2014;18:158–64. https://doi.org/10.6091/ibj.1359.2014.
[67] Lertpiriyapong K, Gamazon ER, Feng Y, Park DS, Pang J, Botka G, et al.
Campylobacter jejuni Type VI Secretion System: Roles in Adaptation to
Deoxycholic Acid, Host Cell Adherence, Invasion, and In Vivo Colonization. PLoS
ONE 2012;7:e42842. https://doi.org/10.1371/journal.pone.0042842.
[68] Bleumink-Pluym NMC, van Alphen LB, Bouwman LI, Wösten MMSM, van Putten
JPM. Identification of a Functional Type VI Secretion System in Campylobacter
jejuni Conferring Capsule Polysaccharide Sensitive Cytotoxicity. PLoS Pathog
2013;9:e1003393. https://doi.org/10.1371/journal.ppat.1003393.
[69] Liaw J, Hong G, Davies C, Elmi A, Sima F, Stratakos A, et al. The Campylobacter
jejuni Type VI Secretion System Enhances the Oxidative Stress Response and
Host Colonization. Front Microbiol 2019;10:2864.
https://doi.org/10.3389/fmicb.2019.02864.
[70] Ghatak S, He Y, Reed S, Strobaugh T, Irwin P. Whole genome sequencing and
analysis of Campylobacter coli YH502 from retail chicken reveals a plasmid-borne
type VI secretion system. Genom Data 2017;11:128–31.
https://doi.org/10.1016/j.gdata.2017.02.005.
[71] Taveirne ME, Dunham DT, Perault A, Beauchamp JM, Huynh S, Parker CT, et al.
Complete Annotated Genome Sequences of Three Campylobacter jejuni Strains
Isolated from Naturally Colonized Farm-Raised Chickens. Genome Announc
2017;5:e01407-16, e01407-16. https://doi.org/10.1128/genomeA.01407-16.
[72] Kersulyte D, Velapatiño B, Mukhopadhyay AK, Cahuayme L, Bussalleu A, Combe J,
et al. Cluster of type IV secretion genes in Helicobacter pylori's plasticity zone. J
Bacteriol 2003;185:3764–72. https://doi.org/10.1128/jb.185.13.3764-3772.2003.
[73] Bacon DJ, Alm RA, Hu L, Hickey TE, Ewing CP, Batchelor RA, et al. DNA sequence
and mutational analyses of the pVir plasmid of Campylobacter jejuni 81-176. Infect
Immun 2002;70:6242–50. https://doi.org/10.1128/iai.70.11.6242-6250.2002.
[74] Batchelor RA, Pearson BM, Friis LM, Guerry P, Wells JM. Nucleotide sequences
and comparison of two large conjugative plasmids from different Campylobacter
species. Microbiology (Reading, Engl) 2004;150:3507–17.
https://doi.org/10.1099/mic.0.27112-0.
763

FIGURE LEGENDS

Figure 1. A) Phylogenetic analysis of *C. jejuni* and *C. coli* strains from Central Chile. The phylogenetic tree is based on cgMLST analyses performed with RIDOM SeqSphere[+] and visualized by iTOL. Sequence types (STs) are shown in colored boxes for each of the *Campylobacter* strains, and the tree branches have been color coded to highlight the *C. jejuni* and *C. coli* strains from clades I, II and III. **B)** Global cgMLST analysis and distribution of clonal complexes of clinical *C. jejuni* and *C. coli* strains. Strains from this study are highlighted in red. cgMLST analysis performed by SeqSphere+ and visualized by iTOL.

Figure 2. Visualization of the pangenome of the 69 *C. jejuni* and 12 *C. coli* strains from Chile. Pangenome analysis was performed using Roary. The core and accessory genomes are highlighted with red and black brackets, respectively. The cgMLST phylogenetic tree of Figure 1A is shown on the left. A Venn diagram highlighting the number of core genes present in *C. jejuni* and *C. coli* strains from clades I and III is shown on the right.

Figure 3. Distribution of antimicrobial resistance genes in *C. jejuni* and *C. coli* strains from Chile. Binary heatmaps show the presence and absence of antimicrobial resistance genes. Colored cells represent the presence of genes, and blank cells represent the absence of genes. The phylogeny of *C. jejuni* and *C. coli* strains is the same as that in Figure 1. Strain names are color coded to highlight phylogenetic groups: *C. jejuni* strains (light green), *C. coli* clade I strains (orange) and *C. coli* clade III strains (magenta).

Figure 4. Distribution of virulence-related genes in *C. jejuni and C. coli* strains. Binary heatmaps show the presence and absence of virulence genes linked to bacterial adhesion (red), invasion (blue), toxins (purple) and specialized protein secretion systems (green). Colored cells represent the presence of genes, and blank cells represent the absence of genes. The phylogeny of *C. jejuni* and *C. coli* strains is the same as that in Figure 1. Strain names are color coded to highlight *C. jejuni* strains (green), *C. coli* clade I strains (orange) and *C. coli* clade III strains (purple).

Figure 5. Comparative genomic analysis of the T6SS and T4SS gene clusters identified in clinical *C. jejuni* and *C. coli* strains. **A)** Genetic organization of the T6SS gene clusters of *C. jejuni* and *C. coli* strains (bold) compared to the cluster of *C. jejuni* strain 108. The genes *tagH*, *tssD*, *tssB* and *tssC* are shown in color. **B)** Genetic organization of the T4SS gene clusters identified in clinical *C. jejuni* strains in comparison to the T4SS gene clusters of the plasmids pTet and pCC31. The *cp33* gene is highlighted in orange. BLASTn alignments were performed and visualized using EasyFig.

Figure 6. Comparative genomic analysis of plasmids encoding T6SS **(A)** and T4SS **(B)** gene clusters in clinical *C. jejuni* strains. Pangenome BLAST analysis of plasmids was performed using the Gview server. T6SS and T4SS gene clusters are highlighted in red brackets.

Figure S1. Distribution of the presence (red) and absence (blue) of virulence-related genes in *C. jejuni* and *C. coli* strains from the study.

Campylobacter species

- ■ C. jejuni
- ■ C. coli Clade I
- ■ C. coli clade II
- ■ C. coli Clade III

STs

829	8941`	3720	38	10204	607	43	1203	51	10197	10201
8464	3573	4053	475	10196	1510	21	1173	52	10198	10203
1556	137	6091	10192	1845	464	883	1359	508	10200	
828	257	8938	10193	468	353	222	50	3997	10202	

Tree scale: 0.1

Tree scale: 0.1

putative lipase

hypothetical protein

hypothetical protein

putative TraG protein

hypothetical protein

hypothetical protein

hypothetical protein

hypothetical protein

hypothetical protein

hypothetical protein

putative nuclease

putative lipase

putative nuclease

hypothetical protein

hypothetical protein

hypothetical protein

VasC protein

putative IcmF protein

putative IcmF protein

DotU domain protein

hypothetical protein

Type VI secretion system protein TssF

hypothetical protein

hypothetical protein

putative nuclease

hypothetical protein

putative phage integrase

T6SS gene cluster

0 kbp
5 kbp
10 kbp
15 kbp
20 kbp
25 kbp
30 kbp
35 kbp
40 kbp
45 kbp
50 kbp
55 kbp
60 kbp
65 kbp
70 kbp
75 kbp
80 kbp

■ pCFSAN093246
■ pMTVDDCj13-2
■ Pangenome

tetO
cpp3
repA
cpp7
cpp9
cpp11
cpp13
cpp14
cpp16
cpp50
cpp49
cpp49
cpp49
cpp47
cpp46
cpp45
cmgD4
cmgD4
cmgD4
cmgB11
cmgB10
cmgB9
cmgB8
cmgB6
ssb1
cmgB3/4
vapD
cpp26
cpp23
cpp22
cpp21
cpp18
cpp17
cpp17
cpp16

0 kbp
5 kbp
10 kbp
15 kbp
20 kbp
25 kbp
30 kbp
35 kbp
40 kbp
45 kbp

T4SS gene cluster

■ pCFSAN093226
■ pCFSAN096297
■ pCC31
■ pTet
■ Pangenome