Genomic analysis of the diversity, antimicrobial resistance and virulence

potential of *Campylobacter jejuni* and *Campylobacter coli* strains from Chile

Veronica Bravo[a,b], Assaf Katz[c], Lorena Porte[d], Thomas Weitzel[d,e], Carmen Varela[d], Narjol

Gonzalez-Escalona[f] and Carlos J. Blondel[b,g]*

[a] Unidad de Microbiología, Escuela de Medicina, Facultad de Ciencias Médicas, Universidad

de Santiago de Chile, Santiago, Chile.

[b] Instituto de Ciencias Biomédicas, Facultad de Medicina y Facultad de Ciencias de la Vida,

Universidad Andrés Bello, 8320000, Santiago, Chile.

[c] Programa de Biología Celular y Molecular, ICBM, Facultad de Medicina, Universidad de

Chile, Santiago, 8380453, Chile.

[d] Laboratorio Clínico, Clínica Alemana de Santiago, Facultad de Medicina Clínica Alemana,

Universidad del Desarrollo, Santiago, Chile.

[e] Instituto de Ciencias e Innovación en Medicina (ICIM), Facultad de Medicina Clínica

Alemana, Universidad del Desarrollo, Santiago, Chile.

[f] Center for Food Safety and Applied Nutrition, Office of Regulatory Science, Division of

Microbiology, U.S. Food and Drug Administration, College Park, Maryland, USA.

[g] Instituto de Ciencias Biomédicas, Facultad de Ciencias de la Salud, Universidad Autónoma

de Chile, Santiago, Chile.

*Corresponding author. carlos.blondel@unab.cl

Mailing address, Instituto de Ciencias Biomédicas, Universidad Andrés Bello. Echaurren

183, Santiago, 8320000, Chile.

ABSTRACT

Campylobacter jejuni and *Campylobacter coli* are the leading cause of human gastroenteritis

in the industrialized world and an emerging threat in developing countries. The incidence of

campylobacteriosis in South America is greatly underestimated, mostly due to the lack of

adequate diagnostic methods. Accordingly, there is limited genomic and epidemiological

data from this region. In the present study, we performed a genome-wide analysis of the

genetic diversity, virulence, and antimicrobial resistance of the largest collection of clinical

C. jejuni and *C. coli* strains from Chile available to date (n=81), collected in 2017-2019 in

Santiago, Chile. Collection which accounts for over a third of the genome sequences

available of clinical strains from South America. cgMLST analysis identified a high genetic

diversity as well as 13 novel STs and alleles in both *C. jejuni* and *C. coli*. Pangenome and

virulome analyses showed a differential distribution of virulence factors, including both

plasmid and chromosomally encoded T6SSs and T4SSs. Resistome analysis predicted

widespread resistance to fluoroquinolones, but low rates of erythromycin resistance. This

study provides valuable genomic and epidemiological data and highlights the need for further

genomic epidemiology studies in Chile and other South American countries to better

understand molecular epidemiology and antimicrobial resistance of this emerging intestinal

pathogen.

Key words: *Campylobacter jejuni*, *Campylobacter coli*, cgMLST, comparative genomics,

foodborne pathogen.

IMPORTANCE

Campylobacter is the leading cause of bacterial gastroenteritis worldwide and an emerging and neglected pathogen in South America. In this study, we performed an in-depth analysis of the genome sequences of 69 *C. jejuni* and 12 *C. coli* clinical strains isolated from Chile, which account for over a third of the sequences from clinical strains available from South America. We identified a high genetic diversity among *C. jejuni* strains and the unexpected identification of clade 3 *C. coli* strains, which are infrequently isolated from humans in other regions of the world. Most strains harbored the virulence factors described for *Campylobacter* and while ~40% of strains harbored mutation in the *gyrA* gene described to confer fluoroquinolone resistance, very few strains encoded the determinants linked to macrolide resistance, currently used for the treatment of campylobacteriosis. Our study contributes to our knowledge of this important foodborne pathogen providing valuable data from South America.

INTRODUCTION

Campylobacteriosis caused by *Campylobacter jejuni* and *Campylobacter coli* has emerged as an important public health problem and is the most common bacterial cause of human gastroenteritis worldwide [1,2]. It is usually associated with a self-limiting illness characterized by diarrhea, nausea, abdominal cramps and bloody stools [1]. Nevertheless, extraintestinal manifestations might cause long-term complications such Miller-Fisher or Guillain-Barré syndrome [1]. Despite the self-limiting course, certain populations such as children under 5 years of age, elderly patients, and immunocompromised patients might suffer severe infections and require antimicrobial treatment [3]. In this context, the World Health Organization (WHO) has included *Campylobacter* as a high priority pathogen due to the worldwide emergence of strains with high level fluoroquinolone resistance [4].

In South America, there is limited data available on the prevalence of *C. jejuni* and *C. coli* in comparison to other enteric bacterial pathogens [1,5,6]. In Chile, the National Laboratory Surveillance Program of the Public Health Institute has reported an incidence rate of campylobacteriosis of 0.1 to 0.6 cases per 100.000 persons [7]. This is low compared to high-income countries, where incidence rates can reach two orders of magnitude higher [1]. Nevertheless, recent reports suggest that the burden of campylobacteriosis in Chile has been greatly underestimated due to suboptimal diagnostic protocols [7]. Indeed, a recent report suggests that *Campylobacter* spp. is emerging as the second most prevalent bacterial cause of human gastroenteritis in Chile [8].

Only few reports have analyzed the genetic diversity and antimicrobial resistance profiles of clinical human strains of *Campylobacter* in Chile [9–13]. Furthermore, these

studies have either used a limited number of strains [13] or lacked the resolution provided by

whole-genome sequence-based typing methods [9–12]. Therefore, larger genomic

epidemiology studies are needed in Chile and more broadly in South America.

We recently made available the whole-genome sequences of 69 *C. jejuni* and 12 *C.*

coli human clinical strains isolated from patients with enteric infection attending Clínica

Alemana in Santiago, Chile, during a 2-year period (2017-2019) [14,15]. This represents the

largest genome collection of clinical *Campylobacter* strains from Chile and one of the few

collections available from South America. In the present study, we performed a genome-

wide analysis of the genetic diversity, virulence potential, and antimicrobial resistance

profiles of these strains. Importantly, we identified a high genetic diversity of both *C. jejuni*

and *C. coli* strains, including 13 novel sequence types (STs) and alleles. Resistome analysis

predicted a widespread resistance to fluoroquinolones, but decreased resistance to

tetracycline in comparison to the reported resistance from other regions of the world, in

addition to the differential distribution of virulence factors, including plasmids and

chromosomally encoded T6SS and T4SS.

MATERIALS AND METHODS

Genome data set, sequencing and reannotation

The genome sequence dataset analyzed in this study is listed in **Table S1** and includes our recently published collection of human clinical strains of *C. jejuni* and *C. coli* from Chile [14,15]. We additionally included the draft whole-genome sequence of *C. jejuni* strain CFSAN096303, from the same collection of isolates. CFSAN096303 was sequenced as described in [14], using the NextSeq sequencer (Illumina) obtaining an estimated genome average coverage of 90X. The genome was assembled using the CLC Genomics Workbench v9.5.2 (Qiagen) following the previously used pipeline [14] and was deposited under accession number WXZQ01000000. Fasta files of all draft genomes were reordered to the reference genomes of the *C. coli* aerotolerant strain OR12 (accID GCF_002024185.1) and *C. jejuni* strain NCTC 11168 (accID GCF_000009085.1) using the Mauve Contig Mover (MCM) from the Mauve package [16]. The ordered genomes were reannotated using Prokka [17] with the same reference files mentioned above and by forcing GenBank compliance. Draft genome sequence reannotations and sequence fasta files are available as a supplementary dataset on Zenodo https://doi.org/10.5281/zenodo.3925206.

MLST and core genome MLST (cgMLST) analysis:

Genome assemblies were mapped against an MLST scheme based on seven housekeeping genes [18] and the 1,343-locus cgMLST scheme [19], using Ridom SeqSphere software (Münster, Germany) [20]. *C. hepaticus HV10* (accID GCF-001687475.2)*, C. upsaliensis DSM 5365* (accID GCF-000620965.1), *C. lari RM2100* (accID GCF-000019205.1), *C. jejuni NCTC 11168* (accID GCF-000009085.1)*, C. jejuni doylei 269.97* (accID GCA-000017485.1)*,* aerotolerant *C. coli OR12* (accID GCF-002024185.1)*, C. coli BIGS0008*

(accID GCA-000314165.1)*, C. coli 76339* (accID GCA-000470055.1)*, and *C. coli*

H055260513 (accID GCA-001491555.1) were used as reference genomes. Phylogenetic

comparison of the 81 *Campylobacter* genome sequences was performed using a neighbor-

joining tree based on a distance matrix of the core genomes of all strains. STs and CCs were

determined after automated allele submission to the *Campylobacter* PubMLST server [18].

Comparative cgMLST analysis with *C. jejuni* and *C. coli* clinical strains worldwide was

performed with genomes available at the NCBI database. Annotations and visualizations

were performed using iTOL v.4 [21].

Virulome, resistome and comparative genomic analysis

A local BLAST database was constructed using fasta files of all sequenced genomes and the

makeblastdb program from BLAST (NCBI) [22], which was screened for the nucleotide

sequence of known pathogenicity genes using BLASTn (version 2.8.1) [22]. A 90% sequence

length and 90% identity threshold were used to select positive matches. The fasta file

containing the nucleotide sequence of virulence genes screened can be found at

https://doi.org/10.5281/zenodo.3925206. The C257T mutation in *gyrA* and mutations

A2058C and A2059G in the genes for 23S RNA were searched manually by BLAST and

sequences were aligned to the reference gene from strain *Campylobacter jejuni* ATCC 33560

using ClustalX [23]. Antimicrobial resistance genes were screened in each *Campylobacter*

genome using the ABRicate pipeline [24], using the Resfinder [25], CARD [26], ARG-

ANNOT [27] and NCBI ARRGD [28] databases. BLAST-based genome comparisons and

visualization of genetic clusters were performed using EasyFig [29].

Pangenome and plasmid screening analysis

Annotated genome sequences were compared using Roary [30], setting a minimum

percentage identity of 90% for BLASTp. Pangenome visualization was performed using

roary_plots.py. Additionally, the file gene_presence_absence.csv produced by Roary was

analyzed using a spreadsheet program to find genes that are shared by subgroups of isolates.

Venn diagrams were constructed using the tool available at

http://bioinformatics.psb.ugent.be/webtools/Venn/. Plasmids screening was performed on

FASTQ files using PlasmidSeeker [31], based on the default bacterial plasmid database. The

inhouse database of known plasmids from *C. coli* or *C. jejuni* and the output table for the

presence/abscence of gene content in the Roary analysis can be found in

https://doi.org/10.5281/zenodo.3925206. Pangenome BLAST analysis was performed using

GView (https://server.gview.ca/) [32].

RESULTS

Phylogenetic analysis of clinical Campylobacter strains.

cgMLST analysis showed high genetic diversity among the 69 *C. jejuni* strains, as

shown by the 14 CCs and 31 distinct STs identified (**Figure 1A**). Almost 93% (n=64) of the

C. jejuni strains belonged to one of the CCs previously described for this species. Among our

strains, CC-21 was the most common (37.6%) and diverse (5 distinct STs) (**Figure 1A, Table**

S1). ST-1359 was the most prevalent ST of CC-21 (48%) of all *C. jejuni* strains in this study

(17.4%) (**Figure 1A and Table S1**). In addition to CC-21, we identified high prevalence

rates of CC-48 (13%) and CC-52 (7.3%). ST-475 from CC-48 was the second most prevalent

ST (11.6%) identified (**Figure 1A**).

cgMLST analysis of the 12 *C. coli* strains from our collection revealed that 83%

(10/12) of the strains belonged to clade 1, while 17% (2/12) belonged to clade 3 and none to

clade 2. Most clade 1 strains (8/10) belonged to CC-828 and included four previously

described STs (828, 829, 1173, and 1556) and one novel ST (10201). Only one strain from

clade 1 belonged to the usually common CC-1150 (strain CFSAN093253 from ST-10203),

and another strain could not be assigned to any clonal complex (strain CFSAN096322 from

ST-10202). Both clade 3 strains belonged to ST-10204.

A global cgMLST analysis comparing our 81 *Campylobacter* genomes with other *C.*

jejuni and *C. coli* genomes available from the NCBI database was performed. The strains of

our collections from Central Chile showed great diversity and did not group with any existing

clusters (**Figure 1B).** The reconstructed phylogenetic trees and associated metadata of

clinical strains from Chile in the global context of *C. jejuni* and *C. coli* sequenced strains can

be interactively visualized on Microreact at https://microreact.org/project/VbEQsZtQD.

Pangenome analysis of Campylobacter strains.

194   To identify the genetic elements that are shared and distinct among the Chilean *C.*

jejuni and *C. coli* strains, we performed pangenome analysis of the 81 strains from both

species (**Figure 2**). The analysis identified a pangenome of 6609 genes for both species and

identified the core genes shared between *C. jejuni* and *C. coli* (532 genes) and the core

genome of each species (1300 genes for *C. jejuni* and 1207 genes for *C. coli*). In addition,

the analysis was able to identify the differences among the core and accessory genomes of

C. coli strains from clade 1 and clade 3. *C. coli* strains from clade 1 harbored 340 core genes

absent from clade 3 core genes, which in turn, harbored 159 core genes not present in clade

1 core genes (**Figure 2**). The estimated core genome for *C. jejuni* in our study (1300 genes)

was close to the 1343 core genes identified in the development of the cgMLST scheme of *C.*

jejuni [19] and in agreement with recent reports [33] (**Table S2**). In addition, a significant

variation in the accessory genome between *C. jejuni* and *C. coli* strains was identified,

confirming the high genetic variability of *Campylobacter*. Accessory genes that are common

to a particular cluster seem to contribute to modifications of carbohydrates and flagella as

well as antibiotic resistance. There is also a proportion of genomes harboring phage

sequences as well as T4SS and T6SS-related genes (see below).

Genomic analysis of the resistome.

To determine the resistome, we performed *in silico* analysis to identify genes

associated with antimicrobial resistance (**Figure 3**) by screening each *Campylobacter*

genome by means of the ABRicate pipeline [24] using the Resfinder, CARD, ARG-ANNOT

and NCBI ARRGD databases. We additionally performed sequence alignments to identify

point mutations in the *gyrA* and 23S rRNA genes. The *cmeABC* operon was present in all *C.*

jejuni and *C. coli* strains analyzed. This operon encodes a common multidrug efflux pump

characterized in different *Campylobacter* species, which mainly confers resistance to

fluoroquinolones and in some cases to macrolides [34]. Additionally, the point mutation in

the quinolone resistance-determining region (QRDR) of the *gyrA* gene, leading to the T86I

substitution in GyrA, was identified in 54.3% of the strains. This mutation confers high levels

of resistance to fluoroquinolones [35].

In contrast, a much lower fraction of the strains harbored resistance markers for other

antimicrobials. 22.2% of the strains harbored the *tetO* gene, which confers tetracycline

resistance in *Campylobacter* spp [36]. Furthermore, we did not detect neither mutation of the

23S rRNA gene nor presence of the *ermB* gene, both associated to macrolide resistance [37].

Finally, we also found a high proportion of strains (79%) harboring the blaOXA-61 gene

among *C. jejuni* and *C. coli* strains. This genetic marker is associated with β-lactam resistance

and has high prevalence in *Campylobacter* strains worldwide [38].

Presence and diversity of virulence gene content among Chilean Campylobacter strains.

To determine the virulence gene content of the *C. jejuni* and *C. coli* strains, we first

assembled a list of 220 potential virulence genes, including genes described in the Virulence

Factor Database [39] and genes reported to contribute to the virulence of *Campylobacter* in

the literature [40–43]. These genes were grouped into five distinct categories (adhesion and

colonization, invasion, motility, secretion systems and toxins) and were used to screen each

of the 81 genomes using BLASTn (**Figure 4**, **Figure S1** and **Table S3** for the full list). Most

of the described factors involved in adhesion and colonization (*cadF*, *racR*, *jlpA, pdlA, dnaJ,*

aphC) were prevalent just among the *C. jejuni* strains and were not associated with any

particular STs (**Figure 4**, **Table S3**).

The genes responsible for the production of the CDT toxin (*cdtA*, *cdtB* and *cdtC*) were

also found in most strains of *C. jejuni* and in every *C. coli* strain tested. The *cdtA* and *cdtB*

genes were not detected in the *C. jejuni* strain CFSAN093229. In contrast, the *ciaB* gene was

identified only in *C. jejuni* (in 98.7% of the strains) (**Figure 4**). The product of this gene is

involved in cecal colonization in chickens and in translocation into host cells in *C. jejuni*

[44,45]. The absence of this gene in *C. coli* might suggest that it does not play a crucial role

in the pathogenesis of this *Campylobacter* species.

The major differences among *C. jejuni* strains were observed for genes located at the

cps locus (Cj1413c -Cj1442) involved in capsule polysaccharide synthesis, the LOS locus

(C8J1074-C8J1095) involved in lipooligosaccharide synthesis, a small set of genes in the O-

linked flagellin glycosylation island (Cj1321-Cj1326) and the T6SS and T4SS gene clusters

(**Figure 4**, **Figure S1 and Table S3**). Additionally, *C. coli* strains carried most of the genes

known to be associated with invasiveness in *C. jejuni*, including the *iamb, flaC* and *ciaB*

genes. Both *C. coli* clades also harbored some genes associated with other steps of infection

such as the *cdt* toxin gene and most of the flagellar biosynthesis and chemiotaxis-related

genes described for *C. jejuni* (**Figure S1**). its relevance in pathogenicity remains unclear [46].

Distribution of the T6SS and T4SS gene clusters.

Our analysis identified the T6SS gene cluster in 9 *C. jejuni* strains and in 1 *C. coli*

strain from clade 1 (**Figure 4**). As shown in **Figure 5A**, whenever present the genetic

structure and the sequence identity of this cluster was highly conserved among *C. jejuni* and

C. coli strains. For strains with genomes that were not closed, we were not able to confirm

whether the *tssI* gene is located within the genetic context of the of the T6SS cluster, since it

was not assembled in the same contig as the rest of the cluster. The *Campylobacter* T6SS

gene cluster is often encoded within the CjIE3 conjugative element in the chromosome of *C.*

jejuni [41], but it can also be found in plasmids [47,48]. Since most of our sequenced strains

correspond to draft genome assemblies, we could not determine whether the T6SSs were

plasmid or chromosomally encoded in most of them. However, we have closed the genomes

of 17 of these strains [15]. In two of these strains (CFSAN093238 and CFSAN093227), T6SS

was inserted in the chromosomal CjIE3 element, while in strain CFSAN093246, it was

plasmid-encoded. Sequence-based analysis of this plasmid showed a high degree of identity

with plasmid pMTVDDCj13-2 (**Figure 6A**), which encodes a complete T6SS gene cluster

[48].

Genome analysis identified a T4SS gene cluster in 12 strains of *C. jejuni* and 1 strain

of *C. coli* clade 1 (**Figure 4**). It harbored the 16 core genes and was correctly assembled in

individual contigs in 11 strains, which allowed the analysis of the genetic context of the

cluster. In each strain, the T4SS cluster had a conserved genetic structure (**Figure 5B**). One

of the main differences between the T4SS gene clusters encoded in plasmids pTet and pCC31

is the sequence divergence of the *cpp33* gene [49]. While *cp33* present in strain

CFSAN096321 showed high identity with the gene harbored in the pTet plasmid, every other

strain possessed high identity to the *cp33* encoded in pCC31 plasmid. Both T4SS have been

involved in bacterial conjugation and are not directly linked to the virulence of *C. jejuni* and

C. coli strains [49]. As shown in **Figure 6B**, plasmids from closed genome strains

CFSAN093226 and CFSAN096297 share significant similarity with both the pCC31 and

pTET plasmids beyond the T4SS gene cluster. For all other strains, BLASTn analysis of each

of our draft genome assemblies identified most of the plasmid genes found in

pCFSAN093226, pCFSAN096297 and pTet-like plasmids, suggesting that all the T4SS gene

clusters identified in these strains are encoded in plasmids as well.

296   **DISCUSSION**

297

298   In South America, Campylobacteriosis is an emerging and neglected foodborne

299   disease. Here, we performed an in-depth genome analysis of the largest collection of clinical

300   *Campylobacter* strains from Chile, which accounts for over 35% of the genome sequences

301   available of clinical strains from South America.

302

303   cgMLST analysis provided insight into key similarities and differences in comparison

304   to the genetic diversity reported for clinical *Campylobacter* strains worldwide. *C. jejuni*

305   strains were highly diverse being the CC-21 the most common. CC-21 is the largest and most

306   widely distributed CC, representing 18.9% of all *C. jejuni* strains submitted to the PubMLST

307   database. The high prevalence of ST-1359 in our study was unexpected, since it is infrequent

308   within the PubMLST database (0.06%, February 2020). However, ST-1359 has been

309   recognized as a prominent ST in Israel [50]. ST-45, the second most prevalent ST worldwide

310   (4.2%), was not identified among our strains. Previous reports from Chile identified four

311   major CCs, including CC-21 (ST-50), CC-48 (ST-475), CC-257 (ST-257), and CC-353 (ST-

312   353) [9,13]. Our study is consistent with these reports, except for the high prevalence of ST-

313   1359. Further studies are needed to determine if ST-1359 represents an emergent ST. Our

314   study also provided the first genome data of clinical *C. coli* strains from Chile. *C. coli* is

315   divided into three genetic clades [51]. Clade 1 strains are often isolated from farm animals

316   and human gastroenteritis cases while clade 2 and 3 strains are mainly isolated from

317   environmental sources [51]. We showed that clade 1 strains from CC-828 were most

318   prevalent, which is consistent with the worldwide distribution of this CC [51]. Interestingly,

319   despite the lower amount of *C. coli* strains isolated in the 2-year period of our study (n=12),

we were able to identify strains from the uncommon clade 3. Therefore, larger genomic

epidemiological studies are needed to determine the prevalence of clade 3 strains in Chile

and South America.

Although most *Campylobacter* infections are self-limiting, antibiotics are indicated

for patients with persistent and severe gastroenteritis, extraintestinal infections, or who are

immunocompromised [1]. The high percentage of *C. jejuni* strains with a mutation of the

gyrA gene, which confers quinolone resistance, is consistent with recent studies from Chile,

reporting ciprofloxacin resistance of 30-60% [9,11–13]. One study reported that most

resistant strains were associated with ST-48 [52], but we did not find an association to any

particular STs. However, all of the strains belonging to ST-1359, the most frequent ST

identified, harbored this mutation. Since fluoroquinolones are not the first choice

antimicrobials for human campylobacteriosis, it has been suggested that high levels of

resistance might be a consequence of their broad use in animal husbandry in Latin America

[53]. These animal production practices might be responsible for the dissemination of

antibiotic resistance genes among Chilean *Campylobacter* strains.

None of the clinical strains harbored the mutation of the 23S rRNA gene nor presence

of the *ermB* gene which confers macrolide resistance, which is consistent with their choice

as first line treatment for *Campylobacter* infections [37]. Tetracycline resistance is

widespread among *Campylobacter* strains worldwide [35]. Recently, the European Union

reported high levels of resistance in *C. jejuni* strains (45.4%) and even higher levels in *C.*

coli (68.3%). Tetracycline is especially used in poultry industry worldwide [54], and might

serve as an important reservoir for resistant strains [55]. Indeed, it was recently reported that

32.3% of *C. jejuni* strains isolated from poultry in Chile were resistant to tetracycline [12].

Our data are consistent with this report, as 22% of the clinical *C. jejuni* strains were resistant.

This potential emergence of resistance highlights the need for a permanent surveillance

program to implement control measures for tetracycline usage in animal production.

While the ability of *Campylobacter* to cause human disease is thought to be

multifactorial, there are several genes associated with the virulence of *Campylobacter*, even

though their role in campylobacteriosis is not fully understood [56]. Genome analysis showed

that clinical *C. jejuni* strains harbored most of the known virulence factors described for the

species. On the contrary, *C. coli* strains lacked most virulence genes described for *C. jejuni*

(**Figure 4**). Since most of our knowledge regarding the virulence of *Campylobacter* comes

from studies performed in *C. jejuni*, this highlights the need for a better understanding of

how *C. coli* strains cause disease.

To date, there is only one T6SS cluster described in *Campylobacter* [40]. This T6SS

has been shown to contribute to host cell adherence, invasion, resistance to bile salts and

oxidative stress [57–59], and required for the colonization of murine [57] and avian [59]

infection models. We identified a complete T6SS gene cluster in 9 *C. jejuni* and in 1 *C. coli*

strain (**Figure 4**). No correlation was found between presence of the T6SS and any particular

ST, which differs from what has been described in Israel, where most T6SS-positive strains

belong to ST-1359 [50]. Although ST-1359 was the most common ST described in this study,

none of the strains carried a T6SS.

Two distinct T4SSs have been described in *Campylobacter*. One T4SS is encoded in

the pVir plasmid of *C. jejuni* and has been shown to contribute to the invasion of INT407

cells and the ability to induce diarrhea in a ferret infection model [42,60]. The second T4SS

is encoded in the pTet and pCC31 plasmids of *C. jejuni* and *C. coli,* which contributes to

bacterial conjugation and is not directly linked to virulence [49]. Each of the T4SS gene

clusters identified in our strains showed a high degree of identity to the pTet and pCC31

T4SSs. Suggesting that they do not directly contribute to virulence, but they could facilitate

horizontal gene transfer events that could lead to increase fitness and virulence.

Altogether, we provide valuable epidemiological and genomic data of the diversity,

virulence and resistance profiles of a large collection of clinical *Campylobacter* strains from

Chile. Further studies are needed to determine the dynamics of transmission of pathogenic

Campylobacter to humans and the potential emergence of new virulence and antimicrobial

resistance markers in order to provide actionable public health data to support the design of

strengthened surveillance programs in Chile and South America.

References

[1]　Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, et al. Global Epidemiology of Campylobacter
384
Infection. Clin Microbiol Rev. 2015;28:687–720.

[2]　World Health Organization, Food and Agriculture Organization of the United Nations, World
Organisation for Animal Health. The global view of campylobacteriosis: report of an expert
consultation, Utrecht, Netherlands, 9-11 July 2012 [Internet]. Geneva: World Health Organization;
2013. Available from: https://apps.who.int/iris/handle/10665/80751.

[3]　Acheson D, Allos BM. Campylobacter jejuni Infections: Update on Emerging Issues and Trends.
Clinical Infectious Diseases. 2001;32:1201–1206.

[4]　World Health Organization E. Global priority list of antibiotic-resistant bacteria to guide research,
discovery, and development of new antibiotics. 2017;7.

[5]　Fernández H. [Campylobacter and campylobacteriosis: a view from South America]. Rev Peru Med
Exp Salud Publica. 2011;28:121–127.

[6]　Pascoe B, Schiaffino F, Murray S, et al. Genomic epidemiology of Campylobacter jejuni associated
with asymptomatic pediatric infection in the Peruvian Amazon. Makepeace BL, editor. PLoS Negl Trop
Dis. 2020;14:e0008533.

[7]　Instituto de Salud Pública de Chile. Vigilancia de laboratorio de Campylobacter spp. Chile, 2005–2013.
Boletín ISP. 2014;4:1–17.

[8]　Lorena Porte, Weitzel T, Barbe M, et al. Routine molecular diagnosis of enteropathogens by the
FilmArray gastrointestinal panel in a microbiological laboratory in Santiago, Chile. 28th European
Congress of Clinical Microbiology and Infectious Diseases. 2018;

[9]　Collado L, Muñoz N, Porte L, et al. Genetic diversity and clonal characteristics of ciprofloxacin-
resistant Campylobacter jejuni isolated from Chilean patients with gastroenteritis. Infection, Genetics
and Evolution. 2018;58:290–293.

[10]   Rivera F N, Bustos B R, Montenegro H S, et al. [Genotyping and antibacterial resistance of
408   Campylobacter spp strains isolated in children and in free range poultry]. Rev Chilena Infectol.
409   2011;28:555–562.

410   [11]   Gonzalez-Hein G, Cordero N, García P, et al. Análisis molecular de la resistencia a fluoroquinolonas y
411   macrólidos en aislados de Campylobacter jejuni de humanos, bovinos y carne de ave. Rev chil infectol.
412   2013;30:135–139.

413   [12]   Lapierre L, Gatica MA, Riquelme V, et al. Characterization of Antimicrobial Susceptibility and Its
414   Association with Virulence Genes Related to Adherence, Invasion, and Cytotoxicity in *Campylobacter*
415   *jejuni* and *Campylobacter coli* Isolates from Animals, Meat, and Humans. Microbial Drug Resistance.
416   2016;22:432–444.

417   [13]   Levican A, Ramos-Tapia I, Briceño I, et al. Genomic Analysis of Chilean Strains of *Campylobacter*
418   *jejuni* from Human Faeces. BioMed Research International. 2019;2019:1–12.

419   [14]   Bravo V, Varela C, Porte L, et al. Draft Whole-Genome Sequences of 51 Campylobacter jejuni and 12
420   Campylobacter coli Clinical Isolates from Chile. Rasko D, editor. Microbiol Resour Announc.
421   2020;9:MRA.00072-20, e00072-20.

422   [15]   Bravo V, Porte L, Weitzel T, et al. Complete genome sequences of 17 clinical Campylobacter jejuni
423   strains from Chile. Microbiology Resource Announcements. :9:e00535-20.

424   [16]   Darling AE, Mau B, Perna NT. progressiveMauve: Multiple Genome Alignment with Gene Gain, Loss
425   and Rearrangement. Stajich JE, editor. PLoS ONE. 2010;5:e11147.

426   [17]   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30:2068–2069.

427   [18]   Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb software, the
428   PubMLST.org website and their applications. Wellcome Open Res. 2018;3:124.

429   [19]   Cody AJ, Bray JE, Jolley KA, et al. Core Genome Multilocus Sequence Typing Scheme for Stable,
430   Comparative Analyses of Campylobacter jejuni and C. coli Human Disease Isolates. Diekema DJ,
431   editor. J Clin Microbiol. 2017;55:2086–2097.

432   [20]   Jünemann S, Sedlazeck FJ, Prior K, et al. Updating benchtop sequencing performance comparison. Nat
433   Biotechnol. 2013;31:294–296.

434   [21]   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments. Nucleic
435   Acids Research. 2019;47:W256–W259.

436   [22]   Altschul SF, Gish W, Miller W, et al. Basic local alignment search tool. J Mol Biol. 1990;215:403–410.

437   [23]   Larkin MA, Blackshields G, Brown NP, et al. Clustal W and Clustal X version 2.0. Bioinformatics.
438   2007;23:2947–2948.

439   [24]   Seemann T. tseemann/abricate [Internet]. 2020 [cited 2020 Jul 2]. Available from:
440   https://github.com/tseemann/abricate.

441   [25]   Zankari E, Hasman H, Cosentino S, et al. Identification of acquired antimicrobial resistance genes.
442   Journal of Antimicrobial Chemotherapy. 2012;67:2640–2644.

443   [26]   Jia B, Raphenya AR, Alcock B, et al. CARD 2017: expansion and model-centric curation of the
444   comprehensive antibiotic resistance database. Nucleic Acids Res. 2017;45:D566–D573.

445   [27]   Gupta SK, Padmanabhan BR, Diene SM, et al. ARG-ANNOT, a New Bioinformatic Tool To Discover
446   Antibiotic Resistance Genes in Bacterial Genomes. Antimicrob Agents Chemother. 2014;58:212–220.

447   [28]   Feldgarden M, Brover V, Haft DH, et al. Validating the AMRFinder Tool and Resistance Gene Database
448   by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a Collection of Isolates.
449   Antimicrob Agents Chemother. 2019;63:e00483-19, /aac/63/11/AAC.00483-19.atom.

450   [29]   Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer. Bioinformatics.
451   2011;27:1009–1010.

452   [30]   Page AJ, Cummins CA, Hunt M, et al. Roary: rapid large-scale prokaryote pan genome analysis.
453   Bioinformatics. 2015;31:3691–3693.

454   [31]   Roosaare M, Puustusmaa M, Möls M, et al. PlasmidSeeker: identification of known plasmids from
455   bacterial whole genome sequencing reads. PeerJ. 2018;6:e4588.

456   [32]   Stothard P, Grant JR, Van Domselaar G. Visualizing and comparing circular genomes using the
457   CGView family of tools. Briefings in Bioinformatics. 2019;20:1576–1582.

458   [33]   Méric G, Yahara K, Mageiros L, et al. A Reference Pan-Genome Approach to Comparative Bacterial
459   Genomics: Identification of Novel Epidemiological Markers in Pathogenic Campylobacter. Bereswill
460   S, editor. PLoS ONE. 2014;9:e92798.

461   [34]   Luangtongkum T, Jeon B, Han J, et al. Antibiotic resistance in *Campylobacter* : emergence,
462   transmission and persistence. Future Microbiology. 2009;4:189–200.

463   [35]   Tang Y, Fang L, Xu C, et al. Antibiotic resistance trends and mechanisms in the foodborne pathogen,
464   *Campylobacter*. Anim Health Res Rev. 2017;18:87–98.

465   [36]   Connell SR, Trieber CA, Dinos GP, et al. Mechanism of Tet(O)-mediated tetracycline resistance.
466   EMBO J. 2003;22:945–953.

467   [37]   Bolinger H, Kathariou S. The Current State of Macrolide Resistance in Campylobacter spp.: Trends and
468   Impacts of Resistance Mechanisms. Schaffner DW, editor. Appl Environ Microbiol. 2017;83:e00416-
469   17, e00416-17.

470   [38]   Alfredson DA, Korolik V. Isolation and expression of a novel molecular class D beta-lactamase, OXA-
471   61, from Campylobacter jejuni. Antimicrob Agents Chemother. 2005;49:2515–2518.

472   [39]   Liu B, Zheng D, Jin Q, et al. VFDB 2019: a comparative pathogenomic platform with an interactive
473   web interface. Nucleic Acids Research. 2019;47:D687–D692.

[40] Redondo N, Carroll A, McNamara E. Molecular characterization of Campylobacter causing human
clinical infection using whole-genome sequencing: Virulence, antimicrobial resistance and phylogeny
in Ireland. Zilch TJ, editor. PLoS ONE. 2019;14:e0219088.

[41] Agnetti J, Seth-Smith HMB, Ursich S, et al. Clinical impact of the type VI secretion system on virulence
of Campylobacter species during infection. BMC Infect Dis. 2019;19:237.

[42] Bacon DJ, Alm RA, Burr DH, et al. Involvement of a Plasmid in Virulence of Campylobacter jejuni 81-
176. Barbieri JT, editor. Infect Immun. 2000;68:4384–4390.

[43] Dasti JI, Tareen AM, Lugert R, et al. Campylobacter jejuni: A brief overview on pathogenicity-
associated factors and disease-mediating mechanisms. International Journal of Medical Microbiology.
2010;300:205–211.

[44] Konkel ME, Kim BJ, Rivera-Amill V, et al. Bacterial secreted proteins are required for the internaliztion
of Campylobacter jejuni into cultured mammalian cells. Mol Microbiol. 1999;32:691–701.

[45] Novik V, Hofreuter D, Galán JE. Identification of Campylobacter jejuni genes involved in its interaction
with epithelial cells. Infect Immun. 2010;78:3540–3553.

[46] Ghorbanalizadgan M, Bakhshi B, Kazemnejad Lili A, et al. A molecular survey of Campylobacter jejuni
and Campylobacter coli virulence and diversity. Iran Biomed J. 2014;18:158–164.

[47] Ghatak S, He Y, Reed S, et al. Whole genome sequencing and analysis of Campylobacter coli YH502
from retail chicken reveals a plasmid-borne type VI secretion system. Genom Data. 2017;11:128–131.

[48] Taveirne ME, Dunham DT, Perault A, et al. Complete Annotated Genome Sequences of Three
Campylobacter jejuni Strains Isolated from Naturally Colonized Farm-Raised Chickens. Genome
Announc. 2017;5:e01407-16, e01407-16.

[49] Batchelor RA, Pearson BM, Friis LM, et al. Nucleotide sequences and comparison of two large
conjugative plasmids from different Campylobacter species. Microbiology (Reading, Engl).
2004;150:3507–3517.

[50]   Rokney A, Valinsky L, Moran-Gilad J, et al. Genomic Epidemiology of Campylobacter jejuni
499   Transmission in Israel. Front Microbiol. 2018;9:2432.

500   [51]   Sheppard SK, Maiden MCJ. The Evolution of *Campylobacter jejuni* and *Campylobacter coli*. Cold
501   Spring Harb Perspect Biol. 2015;7:a018119.

502   [52]   Cody AJ, McCarthy NM, Wimalarathna HL, et al. A Longitudinal 6-Year Study of the Molecular
503   Epidemiology of Clinical Campylobacter Isolates in Oxfordshire, United Kingdom. Journal of Clinical
504   Microbiology. 2012;50:3193–3201.

505   [53]   Fernández H, Pérez-Pérez G. Campylobacter: fluoroquinolone resistance in Latin-American countries.
506   Archivos de medicina veterinaria. 2016;48:255–259.

507   [54]   Van Boeckel TP, Brower C, Gilbert M, et al. Global trends in antimicrobial use in food animals. Proc
508   Natl Acad Sci USA. 2015;112:5649–5654.

509   [55]   Maron D, Smith TJ, Nachman KE. Restrictions on antimicrobial use in food animal production: an
510   international regulatory and economic survey. Global Health. 2013;9:48.

511   [56]   Hansson I, Sandberg M, Habib I, et al. Knowledge gaps in control of *Campylobacter* for prevention of
512   campylobacteriosis. Transbound Emerg Dis. 2018;65:30–48.

513   [57]   Lertpiriyapong K, Gamazon ER, Feng Y, et al. Campylobacter jejuni Type VI Secretion System: Roles
514   in Adaptation to Deoxycholic Acid, Host Cell Adherence, Invasion, and In Vivo Colonization.
515   Bereswill S, editor. PLoS ONE. 2012;7:e42842.

516   [58]   Bleumink-Pluym NMC, van Alphen LB, Bouwman LI, et al. Identification of a Functional Type VI
517   Secretion System in Campylobacter jejuni Conferring Capsule Polysaccharide Sensitive Cytotoxicity.
518   Gaynor EC, editor. PLoS Pathog. 2013;9:e1003393.

519   [59]   Liaw J, Hong G, Davies C, et al. The Campylobacter jejuni Type VI Secretion System Enhances the
520   Oxidative Stress Response and Host Colonization. Front Microbiol. 2019;10:2864.

521　　[60]　　Bacon DJ, Alm RA, Hu L, et al. DNA sequence and mutational analyses of the pVir plasmid of

522　　　　　　Campylobacter jejuni 81-176. Infect Immun. 2002;70:6242–6250.

524 **FIGURES**

Figure 1. Phylogenetic analysis of clinical *C. jejuni* and *C. coli* strains from Chile. A)

Phylogenetic tree is based on cgMLST performed with RIDOM SeqSphere$^+$ and visualized

by iTOL. STs are shown in colored boxes for each strain, and tree branches are color coded

to highlight *C. jejuni* and *C. coli* strains from clades 1, 2 and 3. **B)** Global cgMLST analysis

of clinical *C. jejuni* and *C. coli* strains. Strains from this study are highlighted in red.

Figure 2. Visualization of the pangenome of the *C. jejuni* and *C. coli* strains. Pangenome

analysis was performed using Roary. Core and accessory genomes are highlighted with red

and black brackets, respectively. cgMLST phylogenetic tree of Figure 1A is shown on the

left. A Venn diagram highlighting the number of core genes present in *C. jejuni* and *C. coli*

strains from clades 1 and 3 is shown on the right.

Figure 3. Distribution of antimicrobial resistance genes. Binary heatmaps show the presence

and absence of antimicrobial resistance genes. Colored cells represent the presence of genes.

cgMLST phylogenetic tree of Figure 1A is shown on the left. Strain names are color coded

to highlight phylogenetic groups: *C. jejuni* strains (light green), *C. coli* clade 1 strains

(orange) and *C. coli* clade 3 strains (magenta).

Figure 4. Distribution of virulence-related genes. Binary heatmaps show the presence and

absence of virulence genes. Colored cells represent the presence of genes. cgMLST

phylogenetic tree of Figure 1A is shown on the left. Strain names are color coded to highlight

C. jejuni strains (green), *C. coli* clade 1 strains (orange) and *C. coli* clade 3 strains (purple).

Figure 5. Comparative genomic analysis of T6SS and T4SS gene clusters. **A)** T6SS gene

clusters of *C. jejuni* and *C. coli* strains (bold) compared to the cluster of *C. jejuni* strain 108.

Genes *tagH*, *tssD*, *tssB* and *tssC* are shown in color. **B)** T4SS gene clusters of *C. jejuni* strains

in comparison to the T4SS gene clusters of pTet and pCC31. The *cp33* gene is highlighted

in orange. BLASTn alignments were performed and visualized using EasyFig.

Figure 6. Comparative genomic analysis of plasmids encoding T6SS **(A)** and T4SS **(B)** gene

clusters. Pangenome BLAST analysis was performed using the Gview server. T6SS and

T4SS gene clusters are highlighted in red brackets.

Figure S1. Distribution of the presence (red) and absence (blue) of virulence-related genes

in *C. jejuni* and *C. coli* strains from this study.

Table S1. Metadata of strains used in this study.

Table S2. Frequency and distribution of CCs and STs of strains of this study.

Table S3. Identification of virulence genes. Cells with a red background represent

presence. Empty cells represent absence of the gene in each genome.