Genomic analysis of the diversity, antimicrobial resistance and virulence potential

of *Campylobacter jejuni* and *Campylobacter coli* strains from a private health care

center in Central Chile

Bravo V[1,2], Katz A[3], Porte L[4], Weitzel T[4,5], Varela, C[4], Gonzalez-Escalona N[6], Blondel

CJ[2]*

[1] Unidad de Microbiología, Escuela de Medicina, Facultad de Ciencias Médicas,

Universidad de Santiago de Chile, Santiago, Chile.

[2] Instituto de Ciencias Biomédicas, Facultad de Medicina y Facultad de Ciencias de la

Vida, Universidad Andrés Bello, 8320000, Santiago, Chile.

[3] Programa de Biología Celular y Molecular, ICBM, Facultad de Medicina, Universidad de

Chile, Santiago, 8380453, Chile.

[4] Laboratorio Clínico, Clínica Alemana de Santiago, Facultad de Medicina Clínica

Alemana, Universidad del Desarrollo, Santiago, Chile.

[5] Instituto de Ciencias e Innovación en Medicina (ICIM), Facultad de Medicina Clínica

Alemana, Universidad del Desarrollo, Santiago, Chile.

[6] Center for Food Safety and Applied Nutrition, Office of Regulatory Science, Division of

Microbiology, U.S. Food and Drug Administration, College Park, Maryland, USA.

Key words: *Campylobacter jejuni*, *Campylobacter coli*, cgMLST, comparative genomics,

foodborne pathogen, Chile.

*Corresponding author. carlos.blondel@unab.cl

Mailing address, Instituto de Ciencias Biomédicas, Universidad Andrés Bello. Echaurren

183, Santiago, 8320000, Chile.

ABSTRACT

Campylobacter jejuni and *Campylobacter coli* are the leading causes of human

gastroenteritis in the developed world and an emerging threat in developing countries.

There is increasing evidence that the incidence of campylobacteriosis in Chile and other

South American countries is greatly underestimated, mostly due to the lack of adequate

diagnostic methods. Accordingly, there is limited genomic and epidemiological data from

these countries, leaving an important knowledge gap in our understanding of the

population structure, virulence potential, and antimicrobial resistance profiles of clinical

Campylobacter strains. We recently published the whole-genome sequences of a

collection of 68 *C. jejuni* and 12 *C. coli* strains isolated from patients with enteric infections

during a 2-year period (2017-2019) attending Clinica Alemana in Santiago, Chile. In the

present study, we sequenced an additional *C. jejuni* genome and performed a genome-

wide analysis of the genetic diversity, virulence, and antimicrobial resistance potential of

these 81 strains. cgMLST analysis identified a high genetic diversity of *C. jejuni* as well as

13 novel STs and alleles in both *C. jejuni* and *C. coli*. Pangenome and virulome analyses

showed a differential distribution of virulence factors, including both plasmid and

chromosomally encoded T6SSs and T4SSs. Resistome analysis predicted widespread

resistance to fluoroquinolones, but a low rates of erythromycin resistance. This study

provides valuable genomic and epidemiological data and highlights the need for larger

genomic epidemiology studies to better understand the genetic diversity, virulence

potential and antimicrobial resistance of human clinical *Campylobacter* strains in South

America.

DATA SUMMARY

1. Genome sequences and assemblies of all *C. jejuni* and *C. coli* used in this study have been deposited in public databases (**Table S1**) and most strains have also been published as genome resource announcements [1,2].

2. Bacterial draft genome sequence reannotations and nucleotide sequence fasta files used for virulome and plasmid screening analysis are available as a supplementary dataset on Zenodo (DOI: 10.5281/zenodo.3925206) https://doi.org/10.5281/zenodo.3925206.

3. The reconstructed phylogenetic trees and associated metadata of clinical strains from Chile in the global context of *C. jejuni* and *C. coli* sequenced strains can be interactively visualized on Microreact at https://microreact.org/project/VbEQsZtQD.

INTRODUCTION

Campylobacteriosis caused by *Campylobacter jejuni* and *Campylobacter coli* has emerged as an important public health problem and is the most common bacterial cause of human gastroenteritis worldwide [3–5]. Although the mortality rates associated with campylobacteriosis are low (<0.01% to 8.8%), this foodborne illness has emerged as an important public health threat worldwide due to its high morbidity rates and the emergence of antimicrobial resistance [6,7]. In this context, the European Food Safety Authority (EFSA) has estimated a disease burden and annual cost of 2.4 billion euros [8].

Campylobacteriosis is usually associated with a self-limiting illness characterized by diarrhea, nausea, abdominal cramps and bloody stools [9]. Only a minor percentage of cases are severe or fatal. Nevertheless, extraintestinal manifestations might cause long-term complications such as reactive arthritis, and Miller-Fisher or Guillain-Barré syndrome [10,11]. Despite the self-limiting course, certain populations such as children under 5 years of age, elderly patients, and immunocompromised patients might suffer severe infections and require antimicrobial treatment [9]. In this context, the World Health Organization (WHO) has included *Campylobacter* as a high priority pathogen [7] due to the worldwide emergence of strains with high level fluoroquinolone resistance [12,13].

In South America, there is limited data available on the prevalence of *C. jejuni* and *C. coli* in comparison to other enteric bacterial pathogens [3,14]. Furthermore, the incidence and number of cases reported varies substantially between countries, most probably due to differences in sensitivity of the diagnostic methods used in each surveillance program.

In Chile, the National Laboratory Surveillance Program (DTO. N° 7/2020) of the Public Health Institute of Chile (ISP) has reported a national incidence rate of campylobacteriosis

of 0.1 to 0.6/100.000 [15]. This is quite low compared to other high-income countries,

where incidence rates are two orders of magnitude higher [3]. Nevertheless, recent reports

suggest that the burden of campylobacteriosis in Chile has been greatly underestimated

due to suboptimal diagnostic protocols in both public and private health centers [15]. A

recent study showed a 10-fold increase of the detection rate of *Campylobacter* in human

stool samples after changing from microscopic stool examination by Gram staining (the

routine protocol in Chilean health centers) to culture-dependent techniques [16]. In

another study from the same group, *Campylobacter* spp. emerged as the second most

prevalent bacterial agents after diarrheagenic *Escherichia coli* when a multiplex PCR

based assay (FilmArray® Gastrointestinal Panel) was used [17].

To date, only few reports have analyzed the genetic diversity and antimicrobial resistance

profiles of clinical human isolates of *Campylobacter* in Chile [18–21]. Furthermore, these

studies have either used a limited number of strains or lacked the resolution provided by

sequence-based typing methods such as multilocus sequence typing (MLST). There is

only one report that used whole-genome sequence data (wgMLST) to determine the

genetic diversity of *C. jejuni* strains from the country, which was based on the first 3

genomes of Chilean *C. jejuni* strains. Therefore larger genomic epidemiology studies are

needed [22].

We recently published the whole-genome sequences of 69 *C. jejuni* and 12 *C. coli* human

clinical strains isolated from patients with enteric infection attending Clínica Alemana in

Santiago, Chile, during a 2-year period (2017-2019) [1]. For 17 of the *C. jejuni* strains, we

obtained complete genome assemblies [2]. This represents the largest collection of

genomes from Chilean human clinical *Campylobacter* strains and includes the first *C. coli*

genome in the country [1].

In the present study, we performed a genome-wide analysis of the genetic diversity,

virulence potential, and antimicrobial resistance profiles of these 81 clinical strains.

Importantly, we identified a high genetic diversity of both *C. jejuni* and *C. coli* strains,

including 13 novel sequence types (STs) and alleles. Resistome analysis predicted a

widespread resistance to fluoroquinolones, but a decreased resistance to tetracycline in

comparison to the reported resistance from other regions of the world, in addition to the

differential distribution of virulence factors, including plasmids and chromosomally

encoded T6SS and T4SS.

METHODS

Genome data set, sequencing and reannotation

The genome sequence dataset analyzed in this study is listed in **Table S1** and was

compiled from our recently published collection of human clinical strains of *C. jejuni* and

C. coli from Central Chile [1,2] and available whole-genome sequences of *C. jejuni* and *C.*

coli strains from public databases. We additionally included in this dataset the draft whole

genome sequence of *C. jejuni* strain CFSAN096303, from the same collection of Clinica

Alemana isolates which was sequenced for this study. The final genome composed of 2

contigs was obtained using a combination of long (nanopore) and short (Illumina,

NextSeq) sequencing reads. Briefly, for the short reads the sequencing library was

prepared using 1 ng DNA for the Nextera XT kit according to the manufacturer's

instructions. The strain was sequenced using the NextSeq sequencer (Illumina). We used

a NextSeq 500/550 high-output kit v2.5 (300 cycles) with 2 × 150-bp paired-end chemistry,

according to the manufacturer's instructions, with >200X average coverage. The long

reads were generated in a MinION sequencer (Nanopore Technologies, Oxford, UK). The

sequencing library was prepared using the rapid barcoding sequencing kit (SQK-

RBK004). The library for this strain was pooled with other 9 libraries and run in a FLO-

MIN106 (R9.4.1) flow cell, according to the manufacturer's instructions, for 48 h. The run

was base called live using default settings (Minknow v 19.06.8, Guppy v3.0.7). Default

parameters were used for all software unless otherwise specified. The sequencing outputs

for that run was 7.2 Gb (quality score, 11.46; N50, 6,000 bp; total reads, 569,349) for an

estimated genome average coverage of 60 to 90 X. Reads less than 5,000 bp and quality

scores of <7 were discarded for downstream analysis. To assemble the genome of *C.*

jejuni strain CFSAN096303, the Illumina reads were managed with the CLC Genomics

Workbench v9.5.2 (Qiagen), assessed for quality (Q, >30) with the quality control tool, and

trimmed (adapter trimming, quality trimming, and length trimming) with the trim sequences

tool. The final complete genome sequence for this strain was obtained using a previously

described pipeline [1], except that Flye was used instead of CANU for long read de novo

assembling. The genomes were annotated using the NCBI Prokaryotic Genomes

Automatic   Annotation   Pipeline   (PGAP,

http://www.ncbi.nlm.nih.gov/genome/annotation_prok). Default parameters were used for

all software unless otherwise specified. The NextSeq data was deposited in NCBI under

accession number SRR10859573. The nanopore raw data was deposited in NCBI under

accession number SRR11786978. The final draft genome was deposited under accession

number WXZQ01000000. Fasta files of all draft genomes from the collection of strains

from Chile were reordered to the reference genomes of the *C. coli* aerotolerant strain

OR12 (accID GCF_002024185.1) and *C. jejuni* strain NCTC 11168 (accID

GCF_000009085.1) using the Mauve Contig Mover (MCM) from the Mauve package [23].

The ordered genomes were reannotated using Prokka [24] with the same reference files

mentioned above and by forcing GenBank compliance.

MLST and cgMLST analysis:

For MLST and cgMLST, genome assemblies ere mapped against an MLST scheme based

on seven housekeeping genes [25] and the 1,343-locus cgMLST scheme [26], using

Ridom SeqSphere software (Münster, Germany) [27]. *C. hepaticus HV10* (accID GCF-

001687475.2)*, C. upsaliensis DSM 5365* (accID GCF-000620965.1), *C. lari RM2100*

(accID GCF-000019205.1), *C. jejuni NCTC 11168* (accID GCF-000009085.1*), C. jejuni*

*doylei 269.97 (*accID GCA-000017485.1*),* aerotolerant *C. coli OR12 (*accID *GCF-*

*002024185.1), C. coli BIGS0008 (*accID *GCA-000314165.1), C. coli 76339 (*accID GCA-

000470055.1*),* and *C. coli H055260513* (accID GCA-001491555.1) were used as

reference genomes. Phylogenetic comparison of the 81 *Campylobacter* genome

sequences was performed using a neighbor-joining (NJ) tree based on a distance matrix

of the core genomes of all the strains. STs and clonal complexes (CCs) were determined

after automated allele submission to the PubMLST server for *Campylobacter* MLST

(https://pubmlst.org/campylobacter/). Additionally, to compare the isolates with previously

described *C. jejuni* and *C. coli* isolates worldwide, we performed a similar phylogenetic

analysis using the NCBI database including the isolates reported for the whole Latin

American region. Further annotations and visualizations were performed using iTOL v.4

[28].

Virulome, resistome and comparative genomic analysis

A local BLAST database was constructed using the fasta files of all sequenced genomes

and the makeblastdb program from BLAST (NCBI) [29]. Subsequently, the database was

screened for the nucleotide sequence of known pathogenicity genes using BLASTn

(version 2.8.1) [29]. A threshold of 90% of the original sequence length and 90% identity

was used to select positive matches. A fasta file containing the nucleotide sequence of

Campylobacter pathogenicity genes screened in this work can be found at

https://doi.org/10.5281/zenodo.3925206. Mutations in *gyrA* and 23S rRNA were searched

as indicative of resistance to fluoroquinolones and macrolides, respectively. The

sequences of both genes were screened by BLAST in the local BLAST database and

sequences were aligned to the reference gene from strain Campylobacter jejuni ATCC

33560 using ClustalX [30]. We then manually searched for mutation C257T in *gyrA* and

mutations A2058C and A2059G in the genes for 23S RNA. Antimicrobial resistance genes

were additionally screened in each *Campylobacter* genome by means of the ABRicate

pipeline [31], using the included Resfinder [32], CARD [33], ARG-ANNOT [34] and NCBI

ARRGD [35] databases. BLAST-based genome comparison analysis and visualization of

genetic clusters were performed using EasyFig [36].

Pangenome and plasmid screening analysis

Annotated genome sequences from all sequenced genomes were compared using Roary [37], setting a minimum percentage identity of 90% for BLASTp. Visualization of genes shared between genomes was performed using roary_plots.py. Additionally, the file gene_presence_absence.csv [https://github.com/sanger-pathogens/Roary/tree/master/contrib/roary_plots] produced by Roary was further analyzed using a spreadsheet program to find genes that are shared by subgroups of isolates. Using this information Venn diagrams were constructed using the tool available at http://bioinformatics.psb.ugent.be/webtools/Venn/. The screen for plasmids was performed on the FASTQ files of each sequencing reaction using PlasmidSeeker [38] based on the default bacterial plasmid database. Additional searches were performed using an inhouse database containing known plasmids from *C. coli* or *C. jejuni*. The fasta file for in-house databases can be found in https://doi.org/10.5281/zenodo.3925206. Pangenome BLAST analysis was performed using the GView server (https://server.gview.ca/) [39].

Results and Discussion

Phylogenetic analysis of clinical Campylobacter strains

cgMLST analysis showed high genetic diversity among the 69 *C. jejuni* strains (**Figure**

1A), as shown by the 14 CCs and 31 distinct STs identified. Almost 93% (n=64) of the *C.*

jejuni strains belonged to one of the CCs previously described for this species. Among our

strains, CC-21 was the most common (37.6%) and diverse (5 distinct STs) (**Figure 1A,**

Table S1). This is consistent with other reports, which describe CC-21 as the largest and

most widely distributed clonal complex representing 18.9% of all *C. jejuni* isolates

submitted to the PubMLST database. ST-1359 was the most prevalent ST of CC-21 (48%)

of all strains in this study (17.4%) (**Figure 1A and Table S1**). The high prevalence of ST-

1359 was unexpected, since it is infrequent within the PubMLST database (0.06% by

February 2020). However, ST-1359 has been recognized as a prominent ST in Israel [40].

Other STs within CC-21 included ST-50 (8.7%) and ST-21 (1.45%), which were

predominant in reports from Israel, Belgium, and the United Kingdom [40–42] as well as

worldwide in PubMLST (4.1% and 4.3% respectively). ST-45, the second most prevalent

sequence type worldwide worldwide (4.2%), was not identified among our strains. In

addition to CC-21, we identified high prevalence rates of CC-48 (13%) and CC-52 (7.3%).

ST-475 from CC-48 was the second most prevalent ST (11.6%) identified (**Figure 1A**).

There are only two previous reports of MLST data of clinical *C. jejuni* strains from Chile

[18,22]. Both reports identified four major CCs and STs, including CC-21 (ST-50), CC-48

(ST-475), CC-257 (ST-257), and CC-353 (ST-353). Our study is consistent with these

reports, except for the high prevalence of ST-1359 (CC-21), which has not been previously

reported in Chile. Further epidemiological information is needed to determine if the high

prevalence of these four CCs is specific to Chile and to determine if the high prevalence

of ST-1359 strains in our report is an isolated finding or if it represents an emergent ST in

Chile.

C. coli is divided into three genetic clades [43,44]. Clade 1 is subdivided into two major

clonal complexes, CC-828 and CC-1150, and often isolated from farm animals and human

cases of gastroenteritis [44]. Clades 2 and 3 lack clonal complex substructures and are

mainly isolated from environmental sources [43,44]. cgMLST analysis of the 12 *C. coli*

strains from our collection revealed that 83% (10/12) of the strains belonged to clade 1,

while 17% (2/12) belonged to clade 3 and none to clade 2. Most clade 1 strains (8/10)

belonged to CC-828 and included four previously described STs (828, 829, 1173, and

1556) and one novel ST (10201). Only one strain from clade 1 belonged to the usually

common CC-1150 (strain CFSAN093253 from ST-10203), and another strain could not

be assigned to any clonal complex (strain CFSAN096322 from ST-10202). Both clade 3

strains belonged to ST-10204. Overall, clade 1 strains from CC-828 were most prevalent,

which is consistent with the worldwide distribution of this species. Interestingly, despite

the low incidence of cases of gastroenteritis associated with *C. coli*, in the 2-year period

of our study (only 12 strains in total), we were able to identify strains from the uncommon

clade 3. Considering that these are the first strains of *C. coli* described from Chile and

considering the size of our study, further genomic epidemiological studies are needed to

fully grasp the diversity of the strains from this country.

A global cgMLST analysis comparing our 81 *Campylobacter* genomes with other *C. jejuni*

and *C. coli* genomes available from the NCBI database was performed. The strains of our

collections from Central Chile showed great diversty and did not group with any existing

clusters (**Figure 1B)** (the interactive tree and metadata are available in Microreact, see

data summary).

Pangenome analysis of Campylobacter strains

To identify the genetic elements that are shared and distinct among the Chilean *C. jejuni*

and *C. coli* strains, we performed pangenome analysis of the 81 strains from both species

(**Figure 2**). The analysis identified a pangenome of 6609 genes for both species and

identified the core genes shared between *C. jejuni* and *C. coli* (532 genes) and the core

genome of each species (1300 genes for *C. jejuni* and 1207 genes for *C. coli*). In addition,

the analysis was able to identify the differences among the core and accessory genomes

of *C. coli* strains from clade 1 and clade 3. *C. coli* strains from clade 1 harbored 340 core

genes absent from clade 3 core genes, which in turn, harbored 159 core genes not present

in clade 1 core genes (**Figure 2**). The estimated core genome for *C. jejuni* in our study

(1300 genes) was close to the 1343 core genes identified in the development of the

cgMLST scheme of *C. jejuni* [25] and in agreement with recent reports [45–47] (**Table**

S2). In addition, a significant variation in the accessory genome between *C. jejuni* and *C.*

coli strains was identified, confirming the high genetic variability of *Campylobacter*.

Accessory genes that are common to a particular cluster seem to contribute to

modifications of carbohydrates and flagella as well as antibiotic resistance. There is also

a proportion of genomes harboring phage- and type IV and VI secretion system-related

genes (see below).

Genomic analysis of the resistome of Chilean clinical Campylobacter strains.

Although most *Campylobacter* infections are self-limiting and do not require antibiotic

treatment, the use of antibiotics is indicated for patients with persistent and severe

gastroenteritis, patients with extraintestinal infections, and patients who are
immunocompromised [3].

To determine the resistome of the *C. jejuni* and *C. coli* strains, we performed an *in silico*
analysis to identify genes associated with antimicrobial resistance (**Figure 3**). We
screened each *Campylobacter* genome by means of the ABRicate pipeline [31], using the
Resfinder, CARD, ARG-ANNOT and NCBI ARRGD databases, to identify antimicrobial
resistance genes. We additionally performed sequence alignments to identify point
mutations in *gyrA* and 16S rRNA genes that have been previously described to confer
antimicrobial resistance.

The *cmeABC* operon was present in all *C. jejuni* and *C. coli* strains analyzed. This operon
encodes a common multidrug efflux pump characterized in different *Campylobacter*
species, which mainly confers resistance to fluoroquinolones and macrolides [48,49].
Additionally, the point mutation in the quinolone resistance-determining region (QRDR) of
the *gyrA* gene, leading to the T86I substitution in GyrA, was identified in 54.3% of the
strains. This mutation is the most frequently reported mutation worldwide and confers high
levels of resistance to fluoroquinolones [48,49]. The high percentage of *C. jejuni* strains
with this mutation is consistent with recent studies from Chile, reporting ciprofloxacin
resistance in 30-60% of *C. jejuni* strains [18,20–22]. While Porte *et al.* reported that most
resistant strains were associated with ST-48 and ST-353, in our study, there was no
particular association of this mutation specific STs. However, it is important to note that
100% of strains belonging to the most frequent ST described in this work (ST-1359)
harbored the mutation in the QRDR region. , while most ST-475 isolates (the second most
prevalent ST) not have this mutation. Although fluoroquinolones are not first choice
antimicrobial for human campylobacteriosis, these high levels of resistance might be a

consequence of their broad use in animal husbandry. Fernandez and Perez-Perez

concluded that the primary source of transmission of resistant *Campylobacter* strains to

humans in Latin America is related to the consumption of animal-origin food and

interactions with environmental animal sources [50]. However, further studies are needed

to obtain a better understanding of the relevance of animal production practices in the

dissemination of antibiotic resistance genes among Chilean *Campylobacter* strains.

Regarding macrolide resistance, neither mutation of the 23S rRNA gene nor presence of

the *ermB* gene was detected, which is consistent with previous reports on Chilean *C. jejuni*

strains [18,22]. The low incidence of resistance against this group of antibiotics (especially

azithromycin and erythromycin) makes them the first choice for treatment of

Campylobacter infections [51]. However, further studies are necessary to analyze a higher

number of *C. coli* strains. This is particularly important given that according to EFSA,

macrolide resistance has been mostly detected in this species [52].

In terms of tetracycline resistance, 22.2% of the strains harbored the *tetO* gene, which is

the main genetic factor mediating tetracycline resistance in *Campylobacter* spp [53]. This

gene can be harbored on a plasmid or chromosomally encoded and is widespread among

Campylobacter strains worldwide [53]. Recently, the European Union (EU) reported very

high levels of tetracycline resistance in *C. jejuni* strains isolated from humans (45.4%) and

even higher levels of resistance in *C. coli* (68.3%). This antibiotic is widely used, especially

in the poultry industry worldwide [54], and might serve as an important reservoir for

resistant strains [55]. In fact, Lapierre *et al.* reported that 32.3% of *C. jejuni* isolates from

poultry in Chile were resistant to tetracycline [21]. Our data are consistent with a recent

report from Chile [21] that found that 22% of *C. jejuni* strains of clinical origin were

tetracycline resistant, which suggests that Chilean strains of *Campylobacter* have lower

levels of tetracycline resistance than strains from other countries. This highlights the need

for a permanent surveillance program to implement control measures for tetracycline

usage in animal production to avoid an increase in tetracycline resistance levels in the

human population. Finally, we also found a high proportion of strains (79%) harboring the

blaOXA-61 gene among *C. jejuni* and *C. coli* strains. This genetic marker is associated

with β-lactam resistance and has high prevalence in *Campylobacter* isolates worldwide

[56,57].

Presence and diversity of virulence gene content among Chilean Campylobacter

strains

The ability of *Campylobacter* to cause disease in humans is thought to be multifactorial,

as it lacks several of the key virulence factors used by other enteric bacterial pathogens

to cause disease (e.g., type III secretion systems). Nevertheless, there are several genes

and loci that are associated with the virulence and pathogenic potential of *Campylobacter*,

even though their role in the development of campylobacteriosis is not fully understood

[58]. Furthermore, most of our knowledge regarding the virulence potential of

Campylobacter strains comes from studies performed in *C. jejuni*; therefore, information

regarding the genetic determinants that contribute to the virulence of *C. coli* strains is still

lacking.

To determine the virulence gene content of the *C. jejuni* and *C. coli* strains, we first

assembled a list of 220 potential virulence genes, including genes described in the

Virulence Factor Database [59] and genes reported to contribute to the virulence of

Campylobacter in the literature [38,60–63]. These genes were grouped into five distinct

categories (adhesion and colonization, invasion, motility, secretion systems and toxins)

and were used to screen each of the 81 genomes using BLASTn. The distribution of the

major virulence factors described for *C. jejuni* and *C. coli* is shown in **Figure 4** (see **Figure**

S1 and **Table S3** for the full list).

Most of the described factors involved in adhesion and colonization (*cadF*, *racR*, *jlpA*,

pdlA, dnaJ, aphC) were prevalent among the *C. jejuni* strains and were not associated

with any particular STs (**Figure 4**, **Table S3**). This finding is consistent with other reports

indicating that the prevalence of these factors in clinical strains is very high [63] and shows

the relevance of adherence in the pathogenicity of *C. jejuni*. The genes responsible for the

production of the CDT toxin (*cdtA*, *cdtB* and *cdtC*) were also found in most strains of *C.*

jejuni and in every *C. coli* strain tested. The *cdtA* and *cdtB* genes were not detected in the

C. jejuni strain CFSAN093229. In contrast, the *ciaB* gene was identified only in *C. jejuni*

(in 98.7% of the strains) (**Figure 4**). The product of this gene is involved in cecal

colonization in chickens and in translocation into host cells in *C. jejuni* [64,65]. The

absence of this gene in *C. coli* might suggest that it does not play a crucial role in the

pathogenesis of this *Campylobacter* species. The major differences among *C. jejuni*

strains were observed for genes located at the *cps* locus (Cj1413c -Cj1442) involved in

capsule polysaccharide synthesis, the LOS locus (C8J1074-C8J1095) involved in

lipooligosaccharide synthesis, a small set of genes in the O-linked flagellin glycosylation

island (Cj1321-Cj1326) and the T6SS and T4SS gene clusters (**Figure 4**, **Figure S1 and**

Table S3).

Most virulence genes described for *C. jejuni* were absent in *C. coli* clades 1 and 3 strains,

highlighting the need for a better understanding of how *C. coli* strains cause the disease.

However, *C. coli* strains carried most of the genes known to be associated with

invasiveness in *C. jejuni*, including the *iamb, flaC* and *ciaB* genes. Both *C. coli* clades also

harbored some genes associated with other steps of infection such as the *cdt* toxin gene

and most of the flagellar biosynthesis and chemiotaxis-related genes described for *C.*

jejuni (**Figure S1**). The high prevalence of the *iamb* gene in *C. coli* has also been observed

in other studies, but its relevance in pathogenicity remains unclear [66].

Distribution of the T4SS and T6SS gene clusters.

To date, there is only one T6SS cluster described in *Campylobacter* [60], and it has been

linked to a wide variety of phenotypes, including host cell adherence, cellular invasion and

resistance to bile salt, and oxidative stress [67–69]. In addition, *Campylobacter* T6SS

mutant strains exhibit colonization defects in both mouse [67] and chicken [69] *in vivo*

infection models. The *Campylobacter* T6SS gene cluster is often encoded within the CjIE3

conjugative element in the chromosome of *C. jejuni* [61], but it can also be found in

plasmids [70,71]. Our analysis identified the T6SS gene cluster in 9 *C. jejuni* strains and

in 1 *C. coli* strain from clade 1 (**Figure 4**). As shown in **Figure 5A**, whenever present the

genetic structure and the sequence identity of this cluster was highly conserved among *C.*

jejuni and *C. coli* strains. For strains with genomes that were not closed, we were not able

to confirm whether the *tssI* gene is located within the genetic context of the of the T6SS

cluster, since it was not assembled in the same contig as the rest of the cluster [70].

Since most of our sequenced strains correspond to draft genome assemblies, we could

not determine whether the T6SSs were plasmid or chromosomally-encoded in most of

them. However, in two T6SS-positive strains with closed genomes (CFSAN093238 and

CFSAN093227), T6SS was inserted in the chromosomal CjIE3 element, while in strain

CFSAN093246, it was plasmid-encoded. Sequence-based analysis of this plasmid

showed a high degree of identity with plasmid pMTVDDCj13-2 (**Figure 6A**), which

encodes a complete T6SS gene cluster and was recently sequenced from the
MTVDSCj13 strain of *C. jejuni* [71].

No correlation was found between the T6SS gene cluster and a particular ST or clonal
complex. This is different from what has been described in other countries, such as Israel,
where most T6SS-positive strains belong to ST-1359 [40]. Although ST-1359 was the
most common ST described in this study, none of the strains carried genes encoding
T6SS.

To date, two distinct T4SSs have been described in *Campylobacter*. One of them is
encoded in the pVir plasmid of *C. jejuni* and is homologous to the T4SSs from *Helicobacter*
pylori [72]. The pVir T4SS has been linked to the virulence of *C. jejuni*. T4SS mutants
show reduced invasion capacity of INT407 cells and decreased diarrhea-inducing ability
in a ferret model [62,73]. The second T4SS is encoded in the pTet and pCC31 family of
plasmids of *C. jejuni* and *C. coli*. This distinct T4SS is involved in bacterial conjugation
and is not directly linked to the virulence of *C. jejuni* and *C. coli* strains [74].

We identified a T4SS gene cluster homologous to this second T4SS in 12 strains of *C.*
jejuni and 1 strain of *C. coli* clade 1 (**Figure 4**). It harbored the 16 core genes and was
correctly assembled in individual contigs in 11 strains, which allowed the analysis of the
genetic context of the cluster. As shown in **Figure 5B**, in each strain, the T4SS cluster
had a conserved genetic structure. One of the main differences between the T4SS gene
clusters encoded in plasmids pTet and pCC31 is the sequence divergence of the *cpp33*
gene [74]. Only the *cp33* present in strain CFSAN096321 showed high identity with the
gene harbored in the pTet plasmid. T4SS gene clusters from every other *C. jejuni* and *C.*
coli strain possessed high identity to the *cp33* encoded in pCC31 plasmid.

As shown in **Figure 6B**, plasmids pCFSAN093226 and pCFSAN096297 share significant

similarity with both the pCC31 and pTET plasmids beyond the T4SS gene cluster. For all

other strains, BLASTn analysis of each of our draft genome assemblies identified most of

the plasmid genes found in pCFSAN093226, pCFSAN096297 and pTet-like plasmids,

suggesting that all the T4SS gene clusters identified in these strains are encoded in

plasmids as well.

Conclusions:

The genomes reported in the current study represent 96.4% of the sequenced Chilean

Campylobacter strains and 36.8% of the strains from the South American region. Taken

together, the results of this study provide valuable epidemiological data and relevant

information about the population structure, virulence-associated genes and AMR profiles

of an important number of clinical *Campylobacter* isolates. Further studies are needed to

determine the dynamics of transmission of pathogenic *Campylobacter* from animals to

humans and to monitor the emergence of new virulence and antimicrobial resistance

markers that might represent a threat to the human population. Finally, we expect that

these actionable public health data will support the design of strengthened surveillance

programs, which will be essential for the development of an effective integrated health

system in Chile.

AUTHOR CONTRIBUTIONS

Veronica Bravo: Conceptualization, Methodology, Formal analysis, Investigation, Visualization, Data curation, Writing-Original Draft and review and editing.

Assaf Katz: Methodology, Formal analysis, Investigation, Visualization, Data Curation, Resources, Writing-Original Draft and review and editing.

Carmen Varela: Resources, Writing - Review and editing

Lorena Porte: Resources, Writing - Review and editing.

Thomas Weitzel: Resources, Writing - Review and editing.

Narjol Gonzalez-Escanola: Conceptualization, Methodology, Formal analysis, Resources, review and editing final manuscript

Carlos J Blondel: Conceptualization, Methodology, Formal analysis, Investigation, Visualization, Resources, Data curation, Writing - Review and editing, Funding acquisition, Supervision and Project administration.

CONFLICT OF INTERESTS

The authors declare that there are no conflicts of interest.

FUNDING INFORMATION

This work was funded by the Howard Hughes Medical Institute (HHMI)-Gulbenkian International Research Scholar Grant #55008749, FONDECYT Grant 1201805 (ANID) and REDI170269 (ANID) from CJB. VB is supported by DICYT grant 022001BZ (USACH). AK is supported by FONDECYT Grant 1191074 (ANID). NJG is supported by MCMi Challenge Grants program proposal number 2018-646 and the FDA Foods Program Intramural Funds (FDA employees).

REFERENCES

[1] Bravo V, Varela C, Porte L, Weitzel T, Kastanis GJ, Balkey M, et al. Draft Whole-Genome Sequences of 51 Campylobacter jejuni and 12 Campylobacter coli Clinical Isolates from Chile. Microbiol Resour Announc 2020;9:MRA.00072-20, e00072-20. https://doi.org/10.1128/MRA.00072-20.

[2] Bravo V, Porte L, Weitzel T, Varela C, Blondel C, Gonzalez-Escalona N. Complete genome sequences of 17 clinical Campylobacter jejuni strains from Chile. Microbiology Resource Announcements n.d.:9:e00535-20. https://doi.org/10.1128/MRA.00535-20.

[3] Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, Man SM. Global Epidemiology of Campylobacter Infection. Clin Microbiol Rev 2015;28:687–720. https://doi.org/10.1128/CMR.00006-15.

[4] World Health Organization, Food and Agriculture Organization of the United Nations, World Organisation for Animal Health. The global view of campylobacteriosis: report of an expert consultation, Utrecht, Netherlands, 9-11 July 2012. Geneva: World Health Organization; 2013.

[5] EFSA. The European Union One Health 2018 Zoonoses Report. EFSA Journal 2019;17:e05926. https://doi.org/10.2903/j.efsa.2019.5926.

[6] Eurosurveillance editorial team. European Union Summary Report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food 2012 published. Euro Surveill 2014;19:20748. https://doi.org/10.2807/ese.19.12.20748-en.

[7] World Health Organization E. Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics 2017:7.

[8] EFSA. Scientific Opinion on Campylobacter in broiler meat production: control options and performance objectives and/or targets at different stages of the food chain. EFSA Journal 2011;9:2105. https://doi.org/10.2903/j.efsa.2011.2105.

[9] Acheson D, Allos BM. Campylobacter jejuni Infections: Update on Emerging Issues and Trends. Clinical Infectious Diseases 2001;32:1201–6. https://doi.org/10.1086/319760.

[10] Wijdicks EFM, Klein CJ. Guillain-Barré Syndrome. Mayo Clinic Proceedings 2017;92:467–79. https://doi.org/10.1016/j.mayocp.2016.12.002.

[11] Scallan Walter EJ, Crim SM, Bruce BB, Griffin PM. Incidence of *Campylobacter* -Associated Guillain-Barré Syndrome Estimated from Health Insurance Data. Foodborne Pathogens and Disease 2020;17:23–8. https://doi.org/10.1089/fpd.2019.2652.

[12] Yang Y, Feye KM, Shi Z, Pavlidis HO, Kogut M, J. Ashworth A, et al. A Historical Review on Antibiotic Resistance of Foodborne Campylobacter. Front Microbiol 2019;10:1509. https://doi.org/10.3389/fmicb.2019.01509.

[13] Sproston EL, Wimalarathna HML, Sheppard SK. Trends in fluoroquinolone resistance in Campylobacter. Microbial Genomics 2018;4. https://doi.org/10.1099/mgen.0.000198.

[14] Fernández H. [Campylobacter and campylobacteriosis: a view from South America]. Rev Peru Med Exp Salud Publica 2011;28:121–7. https://doi.org/10.1590/s1726-46342011000100019.

[15] Instituto de Salud Pública de Chile. Vigilancia de laboratorio de Campylobacter spp. Chile, 2005–2013. Boletín ISP 2014;4:1–17.

[16] Porte L, Varela C, Haecker T, Morales S, Weitzel T. Impact of changing from staining to culture techniques on detection rates of Campylobacter spp. in routine stool

samples in Chile. BMC Infect Dis 2016;16:196. https://doi.org/10.1186/s12879-016-
1546-7.
[17]Lorena Porte, Weitzel T, Barbe M, Armijo Y, Vollrath V. Routine molecular diagnosis
of enteropathogens by the FilmArray gastrointestinal panel in a microbiological
laboratory in Santiago, Chile. 28th European Congress of Clinical Microbiology and
Infectious Diseases 2018.
[18]Collado L, Muñoz N, Porte L, Ochoa S, Varela C, Muñoz I. Genetic diversity and clonal
characteristics of ciprofloxacin-resistant Campylobacter jejuni isolated from Chilean
patients with gastroenteritis. Infection, Genetics and Evolution 2018;58:290–3.
https://doi.org/10.1016/j.meegid.2017.12.026.
[19]Rivera F N, Bustos B R, Montenegro H S, Sandoval M M, Castillo N J, Fernández J
H, et al. [Genotyping and antibacterial resistance of Campylobacter spp strains
isolated in children and in free range poultry]. Rev Chilena Infectol 2011;28:555–62.
https://doi.org//S0716-10182011000700008.
[20]González-Hein G, Cordero N, García P, Figueroa G. [Molecular analysis of
fluoroquinolones and macrolides resistance in Campylobacter jejuni isolates from
humans, bovine and chicken meat]. Rev Chilena Infectol 2013;30:135–9.
https://doi.org/10.4067/S0716-10182013000200003.
[21]Lapierre L, Gatica MA, Riquelme V, Vergara C, Yañez JM, San Martín B, et al.
Characterization of Antimicrobial Susceptibility and Its Association with Virulence
Genes Related to Adherence, Invasion, and Cytotoxicity in *Campylobacter jejuni* and
Campylobacter coli Isolates from Animals, Meat, and Humans. Microbial Drug
Resistance 2016;22:432–44. https://doi.org/10.1089/mdr.2015.0055.
[22]Levican A, Ramos-Tapia I, Briceño I, Guerra F, Mena B, Varela C, et al. Genomic
Analysis of Chilean Strains of *Campylobacter jejuni* from Human Faeces. BioMed
Research International 2019;2019:1–12. https://doi.org/10.1155/2019/1902732.
[23]Darling AE, Mau B, Perna NT. progressiveMauve: Multiple Genome Alignment with
Gene Gain, Loss and Rearrangement. PLoS ONE 2010;5:e11147.
https://doi.org/10.1371/journal.pone.0011147.
[24]Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics
2014;30:2068–9. https://doi.org/10.1093/bioinformatics/btu153.
[25]Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics:
BIGSdb software, the PubMLST.org website and their applications. Wellcome Open
Res 2018;3:124. https://doi.org/10.12688/wellcomeopenres.14826.1.
[26]Cody AJ, Bray JE, Jolley KA, McCarthy ND, Maiden MCJ. Core Genome Multilocus
Sequence Typing Scheme for Stable, Comparative Analyses of Campylobacter jejuni
and C. coli Human Disease Isolates. J Clin Microbiol 2017;55:2086–97.
https://doi.org/10.1128/JCM.00080-17.
[27]Jünemann S, Sedlazeck FJ, Prior K, Albersmeier A, John U, Kalinowski J, et al.
Updating benchtop sequencing performance comparison. Nat Biotechnol
2013;31:294–6. https://doi.org/10.1038/nbt.2522.
[28]Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. Nucleic Acids Research 2019;47:W256–9.
https://doi.org/10.1093/nar/gkz239.
[29]Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search
tool. J Mol Biol 1990;215:403–10. https://doi.org/10.1016/S0022-2836(05)80360-2.
[30]Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, et al.
Clustal W and Clustal X version 2.0. Bioinformatics 2007;23:2947–8.
https://doi.org/10.1093/bioinformatics/btm404.
[31]Seemann T. tseemann/abricate. 2020.

[32]Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al.
Identification of acquired antimicrobial resistance genes. Journal of Antimicrobial
Chemotherapy 2012;67:2640–4. https://doi.org/10.1093/jac/dks261.
[33]Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, et al. CARD 2017:
expansion and model-centric curation of the comprehensive antibiotic resistance
database. Nucleic Acids Res 2017;45:D566–73.
https://doi.org/10.1093/nar/gkw1004.
[34]Gupta SK, Padmanabhan BR, Diene SM, Lopez-Rojas R, Kempf M, Landraud L, et
al. ARG-ANNOT, a New Bioinformatic Tool To Discover Antibiotic Resistance Genes
in Bacterial Genomes. Antimicrob Agents Chemother 2014;58:212–20.
https://doi.org/10.1128/AAC.01310-13.
[35]Feldgarden M, Brover V, Haft DH, Prasad AB, Slotta DJ, Tolstoy I, et al. Validating the
AMRFinder Tool and Resistance Gene Database by Using Antimicrobial Resistance
Genotype-Phenotype Correlations in a Collection of Isolates. Antimicrob Agents
Chemother 2019;63:e00483-19, /aac/63/11/AAC.00483-19.atom.
https://doi.org/10.1128/AAC.00483-19.
[36]Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer.
Bioinformatics 2011;27:1009–10. https://doi.org/10.1093/bioinformatics/btr039.
[37]Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, et al. Roary: rapid
large-scale prokaryote pan genome analysis. Bioinformatics 2015;31:3691–3.
https://doi.org/10.1093/bioinformatics/btv421.
[38]Roosaare M, Puustusmaa M, Möls M, Vaher M, Remm M. PlasmidSeeker:
identification of known plasmids from bacterial whole genome sequencing reads.
PeerJ 2018;6:e4588. https://doi.org/10.7717/peerj.4588.
[39]Stothard P, Grant JR, Van Domselaar G. Visualizing and comparing circular genomes
using the CGView family of tools. Briefings in Bioinformatics 2019;20:1576–82.
https://doi.org/10.1093/bib/bbx081.
[40]Rokney A, Valinsky L, Moran-Gilad J, Vranckx K, Agmon V, Weinberger M. Genomic
Epidemiology of Campylobacter jejuni Transmission in Israel. Front Microbiol
2018;9:2432. https://doi.org/10.3389/fmicb.2018.02432.
[41]Elhadidy M, Arguello H, Álvarez-Ordóñez A, Miller WG, Duarte A, Martiny D, et al.
Orthogonal typing methods identify genetic diversity among Belgian Campylobacter
jejuni strains isolated over a decade from poultry and cases of sporadic human
illness. Int J Food Microbiol 2018;275:66–75.
https://doi.org/10.1016/j.ijfoodmicro.2018.04.004.
[42]Cody AJ, McCarthy NM, Wimalarathna HL, Colles FM, Clark L, Bowler ICJW, et al. A
Longitudinal 6-Year Study of the Molecular Epidemiology of Clinical Campylobacter
Isolates in Oxfordshire, United Kingdom. Journal of Clinical Microbiology
2012;50:3193–201. https://doi.org/10.1128/JCM.01086-12.
[43]Sheppard SK, McCarthy ND, Falush D, Maiden MCJ. Convergence of Campylobacter
Species: Implications for Bacterial Evolution. Science 2008;320:237–9.
https://doi.org/10.1126/science.1155532.
[44]Sheppard SK, Didelot X, Jolley KA, Darling AE, Pascoe B, Meric G, et al. Progressive
genome-wide introgression in agricultural *Campylobacter coli*. Mol Ecol
2013;22:1051–64. https://doi.org/10.1111/mec.12162.
[45]Méric G, Yahara K, Mageiros L, Pascoe B, Maiden MCJ, Jolley KA, et al. A Reference
Pan-Genome Approach to Comparative Bacterial Genomics: Identification of Novel
Epidemiological Markers in Pathogenic Campylobacter. PLoS ONE 2014;9:e92798.
https://doi.org/10.1371/journal.pone.0092798.
[46]Biggs PJ, Fearnhead P, Hotter G, Mohan V, Collins-Emerson J, Kwan E, et al. Whole-
Genome Comparison of Two Campylobacter jejuni Isolates of the Same Sequence

Type Reveals Multiple Loci of Different Ancestral Lineage. PLoS ONE
2011;6:e27121. https://doi.org/10.1371/journal.pone.0027121.
[47]Friis C, Wassenaar TM, Javed MA, Snipen L, Lagesen K, Hallin PF, et al. Genomic
Characterization of Campylobacter jejuni Strain M1. PLoS ONE 2010;5:e12253.
https://doi.org/10.1371/journal.pone.0012253.
[48]Luangtongkum T, Jeon B, Han J, Plummer P, Logue CM, Zhang Q. Antibiotic
resistance in *Campylobacter*: emergence, transmission and persistence. Future
Microbiology 2009;4:189–200. https://doi.org/10.2217/17460913.4.2.189.
[49]Shen Z, Wang Y, Zhang Q, Shen J. Antimicrobial Resistance in Campylobacter spp.
In: Schwarz, Cavaco, Shen, editors. Antimicrobial Resistance in Bacteria from
Livestock and Companion Animals, American Society of Microbiology; 2018, p. 317–
30. https://doi.org/10.1128/microbiolspec.ARBA-0013-2017.
[50]Fernández H, Pérez-Pérez G. Campylobacter: fluoroquinolone resistance in Latin-
American countries. Archivos de Medicina Veterinaria 2016;48:255–9.
https://doi.org/10.4067/S0301-732X2016000300002.
[51]Bolinger H, Kathariou S. The Current State of Macrolide Resistance in Campylobacter
spp.: Trends and Impacts of Resistance Mechanisms. Appl Environ Microbiol
2017;83:e00416-17, e00416-17. https://doi.org/10.1128/AEM.00416-17.
[52]Engberg J, Aarestrup FM, Taylor DE, Gerner-Smidt P, Nachamkin I. Quinolone and
macrolide resistance in Campylobacter jejuni and C. coli: resistance mechanisms
and trends in human isolates. Emerging Infect Dis 2001;7:24–34.
https://doi.org/10.3201/eid0701.010104.
[53]Connell SR, Trieber CA, Dinos GP, Einfeldt E, Taylor DE, Nierhaus KH. Mechanism
of Tet(O)-mediated tetracycline resistance. EMBO J 2003;22:945–53.
https://doi.org/10.1093/emboj/cdg093.
[54]Van Boeckel TP, Brower C, Gilbert M, Grenfell BT, Levin SA, Robinson TP, et al.
Global trends in antimicrobial use in food animals. Proc Natl Acad Sci USA
2015;112:5649–54. https://doi.org/10.1073/pnas.1503141112.
[55]Maron D, Smith TJ, Nachman KE. Restrictions on antimicrobial use in food animal
production: an international regulatory and economic survey. Global Health
2013;9:48. https://doi.org/10.1186/1744-8603-9-48.
[56]Alfredson DA, Korolik V. Isolation and expression of a novel molecular class D beta-
lactamase, OXA-61, from Campylobacter jejuni. Antimicrob Agents Chemother
2005;49:2515–8. https://doi.org/10.1128/AAC.49.6.2515-2518.2005.
[57]Marotta F, Garofolo G, di Marcantonio L, Di Serafino G, Neri D, Romantini R, et al.
Antimicrobial resistance genotypes and phenotypes of Campylobacter jejuni isolated
in Italy from humans, birds from wild and urban habitats, and poultry. PLoS ONE
2019;14:e0223804. https://doi.org/10.1371/journal.pone.0223804.
[58]Hansson I, Sandberg M, Habib I, Lowman R, Engvall EO. Knowledge gaps in control
of *Campylobacter* for prevention of campylobacteriosis. Transbound Emerg Dis
2018;65:30–48. https://doi.org/10.1111/tbed.12870.
[59]Liu B, Zheng D, Jin Q, Chen L, Yang J. VFDB 2019: a comparative pathogenomic
platform with an interactive web interface. Nucleic Acids Research 2019;47:D687–
92. https://doi.org/10.1093/nar/gky1080.
[60]Redondo N, Carroll A, McNamara E. Molecular characterization of Campylobacter
causing human clinical infection using whole-genome sequencing: Virulence,
antimicrobial resistance and phylogeny in Ireland. PLoS ONE 2019;14:e0219088.
https://doi.org/10.1371/journal.pone.0219088.
[61]Agnetti J, Seth-Smith HMB, Ursich S, Reist J, Basler M, Nickel C, et al. Clinical impact
of the type VI secretion system on virulence of Campylobacter species during
infection. BMC Infect Dis 2019;19:237. https://doi.org/10.1186/s12879-019-3858-x.

[62] Bacon DJ, Alm RA, Burr DH, Hu L, Kopecko DJ, Ewing CP, et al. Involvement of a Plasmid in Virulence of Campylobacter jejuni 81-176. Infect Immun 2000;68:4384–90. https://doi.org/10.1128/IAI.68.8.4384-4390.2000.

[63] Dasti JI, Tareen AM, Lugert R, Zautner AE, Groß U. Campylobacter jejuni: A brief overview on pathogenicity-associated factors and disease-mediating mechanisms. International Journal of Medical Microbiology 2010;300:205–11. https://doi.org/10.1016/j.ijmm.2009.07.002.

[64] Konkel ME, Kim BJ, Rivera-Amill V, Garvis SG. Bacterial secreted proteins are required for the internaliztion of Campylobacter jejuni into cultured mammalian cells. Mol Microbiol 1999;32:691–701. https://doi.org/10.1046/j.1365-2958.1999.01376.x.

[65] Novik V, Hofreuter D, Galán JE. Identification of Campylobacter jejuni genes involved in its interaction with epithelial cells. Infect Immun 2010;78:3540–53. https://doi.org/10.1128/IAI.00109-10.

[66] Ghorbanalizadgan M, Bakhshi B, Kazemnejad Lili A, Najar-Peerayeh S, Nikmanesh B. A molecular survey of Campylobacter jejuni and Campylobacter coli virulence and diversity. Iran Biomed J 2014;18:158–64. https://doi.org/10.6091/ibj.1359.2014.

[67] Lertpiriyapong K, Gamazon ER, Feng Y, Park DS, Pang J, Botka G, et al. Campylobacter jejuni Type VI Secretion System: Roles in Adaptation to Deoxycholic Acid, Host Cell Adherence, Invasion, and In Vivo Colonization. PLoS ONE 2012;7:e42842. https://doi.org/10.1371/journal.pone.0042842.

[68] Bleumink-Pluym NMC, van Alphen LB, Bouwman LI, Wösten MMSM, van Putten JPM. Identification of a Functional Type VI Secretion System in Campylobacter jejuni Conferring Capsule Polysaccharide Sensitive Cytotoxicity. PLoS Pathog 2013;9:e1003393. https://doi.org/10.1371/journal.ppat.1003393.

[69] Liaw J, Hong G, Davies C, Elmi A, Sima F, Stratakos A, et al. The Campylobacter jejuni Type VI Secretion System Enhances the Oxidative Stress Response and Host Colonization. Front Microbiol 2019;10:2864. https://doi.org/10.3389/fmicb.2019.02864.

[70] Ghatak S, He Y, Reed S, Strobaugh T, Irwin P. Whole genome sequencing and analysis of Campylobacter coli YH502 from retail chicken reveals a plasmid-borne type VI secretion system. Genom Data 2017;11:128–31. https://doi.org/10.1016/j.gdata.2017.02.005.

[71] Taveirne ME, Dunham DT, Perault A, Beauchamp JM, Huynh S, Parker CT, et al. Complete Annotated Genome Sequences of Three Campylobacter jejuni Strains Isolated from Naturally Colonized Farm-Raised Chickens. Genome Announc 2017;5:e01407-16, e01407-16. https://doi.org/10.1128/genomeA.01407-16.

[72] Kersulyte D, Velapatiño B, Mukhopadhyay AK, Cahuayme L, Bussalleu A, Combe J, et al. Cluster of type IV secretion genes in Helicobacter pylori's plasticity zone. J Bacteriol 2003;185:3764–72. https://doi.org/10.1128/jb.185.13.3764-3772.2003.

[73] Bacon DJ, Alm RA, Hu L, Hickey TE, Ewing CP, Batchelor RA, et al. DNA sequence and mutational analyses of the pVir plasmid of Campylobacter jejuni 81-176. Infect Immun 2002;70:6242–50. https://doi.org/10.1128/iai.70.11.6242-6250.2002.

[74] Batchelor RA, Pearson BM, Friis LM, Guerry P, Wells JM. Nucleotide sequences and comparison of two large conjugative plasmids from different Campylobacter species. Microbiology (Reading, Engl) 2004;150:3507–17. https://doi.org/10.1099/mic.0.27112-0.

751 **FIGURE LEGENDS**
752
Figure 1. A) Phylogenetic analysis of *C. jejuni* and *C. coli* strains from Central Chile. The
phylogenetic tree is based on cgMLST analyses performed with RIDOM SeqSphere⁺ and
visualized by iTOL. Sequence types (STs) are shown in colored boxes for each of the
Campylobacter strains, and the tree branches have been color coded to highlight the *C.*
jejuni and *C. coli* strains from clades I, II and III. **B)** Global cgMLST analysis and
distribution of clonal complexes of clinical *C. jejuni* and *C. coli* strains. Strains from this
study are highlighted in red. cgMLST analysis performed by SeqSphere+ and visualized
by iTOL.

Figure 2. Visualization of the pangenome of the 69 *C. jejuni* and 12 *C. coli* strains from
Chile. Pangenome analysis was performed using Roary. The core and accessory
genomes are highlighted with red and black brackets, respectively. The cgMLST
phylogenetic tree of Figure 1A is shown on the left. A Venn diagram highlighting the
number of core genes present in *C. jejuni* and *C. coli* strains from clades I and III is shown
on the right.

Figure 3. Distribution of antimicrobial resistance genes in *C. jejuni* and *C. coli* strains from
Chile. Binary heatmaps show the presence and absence of antimicrobial resistance
genes. Colored cells represent the presence of genes, and blank cells represent the
absence of genes. The phylogeny of *C. jejuni* and *C. coli* strains is the same as that in
Figure 1. Strain names are color coded to highlight phylogenetic groups: *C. jejuni* strains
(light green), *C. coli* clade I strains (orange) and *C. coli* clade III strains (magenta).

Figure 4. Distribution of virulence-related genes in *C. jejuni and C. coli* strains. Binary
heatmaps show the presence and absence of virulence genes linked to bacterial adhesion
(red), invasion (blue), toxins (purple) and specialized protein secretion systems (green).
Colored cells represent the presence of genes, and blank cells represent the absence of
genes. The phylogeny of *C. jejuni* and *C. coli* strains is the same as that in Figure 1. Strain
names are color coded to highlight *C. jejuni* strains (green), *C. coli* clade I strains (orange)
and *C. coli* clade III strains (purple).

Figure 5. Comparative genomic analysis of the T6SS and T4SS gene clusters identified
in clinical *C. jejuni* and *C. coli* strains. **A)** Genetic organization of the T6SS gene clusters
of *C. jejuni* and *C. coli* strains (bold) compared to the cluster of *C. jejuni* strain 108. The
genes *tagH*, *tssD*, *tssB* and *tssC* are shown in color. **B)** Genetic organization of the T4SS
gene clusters identified in clinical *C. jejuni* strains in comparison to the T4SS gene clusters
of the plasmids pTet and pCC31. The *cp33* gene is highlighted in orange. BLASTn
alignments were performed and visualized using EasyFig.
787
Figure 6. Comparative genomic analysis of plasmids encoding T6SS **(A)** and T4SS **(B)**
gene clusters in clinical *C. jejuni* strains. Pangenome BLAST analysis of plasmids was
performed using the Gview server. T6SS and T4SS gene clusters are highlighted in red
brackets.

Figure S1. Distribution of the presence (red) and absence (blue) of virulence-related
genes in *C. jejuni* and *C. coli* strains from the study.

Core *C. jejuni* genes   Accessory *C. jejuni* genes

C. jejuni strains

C. coli strains

Core *C. coli*   +   Core clade 1 genes

Core *C. coli*   +   Core clade 3 genes

Roary matrix (6609 gene clusters)

C. jejuni

C. coli Clade I

737   23   136

8   675

C. coli Clade III

putative lipase
hypothetical protein
hypothetical protein
putative TraG protein
hypothetical protein
putative nuclease
hypothetical protein
hypothetical protein
hypothetical protein
hypothetical protein
hypothetical protein
hypothetical protein
hypothetical protein
putative phage integrase
hypothetical protein
putative nuclease
hypothetical protein
Type VI secretion system protein TssF
hypothetical protein
DotU domain protein
putative IcmF protein
putative IcmF protein
VasC protein
hypothetical protein
hypothetical protein
hypothetical protein
hypothetical protein
putative nuclease

T6SS
gene cluster

0 kbp, 5 kbp, 10 kbp, 15 kbp, 20 kbp, 25 kbp, 30 kbp, 35 kbp, 40 kbp, 45 kbp, 50 kbp, 55 kbp, 60 kbp, 65 kbp, 70 kbp, 75 kbp, 80 kbp

pCFSAN093246
pMTVDDCj13-2
Pangenome

tetO
cpp3
repA
cpp7
cpp9
cpp11
cpp13
cpp14
cpp16
cpp17
cpp17
cpp18
cpp21
cpp22
cpp23
cpp26
vapD
cmgB3/4
ssb1
cmgB6
cmgB8
cmgB9
cmgB10
cmgB11
cmgD4
cmgD4
cmgD4
cpp45
cpp46
cpp47
cpp49
cpp49
cpp49
cpp50
cpp16

T4SS
gene cluster

0 kbp, 5 kbp, 10 kbp, 15 kbp, 20 kbp, 25 kbp, 30 kbp, 35 kbp, 40 kbp, 45 kbp

pCFSAN093226
pCFSAN096297
pCC31
pTet
Pangenome