Genomic resources and toolkits for developmental study of whip
spiders (Amblypygi) provide insights into arachnid genome evolution
and antenniform leg patterning

Guilherme Gainett[1], Prashant P. Sharma[1]

[1]Department of Integrative Biology, University of Wisconsin-Madison, Madison, WI, USA

53706

Correspondence: guilherme.gainett@wisc.edu

Abstract

Background

The resurgence of interest in the comparative developmental study of chelicerates has led to

important insights, such as the discovery of a genome duplication shared by spiders and

scorpions, inferred to have occurred at the most recent common ancestor of

Arachnopulmonata (a clade comprised of the five book lung-bearing arachnid orders).

Nonetheless, several arachnid groups remain understudied in the context of development and

genomics, such as the order Amblypygi (whip spiders). The phylogenetic position of

Amblypygi in Arachnopulmonata posits them as an interesting group to test the incidence of

the proposed genome duplication in the common ancestor of Arachnopulmonata, as well as

the degree of retention of duplicates over 450 Myr. Moreover, whip spiders have their first

pair of walking legs elongated and modified into sensory appendages (a convergence with the

antenna of mandibulates), but the genetic patterning of these antenniform legs has never been

investigated.

Results

We established genomic resources and protocols for cultivation of embryos and gene

expression assays by in situ hybridization to study the development of the whip spider

Phrynus marginemaculatus. Using embryonic transcriptomes from three species of

Amblypygi, we show that the ancestral whip spider exhibited duplications of all ten Hox

genes. We deploy these resources to show that paralogs of the leg gap genes *dachshund* and

homothorax retain arachnopulmonate-specific expression patterns in *P. marginemaculatus*.

We characterize the expression of leg gap genes *Distal-less*, *dachshund-1/2* and *homothorax-*

1/2 in the embryonic antenniform leg and other appendages, and provide evidence that

allometry, and by extension the antenniform leg fate, is specified early in embryogenesis.

Conclusion

This study is the first step in establishing *P. marginemaculatus* as a model for modern

chelicerate evolutionary developmental study, and provides the first resources sampling whip

spiders for comparative genomics. Our results suggest that Amblypygi share a genome

duplication with spiders and scorpions, and set up a framework to study the genetic

specification of antenniform legs. Future efforts to study whip spider development must

emphasize the development of tools for functional experiments in *P. marginemaculatus*.

Keywords: Arachnida | Arachnopulmonata | gene duplication | paralogs | leg gap genes |

sensory biology

Background

In the past 25 years, comparative developmental study of chelicerates has resurged with the

integration of molecular biology and genomics, spearheaded by developmental genetic

investigations of the spiders *Cupiennius salei* and *Parasteatoda tepidariorum* [1-6]. Such

works have revealed several important aspects of chelicerate development, such as the

molecular mechanism of dorso-ventral axis patterning [3,7-9], segmentation of the prosoma

[1,10-13] and opisthosoma [2,13], and specification of prosomal versus opisthosomal fate

[14,15]. A few non-spider chelicerate models have also emerged in recent years and provided

new perspectives on chelicerate development, such as the horseshoe crab *Limulus*

polyphemus (Xiphosura) [16,17] , the mites *Archegozetes longisetosus* [18-21] and

Tetranychus urticae (Acariformes) [22-24], the tick *Rhipicephalus microplus* (Parasitiformes)

[25], the Arizona bark scorpion *Centruroides sculpturatus* (Scorpiones) [26] and the

harvestmen *Phalangium opilio* (Opiliones) [27-30]. Studies in *A. longisetosus* and *C.*

sculpturatus have suggested that changes in Hox gene number and expression domains are

responsible for such phenomena as the reduced segmentation of mites and the supernumerary

posterior appendage identities of scorpions [18,26].

In addition to the insights into chelicerate development, one of the major recent

outcomes of increasing availability of genomic resources in in the group was the discovery of

a whole or partial genome duplication in spiders and scorpions. This genome duplication is

inferred to trace back to the most recent common ancestor of the recently proposed clade

Arachnopulmonata, which includes spiders, scorpions, and three other arachnid orders with

book lungs [6,26,31-33]. This duplication event is also inferred to be independent from the

multiple rounds of whole genome duplication undergone by horseshoe crabs (Xiphosura)

[34,35]. The duplication in spiders and scorpions encompasses several important

developmental genes, such as the homeobox family [6,32]. Excitingly, there is increasing

evidence of divergent expression and function of paralogs, such as in the case of Hox genes,

leg gap genes and Retinal Determination Gene Network homologs [6,26,36-40].

Nevertheless, knowledge on the development of several arachnid orders still remains scarce

or entirely unexplored in the contexts of embryological study and genomic architecture. One

particularly interesting arachnid order whose genomic evolution and developmental biology

remains largely unexplored is Amblypygi. Commonly known as whip spiders, Amblypygi

comprise approximately 220 described species of nocturnal predators that are distributed

primarily in the tropics and subtropics worldwide [41,42]. Amblypygi is part of the clade

Pedipalpi (together with the orders Thelyphonida and Schizomida), which in turn is sister

group to spiders (Araneae) [31,43-45]. Considering the phylogenetic position of Amblypygi

within Arachnopulmonata, a better understanding of Amblypygi genomics is anticipated to

facilitate exploration of the evolutionary outcomes of the genome duplication in

Arachnopulmonata, and better characterizing the extent to which arachnopulmonate orders

retained and specialized the ensuing paralogous genes. Unfortunately, the embryology of the

order Amblypygi is known only from the works of Pereyaslawzewa [46], a brief mention in

Strubell [47] ("Phrynidae"), Gough [48], and the comprehensive study of Weygoldt [49] in

Phrynus marginemaculatus (formerly *Tarantula marginemaculata*).

Beyond the interest in their genomic architecture, whip spiders possess a fascinating biology

and natural history. Whip spiders are emerging as model organisms in behavioral ecology,

learning, and neurophysiology in arachnids [50,51]. Their pedipalps, the second pair of

appendages, are robust raptorial devices used for striking prey, including mostly arthropods,

and even small vertebrates [41,50,52]. However, their most conspicuous characteristic and

namesake is their "whip", a modified antenniform first walking leg that is not used for

locomotion, but instead as a sensory appendage. The antenniform legs are extremely

elongated relative to the other leg pairs, and the tibia and tarsus are pseudo-segmented into

hundreds of articles that harbor an array of chemo-, thermo-, hygro- and mechanoreceptive

sensilla [53-55]. The peripheral circuitry of the antenniform legs is complex, exhibiting

peripheral synapses and giant neurons which are possibly involved in fast sensory responses

[55-57]. The antenniform legs also have been shown to be important for foraging, mating,

and intrasexual contests [58,59]. Notably, concentration of sensory structures, elongation and

pseudo-segmentation of these legs constitute a striking convergence with the antenna of

mandibulate arthropods (Pancrustacea and Myriapoda). Evidence from Hox gene expression

and functional experiments support the view that the antenna of mandibulates is positionally

homologous to the chelicera of chelicerates (deutocerebral appendage) [21,30]. Despite their

different positions along the antero-posterior axis of the body, the serial homology of

antennae (mandibulates) and antenniform legs (whip spiders) to walking legs constitutes a

potentially useful comparison to address whether a striking morphological convergence is

associated with convergence in mechanisms of genetic patterning.

117   Most of the knowledge about the genetics of antenna versus leg fate specification in

arthropods was discovered in the fruit fly *Drosophila melanogaster* and involves the

regulation by leg gap genes and Hox genes. Broadly overlapping domains of leg gap genes

homothorax (*hth*), *dachshund* (*dac*) and *Distal-less* (*Dll*) in the antenna promote antennal fate

through activation of downstream antennal determinants such as *spineless* [60-63] (reviewed

by [64]); leg identity in thoracic appendages is mediated by mutually antagonistic *hth*

(proximal), *dac* (median) and *Dll* (distal) domains, and repression of *hth* by the Hox gene

Antennapedia (*Antp*) [62,65,66]. Ectopic expression of *hth* on the leg discs results in leg-to-

antenna transformation, whereas ectopic *Antp* in the antennal disc results in antenna-to-leg

transformation [62,65,66]. In chelicerates, similar to *D. melanogaster*, the chelicera (antennal

positional homolog) has broadly overlapping domains of *Dll* and *hth*; legs and pedipalps lack

a distal *hth*, and a proximal *Dll* expression domain, indicating a conservation of expression of

positional homologs [28,67-69]. In line with this evidence, knockdown of *hth* in the

harvestmen *Phalangium opilio* results in chelicera-to-leg transformation (and also partial

pedipalp-to-leg transformation) [30]. In contrast to *D. melanogaster* and other insects,

arachnid *Antp* is not expressed in the leg-bearing segments, and knockdown of this gene in

the spider *Parasteatoda tepidariorum* has revealed a role in repressing limbs in the

opisthosoma [14]. Expression of *dac* in a mid-domain of chelicerate pedipalps and legs

suggests a conserved role in mid segment patterning, and, accordingly, knockdown of *dac* in

the harvestmen *P. opilio* and the spider *P. tepidariorum* results in defects on the mid-leg

segments [27,38]. Additionally, loss of a *dac* domain of the chelicera of spiders in

comparison to harvestman has been implicated in the transition from three- to two-segmented

chelicera [27,28].

Towards revitalizing embryological studies in Amblypygi, we generated herein protocols and

comprehensive genomic resources for developmental genetic study of the whip spider

Phrynus marginemaculatus (Fig. 1A–B), resuming the seminal studies begun by Peter

Weygoldt with this promising model species. Leveraging a comprehensive developmental

transcriptome for this species, we first investigated the presence and retention of systemic

gene duplications in Amblypygi predicted from their phylogenetic position in

Arachnopulmonata; to mitigate biases stemming from survey of a data point from a single

exemplar, we included as well the embryonic transcriptomes we recently generated for two

species of the distantly related family Charinidae (*Charinus ioanniticus* and *Charinus*

israelensis) [40,41,70]. As a first step towards elucidating the patterning of antenniform legs,

we investigated a possible role of the leg gap genes *homothorax*, *dachshund* and *Distal-less*

in specifying antenniform leg fate, by characterizing their expression pattern in developing

embryonic stages of *P. marginemaculatus* using newly established procedures for

colorimetric whole mount in situ hybridization.

Results

Overview of embryogenesis in Amblypygi

A characterization of the postembryonic development of *P. marginemaculatus* is available in

the works of Weygoldt [71,72]. The following summary is after Weygoldt [49],

complemented with observations from the present study. We note that the timing of

embryonic stages is approximate, as the temperature in a previous work [49] was kept in a

range of 24–27˚C. Details on the culture and collection of embryos are described in *Methods*.

In our colony, a sample of 18 clutches from different females had an average of 19

eggs (min: 8 eggs; max: 39 eggs). Each egg is ovoid and measures 1.2–1.4 mm at the major

axis. The yolk granules are light green, and fixation turns them light yellow. Upon secreting

the brood sac and eggs into the ventral side of the opisthosoma, the brood sac is transparent,

but gradually hardens and becomes dark brown in the course of two days. Around 6 dAEL, a

blastoderm is formed. At 7–8 dAEL, the yolk is more condensed, a perivitelline space is

formed, cells move into the yolk presumably in the region of the future blastopore. At 8–9

dAEL a blastopore is recognizable on the dorsal surface of the embryo, and a germ disc is

formed. Towards 10 dAEL, the germ band starts to take form, the blastopore region is

displaced posteriorly, and the first signs of segmentation of the prosoma are visible. A

presumable cumulus, a group of migrating cells with dorso-ventral organizing properties, was

reported by Weygoldt (1975) at this stage, but was not investigated by us. At 11-12 dAEL,

the nascent prosomal segments elongate anteriorly and by 13-14 dAEL the segmented germ

band occupies the ventral hemisphere of the egg. The limb buds of prosomal appendages

begin to appear (Fig. 2A). The process of reversion begins, with a sagittal split of the germ

band (Fig. 2A). From this point, until 19 dAEL, the ventral sulcus progressively enlarges

with the dorsal displacement of both halves, and then begin to ventrally fuse again starting

anteriorly (Fig. 2A–E). The head lobes on 15 dAEL have a circular outline and uniform

ectoderm (Fig. 2F). From 15–19 dAEL, ectodermal cells invaginate along the ventral midline

and head lobes (neural precursor cells), which can be seen as evenly spaced dark dots (Fig.

2F–H). Concomitantly, the anterior rim of the head lobes forms semilunar groves (i.e., folds

over itself) (Fig. 2F–H). Opisthosomal segments 1–12 sequentially appear from the posterior

growth zone (Fig. 2I–K). At 19 dAEL, the yolk moves into the caudal papilla (growth zone)

and the opisthosoma flexes ventrally (Fig. 2K, N). The first pair of legs (future antenniform

legs) is already longer than the other appendages in an early limb bud stage (15 dAEL), and

more clearly in subsequent stages (Fig. 2L–N). A protuberance on the proximal end of the

second walking leg pair (L2), the nascent lateral organ (osmoregulatory embryonic organ), is

discernible at 16 dAEL and later stages, and becomes reniform (Fig. 2L–N). From 20–21

dAEL, the embryo begins to secrete a cuticle, which is darker on the fully formed lateral

organ. The head lobes have completely folded onto themselves, and the distal segment of the

chelicera is sclerotized. The embryo then ruptures the egg shell, and is now termed

deutembryo (Fig. 2O–Q). Deutembryos remain in the brood sac for further 70 days without

discernible external changes. The primordia of the eyes are formed towards the end of

deutembryo development (70–106 dAEL), and eye spots of the median and lateral eyes

become visible. From 90-106 dAEL the deutembryos molt into praenymphs, leave the brood

sac and climb onto the adult female's back (Fig. 1B). For a histological description of

gastrulation, of the mesoderm, brain and nervous system, digestive system and gonad

formation, we refer the reader to the exquisite original descriptions of Weygoldt [49].

Phrynus marginemaculatus embryonic transcriptome

In order to provide a comprehensive resource for Amblypygi to facilitate

developmental genetic study and comparative genomics, we assembled an embryonic

transcriptome spanning different embryonic stages. The assembly of *P. marginemaculatus*

transcriptome resulted in 544.372 transcripts composed of 277.432.101 bp and N50 of 440

bp. BUSCO scores, which reflect the presence and number of copies of widely conserved

single-copy genes (BUSCO-Arthropoda; n=1066), indicated 94.2% completeness with 5.9%

of genes duplicated. From the total number of transcripts (including isoforms), we predicted

37,540 peptide sequences with the Trinotate pipeline with a top blastp hit in Swiss-Prot

database. A total of 801 of these peptides (2.1%) had a top hit to non-metazoan proteins, and

presumed microbial contaminants. The total number of predicted proteins is higher in

comparison with the predicted protein set from the genome of the spider *Parasteatoda*

tepidariorum (27,900) and the scorpion *Centruroides sculpturatus* (30,456) [6], which is

attributable to the typical fragmentation of genes in such transcriptomic assemblies.

To investigate the possibility of systemic gene duplication in Amblypygi and further

test the transcriptome completeness, we searched for all ten Hox genes, emphasizing

discovery of paralogs known from the genomes of the spider *Parasteatoda tepidariorum*

(except for *fushi tarazu*) and the scorpion *Centruroides sculpturatus* (except for *Hox3*)

[6,26,32]. One of the *Hox3* copies of *P. tepidariorum* has a highly divergent homeodomain

sequence, constituting an unusual long branch in our phylogenetic analysis (Additional file 1,

Fig. S1). The same is true for the *Hox3* ortholog *bicoid* of *Drosophila melanogaster*

(Additional file 1, Fig. S1). We therefore removed both paralogs from our main analysis. We

found orthologs of all ten Hox genes in *Phrynus marginemaculatus*, namely *labial*,

proboscipedia, *Hox3*, *Deformed*, *Sex combs reduced*, *fushi tarazu*, *Antennapedia*,

Ultrabithorax, *abdominal A* and *Abdominal B* (Fig. 3A–B). We found evidence that all Hox

genes, except *proboscipedia* are assembled in two genes, and the predicted proteins

sequences overlap and have amino acid differences in *P. marginemaculatus* transcriptome

(Fig. 3B). A *Pmar-Dfd* and *Pmar-abdA* were assembled in a third gene, but their non-

overlapping sequences on the alignment and nearly identical amino acid sequence to the

respective paralog 2 indicate that they are fragmented assemblies of the same copy. We

additionally analyzed two embryonic transcriptomes of *Charinus* whip spiders recently

generated by us [40]. For *C. israelensis*, we discovered orthologs of all ten Hox genes, with

two genes for all cases except *ftz* and *Antp* (Fig. 3A–B). A third gene was annotated as *Cisr-

Dfd*, but we interpret it as an assembly error, given its largely non-overlapping sequence with

respect to the other *Cisr-Dfd* sequences. For *C. ioanniticus*, we discovered one gene each for

ftz, *Antp*, *Ubx*, *AbdB*, and two genes each for *Hox3*, *Dfd*, and *Scr*. We did not detect orthologs

of *lab*, *pb* and *AbdB* in *C. ioanniticus* transcriptome (Fig. 3A–B).

These absences almost certainly reflect incompleteness of transcriptomic assemblies,

and cannot be equated with true losses. We therefore consider the union of the Hox gene

counts across the three species as the reconstruction of the whip spider ancestral Hox

complement. Our results suggest that the ancestral Hox complement in Amblypygi consists

of 20 genes (two copies of every Hox gene), a trait they share with spiders and scorpions.

While a genome is not available for whip spiders, the phylogenetic placement of this order, as

well as the pattern of Hox duplication, strongly suggests that whip spiders share the condition

of two Hox clusters observed in spider and scorpion genomes.

Gene expression analyses in *Phrynus marginemaculatus*

All three methods we trialled for fixing embryos of *P. marginemaculatus* yielded embryos

which are suitable for in situ hybridization (ISH). However, fixation of intact embryos in

formaldehyde/heptane phase sometimes results in the vitelline membrane adhering to the

embryo. Upon dissection of the vitelline membrane for ISH, parts of the embryo are

commonly lost. Piercing a hole in the embryo for fixation in formaldehyde achieves a similar

result, and is therefore not recommended. Dissecting the vitelline membrane under PBS and

immediately fixing the embryos in formaldehyde yielded the best results. We did not detect

any difference in the signal-to-background ratio of colorimetric ISH between embryos fixed

for 2 h or 24 h. To test gene expression assays and investigate the genetic patterning of

antenniform legs, we selected the leg gap genes *Distal-less*, *dachshund* and *homothorax*,

which constitute the subset of PD axis patterning genes associated with functional datasets in

other arachnid model species [27,30,73].

Distal-less identification and expression

Similar to other arachnopulmonates and arthropods in general, a single *Distal-less* gene,

Pmar-Dll, was found by similarity searches and its orthology to other metazoan *Distal-less*

genes was confirmed with a phylogenetic analysis (Additional file 2, Fig. S2). *Pmar-Dll* is

strongly expressed in all developing appendages, from early limb bud stage (15 dAEL) and is

localized to the distal portion of the chelicera, pedipalp and legs throughout leg elongation

(Fig. 4A–I; Fig. 5A–L). In 17-18 dAEL embryos, the expression is heterogeneous and forms

bands of stronger expression on the forming podomeres of the pedipalps and legs (Fig. 5H–

L). We did not detect expression on ventral midline (*contra* [28] and[16]), or in the

opisthosoma, and observed only diffuse expression on the head lobes of the stages surveyed.

Expression on L1 of 16-17 dAEL embryos is subtly stronger on the proximal part of the

expression domain (Fig. 5C–C'), but we did not detect clear differences in younger or older

embryonic stages. The proximal and distal boundaries of strong *Distal-less* expression do not

differ between L1 and the other legs, spanning two-thirds of the legs from the tip of the tarsus

and excluding the most proximal third (Fig. 5C, 5I).

In contrast to other arachnid embryos (mygalomorph spiders [69]; araneomorph spiders

[67,73]; Opiliones, Sharma et al. 2012; and mites [20,74]), *Dll* expression was not detected as

a strong domain in the endite of pedipalp, which is very inconspicuous at these embryonic

stages. We were not able to assess *Dll* expression in the pedipalpal endite in deutembryo

stages due to deposition of cuticle.

dachshund identification and expression

dachshund is typically present as a single copy in Arthropoda, but is found independently

duplicated in Arachnopulmonata and Xiphosura [38,75]. We found two genes with high

similarity to *Dmel-dac* and confirmed their identity as dachshund paralogs with a

phylogenetic analysis (Additional file 3, Fig. S3). *Pmar-dac1* and *Pmar-dac2* are each nested

in a clade with paralogs 1 and 2 of other arachnopulmonates, which appears as independent

from horseshoe crab duplications (Additional file 3, Fig. S3).

Pmar-dac1 is strongly expressed as broad median band in the pedipalp and the four pairs of

legs (Fig. 6A–F). This sharp mid band excludes the most proximal and the distal domains of

these appendages (Fig. 7). No staining was detected on the chelicera (Fig. 6A–F; 7A, E).

Additionally, expression occurs in three domains in the head lobes and strongly on the ventral

midline of the prosoma. The continuous expression on the ventral midline stops before the

opisthosoma (Fig. 6A–F).

Pmar-dac2 is expressed as ring at the proximal region of chelicera, pedipalp and legs (Fig.

6G–I). On 16-17 dAEL embryos, a second thinner ring of expression is also present on the

pedipalps and legs (Fig. 6G–I; 7I–L). Expression domains also occur on the neuromere of L1

and pedipalp, being larger on the pedipalpal neuromere (Fig. 6G). Three spot domains also

occur on O1, O2 and O3 segments (Fig. 6H–I).

homothorax identification and expression

Similar to the case with arachnid *dachshund*, *homothorax* is also found duplicated in

arachnopulmonates [32,39]. We annotated two genes in *P. marginemaculatus* transcriptome

and the phylogenetic analysis shows that each copy is nested in two clades with the

arachnopulmonate paralogs (Additional file 4, Fig. S4).

Pmar-hth1 is conspicuously expressed on the developing appendages throughout the stages

studied (15-18 dAEL; Fig. 8–9). On 16-17 dAEL embryos, *Pmar-hth1* is expressed more

strongly on the proximal parts of the chelicera, pedipalp, and legs, and is absent in a small

distal domain (Fig. 9A–D). On 17-18 dAEL, expression in all elongated appendages is

similar, and spans most of the appendage length except for a small distal territory (Fig. 9E-

H). Additionally, three stronger stripe domains occur distal to the coxa of pedipalp and legs.

Expression in L1 and other legs is similar in the studied stages, save for a slightly expanded

distal domain on the proximal tarsus of L1 (Fig. 9G–H). Expression on the ventral ectoderm

and head lobes is weak and homogenous in early stages, and becomes stronger in 17-18

dAEL (not shown)

Pmar-hth2 is expressed in the developing appendages, but is reduced in intensity and

different in pattern in comparison to *Pmar-hth1* (Fig. 8B–B'). *Pmar-hth2* spans the whole

chelicera (Fig. 9I). Expression on the pedipalp spans most of the appendage, excluding only a

distal territory (Fig. 9J). In addition, there is a proximal and a sub terminal strong domain

(Fig. 9J). Expression on L1 and other legs is identical in all stages studies. *Pmar-hth2*

expression on the legs is consistently weaker in comparison to the pedipalp, but likewise

possess a proximal and a distal sub-terminal domain (Fig. 9K–L). Expression on the ventral

ectoderm, opisthosoma and head lobes is ubiquitous (Fig. 8B).

Discussion

Genomic resources for whip spiders corroborate widespread retention of duplicates across

Arachnopulmonata

Changes in Hox gene expression, regulation and targets are linked to changes in

morphology and Hox gene duplications have also been implicated in morphological

innovations [76-79]. Duplication in the Hox cluster is common in the vertebrates, which may

have up to eight Hox clusters (teleost fishes) due to successive rounds of genome

duplications [76]. On the other hand, genome and Hox cluster duplications are considered

rare in Arthropoda [80]. Complete Hox cluster duplication in the phylum has only been

reported in horseshoe crabs (Xiphosura) [34,35,81,82], and in spiders and scorpions

(presumed ancestral in Arachnopulmonata). The phylogenetic placement of Amblypygi

makes this group an opportune system to test the downstream prediction that *P.*

marginemaculatus retains and expresses Hox paralogs shared by spiders and scorpions. In

accordance with this hypothesis, we discovered that the whip spider *P. marginemaculatus*

presents homologs of all ten Hox genes, and all excepting *proboscipedia* appear to be

duplicated, as supported by our alignment and phylogenetic analysis. Our survey in the two

Charinus transcriptomes confirmed the duplication of several Hox genes, and additionally

revealed that two copies of *proboscipedia* (absent in *P. marginemaculatus*) are present at

least in one species. Our results suggest that Amblypygi retain a complete duplicated Hox

complement with 20 genes. Similarly, a recent transcriptomic survey in *Charinus acosta* and

Euphrynichus bacillifer reported duplications of nearly all Hox genes (excepting

proboscipedia), as well as Wnt and Frizzled genes [83]. Taken together with the recent

discovery of duplicated RDGN genes in *Charinus*, our results suggest that Amblypygi retain

ohnologs of numerous developmental patterning genes, as a function of an ancient, shared

whole genome duplication.

A second line of evidence of a shared genome duplication event comes from conserved

expression patterns of leg gap gene paralogs. It has recently been shown that duplicates of leg

patterning genes *extradenticle*, *optomotor blind*, *dachshund* (*dac*) and *homothorax* (*hth*)

present one copy with an expression pattern resembling a stereotypical panarthropod domain,

and a second copy with a divergent expression, with these two sets of expression patterns

shared by a spider and a scorpion [75]. For instance, *dac1* of surveyed arachnopulmonates

retains the plesiomorphic expression pattern of a broad median band in the appendages, the

condition observed in the single-copy *dac* of all non-arachnopulmonate arachnids surveyed to

date [20,28], whereas *dac2* is expressed as a proximal and a medial domain at the patella-

tibia boundary [38,67,69,75]. Similarly, *hth1* retains a plesiomorphic expression pattern on

the proximal-median appendage excluding the tarsus, whereas *hth2* presents a strong short

proximal domain and stripes of expression at podomere boundaries [39,67,69,75]. In the

whip spider *P. marginemaculatus*, we indeed recovered two copies of *dachshund* and

homothorax, with paralogs nested in two clades of arachnopulmonate copies with high nodal

support (Additional file 3, Fig. S3; Additional file 4, Fig. S4). Moreover, *Pmar-dac* and

Pmar-hth have paralog-specific expression patterns shared with spiders and a scorpion:

Pmar-dac1/2 and *Pmar-hth1/2* conserve the plesiomorphic expression pattern of arachnids,

while copy 2 retains the arachnopulmonate-specific pattern. These data, together with the

evidence from Hox gene duplications and the topologies of the *dac* and *hth* gene trees,

provide compelling support for genome duplication as a synapomorphy of

Arachnopulmonata.

A *dachshund2* domain in the two-segmented clasp-knife chelicera of *P. marginemaculatus*

Expression of *Pmar-Dll*, *Pmar-dac1/2* and *Pmar-hth1/2* on the appendages is largely similar

to spiders and scorpions (see above) [38,39,67-69,75]. A notable difference in the chelicera

of the whip spider is the retention of proximal expression of one of the *dachshund* paralogs.

In arachnids with three segmented chelicerae, such as the daddy-long-legs *Phalangium*

opilio, *dac* is expressed in the proximal segment of the appendage [28]. The same is true for

the three-segmented chelicera of scorpions, in which both *dac1* and *dac2* are expressed in the

proximal segment [75]. In the two-segmented chelicera of spiders, *dac1* is not expressed in

the chelicera save for a non-ectodermal faint staining in late embryonic stages [38,67-69,84].

dac2 of spiders is also largely not expressed in the chelicera, save for a small ventral domain

at the base of the chelicera of *Parasteatoda tepidariorum* and *Pholcus phalangioides* [38]. It

has been proposed that the loss of a proximal *dac* domain is associated with the loss of the

proximal segment in the chelicera of spider and other arachnids. This hypothesis is supported

by functional data in the daddy-long-leg *P. opilio*: knockdown of the single-copy *dac* resulted

in the loss of the proximal segment of the chelicera (as well as mid-segment truncations in

pedipalps and legs) [27]. Similar to spiders, the two-segmented chelicera of the whip spider

P. marginemaculatus does not express *Pmar-dac1*. Nonetheless, a clear complete ring of

proximal expression was detected for *Pmar-dac2*.

In contrast to the classic leg gap phenotype incurred by knockdown of the ancestral *dac* copy

in the harvestman, knockdown of *dac2* in the spider *P. tepidari*orum resulted in the loss of

the patella-tibia segment boundary in the pedipalp and walking legs; effects of this

knockdown on the chelicera were not reported by the authors, nor is the function of *dac1*

known in *P. tepidariorum* [38]. Given the interpretation that *dac2* has a derived role in

patterning a segment boundary in the spider, the incidence of the *Pmar-dac2* stripe in the

middle of the whip spider chelicera (and the absence of *Pmar-dac1* in the chelicera) is

tentatively hypothesized here to constitute evidence of a segment boundary patterning

activity that separates the basal and distal cheliceral segment of Amblypygi (rather than

evidence of a proximal segment identity in whip spiders that is homologous to the proximal

cheliceral segment of groups like harvestmen and scorpions). Exploring this hypothesis in

future will require the establishment of RNAinterference tools in *P. marginemaculatus.*

Antenniform leg distal allometry is specified early in development

To investigate a possible role of changes in leg gap genes expression in the patterning

of the antenniform leg pair (L1) of whip spiders, we specifically searched for changes in

expression domain between embryonic antenniform leg (L1) and the posterior legs (L2–4).

The expression patterns of *Pmar-Dll*, *Pmar-dac1/2* and *Pmar-hth1/2* are mostly identical

between the antenniform leg L1 and the posterior legs. We found only a slightly expanded

Pmar-hth1 domain in the distal leg (compare Fig. 9G–H). It is unclear if this difference bears

any functional significance in the differential fate of L1. Similar distal expression dynamics

appear to distinguish the identity of the pedipalp and walking legs of the harvestman, as

knockdown of the single copy of *hth* in *P. opilio* results in homeotic transformation of

pedipalps to legs [30].

Besides the difference in sensory equipment, the antenniform leg pair of adults differ from

the posterior walking legs in being much longer, and presenting numerous articles (pseudo-

segments) on the tibia, metatarsus and especially the tarsus, whereas the walking legs have

pseudo-segments only in the tarsus [41,53,55]. Antenniform legs may be more than 2.5 times

longer than the walking legs, and this allometry in adults is mostly accounted for by the

length added by the distal leg segments, which may attain more than 100 distal articles

(tibiomeres + tarsomeres) [53]. Understanding the timing of the ontogenetic divergence of L1

relative to the walking legs is crucial for circumscribing target stages for investigation of the

genetic patterning of this phenomenon.

Current evidence indicates that this allometry on the antenniform legs takes place both

embryonically and postembryonically. The praenymph, the stage that molts from the

deutembryo and climbs to the mother's back, has the full number of tibiomeres, but not of

tarsomeres. The final number of tarsomeres is added in the next molt to the protonymph

[53,85], and from this moment onwards the subsequent growth is isometric [53,85]. In

embryos of *P. marginemaculatus*, Weygoldt [49] observed that the primordia of L1 are the

first to appear in the germ band and are already longer than the remaining prosomal

appendages, an observation paralleling the allometric growth of the L2 limb bud of the

harvestman *P. opilio* (wherein L2 is the longest leg and also bears the most tarsomeres) [29].

It is presently unclear at what point in development the distal podomeres of antenniform legs

attain their allometry with respect to the proximal podomeres in the whip spider. In other

words, are embryonic legs simply growing longer overall with respect to walking legs, or are

they longer due to distal growth? The expression pattern *dachshund* in *P. marginemaculatus*

early embryos reveals a further interesting aspect of this allometry. The expression of *dac*

(including duplicates) in the legs of arachnids surveyed to date occurs on podomeres

proximal to the tibia, so that the tibia-metatarsus-tarsus territory may be identified as the a

dac-free domain [20,27,38,67,69,75,84]. The distal boundary of *Pmar-dac1* and *Pmar-dac2*

in stages long before podomere formation reveals that the distal domain (*Pmar-dac1*-free

cells) of L1 is already longer with respect to the proximal leg and the other legs (compare

Fig. 7C, D). These data support the inference that the distal allometry, and by extension the

antenniform leg fate, are already specified from the onset of limb bud development.

Conclusion

The study of the embryology of Amblypygi has faced 45 years of hiatus since the seminal

work describing the development of *Phrynus marginemaculatus* [49]. Here, we take the first

steps in establishing *P. marginemaculatus* as a model for modern chelicerate evolutionary

developmental study and for comparative genomics. We provide evidence for a shared

genome duplication in Amblypygi; show that the distal allometry of antenniform legs is

specified early in appendage embryogenesis; and observe that expression patterns of the leg

gap genes *Distal-less*, *dachshund* and *homothorax* do not substantian their role in fate

specification of the sensory legs of whip spiders. Future efforts to uncover the genetic

underpinnings of this fascinating convergence with the mandibulate antenna should prioritize

discovery of leg fate specification mechanisms in early whip spider development. Despite the

establishment of a few promising arachnid models for studying chelicerate embryology, a

pervasive challenge has been the establishment of toolkits for testing gene function

[3,22,27,73]. Therefore, future efforts to study whip spider development must emphasize the

development of tools for functional experiments in *P. marginemaculatus*.

Methods

Establishment of amblypygid colony in the laboratory

Adult individuals of *Phrynus marginemaculatus* were purchased from a commercial supplier

collects animals in the Florida Keys (USA). The maintenance of a colony of *Phrynus*

marginemaculatus (Fig. 1A–B) in the laboratory followed guidelines provided is the original

developmental works of Peter Weygoldt [49,71,72] and notes on behavioral studies on this

species [86-88].While most species of Amblypygi have been reported as aggressive and

better kept in separate containers [41], individuals of *P. marginemaculatus* can be kept in a

stable laboratory colony in large community containers [86]. A maximum of 20 animals may

be allowed to interact and mate freely in plastic boxes (67.3 x 41.9 x 32.4 cm) with a 2 cm

layer of damp coconut fiber saturated with distilled water. The substrate is continuously

dampened with approximately 500 mL of dH_2O on the substrate every week. All animals are

kept in an acclimatized room at 26-28˚C in a controlled reversed 16L:8D light cycle to

simulate summer months in Florida. To prevent flies from reproducing in this humid

environment, the lid of the container is provided with a 10 x 20 cm fine mesh cloth, which

also allows for air circulation. Each container is furnished with pieces of tree bark, egg

carton, and retreats made from polystyrene foam or florist's foam, to provide enough space

for animals to hide and suspend themselves during molting and egg-laying. After about two

weeks of setting up a terrarium, the substrate hosted a stable colony of collembolans

(possibly brought by the tree bark). The co-occurrence of collembolans does not appear to be

disadvantageous in any way. Collembolans are invariably observed feasting on carcasses and

likely contribute to preventing mite infestations (see also [41]). Animals are fed at least twice

a week with nymphs of *Acheta domestica ad libitum.* Carcasses are removed as soon as

detected to prevent the spread of mites. Freshly hatched praenymphs are kept with the

female, and aggregate on the back of the brooding female (Fig. 1B). The next instar, the

protonymph, disperse from the female upon molting. Protonymphs are transferred to

community terraria as described above. Protonymphs and subsequent instars are fed pinheads

of *A. domestica* and *Gryllus bimaculatus ad libitum* three times per week until adulthood.

The first adult individuals from our laboratory-bred animals were detected 8 months after

hatching, which accords to what has been described by Weygoldt [49].

Terraria with adults are inspected daily for freshly molten individuals or ovigerous females,

which are immediately separated from the main colony to avoid aggressive interactions.

Ovigerous females carry the eggs outside the body in a brood sac attached to the ventral side

of the opisthosoma [41]. Eggs are allowed to develop with the female until the desired stage

for developmental work. Approximate stages may be inferred from days after egg laying

(dAEL) at 26°C following the staging in [49]. For obtaining the eggs, females are

anesthetized with CO_2 and immobilized in customized plastic petri dishes (35 x 11 mm) with

a central whole and foam (Fig. 10A–B). With a blunt forceps pair, the pleura of the

opisthosoma embracing the eggs may be gently displaced from the brood sac, and the brood

sac removed as whole (Fig. 10C). The female may be returned to the colony unharmed. The

brood sac easily dissolves when immersed in a phosphate buffer solution (PBS), and eggs

may be cleaned with forceps under PBS. We obtained high mortality in keeping isolated eggs

developing in petri dishes with humid (wet filter paper) or dry in 26˚C incubators, as mold

frequently develops. However, we found out that embryos will survive and develop

immersed in PBS supplemented with kanamycin (1:5000) at 26˚C incubator and even hatch

into deutembryos (~20 dAEL).

Phrynus marginemaculatus transcriptome assembly

We collected embryos of different embryonic stages as described above, and conducted total

RNA extractions of each stage separately: hatched deutembryo (female #5, n=3;

undetermined age, before eyespots), unhatched deutembryo (female #2 n=3; approx. 21

dAEL), and leg elongation stage (female #3, n=2; approx. 16-17 dAEL), late limb bud stage

(female 6, n=1; 16 dAEL) and early limb bud stage (female 6, n=1; 14 dAEL) (voucher

images are available upon request). Embryos were homogenized in Trizol TRIreagent

(Invitrogen) and stored at –80˚C until extraction. The combined total RNA of the five

extractions was submitted to cDNA library preparation and sequencing in an Illumina High-

Seq platform using paired-end short read sequencing strategy (2×100 bp) at the Center for

Gene Expression, Biotechnology Center of University of Wisconsin-Madison (USA). The

quality of the raw reads was assessed with FastQC (Babraham Bioinformatics). De novo

assembly of compiled paired-end reads was conducted with the software Trinity v. 2.6.6 [89],

using Trimmomatic v. 0.36 [90] to remove low quality reads and adaptors. Completeness of

the transcriptome was assessed with the Trinity script '*TrinityStats.pl*', and BUSCO v. 3 [91]

on the longest isoforms using the database 'Arthropoda' and analysis of Hox genes. Whole

transcriptome annotation was conducted following the Trinotate pipeline, by predicting

proteins with TransDecoder v. 5.2.0 [92] and searching best hits with blastx and blastp

against Swiss-Prot and Pfam databases [93].

Gene identification

Hox genes (*labial*, *proboscipedia*, *Hox3*, *Deformed*, *Sex combs reduced*, *fushi tarazu*,

Antennapedia, *Ultrabithorax*, *abdominal A* and *Abdominal B*), *Distal-less*, *dachshund*, and

homothorax orthologs were identified from the embryonic transcriptome of *Phrynus*

marginemaculatus using BLAST searches (tblastn) with peptide queries of the homologs of

Drosophila melanogaster and *Parasteatoda tepidariorum*. The best hits were reciprocally

blasted to NCBI database and further assessed with a phylogenetic analysis.

For the phylogenetic analysis of Hox genes, we selected arthropod terminals with available

genomes and well annotated references: the fruit fly *Drosophila melanogaster* (FlyBase), the

beetle *Tribolium castaneum* (GCA_000002335.3) (Insecta), the centipede *Strigamia*

maritima (GCA_000239455.1) (Myriapoda), the tick *Ixodes scapularis* (GCA_002892825.2),

the mite *Tetranychus urticae* (GCA_000239435.1), the spider *Parastetoda tepidarioum*

(GCA_000365465.2) and the scorpion *Centruroides sculpturatus* (GCA_000671375.2)

(Chelicerata) [6,94-97] (Additional file 5, Table S1). For the analysis of *Distal-less*,

dachshund, and *homothorax* we selected terminals used in Nolan et al. [75] and included the

peptide sequence for *P. marginemaculatus* candidates (Additional file 5, Table S1). Peptides

were predicted with ORF Finder (NCBI), TransDecoder v. 5.5.0 [92], or obtained as CDS

when possible from publicly available NCBI database. Peptide sequences were aligned using

Clustal Omega [98] in the software SeaView v. 4 [99]. Gene trees were inferred under

maximum likelihood using IQTREE [100] with automatic model selection (-m TEST) and

1000 ultrafast bootstrap resampling (-bb 1000). Alignments for the Hox gene analysis are

available in Additional files 6–7. Genes referred throughout text refer to the longest isoform

assembled by Trinity in a contig, and annotated by the procedure detailed above. Paralogs

were discerned from gene fragments using as criteria the presence of largely overlapping

sequences with amino acid differences.

Fixation and in situ hybridization

Collecting freshly laid eggs and keeping them in controlled temperature with the female or

incubator (26°C) allows to select desirable stages for fixation. We fixed embryos for in situ

hybridization with three methods: (1) intact eggs in a phase of 4% formaldehyde/PBS and

heptane overnight, after an established spider fixation protocol [3]; (2) piercing a hole in the

vitelline membrane, and immersing eggs in 4% formaldehyde/PBS overnight; (3) dissecting

the embryo out of the vitelline membrane and fixing in 4% formaldehyde/PBS for periods

varying from 2–24h. All three procedures were conducted in a shaking platform at room

temperature (~22 °C). Following the fixation period, embryos were rinsed several times in 1x

PBS/0.1% Tween-20 (Sigma Aldrich) (PBS-T), and gradually dehydrated into pure methanol.

Embryos were then stored in 20 °C freezer until next procedures.

cDNA was synthetized with SuperScriptIII kit (Thermo Fisher) following the manufacturer's

instructions, using the same total RNA used to generate the embryonic transcriptome.

Riboprobe templates for the genes *Pmar-dac1*, *Pmar-dac2*, *Pmar-hth1*, and *Pmar hth-2* in

situ hybridizations were generated as follows: genes were PCR amplified from cDNA using

gene-specific primers designed with Primer3 v. 4.1.0 [101] and complemented with T7 linker

sequences (5′ -ggccgcgg-3′ ; 5′ - cccggggc-3′), and a second PCR using universal T7 primer

to generate T7 polymerase templates specific for sense (control) and anti-sense (signal)

probes (Additional file 8, Table S2). Sense and antisense riboprobes for *Pmar-Dll* were

generated with T7 and T3 polymerases using a plasmid template. For cloning of *Pmar-Dll*,

gene-specific primers were used to amplify an 848 bp fragment (Additional file 8, Table S2),

which was incorporated to vectors and *E. coli* using TOPO TA cloning kit (Thermo Fisher)

following the manufacturer's instructions, and Sanger sequenced to confirm their identities.

For colorimetric in situ hybridization, we adapted a protocol for the spider *Parasteatoda*

tepidariorum after Akiyama-Oda and Oda (2003). The staining reactions lasted between 2h

and 8h at room temperature using nitro-blue tetrazolium (NBT) and 5-bromo-4-chloro-3'-

indolyphosphate (BCIP) staining reactions (Sigma Aldrich). Embryos were then rinsed in

PBS-T (0.05% Tween-20), counter-stained with 10µg/mL Hoechst 33342 (Sigma-Aldrich,

St. Louis, MO, USA), post-fixed in 4% formaldehyde/PBS-T and stored at 4˚C. Images were

taken in a petri dish with agar under PBS-T, using a Nikon SMZ25 fluorescence

stereomicroscope mounted with either a DS-Fi2 digital color camera (Nikon Elements

software). Appendage mounts were imaged in 70% glycerol/PBS-T in a Zeiss LCM 710

confocal microscope.

Declarations

Ethics approval and consent to participate

Studies with arachnids do not required ethics approval or consent to participate.

Consent for publication

Not applicable.

Availability of Data and Materials

The data generated and analyzed in this study is included in the main article, supplementary

information and available upon request. Raw reads are deposited in NCBI Sequence Read

Archive under accession SRR12232018.

Competing interests

The authors declare having no conflict of interest.

Funding

GG was supported by a Wisconsin Alumni Research Foundation Fall Research Competition

award. Materials for study were supported by an Oscar Franke student competition award to

GG from the International Society of Arachnology. This material is based on work supported

by the National Science Foundation under grant IOS-1552610 to PPS.

Author's contributions

Both authors designed the study, analyzed data, developed protocols and contributed to

writing. GG conducted gene searches, orthology assessment, phylogenetic analysis, in situ

hybridization, assembled figures and drafted the manuscript. Both authors read and approved

the final version of the manuscript.

Acknowledgments

This work is dedicated to Peter Weygoldt, whose pioneer work in the development of

Phrynus marginemaculatus and contributions to the study of Amblypygi biology continue to

inspire the authors of this paper. Comments from two anonymous reviewers improved an

early draft of this manuscript. Sequencing was performed at the University of Wisconsin-

Madison BioTechnology Center. Microscopy was performed at the Newcomb Imaging

Center, Department of Botany, University of Wisconsin-Madison.

662
Author information

Affiliation

Department of Integrative Biology, University of Wisconsin-Madison, Madison, WI, USA

53706

Guilherme Gainett; Prashant P. Sharma

Corresponding author

Correspondence to Guilherme Gainett (guilherme.gainett@wisc.edu)

References
675

1. Damen WGM, Hausdorf M, Seyfarth E-A, Tautz D. A conserved mode of head

segmentation in arthropods revealed by the expression pattern of Hox genes in a spider.
PNAS. National Academy of Sciences; 1998;95:10665–70.

2. Stollewerk A, Schoppmeier M, Damen WGM. Involvement of *Notch* and *Delta* genes in
spider segmentation. Nature. Nature Publishing Group; 2003;423:863–5.

3. Akiyama-Oda Y, Oda H. Early patterning of the spider embryo: a cluster of mesenchymal
cells at the cumulus produces Dpp signals received by germ disc epithelial cells.
Development. 2003;130:1735–47.

4. Akiyama-Oda Y, Oda H. Cell migration that orients the dorsoventral axis is coordinated
with anteroposterior patterning mediated by Hedgehog signaling in the early spider embryo.
Development. 2010;137:1263–73.

5. Schwager EE, Schoppmeier M, Pechmann M, Damen WG. Duplicated Hox genes in the
spider *Cupiennius salei*. Front Zool. 2007;4:10–1.

6. Schwager EE, Sharma PP, Clarke T, Leite DJ, Wierschin T, Pechmann M, et al. The house
spider genome reveals an ancient whole-genome duplication during arachnid evolution. BMC
Biol. 2017;15:62.

7. Akiyama-Oda Y, Oda H. Axis specification in the spider embryo: *dpp* is required for
radial-to-axial symmetry transformation and *sog* for ventral patterning. Development.
2006;133:2347–57.

8. Oda H, Akiyama-Oda Y. Differing strategies for forming the arthropod body plan: lessons
from Dpp, Sog and Delta in the fly *Drosophila* and spider *Achaearanea*. Develop. Growth
Differ. 2008;50:203–14.

9. Pechmann M, Benton MA, Kenny NJ, Posnien N, Roth S. A novel role for *Ets4* in axis
specification and cell migration in the spider *Parasteatoda tepidariorum*. eLife. 2017;6.

10. Schwager EE, Pechmann M, Feitosa NM, McGregor AP, Damen WGM. *hunchback*
functions as a segmentation gene in the spider *Achaearanea tepidariorum*. Curr. Biol.
2009;19:1333–40.

11. Pechmann M, Khadjeh S, Turetzek N, McGregor AP, Damen WGM, Prpic N-M. Novel
function of *Distal-less* as a gap gene during spider segmentation. PLoS Genet.
2011;7:e1002342.

12. Setton EVW, Sharma PP. Cooption of an appendage-patterning gene cassette in the head
segmentation of arachnids. PNAS. 2018;128:201720193–10.

13. Paese CLB, Schoenauer A, Leite DJ, Russell S, McGregor AP. A SoxB gene acts as an
anterior gap gene and regulates posterior segment addition in a spider. eLife. 2018;7:1735.

14. Khadjeh S, Turetzek N, Pechmann M, Schwager EE, Wimmer EA, Damen WGM, et al.
Divergent role of the Hox gene *Antennapedia* in spiders is responsible for the convergent
evolution of abdominal limb repression. PNAS. National Academy of Sciences;
2012;109:4921–6.

15. Pechmann M, Schwager EE, Turetzek N, Prpic N-M. Regressive evolution of the

arthropod tritocerebral segment linked to functional divergence of the Hox gene *labial*.
Proceedings of the Royal Society B: Biological Sciences. 2015;282:20151162–6.

16. Mittmann B, Scholtz G. *Distal-less* expression in embryos of *Limulus polyphemus*
(Chelicerata, Xiphosura) and *Lepisma saccharina* (Insecta, Zygentoma) suggests a role in the
development of mechanoreceptors, chemoreceptors, and the CNS. Dev. Genes Evol.
2001;211:232–43.

17. Blackburn DC, Conley KW, Plachetzki DC, Kempler K, Battelle B-A, Brown NL.
Isolation and expression of *Pax6* and *atonal* homologues in the American horseshoe crab,
Limulus polyphemus. Satir P, Fallon JF, editors. Dev. Dyn. 2008;237:2209–19.

18. Barnett AA, Thomas RH. Posterior Hox gene reduction in an arthropod: *Ultrabithorax*
and *Abdominal-B* are expressed in a single segment in the mite *Archegozetes longisetosus*.
EvoDevo. 2013;4:23.

19. Barnett AA, Thomas RH. The delineation of the fourth walking leg segment is temporally
linked to posterior segmentation in the mite *Archegozetes longisetosus* (Acari: Oribatida,
Trhypochthoniidae). Evolution & Development. 2012;14:383–92.

20. Barnett AA, Thomas RH. The expression of limb gap genes in the mite *Archegozetes
longisetosus* reveals differential patterning mechanisms in chelicerates. Evolution &
Development. 2013;15:280–92.

21. Telford MJ, Thomas RH. Expression of homeobox genes shows chelicerate arthropods
retain their deutocerebral segment. PNAS. 1998;95:10671–5.

22. Khila A, Grbić M. Gene silencing in the spider mite *Tetranychus urticae*: dsRNA and
siRNA parental silencing of the *Distal-less* gene. Dev. Genes Evol. 2007;217:241–51.

23. Grbić M, Khila A, Lee K-Z, Bjelica A, Grbić V, Whistlecraft J, et al. Mity model:
Tetranychus urticae, a candidate for chelicerate model organism. Bioessays. 2007;29:489–96.

24. Dearden PK. Expression of pair-rule gene homologues in a chelicerate: early patterning
of the two-spotted spider mite *Tetranychus urticae*. Development. 2002;129:5461–72.

25. Santos VT, Ribeiro L, Fraga A, de Barros CM, Campos E, Moraes J, et al. The
embryogenesis of the Tick *Rhipicephalus* (*Boophilus*) *microplus*: The establishment of a new
chelicerate model system. genesis. 2013;51:803–18.

26. Sharma PP, Schwager EE, Extavour CG, Wheeler WC. Hox gene duplications correlate
with posterior heteronomy in scorpions. Proc. Biol. Sci. 2014;281:20140661–1.

27. Sharma PP, Schwager EE, Giribet G, Jockusch EL, Extavour CG. *Distal-less* and
dachshund pattern both plesiomorphic and apomorphic structures in chelicerates: RNA
interference in the harvestman *Phalangium opilio* (Opiliones). Evolution & Development.
2013;15:228–42.

28. Sharma PP, Schwager EE, Extavour CG, Giribet G. Evolution of the chelicera: a
dachshund domain is retained in the deutocerebral appendage of Opiliones (Arthropoda,
Chelicerata). Evolution & Development. 2012;14:522–33.

29. Sharma PP, Schwager EE, Extavour CG, Giribet G. Hox gene expression in the
harvestman *Phalangium opilio* reveals divergent patterning of the chelicerate opisthosoma.
Evolution & Development. 2012;14:450–63.

30. Sharma PP, Tarazona OA, Lopez DH, Schwager EE, Cohn MJ, Wheeler WC, et al. A
conserved genetic mechanism specifies deutocerebral appendage identity in insects and
arachnids. Proceedings of the Royal Society B: Biological Sciences. 2015;282:20150698–8.

31. Sharma PP, Kaluziak ST, Pérez-Porro AR, González VL, Hormiga G, Wheeler WC, et al.
Phylogenomic interrogation of Arachnida reveals systemic conflicts in phylogenetic signal.
Mol. Biol. Evol. 2014;31:2963–84.

32. Leite DJ, Baudouin-Gonzalez L, Iwasaki-Yokozawa S, Lozano-Fernandez J, Turetzek N,
Akiyama-Oda Y, et al. Homeobox gene duplication and divergence in arachnids. Mol. Biol.
Evol. 2018;35:2240–53.

33. Sharma PP, Santiago MA, González-Santillán E, Monod L, Wheeler WC. Evidence of
duplicated Hox genes in the most recent common ancestor of extant scorpions. Evolution &
Development. 2015;17:347–55.

34. Kenny NJ, Chan KW, Nong W, Qu Z, Maeso I, Yip HY, et al. Ancestral whole-genome
duplication in the marine chelicerate horseshoe crabs. Heredity. Nature Publishing Group;
2015;116:190–9.

35. Shingate P, Ravi V, Prasad A, Tay B-H, Garg KM, Chattopadhyay B, et al. Chromosome-
level assembly of the horseshoe crab genome provides insights into its genome evolution.
Nature Communications. Springer US; 2020;:1–13.

36. Schomburg C, Turetzek N, Prpic N-M. Candidate gene screen for potential interaction
partners and regulatory targets of the Hox gene labial in the spider *Parasteatoda*
tepidariorum. Development Genes and Evolution; 2020;:1–16.

37. Samadi L, Schmid A, Eriksson BJ. Differential expression of retinal determination genes
in the principal and secondary eyes of *Cupiennius salei* Keyserling (1877). EvoDevo.
BioMed Central; 2015;6:16–7.

38. Turetzek N, Pechmann M, Schomburg C, Schneider J, Prpic N-M. Neofunctionalization
of a Duplicate *dachshund* Gene Underlies the Evolution of a Novel Leg Segment in
Arachnids. Mol. Biol. Evol. 2015;33:109–21.

39. Turetzek N, Khadjeh S, Schomburg C, Prpic N-M. Rapid diversification of *homothorax*
expression patterns after gene duplication in spiders. BMC Evolutionary Biology; 2017;:1–
12.

40. Gainett G, Ballesteros JA, Kanzler CR, Zehms JT, Zern JM, Aharon S, et al. How spiders
make their eyes: Systemic paralogy and function of retinal determination network homologs
in arachnids. bioRxiv. Cold Spring Harbor Laboratory; 2020;129:1143–34.

41. Weygoldt P. Whip Spiders (Chelicerata: Amblypygi). Their Biology, Morphology and
Systematics. Apollo Books; 2000.

42. de Miranda GS, Giupponi APL, Prendini L, Scharff N. Weygoldtia, a new genus of

Charinidae Quintero, 1986 (Arachnida, Amblypygi) with a reappraisal of the genera in the family. Zoologischer Anzeiger - A Journal of Comparative Zoology. Elsevier GmbH; 2018;273:23–32.

43. Giribet G. Current views on chelicerate phylogeny—A tribute to Peter Weygoldt. Zoologischer Anzeiger - A Journal of Comparative Zoology. Elsevier GmbH; 2018;273:7–13.

44. Ballesteros JA, Sharma PP. A critical appraisal of the placement of Xiphosura (Chelicerata) with account of known sources of phylogenetic error. Halanych K, editor. Syst Biol. 2019;33:440–22.

45. Lozano-Fernandez J, Tanner AR, Giacomelli M, Carton R, Vinther J, Edgecombe GD, et al. Increasing species sampling in chelicerate genomic-scale datasets provides support for monophyly of Acari and Arachnida. Nature Communications. Springer US; 2019;:1–8.

46. Pereyaslawzewa S. Développement embryonnaire des phrynes. Annales des sciences naturelles, Zoologie. 1901;13:104–17.

47. Strubell A. Zur Entwicklungsgeschichte der Pedipalpen. Zoologischer Anzeiger. 1892;15:87–93.

48. Gough LH. The development of *Admetus pumilio* Koch; a contribution to the embryology of the Pedipalps. Quarterly Journal of Microscopical Science New Series. 1902;45:595–630.

49. Weygoldt P. Untersuchungen zur Embryologie und Morphologie der Geißelspinne *Tarantula marginemaculata* C. L. Koch (Arachnida, Amblypygi, Tarantulidae). Zoomorphologie. 1975;82:137–99.

50. Chapin KJ, Hebets EA. The behavioral ecology of amblypygids. Journal of Arachnology. American Arachnological Society; 2016;44:1–14.

51. Wiegmann DD, Hebets EA, Gronenberg W, Graving JM, Bingman VP. Amblypygids: Model Organisms for the Study of Arthropod Navigation Mechanisms in Complex Environments? Front. Behav. Neurosci. 2016;10:1–8.

52. Seiter M, Lemell P, Gredler R, Wolff JO. Strike kinematics in the whip spider *Charon* sp. (Amblypygi: Charontidae). The Journal of Arachnology. 2019;47:260–7.

53. Igelmund P. Morphology, sense organs, and regeneration of the forelegs (whips) of the whip spider *Heterophrynus elaphus* (Arachnida, Amblypygi). Journal of Morphology. Wiley-Blackwell; 1987;193:75–89.

54. Foelix RF, Chu-Wang IW, Beck L. Fine structure of tarsal sensory organs in the whip spider *Admetus pumilio* (Amblypygi, Arachnida). Tissue and Cell. 1975;7:331–46.

55. Santer RD, Hebets EA. The Sensory and Behavioural Biology of Whip Spiders (Arachnida, Amblypygi). 1st ed. Spider Physiology and Behaviour. Elsevier Ltd; 2011. pp. 1–64.

56. Foelix R, Troyer D, Igelmund P. Peripheral synapses and giant neurons in whip spiders. Microsc. Res. Tech. 2002;58:272–82.

57. Foelix RF. Occurrence of synapses in peripheral sensory nerves of arachnids. Nature.
1975;254:146–8.

58. Santer RD, Hebets EA. Agonistic signals received by an arthropod filiform hair allude to
the prevalence of near-field sound communication. Proceedings of the Royal Society B:
Biological Sciences. 2008;275:363–8.

59. Santer RD, Hebets EA. Evidence for Air Movement Signals in the Agonistic Behaviour
of a Nocturnal Arachnid (Order Amblypygi). Gronenberg W, editor. PLoS ONE.
2011;6:e22473–6.

60. Dong PDS, Dicks JS, Panganiban G. *Distal-less* and *homothorax* regulate multiple targets
to pattern the *Drosophila* antenna. Development. 2002;129:1967–74.

61. Dong PD, Chu J, Panganiban G. Proximodistal domain specification and interactions in
developing *Drosophila* appendages. Development. 2001;128:2365–72.

62. Casares F, Mann RS. Control of antennal versus leg development in *Drosophila*. Nature.
Nature Publishing Group; 1998;392:723–6.

63. Duncan DM, Burgess EA, Duncan I. Control of distal antennal identity and tarsal
development in *Drosophila* by *spineless-aristapedia*, a homolog of the mammalian dioxin
receptor. Genes Dev. Cold Spring Harbor Lab; 1998;12:1290–303.

64. Setton EVW, March LE, Nolan ED, Jones TE, Cho H, Wheeler WC, et al. Expression
and function of *spineless* orthologs correlate with distal deutocerebral appendage morphology
across Arthropoda. Developmental Biology. Elsevier Inc; 2017;430:224–36.

65. Struhl G. A homoeotic mutation transforming leg to antenna in *Drosophila*. Nature.
1981;292:635–8.

66. Struhl G. Genes controlling segmental specification in the *Drosophila* thorax. PNAS.
National Academy of Sciences; 1982;79:7380–4.

67. Prpic N-M, Damen WM. Expression patterns of leg genes in the mouthparts of the spider
Cupiennius salei (Chelicerata: Arachnida). Dev. Genes Evol. Springer-Verlag; 2004;214:1–7.

68. Prpic N-M, Janssen R, Wigand B, Klingler M, Damen WGM. Gene expression in spider
appendages reveals reversal of *exd/hth* spatial specificity, altered leg gap gene dynamics, and
suggests divergent distal morphogen signaling. Developmental Biology. 2003;264:119–40.

69. Pechmann M, Prpic N-M. Appendage patterning in the South American bird spider
Acanthoscurria geniculata (Araneae: Mygalomorphae). Dev. Genes Evol. 2nd ed. Springer-
Verlag; 2009;219:189–98.

70. Garwood RJ, Dunlop JA, Knecht BJ, Hegna TA. The phylogeny of fossil whip spiders.
BMC Evolutionary Biology; 2017;:1–14.

71. Weygoldt P. Beobachtungen zur fortpflanzungsbiologie und zum verhalten der
geißelspinne *Tarantula marginemaculata* C. L. Koch (Chelicerata, Amblypygi). Z. Morph.
Tiere. Springer-Verlag; 1969;64:338–60.

72. Weygoldt P. Lebenszyklus und postembryonale entwicklung der geibelspinne *Tarantula*
marginemaculata C. L. Koch (Chelicerata, Amblypygi) im laboratorium. Z. Morph. Tiere.
Springer-Verlag; 1970;67:58–85.

73. Schoppmeier M, Damen WGM. Double-stranded RNA interference in the spider
Cupiennius salei: the role of *Distal-less* is evolutionarily conserved in arthropod appendage
formation. Dev. Genes Evol. 2001;211:76–82.

74. Thomas RH, Telford MJ. Appendage development in embryos of the oribatid mite
Archegozetes longisetosus (Acari, Oribatei, Trhypochthoniidae). Acta Zoologica. John Wiley
& Sons, Ltd; 1999;80:193–200.

75. Nolan ED, Santibáñez López CE, Sharma PP. Developmental gene expression as a
phylogenetic data class: support for the monophyly of Arachnopulmonata. Dev. Genes Evol.
Springer Berlin Heidelberg; 2020;230:137–53.

76. Wagner GP, Amemiya C, Ruddle F. Hox cluster duplications and the opportunity for
evolutionary novelties. PNAS. National Academy of Sciences; 2003;100:14603–6.

77. Hughes CL, Kaufman TC. Hox genes and the evolution of the arthropod body plan.
Evolution & Development. 2002;4:459–99.

78. Ronshaugen M, McGinnis N, McGinnis W. Hox protein mutation and macroevolution of
the insect body plan. Nature. Nature Publishing Group; 2002;415:914–7.

79. Kaufman CLHATC. RNAi analysis of *Deformed*, *proboscipedia* and *Sex combs reduced*
in the milkweed bug *Oncopeltus fasciatus*: novel roles for Hox genes in the Hemipteran head.
2000;:1–12.

80. Pace RM, Grbić M, Nagy LM. Composition and genomic organization of arthropod Hox
clusters. EvoDevo. BioMed Central; 2016;7:11–1.

81. Nossa CW, Havlak P, Yue J-X, Lv J, Vincent KY, Brockmann HJ, et al. Joint assembly
and genetic mapping of the Atlantic horseshoe crab genome reveals ancient whole genome
duplication. GigaSci. 2014;3:708–21.

82. Zhou Y, Liang Y, Yan Q, Zhang L, Chen D, Ruan L, et al. The draft genome of
horseshoe crab *Tachypleus tridentatus* reveals its evolutionary scenario and well-developed
innate immunity. BMC Genomics. 4 ed. BioMed Central; 2020;21:137–15.

83. Harper A, Baudouin-Gonzalez L, Schönauer A, Seiter M, Holzem M, Arif S, et al.
Widespread retention of ohnologs in key developmental gene families following whole
genome duplication in arachnopulmonates. bioRxiv. Cold Spring Harbor Laboratory;
2020;160:1067–40.

84. Abzhanov A, Kaufman TC. Homologs of *Drosophila* appendage genes in the patterning
of arthropod limbs. Developmental Biology. 2000;227:673–89.

85. Beck L, Foelix R, Gödeke E, Kaiser R. Morphology, larval development, and hair sensilla
of the antenniform legs of the whip spider *Heterophrynus longicornis* Butler (Arach.,
Amblypygi). Zoomorphologie. Springer-Verlag; 1977;88:259–76.

86. Rayor LS, Taylor LA. Social behavior in amblypygids, and a reassessment of arachnid
social patterns. Journal of Arachnology. 2006;34:399–421.

87. Santer RD, Hebets EA. Tactile learning by a whip spider, *Phrynus marginemaculatus*
C.L. Koch (Arachnida, Amblypygi). J Comp Physiol A. 4 ed. 2009;195:393–9.

88. Fowler-Finn KD, Hebets EA. An Examination of agonistic interactions in the whip spider
Phrynus marginemaculatus (Arachnida, Amblypygi). Journal of Arachnology. 2006;34:62–
76.

89. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length
transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol.
Nature Publishing Group; 2011;29:644–52.

90. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics. 2014;30:2114–20.

91. Waterhouse RM, Seppey M, Simão FA, Manni M, Ioannidis P, Klioutchnikov G, et al.
BUSCO Applications from quality assessments to gene prediction and phylogenomics. Mol.
Biol. Evol. 2017;35:543–8.

92. Bryant DM, Johnson K, DiTommaso T, Tickle T, Couger MB, Payzin-Dogru D, et al. A
tissue-mapped axolotl de novo transcriptome enables identification of limb regeneration
factors. CellReports. 2017;18:762–76.

93. The UniProt Consortium. UniProt: a worldwide hub of protein knowledge. Nucleic Acids
Research. Oxford University Press; 2018;47:D506–15.

94. Gulia-Nuss M, Nuss AB, Meyer JM, Sonenshine DE, Roe RM, Waterhouse RM, et al.
Genomic insights into the *Ixodes scapularis* tick vector of Lyme disease. Nature
Communications. Nature Publishing Group; 2016;7:10507–13.

95. Grbić M, Van Leeuwen T, Clark RM, Rombauts S, Rouzé P, Grbić V, et al. The genome
of *Tetranychus urticae* reveals herbivorous pest adaptations. Nature. Nature Publishing
Group; 2011;479:487–92.

96. Herndon N, Shelton J, Gerischer L, Ioannidis P, Ninova M, Dönitz J, et al. Enhanced
genome assembly and a new official gene set for *Tribolium castaneum*. BMC Genomics;
2020;:1–13.

97. Chipman AD, Ferrier DEK, Brena C, Qu J, Hughes DST, Schröder R, et al. The first
myriapod genome sequence reveals conservative arthropod gene content and genome
organisation in the centipede *Strigamia maritima*. Tyler-Smith C, editor. PLoS Biol. Public
Library of Science; 2014;12:e1002005–24.

98. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable
generation of high-quality protein multiple sequence alignments using Clustal Omega. Mol
Syst Biol. Nature Publishing Group; 2011;7:1–6.

99. Gouy M, Guindon S, Gascuel O. SeaView Version 4: A Multiplatform Graphical User
Interface for Sequence Alignment and Phylogenetic Tree Building. Mol. Biol. Evol.
2010;27:221–4.

100. Nguyen L-T, Schmidt HA, Haeseler von A, Minh BQ. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol. Biol. Evol.
2015;32:268–74.

101. Koressaar T, Remm M. Enhancements and modifications of primer design program
Primer3. Bioinformatics. 2007;23:1289–91.

102. Nong W, Qu Z, Li Y, Barton-Owen T, Wong AYP, Yip HY, et al. Horseshoe crab
genomes reveal the evolutionary fates of genes and microRNAs after three rounds (3R) of
whole genome duplication. bioRxiv. Cold Spring Harbor Laboratory;
2020;6:2020.04.16.045815.

Figure legends

Figure 1: *Phrynus marginemaculatus* as a model to study Amblypygi development. A: adult

male *P. marginemaculatus*, dorsal view. B: adult female P. *marginemaculatus*, carrying

praenymphs on the opisthosoma. The yellow ink is used to mark the females in the colony.

Figure 2: Overview of the embryonic development of *Phrynus marginemaculatus*. Z-stack

automontage pictures of embryos imaged with nuclear staining (Hoechst) fluorescence. A–E:

five embryonic stages, in ventral view, spanning the elongation of appendages, and showing

germ band inversion and development of the ventral midline. F–H: head development, frontal

view. F: undifferentiated head lobes. G: head lobe with neuron precursor cell invaginations.

H: head lobes with anterior rim folding (semi-lunar grooves). I–K: opisthosomal

development, posterior view. I: embryo with one opisthosomal segment. J: embryo with five

opisthosomal segments. K: embryo ventral curved opisthosoma and most opisthosomal

segments formed (number unclear). L–N: lateral organ formation and opisthosomal curving,

lateral view. L: lateral organ as an undifferentiated bud proximal in leg II. M: lateral organ

with a discreet "C" shape, and opisthosomal segments curved up. N: lateral organ in its final

"kidney" shape, and opisthosoma curved ventrally. Insets in L–N show the lateral organ (Lo),

which is outlined with dashed lines in the respective main panel. Dashed lines on the

posterior regions of the embryos in L–N mark the curvature of the opisthosoma. O–Q:

deutembryo, in frontal, ventral and lateral view respectively. Ch: chelicera; Gz: growth zone;

H: head lobe; Iv: neural precursor cell invagination; St: stomodeum. Scale bars: 500 µm.

Figure 3: Hox gene duplications in Amblypygi. A: Schematic representation of the number of

copies of each of the ten Hox genes in the selected arthropod terminals. Dotted squares in the

amblypygids *Phrynus marginemaculatus*, *Charinus israelensis*, and *C. ioanniticus* indicate

absence from the transcriptome. Empty squares in *Parasteatoda* and *Tetranychus* indicate

absence from the genome. Number of Hox genes in *Carcinoscorpius rotundicauda* after [35]

(but see [102] for evidence of additional copies). B: Tree topology inferred from maximum

likelihood analysis of a conserved region (71 amino acid characters) using the same terminals

of the schematics (ln L = -3442.810). Numbers on the notes are ultrafast bootstrap resampling

frequencies (only >80 shown). Species: *Phrynus marginemaculatus* (Pmar); *Charinus*

israelensis (Cisr); *C. ioanniticus* (Cioa); *Parasteatoda tepidariorum* (Ptep); *Centruroides*

sculpturatus (Cscu); *Ixodes scapularis* (Isca); *Tetranychus urticae* (Ttur); *Strigamia maritima*

(Smar); *Drosophila melanogaster* (Dmel); *Tribolium castaneum* (Tcas). *lab*: *labial*; *pb*:

proboscipedia/maxillopedia; *Hox3*: *Hox3/zerknullt/z2*; *Dfd*: *Deformed*; *Scr*: *Sex combs*

reduced; *ftz*: *fushi tarazu*; *Antp*: *Antennapedia/prothorax-less*; *Ubx*: *Ultrabithorax*; *abdA*:

abdominal A; *AbdB*: *Abdominal B*.

Figure 4: *Pmar-Distal-less* (*Pmar-Dll*) colorimetric in situ hybridization. Whole mount bright

field images (Z-stack automontage) overlaid with nuclear staining (Hoechst). Ch: chelicera;

Pp: pedipalp; L1–4: leg 1–4; A/P: anterior/posterior; dAEL: days after egg laying. Scale bars:

500 µm.

Figure 5: *Pmar-Distal-less* (*Pmar-Dll*) colorimetric in situ hybridization, appendage

dissections in lateral view. Distal is to the left. A–F, G–L: bright field. A'–F', G'-L': nuclear

staining (Hoechst), confocal Z-stack maximum projection. A–F: 16–17 dAEL embryo. G–L:

17-18 dAEL embryo. J and J' have been vertically reflected to adjust orientation. Distal is to

the left. Arrowheads in A–F indicate proximal boundary of expression. Arrowheads in G–L

indicate stronger domains of expression. Ch: chelicera; Pp: pedipalp; L1–4: leg 1–4. Scale

bar: 250µm.

1007

1008   Figure 6: *Pmar-dachshund 1* (*Pmar-dac1*) (A–F) and *Pmar-dachshund 2* (*Pmar-dac2*) G–I)

1009   colorimetric in situ hybridization. Whole mount bright field images (Z-stack automontage)

1010   overlaid with nuclear staining (Hoechst). Ch: chelicera; Pp: pedipalp; L1–4: leg 1–4; Lo:

1011   lateral organ; A/P: anterior/posterior; arrowhead: expression domain; dAEL: days after egg

1012   laying. Scale bars: 500 µm.

1013

1014   Figure 7: *Pmar-dachshund 1* (*Pmar-dac1*) (A–H) and *Pmar-dachshund 2* (*Pmar-dac2*) (I–L)

1015   colorimetric in situ hybridization, appendage dissections in lateral view. Distal is to the left.

1016   White arrowhead indicates discreet domains of expression. Asterisk in "E" marks expression

1017   from the lateral margin of the headlobe. Ch: chelicera; Pp: pedipalp; L1, 4: leg 1, 4. dAEL:

1018   days after egg laying. Scale bar: 200 µm.

1019

1020   Figure 8: *Pmar-homothorax 1* (*Pmar-hth1*) (A–A') and *Pmar-homothorax 2* (*Pmar-hth2*) (B–

1021   B') colorimetric in situ hybridization, ventral view. A, B: whole mount bright field images

1022   (Z-stack automontage). A', B': nuclear staining (Hoechst) (Z-stack automontage). Ch:

1023   chelicera; Pp: pedipalp; L1–4: leg 1–4. A/P: anterior/posterior; arrowhead: expression

1024   domain; dAEL: days after egg laying. Scale bar: 500 µm.

1025

1026   Figure 9: *Pmar-homothorax 1* (*Pmar-hth1*) (A–H) and *Pmar-homothorax 2* (*Pmar-hth2*) (I–

1027   L) colorimetric in situ hybridization, appendage dissections in lateral view. Distal is to the

1028   left. White arrowhead indicates discreet domains of expression. Dotted line in "G" marks

1029   expanded distal expression domain. Ch: chelicera; Pp: pedipalp; L1, 4: leg 1, 4. dAEL: days

1030   after egg laying. Scale bar: 200 µm.

1031

1032   Figure 10: Tool for removing the brood sac from whip spiders. A: dissecting dish made of

1033   plastic petri dish and foam. B: detail of an immobilized female ready to have the brood sac

1034   dissected. C: intact brood sac dissected from the female. Scale bars: A: 1 cm (approximate);

1035   B: 5 mm (approximate); C: 1 mm.

1036

1037

Additional file 1, Figure S1: Tree topology inferred from maximum likelihood analysis of a

conserved region (75 amino acid characters) (ln L = -4780.167) using the same terminals of

Figure 3 and including *Parasteatoda tepidariorum Hox3-A* paralog and *Drosophila*

melanogaster bicoid (*Hox3*) (red arrow). Numbers on the notes are ultrafast bootstrap

resampling frequencies (only >80 shown). For abbreviations, see Additional file 5, Table S1.

File format: jpeg.

1044

1045   Additional file 2, Figure S2: Tree topology of *Distal-less* (*Dll*) inferred from maximum

1046   likelihood analysis amino acid sequences (ln L = -15272.780). Numbers on the notes are

1047   ultrafast bootstrap resampling frequencies (only >80 shown). Accession numbers are

1048   available in Additional file 5, Table S1. The terminal for the whip spider *Phrynus*

1049   *marginemaculatus Dll* ortholog is in boldface. File format: jpeg.

1050

Additional file 3, Figure S3: Tree topology of *dachshund* (*dac*) inferred from maximum

likelihood analysis amino acid sequences (ln L = -21699.907). Numbers on the notes are

ultrafast bootstrap resampling frequencies (only >80 shown). The terminals for the whip

spider *Phrynus marginemaculatus Dll* orthologs are in boldface. The original protein

alignment of all terminals (except *P. marginemaculatus*) is from Noland et al. 2020. File

format: jpeg.

1057

Additional file 4, Figure S4: Tree topology of *homothorax* (*hth*) inferred from maximum

likelihood analysis amino acid sequences (ln L = -10459.647). Numbers on the notes are

ultrafast bootstrap resampling frequencies (only >80 shown). The terminals for the whip

spider *Phrynus marginemaculatus hth* orthologs are in boldface. The original protein

alignment of all terminals (except *P. marginemaculatus*) is from Noland et al. 2020. File

format: jpeg.

1064

1065   Additional file 5, Table S1: List of species, accession numbers and abbreviations of terminals

1066   used in the phylogenetic analysis of Hox genes, *Distal-less*, *dachshund* and *homothorax*

1067   homologs. File format: xlsx

1068   Additional file 6: Protein alignment used in the phylogenetic analysis depicted in Figure 3.
1069   File format: fasta.

1070

1071   Additional file 7: Protein alignment used in the phylogenetic analysis depicted in Additional
1072   file 1, Figure S1. File format: fasta.

1073

1074   Additional file 8, Table S2: List of primers for *Pmar-Dll*, *Pmar-dac1/2*, *Pmar hth1/2*, and

1075   cloned fragments for *Pmar-Dll*. File format: xlsx

1076

1077

anterior view ventral view lateral view

A **B** **C** 15th dAEL

Ch
Pp
L1
L2
L3
L4
A ↕ P

D **E** **F** 16th–17th dAEL

Ch
Pp
L1
L2
L3
L4
A ↕ P

G **H** **I** 17th–18th dAEL

Ch
Pp
L1
L2
L3
L4
A ↕ P

Pmar-Dll

500 μm

250µm

| anterior view | ventral view | lateral view |

Pmar-hth1

Pmar-hth2

A

Ch
Pp
L1
L2
L3
L4

A
P

B

Pp
L1
L2
L3
L4

A
P

A'

Ch
Pp
L1
L2
L3
L4

A
P

B'

Pp
L1
L2
L3
L4

A
P

16th–17th dAEL

500 µm