Multiple sounds degrade the frequency representation in monkey inferior colliculus

Authors: Shawn M. Willett[1,2*] and Jennifer M. Groh[1-6]

Affiliations:
[1]Department of Neurobiology
[2]Center for Cognitive Neuroscience
[3]Department of Psychology and Neuroscience
[4]Department of Computer Science
[5]Department of Biomedical Engineering
[6]Duke Institute for Brain Sciences
Duke University
* corresponding author: shawn.willett@duke.edu

Acknowledgements: We are grateful to Stephen Lisberger, Lindsey Glickfeld, Henry Greenside, Patrick Mayo, Christopher Henry, Jeff Mohl, Surya Tokdar, John Pearson, Stephanie Schlebusch and members of the Groh lab for comments on this work. We are also grateful to Valeria Caruso for her help in establishing the behavior paradigm. This work was funded through NIH grants DC016363 and DC013906 to JMG.

Abstract:

How we distinguish multiple simultaneous stimuli is uncertain, particularly given that such stimuli sometimes recruit largely overlapping populations of neurons. One hypothesis is that tuning curves might change to limit the number of stimuli driving any given neuron when multiple stimuli are present. To test this hypothesis, we recorded the activity of neurons in the inferior colliculus while monkeys localized either one or two simultaneous sounds differing in frequency. Although monkeys easily distinguished simultaneous sounds (~90% correct performance), the frequency tuning of inferior colliculus neurons on dual sound trials did not improve in any obvious way. Frequency selectivity was degraded on dual sound trials compared to single sound trials: tuning curves broadened, and frequency accounted for less of the variance in firing rate. These tuning curve changes led a maximum-likelihood decoder to perform worse on dual sound trials than on single sound trials. These results fail to support the hypothesis that changes in frequency response functions serve to reduce the overlap in the representation of simultaneous sounds. Instead these results suggest alternative theories, such as recent evidence of alternations in firing rate between the rates corresponding to each of the two stimuli, offer a more promising approach.

Introduction:

Natural environments contain numerous stimuli. A fundamental question is how the brain simultaneously encodes multiple items, particularly when more than one item falls within a given neuron's receptive field. Two possible (and potentially complementary) solutions to this multiplicity conundrum have been proposed: that receptive fields effectively shrink or shift to limit the number of stimuli within them, and/or that fluctuating activity patterns allow individual neurons to contribute to encoding only one particular stimulus at any given time. There is evidence in favor of the fluctuating activity hypothesis (Caruso et al., 2018), but whether the underlying tuning curves may also change to facilitate encoding multiple stimuli remains unknown. Here, we evaluated whether and how the presence of more than one sound alters the frequency tuning properties of neurons in the inferior colliculus (IC).

The monkey IC is an ideal structure to probe how neurons code concurrent stimuli for two reasons. First, nearly every auditory signal converges to the IC prior to ascending to the thalamus (Aitkin & Phillips, 1984), so information must be preserved at this stage in order to be available to higher areas (for review see *The Inferior Colliculus*, 2005). Causal studies have indicated a role in frequency perception (Pages et al., 2016) and sound localization (Jenkins & Masterton, 1982; Kelly & Kavanagh, 1994). Second, the representations of sound frequency and location in the monkey IC are coarse (Bulkin & Groh, 2011; Groh, Kelly, & Underhill, 2003; Ryan & Miller, 1978; Versnel, Zwiers, & van Opstal, 2009; Zwiers, Versnel, & Van Opstal, 2004). A given pure tone can activate anywhere from 20-80% of frequency modulated sites (Bulkin & Groh, 2011), implying that broadband natural stimuli are likely to be capable of activating an even higher percentage. Neurons tend not to have circumscribed spatial receptive fields but respond monotonically as a function of increasing sound eccentricity over a broad range of space (a hemifield or more, Groh et al., 2003; Zwiers et al., 2004).

In short, the almost complete convergence of ascending auditory signals in the IC, together with the broad frequency and spatial tuning in this structure, suggests that two sounds can potentially recruit similar populations of IC neurons. Nevertheless, humans (Best, Schaik, & Carlile, 2004; Perrott, 1984a, 1984b; Zhong & Yost, 2017) and monkeys (Caruso et al., 2018) are capable of localizing two concurrent sounds, implying that the IC population must overcome these coding constraints.

One possibility is that IC neurons sharpen or shift their frequency tuning curves to limit the degree to which any two stimuli fall within their receptive field. Tuning curves are not

immutable (Weinberger, 1995). For example, spatial tuning in cat auditory cortex is modulated by behavioral state: units show broad selectivity under anesthesia (Imig, Irons, & Samson, 1990; Middlebrooks, Clock, Xu, & Green, 1994; Middlebrooks & Pettigrew, 1981) but become more selective in the awake cat (Mickey & Middlebrooks, 2003), particularly when performing a spatial auditory task (Lee & Middlebrooks, 2011). Surround suppression in the visual system provides another example (Chettih & Harvey, 2019; Hartline & Ratliff, 1957) as do response profile modulations associated with experience (Freedman, Riesenhuber, Poggio, & Miller, 2005) or auditory attention (Connell, Barczak, Schroeder, & Lakatos, 2014).

Could the monkey IC also sharpen tuning curves to facilitate encoding of multiple stimuli? Previous work in the rabbit IC showed concurrent sound presentation broadens spatial tuning (Day, Koka, & Delgutte, 2012). However, it remains to be seen if frequency tuning is affected by the presence of a concurrent sound. The present study investigates how frequency tuning curves are modulated by the presence of an additional sound and if these changes are beneficial to decoding stimulus information from the population.

Single neuron activity was recorded from the IC while monkeys made saccades to one or two simultaneously presented sounds. Monkeys performed at least 86% correct on dual sound trials for all frequency separations. Frequency tuning curves were often altered when a second sound was presented but did not appear to shrink or shift. Rather, most neurons actually broadened their tuning curves. A maximum-likelihood decoder used to decode the stimulus on a held-out trial performed slightly worse when the held-out trial was a dual-sound trial. Thus, the frequency tuning curve modulations due to the presence of an additional sound seem to limit the available stimulus information encoded by the population of IC neurons. We conclude that changes to the frequency tuning properties of IC neurons in the presence of multiple sounds are unlikely to solve the multiplicity problem.

Methods:

General: All procedures were approved by the Duke Institutional Animal Care and Use Committee. Two adult, female, rhesus monkeys participated in this study. General procedures have been previously described in greater detail (Caruso et al., 2018; Groh, Trause, Underhill, Clark, & Inati, 2001). Briefly, under aseptic conditions a head post and a scleral search coil (Judge, Richmond, & Chu, 1980; Robinson, 1963), were surgically implanted. After recovery monkeys were trained to perform a sound localization task as described below. Once the task

was learned, a recording chamber (Crist Instruments) was implanted over the left IC (in both monkeys) based on stereotaxic coordinates and was confirmed after implantation with structural MRI (Groh et al., 2001).

Training and Behavior: During behavioral training and recording sessions monkeys performed single and dual sound localization tasks as shown in figure 1AB. Monkeys were seated in a dark, sound attenuated room with head movements restrained. The illumination of a fixation light (LED) at 0° in both the horizontal and vertical dimensions signified the start of the trial. Monkeys began the trial by acquiring and holding fixation on the light. After a variable duration of fixation (600-700 ms) either a single sound or two simultaneous sounds would be presented while the monkey held fixation. After a variable overlap of fixation and sound presentation (1000-1100 ms) the fixation light was turned off cueing the monkey to saccade to the origin of the sound or sounds. The monkey had a brief temporal window for their eyes to leave the fixation window and enter the first target window. The monkey then had to hold fixation in the target window briefly before initiating a second saccade to the second target window or until the single sound trial ended. The sound remained on until the end of the trial. If the monkeys performed the trial correctly, they were rewarded with a grape juice solution (30% grape juice, 70% water). Reward was delivered through a drinking tube and controlled via a solenoid.

The sound localization task used in this study was designed and implemented with proprietary software (Beethoven, RYKLIN Software Inc) and was adapted from a previous study (Caruso et al., 2018). There were two major adaptations. First, additional sound frequencies were used. Second, sounds in this study are only located at ±12°, meaning on dual sound trials the sounds are 24° apart compared to 30° in Caruso et al. (2018).

Initially, monkeys were trained to saccade to the origin of a sound (e.g. to the location of the speaker). Monkey Y was then trained to saccade to the source of sounds presented in a sequence. Over time the inter-sound-interval was decreased until eventually the sounds were presented simultaneously (Caruso et al., 2018). Monkey Y was trained on all conditions except the control conditions. Monkey N was trained on the final task, including all conditions.

Eye position was acquired through an implanted scleral coil and Riverbend field system. Eye coil outputs were calibrated each day before behavioral and recording sessions. All behavioral data, including eye position data, were saved in Beethoven files for offline analysis.

Sound Stimuli: All sounds consisted of bandpass filtered noise constructed by adding a series of tones. Sounds were frozen for each experimental session; that is for any given experimental session every instance of the sound stimulus of a given frequency involved the same waveform. The center frequencies were equally spaced in 0.19 octave intervals: 420, 500, 595, 707, 841, 1000, 1189, 1414, 1682 and 2000 Hz. These sounds were chosen due to the large proportion of IC neurons responsive at these frequencies (Bulkin & Groh, 2011), increasing the chance that more than one sound would significantly drive any recorded neuron.

All sounds were generated using proprietary software (Beethoven, RYKLIN Software Inc.) and passed through appropriate amplifiers to two speakers matched in their frequency response characteristics. Speakers were positioned at ±12° in the horizontal azimuth and at 0° in elevation at a distance approximately 1 meter from the monkey. Sounds were calibrated using a microphone at the position where the monkey's head would be during the experiment. Sounds were calibrated to be 55 dB SPL when presented in isolation; these stimuli resulted in 60 dB SPL sound reaching the monkey on dual sound trials. The intensity difference is not expected to have substantive effects on neural activity or behavior (Caruso et al., 2018). We kept the sound source signals constant across conditions because in the real-world simultaneously occurring sounds are louder than when the component sounds are presented in isolation. All experiments were conducted in an anechoic chamber; the wall and ceilings were lined with echo attenuating foam (Sonex) and the floor was carpeted.

Across all experimental sessions there were 56 conditions, involving either one or two sounds in combination. For simplicity, we will refer to the sounds in two subsets, "A" and "B". The A sounds were used to assess the frequency tuning curves and consisted of the 8 middle sound frequencies (500 to 1682 Hz). The B sounds flanked the A sounds, at either 420 or 2000 Hz. On single sound trials any one of the 10 (8 A + 2 B) sounds could be presented in isolation at either ± 12°. On dual sound trials, an A sound was presented at either ± 12° and paired with a B sound at the opposite location. The dual and single sound trials accounted for 52 of the 56 conditions (8 A sounds x 3 paired conditions (none, 420 or 2000 Hz) x 2 locations = 48 + 2 B sounds presented in isolation x 2 locations = 56 conditions). To ensure monkeys were actively perceiving both sounds and not saccading to remembered locations in response to more complex spectral content (e.g. the dual sound trials have power in more bands than single sound trials), an added control accounted for the last 4 conditions. The control trial presented a 1189 Hz sound with either B sound at either ±12°, that is to say the 1189 Hz and B sound were targeted to the

same speaker location (2 sounds (1189+420 Hz or 1189+2000 Hz) x 2 locations = 4). Monkeys typically made single saccades on these control trials and reached high performance levels.

To acquire enough trials per condition and to accommodate cell holding times (typically 1-2 hours), the number of conditions per recording session were limited by presenting the A sounds at only one location (either ±12°) each day. Therefore, each recording session consisted of 32 conditions. The eight A sounds presented alone or in combination with either B sound accounted for 24 of the 32 conditions (8 sounds x 3 paired tones x 1 location = 24). To allow counterbalancing, the two B sounds were still presented at both locations (2 sounds x 2 locations = 4). The control conditions were not changed across days and accounted for the last 4 conditions. (2 sounds x 2 locations = 4). However, the control condition was added after monkey Y recordings were completed, therefore monkey Y recording sessions only consist of 28 conditions.

Conditions were randomly interleaved: any given trial had a 45% chance to be a single sound trial, a 45% chance to be a dual sound trial, or a 10% chance to be a control trial (or for monkey Y a 50% chance to be either a single- or dual-sound trial). Within the control and dual sound trial types the likelihood of any condition was equal. The likelihood of any single condition was equal except on 50% of single sound trials the B sounds were presented at the opposite location of the A sounds to counterbalance the location of the single sounds.

Neural Recordings: Neural activity was acquired through a Multichannel Acquisition Processor (MAP system, Plexon Inc). Single and multiunit activity signals were sent to an external speaker for auditory detection of neural activity. Once single units were isolated, using the box method (Plexon SortClient software), spike times were sent to Beethoven and saved for offline analysis.

Prior to each recording session the chamber was cleaned and a grid and turret (Crist Instrument, 1mm spaced grid) was inserted into the chamber. Prior to recordings, grid positions likely to intercept the IC were localized using structural MRI (Groh et al., 2001). A single tungsten micro-electrode (FHC, ~1-5 MΩ) was backloaded into a stainless-steel guide tube. The guide tube was then manually inserted approximately 1 cm into the brain. The electrode was then advanced with a microdrive (NAN Instruments) at 100-200 µm/s until the electrode tip was just outside of the guide tube. This was indicated by a mark on the electrode and confirmed by a change in the hash typical upon electrode entrance into the brain. The depth on the microdive was then zeroed and the electrode was advanced at roughly 8 µm/s while monkeys sat passively listening to sounds of variable frequencies presented every few seconds. Once the expected depth

and/or sound related activity (determined by ear) was encountered in the hash, electrode movement was stopped, and the electrode was allowed to settle for approximately 1 hour. The settling depth typically occurred 22 mm after exiting the guide tube. The electrode was then advanced in 1 µm increments until a cell was well isolated. If no cells were well isolated the electrode was retracted at roughly 8 µm/s.

A total of 105 neurons were recorded (monkey Y right IC, N=45; monkey N right IC, N=59; monkey N left IC, N=1).

Analysis: *Behavior:* All behavioral analysis was done on attempted trials, defined as those in which the monkey successfully completed the fixation epoch and left the fixation window after the go cue. Proportion correct was defined as the # of Correct Trials / # of Attempted Trials

Neural Activity: All analysis of neural activity was done on correct trials and during the first 500 ms after sound onset, i.e. during a period of time when monkeys were fixating (to reduce effects related to eye movements known to occur in the macaque IC (Bulkin & Groh, 2012a, 2012b; Groh et al., 2001; Porter, Metzger, & Groh, 2006, 2007). Neurons were classified as responsive if their spike count 500 ms after sound onset across all tested frequencies was significantly different from their spike count during baseline, the 500 ms prior to sound onset (t-test, $p < 0.05$). Ninety-three of the 105 neurons were classified as responsive. Neurons were classified as frequency selective, or tuned, if their spike counts 500 ms after sound onset were significantly modulated across all 10 sound frequencies (one-way ANOVA, $p < 0.05$). Sixty-three of 105 neurons were classified as frequency selective.

Point Image: The point image refers to the proportion of recorded units that are responsive to a given condition (Capuano & McIlwain, 1981). Neurons were classified as responsive if, for a given condition, their spike count 500 ms after sound onset was significantly different than their spike count 500 ms prior to sound onset (t-test, $p < 0.05$).

d' (d-prime): To investigate how differently a cell responded for A vs B sounds we calculated a sensitivity index or d':

$$d' = \frac{\mu_A - \mu_B}{\sqrt{\frac{1}{2}(\sigma_A^2 + \sigma_B^2)}}$$

Where μ_A and σ_A^2 corresponds to the mean spike count and spike count variance for sound A, respectively, and μ_B and σ_B^2 corresponds to the mean spike count and spike count variance for sound B, respectively. This metric quantifies the separability of the sound A and sound B spike count distributions. We took the magnitude of the d' value (|d'|) since the order in the numerator is arbitrary (the A or B sound could be the first term). A |d'| value of 0 indicates the sound A and sound B spike count distributions are similar and higher values of |d'| indicate the sound A and sound B spike count distributions are more distinguishable.

Equivalent Rectangular Receptive Fields (ERRFs): ERRFs serve as a measure of the width of the frequency receptive field and were calculated in a similar fashion to previous work (Lee & Middlebrooks, 2011). The area under the (non-baseline subtracted) tuning curve was measured with the *trapz()* function in MATLAB, which integrates the area using the trapezoid method. This area was then divided by the peak firing rate of the tuning curve, creating a rectangle with a width dependent upon tuning curve breadth and frequency modulation of peak firing rate.

Maximum-Likelihood Decoding: The decoding algorithm used here was adapted from previous work (Wallisch, 2014) and is similar to other implementations of maximum-likelihood decoding (Day & Delgutte, 2013; Jazayeri & Movshon, 2006). The goal of the decoder was to infer the stimulus that was presented on a set of held-out trials (1 held-out trial of a given condition per cell). The predicted condition was the condition (1 out of 26 possible conditions) that had highest population-log-likelihood. This computation was repeated 1000 times per condition. Since there were not 1000 unique trials per condition for each cell the particular held-out trial on a given repeat was sampled from all possible trials with replacement.

To determine the population-log-likelihood, first the likelihood for each cell was computed given the held-out trials spike count. Second, the weights were determined by finding the average spike count for each condition without the held-out data. Then likelihood for all 26 conditions given the spike count was computed:

$$\ln (Condition\ Likelihood) = \ln (P(spike\ count \mid frequency))$$

This results in a vector with 26 elements for each cell. Each element corresponds to the log-likelihood of that condition for that held-out spike count. Essentially the log-likelihood is given by the difference between the held-out spike count and the mean firing rate across all conditions. These log-likelihood vectors are then summed across all cells to generate a population-log-likelihood:

$$\ln(\text{Population Condition Likelihood}) = \sum_{i=1}^{N} \ln\left(P(\text{spike count}_i \mid \text{stimulus})\right)$$

Where spike count$_i$ corresponds to the *ith* neuron spike count, and N is the number of neurons in the population. Therefore, for each repeat of the decoding computation, the result is a 26-element vector where each element corresponds to the population-log-likelihood for each condition. The predicted condition for that repeat is the element with the largest population-log-likelihood. Importantly, for any repeat the inferred condition could have been one of the 26 conditions (8 A sounds x 3 sound pairs = 24 conditions + 2 B sounds = 26 conditions). We used all responsive units (N = 93) for the decoding analysis.

Figure 1 Task schematic and behavioral performance. (**A**) Monkeys were trained on a sound localization task. Left panel: A time course schematic of the task. Speaker 2 onset and the second saccade and fixation would only occur if it was a dual-sound trial. Right Panel: In the sound localization task the monkey would initiate a trial by fixating an LED. After variable interval fixation (600-700 ms), either 1, or 2 simultaneous, sound(s) were presented from one or two locations. After a variable interval of sound/fixation overlap (1000-1100 ms) the fixation light is turned off, which cued the monkey to report to the origin(s) of the sound(s) via saccade(s). On single sound trials monkeys made a single saccade. On dual-sound trials monkeys made a sequence of saccades to the origin of one sound and then the other. (**B**) The combined proportion of correct single-sound trials (mean = 0.8011 ± 0.0069) and dual-sound trials (mean = 0.8941 ± 0.0060) across both monkey Y and monkey N for 105 behavioral sessions collected during cell recordings. (**C**) Change in performance across both monkeys as a function of the frequency separation between the two sounds. Performance increases as distance between sound frequencies increase (significantly fit by a line with a slope > 0; slope 0.867, $p < 0.05$). Error bars represented standard error of the mean (**B & C**).

Results:

Monkeys could successfully report the locations of all sound combinations used in this study. Across all attempted dual sound trials, monkeys performed roughly 90% correct (Fig. 1B), compared to the 80% correct on single sound trials. This high degree of performance on the dual sound trials was only modestly affected by the frequency separation between the two sounds: monkeys performed at about 87% correct for the smallest frequency separation of 0.19 octaves (Figure 1C), rising slightly to 91% correct for the largest frequency separation of 2 octaves.

This good performance does not appear to be the result of recruiting largely unique sets of neurons to encode each of the two stimuli. Of the N=93 sound-responsive neurons we recorded (identified by t-test, see Methods), 45-70% responded significantly to any given stimulus presented individually (Fig. 2, red curve). The large proportion of responsive units suggests that sounds presented in this experiment recruit overlapping populations of neurons. The low frequency bias (negative slope of the red curve in Fig. 2), is similar to previous macaque IC findings (Bulkin & Groh, 2011).

When a second sound was added, more units responded (Fig. 2, blue and black curves vs. the red curve). This increase in the number of responsive units at the population level should not occur if individual units were sharpening their tuning.

Figure 2 A point image plot, or the proportion of the responsive population that responds to a given frequency. Each point corresponds to the proportion of the responsive population that is significantly responsive compared to baseline for the given condition. A unit was counted as responsive if its firing rate in 0-500 ms after sound onset was significantly different than its firing rate from -500-0 ms before sound onset (t-test, p<0.05). Red corresponds to sounds presented in isolation, while blue and black correspond to dual-sound trials paired with 420 Hz or 2000 Hz, respectively.

We next compared the sensitivity of individual IC neurons to the specific pairs of sounds used. This analysis focused on the roughly 67% (N = 63) of responsive neurons whose spike counts were significantly modulated by frequency (one-way ANOVA, p < 0.05). We quantified the separability of the spike count distributions of the two components of the dual sound conditions (e.g. 500 (A-sound) vs. 420 (B-sound) when either were presented in isolation) using a d' analysis (see Methods).

Figure 3A shows the results for the comparison with a 420 Hz B sound. Units were sorted based on the sum of their |d'| values across conditions, with the units showing the highest aggregate |d'| on top. For the smallest frequency separation (left-most column), few neurons exhibit large d' values: only about a third exhibit a d' value greater than 1 (bottom panel). Figure 3B shows the complementary pattern for the 2000 Hz B sounds. Note that the unit order in two panels are different; both are sorted so the unit with the highest summed |d'| within a panel is at the top. For both panels, a large minority of cells barely differentiated between any of the A and B sounds, as evidenced by the white bands at the bottom of the figure (roughly unit 70

to 93). Overall, this analysis suggests that only a fraction of the recorded populations (roughly the top 20 units in both panels) possess sufficient sensitivity to underlie the improvement in performance due to frequency separation on dual sound trials (Fig. 1C).

Figure 3 A sensitivity index, or d', between the A and B sounds when they are presented in isolation. (A) Top Panel: The magnitude of d' between each of the 8 A sounds and 420 Hz. A higher |d'| corresponds to a larger difference between the spike count distributions of the two sounds. Each row corresponds to a unit. Each column corresponds to a particular sound frequency. Units are sorted by their total |d'| across all 8 A sounds compared to 420 Hz. Bottom Panel: The percent of responsive neurons (N = 93) that have a |d'| value > 1 for each frequency compared to 420 Hz. (**B**) The same as panel (**A**) but compared to 2000 Hz. Note these units are sorted by their total |d'| across all 8 A sounds compared to 2000 Hz. Therefore, unit n in **A** does not necessarily correspond to unit n in panel **B**.

We then evaluated whether neural response patterns change to facilitate the encoding of simultaneously presented sounds. Figure 4 compares the single and dual sound tuning profiles of six example cells. On single sound trials (red traces), neurons could prefer low frequencies (Fig. 4A, E, F), intermediate frequencies (Fig. 4B, D), or high frequencies (Fig. 4C). Concurrent sounds could change tuning in multiple ways. The neurons in figure 4A-C all become less responsive to sounds overall: the responses in the dual sound cases were generally lower than the responses to the "better" of the two single sounds (e.g. black curves fall between the red and blue curves). Some neurons appear to shift their preferred frequency (Fig. 4D), and some neurons appear unaffected by the presence of an additional sound (Fig. 4E, F). Interestingly, some neurons displayed multi-peaked tuning curves that were robust across single and dual sound conditions (Fig. 4F). Qualitatively, it seems that the majority of neurons become less responsive in the presence of simultaneous sounds, but whether this suppression leads to receptive fields that sharpen or broaden requires quantification of tuning curves across the population.

Figure 4 Example modulations of frequency tuning curves due to the presence of an additional sound in 6 different cells. (**A**) The tuning curve across conditions for a cell that displays sharpening in the dual-sound condition. Each point corresponds to the baseline (-200 ms bin w.r.t sound onset) subtracted mean firing rate (500 ms bin w.r.t sound onset). Red points correspond to sounds presented in isolation. The blue point corresponds to the paired sound when presented in isolation. The blue line is a reference line, so the firing rate of the paired sound is visible across the panel. The black points correspond to the dual-sound condition. (**B and C**) The same as panel (**A**) but for two different cells that show a decrease in response and selectivity. (**D**) The same as panel (**A**) but for a cell that shifts its preferred frequency away from the paired sound. (**E & F**) The same as panel (**A**) but for two cells that seem to be unaffected by the presence of an additional sound. Error bars represent standard error of the mean(**A-F**).

To do this, we selected units that were frequency selective to the A sounds when presented alone (N=57, one-way ANOVA, see Methods). The number of frequency-selective units dropped to N = 47 (82% of N=57) when the A sounds were paired with either 420 or 2000 Hz. To evaluate effect size, we compared the F-statistic (a metric of how much frequency modulates spike count) in the dual vs. single conditions (Figure 5A, C). If the neurons were more selective in the dual sound conditions, the F statistic should be higher for these conditions and the data would tend to lie above the line of slope one, but they do not. Figure 5 B and D quantify this further by subtracting the single sound F statistic from the dual sound F statistic for a given neuron. If the neurons were more selective on dual sound than on single sound trials, these distributions would be positively skewed, but they are not. In fact, the negative skew, i.e. worsening frequency selectivity, is significant for the 420-Hz pairing (Figure 5B, mean = -7.989 ± 1.219 S.E.M, $p < 0.05$). Overall, then, neurons' frequency selectivity was either unaffected by a second sound or was reduced by it.

Figure 5 Frequency modulation of firing rate is affected by the presence of an additional sound. (**A**) The F-statistic measured from a one-way ANOVA investigating firing rate across different frequencies (all 8 A sounds). A high F-statistics indicates that there are large differences in firing rate across the different frequencies. The single-sound F-statistics are plotted against the 420 Hz paired F-statistics. Each point corresponds to one unit, out of 93 sound responsive units. Points above the unity line indicate that frequency modulates firing rate more in the 420 Hz paired condition. Points on the unity line indicate there is no change in the modulation of firing rate across the two conditions. Points below the unity line indicate that frequency modulates firing more in the single-sound condition. (**B**) The distribution of change in F-statistic per cell (F-Statistic$_{Single}$ – F-statistic$_{Dual}$). A value of 0 indicates no change across the two conditions, a value greater than 0 indicates the 420 Hz paired condition was more sensitive to frequency and a value less than 0 indicates the single-sound condition was more sensitive to frequency. The single-sound condition was significantly more sensitive to frequency (mean = -7.99 ± 1.22 S.E.M, p < 0.05). The blue vertical line indicates the mean of the distribution. The black vertical line is at 0. (**C**) The same as (**A**) but for compared to the 2000 Hz paired condition. (**D**) The same as (**B**) but compared to the 200 Hz paired condition. There was no difference in the modulation of firing rate across the two conditions (mean = -1.14 ± 0.9 S.E.M, p = 0.21). Note that in (**A & C**) an outlier data point was removed for visualization purposes. In panel (**A**) its position is x =364.5, y = 31.2. In panel (**C**) its position is x = 364.5, y = 396.3.

The preceding analysis considers how the means and variances in firing depend on sound frequency but doesn't directly investigate whether tuning curves change in systematic ways, such as changes in the width or peak (best frequency).

To investigate possible changes in tuning curve width, we implemented the equivalent rectangular receptive field analysis (ERRF) from Lee and Middlebrooks (2011). Briefly, the ERRF divides the area under the tuning curve by the peak firing rate, creating a metric that incorporates the number of frequencies to which the unit responds based in part on the level of activity evoked by those stimuli. For example, many of the cells display a suppression in the dual sound condition (Fig. 4 A-C), yet these suppressions seem to be of different kinds. Generally, suppressions that compress the whole firing rate range (Fig. 4B & C) tend to increase ERRF width while suppressions that affect some conditions but leave the peak untouched (Fig. 4A) tend to decrease the ERRF width.

Figure 6 ERRF width is larger in dual-sound conditions. (**A**) The width of the ERRF for each unit in the single sound condition plotted against the width of the ERRF for each unit plotted in the 420 Hz paired condition. Each point represents a unit, out of 63 frequency modulated cells. The dashed line indicates unity, or ERRFs of equal width in the single sound and 420 Hz paired condition. Points above unity indicate ERRFs of greater width in the dual-sound condition. Points below unity indicate ERRFs of greater width in the single-sound condition. (**B**) The distribution of change in ERRF width per cell (ERRF$_{420}$ – ERRF$_{Single}$). A value of 0 indicates the ERRF width did not change across conditions. Values greater than 0 indicate the ERRF is broader, while values less than 0 indicated the ERRF is narrower. The ERRF width in the 420 Hz dual sound condition is significantly broader (mean = 0.75±0.11 S.E.M, $p < 0.05$). The blue vertical line indicates the mean of the distribution. The black vertical line is at 0. (**C**) The same as (**A**) except for the 2000 Hz paired condition. (**D**) The same as (**C**) except for the 2000 Hz paired condition. The mean in the 2000 Hz paired condition is significantly broader (mean = 0.26 ± 0.11 S.E.M, $p < 0.05$).

To compare tuning curve width between single and dual sound conditions, the distribution of each were plotted against each other (Fig. 6B & D). If the ERRF widths were similar across single and dual sound conditions the distribution would fall upon the unity line (black line in Fig. 6B, D). If the ERRF widths were narrower in the dual sound condition the distribution would be shifted below unity and if the ERRF widths were narrower in the single sound condition the distribution would be shifted above unity. It is quite apparent that ERRF widths are broader in the dual sound conditions, for both high- and low-frequency paired sounds.

To evaluate whether these width changes are significant, the difference between each neuron's dual sound ERRF width and single sound ERRF width was calculated. The 420 Hz paired ERRF widths were significantly broader than the single sound ERRF widths (Fig. 6C); the difference distribution was significantly greater than 0 (mean = 0.75±0.11 S.E.M, $p < 0.05$). Although the effect was smaller, the 2000 Hz paired ERRF widths were also significantly broader than the single sound EERF widths (Fig. 6D); the difference distribution was significantly greater than 0 (mean = 0.26 ± 0.11 S.E.M, $p < 0.05$). In summary, across the population the majority of neurons actually broaden their tuning curve in response to presentation of two simultaneous sounds.

Figure 7 The presence of an additional sound does not cause a large shift in best frequency (BF) across the population of recorded cells. (**A**) The single-sound best frequency plotted against the 420 Hz paired best frequency in the 17 cells that have best frequencies between 595-1414 Hz (grey lines). The bold line indicates the mean best frequency across the population. There is no significant shift in best frequency across the two conditions (single mean = 759.3 ± 76.48 S.E.M vs. dual mean = 937.6 ± 113.6 S.E.M, p = 0.072). (**B**) The same as (**A**) but for the 2000 Hz paired condition. There is no significant shift in best frequency across the two conditions (single mean = 759.3 ± 76.48 S.E.M vs. dual mean = 740.9 ± 75.67 S.E.M, p = 0.56).

Could some neurons shift their tuning curves, exhibiting a different best frequency in response to dual sounds (e.g. Fig 4D)? To investigate if neurons shifted their tuning, we selected neurons with firing rates significantly modulated by frequency and with single sound best frequencies that fell within the middle six A sounds (595-1414 Hz), leaving out the edge cases for which a shift outside the tested range could not in principle be observed (N=17). We then compared their single sound best frequency to their best frequency in the dual sound conditions. Seventeen neurons met these criteria. Figure 7 shows how best frequency changes in the 420 Hz

paired condition (Fig. 7A) or the 2000 Hz paired condition (Fig. 7B). Although, there is a slight shift of the mean best frequency away from the paired sound, it is not significant for either the 420 Hz dual sound (single mean = 759.3 ± 76.48 S.E.M vs. dual mean = 937.6 ± 113.6 S.E.M, p = 0.072) or the 2000 Hz dual condition (single mean = 759.3 ± 76.48 S.E.M vs. dual mean = 740.9 ± 75.67 S.E.M, p = 0.56). These findings suggest that shifting of receptive fields is not a ubiquitous computation within the population and is unlikely to substantially contribute to the encoding of simultaneous sounds.

In short, the modulations of frequency tuning on dual sound trials do not appear to be optimized for the preservation of information about either sound. To determine the impact of multiple sounds on the efficacy of coding stimulus related information, we evaluated how this code might be read out.

We first used a population-maximum-likelihood decoder to infer the condition from the observed spike count on a set of held-out trials (see Methods). Spike counts were assumed to be Poisson-distributed, and neurons were assumed to be independent of each other (a reasonable assumption given they were recorded at different times). For each held-out trial, the log-likelihood for each condition was summed across the population and the condition with the maximum-population-log-likelihood was predicted to be the held-out trials condition. This computation was repeated 1000 times per condition.

Overall, the decoder inferred every condition above chance (chance = 1/26). Figure 8A, C, and E shows the proportion of correctly predicted conditions across the 3 paired sound trial-types, sounds presented in isolation, sounds paired with 420 Hz, or sound paired with 2000 Hz, respectively. Qualitatively, the mean across all sounds presented in isolation (~66.7% correct) was higher than the mean across all 420 Hz paired conditions (~47.1% correct) or the mean across all 2000 Hz paired conditions(~51.8% correct). The likelihood functions show that the confusion in decoding tends to come from proximal conditions (Fig. 8B, D, F). This can be further seen in the confusion matrix (Fig. 9A) where the off diagonals are slightly darker in the dual compared to single sound conditions. This confusion arises due to the increase in likelihood when a low frequency sound is present (420 Hz paired and when 2000 Hz is paired with a low frequency: Fig. 9B), which may stem from the populations low frequency bias (Fig. 2). Ultimately, the changes in the frequency response functions seem to degrade the amount of information concerning each sound, resulting in worse decoding on dual sound trials.

Until now all the analyses presented probed how monkey IC encoding of sound changes in the presence of an additional sound. With the implementation of the maximum-likelihood

decoder, we can now ask if single and dual sounds are encoded with similar neural codes. To investigate if the monkey IC uses a similar code for the single and dual sound conditions, we attempted to decode the dual sound condition using the decoding weights learned from the single sound trials. Each held-out dual sound trial could, therefore, be labeled as one of the 8 A sounds. Performance was markedly worse in the 420 Hz condition when decoded with the single sound weights (accuracy = 16.5%; chance 12.5%) compared to the dual sound weights (accuracy ~ 47.1%; chance ~ 3.846%). This is evident in the confusion matrix (Fig. 10A) which shows 595 Hz is the most predicted stimulus for seven of the eight conditions. The performance in the 2000 Hz condition when decoded with the single sound weights (accuracy = 40.65%; chance = 12.5%) is also worse compared to decoding with the dual sound weights (accuracy ~ 51.8% correct; chance ~3.846%). Unlike the 420 Hz paired condition, the 2000 Hz paired conditions seem to confuse proximal conditions (Fig. 10B). These results show that the IC would need to use different coding strategies in the single and dual sound conditions, implying an IC readout may need to implement a different set of weights depending on the number of sounds present.

Figure 8 Accuracy of population maximum-log-likelihood decoding and mean log-likelihood functions across conditions. (**A**) The accuracy of the population (N = 93) maximum-log-likelihood decoder across each single sound condition. Each point corresponds to the percent of correctly labeled repeats (out of 1000) for each condition. Chance level prediction is 1/26 or ~ 3.85% (**B**) The mean population log-likelihood function for each condition across all potential predicted conditions. Each condition had the chance of being labeled as any one of the 26 single or dual conditions. (**C**) The same as panel (**A**) but for 420 Hz paired conditions. (**D**) The same as panel (**B**) but for 420 Hz paired conditions. (**E**) The same as panel (**A**) but for 2000 Hz paired conditions. (**F**) The same as panel (**B**) but for 2000 Hz paired conditions.

Figure 9 Confusion and mean-likelihood matrices for the decoder. (**A**) The confusion matrix for the decoder. Each column corresponds to the actual condition and each row corresponds to the predicted condition. The color of each square is the proportion of predicted trials (out of 1000) that fell into that bin. The diagonal corresponds to correct predictions. (**B**) A matrix of mean log-likelihood across conditions for the decoder. Each row corresponds to the likelihood function for that condition while each column corresponds to the mean log-likelihood for that condition across all actual conditions (e.g. leftmost column corresponds to the mean log-likelihood for 420 Hz across all conditions).

Figure 10 Decoding the paired condition with the single-sound weights impairs performance. (**A**) The confusion matrix for decoding 420 Hz paired conditions using the weights learned from single-sound trials. Each column corresponds to the actual condition and each row corresponds to the predictable condition. The color of each square is the proportion of predicted trials (out of 1000) that fell into that bin. The diagonal corresponds to correct predictions. (**B**) The same as (**A**) but for decoding the 2000 Hz paired conditions.

Discussion

How the brain simultaneously encodes multiple items, particularly when more than one item falls within a given neuron's receptive field, is an often-overlooked problem in neuroscience. It is assumed the population representation overcomes ambiguity in single unit encoding of stimuli, but the problem remains if the items recruit largely overlapping populations. To address this question, we first verified that monkeys could report the occurrence of two sounds whose center frequencies are separated by as little as 0.19 octaves (and whose bandwidth overlaps). We then investigated how frequency response functions of are affected by the presence of that second, concurrently presented sound. We found that the sounds we presented recruited largely overlapping populations (Fig. 2 & 3), and that on dual sound trials the majority of these cells' firing rates were less modulated by sound frequency (Fig. 5) and had their tuning curves broadened (Fig. 6). There was little evidence of systematic shifts in best frequency (Fig. 7).

Overall, these changes in frequency response functions reduce the information available to the maximum-likelihood decoder, decreasing the decoding accuracy in the dual sound conditions (Fig. 8 & 9). The poor decoding performance when using the single sound weights to decode dual sound conditions (Fig. 10) suggests that a different read-out is needed depending on the number of sounds – a startling possibility given that the brain cannot have prior knowledge of how many sounds are in the environment except by virtue of what it detects via sensory input.

A limitation of this study is that it is not known how many neurons might be needed to support perception of any given stimulus. It is possible that the small proportion of neurons that responded to only one of a pair of sounds were responsible for the monkeys' perceptual abilities.

Our results seem to be in-line with prior work showing degradation of information when there are multiple stimuli presented (Day & Delgutte, 2013; Day et al., 2012; Henry & Kohn, 2020). If this reduction in information scales with number of stimuli it could potentially explain the finding that the number of identifiable sounds saturates at three (Zhong & Yost, 2017). Generally, these results refute the notion that the changes in frequency response functions could be used to overcome the multiplicity problem and support recent evidence of alternations in firing rates between those evoked by individual stimuli (Caruso et al., 2018) may be a more fruitful research direction.

In the present study, we made the simplifying assumption that the time-and-trial-pooled average response of a neuron to a combination of stimuli is reflective of the information that the neuron encodes. If the underlying activity on dual sound trials actually fluctuates between the A-like and B-like response patterns, the time-and-trial-pooled average will be a poor measure of the information present in neural signals. The fact that average responses to AB sounds were generally between the average responses to A and B sounds presented alone supports the possibility that fluctuations may underlie at least a portion of the results observed here. Future work will test this possibility.

In principle, changes in frequency tuning as investigated here and fluctuating activity patterns as investigated previously (Caruso et al., 2018) both have the potential to limit the degree to which a given neuron is faced with the task of encoding more than one stimulus at a time. It remains possible that information-preserving changes in frequency tuning might be more evident when tested with a wider range of sound frequencies, allowing a fuller exploration of frequency tuning curves. However, the granularity of our testing, 0.19 octave spacing, was clearly coarser than monkeys' perceptual abilities, and the overall range, 2 octaves, should have

been adequate to demonstrate an effect if changes in frequency tuning were a major contributor to these perceptual abilities.

References:

Aitkin, L. M., & Phillips, S. C. (1984). Is the inferior colliculus and obligatory relay in the cat auditory system? *Neuroscience Letters, 44*(3), 259-264. doi:https://doi.org/10.1016/0304-3940(84)90032-6

Best, V., Schaik, A. v., & Carlile, S. (2004). Separation of concurrent broadband sound sources by human listeners. *The Journal of the Acoustical Society of America, 115*(1), 324-336. doi:10.1121/1.1632484

Bulkin, D. A., & Groh, J. M. (2011). Systematic mapping of the monkey inferior colliculus reveals enhanced low frequency sound representation. *Journal of Neurophysiology, 105*(4), 1785-1797. doi:10.1152/jn.00857.2010

Bulkin, D. A., & Groh, J. M. (2012a). Distribution of eye position information in the monkey inferior colliculus. *Journal of Neurophysiology, 107*(3), 785-795. doi:10.1152/jn.00662.2011

Bulkin, D. A., & Groh, J. M. (2012b). Distribution of visual and saccade related information in the monkey inferior colliculus. *Frontiers in Neural Circuits, 6*(61). doi:10.3389/fncir.2012.00061

Capuano, U., & McIlwain, J. T. (1981). Reciprocity of receptive field images and point images in the superior colliculus of the cat. *Journal of Comparative Neurology, 196*(1), 13-23. doi:10.1002/cne.901960103

Caruso, V. C., Mohl, J. T., Glynn, C., Lee, J., Willett, S. M., Zaman, A., . . . Groh, J. M. (2018). Single neurons may encode simultaneous stimuli by switching between activity patterns. *Nature Communications, 9*(1), 2715. doi:10.1038/s41467-018-05121-8

Chettih, S. N., & Harvey, C. D. (2019). Single-neuron perturbations reveal feature-specific competition in V1. *Nature, 567*(7748), 334-340. doi:10.1038/s41586-019-0997-6

Connell, M. N., Barczak, A., Schroeder, C. E., & Lakatos, P. (2014). Layer Specific Sharpening of Frequency Tuning by Selective Attention in Primary Auditory Cortex. *The Journal of Neuroscience, 34*(49), 16496. doi:10.1523/JNEUROSCI.2055-14.2014

Day, M. L., & Delgutte, B. (2013). Decoding Sound Source Location and Separation Using Neural Population Activity Patterns. *The Journal of Neuroscience, 33*(40), 15837. doi:10.1523/JNEUROSCI.2034-13.2013

Day, M. L., Koka, K., & Delgutte, B. (2012). Neural encoding of sound source location in the presence of a concurrent, spatially separated source. *Journal of Neurophysiology, 108*(9), 2612-2628. doi:10.1152/jn.00303.2012

Freedman, D. J., Riesenhuber, M., Poggio, T., & Miller, E. K. (2005). Experience-Dependent Sharpening of Visual Shape Selectivity in Inferior Temporal Cortex. *Cerebral Cortex, 16*(11), 1631-1644. doi:10.1093/cercor/bhj100

Groh, J. M., Kelly, K. A., & Underhill, A. M. (2003). A Monotonic Code for Sound Azimuth in Primate Inferior Colliculus. *Journal of Cognitive Neuroscience, 15*(8), 1217-1231. doi:10.1162/089892903322598166

Groh, J. M., Trause, A. S., Underhill, A. M., Clark, K. R., & Inati, S. (2001). Eye Position Influences Auditory Responses in Primate Inferior Colliculus. *Neuron, 29*(2), 509-518. doi:https://doi.org/10.1016/S0896-6273(01)00222-7

Hartline, H. K., & Ratliff, F. (1957). Inhibitory interaction of receptor units in the eye of Limulus. *The Journal of general physiology, 40*(3), 357-376. doi:10.1085/jgp.40.3.357

Henry, C. A., & Kohn, A. (2020). Spatial contextual effects in primary visual cortex limit feature representation under crowding. *Nature Communications, 11*(1), 1687. doi:10.1038/s41467-020-15386-7

Imig, T. J., Irons, W. A., & Samson, F. R. (1990). Single-unit selectivity to azimuthal direction and sound pressure level of noise bursts in cat high-frequency primary auditory cortex. *Journal of Neurophysiology, 63*(6), 1448-1466. doi:10.1152/jn.1990.63.6.1448

The Inferior Colliculus. (2005). (J. A. Winer & C. E. Schreiner Eds. 1 ed.): Springer-Verlag New York.

Jazayeri, M., & Movshon, J. A. (2006). Optimal representation of sensory information by neural populations. *Nature Neuroscience, 9*(5), 690-696. doi:10.1038/nn1691

Jenkins, W. M., & Masterton, R. B. (1982). Sound localization: effects of unilateral lesions in central auditory system. *Journal of Neurophysiology, 47*(6), 987-1016. doi:10.1152/jn.1982.47.6.987

Judge, S. J., Richmond, B. J., & Chu, F. C. (1980). Implantation of magnetic search coils for measurement of eye position: An improved method. *Vision Research, 20*(6), 535-538. doi:https://doi.org/10.1016/0042-6989(80)90128-5

Kelly, J. B., & Kavanagh, G. L. (1994). Sound localization after unilateral lesions of inferior colliculus in the ferret (Mustela putorius). *Journal of Neurophysiology, 71*(3), 1078-1087. doi:10.1152/jn.1994.71.3.1078

Lee, C.-C., & Middlebrooks, J. C. (2011). Auditory cortex spatial sensitivity sharpens during task performance. *Nature Neuroscience, 14*(1), 108-114. doi:10.1038/nn.2713

Mickey, B. J., & Middlebrooks, J. C. (2003). Representation of Auditory Space by Cortical Neurons in Awake Cats. *The Journal of Neuroscience, 23*(25), 8649. doi:10.1523/JNEUROSCI.23-25-08649.2003

Middlebrooks, J. C., Clock, A. E., Xu, L., & Green, D. M. (1994). A panoramic code for sound location by cortical neurons. *Science, 264*(5160), 842. doi:10.1126/science.8171339

Middlebrooks, J. C., & Pettigrew, J. D. (1981). Functional classes of neurons in primary auditory cortex of the cat distinguished by sensitivity to sound location. *The Journal of Neuroscience, 1*(1), 107. doi:10.1523/JNEUROSCI.01-01-00107.1981

Pages, D. S., Ross, D. A., Puñal, V. M., Agashe, S., Dweck, I., Mueller, J., . . . Groh, J. M. (2016). Effects of Electrical Stimulation in the Inferior Colliculus on Frequency Discrimination by Rhesus Monkeys and Implications for the Auditory Midbrain Implant. *The Journal of Neuroscience, 36*(18), 5071. doi:10.1523/JNEUROSCI.3540-15.2016

Perrott, D. R. (1984a). Concurrent minimum audible angle: A re-examination of the concept of auditory spatial acuity. *The Journal of the Acoustical Society of America, 75*(4), 1201-1206. doi:10.1121/1.390771

Perrott, D. R. (1984b). Discrimination of the spatial distribution of concurrently active sound sources: Some experiments with stereophonic arrays. *The Journal of the Acoustical Society of America, 76*(6), 1704-1712. doi:10.1121/1.391617

Porter, K. K., Metzger, R. R., & Groh, J. M. (2006). Representation of Eye Position in Primate Inferior Colliculus. *Journal of Neurophysiology, 95*(3), 1826-1842. doi:10.1152/jn.00857.2005

Porter, K. K., Metzger, R. R., & Groh, J. M. (2007). Visual- and saccade-related signals in the primate inferior colliculus. *Proceedings of the National Academy of Sciences, 104*(45), 17855. doi:10.1073/pnas.0706249104

Robinson, D. A. (1963). A Method of Measuring Eye Movemnent Using a Scieral Search Coil in a Magnetic Field. *IEEE Transactions on Bio-medical Electronics, 10*(4), 137-145. doi:10.1109/TBMEL.1963.4322822

Ryan, A., & Miller, J. (1978). Single unit responses in the inferior colliculus of the awake and performing rhesus monkey. *Experimental Brain Research, 32*(3), 389-407. doi:10.1007/BF00238710

Versnel, H., Zwiers, M. P., & van Opstal, A. J. (2009). Spectrotemporal Response Properties of Inferior Colliculus Neurons in Alert Monkey. *The Journal of Neuroscience, 29*(31), 9725. doi:10.1523/JNEUROSCI.5459-08.2009

Wallisch, P. (2014). Chapter 21 - Neural Decoding I: Discrete Variables. In P. Wallisch, M. E. Lusignan, M. D. Benayoun, T. I. Baker, A. S. Dickey, & N. G. Hatsopoulos (Eds.), *MATLAB for Neuroscientists (Second Edition)* (pp. 329-336). San Diego: Academic Press.

Weinberger, N. M. (1995). Dynamic Regulation of Receptive Fields and Maps in the Adult Sensory Cortex. *Annual Review of Neuroscience, 18*(1), 129-158. doi:10.1146/annurev.ne.18.030195.001021

Zhong, X., & Yost, W. A. (2017). How many images are in an auditory scene? *The Journal of the Acoustical Society of America, 141*(4), 2882-2892. doi:10.1121/1.4981118

Zwiers, M. P., Versnel, H., & Van Opstal, A. J. (2004). Involvement of Monkey Inferior Colliculus in Spatial Hearing. *The Journal of Neuroscience, 24*(17), 4145. doi:10.1523/JNEUROSCI.0199-04.2004