Class A Penicillin-Binding Protein-mediated cell wall synthesis promotes structural integrity during peptidoglycan endopeptidase insufficiency

Shannon G. Murphy[1,2], Andrew N. Murtha[2], Ziyi Zhao[1*], Laura Alvarez[3], Peter Diebold[2], Jung-Ho Shin[1], Michael S. VanNieuwenhze[4], Felipe Cava[3] and Tobias Dörr[1,2,5#]

[1] Weill Institute for Cell and Molecular Biology, Cornell, University, Ithaca, NY 14853, USA

[2] Department of Microbiology, Cornell University, Ithaca NY 14853, USA

[3]The Laboratory for Molecular Infection Medicine Sweden (MIMS), Department of Molecular Biology. Umeå University, Umeå, Sweden.

[4]Department of Molecular and Cellular Biochemistry and Department of Biology, Indiana University, Bloomington, IN, USA

[5] Cornell Institute of Host-Microbe Interactions and Disease, Cornell University, Ithaca NY 14853, USA

*present address:

Department of Biochemistry and Biophysics, University of California, San Francisco, San Francisco, CA 94158, USA.

[#] address correspondence to tdoerr@cornell.edu

Running title: Endopeptidase depletion effects on cell wall synthesis

Keywords: Peptidoglycan, autolysin, endopeptidase, M23, LysM, Penicillin-binding Protein, mreB

Abstract

The bacterial cell wall is composed primarily of peptidoglycan (PG), a poly-aminosugar that is essential to sustain cell shape, growth and cellular structural integrity. PG is synthesized by two different types of PG synthase complexes (class A Penicillin-binding Proteins [PBP]s/Lpos and Shape, Elongation, Division, Sporulation [SEDS]/class B PBP pairs) and degraded by 'autolytic' enzymes (*e.g.*, endopeptidases, EPs) to accommodate growth processes. It is thought that autolsyin activity (and particulary the activity of EPs) is required for PG synthesis and incorporation by creating gaps that are patched and paved by PG synthases, but the exact relationship between autolysins and the separate synthesis machineries remains incompletely understood. Here, we have probed the consequences of EP depletion for PG synthesis in the diarrheal pathogen *Vibrio cholerae.* We found that EP depletion resulted in severe morphological defects, increased cell mass, and a decline in viability, but continuing (yet aberrant) incorporation of cell wall material. Mass increase and cell wall incorporation proceeded in the presence of Rod system inhibitors, but was abolished upon inhibition of aPBPs. However, the Rod system remained functional (i.e., exhibited sustained directed motion) even after prolonged EP depletion, apparently without effectively inserting significant PG material. Lastly, heterologous expression of an EP from *Neisseria gonorrhoeae* could fully complement growth and morphology of an EP-insufficient *V. cholerae.* Overall, our findings suggest that in *V. cholerae,* only the Rod system absolutely requires endopeptidase activity (but not necessarily direct interaction with EPs) for productive PG incorporation, whereas aPBPs are able to engage in sacculus construction even during severe EP insufficiency.

Importance

Synthesis and turnover of the bacterial cell wall must be tightly co-ordinated to avoid structural integrity failure and cell death. Details of this coordination are poorly understood, particularly if and how cell wall turnover enzymes ("autolysins", *e.g.*, endopeptidases, EPs) are required for activity of the different cell wall synthesis machines, the Rod system and the class A penicillin-binding proteins (aPBPs). Our results suggest that in *Vibrio cholerae*, endopeptidases are required only for cell expansion mediated by the Rod system, while the aPBPs maintain structural integrity during EP insufficiency. Overall, our results imply a complex relationship between cell wall synthesis and cleavage and suggest that aPBPs are more versatile than the Rod system in their ability to repair cell wall gaps formed by autolysins other than the major EPs, adding to our understanding of the co-ordination between autolysins and cell wall synthases.

Introduction

Most bacteria elaborate a cell wall composed primarily of peptidoglycan (PG), which consists of polymerized N-acetyl glucosamine-N-acetyl muramic acid (poly-GlcNAc-MurNAc) dimers. These polymerized strands are covalently linked to each other via their oligopeptide side stems extending from the MurNAc residues; the degree of crosslinking varies with bacterial species and growth conditions (1-3). As such, PG encases the cell in a net-like structure that functions to maintain the high intracellular pressure accumulating in most bacteria and thus to prevent the cell from lysing. In concert with maintenance of

structural integrity, PG has to accommodate growth processes (cell elongation and size

expansion) and is therefore constantly degraded and resynthesized (4-6).

In many rod-shaped Gram-negative bacteria, cell wall synthesis during cell elongation is

mediated by two separate types of cell wall synthase complexes: the Rod complex (which

includes the glycosyltransferase RodA in conjunction with a class B Penicillin-Binding

Protein [bPBP] and accessory proteins) and the class A PBPs in conjunction with their

lipoprotein activators (7-12). The differential physiological roles of these seemingly

redundant systems have only recently been begun to be dissected (13, 14), but remain

incompletely understood.

Cell wall degradation, on the other hand, is mediated by a plethora of so-called

"autolysins", *i.e.*, enzymes with the capability to break bonds in the PG sacculus (15-17).

Members of one such group of autolysins, the endopeptidases (EPs), cleave the

oligopeptide crosslinks between PG strands, presumably to allow for insertion of new PG

material during cell elongation (18-21). To ensure structural integrity, EP-mediated cell

wall cleavage and Rod- and/or aPBP-mediated resynthesis should logically be tightly

coordinated, and this has indeed been demonstrated for cell elongation in Gram-positive

bacteria (22-25). When the putative coordination is perturbed (*e.g.*, after exposure to a

cell wall synthesis inhibitor), PG structural integrity often catastrophically fails and cells

die (26); this is one of the reasons why cell wall synthesis inhibitors (e.g., the β-lactams)

rank highly among our most powerful antimicrobials (27). EPs in particular are a double-

edged sword as they can both promote cell wall synthesis (28) and play major roles in

cell wall cleavage after beta lactam exposure (29, 30). However, how EPs are regulated

has only begun to be unravelled (31-34), and at least in Gram-negative bacteria we lack

a complete understanding of how EP cleavage activity relates to PG synthesis by the two

distinct cell wall synthase complexes.

Several models have been advanced to explain coordination of synthesis and

degradation, with a prominent model being a "make before break" mechanism, where a

nascent PG layer scaffold is elaborated parallel to an existing one, followed by cleavage

of the old material that has been relieved of its critical structural function through this load-

bearing stabilizer of new PG (35, 36). Alternatively, PG might be able to sustain several

cleavage events without experiencing catastrophic structural failure, obviating the need

for any coordination between synthesis and degradation for as long as the Rod system

and/or aPBPs are efficient enough in recognizing gaps in PG, *e.g.*, through interaction

with their cognate OM-localized activators in case of the aPBPs (a "break before make"

model).

Here, we show that in the cholera pathogen *Vibrio cholerae,* EP activity is not required

for cell wall synthesis *per se*. During EP insufficiency, growth and PG accumulation

continue in the presence of Rod system inhibitors but abruptly stop upon inhibition of

aPBPs. However, the Rod system continues directed motion for extended periods of EP

depletion. Lastly, a heterologously expressed EP can fully complement growth and

morphology of an EP-deficient *V. cholerae* strain. Our data thus suggest that aPBPs do

not require wild-type levels of crosslink cleavage for productive PG incorporation, while

EP activity is required for the Rod system to contribute significantly to cell wall growth.

Our cross-species complementation experiments intriguingly raise the possibility that

direct co-ordination between EPs and cell wall synthases might not be necessary at all,

at least under standard laboratory growth conditions.

Results

Cell wall incorporation and mass increase continue during endopeptidase

***insufficiency in* V. cholerae**

Endopeptidase depletion was previously shown to preclude insertion of new cell wall

material in *E. coli,* resulting in rapid cell lysis (21)*.* In contrast, we have noticed during EP

depletion experiments with *Vibrio cholerae* that the cholera pathogen did not lyse, even

in the absence of all 6 of its major D,D-EPs. This $\Delta 6$ endo strain ($\Delta shyA\ \Delta shyB\ \Delta shyC$

$\Delta vc1537\ \Delta vca0843,\ \Delta vca1043$ P_{IPTG}:*shyA*), has the remaining, conditionally essential EP

ShyA under control of an IPTG-inducible promoter and is thus suitable for depletion

experiments. Upon growing the $\Delta 6$ endo strain in the absence of inducer (reducing ShyA

to less than 10 % of initial levels after ~ 2h, **Fig. S1**), mass increase (measured by OD_{600})

continued at a rate similar to cells where *shyA* expression was induced by IPTG (**Fig.**

1A). When plated on solid media containing inducer, however, we observed a slight

decrease in cfu/mL over the depletion timecourse. (**Fig. S2A**), Thus, the ability to recover

and form colonies on a plate decreases during EP insufficiency, albeit without affecting

cell mass increase (OD_{600}). PG architecture analysis revealed that as expected, $\Delta 6$ endo

accumulated more PG crosslinks after depletion (38.5 % crosslinking after ShyA depletion

compared to 29.3 % in the WT) presumably due to the lack of EP cleavage activity (**Fig.**

S2CD). The increase in crosslinking provided additional evidence that functional EP

availability was highly limited under our depletion conditions. The analysis further

revealed a 65% increase in trimer formation, as well as a 32% increase in the amount of

anhydro residues upon ShyA depletion (**Fig. S2CD**). Additionally deleting the genes

encoding PBP4, PBP7 and VC1269 (which have predicted EP activity but are, based on

E. coli, not required for growth and cell elongation (37, 38)) did not appreciably affect

mass increase (except for a slight decrease in final yield after 6 h and a more pronounced

drop in cfu/mL) (**Fig. S2B**), demonstrating that the mass increase phenotype did not

simply reflect the ability of these putative EPs to substitute for ShyA.

We have previously shown that EP depletion in the Δ6 endo strain results in a dramatic

increase in cell diameter and ultimately the generation of giant, bulky and contorted cells

(34). Here, we asked to what extent these enlarged cells accumulated a PG cell wall. To

probe this, we cultured Δ6 endo in the presence of a fluorescent D-amino acid-derivative

(HADA) as a cell wall stain (39). Addition of HADA to ShyA-replete Δ6 endo cells resulted

in an even distribution of staining along the cell wall (**Fig. 1B**), as expected from wt cell

wall synthesis. In contrast, depleting ShyA resulted in a strikingly different pattern, where

large patches of HADA-reactive material accumulated throughout the cell. In principle,

these patches could be a remnant of incompletely-degraded cell wall material synthesized

before ShyA was completely depleted, or they could reflect the activity of L,D-

transpeptidases (which are able to incorporate HADA into the cell wall independent of cell

wall synthesis (39, 40)). We thus repeated the staining experiment in a Δ6 endo strain

lacking all L,D-transpeptidases (Δ*ldtAB*) , and under more stringent depletion conditions.

Following a 2 h depletion we added HADA for an additional hour; this still revealed an

accumulation of PG patches, strongly suggesting that PG synthesis and incorporation

continue under these conditions, albeit in an aberrant, non-directional way (**Fig. S3**).

Quantification of PG confirmed and expanded these observations – after 2 h of ShyA

depletion, cells accumulated ~18-fold more PG than ShyA-replete cells (when normalized

to OD_{600}) (**Fig. 1C**). Since these cells did not divide (**Fig. S2A**), PG accumulation was not

correlated with an increase in cell numbers, but suggested a buildup of cell wall in the

invididual, drastically enlarged ShyA-depleted cells. Consistent with a higher cell wall

content, ShyA-depleted $\Delta 6$ endo cells were almost 10-fold more resistant to osmotic

shock treatment (**Fig. 1D**). Thus, ShyA-depleted $\Delta 6$ endo cells not only incorporate PG,

but do so to higher levels than the WT, possibly reflecting the lack of EP-initiated turnover

processes. Similar observations have been made in autolysin-inactivated *B. subtilis,* a

Gram-positive bacterium (25, 41, 42)*.* While we cannot rule out that residual ShyA

remains in the cell following depletion (at levels too low to detect above background of

the non-specific band we observed via Western Blot in the same size range as ShyA, **Fig.**

S1), we can at a minimum conclude that *wild-type levels* of EPs are not necessary to

facilitate mass increase and incorporation of PG *per se*, but are essential for cell division

and likely key for the proper, directional integration of PG into the sacculus of *V. cholerae*.

Cell wall incorporation and mass increase in EP-deficient cells rely primarily on

aPBPs

We next addressed whether EP insufficiency affected the two cell wall synthesis

machines, the Rod system and the aPBPs differentially. To this end, we repeated our $\Delta 6$

endo depletion experiment in the presence of MP265 (an inhibitor of MreB (43)) or

moenomycin (an aPBP glycosyltransferase inhibitor (44)). Under ShyA-replete

conditions, mass increase proceeded at similar rates for both antibiotics and the untreated

control, while cfu/mL plateaued (moenomycin at 10 µg/mL, 8 x MIC) or decreased 10- 20-

fold (MP265 at 200 µM, 15 x MIC) in the presence of antibiotic (**Fig. 2A**, **Fig. S4A**). The

continued OD_{600} increase upon antibiotic exposure is consistent with our previous

observations that *V. cholerae* (as well as many clinically significant Gram-negative

pathogens) is remarkably tolerant to inhibitors of cell wall synthesis and forms cell wall-

deficient spheroplasts (in the presence of aPBP inhibitors) or spheroid cells containing

cell wall material (MP265) upon exposure to such agents (30, 45). Importantly, both

sphere cell types continue to increase in mass (30, 45), but fail to divide. Thus, OD_{600}

continues to increase while cfu/mL stagnates or declines.

Upon ShyA depletion, mass increase and HADA incorporation continued in the presence

of MP265 (at a similar rate compared to ShyA-replete conditions) (**Fig. 2B, Fig. S3**),

suggesting that the Rod system was not required for cell expansion during EP

insufficiency. Consistent with the OD_{600} data, visual inspection using phase microscopy

revealed intact, spheroid cells under MP265 exposure conditions when ShyA was

expressed (**Fig. 2C**). When ShyA was depleted, MP265-treated cells qualitatively

exhibited a rounder morphology than those in the untreated control, but were intact and

enlarged. Importantly, the mass increase, HADA incorporation and morphological

aberrations were recapitulated when another Rod system inhibitor, the bPBP2 inhibitor

mecillinam, was used (**Fig. S4C-F**).

In striking contrast to Rod system inhibition, moenomycin exposure completely abrogated growth of ShyA-depleted Δ6 endo cells (**Fig. 2B**). This coincided with accumulation of small cells and debris (indicative of lysis) (**Fig. 2C**), notably without strong HADA incorporation (**Fig. S3**). In addition, cells declined rapidly in viability in early stages (consistent with our previous observations (30)), though ultimately exhibited levels of survival similar to untreated or MP265-treated, ShyA-depleted Δ6 endo cells (**Fig. S4**). In summary, our data suggest that during EP-insufficiency, the aPBPs are essential for both mass increase and sustained PG incorporation, while the Rod system is not absolutely required.

MreB movement continues in EP-insufficient cells

The Rod-system, in conjunction with the actin homolog MreB, deposits new cell wall material during cell elongation while performing a rotational movement around the cell, apparently driven by aPBP-independent cell wall synthesis (46-48). Since the Rod system did not appear to contribute to PG synthesis during EP insufficiency, we asked whether EP depletion resulted in immobile Rod-complexes, similar to what has been observed during inhibition of cell wall synthesis (47). We constructed a functional (**Fig. S5**) mreBmsfGFP sandwich fusion in a Δ6 endo background and measured mreBmsfGFP velocity using epifluorescence and Total Internal Reflection Fluorescence (TIRF) microscopy. As a positive control, we confirmed that MP265 stopped MreB movement (**Fig. 3A**), as expected from what has been reported in *E. coli*. Mean square discplacement values indicated mixed populations of diffusive MreB particles and those

exhibiting directed motion under both ShyA replete and depleted conditions (**Fig. S6**).

Interestingly, MreB movement continued even after 3 h of ShyA depletion (**Fig. 3A**). albeit

at reduced velocity (decreasing from ~70 nm/s to ~ 40 nm/s) compared to ShyA-replete

conditions (**Fig. 3B**). Our estimates of MreB velocity under ShyA replete conditions were

higher than what has been reported previously for other bacteria (55 nm/s for *B. subtilis*

(49) and 10 nm/s for *E. coli* (47)), perhaps reflecting species-specific differences or

different properties of our sandwich fusion. Interestingly, the average size and number of

MreB clusters also increased under ShyA depletion conditions (**Fig. 3C-D**), suggesting

that EP depletion might affect Rod complex assembly dynamics. We conclude that similar

to what has been observed in *B. subtilis* (22), EP insufficiency does not result in

immediate inactivation of the Rod system, but changes its velocity and potentially the

stoichiometry of its assembly.

Complementation of EP-insufficiency in* V. cholerae *by expression of heterologous

EPs

So far, our results suggested that during EP insufficiency, aPBPs continue to synthesize

PG and the Rod system may remain functional, yet does not insert new PG material into

the sacculus. This might suggest that the Rod system requires a physical association with

one or more EPs for insertion of nascent PG material. Alternatively, EPs might catalyze

PG insertion independently, *e.g.*, through recognition of intrinsic PG substrate cues. To

gain a better understanding of the necessity for a physical interaction, we conducted

cross-species complementation experiments. To this end, we heterologously expressed

EPs from other bacteria (MepM from *E. coli* [henceforth "MepM$_{Eco}$"] and NGO_1686

[henceforth "MepM$_{Ngo}$"] from *Neisseria gonorrhoeae*) in Δ6 endo and observed their ability

to restore growth. Heterologously expressed EPs (particularly from the distantly related

N. gonorrhoeae, a BLAST alignment indicated 28 % identity to ShyA) are unlikely to

interact with any native *V. cholerae* enzymes, and should thus allow us to isolate their EP

activity from the interaction networks they might be embedded in. We expressed

arabinose-inducible EPs in Δ6 endo and measured differential growth in the presence of

IPTG (ShyA expression) vs. arabinose (heterologous EP expression). We found that

neither wild-type MepM$_{Eco}$ nor MepM$_{Ngo}$ were able to rescue growth of a Δ6 endo during

ShyA depletion conditions (**Fig. 4A, 5A**). However, we recently demonstrated that EPs

from diverse organisms (including *E. coli* and *N. gonorrhoeae*) are produced in an inactive

form due to the inhibitory function of their domain 1 and likely activated *in vivo* by an

unknown mechanism (34). Heterologously expressed enzymes are likely not subject to

this activation pathway in *V. cholerae* (especially if the activator is a protein) and we thus

instead expressed EP mutant versions with their inhibitory domain 1 deleted, which are

expected to be constitutively active. Surprisingly, both MepM$_{Eco}$^{ΔDom1} and MepM$_{Ngo}$^{ΔDom1}

fully complemented growth of the Δ6 endo strain to a similar degree as the native ShyA

(**Fig. 4A, 5A**). Upon visual inspection of Δ6 cells that rely on MepM$_{Eco}$^{ΔDom1} for growth

(ara+ condition), however, we noticed very few rod-shaped and a majority of sphere-

shaped cells (**Fig. 4B**). Thus, heterologous expression of MepM$_{Eco}$^{ΔDom1} can support

growth, but not wild-type shape of a Δ6 endo strain. In striking contrast, complementation

with MepM$_{Ngo}$^{ΔDom1} (but not its active site mutant derivative H373A) promoted both growth

(**Fig. 5A**) and the generation of rod-shaped cells (**Fig. 5B**). We sought to confirm that this

apparent complementation of rod shape was still dependent on MepM$_{Ngo}$^{ΔDom1} (rather than

a mutation derepressing *shyA* in Δ6 endo). Thus, we plated all strains on agar containing

IPTG, arabinose, or no inducer at the end of the experiments where we visualized cells

relying on MepM$_{Ngo}^{\Delta Dom1}$. All strains had the same low level of spontaneous supressors

able to grow in the absence of inducer (**Fig. S7**), confirming that the majority of the rod-

shaped cells observed when only MepM$_{Ngo}^{\Delta Dom1}$ was expressed are not suppressors. In

summary, these data demonstrate that heterologous expression of an activated EP can

be sufficient to restore both growth and proper cell shape to Δ6 endo cells.

Discussion

Bacteria must maintain a careful balance between cell wall cleavage and synthesis to

promote cell elongation/division, but the exact relationship between the two cell wall

synthases (Rod system vs. aPBPs) and cell wall hydrolases (*e.g.*, endopeptidases) is

poorly understood, at least in Gram-negative bacteria. Here, we have used EP depletion

and chemical inactivation experiments to dissect the interplay between cell wall cleavage

and synthesis in the cholera pathogen *V. cholerae.* Our key observation is that in *V.*

cholerae, cell wall synthesis and cell expansion (but not cell division) continue upon EP

depletion. This poses an apparent contradiction to data obtained in *E. coli,* where cell wall

incorporation was drastically reduced after EP depletion and cells started to lyse (21).

While ostensibly fundamental aspects of the coordination between cell wall synthesis and

cleavage may simply not be as well-conserved as one might expect, these observations

might also reflect species-specific differences in EP-independent cell wall turnover rates,

and not necessarily the consequences of EP depletion *per se*. It is possible that lysis

under EP-insufficient conditions in *E. coli* reflects generally higher PG degradation rates

(*E. coli*, for example encodes three amidases (50), while *V. cholerae* possesses only one

(51)). This would mask the underlying continued incorporation of new cell wall material in

the absence of EPs. Importantly, EP depletion in *E. coli* did result in a cell volume increase

prior to lysis (21), also supporting at least a transient continuation of PG synthesis during

EP insufficiency in this species.

Cell wall expansion during EP-insufficiency was surprising, since presumably any form of

cell wall synthesis that promotes the degree of cell expansion we observed in EP-deficient

V. cholerae should require some form of cleavage, likely catalyzed by other autolysins

(*e.g.* the amidase or LTGs). The incisions resulting from such cleavage, and/or the

autolysin(s) involved appear to be of limited utility to the Rod-system, but can be exploited

by the aPBPs. This suggests that aPBPs are more versatile in recognizing a variety of

cell wall cuts (independent of an actual physical connection with EPs), while the Rod

system absolutely requires either EP-mediated cleavage, or a physical association with

EPs (see a more detailed discussion below) for productive PG incorporation. These

interpretations are in line with several recent observations in *E. coli, i.e.,* aPBPs have

more of a damage repair function than a functional role in promoting cell elongation (14)

and that upregulated EP activity promotes aPBP function (28), likely indirectly through the

creation of PG incisions that allow for an interaction between aPBPs and their OM-

localized activators. Thus, EP cleavage is not strictly necessary for, but can promote,

aPBP activity. It is possible that under EP-insufficient conditions, the lytic

transglycosylases (the other major group of cell wall cleavage enzymes that cut the

polysaccharide backbone of PG (52)) create large open areas in PG that can be

recognized, and patched, by aPBPs; LTG activity would be consistent with the increase

in anhydro "caps" we observe in ShyA-depleted Δ6 endo PG.

The observation (consistent with what has been shown in *B. subtilis* (22)) that MreB

continues directed movement at least for some time during EP insufficiency suggests that

the Rod system does not actually require EP activity for assembly and RodA's

glycosyltransferase activity (which likely drives MreB movement). Similar to what has

been proposed for *B. subtilis,* we thus consider the "make-before-break" model as

proposed by Höltje/Koch (35, 36) a likely scenario for *V. cholerae* cell elongation via the

Rod system. In this model, the Rod system creates a second layer of PG that is

incorporated via EPs during or after synthesis. Generation of this second layer could at

first proceed independently of wild-type EP activity, but incorporation into the growing

sacculus would require crosslink cleavage.

As mentioned above, our data suggest that the Rod system requires EP activity, either

through physical association or recognition of EP cut sites in PG. Our cross-species

complementation experiments with an activated *N. gonorrhoeae* EP suggest that a

physical association might not be strictly necessary, unless the heterologously expressed

enzyme does somehow directly interact with the *V. cholera*e Rod system. We thus

consider a model where rather than (or in addition to) co-ordinating with cell wall

synthases directly, EPs can somehow specifically recognize and preferentially cleave old

PG that is adjacent to nascent PG.

An important caveat, however, is that the Δ6 endo strain still maintains a copy of *shyA*

under IPTG control. While the lac promoter is tightly repressed in the absence of inducer,

a small number of molecules under its control might still be produced (53). ShyA is

produced as an inactive precursor and the signal for activation is unkown (34). It is

conceivable that complementation with a heterologously expressed EP might somehow

enhance activation of this leaky background of ShyA molecules, *e.g.* if there is a positive

feedback loop between cell wall cleavage and native EP activation.

Taken together, our data suggest that two main cell wall synthases, the aPBPs and the

Rod system have differential relationships with autolysins, and especially

endopeptidases. While the Rod system likely relies on a "make-before-break" strategy,

the aPBPs seem capable of the reverse, "break-before-make", i.e. the ability to efficiently

recognize and patch holes in the cell wall sacculus. As such, our data also provide

additional support for a functional independence of the aPBPs and the Rod system (14,

54), at least during cell elongation.

Acknowledgements

This work was supported by the National Institutes of Health/NIGMS through

R01GM130971 to TD. Research in the Cava lab is supported by MIMS, the Knut and

Alice Wallenberg Foundation (KAW), the Swedish Research Council and the Kempe

Foundation. Research in the VanNieuwenhze lab is supported by the NIH through R01

GM113172 and R35 GM136365.

Materials and Methods

Bacterial growth conditions.

Cells were grown by shaking (200 rpm) at 37°C in 5 mL of LB unless otherwise indicated. Antibiotics, when appropriate were used as the following concentrations: streptomycin, 200 µg ml^{-1}; ampicillin, 100 µg ml^{-1}; chloramphenicol, 5 µg ml^{-1}; moenomycin, 10 µg ml^{-1}; MP265, 300 µM; and mecillinam,10 µg ml^{-1}. IPTG (200 µM) and arabinose (0.2%) were added for induction of P$_{iptg}$ and P$_{ara}$ promoters, respectively.

Plasmid and strain construction.

All bacterial strains and oligonucleotides used in this study are summarized in Table S1 and Table S2, respectively. All *Vibrio cholerae* strains are derivatives of El Tor strains N16961 (55) or E7946 (56), the latter was used for chitin-induced transformation. Δ6 endo construction is reported elsewhere (30).

Other strains were constructed by chitin-induced transformation of linear PCR products as described in (57). A chloramphenicol resistance cassette insertion into the gene *vc1807* (a well-established neutral locus) was used as the primary selector. The transforming fragment for vc1807::chl was constructed by amplifying upstream and downstream homology regions using primers PD079/PD097 and PD098/PD082, respectively. The chl gene coding for chloramphenicol acetyl transferase was amplified from pBAD33 (58) with primers PD095/PD096 and fused with the flanking homologies of vc1807 via isothermal assembly. For antibiotic resistance gene swapping, a vc1807::trim allele was also produced by amplifying upstream (using primers TDP597/598) and

downstream (primers TDP601/602) homologies of vc1807 and fusing them with a trimR

cassette amplified from V. cholera Haiti (59) (primers TDP599/600) using SOE PCR with

primers TDP603/604.

To construct a functional MreB-msfGFP-MreB sandwich fusion, upstream (primers

PD056/PD074) and downstream (primers PD071/PD057) homologies were amplified

from the *V. choerae* genome and fused via isothermal assembly with msfGFP (amplified

with primers PD054/PD055). Analogous to a published E. coli MreB-msfGFP sandwich

fusion (47), we replaced glycine 228 of MreB with this msfGFP. To enhance the probability

of success of finding a functional fusion, we used semi-degenerate primers to generate a

library of possible linker sequences. Flanking homologies, MreB and msfGFP were first

fused using isothermal assembly (60) and then amplified using nesting primers

PD104/PD105.The resulting upstream-MreB-linker-msfGFP-linker-MreB-downstream

PCR fragments were transformed into E7946 using chitin transformation with vc1807::chl

as the primary selector. 96 colonies were tested for growth rate and clone M2C was

chosen for further experiment due to its wild-type growth behavior. The linkers of this

fusion construct were sequenced (coding for DGVGG upstream of msfGFP and GTPIP

downstream).

Δ8 was constructed by transforming endopeptidase deletion PCR products into a parental

mreB::mreBmsfGFP ΔlacZ:P$_{IPTG}$:*shyA* strain. Deletion scars were amplified from Δ6 endo

and introduced in two steps into this parental background via chitin transformation. The

following primers were used to amplify the EP deletion fragments: *shyA* (TDP577/578),

shyC (TDP581/582), *shyB* (TDP579/580), *vc1537* (TDP583/584), *vc0843* (tagE1)

(TDP587/588), *vca1043* (tagE2) (TDP585/586). PBP4 and PBP7 deletions were

introduced into Δ6 endo by amplifying PCR fragments with upstream and downstream

homologies fused by a linker for chitin-mediated transformation using the following

primers: PBP4 upstream homology (TDP680/TDP681), downstream homology

TDP682/TDP683, fused using SOE PCR with nesting primers TDP691/692. PBP7

upstream homology (TDP676/TDP677), downstream homology TDP678/TDP679, fused

using SOE PCR with nesting primers TDP693/694.

Δ8 exhibited a very low transformation efficiency and we thus introduced the Δvc1269

deletion using homologous recombination with a suicide plasmid pCVD442 as described

(61). In brief, upstream and downstream homologies of vc1269 were amplified using

primers TD810/TD811 and TD812/TD813. These fragments were cloned into Xba1-

digested pCVD442 using isothermal assembly. pCVD442Δvc1269 was then introduced

into Δ8 endo via biparental mating (using SM10 as a donor strain) by mixing 10 µL of

each donor and recipient, followed by 6 h incubation at 37 ˚C, followed by selection for

single crossover strains and selection against the donor strain by plating on LB plates

containing carbenicillin (100 µg/mL), streptomycin (200 µg/mL) and IPTG (200 µM). A

single colony from the first crossover plate was then picked and streaked out on a plate

containing sucrose (10 %), streptomycin (200 µg/mL) and IPTG (200 µM). This plate was

incubated at ambient temperature for 3 days, after which 16 colonies were tested for the

correct knockout construct using vc1269 flanking primers TD814/TD815.

All plasmids were built using isothermal assembly (60). Genes were cloned into

pBADmob (a mobile pBAD33 derivative) using the following primer pairs: MepM$_{ECO}$,

TDP1342/TDP1340; MepM$_{ECO}$$^{\Delta dom1}$ TDP1341/TDP1340, MepM$_{NGO}$, SM861/862; and

MepM$_{NGO}$$^{\triangle D1}$, SM859/SM860. MepM$_{NGO}$$^{\triangle D1}$ was cloned into pTD101 (a *lacZ*

chromosomal delivery vector) using the primer pair SM991/SM992 and the pBAD

construct as a template for amplification. Point mutations were introduced into pBAD

plasmids carrying NGO1686 via Q5 site-directed mutagenesis (NEB, Ipswitch, MA, cat

#E0554S) with the following primer pairs: H373A (TDP1652/TDP1653) and E131R

(TD1368/TD1369). Plasmids were conjugated into *V. cholerae* using donor E. coli strains

(SM10 lamda pir or MFD lamda pir).

Phase contrast microscopy and HADA staining.

Cells were harvested (2 min at 12,000 rpm), spotted on a 0.8% agarose pad containing

PBS and imaged on a Leica Dmi8 inverted microscope. For HADA experiments, Δ6 endo

cells were grown in the presence of 50 µM HADA (3-[[(7-Hydroxy-2-oxo-2*H*-1-

benzopyran-3-yl)carbonyl]amino]-D-alanine hydrocholoride), washed once by pelleting 2

min at 12,000 rpm and resuspending in fresh LB). HADA stain was imaged in the DAPI

channel (395 nm [excitation]/460 [emission]) at 1 s exposure.

Endopeptidase depletion experiments

EP depletion strains were grown overnight in LB broth containing 200 µM IPTG. The next

day, cells were washed 2x by pelleting (2 minutes at 12000 rpm) and resuspending in LB

broth without inducer. Cells were then diluted 100-fold into fresh LB containing either 200

µM IPTG (IPTG+) or no inducer (IPTG-). Where indicated, antibiotics were used at 10

µg/mL (moenomycin, mecillinam) or 200 µM (MP265).

Single particle tracking by TIRF imaging

The Δ8 endo mreB::mreBmsfGFP^{sw} strain with chromosomally expressed MreB-msfGFP

was grown shaking at 37°C in LB medium supplemented with 100µM IPTG overnight.

The saturated cells were diluted (1:100) into fresh LB in two groups (with 100µM IPTG for

ShyA expression or without 100µM IPTG for ShyA depletion). After 2 hours of shaking

(220 rpm) incubation at 37°C, cells were harvested and spotted on a 0.8% agarose pad

containing M9 medium. Time-lapse TIRF imaging was performed on a Zeiss Elyra

equipped with an inverted Axio Observer.Z1 microscope and a 100x 1.46 oil objective.

The objective was heated at 37°C during imaging acquisition. The exposure time was 100

ms and inter-frame intervals were 2 s over a 2-min recording. The movement of MreB-

msfGFP was analyzed using single particle tracking software ImageJ TrackMate (62) and

MATLAB msdanalyzer (63).

The mean square displacements (MSD) of particles trajectories were calculated using

the msdanalyzer package and the motion types were analyzed through log-log fitting

(63). By setting the R^2 coefficient > 0.8, individual MSD curves were fitted and the

values of anomalous diffusion coefficient (α) indicates that MreB particles exhibit a mix

of dynamic behaviors (confined diffusion, $0.1 \leq \alpha < 0.9$; simple diffusion, $0.9 \leq \alpha < 1.1$;

directed motion, $\alpha \geq 1.1$) (64).

Peptidoglycan analysis

PG samples were analyzed as described previously (65). Briefly, 50 mL cultures of Δ6 endo were grown to early/mid exponential phase with or without IPTG (200 µM) for 2h, harvested and boiled in 5% SDS for 1 h. Sacculi were repeatedly washed by ultracentrifugation (110,000 rpm, 10 min, 20ºC) with MilliQ water until SDS was totally removed. Samples were treated with 20 µg Proteinase K (1 h, 37 °C) for Braun's lipoprotein removal, and finally treated with muramidase (100 µg/mL) for 16 hours at 37 ºC. Muramidase digestion was stopped by boiling and coagulated proteins were removed by centrifugation (14,000 rpm, 10 min). For sample reduction, the pH of the supernatants was adjusted to pH 8.5-9.0 with sodium borate buffer and sodium borohydride was added to a final concentration of 10 mg/mL. After incubating for 30 min at room temperature, the samples pH was adjusted to pH 3.5 with orthophosphoric acid.

UPLC analyses of muropeptides were performed on a Waters UPLC system (Waters Corporation, USA) equipped with an ACQUITY UPLC BEH C18 Column, 130Å, 1.7 µm, 2.1 mm X 150 mm (Waters, USA) and a dual wavelength absorbance detector. Elution of muropeptides was detected at 204 nm. Muropeptides were separated at 45°C using a linear gradient from buffer A (formic acid 0.1% in water) to buffer B (formic acid 0.1% in acetonitrile) in an 18-minute run, with a 0.25 ml/min flow.

Relative total PG amount was calculated by comparison of the total intensities of the chromatograms (total area) from three biological replicas normalized to the same OD600 and extracted with the same volumes. Muropeptide identity was confirmed by MS/MS analysis, using a Xevo G2-XS QTof system (Waters Corporation, USA). Quantification of

muropeptides was based on their relative abundances (relative area of the corresponding

peak) normalized to their molar ratio.

Western Blotting

Whole cell lysates (15 µg) were resolved by 10% SDS-PAGE and the proteins were

transferred to a PVDF membrane using a semi-drying transfer system (iBlot 2, Invitrogen).

The membrane was then blocked overnight with blocking solution containing 4% milk (dry

milk dissolved in 20 mM Tris-HCl (pH 7.8), 150 mM NaCl, 0.1% Triton X-100). Next day,

the membrane was incubated with anti-ShyA polyclonal antibody (1: 5,000, produced by

Pocono Rabbit Farm & Laboratory, PA) for two hours and then washed twice with 1xTBST

(20 mM Tris-HCl (pH7.8), 150 mM NaCl, 0.1% Triton X-100). The washed membranes

were then incubated with anti-rabbit secondary antibody (1:15,000, Li-Cor cat# 926-

32211) for 1 hour. Membranes were then washed three times with 1xTBST, scanned on

an Odyssey CLx imaging device (LI-COR Biosciences) and visualized using Image

Studio™ Lite Ver 5.2 software (Li-Cor) for signal quantification.

Figures

Figure 1. **Cell mass increase and cell wall incorporation continue during EP insufficiency. (A)** Overnight cultures of Δ6 endo (ΔshyABC Δvc1537 ΔtagE1,2

P$_{IPTG}$:*shyA*) grown in the presence of IPTG (200 μM) were washed twice and diluted

100fold into growth medium with (+IPTG) or without (-IPTG) inducer. At the indicated time

points, OD$_{600}$ was measured. Data are mean of six biological replicates, error bars

represent standard deviation. **(B)** Δ6 endo was treated as described in **(A)** in the presence

of HADA (100 μM). After 3 h, cells were washed twice and then imaged. **(C)** Relative PG

content of Δ6 endo was measured via UPLC analysis (see methods for details) after 2

hours of growth in the presence (IPTG +) or absence (IPTG -) of inducer. Error bars

represent standard deviation of 3 biological replicates. **(D)** Cells were treated as

described in **(A).** After 3 h of growth in the presence or absence of inducer, cells were

pelleted and resuspended in 20 mM NaCl (osmotic shock treatment) for 5 min. Shock

treatment was stopped by adding PBS to 180 mM. % survival is cfu/mL before treatment

divided by cfu/mL after treatment. Raw data points of 3 biological replicates are shown.

(C-D) Asterisks denote statistical difference via unpaired t-test (****, $p < 0.0001$; ***, $p <$

0.001, ** $p < 0.01$, * < 0.05). Scale bars, 5 μm

Figure 2. Cell mass increase during EP insufficiency relies on aPBP activity

Δ6 endo grown overnight in IPTG (200 µM) was washed twice and diluted 100-fold into

fresh medium containing either IPTG **(A)** or no IPTG (**B**) and either no antibiotic, the aPBP

inhibitor moenomycin (moeno, 10 µg/mL, 8x MIC) or the MreB inhibitor MP265 (300 µM,

15 x MIC). At the indicated time points, OD_{600} was measured; after 3 h of growth, cells

were also imaged (**C**). Data are averages of six biological replicates, error bars represent

standard deviation. Scale bar, 5 µm

Figure 3. MreB movement continues during EP insufficiency

Δ6 endo (**A**) or Δ8 endo (**B-D**) expressing an mreBmsfGFP^{sw} fusion from its native

chromosomal locus was diluted from an overnight culture grown in the presence of IPTG

into growth medium without inducer. After 3 hours, cells were imaged using

epifluorescence microscopy (**A**) or TIRF (**B-D**). MreB movement was analyzed using Fiji

(TrackMate). A representative single moving MreB focus track is shown in (**A**) (frames

are 2.5 s apart). Kymographs from 3 representative cells are also shown, in addition to

kymographs obtained from mobile MreB foci in the presence of inducer (+IPTG) and in

the presence of the MreB inhibitor MP265. (**B-D**) TIRF was used to assess MreB focus

velocity, mean cluster size and mean cluster number. Numbers above bar graphs

represent the number of imaged cells, error bars represent standard deviation.

Differences between IPTG+ and IPTG – were significant for all three graphs (t-test) at p

= 0.017 (velocity), 0.021 (cluster size) and 0.028 (cluster number), respectively.

Figure 4. Cross-species complementation of Δ6 endo phenotypes with an EP from

***Escherichia coli*.**

Δ6 endo carrying pBAD33 (arabinose-inducible) encoding MepM$_{Eco}$ or its Δdomain 1

derivative, was diluted and spot-plated on medium containing either IPTG (200 μM, ShyA

expressed), arabinose (0.2 %, heterologous EP expressed) or no inducer. Plates were

incubated at 37 ˚C for 24 hours and then imaged. (**C**) Δ6 endo carrying the indicated EP

under control of an arabinose-inducible promoter was grown without IPTG (chromosomal

ShyA off) and with arabinose (pBAD33-encoded EP on) for 3 hours and then imaged.

Scale bar, 5 μm

Figure 5. Cross-species complementation of Δ6 endo phenotypes with an EP from
Neisseria gonorrhoeae. Δ6 endo carrying pBAD33 (arabinose-inducible) expressing
MepM$_{Ngo}$ or MepM$_{Ngo}^{\Delta Dom1}$ was diluted and spot-plated on medium containing either IPTG
(200 µM, ShyA expressed), arabinose (0.2 %, MepM$_{Ngo}$) or no inducer. Plates were

584   incubated at 37 ˚C for 24 hours and then imaged. (**C**) Δ6 endo carrying the indicated EP

585   under control of an arabinose-inducible promoter was grown without IPTG (chromosomal

586   ShyA off) and with arabinose (pBAD33-encoded EP on) for 3 hours and then imaged. ss,

587   DsbA signal sequence.

Expected mass of ShyA w/o signal sequence: 46 kDa

Figure S1. ShyA levels after depletion in Δ6 endopeptidases mutant strains. (A) Wild

type, IPTG inducible ShyA in *Δ6* endopeptidases mutant and Δ*shyA* mutant strains were

grown overnight in LB broth containing 200 μM IPTG, then washed three times with fresh

LB. Cells were then inoculated 100fold into 150 ml pre-warmed LB medium without IPTG

for ShyA depletion and samples were collected at indicated time points. For the Western

Blotting, each 15 μg cell extracts were separated on 10% SDS-PAGE gels and subjected

600   to Western Blot analysis suing ShyA polyclonal antibody **(B)** ShyA band intensities were

601   quantified and subtracted with intensity value of non-specific background band detected

602   in the *ΔshyA* mutant. Residual ShyA protein levels were normalized to non-depleted ShyA

603   at time point T0 (100%).

Figure S2. Effects of EP depletion on growth, survival and PG composition.

N16961 $\Delta 6$ endo or SAD30 $\Delta 9$ endo (mreB::mreBmsfGFP^{sw}) was grown overnight in IPTG (200 µM), washed twice, and diluted 100-fold into fresh medium with or without inducer. At the indicated time points, OD$_{600}$ (**A**) was measured via spectrophotometry and cells were diluted serially onto LB IPTG (200 µM) plates to determine colony forming units per mL (**B**). Data are averages of 3 biological replicates, error bars represent standard deviation. (**C**) Chromatogram of PG composition of N16961 $\Delta 6$ endo cells harvested after 2 hours with (+) or without (-) IPTG (200 µM). (**D**) The table summarizes the relative molar abundance (%) of monomers, dimers, trimers shown in the chromatogram. Data regarding the % of crosslinkage (proportion of crosslinked peptide side chains, calculated on dimers and trimers content) is also included. Anhydro muropeptides (with a residue of (1-6 anhydro) N-acetyl muramic acid) are the terminal subunits of the sugar chains and hence used to calculate the chain length. Values are mean of three biological replicas. Percent change was calculated relative to the IPTG-treated sample and p-values were generated using a multiple comparisons t-test (****, $p < 0.0001$; ***, $p < 0.001$, ** $p < 0.01$, * < 0.05).

Figure S3. Continued PG incorporation upon Rod system inhibition.

An overnight culture of Δ6 endo Δ*ldtAB* was diluted into medium without inducer. After 2 hours of ShyA depletion, HADA (100 μM) was added for another 1 hour. Cells were then washed twice and imaged. For the antibiotic experiments, MP265 (200 μM, 15 x MIC) or moenomycin (moeno, 10 μg/mL, 8x MIC) was added for 1 h after the 2 h initial depletion, followed by 1 h addition of HADA. Scale bar, 5 μm.

10 µg/mL mecillinam

Figure S4. Mass increase during EP insufficiency relies on aPBPs, but not the Rod

system. Δ6 endo was grown overnight in IPTG (200 μM), washed twice, and diluted 100-

fold into fresh medium containing either IPTG **(A)** or no IPTG (**B**) and either no antibiotic,

the aPBP inhibitor moenomycin (moeno, 10 μg/mL, 8x MIC) or the MreB inhibitor MP265

(300 μM, 15 x MIC). At the indicated time points, cells were diluted serially and tittered

onto LB plates containing IPTG (200 μM). Data are averages of six biological replicates,

error bars represent standard deviation. (**C-E)** In a similar experiment, Δ6 endo was

treated with mecillinam (10 μg/mL, 20x MIC). At the indicated time points, OD_{600} (**C-D**)

was measured via spectrophotometry and cells were harvested and spotted on a 0.8%

agarose pad containing PBS for phase contrast microscopy (**E**). (**F**) The cell wall was

stained using HADA as described for **Fig. 1.** Scale bar, 5 μm.

Figure S5. Growth of the mreBmsfGFP strain compared to WT.

Wild-type SAD30 and mreBmsfGFP-containing derivative were grown overnight in LB.

Cells were diluted 1000-fold into fresh medium and 200-μl of each was loaded into a 100-

well plate. Growth of each culture was monitored by optical density at 600 nm (OD_{600}) in

a Bioscreen C plate reader (Growth Curves America).

A

IPTG +   IPTG -

B

IPTG +	alpha		ROI-1	ROI-3
confined motion	<0.9		34	42
simple diffusion	0.9≤ <1.1		10	19
directed motion	≥1.1		27	20
		total tracks (good fit)	71	81
		directed%	38.03%	24.69%

IPTG-	alpha		ROI-1	ROI-2
confined motion	<0.9		55	72
simple diffusion	0.9≤ <1.1		36	47
directed motion	≥1.1		56	89
		total tracks (good fit)	147	208
		directed%	38.10%	42.79%

Figure S6. Mean square displacement analysis. Δ6 endo mreBmsfGFP^{sw} was grown

with or without IPTG and imaged using TIRF microscopy. **(A)** Example MSD curves for 2

regions of interest (ROIs) for each, the IPTG+ and IPTG- condition, **(B)** alpha values and

% of MreBmsfGFP patches exhibiting directed motion.

Figure S7. Low suppressor background in Δ6 endo cells

Δ6 endo carrying pBAD33 (arabinose-inducible) expressing the indicated contructs

("pBAD" column) were diluted into fresh medium containing either no inducer, IPTG or

arabinose and grown for 3 hours ("experiment" column). Cells were then spot-plated on

medium containing either IPTG (200 μM, ShyA expressed), arabinose (0.2 %, MepM$_{Ngo}$

expressed) or no inducer. Plates were incubated at 37 ˚C for 24 hours and then imaged.

References

1. Espaillat A, Forsmo O, El Biari K, Bjork R, Lemaitre B, Trygg J, Canada FJ, de
Pedro MA, Cava F. 2016. Chemometric Analysis of Bacterial Peptidoglycan
Reveals Atypical Modifications That Empower the Cell Wall against Predatory
Enzymes and Fly Innate Immunity. J Am Chem Soc 138:9193-204.

2. Vollmer W, Bertsche U. 2008. Murein (peptidoglycan) structure, architecture and biosynthesis in Escherichia coli. Biochim Biophys Acta 1778:1714-34.

3. Vollmer W, Blanot D, de Pedro MA. 2008. Peptidoglycan structure and architecture. FEMS Microbiol Rev 32:149-67.

4. Dik DA, Fisher JF, Mobashery S. 2018. Cell-Wall Recycling of the Gram-Negative Bacteria and the Nexus to Antibiotic Resistance. Chem Rev 118:5952-5984.

5. Park JT, Uehara T. 2008. How bacteria consume their own exoskeletons (turnover and recycling of cell wall peptidoglycan). Microbiol Mol Biol Rev 72:211-27, table of contents.

6. Uehara T, Park JT. 2008. Growth of Escherichia coli: significance of peptidoglycan degradation during elongation and septation. J Bacteriol 190:3914-22.

7. Meeske AJ, Riley EP, Robins WP, Uehara T, Mekalanos JJ, Kahne D, Walker S, Kruse AC, Bernhardt TG, Rudner DZ. 2016. SEDS proteins are a widespread family of bacterial cell wall polymerases. Nature 537:634-638.

8. Paradis-Bleau C, Markovski M, Uehara T, Lupoli TJ, Walker S, Kahne DE, Bernhardt TG. 2010. Lipoprotein cofactors located in the outer membrane activate bacterial cell wall polymerases. Cell 143:1110-20.

9. Typas A, Banzhaf M, van den Berg van Saparoea B, Verheul J, Biboy J, Nichols RJ, Zietek M, Beilharz K, Kannenberg K, von Rechenberg M, Breukink E, den Blaauwen T, Gross CA, Vollmer W. 2010. Regulation of peptidoglycan synthesis by outer-membrane proteins. Cell 143:1097-109.

10. Emami K, Guyet A, Kawai Y, Devi J, Wu LJ, Allenby N, Daniel RA, Errington J. 2017. RodA as the missing glycosyltransferase in Bacillus subtilis and antibiotic discovery for the peptidoglycan polymerase pathway. Nat Microbiol 2:16253.

11. Dorr T, Lam H, Alvarez L, Cava F, Davis BM, Waldor MK. 2014. A novel peptidoglycan binding protein crucial for PBP1A-mediated cell wall biogenesis in Vibrio cholerae. PLoS Genet 10:e1004433.

12. Dorr T, Moll A, Chao MC, Cava F, Lam H, Davis BM, Waldor MK. 2014. Differential requirement for PBP1a and PBP1b in in vivo and in vitro fitness of Vibrio cholerae. Infect Immun 82:2115-24.

13. Dion MF, Kapoor M, Sun Y, Wilson S, Ryan J, Vigouroux A, van Teeffelen S, Oldenbourg R, Garner EC. 2019. Bacillus subtilis cell diameter is determined by the opposing actions of two distinct cell wall synthetic systems. Nat Microbiol 4:1294-1305.

14. Vigouroux A, Cordier B, Aristov A, Alvarez L, Ozbaykal G, Chaze T, Oldewurtel ER, Matondo M, Cava F, Bikard D, van Teeffelen S. 2020. Class-A penicillin binding proteins do not contribute to cell shape but repair cell-wall defects. Elife 9.

15. Vermassen A, Leroy S, Talon R, Provot C, Popowska M, Desvaux M. 2019. Cell Wall Hydrolases in Bacteria: Insight on the Diversity of Cell Wall Amidases, Glycosidases and Peptidases Toward Peptidoglycan. Front Microbiol 10:331.

16. Lee TK, Huang KC. 2013. The role of hydrolases in bacterial cell-wall growth. Curr Opin Microbiol 16:760-6.

17. Do T, Page JE, Walker S. 2020. Uncovering the activities, biological roles, and regulation of bacterial cell wall hydrolases and tailoring enzymes. J Biol Chem 295:3347-3361.

18. Vollmer W. 2012. Bacterial growth does require peptidoglycan hydrolases. Mol Microbiol 86:1031-5.

19. Hashimoto M, Ooiwa S, Sekiguchi J. 2012. Synthetic lethality of the lytE cwlO genotype in Bacillus subtilis is caused by lack of D,L-endopeptidase activity at the lateral cell wall. J Bacteriol 194:796-803.

20. Dorr T, Cava F, Lam H, Davis BM, Waldor MK. 2013. Substrate specificity of an elongation-specific peptidoglycan endopeptidase and its implications for cell wall architecture and growth of Vibrio cholerae. Mol Microbiol 89:949-62.

21. Singh SK, SaiSree L, Amrutha RN, Reddy M. 2012. Three redundant murein endopeptidases catalyse an essential cleavage step in peptidoglycan synthesis of Escherichia coli K12. Mol Microbiol 86:1036-51.

22. Meisner J, Montero Llopis P, Sham LT, Garner E, Bernhardt TG, Rudner DZ. 2013. FtsEX is required for CwlO peptidoglycan hydrolase activity during cell wall elongation in Bacillus subtilis. Mol Microbiol 89:1069-83.

23. Brunet YR, Wang X, Rudner DZ. 2019. SweC and SweD are essential co-factors of the FtsEX-CwlO cell wall hydrolase complex in Bacillus subtilis. PLoS Genet 15:e1008296.

24. Carballido-Lopez R, Formstone A, Li Y, Ehrlich SD, Noirot P, Errington J. 2006. Actin homolog MreBH governs cell morphogenesis by localization of the cell wall hydrolase LytE. Dev Cell 11:399-409.

25. Dominguez-Cuevas P, Porcelli I, Daniel RA, Errington J. 2013. Differentiated roles for MreB-actin isologues and autolytic enzymes in Bacillus subtilis morphogenesis. Mol Microbiol 89:1084-98.

26. Tomasz A. 1979. The mechanism of the irreversible antimicrobial effects of penicillins: how the beta-lactam antibiotics kill and lyse bacteria. Annu Rev Microbiol 33:113-37.

27. Schneider T, Sahl HG. 2010. An oldie but a goodie - cell wall biosynthesis as antibiotic target pathway. Int J Med Microbiol 300:161-9.

28. Lai GC, Cho H, Bernhardt TG. 2017. The mecillinam resistome reveals a role for peptidoglycan endopeptidases in stimulating cell wall synthesis in Escherichia coli. PLoS Genet 13:e1006934.

29. Kitano K, Tuomanen E, Tomasz A. 1986. Transglycosylase and endopeptidase participate in the degradation of murein during autolysis of Escherichia coli. J Bacteriol 167:759-65.

30. Dorr T, Davis BM, Waldor MK. 2015. Endopeptidase-mediated beta lactam tolerance. PLoS Pathog 11:e1004850.

31. Singh SK, Parveen S, SaiSree L, Reddy M. 2015. Regulated proteolysis of a cross-link-specific peptidoglycan hydrolase contributes to bacterial morphogenesis. Proc Natl Acad Sci U S A 112:10956-61.

32. Banzhaf M, Yau HC, Verheul J, Lodge A, Kritikos G, Mateus A, Cordier B, Hov AK, Stein F, Wartel M, Pazos M, Solovyova AS, Breukink E, van Teeffelen S, Savitski MM, den Blaauwen T, Typas A, Vollmer W. 2020. Outer membrane lipoprotein NlpI scaffolds peptidoglycan hydrolases within multi-enzyme complexes in Escherichia coli. EMBO J 39:e102246.

33. Srivastava D, Seo J, Rimal B, Kim SJ, Zhen S, Darwin AJ. 2018. A Proteolytic Complex Targets Multiple Cell Wall Hydrolases in Pseudomonas aeruginosa. mBio 9.

34. Shin JH, Sulpizio AG, Kelley A, Alvarez L, Murphy SG, Fan L, Cava F, Mao Y, Saper MA, Dorr T. 2020. Structural basis of peptidoglycan endopeptidase regulation. Proc Natl Acad Sci U S A 117.

35. Koch AL. 1998. The three-for-one model for gram-negative wall growth: a problem and a possible solution. FEMS Microbiol Lett 162:127-34.

36. Höltje JV. 1993. "Three for one" - a Simple Growth Mechanism that Guarantees a Precise Copy of the Thin, Rod-Shaped Murein Sacculus of *Escherichia coli*. *In* de Pedro MA, Höltje JV, Löffelhardt W (ed), Bacterial Growth and Lysis. Springer, Boston, MA.

37. Chodisetti PK, Reddy M. 2019. Peptidoglycan hydrolase of an unusual cross-link cleavage specificity contributes to bacterial cell wall synthesis. Proc Natl Acad Sci U S A 116:7825-7830.

38. Denome SA, Elf PK, Henderson TA, Nelson DE, Young KD. 1999. Escherichia coli mutants lacking all possible combinations of eight penicillin binding proteins: viability, characteristics, and implications for peptidoglycan synthesis. J Bacteriol 181:3981-93.

39. Kuru E, Hughes HV, Brown PJ, Hall E, Tekkam S, Cava F, de Pedro MA, Brun YV, VanNieuwenhze MS. 2012. In Situ probing of newly synthesized peptidoglycan in live bacteria with fluorescent D-amino acids. Angew Chem Int Ed Engl 51:12519-23.

40. Cava F, de Pedro MA, Lam H, Davis BM, Waldor MK. 2011. Distinct pathways for modification of the bacterial cell wall by non-canonical D-amino acids. EMBO J 30:3442-53.

41. Culp EJ, Waglechner N, Wang W, Fiebig-Comyn AA, Hsu YP, Koteva K, Sychantha D, Coombes BK, Van Nieuwenhze MS, Brun YV, Wright GD. 2020. Evolution-guided discovery of antibiotics that inhibit peptidoglycan remodelling. Nature 578:582-587.

42. Blackman SA, Smith TJ, Foster SJ. 1998. The role of autolysins during vegetative growth of Bacillus subtilis 168. Microbiology 144 (Pt 1):73-82.

43. Takacs CN, Poggio S, Charbon G, Pucheault M, Vollmer W, Jacobs-Wagner C. 2010. MreB drives de novo rod morphogenesis in Caulobacter crescentus via remodeling of the cell wall. J Bacteriol 192:1671-84.

44. Ostash B, Walker S. 2010. Moenomycin family antibiotics: chemical synthesis, biosynthesis, and biological activity. Nat Prod Rep 27:1594-617.

45. Cross T, Ransegnola B, Shin JH, Weaver A, Fauntleroy K, VanNieuwenhze MS, Westblade LF, Dorr T. 2019. Spheroplast-Mediated Carbapenem Tolerance in Gram-Negative Pathogens. Antimicrob Agents Chemother 63.

46. Dominguez-Escobar J, Chastanet A, Crevenna AH, Fromion V, Wedlich-Soldner R, Carballido-Lopez R. 2011. Processive movement of MreB-associated cell wall biosynthetic complexes in bacteria. Science 333:225-8.

47. van Teeffelen S, Wang S, Furchtgott L, Huang KC, Wingreen NS, Shaevitz JW, Gitai Z. 2011. The bacterial actin MreB rotates, and rotation depends on cell-wall assembly. Proc Natl Acad Sci U S A 108:15822-7.

807   48.   Garner EC, Bernard R, Wang W, Zhuang X, Rudner DZ, Mitchison T. 2011.
808   Coupled, circumferential motions of the cell wall synthesis machinery and MreB
809   filaments in B. subtilis. Science 333:222-5.

810   49.   Billaudeau C, Chastanet A, Yao Z, Cornilleau C, Mirouze N, Fromion V, Carballido-
811   Lopez R. 2017. Contrasting mechanisms of growth in two model rod-shaped
812   bacteria. Nat Commun 8:15370.

813   50.   Heidrich C, Templin MF, Ursinus A, Merdanovic M, Berger J, Schwarz H, de Pedro
814   MA, Holtje JV. 2001. Involvement of N-acetylmuramyl-L-alanine amidases in cell
815   separation and antibiotic-induced autolysis of Escherichia coli. Mol Microbiol
816   41:167-78.

817   51.   Moll A, Dorr T, Alvarez L, Chao MC, Davis BM, Cava F, Waldor MK. 2014. Cell
818   separation in Vibrio cholerae is mediated by a single amidase whose action is
819   modulated by two nonredundant activators. J Bacteriol 196:3937-48.

820   52.   Dik DA, Marous DR, Fisher JF, Mobashery S. 2017. Lytic transglycosylases:
821   concinnity in concision of the bacterial cell wall. Crit Rev Biochem Mol Biol 52:503-
822   542.

823   53.   Cai L, Friedman N, Xie XS. 2006. Stochastic protein expression in individual cells
824   at the single molecule level. Nature 440:358-62.

825   54.   Cho H, Wivagg CN, Kapoor M, Barry Z, Rohs PDA, Suh H, Marto JA, Garner EC,
826   Bernhardt TG. 2016. Bacterial cell wall biogenesis is mediated by SEDS and PBP
827   polymerase families functioning semi-autonomously. Nat Microbiol 1:16172.

828   55.   Heidelberg JF, Eisen JA, Nelson WC, Clayton RA, Gwinn ML, Dodson RJ, Haft
829   DH, Hickey EK, Peterson JD, Umayam L, Gill SR, Nelson KE, Read TD, Tettelin
830   H, Richardson D, Ermolaeva MD, Vamathevan J, Bass S, Qin H, Dragoi I, Sellers
831   P, McDonald L, Utterback T, Fleishmann RD, Nierman WC, White O, Salzberg SL,
832   Smith HO, Colwell RR, Mekalanos JJ, Venter JC, Fraser CM. 2000. DNA
833   sequence of both chromosomes of the cholera pathogen Vibrio cholerae. Nature
834   406:477-83.

835   56.   Mekalanos JJ. 1983. Duplication and amplification of toxin genes in Vibrio
836   cholerae. Cell 35:253-63.

837   57.   Dalia AB. 2018. Natural Cotransformation and Multiplex Genome Editing by
838   Natural Transformation (MuGENT) of Vibrio cholerae. Methods Mol Biol 1839:53-
839   64.

840   58.   Guzman LM, Belin D, Carson MJ, Beckwith J. 1995. Tight regulation, modulation,
841   and high-level expression by vectors containing the arabinose PBAD promoter. J
842   Bacteriol 177:4121-30.

843   59.   Chin CS, Sorenson J, Harris JB, Robins WP, Charles RC, Jean-Charles RR,
844   Bullard J, Webster DR, Kasarskis A, Peluso P, Paxinos EE, Yamaichi Y,
845   Calderwood SB, Mekalanos JJ, Schadt EE, Waldor MK. 2011. The origin of the
846   Haitian cholera outbreak strain. N Engl J Med 364:33-42.

847   60.   Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO. 2009.
848   Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat
849   Methods 6:343-5.

850   61.   Donnenberg MS, Kaper JB. 1991. Construction of an eae deletion mutant of
851   enteropathogenic Escherichia coli by using a positive-selection suicide vector.
852   Infect Immun 59:4310-7.

62.   Tinevez JY, Perry N, Schindelin J, Hoopes GM, Reynolds GD, Laplantine E,
Bednarek SY, Shorte SL, Eliceiri KW. 2017. TrackMate: An open and extensible
platform for single-particle tracking. Methods 115:80-90.
63.   Tarantino N, Tinevez JY, Crowell EF, Boisson B, Henriques R, Mhlanga M, Agou
F, Israel A, Laplantine E. 2014. TNF and IL-1 exhibit distinct ubiquitin requirements
for inducing NEMO-IKK supramolecular structures. J Cell Biol 204:231-45.
64.   Bacher CP, Reichenzeller M, Athale C, Herrmann H, Eils R. 2004. 4-D single
particle tracking of synthetic and proteinaceous microspheres reveals preferential
movement of nuclear particles along chromatin - poor tracks. BMC Cell Biol 5:45.
65.   Alvarez L, Hernandez SB, de Pedro MA, Cava F. 2016. Ultra-Sensitive, High-
Resolution Liquid Chromatography Methods for the High-Throughput Quantitative
Analysis of Bacterial Cell Wall Chemistry and Structure. Methods Mol Biol 1440:11-
27.
866