The Internal Structure of Metacommunities

Authors and Affiliations:

Mathew A. Leibold, Department of Biology, University of Florida. 32611

mleibold@ufl.edu

F. Javiera Rudolph, Department of Biology, University of Florida.

javiera.av@ufl.edu

F. Guillaume Blanchet, Département de Biologie, Université de Sherbrooke, 2500 Boulevard

Université, Sherbrooke, Québec, J1K 2R1, Canada.

 guillaume.blanchet@usherbrooke.ca

Luc De Meester, Laboratory of Aquatic Ecology, Evolution and Conservation, University of

Leuven, 3000 Leuven, Belgium; Leibniz Institut für Gewässerökologie und Binnenfischerei

(IGB), 12587 Berlin, Germany; Institute of Biology, Freie Universität Berlin, Königin-Luise-

Strasse 1-3, 14195 Berlin, Germany; Berlin-Brandenburg Institute of Advanced Biodiversity

Research (BBIB), Königin-Luise-Str. 2-4, 14195 Berlin, Germany

Luc.DeMeester@kuleuven.be

Dominique Gravel, Département de Biologie, Université de Sherbrooke, 2500 Boulevard

Université, Sherbrooke, Québec, J1K 2R1, Canada.

Dominique.Gravel@USherbrooke.ca

Florian Hartig, Theoretical Ecology, University of Regensburg, Germany, florian.hartig@ur.de

Pedro Peres-Neto, Department of Biology, Concordia University.

pedro.peres-neto@concordia.ca

Lauren Shoemaker, Botany Department, University of Wyoming.

25   lshoema1@uwyo.edu
26
27   Jonathan M. Chase, German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena
28   Leipzig; Department of Computer Sciences, Martin Luther University Halle-Wittenberg;
29   jonathan.chase@idiv.de
30

Running title: Internal Metacommunity Structure

Keywords: Interspecific competition, site dynamic theory, hierarchical community models,

metacommunity paradigms, variation partitioning.

Data accessibility statement: Should this manuscript be accepted, the data and code supporting

the results will be archived in an appropriate public repository such as Github and will be

mirrored at Zendo or similar locations.

Abstract

Current analyses of metacommunity data largely focus on global attributes across the entire

metacommunity, such as mean alpha, beta, and gamma diversity, as well as the partitioning of

compositional variation across all species and sites. This view neglects that different species and

sites in the landscape can vary widely in how they contribute to these metacommunity-wide

attributes as a function of traits and site attributes. We argue that the study of this internal

structure can help define and describe the complex and interactive relations between process and

pattern in metacommunities. To illustrate this general idea, we create synthetic data using a

simple colonization-extinction metacommunity model, and quantify variation between species

and sites (what we call the 'internal structure' of metacommunities) using Joint Species

Distribution Models. We find that this perspective on internal structure of metacommuniites

provides useful information about the distinct ways that different species and different sites

contribute to metacommunity attributes. We conclude with some discussion about how these

realistic complexities regarding internal structure can be incorporated into a more cohesive

metacommunity theory.

Introduction

Recent developments in community ecology indicate that the field is undergoing an important

renaissance in both its concepts and tools. One of the more exciting and important elements of

this renaissance is in the use of the metacommunity concept, which recognizes the feedback

between local communities and the broader-scale regional biota (Hanski and Gilpin 1991,

Leibold et al. 2004 and reviewed in Leibold and Chase 2017). Earlier work focused on specific

scenarios where such feedbacks could occur (e.g., Levins and Culver 1971, Horn and MacArthur

1974, Levin 1974, Sloan-Wilson 1992, Leibold 1998, Hubble 2001, Amarasekare and Nisbet

2001), which was synthesized into a useful categorization of the different ways in which

metacommunities could be structured (Leibold et al. 2004). However, it is apparent that there is

a much more complex and nuanced spectrum of possibilities regarding the mechanisms and

processes underlying the structure of metacommunities (Leibold and Chase 2017). Ongoing

developments, including both more sophisticated theoretical (e.g., Shoemaker and Melbourne

2016, Fournier et al. 2017, Ovaskainen et al. 2019, Thompson et al. 2020) and analytical (e.g.,

Legendre and DeCaceres 2013, Hui et al. 2013, Ovaskainen et al. 2017, Ohlman et al. 2018,

Jabot et al. 2020) approaches, promise a new level of understanding and synthesis of the

regional-local community-level feedbacks. It seems increasingly apparent that understanding the

feedbacks between local communities and the regional biota is a major challenge for community

ecology in general (see Leibold and Chase 2017) as well as for the application of these insights

to applied issues of environmental and health concerns (e.g., Bengtsson 2009, Schiesari et al.

2019, Miller et al. 2019, Brown and Barney 2020).

While important progress has been made in the study of metacommunities, most work tends to

assume that processes act similarly on all species and sites in a metacommunity. However,

variation among species and among sites can strongly influence metacommunity-level properties

of biotas. This is particularly relevant to local-regional feedbacks because species and sites vary

across metacommunities. For instance, the spatial structure of environmental features among

sites almost always changes different explicitly determined landscapes due to spatial contingency

(Peres-Neto et al. 2012). In contrast with many previous studies (e.g., Blanchard et al. 2020,

Jabot 2020), we emphasize that different species can be heterogenous in how they contribute to

metacommunity level properties and that different sites can also be heterogenous in how they

contribute to these patterns (but see Pandit et al. 2009, Legendre and De Cáceres 2013). Using

this concept as a starting point, we highlight the importance of understanding the 'internal

structure' of metacommunities, which can help resolve more complex dynamics that could not be

resolved without taking this different perspective. Community assembly in metacommunities is a

complex process that involves mechanisms related to environmental effects, species interactions,

dispersal, and stochasticity as they act on a regionally-defined pool of species within a specific

landscape (e.g. Vellend 2010, 2015, Weiher et al. 2011). Current analytical tools for inferring

the effects of these mechanisms from metacommunity patterns focus on global metrics that

describe the entire metacommunity, such as diversity metrics and coexistence patterns and

variation partitioning analysis (e.g., Borcard et al. 1992, Gotelli and McCabe 2002, Leibold and

Mikkelson 2002). Recent efforts have used several of these global metrics to separate

metacommunity processes (Ovaskainen et al. 2019, Guzman et al. 2020). While these

approaches provide insights into the processes that drive species distributions and determine their

levels of interaction within metacommunities, they only characterize them in very broad terms at

the larger metacommunity scale, which we consider to be the 'external' structure of

metacommunities. Here, we focus on the 'internal structure' instead and thus on resolving how

individual species and sites or patches contribute to such broad patterns by dissecting their

individual contribution to the global metacommunity pattern (see also Fournier et al. 2017,

Suzuki and Economo 2021).

To illustrate the advantages of studying the internal structure of metacommunities, we create

synthetic data from a process-based metacommunity model of competition-extinction dynamics.

We evaluate species-level and site-level variation using joint species distribution models

(JSDMs; see review in Warton et al. 2016). We find that there can be tremendous heterogeneity

between the contributions of individual species and sites to overall metacommunity structure;

and that some of this heterogeneity can be identified by JSDMs and be related to attributes such

as dispersal and species associations for species, or environmental uniqueness for sites.

Previous work has shown that particular examples of such effects can occur but here we seek a

more general approach. For example, Pandit et al. (2009) showed that species can be

heterogenous in their responses to environmental and stochastic factors depending on their

degree of habitat specificity. Others have argued that JSDMs can provide important insights into

the drivers of such variation in species distributions ((Hui et al. 2013, 2016, Pollock et al. 2014,

Ovaskainen et al. 2017 see also Ovaskainen and Abrego 2020; but see Poggiato et al. 2021).

Similarly, the heterogeneity among sites has long been identified as driving individual species

distributions (see Guisan and Thulliers 2005, Soberon and Peterson 2005, Elith and Leathwick

2006) as well as driving overall variation among sites in global metrics of community structure

(e.g. diversity patterns, etc.) as characterized by the field of landscape ecology (Turner et al.

2001). Further suggestions that individual sites might vary in how they contribute to

metacommunity dynamics include the concept of 'keystone communities' (Mouquet et al. 2012,

Resetarits et al. 2017, Yang et al 2020) and metacommunity approaches to spatial networks.

These disparate approaches (species vs sites) to metacommunities are likely closely related to

each other and can be linked by the emerging methodologies of methods such as JSDMs to a

more nuanced metacommunity ecology that recognizes a plurality of mechanisms and processes

underlying community assembly. Nevertheless, we discuss some remaining important

challenges to resolve in making process-pattern linkages in metacommunities (see also Poggiato

et al. 2021, Miele et al. 2021).

134
135
Quantifying the link between process and pattern using a simple metacommunity

simulation and refined statistical approach

To test and exemplify our ability to infer individual species and site contributions, we

simulate data from a process-based model, which allows us to create observations with full

knowledge about the underlying mechanisms. Our process-based model is based on a spatial

implementation of spatially implicit site occupancy models (e.g., Levins and Culver 1971, Horn

and MacArthur 1972, Levin 1974, Hastings 1980, Hanski 1991) to describe dynamics in

heterogeneous metacommunities. We focused here on a model for predicting presence-absence

(and not abundance; but see Supporting Information) because it is the most widely available type

of empirical data for metacommunity analysis. For each species in each patch, we model

occupancy using two key equations (details in the Supporting Information). The first of these

describes the colonization of patch z by species i during a discrete time interval, Δt:

$$P(X_{i,z,t+\Delta t} = 1|X_{i,z,t} = 0) = I_{i,z,t}S_{i,z,t}C_{i,z,t} \tag{1}$$

where $X_{i,z,t}$ is a stochastic variable representing the occurrence of species i at location z at time

t, $I_{i,z,t}$ is the number of immigrants of species, $S_{i,z,t}$ is the effect of environmental filtering on the

probability of establishing a viable local population, and $C_{i,z,t}$ is the effect of ecological

interactions on the establishment probability. Second, we consider the alternative possibility: the

extinction of species i in patch z during the time interval Δt:

$$P(X_{i,z,t+\Delta t} = 1|X_{i,z,t} = 0) = M_{i,z,t}E_{i,z,t} \tag{2}$$

where $M_{i,z,t}$ and $E_{i,z,t}$ are the responses of the extinction probability to the local environment and

to ecological interactions, respectively.

At steady state the solution to this model is:

$$\log \frac{\hat{p}^{i,z}}{1-\hat{p}^{i,z}} = \log I^{i,z} + \log \frac{S^{i,z}}{M^{i,z}} + \log \frac{C^{i,z}}{E^{i,z}}$$ (3)

where $\hat{P}^{iz}$ is the expected probability that site z is occupied by species i. This formulation

assumes that immigration($I^{i,z}$), 'environmental selection' ($S^{i,z}$ and $M^{i,z}$) and interactions

($C^{i,z}$ and $E^{i,z}$) can be separated into distinct effects that do not interact.

Equation 3 can be analyzed by a JSDM to separate the contributions of these effects into spatial

effects (driven by immigration), environmental filtering (driven by abiotic selection) and species

co-distribution unrelated to either space or environment, with an additional fraction quantifying

residuals resulting from stochasticity in the case of a finite number of patches (see also

Shoemaker et al. 2020). Furthermore, the likelihood of every observation can be marginalized

over each species (by summing the likelihoods for a given species across all patches) to describe

the variation among species. Alternatively, the likelihood can be marginalized by sites (by

summing the likelihoods for a patch across all species) to describe the variation across the

metacommunity landscape. Doing so, we can quantify the importance of environment, species

co-distribution, and space for predicting metacommunity structure as a whole, as well as their

importance for predicting the presence-absence (or, in theory, the abundance) of individual

species or community composition at individual patches.

Analyzing the inferred quantities, we can study the internal structureof the metacommunity. This

can clarify how these key processes work together in a diverse and complex metacommunity to

explain species distributions within the metacommunity (Figure 1). We also emphasize that one

main advantage of the proposed framework is to allow ecologists to identify the contributions of

individual communities and species which are often units of personal interest and knowledge

(e.g., natural history of species, place-based ecological knowledge).

To illustrate the utility of this approach in a more realistic framework that includes

stochasticity and spatially explicit landscapes, we implemented the key processes of drift,

environmental filtering, dispersal, and species interactions (Vellend 2010, 2016) in a flexible

simulation version of the model (described in more details in the Supporting Information). The

simulation model allows us to vary each process separately for each species in a heterogeneous

spatially explicit landscape. It simulates the dynamics of a metacommunity across a set of

patches and generates a spatial network that specifies the connectivity among patches. The state

variables of the simulation are the occupancy of each species in every patch (i.e.

presence/absence, though future implementations could also address abundance data, e.g.

Rybicki et al. 2018, Ovaskainen et al. 2019, Thompson et al. 2020). Each patch can be colonized

from nearby patches depending on their location in the landscape, dispersal rate of the species

and proximity of extant populations in neighboring patches. Each species in each patch is subject

to extinctions that reflect demographic and/or environmental stochasticity. Patches can differ in

local environmental conditions that differentially influence baseline colonization and extinction

probabilities. Species interactions are modeled in two ways. First, the presence of other species

in a patch can modify baseline colonization probability (a reduction in the case of competition).

Second, co-occurring species can modify baseline extinction probability (an increase in the case

of competition).

We next apply a JSDM to the resulting distribution of species among patches.

Specifically, we use the HMSC R package (Ovaskainen et al. 2017, Ovaskainen and Nerea 2020)

that models species distributions as a function the environment, spatial autocorrelation and

species co-distributions. After fitting the model using HMSC, we partition variation between the

model processes into four components (or fractions) using an approach akin to classic variation

partitioning (Borcard et al. 1992; Peres-Neto et al. 2006). Details about how this type of variation

partitioning is computed through HMSC is given in the Supporting Information. Specifically, the

variation was partitioned to quantify the effects of environment (labeled [E]), spatial patterning

(labeled [S]), co-distribution among species (labeled [C]) and residual (unexplained) variation

that cannot be attributed to any of the three previously mentioned fractions (i.e. sets of

predictors). This is expressed as $1-R^2$, where R^2 is the proportion of variation explained by the

model and includes fractions [E], [S], and [C].

Compared to the classical method of variation partitioning (Borcard et al. 1992, Peres-

Neto et al. 2006), this approach provides two major advances. The first is that we can infer a

third fraction that quantifies co-variation (or co-distribution) among species using latent

variables (fraction [C]), which is distinct from either the environment effect (fraction [E]) or the

spatial effects (fraction [S]). In our model this fraction reflects the consequences of species

associations rather than biotic interactions, and we emphasize that this interpretation must be

done with caution (see Dormann et al. 2018 and Blanchet et al. 2020 for critical reviews of this

issue). In nature, this latent fraction may also reflect other sources of variation such as those

from unmeasured environmental variables. Similar points have been raised in metacommunity

analysis regarding the interpretation of model variation due to spatial autocorrelation as these

could be related to spatialized environmental effects due to unmeasured environmental factors

(Peres-Neto et al. 2012). The second advance is that all fractions are calculated at the species

level, where species-specific variation detail the internal structure of the metacommunity from

the perspective of each individual species. Fractions of the variation at the metacommunity level

are then calculated by averaging across all species to obtain metacommunity (global) level

descriptors as those obtained from variation partitioning. This allows us to subsequently

examine potential drivers (environment, space and joint co-distribution) underlying the variation

in these different effects among species. In addition, we implemented a partitioning scheme to

quantify how each site separately contributes to different drivers as described in the Supplement

Information. This way to partition variation among sites informs us about the individual

contribution each site has to variation in the data for each set of drivers. The ability to resolve

species-specific and site-specific components means that we can study the 'internal structure' of

the metacommunity in ways that were absent in classic variation partitioning.

Although the analytical model described in equations 1-3 suggests that making links

between processes and patterns using JSDMs are possible, we wished to evaluate if this was also

likely in less idealized situations such as those used in our metacommunity model. To do this,

we simulated a number of scenarios that vary the strength of environmental selection, dispersal

and competition. Comparing scenarios with varying niche breadth and competition (scenarios

A-D, Fig. 2 and Fig 3), and a more complex case where species compete and vary in both

dispersal and in their responses to the environment (scenario G, Figure 4, scenarios E and F in

the SI) highlight how HMSC provides an avenue for distinguishing between underlying

processes based on abundances of species across metacommunity patches. Using our

framework, our goal here is to illustrate how links between pattern and process might be made in

metacommunities. In doing so, we leave a more extensive and systematic evaluation of the

model's components (e.g., performance of JSDMs under multiple complex scenarios) for future

work (but see Ovaskainen and Nerea 2020).

Simulation experiments

In a first set of simulations, we considered a situation where species had distinct

environmental optima along an environmental gradient and had limited dispersal (Figure 2,

Table 1). We contrasted the case where the environmental niches were narrow (steep changes in

baseline colonization success and extinction rates with small deviations in environment) with the

case with identical optima, but with wide environmental niches (much weaker changes in

colonization and extinction with environmental value). As expected, we find that these

differences in environmental niche breadth had strong effects on the relative importance of

environmental filtering (fraction [E]) versus spatial patterning (fraction [S]). Specifically, we

find stronger spatial patterning when niche breadths were broad and stronger environmental

filtering when niches were narrower (Figure 2, Table 1). We also found that the R^2 values were

higher for the case with narrow niches than with wide niches. Finally, we found non-zero

(though relatively weak) variance components for co-distributions (fraction [C]) in both cases,

especially when niches were broad even though our analytical model would predict the absence

of such variation components since colonization and extinctions were not affected by species

interactions in these simulations.

We next simulated metacommunities with identical parameters as above, except with added

interspecific competition effects (Figure 3, Table 1). As in the case without species interactions

(compare with Figure 2), narrow niches enhanced the relative strength of environmental filtering

(fraction [E]) and reduced spatial patterning (fraction [S]) when compared to wide niches. In

these simulations, however, the co-distribution components (fraction [C]) were much more

substantial than without species interactions. We also found that adding interspecific

competition substantially increased the total amount of variation explained (R^2) by the model

(i.e. due to the joint component of co-distribution).

We conducted a number of other simulations to explore if that describe interspecific

variation on environment (fraction [E]), space (fraction [S]), and co-distributions (fraction [C])

depend on the dispersal, niches breadth, and interactions. Illustrative examples are shown in the

supplemental information and summary statistics are shown in Table 1. In Figure 4, we present

the results from one of these examples that includes heterogenous dispersal to show how the

internal structure can reveal how dispersal variation affects species distributions. We found that

one could distinguish species by the degree to which their distributions are related to

environment (fraction [E]), space (fraction [S]) and co-distributions (fraction [C]) (Figure 5a)

and we found that this could be related to their traits (i.e., species optima in our simulation

framework). Species with higher dispersal ability and more specialized environmental niche

positions had distributions better predicted by the environment than those that were dispersal

limited and had distributions that presented a higher level of spatial autocorrelation (fraction [S]).

Species with optima closer to the middle of the environmental gradient also had a larger fraction

[C] than those with more extreme optima.

Sites also differed in how their species composition was related to environmental

(fraction [E]) and spatial effects (fraction [S]) as well as co-distributions (fraction [C] - Figure

4b). Some sites tended to be occupied by locally dominant species (in the lower left of the

ternary plot, nearer to fraction [E]), while others were occupied by species found in nearby sites

(lower right of the ternary plot, nearer to fraction [S]). Some sites were also occupied by

combinations of species that were differentially associated with each other regardless of

environment or dispersal (upper apex of the ternary plot, nearer to fraction [C]). As can be seen

in Figure 4b, there were also a wide range of intermediate conditions. A major driver of this

variation is the local environmental condition, especially in relation to how distinct the local

environment is from the overall mean environment of the metacommunity (Figure 5b).

We further investigated the structure of the species co-distribution (fraction [C]). This

covariation can be directly attributable to species interactions because we explicitly model the

processes underlying metacommunity dynamics. However, even in our model, species co-

distribution may not directly link to pairwise interaction coefficients, but rather may emerge as a

complex relationship between species interactions and environmental conditions (Cazelles et al.

2015, Blanchet et al. 2020). To illustrate this, we show the co-distribution among species as a

heat map separately for each of the five individual simulations presented in Figure 4 and

compared them to the actual interaction matrix that describes interspecific competition in our

model (Figure 6; similar heat maps obtained with the other scenarios are shown in the

Supplement Information). Despite the fact that the same interaction matrix was used for all five

of these simulations, the resulting co-distribution patterns are inconsistent in their details.

However, these matrices show that there is consistency in several features of the co-distribution

pattern. For example, they all share the predominance of strong negative correlations along the

main diagonal that match the interaction matrix we used. They also share a strong

'checkerboard' pattern with alternating negative and positive co-distributions between species

when these are ranked against their environmental optima. Given the simple scheme of species

interactions we used (Fig. 5 and SI), these results are consistent with the predictions that direct

interactions are stronger than indirect ones and tend to weaken with the number of links in

indirect chains even if the details of these effects are less predictable.

Discussion

While metacommunity ecology has made great progress in the past decades, the assumptions that species and sites were relatively homogeneous in their underlying processes (reviewed in Leibold and Chase 2017) have limited the applicability of metacommunity theory to more realistic species pools within complex landscapes. Here, by combining a tractable process-based model with emerging analytical methods, we provide a general quantitative approach that accounts for multiple interacting processes that may operate differently among species or in different parts of landscapes.

Although there are some important challenges to consider, our study illustrates important insights about the internal structure of metacommunities, including:

1) Variation partitioning using JSDMs (here implemented using HMSC) can be linked to basic processes of community assembly at the species and sites levels (e.g. environmental selection, dispersal, biotic selection, and drift).

2) Quantifying co-distributions of species in metacommunities can improve predictive ability even when the processes that generate these distributions are complex (stochasticity, complex spatial landscapes, and species interactions).

3) Species can have distributions that vary in the degree to which they are determined by combinations of the basic community assembly processes depending on features of their ecology (e.g. dispersal and environmental preferences); and

4) The predominant assembly processes that determine local communities can differ among adjacent sites in a metacommunity (e.g. sites that are occupied by species most fit for environmental conditions vs sites occupied by species in nearby sites that may differ in environmental conditions).

It is important to emphasize that there remain some substantial challenges in moving

forward with the overall approach we advocate in this paper. This includes some technical

issues, such as the estimation of parameters and interpretation of results in more complex

metacommunity models in JSDMs, as well as conceptual ones, such as accounting for other

processes such as speciation, local adaptation, historical biogeography. Nevertheless, we see that

such endeavor will be fruitful, allowing a deeper understanding of ecological dynamics in more

realistic, but necessarily complex, spatial landscapes.

Our analytical framework simplifies several potentially complex processes (e.g. non-

linearities and interactive effects of mechanisms) into an approximation involving colonization-

extinction dynamics. It is possible that more realistic and complex mechanisms driving these

processes will weaken associations between pattern and process or create biases in the

partitioning of the variation revealed by JSDMs. However, the developments of JSDMs are still

progressing, and we anticipate that future developments will solve some of these problems (see

Wilkinson et al. 2020).

The co-distribution component of the JSDMs (fraction [C]) is particularly concerning.

We find that this component can be biased, especially when species have broad environmental

niches (Figure 2). It is widely recognized that there are particular challenges in quantifying and

interpreting species co-distributions (reviewed by Blanchet et al. 2020). Perhaps the most

obvious challenge is that this component can reflect any process that is inadequately quantified

by the environmental (fraction [E]) and spatial components (fraction [S]). In addition to species

interactions, this would include, for example, unmeasured environmental factors (see Blanchet et

al. 2020) or inadequately quantified landscape attributes. Teasing apart the effects of species

interactions from these confounding factors should thus be a major focus for future work.

Nevertheless, it is important to understand that including the co-distribution component in our

analyses allows us to account for them, rather than lumping them with residual variation where

they have likely given a greatly exaggerated impression of stochasticity.

The basic framework we used to address the process to pattern links is most transparent

for mechanisms that focus on interspecific competition because of the analogies with

evolutionary genetics (Vellend 2010, 2016). However, species interactions in metacommunities

are much more variable and include consumer-resource interactions, mutualisms, and facilitative

interactions. Although such interactions can easily be incorporated in simulations, the

interpretation that might link process to pattern in such cases are likely to become more complex.

Likewise, future work could include local (co-)evolutionary dynamics (see Urban et al. 2020)

and historical effects of biogeography and speciation (e.g. Leibold and Chase 2018, Overcast et

al. 2020). Here, we have also retained a simple two-level perspective on spatial scale (local

discrete sites in a broader regional landscape). It is increasingly apparent that metacommunity

dynamics occur over multiple nested scales and that habitats can be continuous and/or nested,

rather than discretely patchy (e.g. Munkenmuller et al. 2012, Rybicki et al 2018, Ovaskainen et

al. 2019, Viana and Chase 2019, König et al. 2021)Refining our approach to address multiple

spatial scales is a logical next step.

Finally, it is increasingly clear that temporal dynamics of community change in

metacommunities can provide critical insights about the mechanisms that drive metacommunity

patterns (e.g. Jabot et al. 2020, Blanchard et al. 2020, Guzman et al. 2021). We imagine future

work on the internal structure of metacommunities as being very amenable to incorporating

temporal changes (see for example, Ovaiskainen et al. 2017 for an initial step in this direction).

Here we have focused on purely spatial approaches because there are still remarkably few studies

that might permit sufficiently structured data to permit temporal analyses and because the

limitations and challenges of such analyses are not yet clear.

We have argued that a focus on the internal structure of metacommunities, by examining

site-specific and species-specific variation components, can enhance our study of the links

between process and pattern in the distributions of species and the occupancies of sites. We also

argue that continued work on this focus is essential if metacommunity ecology is to address the

dynamics and structure of realistic metacommunities that have typically high biodiversity and

occur in complex landscapes. This focus on internal structure represents a shift from traditional

approaches that used descriptors of overall variation components at the metacommunity scale

that generalizes previous ad hoc approaches to similar internal variation in metacommunity

patterns (e.g. Pandit et al. 2009, Legendre and De Cáceres 2013). The dynamics and structure of

distributions of realistically diverse species in a realistically structured landscape of sites likely

involves the interaction of community assembly processes including environmental filtering,

dispersal, and drift, and these are unlikely to be adequately described by simple metacommunity

level metrics (see Ovaskainen et al. 2019). Consequently, dissecting the internal structure of

metacommunities on the basis of species and site contributions could provide key insights into

the processes underlying metacommunity assembly. Such insights might be particularly useful

in the management of landscapes and metacommunities for conservation purposes since they

focus on particular units (species or sties) that are often the focus of concern in such cases.

Acknowledgments: This paper emerged from workshops funded with the support (to JMC)

from the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig funded

by the German Research Foundation (FZT 118). MAL would like to acknowledge the

Alexander von Humboldt Foundation for a Research Award that helped fund this work. LDM

acknowledges KU Leuven Research Fund project C/2017/002 and FWO projects G0B9818 and

G0C3818. We also thanks F. Engel, C. Rakowski, K. Taylor, R. Pelinson, X. Zhao, L. Juen, T.

Michelan and A. Rudolf for comments on earlier versions of the manuscript.

References

Amarasekare, P. , & Nisbet R. M. (2001). Spatial heterogeneity, source-sink dynamics, and the

local coexistence of competing species. *The American Naturalist,* 158, 572-584.

Bengtsson J. (2009). Applied (meta)community ecology: diversity and ecosystem services at the

intersection of local and regional processes. In: Verhoef HA and Morin PJ (Eds),

Community Ecology: Processes, Models, and Applications. Oxford, UK: Oxford

University Press.

Blanchard, G., Birnbaum, P., & Munoz, F. (2020). Extinction–immigration dynamics lag behind

environmental filtering in shaping the composition of tropical dry forests within a

changing landscape. *Ecography,* 43, 869-881.

Blanchet, F.G., Cazelles, K., & Gravel, D. (2020). Co-occurrence is not evidence of ecological

interactions. *Ecology Letters*, 23, 1050–1063.

Borcard, D., Legendre, P., & Drapeau, P. (1992). Partialling out the spatial component of

ecological variation. *Ecology*, 73, 1045–1055.

Boulangeat, I., Gravel, D., & Thuiller, W. (2012). Accounting for dispersal and biotic

interactions to disentangle the drivers of species distributions and their abundances. *Ecology*

Letters, 15, 584-593.

Brown, B.L., & Barney, J.N. (2021). Rethinking Biological Invasions as a Metacommunity

Problem. *Frontiers in Ecology and Evolution,* 8, 584701.

Brown, B.L., Sokol, E.R., Skelton, J., & Tornwall, B. (2017). Making sense of

metacommunities: dispelling the mythology of a metacommunity typology. *Oecologia*, 183,

643–652.

Cottenie, K. (2005). Integrating environmental and spatial processes in ecological community

dynamics: Meta-analysis of metacommunities. *Ecology Letters*, 8, 1175–1182.

Dormann, C.F., Bobrowski, M., Dehling, M. et al. (2018). Biotic interactions in species

distribution modelling: ten questions to guide interpretation and avoid false conclusions.

Global Ecology and Biogeography, 27,1004-1016.

Elith, J., & Leathwick, J.R. (2009). Species Distribution Models: Ecological Explanation and

Prediction Across Space and Time. *Annual Review of Ecology, Evolution, and Systematics*,

40, 677–697.

Fournier, B., Mouquet, N., Leibold, M.A., & Gravel, D. (2017). An integrative framework of

coexistence mechanisms in competitive metacommunities. *Ecography*, 40, 630–641.

Gotelli, N.J., & McCabe, D.J. (2002). A meta-analysis of J. M. Diamond's assembly rules

model. *Ecology*, 83, 2091-2096.

Gravel, D., Canham, C.D., Beaudet, M., & Messier, C. (2006). Reconciling niche and

neutrality: the continuum hypothesis. *Ecology Letters*, 9, 399–409.

Gravel, D., & Massol, F. (2020). Toward a general theory of metacommunity ecology. *In*:

McCann, K.S. & Gellner, G. (eds.). Theoretical Ecology. Oxford University Press. *In press.*

Guisan, A., & Thuiller, W. (2005). Predicting species distribution: offering more than simple

habitat models. *Ecology Letters*, 8, 993-1009.

Guzman, L.M., Germain, R.M., Forbes, C., Straus, S., O'Connor, M. I., Gravel, D. Srivastava,

D.S., & Thompson, P.L. (2019). Towards a multi-trophic extension of metacommunity

ecology. *Ecology Letters,* 22, 19-33.

Guzman, L.M., Thompson, P.L., Viana, D.S., Vanshoenwinkel, B., Horvath, Z., Ptacnik, R.,

Jeliazkov, A., Gascon, S., Lemmens, P., Anton-Pardo, M., Langenheder, S., De Meester, L.,

& Chase, J.M. (2021). Disentangling metacommunity processes using multiple metrics in

space and time. https://www.biorxiv.org/content/10.1101/2020.10.29.361303v1

Hanski, I. (1991). Single-species metapopulation dynamics: concepts, models and observations.

Biological Journal of the Linnean Society of London. 42, 17-38.

Hanski, I., & M. Gilpin. (1991). Metapopulation dynamics: brief history and conceptual

domain. *Biological Journal of the Linnean Society of London. 42, 3-16.*

Hastings, A. (1980). Disturbance, coexistence, history, and competition for space. *Theoretical*

Population Biology, 18, 363–373.

Horn, H.S., & MacArthur, R.H. (1972). Competition among fugitive species in a harlequin

environment. *Ecology*, 53, 749–752.

Hubbel, S. P. (2001). The Unified Neutral theory of Biodiversity and Biogeography. Princeton

University Press. Princeton, N.J.

Hui, F.K.C., Warton, D.I., Forster, S.D., & Dunston, P.K. (2013). To mix or not to mix:

comparing the predictive performance of mixture models vs. separate species distribution

models. *Ecology,* 94, 1913-1919.

Hui, F.K.C. (2016). boral – Bayesian Ordination and Regression Analysis of Multivariate

Abundance Data in r. *Methods in Ecology and Evolution*, 7, 744–750.

Jabot, F., Laroche, F., Massol, F., Arthaud, F., Crabot, J., Dubart, M., Blanchet, S., Munoz, F.,

David, P., & Datry, T. (2020). Assessing metacommunity processes through signatures in

spatiotemporal turnover of community composition. *Ecology Letters*, 23, 1330-1339.

König, C., Wuest, R.O., Graham, C.H., Karger, D.N., Sattler, T., Zimmerman, N.E. & Zurell, D.

(2021). Scale dependence of joint species distribution models challenges interpretation of

biotic interactions. *Journal of Biogeography, in press.*

Lawton, J.H. (1999). Are there general laws in ecology? *Oikos*, 84, 177.

Legendre, P., & De Cáceres, M. (2013). Beta diversity as the variance of community data:

dissimilarity coefficients and partitioning. *Ecology Letters*, 16, 951-963.

Leibold, M.A. (1998). Similarity and coexistence in regional biotas. *Evolutionary Ecology,* 12,

95-110.

Leibold, M.A., & Chase J.M. (2018). *Metacommunity ecology*. Princeton University Press,

Princeton.

Leibold, M.A., Economo, E.P., & Peres-Neto, P. (2010). Metacommunity phylogenetics:

separating the roles of environmental filters and historical biogeography. *Ecology Letters*, 13,

1290–1299.

Leibold, M.A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J.M., Hoopes, M.F., *et al.*

(2004). The metacommunity concept: a framework for multi-scale community ecology.

Ecology Letters, 7, 601–613.

Leibold, M.A., & McPeek, M.A., (2006). Coexistence of the niche and neutral perspectives in

community ecology. *Ecology*, 87, 1399-1410.

Leibold, M.A., & Mikkelson, G.M. (2002). Coherence, species turnover, and boundary

clumping: elements of meta-community structure. *Oikos*, 97, 237–250.

Levin, S.A. (1974). Dispersion and population interactions. *American Naturalist*, 108, 207-228.

Levins, R., & Culver, D. (1971). Regional coexistence of species and competition between rare

species. *Proceedings of the National Academy of Sciences, U.S.A.*, 68, 1246-1248.

Lindstrom, E.S., & Langenheder, S., (2012). Local and regional factors influencing bacterial

community assembly. *Environmental Microbiology Reports*, 4, 1-9.

Logue, J.B., Mouquet, N., Peter, H., & Hillebrand, H. (2011). Empirical approaches to

metacommunities: a review and comparison with theory. *Trends in Ecology & Evolution*, 26,

482–491.

Miller, E.T., & Bohannan, B. (2019). Life between patches: Incorporating microbiome biology

alters the predictions of metacommunity models. *Frontiers in Ecology and Evolution,* 7, 276.

Mouquet, N., Gravel, D., Massol, F., & Calcagno V. (2012). Extending the conept of keystone

species to communities and ecosystems. *Ecology Letters*, 16, 1-8.

Münkemüller, T., de Bello, F., Meynard, C. N., Gravel, D., Lavergne, S., Mouillot, D.,

Mouquet, N., & Thuiller, W. (2012). From diversity indices to community assembly

processes: a test with simulated data. *Ecography*, 35, 468–480.

Ohlman, M., Mazel, F., Chalmandrier L., Bec, S., Coissac, E., Gielly, L., Pansu, J., Schilling,

V., Taberlet, P., Zinger, L, Chave, J., & Thuiller, W. (2018). Mapping the imprint of biotic

interactions on b-diversity. *Ecology Letters,* 21, 1660-1660.

Ovaskainen, O., & Abrego, N., (2020). Joint Species Distribution Modelling

With Applications in R. Cambridge University Press.

Ovaskainen, O., Tikhonov, G., Norberg, A., Guillaume Blanchet, F., Duan, L., Dunson, D., *et*

al. (2017). How to make more out of community data? A conceptual framework and its

implementation as models and software. *Ecology Letters*, 20, 561–576.

Ovaskainen, O., Rybicki, J., & Abrego, N. (2019). What can observational data reveal about

metacommunity processes? *Ecography*, 42, 177-1886.

Overcast, I., Ruffley, M., Rosindell, J., Harmon, L., Borges, P.A.V., Emerson, B.C., Etienne,

R.S., Gillespie, R., Krehenwinke, H., Mahler, D.L., Massol, F., Parent, C.E., Patino, J., Peter,

B., Week, B., Wagner, C., Hickerson, M.J., & Rominger A. (2020). A unified model of

species abundance, genetic diversity, and functional diversity reveals the mechanisms

structuring ecological communities. bioRxiv 10.1101/2020.01.30.927236.

Pandit, S.N., Kolasa, J., & Cottenie, K. (2009). Contrasts between habitat generalists and

specialists: an empirical extension to the basic metacommunity framework. *Ecology,* 90,

2253–2262.

Poggiato, G., Munkenuller, T., Bruystova, D., Arbel, J., Clark, J.S., & Thuiller, W. (2021). On

the interpretation of Joint Modelling in community ecology. *Trends in Ecology and*

Evolution. (in press).

Peres-Neto, P.R., Legendre, P., Dray, S., & Borcard, D. (2006). Variation partitioning of species

data matrices: Estimation and comparison of fractions. *Ecology*, 87, 2614–2625.

Peres-Neto, P.R., Leibold, M.A., & Dray, S. (2012). Assessing the effects of spatial contingency

and environmental filtering on metacommunity phylogenetics. *Ecology*, 93, S14–S30.

Pollock, L.J., Tingley, R., Morris, W.K., Golding, N., O'Hara, R.B., Parris, K.M., et al. (2014).

Understanding co-occurrence by modelling species simultaneously with a Joint Species

Distribution Model (JSDM). *Methods in Ecology and Evolution*, 5, 397–406.

Resetarits, E.J., Cathey, S.E., & Leibold M.A. (2017). Testing the keystone community concept:

effects of landscape, patch removal, and environment on metacommunity structure. *Ecology,*

99, 57-67.

Ricklefs, R.E. (1987). Community diversity: Relative roles of local and regional processes.

Science, 235, 167–171.

Rybicki, J., Abrego, N., & Ovaskainen, O., (2019). Habitat fragmentation and species diversity

in competitive communities. *Ecology Letters*, 23, 506-517.

Schiesari, L., Matias, M.G., Prado, P.I., Leibold M.A., Albert, C.H., Howeth, J.G., Leroux S.H.,

Pardini, R., Siqueira, T., Brancalion, P.H.S., Cabeza, M., Coutinho, R.M. Diniz-Filho,

J.A.D.F., Fournier, B. Lahr, D.J.G., Lewinsohn, T.M., Martins, A., Morsello, C., Peres-Neto,

P.R., Pillar, V.D., & Vazques, D.P. (2019). *Perspectives in Ecology and Conservation*, 17,

172-181.

Shoemaker, L.G., & Melbourne B.A. (2016). Linking metacommunity paradigms to spatial

coexistence mechanisms. *Ecology,* 97, 2436–2446.

Shoemaker, L.G., Sullivan, L.L., Donohue, I., Cabral, J.S., Williams, R.J. Mayfield M.M.,

Chase, J.M., Hcu, C. Harpole, W.S., Huth, A., HilleRisLambers, J., James, A.R.M., Kraft,

N.J.B., May, F., Muthukrishnan, R., Satterlee, S., Taubert, F., Wang, X., Wiegand, T.,

Yang,Q., & Abbott, K.S. (2020). Integrating the underlying structure of stochasticity into

community ecology. *Ecology*, 101, e02922.

Sloan-Wilson, D. (1992). Complex Interactions in Metacommunities, with Implications for

Biodiversity and Higher Levels of Selection. *Ecology,* 73. 1984-2000.

Soberon, J., & Peterson, A.T. (2005). Interpretation of models of fundamental ecological niches

and species distributional areas. *Biodiversity Informatics*, 2, 1-10.

Suzuki, Y., & Economo, E.P. (2021). From species sorting to mass effects: spatial network

structure mediates the shift between metacommunity archetypes. *Ecography,* 44, 1-12.

Thompson, P.L., Guzman, L.M., De Meester, L., Horvath, Ptacnik, R., Vanschoenwinkle, B.,

Viana, D.S., & Chase, J.M. (2019). A process-based framework for metacommunity ecology.

Ecology Letters, 9, 1314-1329.

Turner MG. (2005). Landscape ecology: what is the state of the science? *Annual Reviews of*

Ecology, Evolution and Systematics, 36, 319–44.

Urban, M.C., Strauss, S.Y., Pelletier, F., Palkovacs, E.P., Leibold M.A., Hendry A.P., De

Meester, L., Carlson, S.M., Angert, A.L., & Giery, S.T. (2020). Evolutionary origins for

ecological patterns. *Proceedings of the National Academy of Sciences, U.S.A.*, in press.

Vellend, M. (2010). Conceptual synthesis in community ecology. *The Quarterly Review of*

Biology, 85, 183–206.

Vellend, M. (2016). The theory of ecological communities. Princeton University Press.

Viana, D.S., & Chase, J.M. (2019). Spatial scale modulates the inference of metacommunity

assembly processes. *Ecology,* 100, e02576.

Warton, D. I., Blanchet, F.G., O'Hara, R.B., Ovaskainen, O., Taskinen, S., Walker, S.C. and

Hui, F.K.C. (2015). So Many Variables: Joint Modeling in Community Ecology. *Trends in*

Ecology & Evolution 30, 766–779.

Weiher, E., Freund, D, Bunton, T.,Stefanski, A., Lee, T., & Bentivenga, S. (2011). Advances,

challenges and a developing synthesis of ecological community assembly theory.

Philosophical Transactions of the Royal Society: B Biological Sciences, 366, 2403-2413.

Wilkinson, D.P., Golding, N., Guillera-Arroita, G., Tingley, R., & McCarthy, M.A. (2020).

Defining and evaluating predictions of joint species distribution models. *Methods in Ecology*

and Evolution, 12, 494-404.

Wiegand, T., Uriarte, M., Kraft, N.J.B., Shen, G., Wang, X., & He, F. (2017). Spatially explicit

metrics of species diversity, functional diversity, and phylogenetic diversity: Insights into

plant community assembly processes. *Annual Review of Ecology, Evolution, and Systematics*,

48, 329–351.

Yang, X, Tan, J., Sun, K.H., & Jiang, L. Experimental demonstration of the importance of

keystone communities for maintaining metacommunity biodiversity and ecosystem

functioning. *Oecologia*, 193, 437-447.

608 Table 1: Summary of metacommunity level variation components for the seven different

609 scenarios modeled in this study.

Scenario	Corresponding Figure	Fractions			
		E (SD)	S (SD)	C (SD)	Residuals $1-R^2$ (SD)
A	2, upper panels	0.75 (0.11)	0.026 (0.019)	0.044 (0.062)	0.18 (0.11)
B	2, lower panels	0.019 (0.015)	0.15 (0.037)	0.019 (0.036)	0.81 (0.04)
C	3, upper panels	0.42 (0.31)	0.21 (0.2)	0.14 (0.16)	0.23 (0.12)
D	3, lower panels	0.018 (0.026)	0.35 (0.3)	0.28 (0.33)	0.35 (0.33)
E	Supplement	0.63 (0.21)	0.067 (0.069)	0.13 (0.15)	0.18 (0.089)
F	Supplement	0.74 (0.097)	0.035 (0.025)	0.047 (0.057)	0.18 (0.097)
G	4	0.5 (0.17)	0.08 (0.072)	0.2 (0.16)	0.22 (0.12)

Figure 1: A summary of the metacommunity problem. Species distributions, denoted by the

species-by-sites **L** matrix, are the outcome of drift, selection, dispersal, and speciation. These

basic processes can be influenced by species interactions, food web structure, biogeography,

phylogeny and micro-evolution. Metacommunity theory mainly focuses on drift, selection and

dispersal. We view previous approaches based on the four archetypes of Leibold et al. (2004) as

being much more indirect and idealized. Instead, we call for a more direct evaluation of how the

basic processes affect the **L** matrix, and how to dissect the consequences to the distributions of

different species and the occupancy of different sites, for example by using a JSDM to identify

main effects and interspecific variability in the importance of unstructured, biotic,

environmental, and spatial effects on **L**. This approach allows us to recognize and address the

effects of heterogeneities among species and among patches on the overall structure of the

metacommunity.

Figure 2: Ternary plots describing the three components of metacommunity internal structure for two different simulation scenarios with no species interactions (independent metapopulations): The upper panels correspond to narrow environmental niches whereas lower panels correspond to wide environmental niches. Each dot represents a species (left panels) or a site (right panels). The size of the symbol is proportional to the R^2 of the model (note the different scales used for species and sites) and the location indicates the proportion of explained variation attributed to environmental factors (E - lower left), spatial effects (S - lower right) and remaining co-distributions (C - upper apex) (see SI for details). In the species panels (left side) different symbols indicate different replicate simulations; generally, these indicate that the distribution of species responses are variable within replicates but that the overall variation among replicates are

repeatable. In the site panels (right side), the shading indicates how central (lighter) or extreme

(darker) the local environmental conditions are on the gradient; these also show substantial

variation but indicate that more extreme environmental conditions increase the effects of local

environment on occupancy patterns than more central conditions.

Figure 3: Ternary plots describing the three components of metacommunity internal structure for

different simulation scenarios with competition among the species. Notation is the same as in

Figure 2. The upper panels correspond to narrow environmental niches whereas the lower panels

correspond to wide environmental niches. Left-hand panels show variation components for

different species whereas panels on the right-hand side of the figure correspond to variation

components for different sites.

Figure 4: Ternary plots for species (left panel) and sites (right panel) for simulations with species

that differ in environmental position along the gradient and dispersal ability. The size of the

symbol indicates the R^2 of the model for each species or site). In the left panel (species) the

color indicates the preferred local environmental conditions for species (yellow for species that

prefer centrally located environmental conditions, purple or magenta for species with more

extreme environmental optima). The symbol indicates the dispersal rate of the species (circles

are more dispersal limited, squares are least dispersal limited and triangles are intermediate). In

the right-hand panel the color indicates the degree of deviation from centrality along the

environmental gradient (as in Figure 2).

Figure 5: Effects of species traits (i.e. species optima; left panel) and site attributes (right panel)

on the environmental fraction of variation in species distributions and site occupancy. A) Higher

dispersal ability and lower niche centrality (i.e. greater deviation from mean niche value)

enhance the degree to which different species (individual symbols) have distributions that

correlate with environmental variation. B) Sites that differ more from the mean environmental

value (environmental deviation) are more likely to be occupied by species with niche traits that

are locally favored.

Figure 6: Comparisons of the interaction matrix (Simulation) with the co-distribution of species in five replicate runs (Iteration 1-5) of the scenario with interspecific variation in dispersal and competition among species. In each panel, species are ranked by the position of their environmental optima along the environmental gradient. The co-distributions are shown as heat maps with the strength of the covariation proportional to the intensity of color and the color indicating negative (green) or positive (gray) covariation among pairs of species. These can be compared to the pattern of direct species interactions (left panel called Simulations). The Xs denote no significant association although the color indicates the trend.

687 Mathew A. Leibold, Javiera Rudolph, Pedro Peres-Neto, Dominique Gravel, Luc De Meester,
688 Lauren Showmaker, Florian Hartig, F. Guillaume Blanchet, Jonathan M. Chase

Table of Contents

Description of Model

Patch locations and environmental variables

In our model, the metacommunity consists of N patches distributed over a spatially
heterogeneous landscape, with multiple environmental variables (although, the current
simulations only have one environmental variable D across 1000 patches) that could either be
randomly distributed or spatially autocorrelated. Each patch has a set of coordinates in a two-
dimensional space, and all possible coordinates are feasible such that this is a continuous space
model that is not restricted to a lattice or some other kind of regular spatial arrangement of
spatial units. A patch may be empty or be occupied by a single or by several species. We define
$X_{i,z,t}$ as a stochastic variable representing the occurrence of species i at location z and time t.
Occurrence, $X_{i,z,t}$, takes a value of 1 when species i is present and a value of 0 when it is
absent. Similarly, we define $\mathbf{Y}_{z,t} = X_{1,z,t}, X_{2,z,t}, \dots, X_{R,z,t}$ as a vector containing the presence-
absence of each species from the regional pool R.

709
The model only tracks patch occupancy (not population densities). Spatial dynamics occurs
because of colonization events, in both empty patches and patches that are occupied by other
species, and because of extinction events. The emerging species co-distributions are a result of
a dynamic balance between these events. Ecological interactions can impact either or both the
colonization and the extinction probabilities. For instance, the presence of a competitor pre-
empting a patch can reduce the colonization probability by another competitor. Alternatively,
the presence of a competitor in a patch could increase the extinction probability of another

species. Similarly, the environment could influence both the colonization and the extinction
probabilities.

Patch Colonization

We consider a discrete-time Markovian process to represent the dynamics of presence-
absence of all species and we incorporate the effect of dispersal, environmental filtering and
ecological interactions in such a way that we could cover all possible scenarios wherein species
differ in any combination of these mechanisms and processes. We can include interspecific
competition along with other types of spatial dynamics such as predator-prey interactions
(Gravel et al. 2011), priority effects (Shurin et al. 2004), or mutualistic interactions
(e.g. Gilarranz et al. 2015). In this paper, we focused on competition only though. Following a
colonization event from time t to $t + \Delta$ corresponds to:

$$P(X_{i,z,t+\Delta t} = 1 \mid X_{i,z,t} = 0) = I_{i,z,t} S_{i,z,t} C_{i,z,t}$$

where $I_{i,z,t}$ is the number of immigrants of species i reaching patch z at time t, $S_{i,z,t}$ is the
effect of environmental filtering on the probability of establishing a viable local population and
$C_{i,z,t}$ is the effect of ecological interactions on the establishment probability. We note that
because we represent a stochastic process, the product of these three functions has to be
bounded between 0 and 1. We consequently define these quantities:

The effect of immigration is given by:

$$I_{i,z,t} = \frac{\sum k(z,\omega) X_{i,\omega,t}}{\sum k(z,\omega)}$$

which is a weighted average of the occurrence probability of species i in the neighborhood
of z. The function $k(z, \omega)$ is a dispersal kernel that depends on the location of patch z and
the neighborhood ω. For convenience, we considered an exponential function of the
Euclidean distance between localities. We added to the kernel a low distance and
neighborhood-independent constant m to account from immigration from outside the
simulated metacommunity. This assumption is required to prevent total extinction by drift
under pure neutral dynamics.

The effect of the environment is given by a product of the establishment performance over
all environmental variables E_n:

$$S_{i,z,t} = \prod f(E_{n,z} \, \mu_{i,n} \, \sigma_{i,n})$$

748   In our simulations, for convenience, we considered that the function f has a quadratic form for
749   all species and all environmental variables, though the model is flexible and general enough to
750   consider other (non-linear) responses that could also differ among species.

Ecological interactions on establishment probability

To incorporate all possible ecological interactions, we started by representing the interaction
network by a community matrix $\mathbf{A}$ of R species. The elements α_{ij} of $\mathbf{A}$ quantify the effect of
species j on the dynamics of species i. When α_{ij} is negative, the colonization probability of
species i decreases and/or its extinction probability increases when j is found locally. Inversely,
when α_{ij} is positive, the colonization probability increases and/or the extinction probability
decreases. To account for the cumulative effects of local interactions on transition probabilities,
we made colonization and extinction probabilities community dependent. As explained above,
at a time t, the $\mathbf{Y}_{z,t}$ vector gives the local assemblages. We calculated the sum of interactions at
any time and for each species as $v = \mathbf{A}_{z,t}\mathbf{Y}_{z,t}$. Our approach can be interpreted as a spatial
analogue to the generalized Lotka–Volterra model because it takes into account the impact of
the whole network of interactions on each species dynamics and can deal with any type of
interaction. We now define the function:

764
$$C_{i,z,t} = g(v_i, z, t)$$

765   representing the total effect of ecological interactions on the colonization probability. For
766   convenience, we will use a sigmoid function, with g ranging between c_{min} at high negative
767   interactions and c_{max} at high positive interactions, where c_{max} should be interpreted as the
768   maximal colonization probability when the environmental conditions are optimal and there are
769   no dispersal limitations.

Patch Extinction

771   The definition of the extinction probability follows exactly the same rules as for colonization,
772   except that extinction is independent of the neighborhood composition. We follow the same
773   logic to define the effect of ecological interactions and of variation in the environment.
774   Consequently, we get the Markovian process:

775
$$P\big(X_{i,z,t+\Delta t} = 1 \,\big|\, X_{i,z,t} = 0\big) = M_{i,z,t}E_{i,z,t}$$

where $M_{i,z,t}$ and $E_{i,z,t}$ are the responses of the extinction probability to the local
environment and to ecological interactions, respectively. The difference with the
colonization functions defined in the previous section is that the extinction probability
must be larger when interactions are negative and smaller when they are positive. In

addition, the extinction rate should be minimal (instead of maximal) at environmental
optimum.

Interpretation

784   To interpret the model, note that, at steady state, for each species, we obtain the expected
785   occurrence probability $(\hat{P})$ at each site as:

$$\frac{\hat{P}^{iz}}{1 - \hat{P}^{iz}} = \frac{I^{iz} \cdot S^{iz} \cdot C^{iz}}{M^{iz} \cdot E^{iz}}$$

787   After a log transformation, this yields:

$$\log\left(\frac{\hat{P}^{iz}}{1 - \hat{P}^{iz}}\right) = \log(I^{iz}) + \log\left(\frac{S^{iz}}{M^{iz}}\right) + \log\left(\frac{C^{iz}}{E^{iz}}\right)$$

This last equation can be interpreted as a macroscopic description of the expected species
distributions pattern (Thuiller et al. 2013). In this formulation, $\log(I)$ describes the
tendency of a patch to resemble other nearby patches due to the spatial contagion by
dispersal, $\log\left(\frac{S}{M}\right)$ describes the tendency of sites to be occupied by species with similar
fitness responses to environmental gradients, and $\log\left(\frac{C}{E}\right)$ describes the remaining
influence of other species on co-occurrence due to interactions among species. The values
for these indices will depend on what choices are made for the components of eq. 1 (see
Supporting Information for details on how we implemented this simulations model).

This modeling framework can represent the classical archetypes but also permits more
intricate (and likely far more realistic) metacommunity scenarios and predictions. For
example, we could use the model to examine how species traits (and environmental
context) link to metacommunity dynamics. Moreover, continuous mixtures of different
metacommunity extremes (archetypes) can be represented by appropriate parameter
choices for dispersal, competitive abilities, and environmental preferences. For instance,
species sorting would require a relatively large colonization to extinction ratio along with
species-specific environmental requirements and regional similarity (sensu Mouquet and
Loreau, 2002). Alternatively, coexistence within competition-colonization trade-offs
requires species to have similar responses to the environment and appropriate
heterogeneities in the I, C and E functions, but no environmental preferences.

The implemented mechanisms in the simulation model can be partially mapped onto
variation partitioning components. For instance, at equilibrium, we could expect dispersal

limitation (the $\log(I)$ term in equation 3) to create positive spatial autocorrelation at the

dispersal scale (the [S/E] fraction in variation partitioning, i.e., spatial variation

independent of environmental selection). Environmental selection (the $\log\left(\frac{S}{M}\right)$ term in the

last equation) should lead to a correlation between composition and environment (the

[E/S] fraction in variation partitioning). The last term in equation 3, however, describing

the effect of interactions on distribution (the $\log\left(\frac{C}{E}\right)$), is novel and has no equivalent in the

context of classical variation partitioning.

There are some interesting properties to point out regarding our proposed variation

partitioning scheme. First, by considering the combined effects of environmental selection,

dispersal and interactions, the final residuals (unexplained sources of variation) in the

model leading to this new partition variation scheme is (in principle) solely related to non-

spatialized independent species variation. Second, in our variation partitioning, the

interaction component is due to species co-variation (i.e., a joint component among species

distributions). In empirical community data, however, this interpretation can only be made

if all the environmental variation (predictors) underlying environmental selection in

empirical community data has been incorporated (as pointed out in the main manuscript).

If not, then the spatial and species interaction components could be measuring variation

related to unmeasured environmental variables that are either spatialized (i.e.

characterized by the spatial component in variation partitioning) or shared among species

(i.e. joint component).

Description of the Statistical Framework

Hierarchical Community Models

In their simplest form, Hierarchical Community Models (HCMs) resemble standard species

distribution models that regress species presences/absences against environmental predictors

(i.e., logit link). However, to reduce model complexity, HCMs assume that all species in a

metacommunity will react to environmental heterogeneity following a similar response

function (e.g., linear vs quadratic or Gaussian). The same assumption is made in common

variation partitioning (see Peres-Neto et al. 2006). To model the spatial component (i.e., due to

spatialized dispersal), either spatial variables such as Moran's eigenvectors maps (MEM, Dray et

al. 2006) or spatially auto-correlated latent variables (Ovaskainen et al. 2016b) can be

incorporated to the model. To account for biotic interactions, non-spatially auto-correlated

latent variables are used. If we use a linear specification approach (here, this can also include

quadratic terms that capture Gaussian responses to environment as imposed in our model), we

can write:

$$\mathbf{L}^{zi} = \mathbf{X}^{zk}\mathbf{B}^{ki} + \boldsymbol{\epsilon}^{zi}$$

with

$$\mathbf{B}^{ki} \tilde{} \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

where $\mathbf{L}^{zi}$ is the presence (or absence) of species i (out of m species) at patch z (out of n
patches), $\mathbf{X}^{zk}$ is the value of the environmental variable k (out of p variables) at site z, $\mathbf{B}$ is
a matrix of regression parameters, $\boldsymbol{\mu}$ is a vector of length p that describes the mean
environmental response of all species, Σ is a $p \times p$ covariance matrix that describes how
species vary (diagonal) and co-vary (off-diagonal) around the mean environmental
response (Ovaskainen and Soininen 2011), and $\boldsymbol{\epsilon}^{zi}$ is a residual value. Estimating species
parameters hierarchically around a community mean reduces the degrees of freedom and
makes the model easier to fit with limited data. Note that both $\boldsymbol{\mu}$ and Σ can be further
informed or constrained by species traits or phylogeny if desired (Ovaskainen et al. 2017).
To account for biotic interactions, we consider latent variables $\mathbf{H}^{zl}$ (where l refers to a
latent variable measured at site z) and their associated parameters $\boldsymbol{\Lambda}^{li}$ (Ovaskainen et
al. 2016a). This yields:

$$\mathbf{L}^{zi} = \mathbf{X}^{zk}\mathbf{B}^{ki} + \mathbf{H}^{zl}\boldsymbol{\Lambda}^{li} + \boldsymbol{\epsilon}^{zi}$$

Note that it is not necessary to always include all of these components in one model; they
can be considered in any combination deemed relevant for a particular question. In this
paper, we used Moran's Eigenvector Maps (MEMs; Dray et al. 2006), a powerful and
commonly used method to model spatial autocorrelation in statistical models involving
species distributions.

Calculating Variation Partitioning for the HCM

As in any generalized linear mixed effect model, we can now partition the explained
variation into different components, notably environmental heterogeneity, space, co-
distribution (biotic interactions), and unexplained variation (Figure S1). To estimate the
contributions of each of these four fractions for each species, we calculated semi-partial
coefficients of determination (i.e., based on Type III sum-of-squares as specified in Peres-
Neto et al. 2006) using the implementation suggested by Tjur (2009) as being more
appropriate for presence-absence data (i.e., logit link) than the traditional variation
partitioning based on an identity link. To adjust for the number of variables used to
quantify each fraction of the variation partitioning analysis, we applied the adjustment to

the coefficient of determination proposed by Gelman and Pardoe (2006) in the variation
partitioning analysis, which is designed for hierarchical models. As shown in Figure S1, the
different fractions were combined so that a unique value was associated to environment
(fractions [a], [d]/2, [f] and [g]/2), co-distribution (fraction [c]), space (fractions [b], [d]/2,
[e] and [g]/2) and the unexplained portion of the variation (fraction [h]). Latent variables
are quite powerful to isolate structure in the data. As such, in the calculation of the
variation partitioning, latent variables will capture almost all (if not all) variation
associated to the environment and space, giving an artificial inflation of the overlapping
partitions between co-distribution and environment and co-distribution and space. For this
reason, all partitions overlapping with co-distribution (fractions [e], [f] and [g]) were
assigned to either environment (fractions [f] and [g]) or space (factions [e] and [g]). In this
calculation, a unique measure of explained variation (akin to adjusted R^2) is associated to
co-distribution (fraction [c]) but this is not the case for environment and space. To
associate a unique value to environment and space, and represent the results as we did in
Figure 2 and 3 (main manuscript), we divided the fractions overlapping environment and
space between these two components. As such, the sum of fractions [a], [f], half of fraction
[d] and half of fraction [g] were used to measure the effect of the environment while
fractions was considered [b], [e], half of fraction [d] and half of fraction [g] were used to
measure the effect of space. This scheme in which half of common variation is assigned to
two or more common components is commonly used in hierarchical partitioning (Chevan &
Sutherland 1991).

904 .

905   Supplementary Figure 1 – variation partitioning scheme used to estimate the importance of
906   each matrix of predictors.

Calculation of the coefficient of determination

1) Classic coefficient of determination

909   The coefficient of determination, R^2, that was partitioned in the variation partitioning analysis
910   (Appendix XX) is calculated for any given species j as:

$$R_j^2 = 1 - \frac{\sum_{i=1}^{n}(y_{ij} - \hat{y}_{ij})^2}{\sum_{i=1}^{n}(y_{ij} - \overline{y}_{ij})^2}$$

912   where y_{ij} is the data (presence-absence) associated with species j (out of p species) at site i
913   (out of n sites), $\hat{y}_{ij}$ is the model (predicted value) associated to species j at site i and $\overline{y}_j$ is the
914   average of the data (i.e., sum of presences divided by n) for species j across all sites.

2) Community-level coefficient of determination

916   Although having an R_j^2 for each species j can be highly informative and is part of our framework
917   on the internal structure of metacommunities, it can be also useful to estimate the contribution
918   of single communities R^2 to the entire metacommunity. This is obtained by averaging all R_j^2:

$$^C R^2 = \frac{\sum_{j=1}^{p} R_j^2}{p}$$

920   where the $^C R^2$ is the community-level R^2.

3) Site contribution to the coefficient of determination

922   In the paper, we use the contribution of each site to R_j^2 to present how each site contributes
923   differently to the environment, space and co-distribution for the community. The calculation of
924   the site i contribution to the R_j^2, is calculated as:

$$R_{ij}^2 = \frac{1}{n} - \frac{(y_{ij} - \hat{y}_{ij})^2}{\sum_{j=1}^{p}(y_{ij} - \overline{y}_{ij})^2}$$

The first part of the equation where the 1 of the classic R_j^2 is divided by n is included to make
sure that if we sum all R_{ij}^2 across all sites for species j, the resulting value equals to R_j^2.

More importantly, what can be noticed is that by calculating R_{ij}^2, the contribution of sites to
each species R_j^2, we obtain a matrix that has the same dimension as the site by species
matrix (an $n \times p$ matrix). Using this matrix, if we sum across all sites, we obtain the R_j^2.

However, if we average across the species for site i we obtain the site's contribution to the

community-level $^C R^2$ or $^C R_i^2$.

The amount of variation expressed by the R_j^2, the $^C R^2$, the R_{ij}^2 or the $^C R_i^2$ can all be

partitioned in its environmental, spatial and co-distribution component following the

procedure presented in the section "Calculating Variation Partitioning for the HCM" above.

Parameterization and Simulation Scenarios

We simulated metacommunity dynamics with a landscape of 1000 patches over 200 time steps

and an initial occupancy of 0.8. Patches were placed randomly in a two-dimensional plane with

coordinates drawn from a uniform distribution with a minimum of 0 and a maximum of 1. The

environment varied spatially, with values drawn from a random distribution between 0 and 1.

In the specific simulations we studied in the paper, colonization was the only component of the

species that were affected by the environment (i.e. $E_{i,z,t} = 1$). Specifically, colonization reacted

to the environment following a quadratic curve.

For all scenarios considered, we simulated 12 species. Niche optimums for the species were

evenly distributed between 0.1 and 0.9 while niche breadth was set to 0.8 for simulations

with narrow niches (scenarios A, B, E, F, G), and to 2 for simulations where niche was

assumed to be broad (scenarios C and D). For dispersal, we considered an exponential

dispersal kernel, with a distance-independent immigration probability of 0.001 and an α

parameter of 0.05. For scenario G, where we have variable dispersal kernels (Figure 3), α

was 0.01 for 1/3 of species, 0.05 for 1/3 of species, and 0.1 for the other 1/3 of species. We

used a sigmoid function to relate the total number of interactions with colonization and

extinction coefficients following the implementation by Cazelles et al. (2016). Colonization

probability in the absence of interactions was set at 0.4, which tends to zero as negative

interactions tend to infinity, while it asymptotes at a 1 with infinite positive interactions.

All other aspects of the colonization-interaction curve were the same for all scenarios.

Similarly, extinction in the absence of interactions was set at 0.025, and tended to 1 with

infinite negative interactions, while its asymptote tended to 0 with infinite positive

interactions. In both cases, the parameter setting the shape of the sigmoid function was set

to 0 for the scenarios without competition (scenarios A, B, and F), and 1.5 in the presence

of competition (scenarios C, D, E, G) . If there were interactions, then a focal species only

interacted with the two species that had the closest niches. For all scenarios, five sets of

metacommunities were simulated and analyzed to obtain the results found in Figures 2, 3,

and 4 in the main text, and supplementary figures.

All scenarios have been implemented in R and the project's repository can be found here:

github.com/javirudolph/testingHMSC

Supplementary Figure 2. Species interactions for all scenarios.

Supplementary Figure 3. In which we have half of the species with interactions and the
other without.

Supplementary Figure 4. In which we change dispersal only, α was 0.01 for 1/3 of species,
0.05 for 1/3 of species, and 0.1 for the other 1/3 of species.

Example of metacommunity simulation functions and analyses

974
975

The following example of code shows the overall processes involved in the metacommunity
simulation for our model. This example shows the scenario for 20 patches and one
environmental variable. The model gives the option for a random or spatially aggregated
structure for the patches. In the aggregated case, we determined four clusters, denoted by
Nclusters in the code below. The value of the environmental variable for each patch is shown
with the color hue. In this case, the environmental variable is randomly distributed.

```r
set.seed(227)
# Random XY coordinates
# Each coordinate is drawn from a random uniform distribution
get_XY = function(N) cbind(runif(N),runif(N))

# Aggregation of XY coordinates
get_XY_agg = function(N, Nclusters, sd_xy) {

  Xclust = runif(Nclusters)
  Yclust = runif(Nclusters)

  X = rnorm(N, rep(Xclust,N/Nclusters), sd_xy)
  Y = rnorm(N, rep(Yclust,N/Nclusters), sd_xy)

  cbind(X,Y)
}

# Random uniform environmental values
get_E = function(D, N) matrix(runif(D*N), nr = N, nc = D)
```

Here, we set N, the number of patches to 20, and D, the number of environmental variables to
one.

```r
N <- 20
D <- 1
```

```
1005
1006    rXY <- get_XY(N)
1007    agXY <- get_XY_agg(N, 4, 0.02)
1008    E <- get_E(D = D, N = N)
```

1009   The following figure shows a side by side comparisson between random and aggregated
1010   patches obtained from our functions.

1011

Initial Occupancy

The toy model allows for setitng the initial occupancy in the metacommunity. For example, to
create the initial conditions, $t = 0$, of presence absence, species occupancy is drawn from a
random uniform distribution, and values smaller than 0.8 are considered as species presence.
Patches or locations z are represented by rows in the matrix, whereas each species is a column.
Each cell in the matrix is $X_{i,z,t}$ and each row is $\mathbf{Y}_{z,t}$ for $t = 0$. The following figure shows three
different iterations of this process, where areas in black represent occupancy = 1, and white
denotes an absence.

```
1020    #Get your initial conditions:
1021    R <- 8
1022    Y0 = matrix(0, nrow = N, ncol = R)
1023    rand = matrix(runif(N*R), nr = N, nc = R)
1024    Y0[rand < 0.8] = 1
1025
1026    ggplot(melt(Y0), aes(x = Var1, y = Var2, fill = value)) +
1027      geom_tile() +
1028      scale_fill_gradient(low = "white", high = "black")+
1029      labs(x = "Plot number", y = "Species", subtitle = "Iteration 1") +
1030      theme_bw() -> A
1031
1032    # Second iteration for the same initial conditions
1033    Y0 = matrix(0, nrow = N, ncol = R)
1034    rand = matrix(runif(N*R), nr = N, nc = R)
1035    Y0[rand < 0.8] = 1
1036
1037    ggplot(melt(Y0), aes(x = Var1, y = Var2, fill = value)) +
1038      geom_tile() +
1039      scale_fill_gradient(low = "white", high = "black")+
1040      labs(x = "Plot number", y = ""  , subtitle = "Iteration 2") +
1041      theme_bw() -> B
```

```
1042
1043    # Third iteration
1044    Y0 = matrix(0, nrow = N, ncol = R)
1045    rand = matrix(runif(N*R), nr = N, nc = R)
1046    Y0[rand < 0.8] = 1
1047
1048    ggplot(melt(Y0), aes(x = Var1, y = Var2, fill = value)) +
1049      geom_tile() +
1050      scale_fill_gradient(low = "white", high = "black")+
1051      labs(x = "Plot number", y = "", subtitle = "Iteration 3") +
1052      theme_bw() -> C
1053
1054    # Set the three iterations to be displayed side by side
1055    ggarrange(A, B, C, ncol = 3, common.legend = TRUE, legend = "none")
```

1056

When we consider the immigration component, we need to also consider the connectivity matrix. In the code below, *I_f* calculates the probabilty of immigration for each species, based on the occupancy matrix and the dispersal kernel, K. The argument *K* is the connectivity matrix. The argument *XY* corresponds to the patch coordinates, whereas *alpha* is the dispersal parameter associated to the exponential distribution used for dispersal. It can be computed with the following function:

```
1063    # Compute the propagule pressure
1064    I_f = function(Y, K, m) I = (1-m)*(K%*%Y)/(K%*%matrix(1,nr=N,nc=R)) + m
```

The arguments for this function are: *Y, K, m*. We calculated $\mathbb{Y}$, species presence or absence, in the previous section with the case for initial conditions. Argument *m* is set in the parameters as a value *m = 0.001* and the connectivity matrix *K* is calculated below.

```
1068    # Compute the connectivity matrix
1069    get_K = function(XY, alpha) {
1070        N = nrow(XY)
1071        distMat = as.matrix(dist(XY, method = "euclidean", upper = T, diag = T))
1072        ConMat = exp(-1/alpha*distMat)
1073        diag(ConMat) = 0
1074        return(ConMat)
1075    }
```

1076   As an example, using the aggregated XY coordinates for 20 patches and our initial occupancy
1077   matrix with 8 species ,we can see the connectivity between patches, and can calculate the
1078   contribution of immigration from each species to each patch.

```
1079    # We can use the aggregated XY coordinates for this example:
1080    XY <- agXY
1081
1082    # Connectivity matrix
1083    alpha <- 0.005
1084    K <- get_K(XY, alpha)
1085    plot_K <- cbind(rownames(K), stack(as.data.frame(K)))
1086    names(plot_K) <- c("X", "Fill", "Y")
1087    plot_K %>%
1088      as_data_frame() %>%
1089      mutate(X = factor(X, levels = c(1:20)),
1090             Y = factor(Y, levels = c(1:20))) -> plot_K
1091    ggplot(plot_K, aes(x = X, y = Y, fill = Fill)) +
1092      geom_raster() +
1093      scale_fill_gradient(low = "black", high = "white") +
1094      #scale_fill_viridis_c() +
1095      labs(title = "Connectivity among patches") +
1096      theme(legend.title = element_blank(),
1097            axis.text.x = element_text(angle = 90)) -> A
1098    # Immigration
1099    m <- 0.001
1100    Y <- Y0
1101    I <- I_f(Y, K, m)
1102    rastPlot(I, title = "I - Immigration to each plot", x = "Patches", y = "Species") +
1103      theme(legend.title = element_blank()) -> plot_I
1104    ggarrange(A, plot_I, ncol = 2)
```

1105

1106   The effect of the environment on each species, depending on each species niche optima, is
1107   computed in the following code section. The argument E corresponds to the vector of values
1108   for the environmental variable in each patch. The other two arguments in this function are
1109   the niche optima (u_s) for each species and niche breadth(s_c).

```
1110    # Compute the local performance of propagules
1111    S_f_quadratic <- function(E, u_c, s_c) {
1112      R <- ncol(u_c)
1113      N <- nrow(E)
1114      D <- ncol(E)
1115      S <- matrix(1, nr = N, nc = R)
```

```
1116    for(i in 1:D){
1117      optima <- matrix(u_c[i,],nrow = N,ncol = R,byrow = TRUE)
1118      breadth <- matrix(s_c[i,],nrow = N,ncol = R,byrow = TRUE)
1119      S <- S * ((-1 / (breadth/2)^2) * (E[,i] - optima)^2 + 1)
1120      S <- ifelse(S < 0, 0 , S)
1121
1122    }
1123    return(S)
1124  }
1125
1126  # Understood as niche optima for each species, for each environmental variable
1127    u_c = matrix(nr = D, nc = R)
1128    u_c[1,] = seq(0.1,0.9, length=R)
1129  # Understood as niche breadth
1130    s_c = matrix(0.2, nr = D, nc = R)
1131
1132  # Local performance, colonization
1133  S <- S_f_quadratic(E, u_c, s_c)
1134  plot_S <- rastPlot(S, title = "S - Local performance colonization", x = "Patches", y = "Specie
1135  s") +
1136    guides(fill = guide_colorbar(title = ""))
1137  plot_S
```

When incorporating species interactions into the toy model, we use the following interaction matrix **A**, where the colored black sections show species with potential of interacting:

```
1142    # # Interaction matrix
1143    A = matrix(0,nr=R,nc=R)
1144    d = as.matrix(dist(c(1:R),upper=TRUE,diag=T))
1145    A[d<=1] = -1
1146    diag(A) = 0
1147
1148    plot_A <- rastPlot(A, title = "A - Interaction matrix", x = "species", y = "species") +
1149      scale_fill_gradient(low = "black", high = "grey") +
1150    theme(legend.position = "none")
```

```
1151    # Compute the sum of ecological interactions for every location and every species
1152    sum_interactions = function (A, Y) t(A%*%t(Y))
1153    # this is considered to be "v"
1154    v <- sum_interactions(A, Y)
1155
1156    plot_v <- rastPlot(v, title = "v - Sum of interactions", x = "patches", y = "species") +
1157      guides(fill = guide_colorbar(title = "")) +
1158      scale_fill_gradient()
1159
1160    ggarrange(plot_A, plot_v, ncol = 2)
```

1161

1162   We compute the effect of ecological interactions on colonization probability with the
1163   function below.The arguments for this function are v as the resulting matrix from the sum
1164   of interactions, d_c as the sensitivity to interactions, c_0 and c_{max} as the colonization
1165   parameters:

```
1166    C_f = function(v, d_c, c_0, c_max) c_max*(1 +(1/c_0 - 1)*exp(-v*d_c))^-1
1167
1168      # # Colonization function
1169      c_0 = rep(0.4, R) # Colonization at 0 interactions
1170      c_max = rep(1, R) # Colonization at max interactions
1171
1172      # # Sensitivity to interactions
1173      d_c = 0.2
1174
1175    C <- C_f(v, d_c, c_0, c_max)
1176    plot_C <- rastPlot(C, title = "C - Interactions on colonization", x = "Patches", y = "Species
1177    ") +
1178      guides(fill = guide_colorbar(title = "")) +
1179      scale_fill_gradient()
1180    plot_C
```

1181

1182   With all the components calculated, we can now compute the colonization probability

1183 $P(X_{i,z,t+\Delta t} = 1 \mid X_{i,z,t} = 0) = I_{i,z,t} S_{i,z,t} C_{i,z,t}$

1184

1185   The following function calculates the effect of the environment on the extinction, with E

1186   being the environmental variable, and u_e and u_s being species level effect and the

1187   assymptote.

```
1188   M_f = function(E, u_e, s_e) {
1189       R = ncol(u_e)
1190       N = nrow(E)
1191       D = ncol(E)
1192       M = matrix(1, nr = N, nc = R)
1193       for(i in 1:D){
1194         M = M*(1-exp(-(E[,i]-matrix(u_e[i,],nr=N,nc=R,byrow=TRUE))^2 / matrix(s_e[i,],nr=N,nc=R,
1195   byrow=TRUE)^2))
1196       }
1197       return(M)
1198   }
1199
1200   # Set the function arguments
1201     # # Effect of the environment on extinction
1202     u_e = matrix(nr = D, nc = R)
1203     u_e[1,] = c(rep(0.5, R-1), 0.05) # One species having a lower level of extinction from envir
1204   onmental effect
1205     s_e = matrix(Inf, nr = D, nc = R)
1206
1207     #u_e
1208     #head(s_e)
```

The following shows the effect of ecological interactions on extinction, using the same v matrix calculated above.

```
1211   E_f = function(v, d_e, e_0, e_min) {
1212
1213     e_min_mat = matrix(e_min, nr = N, nc = R, byrow=TRUE)
1214
1215     e_min_mat+(1/(1-e_min_mat)+(1/(e_0-e_min_mat)-1/(1-e_min_mat))*exp(d_e*v))^-1
1216
1217   }
1218
1219
1220   #With the arguments computed as:
1221
1222
1223   # # Extinction function
1224     #e_0 = c(rep(0.025, R-1), 0.5) # Extinction at 0 interactions, with one species having a hig
1225   her value.
1226   e_0 = rep(0.025, R)
1227   e_min = rep(0, R) # Exinction at max interactions
1228
1229     # # Sensitivity to interactions
1230
1231     d_e = 0
```

We can now compute the probability of extinction $P\left(X_{i,z,t+\Delta t} = 1 \,\middle|\, X_{i,z,t} = 0\right) = M_{i,z,t}E_{i,z,t}$

The figure shows these as identical graphs, since we have made all species have the same probability of extinction and the environment not having an effect on extinction.

1235   The way
1236   parameters are set, the extinction component is the same for all species in every patch.

Testing and changes

```
# Perform the test
delta <- matrix(0, nr = N, nc = R)
rand <- matrix(runif(N*R), nr = N, nc = R)
delta[Y == 0 & rand < P_col] <- 1

# Perform the test
rand = matrix(runif(N*R), nr = N, nc = R)
delta[Y == 1 & rand < P_ext] = - 1
```

1246

1247   References:

1248   Chevan, A., & Sutherland, M. (1991). Hierarchical partitioning. *American Statistician*, 45, 90-
1249   96.

Dray, S., P. Legendre, & Peres-Neto, P.R. (2006). Spatial modeling: a comprehensive framework for principal coordinate analysis of neighbor matrices (PCNM). *Ecological Modelling*, 196, 483-493.

Efron, B. (1978). Regression and ANOVA with zero-one data: Measures of residual Variation. *Journal of the American Statistical Association*, 73, 113–121.

Gelman, A., & Pardoe, I. (2006). Bayesian Measures of Explained Variance and Pooling in Multilevel (Hierarchical) Models. *Technometrics*, 48, 241–251.

Gilarranz, L. J., Sabatino, M., Aizen, M. A., & Bascompte, J. (2015). Hot spots of mutualistic networks. *Journal of Animal Ecology*, 84, 407-413.

Gravel, D., Massol, F., Canard, E., Mouillot, D. & Mouquet, N. (2011). Trophic theory of island biogeography. *Ecology Letters*, 14, 1010–1016.

Ovaskainen, O., Abrego, N., Halme, P. & Dunson, D. (2016a). Using latent variable models to identify large networks of species-to-species associations at different spatial scales. *Methods in Ecology and Evolution*, 7, 549–555.

Ovaskainen, O., Hottola, J. & Siitonen, J. (2010). Modeling species co-occurrence by multivariate logistic regression generates new hypotheses on fungal interactions. *Ecology*, 91, 2514–2521.

Ovaskainen, O., Roy, D.B., Fox, R. & Anderson, B.J. (2016b). Uncovering hidden spatial structure in species communities with spatially explicit joint species distribution models. *Methods in Ecology and Evolution*, 7, 428–436.

Ovaskainen, O., & Soininen, J. (2011). Making more out of sparse data: hierarchical modeling of species communities. *Ecology*, 92, 289–295.

Ovaskainen, O., Tikhonov, G., Norberg, A., Guillaume Blanchet, F., Duan, L., Dunson, D., et al. (2017). How to make more out of community data? A conceptual framework and its implementation as models and software. *Ecology Letters*, 20, 561–576.

1284

1285 Peres-Neto, P.R., Legendre, P., Dray, S., & Borcard, D. (2006). Variation partitioning of species
1286 data matrices: Estimation and comparison of fractions. *Ecology*, 87, 2614–2625.

1287

1288 Shurin, J.B., Amarasekare, P., Chase, J.M., Holt, R.D., Hoopes, M.F. & Leibold, M.A. (2004).
1289 Alternative stable states and regional community structure. *Journal of Theoretical Biology*, 227,
1290 359–368.

1291

1292 Tjur, T. (2009). Coefficients of determination in logistic regression models-A new proposal: the
1293 coefficient of discrimination. The *American Statistician*, 63, 366–372.

1294