No Evidence for Long-range Male Sex Pheromones in Two Malaria Mosquitoes

Serge Bèwadéyir Poda[1,2,3], Bruno Buatois[4], Benoit Lapeyre[4], Laurent Dormont[4], Abdoulaye Diabaté[1], Olivier

Gnankiné[2], Roch K. Dabiré[1], Olivier Roux[1,3*]

1.   *Institut de Recherche en Sciences de la Santé (IRSS), Bobo-Dioulasso, Burkina Faso.*

2.   *Laboratoire d'Entomologie Fondamentale et Appliquée (LEFA), Unité de Formation et de Recherche en*

Sciences de la Vie et de la Terre (UFR-SVT), Université Joseph KI-ZERBO, Ouagadougou, Burkina Faso.

3.   *MIVEGEC, Univ. Montpellier, IRD, CNRS, Montpellier, France.*

4.   *CEFE, Univ. Montpellier, CNRS, EPHE, IRD, Univ. Paul Valéry Montpellier 3, Montpellier, France.*

*corresponding author

Olivier Roux: olivier.roux@ird.fr

Running head: Sex pheromones in malaria mosquitoes

Abstract - Cues involved in mate seeking and recognition prevent hybridization and can be involved in speciation

processes. In malaria mosquitoes, females of the two sibling species *Anopheles gambiae s.s.* and *An. coluzzii* mate

in monospecific male swarms and hybrids are rare. Long-range sex pheromones driving this behavior have been

debated in literature but to date, no study has proven their existence or their absence. Here, we attempted to bring

to light their existence. To put all the odds in our favor, we used different chemical ecology methods such as

behavioral and electrophysiological assays as well chemical analyses, and we worked with mosquitoes at their

optimal physiological mating state *i.e.* with swarming males during their natural swarming windows. Despite all

our efforts, our results support the absence of long-range sex pheromones involved in swarm detection and

recognition by females. We briefly discuss the implications of this finding in ecology, evolution and for control

strategies.

 INTRODUCTION

Cues involved in mate seeking and mate recognition prevent hybridization and can be involved in speciation

processes. In insects, these cues can be visual, acoustical or chemical with a short or a long-range action and are

highly species specific (Alexander et al. 1997; Clements 1999).

In mosquitoes, many species mate in flight, within individual aggregations called "mating swarms"

(Gibson 1945; Downes 1969; Savolainen 1978; Shelly and Whittier 1997; Sivinski and Petersson 1997). In malaria

mosquitoes of the *Anopheles gambiae* complex, mating takes place at sunset in monospecific swarms containing

a few males to thousands of them in which virgin conspecific females come to find a mate (Charlwood and Jones

1980; Marchand 1984; Charlwood et al. 2002; Diabaté et al. 2003, 2009; Howell and Knols 2009; Sawadogo et al.

2014). In West Africa, *An. coluzzii* and *An. gambiae s.s.* are often found in sympatry but they form distinct swarms

spatially segregated and hybrids are rare (≈1%) (della Torre et al. 2001; Tripet et al. 2001; della Torre et al. 2005;

Diabaté et al. 2006; Costantini et al. 2009; Diabaté et al. 2009; Sawadogo et al. 2013, 2014). No evidence for

selection against hybrids was found and spermatheca analyses showed that these species mostly mate assortatively

(Persiani et al. 1986; Tripet et al. 2001; Diabaté et al. 2005, 2007; Hahn et al. 2012; Pombi et al. 2017). This

suggests that reproductive isolation between these two sibling species is achieved by strong pre-mating

reproductive barriers (della Torre et al. 2001; Diabaté et al. 2007; Lehmann and Diabaté 2008). In addition, since

females usually only mate once in their lives (Clements 1992), errors in the choice of mate should be costly and

fall under negative selection. Consequently, one would expect to find specific cues that lead females to conspecific

male swarms. However, the way females are attracted to swarms is unknown.

In species of the *An. gambiae* complex, several cues have been identified to play a role in bringing sexes

together. Acoustic cues were shown to be involved in close-range recognition, the male and the female adjusting

their respective wing-beat frequencies to converge on a shared harmonic frequency (Gibson et al. 2010; Pennetier

et al. 2010). However, it was recently demonstrated that females of some *Anopheles* species are able to detect

swarm sounds only in a very close vicinity. Thus, they are unable to use swarm sound to locate or identify swarms

at long range (Feugère et al. 2021). Visual cues also play an important role in species segregation with *An. coluzzii*

males swarming over contrasted visual ground markers (marker, hereafter) and *An. gambiae* males swarming over

bare ground (Diabaté et al. 2009). However, a recent work showed that, like *An. coluzzii*, *An. gambiae* males also

use visual markers but rather to locate their swarm at a distance from the marker (Poda et al. 2019). Moreover,

females also use these markers to form swarms in the absence of males suggesting that females may use these

markers to join the swarm location (Poda et al. 2019). Nevertheless, the distance between two heterospecific

swarms using the same visual ground marker can be about 2 m in semi-field conditions (Poda et al. 2019) and it is

still unknown from which distance mosquitoes can see such markers, suggesting that females could cross the

location of heterospecific swarms by accident. Chemical cues have also been under investigation in *Anopheles*

species. Heptacosane, a cuticular hydrocarbon, enhances the interaction between males and females (Wang et al.

2021). This compound, however, can only be perceived by contact and can consequently be involved only in mate

recognition during courtship and to stimulate acceptance by females. Moreover, the absence of assortative mating

in confined heterospecific males and females in laboratory cages or indoor swarms (Dao et al. 2008), suggests that

close-range mating cues, if they exist, cannot ensure total reproductive isolation by themselves. Species isolation

is thus likely to occur through long-range and specific swarm recognition cues acting as a first barrier in pre-mating

processes which prevent hybridization by limiting contact between sexes of the different species.

To our knowledge, volatile sex pheromones in the *An. gambiae* complex have never been brought to light

without ambiguity. Charlwood et al. (2002) reported an absence of response of *An. gambiae* males in natural

swarms to squashed females on filter paper or to living females in a net cage. However, in natural conditions,

females are the ones attracted to male aggregation sites. Consequently, males should be the ones that emit long-

range pheromones and should be the attractive sex. Nevertheless, a laboratory study failed to demonstrate any

attractiveness of dead males to virgin females in a Y-tube olfactometer (Gomulski 1988). However, the difficulties

in highlighting the existence of such pheromones might be due to an emission in very low quantities and/or

exclusively during swarming. Indeed, in sandfly species, which also mate in large aggregations of males, it was

shown that the concentration of male-produced sex pheromones is greater in larger swarms, and hence, the chances

for individual males of mating with a conspecific female are also greater (Kelly and Dye 1997; Bray et al. 2010).

Similarly, Diabaté et al. (2011) showed that *An. coluzzii* females were more frequently attracted to large swarms,

suggesting a potential additive effect of cues released by males. Males gathering in large swarms may increase

their "detectability" in the female's olfactory landscape (Shelly and Whittier 1997). In addition, in many insects,

pheromone release and receptivity were shown to respond to diel rhythms or even strict time windows of only a

few minutes or hours (Bjostad et al. 1980; Merlin et al. 2007; Rund et al. 2013; Levi-Zada et al. 2014) or even to

be dependent of the presence of conspecifics (Andersson et al. 2007; Robledo and Arzuffi 2012). Nevertheless, a

recent work by Mozūraitis et al. (2020) and concomitant to a first version of the present work (Poda et al. 2021),

highlighted the existence of five volatile compounds potentially emitted by *Anopheles* males. According to the

authors, these might be involved in aggregation behavior, attracting both males and females, and increasing the

insemination rate. However, though an interesting biological activity was described by the authors, the claim that

these compounds are male swarming aggregation pheromones must be considered with caution and it requires

further investigations. Indeed, these five compounds (acetoin, sulcatone, octanal, nonanal and decanal) are very

frequently found in nature and, as reported by the authors, have been shown to have a biological activity in

Anopheles mosquitoes in very different contexts from the reproductive behavior. They were repeatedly found in

human and animal body odor (Verhulst et al. 2010; Pandey and Kim 2011; Dormont et al. 2013b, a; Tchouassi et

al. 2013; McBride et al. 2014), breath (Poli et al. 2010; Calenic and Amann 2014; Filipiak et al. 2014; Cainap et

al. 2020), and in host-plants (Dekel et al. 2019). These represent the main sources of food (blood and sugar meals,

respectively) for *Anopheles* mosquitoes. As a matter of fact, octanal, nonanal and decanal are part of a blend used

in some traps to mimic mammalian host odor for several species of mosquito vectors (Tchouassi et al. 2013;

Nyasembe et al. 2014). Some of these compounds were also found associated to oviposition sites (Suh et al. 2016;

Wondwosen et al. 2016, 2018) and to ambient air (Kostiainen 1995; Kruza et al. 2017). In addition, sulcatone is

thought to be responsible for discrimination between humans and animals in human-seeking mosquitoes (McBride

et al. 2014). Thus, it is not surprising that they trigger a strong biological/flight activity in mosquitoes.

Nevertheless, the discovery of sex pheromones or a highly attractive blend in malaria mosquitoes would

be a precious step toward the development of new control and monitoring strategies. Furthermore, sex or

aggregation pheromones could help in designing sexually competitive mosquitoes in sterile insect and gene drive

techniques. Here, we investigated the existence of long-range sex pheromones in *An. gambiae* and *An. coluzzii*

mosquitoes, that may allow the females to detect, recognize and track conspecific male swarms. On the premise

that such pheromones could be produced only by males during swarming windows and to put all the odds in our

favor, we used different chemical ecology methods, always with living swarming mosquitoes during their natural

swarming windows. First, we investigated the long-range behavioral response of females exposed to the volatile

blend from male swarms in an olfactometer. Second, we collected and analyzed volatile organic compounds

(VOCs) with different methods on both laboratory-induced swarms and natural swarms. And third, we tested for

an antenna-electrophysiological response of females to male swarm VOCs. As much as possible, we used both

recently colonized mosquitoes and large experimental set-ups to ensure males produced a free swarming behavior.

In addition, we replicated the main experiment by Mozūraitis et al. (2020) to search for the five compounds. We

specifically added two new controls, one to check if these compounds were male specific, and the other to discard

a potential environmental/laboratory pollution.

119   METHODS AND MATERIALS

Mosquitoes

Mosquitoes used in behavioral experiments and for both VOC collection and electrophysiological analyses were

from colonies raised from wild gravid females collected in inhabited human dwellings in Burkina Faso (West

Africa). *Anopheles coluzzii* were collected in 2017 in Bama and *An. gambiae* in 2015 in Soumousso. Bama is a

village located in a rice-growing area located 30 km North of Bobo-Dioulasso (11°24'14"N; 04°24'42"W) where

previous studies showed that the *Anopheles* population is almost exclusively composed of *An. coluzzii* (Mosqueira

et al. 2015; Poda et al. 2018). Soumousso is a typical Guinean savannah village located 30 km North-East of Bobo-

Dioulasso (11°00'46"N, 4°02'45"W) where *An. coluzzii* and *An. gambiae* coexist, with a predominance of the

latter (Diabaté et al. 2004, 2006). Gravid females were placed individually in oviposition cups containing tap

water. After oviposition, females were identified to species by routine PCR-RFLP (Santolamazza et al. 2008). The

larvae were gathered according to their species and reared in tap water, fed with Tetramin® Baby Fish Food

(Tetrawerke, Melle, Germany) *ad libitum*. Adult mosquitoes were held in $30 \times 30 \times 30$ cm mesh-covered cages

and provided with a 5% glucose solution *ad libitum*. Insectarium conditions were maintained at 27 ± 2 °C, $70\pm10\%$

RH and 12L:12D. The colonies were refreshed twice a year with F1 from mosquito females caught in the wild.

Mosquitoes used in the electrophysiological study were transferred as eggs from Burkina Faso to France.

On their arrival, eggs were allowed to hatch in osmosed water and the larvae were fed with Tetramin® Baby Fish

Food *ad libitum*. Adult mosquitoes were held in $20 \times 20 \times 20$ cm mesh-covered cages and provided with honey

diluted at 5% *ad libitum*. Females were fed with rabbit blood on a PS6 Power Unit (Hemotek, Blackburn, UK) for

egg production. Mosquitoes were reared in a laboratory climate chamber KBF-S720 (BINDER Gmbh, Tuttlingen,

Germany) at 27 ± 2 °C, $70\pm10\%$ RH and 12L:12D with a sunset time programmed at 3:00pm (synchronization of

swarming/mating time and electrophysiological test time to ensure an optimal receptivity of females). Mosquitoes

used for the experiments were sexed early after emergence and sexes were kept in separate rearing cages to prevent

mating.

Mosquitoes used to replicate the experiment by Mozūraitis et al. (2020) were from a 15-year-old colony

of *An. gambiae* (Kisumu strain) reared in the IRD laboratory of Montpellier. They were maintained at 27 ± 2 °C,

$80 \pm 10\%$ RH, with a photoperiod cycle of 12h light: 12h dark and reared as described above.

Long-range behavioral response of virgin females to swarm volatile organic compounds

Olfactometer setup - Bioassays were conducted in a dual-port olfactometer originally designed to study host

preference in the *An. gambiae* complex (Lefèvre et al. 2009, 2010; Vantaux et al. 2015; Nguyen et al. 2017). The

odor source container was modified to enclose male swarms in boxes made of transparent plexiglass (L × W × H:

$60 \times 60 \times 120$ cm; "swarming boxes" hereafter). Each swarming box was connected by a PVC air vent hoses (L ×

Ø: 600×10 cm, W3-65014-HQ4, HQ, USA) to a collecting glass box (L × W × H: $32 \times 33.5 \times 44$ cm) which was

linked to a mesh-covered releasing cage (L × W × H: $50 \times 40 \times 40$ cm) by a glass tube (L × Ø: 60×10 cm) (Fig.

1). A custom-made electric fan was located at the mid-length of each air vent hose and drew air from the swarming

boxes (odor-sources) to the releasing cage, providing an odor-laden air current against which mosquitoes in the

releasing cage were induced to fly. The air flow was controlled thanks to two mechanisms; first, a power regulator

(HQ-Power PS1502A, Velleman, Gavère, Belgium) connected to fans; and second, iris dampers (10 cm Ø; CIR

D100, France air, France) connecting the air vent hoses to the collecting boxes. The openings of the air vent hoses

on both the swarming and collecting boxes sides were covered with nets to prevent mosquitoes from flying into

the air vent hoses. The swarming boxes (odor-sources) were located side-by-side outdoors and the olfactometer

inside a room (Fig. 1). The air speed in the releasing cage was regulated at 18 ± 2 cm.s^{-1} using an anemometer

(Model 425, Testo, Forbach, France) and the room temperature and relative humidity were set at 27 ± 2 °C and

$80\pm10\%$ RH thanks to an air conditioner and a humidifier (Defensor 505, Condair, Croissy-Beaubourg, France),

respectively.

Bioassays

Test preparation: In the morning, about 8 hours before the time of the test, 100 2- to 3-day-old virgin

females of *An. coluzzii* and *An. gambiae* were colored with two different colored powders (Luminous Powder Kit,

BioQuip, Rancho Dominguez, California, USA) according to the species. In addition, about 500 4- to 5-day-old

An. coluzzii or *An. gambiae* males were released into a swarming box to allow them to acclimate. Both males and

females were provided with a 10% glucose solution and were kept under insectary conditions until the time of the

test. To avoid biases during the test due to both humidity and odor as a result of manipulation or bacterial

proliferation, the empty swarming box (control box) was also provided with glucose.

Test execution: Bioassays were carried out at sunset, when males naturally swarm. About 30 min before

sunset time, the swarming boxes were moved outside, the cups containing the glucose pads were removed and

visual markers made of a 20×20 cm black cloth were placed either under or next to the boxes for *An. coluzzii* or

An. gambiae, respectively (see Poda et al. 2019). These markers allowed to trigger swarming behavior and to

stabilize swarms in the middle of the box. For the same purpose, we provided an artificial twilight horizon made

of a 40 W incandescent bulb (2,500 K) located on the floor between a white wall and a 50 cm high black horizon

(Facchinelli et al. 2015; Niang et al. 2019; Poda et al. 2019). Inside, about 10 min before introducing females, the

fans were switched on in order to purge the air vent hoses and the air flow was set at 18 ± 2 cm.s^{-1}. The glass tube

openings on the releasing cage side were covered with nets to prevent female mosquitoes from flying into the

collecting boxes before the test began. The complete olfactometer, except the end of the releasing cage, was

covered with a white cloth to eliminate visual bias during the test and to provide a diffused light. Sunset light was

allowed to enter the olfactometer room until dark and an additional artificial twilight made of a 40 W incandescent

bulb (2,500K) projected on the wall of the room on the side of the collecting boxes was also provided. Thirty

minutes before sunset, about 100 females of each species (N=200, 1 replicate) were released simultaneously into

the releasing cage of the olfactometer allowing them to acclimate.

When the males started to swarm (*i.e.* flying in loops at a constant location within the swarming boxes,

see Downes 1969; Poda et al. 2019), the nets covering the glass tube openings and preventing access of females

to the collecting boxes were removed thanks to threads that were attached to the nets. The females were allowed

to respond for 20 min (swarm duration), then the glass tube openings were covered again with nets and females

that reached the upwind collecting boxes were removed with a mouth aspirator, killed by freezing and identified

according to their coloration (species) under a binocular (LEICA S6E) and counted. The remaining mosquitoes

inside the releasing cage were also removed. After each test (1 per day), the olfactometer was cleaned with 95%

ethanol to remove odor contaminants. All materials were handled with gloves to avoid contamination with skin

compounds.

Test combinations: Four combinations of choice tests were performed: i) *An. coluzzii* male swarm *vs.*

empty box (hereafter, *An. coluzzii* test); ii) *An. gambiae* male swarm *vs.* empty box (hereafter, *An. gambiae* test),

iii) *An. coluzzii vs. An. gambiae* male swarms (hereafter, *An. coluzzii vs. An. coluzzii* test) and iv) empty *vs.* empty

box (hereafter, control test). We ran 12 replicates per combination and females were used only once. Males were

used twice in two consecutive days. In that case, a 10% glucose solution was introduced into the two swarming

boxes after the test and boxes were kept under insectarium conditions until the next day. The swarming boxes were

cleaned with ethanol between each batch of males. To avoid biases, the matching of species and colors was

switched between each test, and we alternated the treatments (mosquitoes or control) between the swarming boxes

and the right and left arm of the olfactometer. We assessed the instrumental and arm bias through an empty *vs.*

empty box choice test (control test). The olfactometer did not present any symmetrical biases (Fig. S1).

Collection of swarm volatile organic compounds (VOCs)

Because compounds can be emitted at a given time and only produced in tiny quantities at individual scale, we chose to work directly on swarms. Three different sampling methods were used; two methods were run in a laboratory setup (Niang et al. 2019) and one in the field.

Dynamic headspace collection in the laboratory

VOC collection setup: The collection of swarm VOCs was performed in a room specially designed to stimulate *Anopheles* swarming behavior thanks to a set of visual stimuli (see Niang et al. (2019) for details). The headspace volume consisted of a $50 \times 50 \times 50$ cm transparent plexiglass box (extraction box). Such a volume was a trade-off between a volume large enough to allow males to display a swarming behavior without too much constraints and a volume small enough to be sucked up in a reasonable time. Entrance and exit flow rates were maintained by two pumps and regulated by flowmeters. The entrance flow was higher than the exit flow ensuring that the system was continuously purged to compensate for the inevitable leaks, and that no contaminated outside air would enter the system. At the entrance side, ambient air was purified in a glass charcoal filter and then humidified passing through deionized water. Exit flow passed through an odor trap to adsorb VOCs. Three different odor traps were used for the chemical analyses and electrophysiological analyses. Tenax-TA/Carbotrap sorbent stainless steel 6mm diameter tubes (Gerstell, Mülheim, Germany) and Porapak-Q sorbent VCT glass tubes (ARS, Gainesville, Florida, USA) were used for chemical analyses. Porapak-Q sorbent VCT tubes and home-made sorbent Micro-traps were used for electrophysiological analysis. Micro-traps were constituted by ChromatoProbe® quartz microvials of Varian Inc. (length: 15 mm; inner diameter: 2 mm), previously cut closed-end and filled with 3 mg of a 1:1 mix of Tenax-TA and Carbotrap® (60–80 and 20–40 mesh, respectively; Sigma Aldrich, Munich, Germany)(See Dormont et al. 2013b). Flow rates and VOC collection durations were dependent on the odor trap (See Table 1).

VOC collection: About 500 4- to 6-day-old males of *An. gambiae* or *An. coluzzii* were transferred into the extraction box about 30 min before swarming time. Clean air was pushed into the box at a rate of 10 L.min^{-1} to purge the box from both human and mosquito container odors before connecting it to the odor trap. Five minutes after males started to swarm (*i.e.* flying in loops at a constant location within the box), the air flow was adjusted at the required rate and the odor trap connected. Forty minutes later, swarming behavior was stopped by turning off the light and the VOC collection continued until the required collection time was reached (see Table 1). After

VOC collection, samples were stored at 4 °C until analysis. After males were removed, the box was cleaned with

95% ethanol and then flushed with clean air for at least 5 hours to remove the odor contaminants left from the

previous extraction. A total of 30 swarm extracts were collected; 17 *An. coluzzii* swarms (5, 8 and 4 with

Tenax/Carbotrap tubes, Porapak-Q VCT tubes and Micro-traps, respectively) and 13 *An. gambiae* swarms (6 and

7 with Tenax/Carbotrap tubes, and Porapak-Q VCT tubes, respectively). Control consisted of a clean empty box.

Before they were sent and used in Burkina Faso, adsorbents were cleaned as follows. Tenax-TA/Carbotrap tubes

were heated at 250 °C for 30 min under a 30 mL.min^{-1} flow then sealed. Porapak-Q VCT tubes were eluted with 3

mL of hexane then packed individually in a nalophan bag. Chromatoprobes were heated for 2 h at 270 °C by 100

chromatoprobes under a 100 mL.min^{-1} nitrogen flow, then packed in a glass vial with a Teflon cap.

VOC collection from natural swarms - VOCs were collected from a natural swarm of *An. coluzzii* in the village

of Bama. Static sorptive collections of volatiles from a non-enclosed swarm were performed with Twisters® (100

µm PDMS stir bar; Gerstel, Mülheim, Germany; Bicchi et al. 2000; Tienpont et al. 2000). The swarm flew about

2.5 to 3.5 m above the ground. Thus, we used a 1.5 m long glass stick (Glaswarenfabrik Karl Hecht GmbH & Co,

Germany) with a metallic push pin head glued at the tip and covered with nalophan to introduce the magnetic

twisters directly into the swarm. Control twisters were placed about 3 m away upwind from the swarm. VOC

collection lasted for the whole swarm duration (about 20-25 min). We chose a swarm far from habitations and

livestock to limit odor pollutions, containing a large number of males (more than 1,000) and attractive for females

(observation of a large number of couples). After VOC collections, twisters were individually placed into glass

vials and stored at 4 °C until analysis. Five replicates were performed on the same swarm on different days.

Solvent extraction - About 1,000 4- to 6-day-old males of *An. gambiae* or *An. coluzzii* were introduced into a 30

$\times 30 \times 30$ cm cage and kept under insectarium conditions. The cage was made of unpainted metallic frames covered

by white nets and the bottom was covered with a white paper. Thirty minutes before sunset time, the cage was

placed outside and mosquitoes were observed. A 5×5 cm black cloth was placed in or out of the cage according

to the species as described above to stimulate swarming behavior. About 5 min after mosquitoes started to swarm,

the entire cage was quickly placed into a freezer at -20° C for about 10 min. Then, mosquitoes were transferred

into a 20 ml glass tube and covered with a 1:1 mix of Hexane and Dichloromethane. The tube was sealed and

mosquitoes were kept in the solution for 24h at 27±2 °C. Then, the solution was filtered on glass wool to remove

mosquito scales and stored at 4° C until analysis. Three different controls were made: i) with 1- to 2-day-old virgin

males (young male control), ii) with 2- to 3-day-old virgin females (female control) and iii) with solvent mix alone

(blank control). A total of 8 swarm extracts were done; 4 of *An. coluzzii* and 4 of *An. gambiae* and 1 replicate of

each type of control per species.

VOC collection with SPME fiber – We replicated the experiment from the study by Mozūraitis et al. (2020) in

which they collected mosquito headspace with a SPME fiber in a 1L bottle. However, because most methods used

to introduce mosquitoes into a bottle may potentially result in entering pollutants as well, we added a

supplementary control. One way to easily introduce the mosquitoes is to blow them from a mouth aspirator into

the bottle. This classic method may bring breath volatiles within the recipient, which can contain the five

compounds described in Mozūraitis et al. (2020) (Poli et al. 2010; Calenic and Amann 2014; Filipiak et al. 2014;

Cainap et al. 2020). Consequently, we added a "breath control". In addition, we also replicated the experiment

with females to check for sex specificity. In the morning, the two glass bottles were cleaned with acetone and dried

under a clean air flow, then closed with nalophan. Three hours before swarming time (5pm), we collected the

headspace of empty bottles with polydimethylsiloxane/divinylbenzene-coated SPME fibers for 1h (step 1: empty

bottle). Then, we simulated the introduction of mosquitoes by blowing 3 times for 3 seconds with a mouth aspirator

into the bottles. These were then closed with Nalophan (6pm) and the head space of the breath of the experimenter

was collected for 1h with SPME fibers (step 2: breath control). We then introduced 60-70, 5- to 7-day-old unmated

males in each bottle with the mouth aspirator, closed the bottles with Nalophan (7pm) and let the mosquitoes

acclimate for 1h. During the acclimation period, we simulated a light transition from photophase to scotophase to

trigger crepuscular flight. Then, when males started to fly (8pm), SPME fibers were introduced for 1h (step 3:

"swarm" extract). Six different fibers were used and cleaned at 250 °C during 5 min in the injector of a gas-

chromatograph (HP 6890 Series PLUS, Agilent, Santa Clara, USA) just before use. Each fiber was always used

for the same type of extract and in the same bottle. Before the experiment, we made sure that the fibers had similar

sensitivity by exposing them to a mix of the five compounds. Only minor quantitative differences were detected,

and this was considered during subsequent statistics. Secondly, we performed the same 3 steps but with 60-70, 3-

to 4-day-old virgin females instead of males. And finally, as during the introduction of mosquitoes, we blew a

second time into the bottles and repeated the three steps of headspace collection but this time, instead of introducing

mosquitoes in the 3rd step, we blew again into the bottles (breath ×2) to check for potential accumulation of breath

compounds. We kept track of the day of the experiment, the bottles and fibers identity for statistical analyses.

Chemical analyzes

Samples obtained on Porapak-Q VCT tubes were eluted with 150 µl of Hexane and injected (1 µl) into a Gas Chromatograph coupled with Mass Spectrometer (GC-MS). Those obtained on Tenax/Carbotrap tubes, twisters or SPME fibers were directly thermo-desorbed into GC-MS. Solvent extracts were injected as is or concentrated under a gentle stream of nitrogen. All the analyses were run at the "Platform for Chemical Analyses in Ecology" (PACE), the technical facilities of the LabEx CeMEB (Centre Méditerranéen pour l'Environnement et la Biodiversité, Montpellier, France).

Liquid analysis (Porapak elutions and solvent extractions) - Liquids were analysed with a GC-MS QP2010 Plus (Shimadzu, Kyoto, Japan). Mass spectra were recorded in electronic impact mode (EI) at 70eV over a m/z mass range from 38 to 350. The temperature of the transfer line and the ion source were programmed to 250 °C and 200 °C, respectively. The injections were made with an injector temperature of 250 °C, and a 1:4 split mode ratio. Analyses were performed using a 30 m × 0.25 mm × 0.25 µm Optima 5-MS (Macherey-Nagel, Düren, Germany) fused silica capillary column with a constant helium flow set close to 1 ml.min^{-1}. The oven temperature was programmed as follows: 40 °C (held 5 min) to 250 °C at 6 °C.min^{-1}, and finally to 300 °C at 14 °C.min^{-1} and held 2 min. GC-MS Solution software (Shimadzu, Kyoto, Japan) was used for data processing of these analyses, with the NIST 2011 as spectrum database.

Tenax/Carbotrap tubes and twister analysis - These sorbents were analysed as described by Souto-Vilarós et al. (2018). Samples were analysed using a gas chromatograph (GC, Trace™ 1310, Thermo Scientific™ Milan, Italy) coupled with a mass spectrometer (MS, ISQ™ QD Single Quadrupole, Thermo Scientific™ Milan, Italy). The column used was an Optima 5-MS, the same as for liquid analysis. Absorbent traps were handled with a Multi Purpose Sampler (Gerstell, Mülheim, Germany) and desorbed with a double stage desorption system, composed of a Thermal Desorption Unit (TDU) and a Cold Injection System (CIS) (Gerstell, Mülheim, Germany). The sorbents were desorbed in splitless mode at 250 °C on a trap cooled at −80 °C by liquid nitrogen. Then, the trap was heated to 250 °C with a 1:4 split ratio to inject the compounds in the column. The carrier gas used was helium at 1 ml.min^{-1}. Oven temperature program was as follows: held at 40 °C for 3 min., then increased to 220 °C at 5 °C.min^{-1} and finally to 250 °C at 10 °C.min^{-1}, and held for 2 min. Mass spectra were recorded in electronic impact mode (EI) at 70eV over a m/z mass range from 38 to 350. The temperature of the transfer line and the ion source

were programmed to 250 °C and 200 °C, respectively. Xcalibur™ software (Thermo Scientific™, Milan, Italy)

was used for data processing and the NIST 2011 as spectrum database.

SPME fiber analysis – We analyzed SPME fibers as described in Mozūraitis et al. (2020) with a GC-MS QP2010-

SE (Shimadzu, Kyoto, Japan). Volatiles were desorbed into the injector at 225 °C for 1 min in split mode (1:4

ratio). Analyses were performed using a 30 m × 0.25 mm × 0.25 µm J&W DB-Wax silica capillary column

(Agilent, Santa Clara, USA) with a constant helium flow at 1 ml.min^{-1}. The oven temperature was programmed as

follows: 40 °C (held 1 min.) to 150 °C at 5 °C.min^{-1}, and finally to 220 °C at 20 °C.min^{-1} and held 9 min. Mass

spectra were recorded in electronic impact mode (EI) at 70eV over a m/z mass range from 38 to 350. The

temperature of the transfer line and the ion source were 250 °C and 200 °C, respectively. GC-MS Solution software

(Shimadzu, Kyoto, Japan) was used for data processing of these analyses.

Electrophysiological analysis

Electrophysiological activity of female olfactive detection, on the male swarm extracts, was tested with an Electro-

Antennografic Detection system, coupled with Gas Chromatography (GC-EAD) at the PACE. Thirty minutes

before the mosquito scotophase, both 4- to 6-day-old males and 2- to 3-day-old virgin females were placed

separately in cardboard cups (Ø = 75 mm, h = 100 mm), provided with a 5% honey solution and transferred from

the insectarium to the electrophysiology laboratory at PACE. At mating time, mosquitoes were placed in the dark.

Males were checked for erected antennae as a proxy to ensure that change in the light-dark cycle, transport and

electrophysiology laboratory conditions did not alter mosquito motivation for mating. Two minutes after male

antennae were erected, a female was removed from the cardboard for the antennal preparation.

The female's head was gently excised with a pair of surgical scissors and mounted between two glass

capillary tubes of 76 mm in length and 1.12 mm in diameter (World Precision Instrument, Sarasota, USA), pulled

and cut using a vertical micropipette-puller (P-30 model, World Precision Instruments, Hertfordshire, United

Kingdom) filled with insect Ringer solution (6.0 g.l^{-1} NaCl, 0.4 g.l^{-1} KCl, 0.27 g.l^{-1} CaCl$_2$ and 3.20 g.l^{-1} NaC$_3$H$_5$O$_3$)

and connected to silver wires. The indifferent electrode was inserted into the back (the foramen) of the isolated

head and the intact tip of one antenna was inserted into the recording electrode which was connected to an Electro-

Antennography Detector setup (EAD, Syntech IDAC-2, Kirchzarten, Germany). One end of the glass electrodes

was pulled to a fine point which was then broken to ensure a close fit with the female's head and antenna.

The above EAD setup was linked to a Gas Chromatograph-Flame Ionization Detector (GC-FID, CP-3800,

Varian, Palo Alto, USA) equipped with an Optima 5-MS capillary column, the same as for the GC-MS analyses.

Liquid injections after Porapack solvent elution were made in a 1079 PTV injector (Programmed Temperature

Vaporizator) at 250 °C with a 1:4 split ratio. Microtraps were desorbed in the same injector using a ChromatoProbe

sample introduction device (Varian, Palo Alto, USA) with the same split ratio and temperature programmed as

follows: increased from 40 °C to 250 °C at 200 °C.min^{-1} and held for 5 min. Oven temperature was held at 50 °C

for 0.40 min, then increased to 180 °C at 10 °C.min^{-1}, and finally to 220 °C at 12 °C.min^{-1}, for a run duration less

than 17 min. The carrier gas used was helium at 1 ml.min^{-1}. The effluent was split equally into two deactivated

fused silica capillary columns (100 cm x 0.25 mm), one leading to the FID (270 °C) and the other into a heated

EAD port (200 °C) (transfer line: Syntech, Kirchzarten, Germany) and leaded to antenna. The GC effluent for

antenna was mixed with charcoal-filtered, humidified airflow (300 ml.min^{-1}). StarWorkstation 6.41 software

(Varian, Palo Alto, USA) was used for data processing.

Different types of male swarm extracts were tested on the antennae of several conspecific females.

Antennae of 19 *An. coluzzii* females were tested with solvent extracts, Porapak-Q VCT tube extracts and Micro-

trap extracts (8, 9 and 2 females, respectively) and 20 *An. gambiae* females were tested with solvent and Porapak-

Q VCT tube extracts (10 females each). Before GC-EAD analysis of the VOC extracts, we checked the living

antenna for good activity using a negative (solvent) and a positive (mix of many VOCs) stimulus solutions. We

proceeded as follows: 5µl of each solution was applied to a 1 cm × 2 cm filter paper contained in a Pasteur pipette.

The pipette was then placed in an air pump system and the volatiles were directly sent to the antennal preparation

with a pulse duration of 0.5 s and a flow rate of 890 ml.min^{-1} regulated by a CS-55 Stimulus Controller (Syntech,

Hilversum, Netherlands).

379   Identification of specific and active VOCs

For all the chemical analysis, we compared the chromatograms of swarming mosquito extracts with those

of their corresponding controls to search for any qualitative or quantitative differences. When such a difference

was pinpointed, we matched the mass spectra of the compounds of interest with the NIST 2011 MS library. A

solution of n-alkanes (Alkanes standard solution, 04070, Sigma Aldrich, Darmstadt, Germany) was also injected

to calculate the linear retention index (LRI) of these compounds of interest (Zellner et al. 2008) and their LRIs

were compared with those reported in the literature.

In the electrophysiological analysis, we studied the chromatograms in a range of LRI between 800 and

1700, including compounds present at very low and trace amounts. When a depolarization was observed in the

electrophysiological analysis, a section of 20 points of LRI was studied around the compound's LRI on the GC-

MS trace. Comparison of the samples with their respective controls collected on the same day, allowed to subtract

potential contaminants from the samples. Non-natural compounds, such as industrial chemicals and/or compounds

not naturally produced by living organisms (Charpentier et al. 2012), were considered as contaminants.

We also particularly searched for the presence of acetoin, sulcatone, octanal, nonanal and decanal reported

by Mozūraitis et al. (2020). With that aim, we defined a LRI window of 20 points on both sides of the LRI of these

compounds on the bases of injected standards (for SPME fibers; acetoin, sulcatone, octanal, nonanal and decanal,

≥98%, 99%, 99%, ≥98% and ≥98% GC, respectively, Sigma-Aldrich, Darmstadt, Germany) or literature (for other

extraction methods). Then, we searched for their specific ions at those LRI as follows: (LRI/ion) 711/88, 981/108,

998/84, 1100/98, and 1201/112 for acetoin, sulcatone, octanal, nonanal and decanal, respectively. Then, when a

peak corresponding to both LRI and specific ion was found, we compared its spectra with the NIST 2011 MS

library before quantification with integration units.

Statistical analysis

Both the activation rate and choice were analyzed using binomial Generalized Linear-Mixed Models

(GLMM, "*glmer*" function in "*lme4*" package). Tested combinations (four levels: *An. coluzzii*, *An. gambiae*, *An.*

coluzzii vs. An. gambiae and control tests), female species (two levels: *An. coluzzii* and *An. gambiae*) and their

interaction were considered as fixed effects, and replicates as random effects. All analyses were performed using

R (version 3.4.0).

We analyzed separately the quantities using integration units of acetoin, sulcatone, octanal, nonanal and

decanal. When pairing between the controls and the mosquito extracts was possible (with SPME and twister) and

to account for repeated measurements, we used Gaussian GLMMs. In those cases, the extract (3 levels for SPME

fibers: empty bottles, breath and flying mosquitoes (either male or females or breath ×2) and 2 levels for twisters:

control and swarm) was considered as fixed effect. The replicates (i.e. the day for twisters and the SPME) and

fiber nested within the bottle for SPME fibers were considered random effects. Considering the SPME fibers as a

random effect made it possible to account for their minor differences of sensitivity in the model. When pairing

between the controls and the mosquito extracts was not possible (Tenax-TA/Carbotrap and Porapak-Q VCT tubes),

Gaussian GLMs were used. The extract (empty box, *An. coluzzii* swarm or *An. gambiae* swarm) was considered

as fixed effect. In some occasions, some compounds were not found in the extracts or only in two samples.

Consequently, it was not possible to perform statistical analyses and we considered the extracts as not significantly

different. No statistical analysis was performed with liquid extracts due to a small number of replicates. For model

selection, we used the stepwise removal of terms, followed by likelihood ratio tests. Term removals that

significantly reduced explanatory power ($P <0.05$) were retained in the minimal adequate model. All percentages

are provided with their 95% confident interval.

423   RESULTS

Behavioral assays

Over 12 replicates of c.a. 200 females for each of the four combinations, a total of 8,918 females were

tested (4,423 *An. coluzzii* and 4,495 *An. gambiae*) among which 3,591 flew upwind into the collecting boxes

(activation rate: 40.2±1%). There was a significant interaction between female species and the four combinations

of swarm boxes tested ($\chi^2_3 = 22.19$, $P <0.001$). Species subset analyses showed that the different test combinations

had no effect on the *An. coluzzii* activation rate ($\chi^2_3 = 5.72$, $P=0.12$; Fig. 2). However, in *An. gambiae*, there was

a significant effect of the test combinations on the female activation rate ($\chi^2_2 = 14.94$, $P = 0.001$; Fig. 2) with a

higher activation rate when females were exposed simultaneously to both *An. coluzzii* and *An. gambiae* male

swarms than when exposed to an *An. coluzzii* swarm alone ($z = 4.01$, $P <0.001$) or to the control (two empty boxes:

$z = 2.57$, $P = 0.04$). Nonetheless, no difference was found when females were exposed to an *An. gambiae* male

swarm alone ($z = -1.49$, $P = 0.44$). Neither the *An. coluzzii* swarms, nor the *An. gambiae* swarms were attractive

for activated females (female species: $\chi^2_1 = 2.45$, $P = 0.11$; choice test combinations: $\chi^2_2 = 1.15$, $P = 0.56$ and

female species × choice test combination interaction: $\chi^2_2 = 0.34$, $P = 0.84$; Fig. 3).

Chemical analyses

No qualitative or quantitative differences were detected between samples of swarming males and their

relative controls for both species obtained from the Porapak-Q, Twisters® and the liquid extracts (Figs S2 & S3).

Minor quantitative differences were detected with the Tenax/Carbotrap tubes. However, mass spectra showed that

these peaks were pollutants, such as silicones, BTX (mono-aromatic compounds like benzene, toluene and xylene),

alkanes which were all emitted by the box containing the mosquitoes. Quantitative differences including linear

organic acids were also detected, but they were not reproducible.

During the experiment replicating the protocol by Mozūraitis et al. (2020), both males and females started

to fly at the expected time (8pm) but due to the small volume it was not possible to determine if the group was a

swarm. Analyses of SPME fibers in search for acetoin, sulcatone, octanal, nonanal and decanal showed that acetoin

was absent from almost all the male samples (Fig. 4A), but was present on three occurrences in female extracts

(over six) (Fig. 4B). Sulcatone, octanal, nonanal and decanal were found in almost all types of extracts including

controls. However, we did not find significant increased quantities of the five compounds in "swarming" mosquito

extracts compared to the breath controls (Fig. 4). The control to test the effect of blowing twice in the bottle showed

that the headspaces of "breath" and "breath ×2" were very similar (Fig. S4), demonstrating an absence of

accumulation of breath odor between the breath control and the introduction of mosquitoes.

A posteriori, we also quantified acetoin, sulcatone, octanal, nonanal and decanal in our previous GC-MS

analyses in search for similar variations as described in Mozūraitis et al. (2020). We did not find acetoin in our

chromatograms. Indeed, as we used a low polarity Optima 5-MS column, acetoin (if present) was eluted before

the LRI 800. In the samples obtained from natural swarms of *An. coluzzii* with the twister method, sulcatone and

octanal were absent. The quantities of both nonanal and decanal were not significantly higher in the swarm than

in the control (Fig. 5A). In laboratory swarm samples, sulcatone, octanal, nonanal and decanal were found

inconsistently. Nevertheless, swarms did not show higher quantities of these compounds compared to the empty

box (Fig. 5B) excepted in samples obtained with the Porapack tubes (Fig. 5C). However, this difference was not

statistically significant. Whatever, the method used, we can note a high quantitative variability between the

samples. This was true for both the control and the mosquito samples.

Electrophysiology analyses

The male swarm VOCs induced a few weak and poorly reproducible depolarizations on the antennae of conspecific

females. In *An. coluzzii*, 3 females out of 9 weakly responded to a single Porapak-Q extract at the linear retention

index (LRI) 1371 (retention time (RT) 12.85 min), and 2 females out of 8 responded to solvent extracts at LRI

921, 978 and 1179 (RT 6.40, 7.24 and 10.23 min, respectively, Fig. 6). However, these responses did not

correspond to noticeable peaks in the GC detector of the GC-EAD apparatus. Successive GC-MS analyses of the

same porapak-Q and liquid extracts did not allow to detect and identify any specific peak in relation to antennae

responses (Figs S2b & S3b). The peaks found at LRIs that elicited a response in *An. coluzzii* females were identified

as environmental pollutants and were present in both swarming mosquitoes and control samples. No reproducible

response was recorded from *An. coluzzii* females tested with Micro-traps extracts. In *An. gambiae*, no reproducible

antennae response was recorded from the 10 females tested with either Porapak-Q or liquid extracts.

477   DISCUSSION

For the past decades, several authors have suggested the existence of long-range sex pheromones in the attraction

of *An. gambiae s.l.* females to male swarms (Charlwood et al. 2002; Tripet et al. 2004; Diabaté et al. 2011; Poda

et al. 2019) but to date, no reliable experimental evidence is available. Here, we used behavioral, physiological

and detailed chemical approaches to attempt to highlight the existence of such compounds. However, despite a

large set of methodologies and working hypotheses to maximize the chances of highlighting the presence of

pheromones (*i.e.* timing, physiological state, natural behavior, recently colonized mosquitoes), our study failed to

provide evidence of the presence of long-range male sex pheromones in both *An. coluzzii* and *An. gambiae*. We

also replicated an experimental setup that recently showed an increase in quantity of five compounds in

"swarming" males compared to controls. However, unlike in Mozūraitis et al. (2020), we did not find any

significant difference with our controls.

During our behavioral tests in the olfactometer, females were not attracted to air currents passed over

male swarms regardless of the species. During the experiment, the males released in the large plexiglass boxes and

exposed to natural light started to fly randomly at sunset. Then, about 150-200 males out of the 500 released

(<50%) gathered in the upper half of the box flying in loops without touching the box walls. They also reacted to

the movements of the visual marker, meaning they were indeed swarms. The other males flew randomly in the

box, bouncing against the walls. Females showed a good activation rate but they did not show a particular choice

for any arm, containing a swarm or not. Nevertheless, *An. gambiae* females showed an intriguing higher activation

rate when exposed simultaneously to both a conspecific and a heterospecific swarm compared to a single *An.*

coluzzii swarm. This was, however, probably without biological significance as no difference was detected in the

test providing *An. gambiae* females with a choice between the control and *An. gambiae* swarm, *vs.* a test offering

two controls.

Our behavioral result was consistent with our physiological study. Indeed, despite the fact that we checked

for both behavioral receptivity of mosquitoes and for receptivity of mounted antennae, no consistent antennal

depolarization was observed in females across assays when exposed to swarm extracts. This suggests a lack of

response by females to any volatile chemicals present in our swarm extracts.

Chemical analyses were also negative and no compounds specific for male swarms could be detected

whatever the method used. Moreover, unlike in Mozūraitis et al. (2020), the quantities of acetoin, sulcatone,

octanal, nonanal and decanal were found inconsistently across swarming mosquito samples and not in larger

quantities compared to controls, making it difficult to support the assumption according to which they could

actually be emitted by males. These divergent results could be explained by the fact that, unlike in the experiments

of Mozūraitis et al. (2020), we used a control which considered the potential introduction of pollutions at the same

time as mosquitoes. This showed that breath was responsible for most of the variability. The most convincing

result showing that sulcatone, octanal, nonanal and decanal are probably not pheromones emitted by males is the

one obtained with the twister method. Indeed, twisters are coated with the same adsorbent phases as SPME fibers

used both in this study and in the one by Mozūraitis et al. (2020). According to the manufacturer, twisters are up

to a thousand times more sensitive than SPME fibers partly due to a larger sorbent volume. Moreover, we exposed

the twisters directly in a natural swarm formed by more than 1 000 males (probably up to 6 000 males according

to the estimation of trained technical staffs). Despite this sensitivity, the number of mosquitoes and the natural

biological context, sulcatone and octanal were not found and both nonanal and decanal had similar quantities in

the swarm and outside the swarm (control twister placed 3 m upwind from the swarm). In addition, the high

quantitative variability found in the laboratory and the fact that these compounds are frequently found in controls

suggest that they could be laboratory and/or human pollutions that are difficult to control for.

Our results contrast with those of Mozūraitis et al. (2020) probably for several reasons. First, in their

laboratory experiments, they reported a simple flight activity instead of a swarming activity in which males should

fly in regular loops with erected antennae (Downes 1969; Poda et al. 2019). This can be explained by the absence

of adequate visual stimuli such as ground markers which are mandatory to trigger swarming behavior (Charlwood

and Jones 1980; Marchand 1984; Gibson 1985; Facchinelli et al. 2015; Niang et al. 2019; Poda et al. 2019). Second,

this flight activity was extraordinary long for a swarming flight, exceeding 200 min in the presence of the five

compounds. Indeed, swarming behavior is known to last for only 20-30 min in nature (Charlwood and Jones 1980;

Marchand 1984; Sawadogo et al. 2013, 2014; Bimbilé Somda et al. 2018) and only up to 60 min in artificial

conditions (Charlwood and Jones 1980; Poda et al. 2019). In the latter, the number of males in swarms decreased

over time while the others flew randomly, bouncing on the flight arena walls (Poda et al. 2019). Indeed, swarming

is an activity with a high energy demand, consuming 50% of sugar and glycogen reserves in 25 min (Maïga et al.

2012, 2014). Mosquitoes probably switched to a more random flight to try to escape and search for a sugar meal

to refuel their reserves. This behavior can be stimulated by acetoin, sulcatone, octanal, nonanal and decanal, as

they are frequently emitted from nectar sources and fermented sugar (Goodrich et al. 2006; Schiestl 2010; Dekel

et al. 2019). Finally, they also reported that these five compounds increased insemination rates in five different species

in semi-field cages. However, instead of stopping the experiment after the swarming time, they collected the females in

the morning. As they left both a dark box in the arena as resting site and a sugar source, it is likely that males and females

also mated in or around these two resources during the night. Indeed, it was shown that mating can occur indoors, outside

of swarming times and locations (Dao et al. 2008). The boxes and sugar sources with these five compounds were

probably more attractive than the control ones gathering more mosquitoes and thus inducing more inseminations.

Independently, our methods can also have weaknesses. Even if the number of swarming males was

biologically relevant compared to natural swarms, a higher number of males could be needed in experimental

setups, or the sensitivity thresholds of our chemical and electrophysiological apparatus are below the ability of

insects to detect low amounts of compounds. In addition, although the air flow used in our VOC collection in

laboratory was not unconventional, it might have been too high inducing a leak of compounds of interest. Likewise,

the air quality could have masked some compounds emitted in minute quantities even though our behavioral

experiment and VOC collection in natural swarms led to negative results too.

However, altogether and keeping in mind that an absence of evidence is not an evidence of absence, our

results support the absence of long-range sex pheromones emitted by male swarms. However, further

investigations are needed and complementary methodologies such as electro-physiology on palps (Lu et al. 2007;

Iatrou and Biessmann 2008; Pitts et al. 2011; Guidobaldi et al. 2014), molecular analyses of olfactory protein

expression on adequate physiological status, or real-time chemical analyses of swarm volatiles with a sensitive

apparatus such as a proton-transfer-reaction mass spectrometry could be options. The eventuality that organic

compounds with low volatility might be involved in mating process other than long-range attraction cannot be

discarded.

The question of how *Anopheles* females seek male swarms is still open. The lack of post-mating barrier

(Persiani et al. 1986; Diabaté et al. 2005, 2007; Hahn et al. 2012; Pombi et al. 2017) and the low rate of hybrids in

this geographical area (della Torre et al. 2001, 2005; Tripet et al. 2001; Diabaté et al. 2006) necessarily involve a

reasonably distant process which prevents females from entering heterospecific swarms. As acoustic cues have

shown some limitations at long range (Feugère et al. 2021), and because long-range chemical cues are still

disputable, visual cues such as ground markers (Diabaté et al. 2009; Sawadogo et al. 2014; Poda et al. 2019) could

be good candidates, but their effective range and level of specificity is poorly known.

Declarations

Funding

This work was funded by a grant from the Agence Nationale de la Recherche (ANR-15-CE35-0001-01) awarded

to O.R. S.B.P. received financial support through a doctoral fellowship from the Institut de Recherche pour le

Développement (IRD).

Competing interests

We have no competing interests.

Availability of data and material

The raw datasets are available on Zenodo (https://doi.org/10.5281/zenodo.4719568).

Code availability

Not applicable.

Authors' contributions

O.R. conceived the study. O.R., S.B.P., B.B., B.L. and L.D. designed the chemical and electrophysiological

experiments. O.R. and S.B.P. performed chemical extractions and SBP and BB performed the chemical analysis.

S.B.P. and B.L. performed the electrophysiological experiments. O.R. and S.B.P. designed olfactometric

experiments and S.B.P. performed data collection. S.B.P. and O.R. performed statistical analyses. S.B.P., O.R.,

B.B., B.L. drafted the manuscript and L.D., O.G., A.D, and R.K.D critically revised the manuscript. All authors

revised the manuscript, gave final approval for publication and are accountable for the work performed therein.

Acknowledgements

We thank David Sanou for his assistance in the field, Stephane Somda, Sofan Somé, Bethsabee Scheid, Marie

Rossignol and Carole Ginibre for their help in mosquito rearing and Heidi Lançon for proofreading the paper.

591   REFERENCES

Alexander RD, Marshall DC, Cooley JR (1997) Evolutionary perspectives on insect mating. In: Choe JC, Crespi

BJ (eds) The evolution of mating systems in insects and Arachnids. Cambridge University Press, pp 4–31

Andersson J, Vongvanich N, Wiklund C, et al (2007) Male sex pheromone release and female mate choice in a

butterfly. J Exp Biol 210:964–970. https://doi.org/10.1242/jeb.02726

Bicchi C, Cordero C, Iori C, et al (2000) Headspace Sorptive Extraction (HSSE) in the headspace analysis of

aromatic and medicinal plants. J High Resolut Chromatogr 23:539–546. https://doi.org/10.1002/1521-

4168(20000901)23:9<539::AID-JHRC539>3.0.CO;2-3

Bimbilé Somda NS, Poda BS, Sawadogo PS, et al (2018) Ecology of reproduction of *Anopheles arabiensis* in an

urban area of Bobo-Dioulasso, Burkina Faso (West Africa): monthly swarming and mating frequency and

their relation to environmental factors. PLoS One 13:e0205966.

https://doi.org/10.1371/journal.pone.0205966

Bjostad LB, Gaston LK, Shorey HH (1980) Temporal pattern of sex pheromone release by female *Trichoplusia*

ni. J Insect Physiol 26:493–498. https://doi.org/10.1016/0022-1910(80)90122-5

Bray DP, Alves GB, Dorval ME, et al (2010) Synthetic sex pheromone attracts the leishmaniasis vector

Lutzomyia longipalpis to experimental chicken sheds treated with insecticide. Parasit Vectors 3:1–11.

https://doi.org/10.1186/1756-3305-3-16

Cainap C, Pop LA, Balacescu O, Cainap SS (2020) Early diagnosis and screening in lung cancer. Am J Cancer

Res 10:1993–2009

Calenic B, Amann A (2014) Detection of volatile malodorous compounds in breath: current analytical

techniques and implications in human disease. Bioanalysis 6:357–376. https://doi.org/10.4155/bio.13.306

Charlwood JD, Jones MDR (1980) Mating in the mosquito, *Anopheles gambiae s.l.* II. Swarming behaviour.

Physiol Entomol 5:315–320. https://doi.org/10.1111/j.1365-3032.1980.tb00241.x

Charlwood JD, Pinto J, Sousa CA, et al (2002) The swarming and mating behaviour of *Anopheles gambiae s.s.*

(Diptera: Culicidae) from São Tomé Island. J Vector Ecol 27:178–183

Charpentier MJE, Barthes N, Proffit M, et al (2012) Critical thinking in the chemical ecology of mammalian

communication: roadmap for future studies. Funct Ecol 26:769–774. https://doi.org/10.1111/j.1365-

2435.2012.01998.x

Clements AN (1999) The biology of mosquitoes. Volume 2: sensory, reception and behaviour. CABI Publishing,

Wallingford, UK

Clements AN (1992) The biology of mosquitoes: development, nutrition and reproduction. Chapman & Hall,

London

Costantini C, Ayala D, Guelbeogo MW, et al (2009) Living at the edge: biogeographic patterns of habitat

segregation conform to speciation by niche expansion in *Anopheles gambiae*. BMC Ecol 9:16.

https://doi.org/10.1186/1472-6785-9-16

Dao A, Adamou A, Yaro AS, et al (2008) Assessment of alternative mating strategies in *Anopheles gambiae*:

does mating occur indoors? J Med Entomol 45:643–52

Dekel A, Yakir E, Bohbot JD (2019) The sulcatone receptor of the strict nectar-feeding mosquito *Toxorhynchites*

amboinensis. Insect Biochem Mol Biol 111:103174. https://doi.org/10.1016/j.ibmb.2019.05.009

della Torre A, Fanello C, Akogbeto M, et al (2001) Molecular evidence of incipient speciation within *Anopheles*

gambiae s.s. in West Africa. Insect Mol Biol 10:9–18. https://doi.org/10.1046/j.1365-2583.2001.00235.x

della Torre A, Tu Z, Petrarca V (2005) On the distribution and genetic differentiation of *Anopheles gambiae* s.s.

molecular forms. Insect Biochem Mol Biol 35:755–769. https://doi.org/10.1016/j.ibmb.2005.02.006

Diabaté A, Baldet T, Brengues C, et al (2003) Natural swarming behaviour of the molecular M form of

Anopheles gambiae. Trans R Soc Trop Med Hyg 97:713–716.

https://doi.org/10.1371/journal.pone.0004549

Diabaté A, Brengues C, Baldet T, et al (2004) The spread of the Leu-Phe kdr mutation through *Anopheles*

gambiae complex in Burkina Faso: genetic introgression and de novo phenomena. Trop Med Int Heal

9:1267–1273. https://doi.org/10.1111/j.1365-3156.2004.01336.x

Diabaté A, Dabiré KR, Kim HE, et al (2005) Larval development of the molecular forms of *Anopheles gambiae*

(Diptera: Culicidae) in different habitats: a transplantation experiment. J Med Entomol 42:548–553

Diabaté A, Dabiré KR, Millogo N, Lehmann T (2007) Evaluating the effect of postmating isolation between

molecular forms of *Anopheles gambiae* (Diptera: Culicidae). J Med Entomol 44:60–64

Diabaté A, Dabiré RK, Kengné P, et al (2006) Mixed swarms of the molecular M and S forms of *Anopheles*

gambiae (Diptera: Culicidae) in sympatric area from Burkina Faso. J Med Entomol 43:480–483.

https://doi.org/10.1603/0022-2585(2006)43[480:msotmm]2.0.co;2

Diabaté A, Dao A, Yaro AS, et al (2009) Spatial swarm segregation and reproductive isolation between the

molecular forms of *Anopheles gambiae*. Proc R Soc B Biol Sci 276:4215–4222.

https://doi.org/10.1098/rspb.2009.1167

Diabaté A, Yaro AS, Dao A, et al (2011) Spatial distribution and male mating success of *Anopheles gambiae*

swarms. BMC Evol Biol 11:184. https://doi.org/10.1186/1471-2148-11-184

Dormont L, Bessière J-M, Cohuet A (2013a) Human skin volatiles: a review. J Chem Ecol 39:569–578.

https://doi.org/10.1007/s10886-013-0286-z

Dormont L, Bessiere JM, McKey D, Cohuet A (2013b) New methods for field collection of human skin volatiles

and perspectives for their application in the chemical ecology of human-pathogen-vector interactions. J

Exp Biol 216:2783–2788. https://doi.org/10.1242/jeb.085936

Downes JA (1969) The swarming and mating flight of Diptera. Annu Rev Entomol 14:271–298.

https://doi.org/10.1146/annurev.en.14.010169.001415

Facchinelli L, Valerio L, Lees RS, et al (2015) Stimulating *Anopheles gambiae* swarms in the laboratory:

application for behavioural and fitness studies. Malar J 14:271. https://doi.org/10.1186/s12936-015-0792-2

Feugère L, Gibson G, Manoukis NC, Roux O (2021) Mosquito sound communication: are male swarms loud

enough to attract females? J R Soc Interface 18:20210121. https://doi.org/10.1098/rsif.2021.0121

Filipiak W, Filipiak A, Sponring A, et al (2014) Comparative analyses of volatile organic compounds (VOCs)

from patients, tumors and transformed cell lines for the validation of lung cancer-derived breath markers. J

Breath Res 8:027111. https://doi.org/10.1088/1752-7155/8/2/027111

Gibson G (1985) Swarming behaviour of the mosquito *Culex pipiens quinquefasciatus*: a quantitative analysis.

Physiol Entomol 10:283–296. https://doi.org/10.1111/j.1365-3032.1985.tb00049.x

Gibson G, Warren B, Russell IJ (2010) Humming in tune: sex and species recognition by mosquitoes on the

wing. J Assoc Res Otolaryngol 11:527–540. https://doi.org/10.1007/s10162-010-0243-2

Gibson NHE (1945) On the mating swarms of certain Chironomidae (Diptera). Trans R Entomol Soc London

95:263–294. https://doi.org/10.1111/j.1365-2311.1945.tb00263.x

Gomulski L (1988) Aspects of mosquito mating behaviour. University of London

Goodrich KR, Zjhra ML, Ley CA, Raguso RA (2006) When flowers smell fermented: the chemistry and

ontogeny of yeasty floral scent in Pawpaw (*Asimina triloba*: Annonaceae). Int J Plant Sci 167:33–46.

https://doi.org/10.1086/498351

Guidobaldi F, May-Concha IJ, Guerenstein PG (2014) Morphology and physiology of the olfactory system of

blood-feeding insects. J. Physiol. Paris 108:96–111

Hahn MW, White BJ, Muir CD, Besansky NJ (2012) No evidence for biased co-transmission of speciation

Islands in *Anopheles gambiae*. Philos Trans R Soc B Biol Sci 367:374–384.

https://doi.org/10.1098/rstb.2011.0188

Howell PI, Knols BGJJ (2009) Male mating biology. Malar J 8:S8. https://doi.org/10.1186/1475-2875-8-S2-S8

Iatrou K, Biessmann H (2008) Sex-biased expression of odorant receptors in antennae and palps of the African

malaria vector *Anopheles gambiae*. Insect Biochem Mol Biol 38:268–274.

https://doi.org/10.1016/j.ibmb.2007.11.008

Kelly DW, Dye C (1997) Pheromones, kairomones and the aggregation dynamics of the sandfly *Lutzomyia*

longipalpis. Anim Behav 53:721–731. https://doi.org/10.1006/anbe.1996.0309

Kostiainen R (1995) Volatile organic compounds in the indoor air of normal and sick houses. Atmos Environ

29:693–702

Kruza M, Lewis AC, Morrison CG, Carslaw N (2017) Impact of surface ozone interactions on indoor air

chemistry: a modeling study. Indoor Air 27:1001–1011. https://doi.org/10.1111/ina.12381

Lefèvre T, Gouagna LC, Dabire KR, et al (2009) Evolutionary lability of odour-mediated host preference by the

malaria vector *Anopheles gambiae*. Trop Med Int Heal 14:228–236. https://doi.org/10.1111/j.1365-

3156.2009.02206.x

Lefèvre T, Gouagna LC, Dabiré KR, et al (2010) Beer consumption increases human attractiveness to malaria

mosquitoes. PLoS One 5:e9546. https://doi.org/10.1371/journal.pone.0009546

Lehmann T, Diabaté A (2008) The molecular forms of *Anopheles gambiae*: a phenotypic perspective. Infect

Genet Evol 8:737–746. https://doi.org/10.1016/j.meegid.2008.06.003

Levi-Zada A, Fefer D, David M, et al (2014) Diel periodicity of pheromone release by females of *Planococcus*

citri and *Planococcus ficus* and the temporal flight activity of their conspecific males. Naturwissenschaften

101:671–678. https://doi.org/10.1007/s00114-014-1206-y

Lu T, Qiu YT, Wang G, et al (2007) Odor coding in the maxillary palp of the malaria vector mosquito *Anopheles*

gambiae. Curr Biol 17:1533–1544. https://doi.org/10.1016/j.cub.2007.07.062

Maïga H, Dabiré RK, Lehmann T, et al (2012) Variation in energy reserves and role of body size in the mating

system of *Anopheles gambiae*. J Vector Ecol 37:289–297

Maïga H, Niang A, Sawadogo SP, et al (2014) Role of nutritional reserves and body size in *Anopheles gambiae*

males mating success. Acta Trop 132S:S102–S107. https://doi.org/10.1016/j.actatropica.2013.08.018

Marchand RP (1984) Field observations on swarming and mating in *Anopheles gambiae* mosquitoes in Tanzania.

Netherlands J Zool 34:367–387

McBride CS, Baier F, Omondi AB, et al (2014) Evolution of mosquito preference for humans linked to an

odorant receptor. Nature 515:222–227. https://doi.org/10.1038/nature13964.Evolution

Merlin C, Lucas P, Rochat D, et al (2007) An antennal circadian clock and circadian rhythms in peripheral

pheromone reception in the moth *Spodoptera littoralis*. J Biol Rhythms 22:502–514.

https://doi.org/10.1177/0748730407307737

Mosqueira B, Soma DD, Namountougou M, et al (2015) Pilot study on the combination of an organophosphate-

based insecticide paint and pyrethroid-treated long lasting nets against pyrethroid resistant malaria vectors

in Burkina Faso. Acta Trop 148:162–169. https://doi.org/10.1016/j.actatropica.2015.04.010

Mozūraitis R, Hajkazemian M, Zawada JW, et al (2020) Male swarming aggregation pheromones increase

female attraction and mating success among multiple African malaria vector mosquito species. Nat Ecol

Evol 4:1395–1401. https://doi.org/10.1038/s41559-020-1264-9

Nguyen PL, Vantaux A, Hien DFS, et al (2017) No evidence for manipulation of *Anopheles gambiae*, *An.*

coluzzii and *An. arabiensis* host preference by *Plasmodium falciparum*. Sci Rep 7:9415.

https://doi.org/10.1038/s41598-017-09821-x

Niang A, Nignan C, Poda BS, et al (2019) Semi-field and indoor setups to study malaria mosquito swarming

behavior. Parasit Vectors 12:446. https://doi.org/doi.org/10.1186/s13071-019-3688-0

Nyasembe VO, Tchouassi DP, Kirwa HK, et al (2014) Development and assessment of plant-based synthetic

odor baits for surveillance and control of malaria vectors. PLoS Negl Trop Dis 9:e89818.

https://doi.org/10.1371/journal.pone.0089818

Pandey SK, Kim K (2011) Human body-odor components and their determination. Trends Anal Chem 30:784–

796. https://doi.org/10.1016/j.trac.2010.12.005

Pennetier C, Warren B, Dabiré KR, et al (2010) "Singing on the wing" as a mechanism for species recognition in

the malarial mosquito *Anopheles gambiae*. Curr Biol 20:131–136.

https://doi.org/10.1016/j.cub.2009.11.040

Persiani A, Dideco MA, Petrangeli G (1986) Osservzioni di laboratorio su polimorfismi da inversione originati

da incroci tra popolazioni diverse di *Anopheles gambiae s.s.* Ann Dell'Istituto Super Di Sanita 22:221–224

Pitts RJ, Rinker DC, Jones PL, et al (2011) Transcriptome profiling of chemosensory appendages in the malaria

vector *Anopheles gambiae* reveals tissue- and sex-specific signatures of odor coding. BMC Genomics

12:271. https://doi.org/10.1186/1471-2164-12-271

Poda SB, Buatois B, Lapeyre B, et al (2021) No evidence for long-range male sex pheromones in two malaria

mosquitoes. bioRxiv Prepr. https://doi.org/10.1101/2020.07.05.187542

Poda SB, Nignan C, Gnankiné O, et al (2019) Sex aggregation and species segregation cues in swarming

mosquitoes: role of ground visual markers. Parasites and Vectors 12:589. https://doi.org/10.1186/s13071-

019-3845-5

Poda SB, Soma DD, Hien A, et al (2018) Targeted application of an organophosphate-based paint applied on

windows and doors against *Anopheles coluzzii* resistant to pyrethroids under real life conditions in Vallée

du Kou, Burkina Faso (West Africa). Malar J 17:136. https://doi.org/10.1186/s12936-018-2273-x

Poli D, Goldoni M, Corradi M, et al (2010) Determination of aldehydes in exhaled breath of patients with lung

cancer by means of on-fiber-derivatisation SPME-GC/MS. J Chromatogr B 878:2643–2651.

https://doi.org/10.1016/j.jchromb.2010.01.022

Pombi M, Kengne P, Gimonneau G, et al (2017) Dissecting functional components of reproductive isolation

among closely related sympatric species of the *Anopheles gambiae* complex. Evol Appl 00:1–19.

https://doi.org/10.1111/eva.12517

Robledo N, Arzuffi R (2012) Influence of host fruit and conspecifics on the release of the sex pheromone by

Toxotrypana curvicauda males (Diptera: Tephritidae). Environ Entomol 41:387–391.

https://doi.org/10.1603/en11037

Rund SSC, Bonar NA, Champion MM, et al (2013) Daily rhythms in antennal protein and olfactory sensitivity in

the malaria mosquito *Anopheles gambiae*. Sci Rep 3:2494. https://doi.org/10.1038/srep02494

Santolamazza F, Mancini E, Simard F, et al (2008) Insertion polymorphisms of SINE200 retrotransposons within

speciation islands of *Anopheles gambiae* molecular forms. Malar J 7:163. https://doi.org/10.1186/1475-

2875-7-163

Savolainen E (1978) Swarming in Ephemeroptera: the mechanism of swarming and the effects of illumination

and weather. Ann Zool Fennici 15:17–52

Sawadogo PS, Costantini C, Pennetier C, et al (2013) Differences in timing of mating swarms in sympatric

populations of *Anopheles coluzzii* and *Anopheles gambiae s.s.* (formerly *An. gambiae* M and S molecular

forms) in Burkina Faso, West Africa. Parasit Vectors 6:275. https://doi.org/10.1186/1756-3305-6-275

Sawadogo PS, Namountougou M, Toé KH, et al (2014) Swarming behaviour in natural populations of *Anopheles*

gambiae and *An. coluzzii*: review of 4 years survey in rural areas of sympatry, Burkina Faso (West Africa).

Acta Trop 130:24–34. https://doi.org/10.1016/j.actatropica.2013.10.015

Schiestl FP (2010) The evolution of floral scent and insect chemical communication. Ecol Lett 13:643–656.

https://doi.org/10.1111/j.1461-0248.2010.01451.x

Shelly TE, Whittier TS (1997) Lek behavior of insects. In: Choe JA, Crespi JB (eds) The evolution of mating

systems in insects and Arachnids. Cambridge University Press, New York, pp 294–309

Sivinski JM, Petersson E (1997) Mate choice and species isolation in swarming insects. In: Choe JA, Crespi JB

(eds) The evolution of mating systems in insects and Arachnids. Cambridge University Press, New York,

pp 294–309

Souto-Vilarós D, Proffit M, Buatois B, et al (2018) Pollination along an elevational gradient mediated both by

floral scent and pollinator compatibility in the fig and fig-wasp mutualism. J Ecol 106:2256–2273.

https://doi.org/10.1111/1365-2745.12995

Suh E, Choe D, Saveer AM, Zwiebel LJ (2016) Suboptimal larval habitats modulate oviposition of the malaria

vector mosquito *Anopheles coluzzii*. PLoS One 11:e0149800.

https://doi.org/10.1371/journal.pone.0149800

Tchouassi DP, Sang R, Sole CL, et al (2013) Common host-derived chemicals increase catches of disease-

transmitting mosquitoes and can improve early warning systems for rift valley fever virus. PLoS Negl

Trop Dis 7:e2007. https://doi.org/10.1371/journal.pntd.0002007

Tienpont B, David F, Bicchi C, Sandra P (2000) High capacity headspace sorptive extraction. J Microcolumn

Sep 12:577–584. https://doi.org/10.1002/1520-667X(2000)12:11<577::AID-MCS30>3.0.CO;2-Q

Tripet F, Dolo G, Traoré S, Lanzaro GC (2004) The "wingbeat hypothesis" of reproductive isolation between

members of the *Anopheles gambiae* complex (Diptera: Culicidae) does not fly. J Med Entomol 41:375–

384. https://doi.org/10.1603/0022-2585-41.3.375

Tripet F, Touré YT, Taylor CE, et al (2001) DNA analysis of transferred sperm reveals significant levels of gene

flow between molecular forms of *Anopheles gambiae*. Mol Ecol 10:1725–1732.

https://doi.org/10.1046/j.0962-1083.2001.01301.x

Vantaux A, Hien DFS, Yameogo B, et al (2015) Host-seeking behaviors of mosquitoes experimentally infected

with sympatric field isolates of the human malaria parasite *Plasmodium falciparum*: no evidence for host

manipulation. Front Ecol Evol 3:86. https://doi.org/10.3389/fevo.2015.00086

Verhulst NO, Andriessen R, Groenhagen U, et al (2010) Differential attraction of malaria mosquitoes to volatile

blends produced by human skin bacteria. PLoS One 5:e15829.

https://doi.org/10.1371/journal.pone.0015829

Wang G, Diabaté A, Liu J, et al (2021) Clock genes and environmental cues coordinate *Anopheles* pheromone

synthesis, swarming, and mating. Science (80-) 371:411–415. https://doi.org/10.1126/science.abd4359

Wondwosen B, Birgersson G, Seyoum E, et al (2016) Rice volatiles lure gravid malaria mosquitoes, *Anopheles*

arabiensis. Sci Rep 6:37930. https://doi.org/10.1038/srep37930

Wondwosen B, Birgersson G, Tekie H, et al (2018) Sweet attraction: sugarcane pollen-associated volatiles

attract gravid *Anopheles arabiensis*. Malar J 17:90. https://doi.org/10.1186/s12936-018-2245-1

Zellner B d'Acampora, Bicchi C, Dugo P, et al (2008) Linear retention indices in gas chromatographic analysis:

a review. Flavour Fragr J 23:297–314. https://doi.org/10.1002/ffj

 Figure legends

Fig. 1 Schematic of the olfactometer set up. S: swarm; M: marker. Not at scale.

Fig. 2 Mosquito activation rate, expressed as the proportion of females caught in both collecting boxes out of the

total number released females for each of the four tested combinations. Different letters indicate difference at

$P<0.05$.

Fig. 3 Mosquito choice, expressed as the proportion of females caught in one or the other collecting box out of the

total number of "activated" females for each test.

Fig. 4 Quantities (integration units) of acetoin, sulcatone, octanal, nonanal and decanal contained in the headspace

of the empty bottles, the breath of the manipulator and flying males (A) and females (B) of *Anopheles gambiae*

Kisumu and collected with SPME fibers. The intensity values correspond to the counts related to the abundance

of the specific ions representative of each molecule formed in the mass spectrometer and correspond to the amount

of compound analyzed. The box plots indicate the median (wide horizontal bars), the 25th and 75th percentiles

(squares), and the minimum and maximum values (whiskers). The black dots represent outliers and grey dots raw

data. NS=not-significant; * =$P < 0.05$; ** =$P < 0.01$; *** =$P < 0.001$.

Fig. 5 Quantities (integration units) of sulcatone, octanal, nonanal and decanal contained in a natural swarm

sampled with twisters (A); in the headspace of the empty box, swarming *Anopheles coluzzii* males and *An. gambiae*

males sampled with Tenax/Carbotrap tubes (B); or collected with Porapack tubes (C). The intensity values

correspond to the counts related to the abundance of the specific ions representative of each molecule formed in

the mass spectrometer and correspond to the amount of compound analyzed. The box plots indicate the median

(wide horizontal bars), the 25th and 75th percentiles (squares), and the minimum and maximum values (whiskers).

The black dots represent outliers and grey dots raw data. NS=not-significant; * =$P < 0.05$; ** =$P < 0.01$; *** =P

< 0.001. NA: The analytical procedure for these analyses did not allow detection of acetoin.

Fig. 6 Mean electrophysiological responses of female *An. coluzzii* antennae to male swarm solvent extracts,

recorded by coupled gaz chromatography – electroantennography detection (GC-EAD). Dashed lines indicate

retention times (and related LRI) for which the most significant EAG responses were recorded. Only 2 females in

8 tested produced this pattern.

Table 1: Air inlet and outlet rates, and VOC collection duration according to the type of trap used.

	Air flow rate (L.min^{-1})		VOC collection duration (hour)
	Air inlet	Air outlet	
Tenax/Carbotrap tube	2.5	2.0	2.0
Porapak-Q VCT tube	1.5	1.0	2.5
Micro-trap	0.9	0.4	5.0

Fig. S1 Test for olfactometer symmetry. Female choice, expressed as the proportion of female mosquitoes caught

in one or the other collecting box out of the total number retrieved from both collecting boxes for the test with

empty boxes (control *vs.* control combination).

Fig. S2 A. Chromatograms obtained with eluted Porapak extracts from headspaces of the control, an *An. coluzzii*

swarm and an *An. gambiae* swarm (linear retention index (LRI) 800 to 1700). **B.** Enlarged view of the retention

time (RT) interval containing the LRI at which *An. coluzzii* antennal responses were detected in GC-EAD. Doted

white lines correspond to the theoretical RT of Sulcatone, octanal, nonanal and decanal. No difference was detected

between the control and the swarm extracts. All the peaks correspond to emissions born from the material.

Fig. S3 A. Chromatograms obtained with the desorption of twisters obtained with the static sorptive collection

methods under natural conditions in the field (linear retention index (LRI) 800 to 1700). The twister was either

located within a non-enclosed swarm of *An. coluzzii* or 3 m away upwind (control). **B.** Enlarged view of the

retention time (RT) interval containing the LRI at which *An. coluzzii* antennal responses were detected in GC-

EAD. Doted white lines correspond to the theoretical RT of sulcatone, octanal, nonanal and decanal. When in

italic, the compound was absent. No difference was detected between the control and the swarm extracts. All the

peaks correspond to environmental emissions.

Fig. S4 Quantities (integration units) of acetoin, sulcatone, octanal, nonanal and decanal contained in the headspace

of the empty bottles, the breath of the manipulator and the breath of the manipulator after blowing for the second

time (breath ×2) sampled with SPME fibers. The intensity values correspond to the counts related to the abundance

of the specific ions representative of each molecule formed in the mass spectrometer and correspond to the amount

of compound analyzed. The box plots indicate the median (wide horizontal bars), the 25th and 75th percentiles

(squares), and the minimum and maximum values (whiskers). The black dots represent outliers and grey dots raw

data. NS=not-significant; * =$P < 0.05$; ** =$P < 0.01$.

Treatment combinations