No Evidence for Long-range Male Sex Pheromones in Two Malaria Mosquitoes

Serge Bèwadéyir Poda[1,2,3], Bruno Buatois[4], Benoit Lapeyre[4], Laurent Dormont[4], Abdoulaye Diabaté[1], Olivier

Gnankiné[2], Roch K. Dabiré[1], Olivier Roux[1,3*]

1.   *Institut de Recherche en Sciences de la Santé (IRSS), Bobo-Dioulasso, Burkina Faso.*

2.   *Laboratoire d'Entomologie Fondamentale et Appliquée (LEFA), Unité de Formation et de Recherche en*

Sciences de la Vie et de la Terre (UFR-SVT), Université Joseph KI-ZERBO, Ouagadougou, Burkina Faso.

9   3.   *MIVEGEC, IRD, CNRS, Univ. Montpellier, Montpellier, France.*

4.   *CEFE, Univ. Montpellier, CNRS, EPHE, IRD, Univ. Paul Valéry Montpellier 3, Montpellier, France.*

*corresponding author

Olivier Roux: olivier.roux@ird.fr

Running head: Sex pheromones in malaria mosquitoes

Abstract - Cues involved in mate seeking and recognition prevent hybridization and can be involved in

speciation processes. In malaria mosquitoes, females of the two sibling species *Anopheles gambiae s.s.* and *An.*

coluzzii mate in monospecific male swarms and hybrids are rare. Long-range sex pheromones driving this

behavior have been debated in literature but to date, no study has proven their existence or their absence. Here,

we attempted to bring to light their existence. To put all the odds in our favor, we used different chemical

ecology methods such as behavioral and electrophysiological assays as well chemical analyses, and we worked

with mosquitoes at their optimal physiological mating state *i.e.* with swarming males during their natural

swarming windows. Despite all our efforts, our results support the absence of long-range sex pheromones

involved in swarm detection and recognition by females. We briefly discuss the implications of this finding in

ecology, evolution and for control strategies.

Key Words - *Anopheles*, male swarm, mating behavior, mosquito control.

Cues involved in mate seeking and mate recognition prevent hybridization and can be involved in speciation

processes. In insects, these cues can be visual, acoustical or chemical with a short or a long-range action and are

highly species specific (Alexander et al. 1997; Clements 1999).

In mosquitoes, many species mate in flight, within individual aggregations called "mating swarms"

(Gibson 1945; Downes 1969; Savolainen 1978; Shelly and Whittier 1997; Sivinski and Petersson 1997). In

malaria mosquitoes of the *Anopheles gambiae* complex, mating takes place at sunset in monospecific swarms

made of a few males to thousands of them in which virgin conspecific females come to find a mate (Charlwood

and Jones 1980; Marchand 1984; Charlwood et al. 2002; Diabaté et al. 2003, 2009; Howell and Knols 2009;

Sawadogo et al. 2014). In West Africa, *An. coluzzii* and *An. gambiae s.s.* are often found in sympatry (Diabaté et

al. 2006; Costantini et al. 2009; Sawadogo et al. 2013) but they form distinct swarms spatially segregated

(Diabaté et al. 2009; Sawadogo et al. 2014) and hybrids are rare (≈1%) (della Torre et al. 2001, 2005; Tripet et

al. 2001; Diabaté et al. 2006). No evidence for selection against hybrids was found (Persiani et al. 1986; Diabaté

et al. 2005, 2007; Hahn et al. 2012; Pombi et al. 2017) and spermatheca analyses showed that these species

mostly mate assortatively (Tripet et al. 2001). This suggests that reproductive isolation between these two sibling

species is achieved by strong pre-mating reproductive barriers (della Torre et al. 2001; Diabaté et al. 2007;

Lehmann and Diabaté 2008). In addition, since females usually only mate once in their lives (Clements 1992),

errors in the choice of mate should be costly and fall under negative selection. Consequently, one would expect

to find specific cues that lead females to conspecific male swarms. However, the way females are attracted to

swarms is unknown.

In species of the *An. gambiae* complex, several cues have been identified to have a role in bringing

sexes together. Acoustic cues were shown to be involved in close-range recognition, the male and the female

adjusting their respective wing-beat frequencies to converge on a shared harmonic frequency (Gibson et al.

2010; Pennetier et al. 2010). However, it was recently demonstrated that females of some *Anopheles* species are

able to detect swarm sounds only in a very close vicinity. Thus, they are unable to use swarm sound to locate or

identify swarms at long range (Feugère et al. 2021). Visual cues also play an important role in species

segregation with *An. coluzzii* males swarming over contrasted visual markers and *An. gambiae* males swarming

over bare ground (Diabaté et al. 2009). However, a recent work showed that, like *An. coluzzii*, *An. gambiae*

males also use visual markers but rather to locate their swarm at a distance from the marker (Poda et al. 2019).

Moreover, females also use these markers to form swarms in the absence of males suggesting that females may

use these markers to join the swarm location (Poda et al. 2019). Nevertheless, the distance between two

heterospecific swarms using the same marker can be about 2 m in semi-field conditions (Poda et al. 2019) and it

is still unknown from which distance mosquitoes can see these markers, suggesting that females could cross the

location of heterospecific swarms by accident. Chemical cues have also been under investigation in *Anopheles*

species. Heptacosane, a cuticular hydrocarbon, enhances the interaction between males and females (Wang et al.

2021). This compound, however, can only be perceived by contact and can consequently be involved only in

mate recognition during courtship and to stimulate acceptance by females. Moreover, the absence of assortative

mating in confined heterospecific males and females in laboratory cages or indoor swarms (Dao et al. 2008),

suggests that close-range mating cues, if they exist, cannot ensure total reproductive isolation by themselves.

Species isolation is thus likely to occur through long-range and specific swarm recognition cues acting as a first

barrier in pre-mating processes which prevent hybridization by limiting contact between sexes of the different

species.

To our knowledge, volatile sex pheromones in the *An. gambiae* complex have never been brought to

light without ambiguity. Charlwood et al. (2002) reported an absence of response of *An. gambiae* males in

natural swarms to squashed females on filter paper or to living females in a net cage. However, in natural

conditions, females are the ones attracted to male aggregation sites. Consequently, males should be the ones that

emit long-range pheromones and should be the attractive sex. Nevertheless, a laboratory study failed to

demonstrate any attractiveness of dead males to virgin females in a Y-tube olfactometer (Gomulski 1988).

However, the difficulties in highlighting the existence of such pheromones might be due to an emission in very

low quantities and/or exclusively during swarming. Indeed, in sandfly species, which also mate in large

aggregations of males, it was shown that the larger the swarm, the greater the concentration of male-produced

sex pheromones, and hence, the greater the chances for individual males to mate with a conspecific female

(Kelly and Dye 1997; Bray et al. 2010). Similarly, Diabaté et al. (2011) showed that *An. coluzzii* females were

more frequently attracted to large swarms, suggesting a potential additive effect of cues released by males. Males

gathering in large swarms may increase their "detectability" in the female's olfactory landscape (Shelly and

Whittier 1997). In addition, in many insects, pheromone release and receptivity were shown to respond to diel

rhythms or even strict time windows of only a few minutes or hours (Bjostad et al. 1980; Merlin et al. 2007;

Rund et al. 2013; Levi-Zada et al. 2014) or even to be dependent of the presence of conspecifics (Andersson et

al. 2007; Robledo and Arzuffi 2012). Nevertheless, a recent work by Mozūraitis et al. (2020) and concomitant to

a first version of the present work (Poda et al. 2020), highlighted the existence of five volatile compounds

potentially emitted by *Anopheles* males. According to the authors, these might be involved in aggregation behavior, attracting both males and females, and increasing the insemination rate. However, though an interesting biological activity was described by the authors, the claim that these compounds are male swarming aggregation pheromones must be considered with caution and it requires further investigations. Indeed, these five compounds (acetoin, sulcatone, octanal, nonanal and decanal) are very frequently found in nature and, as reported by the authors, have been shown to have a biological activity in *Anopheles* mosquitoes in very different contexts from the reproductive behavior. They were repeatedly found in human and animal body odor (Verhulst et al. 2010; Pandey and Kim 2011; Dormont et al. 2013b, a; Tchouassi et al. 2013; McBride et al. 2014), breath (Poli et al. 2010; Calenic and Amann 2014; Filipiak et al. 2014; Cainap et al. 2020), and in host-plants (Dekel et al. 2019). These represent the main sources of food (blood and sugar meals, respectively) for *Anopheles* mosquitoes. As a matter of fact, octanal, nonanal and decanal are part of a blend used in some traps to mimic mammalian host odor for several species of mosquito vectors (Tchouassi et al. 2013; Nyasembe et al. 2014). Some of these compounds were also found associated to oviposition sites (Suh et al. 2016; Wondwosen et al. 2016, 2018) and to ambient air (Kostiainen 1995; Kruza et al. 2017). In addition, sulcatone is thought to be responsible for discrimination between humans and animals in human-seeking mosquitoes (McBride et al. 2014). Thus, it is not surprising that they trigger a strong biological/flight activity in mosquitoes.

Moreover, in their laboratory experiments, Mozūraitis et al. (2020) reported a simple flight activity instead of a real swarming activity in which males should fly in regular loops with erected antennae (Downes 1969; Poda et al. 2019). This can be explained by the absence, in their experimental design, of adequate visual stimuli such as ground markers which are mandatory to trigger swarming behavior (Charlwood and Jones 1980; Marchand 1984; Gibson 1985; Facchinelli et al. 2015; Niang et al. 2019; Poda et al. 2019). Moreover, the flight activity was extraordinary long for a swarming flight, exceeding 200 min in the presence of the five compounds. Indeed, swarming behavior is known to last for only 20-30 min in nature (Charlwood and Jones 1980; Marchand 1984; Sawadogo et al. 2013, 2014; Bimbilé Somda et al. 2018) and only up to 60 min in artificial conditions (Charlwood and Jones 1980; Poda et al. 2019). In the latter, the number of males in swarms decreased over time while the others flew randomly, bouncing on the flight arena walls (Poda et al. 2019). Indeed, swarming is an activity with a high energy demand, consuming 50% of sugar and glycogen reserves in 25 min (Maïga et al. 2012, 2014). Mosquitoes probably switched to a more random flight to try to escape and search for a sugar meal to refuel their reserves. This behavior can be stimulated by the five compounds, as they are frequently emitted from nectar sources (Dekel et al. 2019).

Finally, Mozūraitis et al. (2020) also reported that the five compounds increased insemination rates in
five different species in semi-field cages. However, instead of stopping the experiment after the swarming time,
they collected the females in the morning. As they left both a dark box in the arena as resting site and a sugar
source, it is likely that males and females also mated in or around these two resources during the night. Indeed, it
was shown that mating can occur indoors, outside of swarming times and locations (Dao et al. 2008). The boxes
and sugar sources with the five compounds were probably more attractive than the control ones gathering more
mosquitoes and thus inducing more inseminations.

Nevertheless, the discovery of sex pheromones or of a highly attractive blend in malaria mosquitoes
would be a precious step toward the development of new control and monitoring strategies. Furthermore, sex or
aggregation pheromones could help in designing sexually competitive mosquitoes in sterile insect and gene drive
techniques. Here, we attempted to bring to light the existence of long-range sex pheromones in *An. gambiae* and
An. coluzzii mosquitoes, that may allow the females to detect, recognize and track conspecific male swarms. On
the hypothesis that such pheromones could be produced only by males during swarming windows and to put all
the odds in our favor, we used different chemical ecology methods always with living swarming mosquitoes
during their natural swarming windows. As much as possible, we used both recently colonized mosquitoes and
large experimental set-ups to ensure males produced a free swarming behavior. In addition, we replicated the
main experiment by Mozūraitis et al. (2020) to search for the five compounds. We specifically added two new
controls, one to check if these compounds were male specific, and the other to discard a potential
environmental/laboratory pollution origin.

140   METHODS AND MATERIALS

Mosquitoes

Colonies originated from wild gravid females collected in inhabited human dwellings from Burkina Faso, in
2017 in Bama for *An. coluzzii* and in 2015 in Soumousso for *An. gambiae*. Bama is a village located in a rice-
growing area located 30 km North of Bobo-Dioulasso (11°24'14"N; 04°24'42"W) where previous studies
showed that the *Anopheles* population is almost exclusively composed of *An. coluzzii* (Mosqueira et al. 2015;
Poda et al. 2018). Soumousso is a typical Guinean savannah village located 30 km North-East of Bobo-
Dioulasso (11°00'46"N, 4°02'45"W) where *An. coluzzii* and *An. gambiae* coexist, with a predominance of the
latter (Diabaté et al. 2004, 2006). Gravid females were placed individually in oviposition cups containing tap
water. After oviposition, females were identified to species by routine PCR-RFLP (Santolamazza et al. 2008).

The larvae were gathered according to their species and reared in tap water, fed with Tetramin® Baby Fish Food

(Tetrawerke, Melle, Germany) *ad libitum*. Adult mosquitoes were held in 30 x 30 x 30 cm mesh-covered cages

and provided with a 5% glucose solution *ad libitum*. Insectarium conditions were maintained at 27±2 °C,

70±10% RH and 12L:12D. The colonies were repeatedly refreshed with F1 from mosquito females caught in the

wild.

Mosquitoes used in the electrophysiological study were transferred as eggs from Burkina Faso to

France. On their arrival, eggs were allowed to hatch in osmosed water and the larvae were fed with Tetramin®

Baby Fish Food *ad libitum*. Adult mosquitoes were held in 20 x 20 x 20 cm mesh-covered cages and provided

with honey diluted at 5% *ad libitum*. Females were fed with rabbit blood on a PS6 Power Unit (Hemotek,

Blackburn, UK) for egg production. Mosquitoes were reared in a laboratory climate chamber KBF-S720

(BINDER Gmbh, Tuttlingen, Germany) at 27±2 °C, 70±10% RH and 12L:12D with a sunset time programmed

at 3:00pm (synchronization of swarming/mating time and electrophysiological test time to ensure an optimal

receptivity of females). Mosquitoes used for the experiments were sexed early after emergence and sexes were

kept in separate rearing cages to prevent mating.

Mosquitoes used to replicate the experiment by Mozūraitis et al. (2020) were from a 15-year-old colony

of *An. gambiae* (Kisumu strain) reared in the IRD laboratory of Montpellier. They were maintained at

27□±□2 °C, 80□±□10% RH, with a photoperiod cycle of 12h light: 12h dark and reared as described above.

Behavioral assays

Olfactometer setup - Bioassays were conducted in a dual-port olfactometer originally designed to study host

preference in the *An. gambiae* complex (Lefèvre et al. 2009, 2010; Vantaux et al. 2015; Nguyen et al. 2017). The

odor source end was modified to enclose male swarms in boxes made of transparent plexiglass (L x W x H: 60 x

60 x 120 cm; "swarming boxes" hereafter). Each swarming box was connected by a PVC air vent hoses (L x Ø:

600 x 10 cm, W3-65014-HQ4, HQ, USA) to a collecting glass box (L x W x H: 32 x 33.5 x 44 cm) which was

linked to a mesh-covered releasing cage (L x W x H: 50 x 40 x 40 cm) by a glass tube (L x Ø: 60 x 10 cm) (Fig.

1). A custom-made electric fan was located at the mid-length of each air vent hose and drew air from the

swarming boxes (odor-sources) to the releasing cage, providing an odor-laden air current against which

mosquitoes in the releasing cage were induced to fly. The air flow was controlled thanks to two mechanisms;

first, a power regulator (HQ-Power PS1502A, Velleman, Gavère, Belgium) connected to fans; and second, iris

dampers (10 cm Ø; CIR D100, France air, France) connecting the air vent hoses to the collecting boxes. The

openings of the air vent hoses on both the swarming and collecting boxes sides were covered with nets to prevent

mosquitoes from flying into the air vent hoses. The swarming boxes (odor-sources) were located side-by-side

outdoors and the olfactometer inside a room (Fig. 1). The air speed in the releasing cage was regulated at 18 ± 2

cm.s^{-1} using an anemometer (Model 425, Testo, Forbach, France) and the room temperature and relative

humidity were set at 27 ± 2 °C and $80\pm10\%$ RH thanks to an air conditioner and a humidifier (Defensor 505,

Condair, Croissy-Beaubourg, France), respectively.

Bioassays - Bioassays were performed to study the long-range behavioral response of females to whole volatile

organic compounds (VOCs) associated with male swarming activity. For each replicate, about 100 2- to 3-day-

old virgin females of *An. coluzzii* and *An. gambiae* were used simultaneously. Females were colored the morning

of the test with two different colored powders (Luminous Powder Kit, BioQuip, Rancho Dominguez, California,

USA) according to species. The matching between the species and the colors was switched between each test.

About 500 4- to 5-day-old *An. coluzzii* or *An. gambiae* males were released into a swarming box in the morning

about 8 hours before the time of the test to allow them to acclimate. Both males and females were provided with

a 10% glucose solution and were kept under insectary conditions until the time of the test. To avoid odor bias

during the test, the empty box (control) was also provided with glucose.

Bioassays were carried out at sunset, when males naturally swarm. About 30 min before sunset time, the

swarming boxes were moved outside, the cups containing the glucose pads were removed and a visual marker

made of a 20 x 20 cm black cloth were placed either under or next to the boxes for *An. coluzzii* or *An. gambiae*,

respectively (see Poda et al. 2019). These markers allowed to trigger swarming behavior and helped to stabilize

swarms in the middle of the box. For the same purpose, we provided an artificial twilight horizon made of a 40

W incandescent bulb (2,500 K) located on the floor between a white wall and a 50 cm high black horizon

(Facchinelli et al. 2015; Niang et al. 2019; Poda et al. 2019). Inside, females of the two species were released in

equal numbers into the releasing cage of the olfactometer allowing them to acclimate for 30 min before sunset.

Previously, the fans were switched on in order to purge the air vent hoses and then the air flow was set at 18 ± 2

cm.s^{-1}. The glass tube openings on the releasing cage side were covered with nets to prevent female mosquitoes

from flying into the collecting boxes before the test began. The complete olfactometer, except the end of the

releasing cage, was covered with a white cloth to eliminate visual bias during the test and to provide a diffused

light. Sunset light was allowed to enter the olfactometer room until dark and an additional artificial twilight

made of a 40 W incandescent bulb (2,500K) projected on the wall of the room on the side of the collecting boxes

was also provided.

When the males started to swarm (*i.e*. flying in loops at a constant location within the swarming boxes,

see Downes 1969; Poda et al. 2019), the nets covering the glass tube openings and preventing access of females

to the collecting boxes were removed thanks to threads that were attached to the nets. The females were allowed

to respond for 20 min (swarm duration), then the glass tube openings were covered again with nets and females

that reached the upwind collecting boxes were removed with a mouth aspirator, killed by freezing and identified

according to their coloration (species) under a binocular (LEICA S6E) and counted. The remaining mosquitoes

inside the releasing cage were also removed. After each test (1 per day), the olfactometer was cleaned with 95%

ethanol to remove odor contaminants. All materials were handled with gloves to avoid contamination with skin

compounds.

Four combinations of choice tests were performed: i) *An. coluzzii* male swarm *vs*. empty box (hereafter,

An. coluzzii test); ii) *An. gambiae* male swarm *vs*. empty box (hereafter, *An. gambiae* test), iii) *An. coluzzii vs*.

An. gambiae male swarms (hereafter, *An. coluzzii vs*. *An. coluzzii* test) and iv) empty *vs*. empty box (hereafter,

control test). Each combination was repeated 12 times and females were used only once. Males were used twice

in two consecutive days. In that case, a 10% glucose solution was introduced into the two swarming boxes after

the test and boxes were kept under insectarium conditions until the next day. The swarming boxes were cleaned

with ethanol between each batch of males. The treatments (mosquitoes or control) between the swarming boxes

and the right and left arm of the olfactometer were alternated to avoid cage and side effects, respectively. We

assessed the instrumental and side bias through an empty *vs*. empty box choice test (control test). The

olfactometer did not present any symmetrical biases (Fig. S1).

Collection of the volatile organic compounds (VOCs), chemical and electrophysiological analysis.

Because compounds can be emitted at a given time and only produced in tiny quantities at individual scale, we

chose to work directly on swarms. Three different sampling methods were used in Burkina Faso; two methods

were run in a laboratory setup (Niang et al. 2019) and one in the field.

Dynamic headspace - The headspace volume consisted of a 50 x 50 x 50 cm transparent plexiglass box. Such a

volume was a trade-off between a volume large enough to allow males to display a swarming behavior without

too much constraints and a volume small enough to be sucked up in a reasonable time. Entrance and exit flow

rates were maintained by two pumps and regulated by flowmeters. The entrance flow was higher than the exit

flow ensuring that the system was continuously purged to compensate for the inevitable leaks, and that no

contaminated outside air would enter the system. At the entrance side, air was purified in a glass charcoal filter

and then humidified passing through deionized water. Exit flow passed through an odor trap to adsorb VOCs.

Three different odor traps were used for the chemical analyses and electrophysiological analyses. Tenax-

TA/Carbotrap sorbent stainless steel 6mm diameter tubes (Gerstell, Mülheim, Germany) and Porapak-Q sorbent

VCT glass tubes (ARS, Gainesville, Florida, USA) were used for chemical analyses. Porapak-Q sorbent VCT

tubes and home-made sorbent Micro-traps were used for electrophysiological analysis. Micro-traps were

constituted by ChromatoProbe® quartz microvials of Varian Inc. (length: 15 mm; inner diameter: 2 mm),

previously cut closed-end and filled with 3 mg of a 1:1 mix of Tenax-TA and Carbotrap® (60–80 and 20–40

mesh, respectively; Sigma Aldrich, Munich, Germany)(See Dormont et al. 2013b). Flow rates and extraction

durations were dependent on the odor trap (See Table 1). The entire setup was placed in a room specially

designed to stimulate *Anopheles* swarming behavior thanks to a set of visual stimuli (i.e. lights and ground

marker, see Niang et al. (2019) for details).

About 500 4- to 6-day-old males of *An. gambiae* or *An. coluzzii* were transferred into the box about 30

min before swarming time. Clean air was pushed into the box at a rate of 10 L.min^{-1} to purge the box from both

human and mosquito container odors before connecting it to the odor trap. Five minutes after males started to

swarm (*i.e.* flying in loops at a constant location within the box), the air flow was adjusted at the required rate

and the odor trap connected. Forty minutes later, swarming behavior was stopped by turning off the light and the

VOC collection continued until the required collection time was reached (see Table 1). After extraction, samples

were stored at 4° C until analysis. After males were removed, the box was cleaned with 95% ethanol and then

flushed with clean air for at least 5 hours to remove the odor contaminants left from the previous extraction. A

total of 30 swarm extracts were collected; 17 *An. coluzzii* swarms (5, 8 and 4 with Tenax/Carbotrap tubes,

Porapak-Q VCT tubes and Micro-traps, respectively) and 13 *An. gambiae* swarms (6 and 7 with

Tenax/Carbotrap tubes, and Porapak-Q VCT tubes, respectively). Control consisted of a clean empty box. Before

they were sent and used in Burkina Faso, adsorbents were carefully conditioned and checked for residual

compounds and potential pollutions in a gas chromatograph, then sealed (Tenax-TA/Carbotrap tubes), packed in

a glass vial with a Teflon cap (ChromatoProbes) or in a nalophan bag (Porapak-Q VCT tubes).

VOC collection from natural swarms - VOCs were collected from a natural swarm of *An. coluzzii* in the village

of Bama. Static sorptive extractions of volatiles from a non-enclosed swarm were performed with Twisters®

(100 µm PDMS stir bar; Gerstel, Mülheim, Germany; Bicchi et al. 2000; Tienpont et al. 2000). The swarm flew

about 2.5 to 3.5 m above the ground. Thus, we used a 1.5 m long glass stick (Glaswarenfabrik Karl Hecht GmbH

& Co, Germany) with a metallic push pin head glued at the tip and covered with nalophan to introduce the

magnetic twisters directly into the swarm. Control twisters were placed about 3 m away upwind from the swarm.

Extraction lasted for the whole swarm duration (about 20-25 min). We chose a swarm apart from habitations and

livestock to limit odor pollutions, containing a large number of males (more than 1,000) and being attractive for

females (observation of a large number of couples). After VOC collections, twisters were individually placed

into glass vials and stored at 4 °C until analysis. Five replicates were performed on the same swarm at different

days.

Solvent extraction - About 1,000 4- to 6-day-old males of *An. gambiae* or *An. coluzzii* were introduced into a 30

x 30 x 30 cm cage and kept under insectarium conditions. The cage was made of unpainted metallic frames

covered by white nets and the bottom was covered with a white paper. Thirty minutes before sunset time, the

cage was placed outside and mosquitoes were observed. A 5 x 5 cm black cloth was placed in or out of the cage

according to the species as described above to stimulate swarming behavior. About 5 min after mosquitoes

started to swarm, the entire cage was quickly placed into a freezer at -20° C for about 10 min. Then, mosquitoes

were transferred into a 20 ml glass tube and covered with a 1:1 mix of Hexane and Dichloromethane. The tube

was sealed and mosquitoes were kept in the solution for 24h at 27±2 °C. Then, the solution was filtered on glass

wool to remove mosquito scales and stored at 4° C until analysis. Three different controls were made: i) with 1-

to 2-day-old virgin males (young male control), ii) with 2- to 3-day-old virgin females (female control) and iii)

with solvent mix alone (blank control). A total of 8 swarm extracts were done; 4 of *An. coluzzii* and 4 of *An.*

gambiae and 1 replicate of each type of control per species.

SPME extraction – We replicated the experiment from the study by Mozūraitis et al. (2020) in which they

collected mosquito headspace with a SPME fiber in a 1L bottle. However, because most methods used to

introduce mosquitoes into a bottle may potentially result in entering pollutants as well, we added a

supplementary control. One way to easily introduce the mosquitoes is to blow them from a mouth aspirator into

the bottle. This classic method may bring breath volatiles within the recipient, which can contain the five

compounds described in Mozūraitis et al. (2020) (Poli et al. 2010; Calenic and Amann 2014; Filipiak et al. 2014;

Cainap et al. 2020). Consequently, we added a "breath control" which is mandatory to discriminate compounds

from the experimenter's breath from those actually emitted by mosquitoes. In addition, we also replicated the

experiment with females to check for sex specificity. In the morning, the two glass bottles were cleaned with

acetone and dried under a clean air flow, then closed with nalophan. Three hours before swarming time (5pm),

we collected the headspace of empty bottles with polydimethylsiloxane/divinylbenzene-coated SPME fibers for

1h (step 1: empty bottle). Then, we simulated the introduction of mosquitoes by blowing 3 times for 3 seconds

with a mouth aspirator into the bottles. These were then closed with Nalophan (6pm) and the head space of the

breath of the experimenter was collected for 1h with SPME fibers (step 2: breath control). We then introduced

60-70, 5- to 7-day-old unmated males in each bottle with the mouth aspirator, closed the bottles with Nalophan

(7pm) and let the mosquitoes acclimate for 1h. During the acclimation period, we simulated a light transition

from photophase to scotophase to trigger crepuscular flight. Then, when males started to fly (8pm), SPME fibers

were introduced for 1h (step 3: "swarm" extract). Six different fibers were used and cleaned at 250 °C during 5

min in the injector of a gas-chromatograph (HP 6890 Series PLUS, Agilent, Santa Clara, USA) just before use.

Each fiber was always used for the same type of extract and in the same bottle. Before the experiment, we made

sure that the fibers had similar sensitivity by exposing them to a mix of the five compounds. Only minor

quantitative differences were detected, and this was considered during subsequent statistics. Secondly, we

performed the same 3 steps but with 60-70, 3- to 4-day-old virgin females instead of males. And finally, as

during the introduction of mosquitoes, we blew a second time into the bottles and repeated the three steps of

headspace extraction but this time, instead of introducing mosquitoes in the 3rd step, we blew again into the

bottles (breath x2) to check for potential accumulation of breath compounds. We kept track of the day of the

experiment, the bottles and fibers identity for statistical analyses.

Chemical analyzes

Samples were analysed at the "Platform for Chemical Analyses in Ecology" (PACE), technical facilities of the

LabEx CeMEB (Centre Méditerranéen pour l'Environnement et la Biodiversité, Montpellier, France). Those

obtained on Porapak-Q VCT tubes were eluted with 150 µl of Hexane and injected (1 µl) into a Gas

Chromatograph coupled with Mass Spectrometer (GC-MS). Those obtained on Tenax/Carbotrap tubes, twisters

or SPME fibers were directly thermo-desorbed into GC-MS. Solvent extracts were injected as is or concentrated

under a gentle stream of nitrogen.

Liquid analysis (Porapak elutions and solvent extractions) - Liquids were analysed with a GC-MS QP2010 Plus (Shimadzu, Kyoto, Japan). Mass spectra were recorded in electronic impact mode (EI) at 70eV over a m/z mass range from 38 to 350. The temperature of the transfer line and the ion source were programmed to 250 °C and 200 °C, respectively. The injections were made with an injector temperature of 250 °C, and a 1:4 split mode ratio. Analyses were performed using a 30 m x 0.25 mm x 0.25 µm Optima 5-MS (Macherey-Nagel, Düren, Germany) fused silica capillary column with a constant helium flow set close to 1.0 ml.min^{-1}. The oven temperature was programmed as follows: 40 °C (held 5 min.) to 250 °C at 6 °C.min^{-1}, and finally to 300 °C at 14 °C.min^{-1} and held 2 min. GC-MS Solution software (Shimadzu, Kyoto, Japan) was used for data processing of these analyses, with the NIST 2011 as spectrum database.

Tenax/Carbotrap tubes and twister analysis - These sorbents were analysed as described by Souto-Vilarós et al. (2018). Samples were analysed using a gas chromatograph (GC, Trace™ 1310, Thermo Scientific™ Milan, Italy) coupled with a mass spectrometer (MS, ISQ™ QD Single Quadrupole, Thermo Scientific™ Milan, Italy). The column used was an Optima 5-MS, the same as for liquid analysis. Absorbent traps were handled with a Multi Purpose Sampler (Gerstell, Mülheim, Germany) and desorbed with a double stage desorption system, composed of a Thermal Desorption Unit (TDU) and a Cold Injection System (CIS) (Gerstell, Mülheim, Germany). The sorbents were desorbed in splitless mode at 250 °C on a trap cooled at −80 °C by liquid nitrogen. Then, the trap was heated to 250 °C with a 1:4 split ratio to inject the compounds in the column. The carrier gas used was helium at 1 ml.min^{-1}. Oven temperature program was as follow : held at 40 °C for 3 min., then increased to 220 °C at 5 °C.min^{-1} and finally to 250 °C at 10 °C.min^{-1}, and held for 2 min. Mass spectra were recorded in electronic impact mode (EI) at 70eV over a m/z mass range from 38 to 350. The temperature of the transfer line and the ion source were programmed to 250 °C and 200 °C, respectively. Xcalibur™ software (Thermo Scientific™, Milan, Italy) was used for data processing and the NIST 2011 as spectrum database.

SPME fiber analysis – We analyzed SPME fibers as described in Mozūraitis et al. (2020) with a GC-MS QP2010-SE (Shimadzu, Kyoto, Japan). Volatiles were desorbed into the injector at 225 °C for 1 min in split mode (1:4 ratio). Analyses were performed using a 30 m x 0.25 mm x 0.25 µm J&W DB-Wax silica capillary column (Agilent, Santa Clara, USA) with a constant helium flow at 1 ml.min^{-1}. The oven temperature was programmed as follows: 40 °C (held 1 min.) to 150 °C at 5 °C.min^{-1}, and finally to 220 °C at 20 °C.min^{-1} and

held 9 min. Mass spectra were recorded in electronic impact mode (EI) at 70eV over a m/z mass range from 38

to 350. The temperature of the transfer line and the ion source were 250 °C and 200 °C, respectively. GC-MS

Solution software (Shimadzu, Kyoto, Japan) was used for data processing of these analyses. We injected a

solution of standards made of the five compounds (acetoin, sulcatone, octanal, nonanal and decanal, ≥98%GC,

99%, 99%, ≥98% and ≥98%GC, respectively, Sigma-Aldrich, Darmstadt, Germany) to identify both their

specific ions and their retention time.

Electrophysiological analysis

Electrophysiological activity of female olfactive detection, on the male swarm extracts, was tested with an

Electro-Antennografic Detection system, coupled with Gas Chromatography (GC-EAD) at the PACE. Thirty

minutes before the mosquito scotophase, both 4- to 6-day-old males and 2- to 3-day-old virgin females were

placed separately in cardboard cups (Ø = 75 mm, h = 100 mm), provided with a 5% honey solution and

transferred from the insectarium to the electrophysiology laboratory at PACE. At "mosquito sunset time",

mosquitoes were placed in dark. Males were checked for erected antennae, ensuring that change in light and dark

rhythm, transport and electrophysiology laboratory conditions did not alter their physiological state for mating.

Two minutes after male antennae were erected, a female was removed from the cardboard for the antennal

preparation.

The female's head was gently excised with a pair of surgical scissors and mounted between two glass

capillary tubes of 76 mm in length and 1.12 mm in diameter (World Precision Instrument, Sarasota, USA),

pulled and cut using a vertical micropipette-puller (P-30 model, World Precision Instruments, Hertfordshire,

United Kingdom) filled with insect Ringer solution (6.0 g.l^{-1} NaCl, 0.4 g.l^{-1} KCl, 0.27 g.l^{-1} CaCl$_2$ and 3.20 g.l^{-1}

NaC$_3$H$_5$O$_3$) and connected to silver wires. The indifferent electrode was inserted into the back (the foramen) of

the isolated head and the intact tip of one antenna was inserted into the recording electrode which was connected

to an Electro-Antennography Detector setup (EAD, Syntech IDAC-2, Kirchzarten, Germany). One end of the

glass electrodes was pulled to a fine point which was then broken to ensure a close fit with the female's head and

antenna.

The above EAD setup was linked to a Gas Chromatograph-Flame Ionization Detector (GC-FID, CP-

3800, Varian, Palo Alto, USA) equipped with an Optima 5-MS capillary column, the same as for the GC-MS

analyses. Liquid injections after Porapack solvent elution were made in a 1079 PTV injector (Programmed

Temperature Vaporizator) at 250 °C with a 1:4 split ratio. Microtraps were desorbed in the same injector using a

ChromatoProbe sample introduction device (Varian, Palo Alto, USA) with the same split ratio and temperature

programmed as follows: increased from 40 °C to 250 °C at 200 °C.min^{-1} and held for 5 min. Oven temperature

was held at 50 °C for 0.40 min, then increased to 180 °C at 10 °C.min^{-1}, and finally to 220 °C at 12 °C.min^{-1}, for

a run duration less than 17 min. The carrier gas used was helium at 1 ml.min^{-1}. The effluent was split equally

into two deactivated fused silica capillary columns (100 cm x 0.25 mm), one leading to the FID (270 °C) and the

other into a heated EAD port (200 °C) (transfer line: Syntech, Kirchzarten, Germany) and leaded to antenna. The

GC effluent for antenna was mixed with charcoal-filtered, humidified airflow (300 ml.min^{-1}). StarWorkstation

6.41 software (Varian, Palo Alto, USA) was used for data processing.

396   Different types of male swarm extracts were tested on several conspecific females' antennae. Antennae

of 19 *An. coluzzii* females were tested with solvent extracts, Porapak-Q VCT tube extracts and Micro-trap

extracts (8, 9 and 2 females, respectively) and 20 *An. gambiae* females were tested with solvent and Porapak-Q

VCT tube extracts (10 females each). Before GC-EAD analysis of the VOC extracts, we checked the living

antenna for good activity using a negative (solvent) and a positive (mix of many VOCs) stimulus solutions. We

proceeded as follows: 5μl of each solution was applied to a 1 cm x 2 cm filter paper contained in a Pasteur

pipette. The pipette was then placed in an air pump system and the volatiles were directly sent to the antennal

preparation with a pulse duration of 0.5 s and a flow rate of 890 ml.min^{-1} regulated by a CS-55 Stimulus

Controller (Syntech, Hilversum, Netherlands).

Search for specific and active compounds

407   For all the chemical and electrophysiological analyses (Liquid, Tenax/Carbotrap tubes, Micro-traps and

408   twister analysis), a solution of n-alkanes (Alkanes standard solution, 04070, Sigma Aldrich, Darmstadt,

409   Germany) was injected to calculate the linear retention index (LRI) of the compounds of interest (Zellner et al.

410   2008). To identify the compounds, we matched the mass spectra with the NIST 2011 MS library, and the LRIs

411   with those reported in the literature.

412   In the electrophysiological analysis, we studied the chromatograms in a range of LRI between 800 and

1700, including compounds present at very low and trace amounts. When a depolarization was observed in the

electrophysiological analysis, a section of 20 points of LRI was studied around the compound's LRI on the GC-

MS trace. Comparison of the samples with their respective controls collected on the same day, allowed to

subtract potential contaminants from the samples. Non-natural compounds, such as industrial chemicals and/or

compounds not naturally produced by living organisms (Charpentier et al. 2012), were considered as
contaminants.

We also particularly searched for the presence of acetoin, sulcatone, octanal, nonanal and decanal
reported by Mozūraitis et al. (2020). With that aim, we defined a LRI window of 20 points on both sides of the
LRI of these compounds on the bases of injected standards (for SPME fibers) or literature (for other extraction
methods). Then, we searched for their specific ions at those LRI as follows: (LRI/ion) 711/88, 981/108, 998/84,
1100/98, and 1201/112 for acetoin, sulcatone, octanal, nonanal and decanal, respectively. Then, when a peak
corresponding to both LRI and specific ion was found, we compared its spectra with the NIST 2011 MS library
before quantification with integration units.

Statistical analysis

All analyses were performed using R (version 3.4.0). Both the activation rate and choice were analyzed
using binomial Generalized Linear-Mixed Models (GLMM, "*glmer*" function in "*lme4*" package). Tested
combinations (four levels: *An. coluzzii*, *An. gambiae*, *An. coluzzii vs. An. gambiae* and control tests), female
species (two levels: *An. coluzzii* and *An. gambiae*) and their interaction were considered as fixed effects, and
replicates as random effects.

We analyzed separately the quantities using integration units of acetoin, sulcatone, octanal, nonanal and
decanal. When pairing between the controls and the mosquito extracts was possible (with SPME and twister) and
to account for repeated measurements, we used Gaussian GLMMs. In those cases, the extract (3 levels for SPME
fibers: empty bottles, breath and flying mosquitoes (either male or females or breath x2) and 2 levels for twisters:
control and swarm) was considered as fixed effect. The replicates (i.e. the day for twisters and the SPME) and
fiber nested within the bottle for SPME fibers were considered random effects. Considering the SPME fibers as a
random effect made it possible to account for their minor differences of sensitivity in the model. When pairing
between the controls and the mosquito extracts was not possible (Tenax-TA/Carbotrap and Porapak-Q VCT
tubes), Gaussian GLMs were used. The extract (empty box, *An. coluzzii* swarm or *An. gambiae* swarm) was
considered as fixed effect. In some occasions, some compounds were not found in the extracts or only in two
samples. Consequently, it was not possible to perform statistical analyses and we considered the extracts as not
significantly different. No statistical analysis was performed with liquid extracts due to a small number of
replicates. For model selection, we used the stepwise removal of terms, followed by likelihood ratio tests. Term

removals that significantly reduced explanatory power (P <0.05) were retained in the minimal adequate model.

All percentages are provided with their 95% confident interval.

449   RESULTS

Behavioral assays

A total of 8,918 females were tested (4,423 *An. coluzzii* and 4,495 *An. gambiae*) among which 3,591 flew

upwind into the collecting boxes (activation rate: 40.26±1.01%). There was a significant interaction between

female species and the four combinations of swarm boxes tested (χ_3^2= 22.19, P <0.001). Species subset analyses

showed that the different test combinations had no effect on the *An. coluzzii* activation rate (χ_3^2 = 5.72, P=0.12;

Fig. 2). However, in *An. gambiae*, there was a significant effect of the test combinations on the female activation

rate (χ_2^2 = 14.94, P = 0.001; Fig. 2) with a higher activation rate when females were exposed simultaneously to

both *An. coluzzii* and *An. gambiae* male swarms than when exposed to an *An. coluzzii* swarm alone (z = 4.01, P

<0.001) or to the control (two empty boxes: z = 2.57, P = 0.04). Nonetheless, no difference was found when

females were exposed to an *An. gambiae* male swarm alone (z = -1.49, P = 0.44). Neither the *An. coluzzii*

swarms, nor the *An. gambiae* swarms were attractive for activated females (female species: χ_1^2 = 2.45, P = 0.11;

choice test combinations: χ_2^2 = 1.15, P = 0.56 and female species x choice test combination interaction: χ_2^2 =

0.34, P = 0.84; Fig. 3).

Chemical analyses

All the peaks observed in the volatile profiles from male swarms were also present in control samples for both

species (Figs S2 & S3). GC-MS traces obtained with the Tenax/Carbotrap tubes showed peaks identified as

pollutants, such as silicones, BTX (mono-aromatic compounds like benzene, toluene and xylene), alkanes which

were all emitted by the box containing the mosquitoes. Only few differences, including linear organic acids,

were noted but not reproducible. GC-MS traces from the Porapak-Q samples, Twisters® and the liquid extracts

were similar to their respective control.

During the experiment replicating the protocol by Mozūraitis et al. (2020), both males and females

started to fly at the expected time (8pm) but due to the small volume it was not possible to determine if the group

was a swarm. Analyses of SPME fibers in search for acetoin, sulcatone, octanal, nonanal and decanal showed

that acetoin was absent from almost all the samples with only one occurrence in a "swarming" male extract and

in a breath control (over seven) (Fig. 4A), and on three occurrences in female extracts (over six) (Fig. 4B).

Sulcatone, octanal, nonanal and decanal were found in all types of extracts including controls. Sulcatone was

absent from two breath extracts, and both sulcatone and octanal were absent from one female extract while they

were present in their corresponding empty bottle controls. We did not find significant increased quantities of the

five compounds in "swarming" mosquitoes extracts compared to the breath controls (Fig. 4). The control to test

the effect of blowing twice in the bottle showed that the headspaces of "breath" and "breath x2" were very

similar (Fig. S4), demonstrating an absence of accumulation of breath odor between the breath control and the

introduction of mosquitoes.

A posteriori, we also quantified acetoin, sulcatone, octanal, nonanal and decanal in our previous GC-

MS analyses in search for similar variations as described in Mozūraitis et al. (2020). We did not find acetoin in

our chromatograms. Indeed, as we used a low polarity Optima 5-MS column, acetoin (if present) was eluted

before the LRI 800. In the samples obtained from natural swarms of *An. coluzzii* with the twister method,

sulcatone and octanal were absent. The quantities of both nonanal and decanal were not significantly higher in

the swarm than in the control (Fig. 5A). In laboratory swarm samples, octanal was absent from both *An. coluzzii*

and *An. gambiae* headspace obtained with the Tenax/Carbotrap tubes while it was present on two occasions in

the empty box. Sulcatone, nonanal and decanal were present in both the empty box and in swarms. However,

swarms did not show higher quantities of these compounds compared to the empty box (Fig. 5B). The samples

obtained with the Porapack tubes, showed that octanal, nonanal and decanal tended to have higher quantities in

both *An. coluzzii* and *An. gambiae* swarms than in the empty box (Fig. 5C). However, this difference was not

statistically significant and only two samples were available for the empty box limiting the reliability of the

analysis. Whatever, the method used, we can note a high quantitative variability between the samples. This was

true for both the control and the mosquito samples.

Electrophysiology analyses

The male swarm VOCs induced a few weak and poorly reproducible depolarizations on conspecific females'

antennae. In *An. coluzzii*, 3 females out of 9 weakly responded to a single Porapak-Q extract at the linear

retention index (LRI) 1371 (retention time (RT) 12.85 min), and 2 females out of 8 responded to solvent extracts

at LRI 921, 978 and 1179 (RT 6.40, 7.24 and 10.23 min, respectively, Fig. 6). However, these responses did not

correspond to noticeable peaks in the GC detector of the GC-EAD apparatus.

Successive GC-MS analyses of the same porapak-Q and liquid extracts did not allow to detect and

identify any specific peak in relation to antennae responses (Figs S2b & S3b). The peak at LRI 1371 that elicited

a response in *An. coluzzii* females was identified as an alkane family compound (m/z: 57, 43, 71, 85, 99) also

present in the control sample. In liquid extracts, the same type of LRI section was studied, LRI 921 was

identified as Hexane-2,5-dione (m/z: 43, 99, 71, 57, 114), a chemical pollutant. The section studied for the LRI

978 and 1179 showed no differences with the control sample.

No reproducible response was recorded from *An. coluzzii* females tested with Micro-traps extracts. In

An. gambiae, no reproducible antennae response was recorded from the 10 females tested with either Porapak-Q

or liquid extracts.

DISCUSSION

For the past decades, several authors have suggested the existence of long-range sex pheromones in the attraction

of *An. gambiae s.l.* females to male swarms (Charlwood et al. 2002; Tripet et al. 2004; Diabaté et al. 2011; Poda

et al. 2019) but to date, no reliable experimental evidence has been available. Here, we used behavioral,

physiological and detailed chemical approaches to attempt to highlight the existence of such compounds.

However, despite a large set of methodologies and working hypotheses to maximize the chances of highlighting

the presence of pheromones (*i.e.* timing, physiological state, natural behavior, recently colonized mosquitoes),

our study failed to provide evidence of the presence of long-range male sex pheromones in both *An. coluzzii* and

An. gambiae. We also replicated an experimental setup that recently showed an increase in quantity of five

compounds in "swarming" males compared to controls. However, unlike in Mozūraitis et al. (2020), we did not

find any significant difference with our controls.

Our behavioral tests in the olfactometer did not reveal any attraction effect of both hetero- and

conspecific male swarms to females. During the experiment, the males released in the large Plexiglas boxes and

exposed to natural light started to fly randomly at sunset. Then, about 150-200 males out of the 500 released

(<50%) gathered in the upper half of the box flying in loops without touching the box walls. They also reacted to

visual marker movements, meaning they were indeed swarms. The other males flew randomly in the box,

bouncing against the walls. Females showed a good activation rate but they did not show a particular choice for

any arm, containing a swarm or not. Nevertheless, *An. gambiae* females showed an intriguing higher activation

rate when exposed simultaneously to both a conspecific and a heterospecific swarm compared to a single *An.*

coluzzii swarm. This was, however, probably without biological significance as no difference was detected in the

test providing *An. gambiae* females with a choice between the control and *An. gambiae* swarm, *vs.* a test offering

two controls.

This result was in agreement with our physiological study. Indeed, despite the fact that we checked for

both behavioral receptivity of mosquitoes and for receptivity of mounted antennae, no consistent antennal

depolarization was observed in females across assays when exposed to swarm extracts. This supports a lack of

detection by females of any volatile chemicals present in our swarm extracts.

Chemical analyses were also negative and no compounds specific for male swarms could be detected

whatever the method used. Moreover, unlike in Mozūraitis et al. (2020), the quantities of acetoin, sulcatone,

octanal, nonanal and decanal were found inconsistently across swarming mosquito samples and not in larger

quantities compared to controls, making it difficult to support the assumption according to which they could

actually be emitted by males. These divergent results could be explained by the fact that, unlike in the

experiments of Mozūraitis et al. (2020), we used a control which considered the potential introduction of

pollutions at the same time as mosquitoes. This showed that breath was responsible for most of the variability.

The most convincing result showing that sulcatone, octanal, nonanal and decanal are probably not pheromones

emitted by males is the one obtained with the twister method. Indeed, twisters are coated with the same

adsorbent phases as SPME fibers used both in this study and in the one by Mozūraitis et al. (2020). According to

the manufacturer, twisters are up to a thousand times more sensitive than SPME fibers partly due to a larger

sorbent volume. Moreover, we exposed the twisters directly in a natural swarm formed by more than 1 000

males (probably up to 6 000 males according to the estimation of trained technical staffs). Despite this

sensitivity, the number of mosquitoes and the natural biological context, sulcatone and octanal were not found

and both nonanal and decanal had similar quantities in the swarm than outside the swarm (control twister placed

3 m upwind from the swarm). In addition, the high quantitative variability found in the laboratory and the fact

that these compounds are frequently found in controls suggest that they could be laboratory and/or human

pollutions that are difficult to control for.

Independently, our methods can have weaknesses, such as the sensitivity thresholds of our apparatus

which are below the ability of insects to detect low amounts of compounds. However, all together and keeping in

mind that an absence of evidence is not an evidence of absence, our results support the absence of long-range sex

pheromones emitted by male swarms. However, further investigations are needed and complementary

methodologies such as electro-physiology on palps (Lu et al. 2007; Iatrou and Biessmann 2008; Pitts et al. 2011;

Guidobaldi et al. 2014), molecular analyses of olfactory protein expression on adequate physiological status, or

real time chemical analyses of swarm volatiles with a sensitive apparatus such as a proton-transfer-reaction mass

spectrometry could be options.

The question of how *Anopheles* females seek male swarms is still open. The lack of post-mating barrier

(Persiani et al. 1986; Diabaté et al. 2005, 2007; Hahn et al. 2012; Pombi et al. 2017) and the low rate of hybrids

in this geographical area (della Torre et al. 2001, 2005; Tripet et al. 2001; Diabaté et al. 2006) necessarily

involve a reasonably distant process which prevents females entering heterospecific swarms. As acoustic cues

have shown some limitations at long range (Feugère et al. 2021), and because long-range chemical cues are still

disputable, visual cues such as ground markers (Diabaté et al. 2009; Sawadogo et al. 2014; Poda et al. 2019)

could be good candidates, but their effective range and level of specificity is poorly known.

From an evolutionary and ecological point of view the absence of long-range sex pheromones is of

upmost importance. It demonstrates that the ecological barriers that allow these two species to mate assortatively

are probably not of chemical origin. Regarding control strategies, thanks to our findings, research programs

working on the sterile insect technique (SIT) or genetically-modified mosquitoes (GMM) will avoid to focus on

the eventuality that the induced modifications may alter long-range sex pheromones making their mosquitoes

less competitive than wild individuals.

Declarations

Funding

This work was funded by a grant from the Agence Nationale de la Recherche (ANR-15-CE35-0001-01) awarded

to O.R. S.B.P. received financial support through a doctoral fellowship from the Institut de Recherche pour le

Développement (IRD).

Competing interests

We have no competing interests.

Availability of data and material

The raw datasets are available from the corresponding author upon reasonable request.

Code availability

Not applicable.

Authors' contributions

O.R. conceived the study. O.R., S.B.P., B.B., B.L. and L.D. designed the chemical and electrophysiological

experiments. O.R. and S.B.P. performed chemical extractions and SBP and BB performed the chemical analysis.

S.B.P. and B.L. performed the electrophysiological experiments. O.R. and S.B.P. designed olfactometric

experiments and S.B.P. performed data collection. S.B.P. and O.R. performed statistical analyses. S.B.P., O.R.,

B.B., B.L. drafted the manuscript and L.D., O.G., A.D, and R.K.D critically revised the manuscript. All authors

revised the manuscript, gave final approval for publication and are accountable for the work performed therein.

Acknowledgements

We thank David Sanou for his assistance in the field, Stephane Somda, Sofan Somé, Bethsabee Scheid, Marie

Rossignol and Carole Ginibre for their help in mosquito rearing and Heidi Lançon for proofreading the paper.

607   REFERENCES

Alexander RD, Marshall DC, Cooley JR (1997) Evolutionary perspectives on insect mating. In: Choe JC, Crespi

BJ (eds) The evolution of mating systems in insects and Arachnids. Cambridge University Press, pp 4–31.

Andersson J, Borg-Karlson AK, Vongvanich N, Wiklund C (2007) Male sex pheromone release and female mate

choice in a butterfly. J Exp Biol 210:964–970.

Bicchi C, Cordero C, Iori C, et al (2000) Headspace Sorptive Extraction (HSSE) in the headspace analysis of

aromatic and medicinal plants. J High Resolut Chromatogr 23:539–546.

Bimbilé Somda NS, Poda BS, Sawadogo PS, et al (2018) Ecology of reproduction of *Anopheles arabiensis* in an

urban area of Bobo-Dioulasso, Burkina Faso (West Africa): monthly swarming and mating frequency and

their relation to environmental factors. PLoS One 13:e0205966.

Bjostad LB, Gaston LK, Shorey HH (1980) Temporal pattern of sex pheromone release by female *Trichoplusia*

ni. J Insect Physiol 26:493–498.

Bray DP, Alves GB, Dorval ME, et al (2010) Synthetic sex pheromone attracts the leishmaniasis vector

Lutzomyia longipalpis to experimental chicken sheds treated with insecticide. Parasit Vectors 3:1–11.

Cainap C, Pop LA, Balacescu O, Cainap SS (2020) Early diagnosis and screening in lung cancer. Am J Cancer

Res 10:1993–2009.

Calenic B, Amann A (2014) Detection of volatile malodorous compounds in breath: current analytical

techniques and implications in human disease. Bioanalysis 6:357–376.

Charlwood JD, Jones MDR (1980) Mating in the mosquito, *Anopheles gambiae s.l.* II. Swarming behaviour.

Physiol Entomol 5:315–320.

Charlwood JD, Pinto J, Sousa CA, et al (2002) The swarming and mating behaviour of *Anopheles gambiae s.s.*

(Diptera: Culicidae) from São Tomé Island. J Vector Ecol 27:178–183.

Charpentier MJE, Barthes N, Proffit M, et al (2012) Critical thinking in the chemical ecology of mammalian

communication: roadmap for future studies. Funct Ecol 26:769–774.

Clements AN (1999) The biology of mosquitoes. Volume 2: sensory, reception and behaviour. CABI Publishing,

Wallingford, UK

Clements AN (1992) The biology of mosquitoes: development, nutrition and reproduction. Chapman & Hall,

London

Costantini C, Ayala D, Guelbeogo WM, et al (2009) Living at the edge: biogeographic patterns of habitat

segregation conform to speciation by niche expansion in *Anopheles gambiae*. BMC Ecol 9:1–27.

Dao A, Adamou A, Yaro AS, et al (2008) Assessment of alternative mating strategies in *Anopheles gambiae*:

does mating occur indoors? J Med Entomol 45:643–52.

Dekel A, Yakir E, Bohbot JD (2019) The sulcatone receptor of the strict nectar-feeding mosquito *Toxorhynchites*

amboinensis. Insect Biochem Mol Biol 111:103174.

della Torre A, Fanello C, Akogbeto M, et al (2001) Molecular evidence of incipient speciation within *Anopheles*

gambiae s.s. in West Africa. Insect Mol Biol 10:9–18.

della Torre A, Tu Z, Petrarca V (2005) On the distribution and genetic differentiation of *Anopheles gambiae s.s.*

molecular forms. Insect Biochem Mol Biol 35:755–769.

Diabaté A, Baldet T, Brengues C, et al (2003) Natural swarming behaviour of the molecular M form of

Anopheles gambiae. Trans R Soc Trop Med Hyg 97:713–716.

Diabaté A, Brengues C, Baldet T, et al (2004) The spread of the Leu-Phe *kdr* mutation through *Anopheles*

gambiae complex in Burkina Faso: genetic introgression and de novo phenomena. Trop Med Int Heal

9:I267–I273.

Diabaté A, Dabiré KR, Kim HE, et al (2005) Larval development of the molecular forms of *Anopheles gambiae*

(Diptera: *Culicidae*) in different habitats: a transplantation experiment. J Med Entomol 42:548–553

Diabaté A, Dabiré KR, Millogo N, Lehmann T (2007) Evaluating the effect of postmating isolation between

molecular forms of *Anopheles gambiae* (Diptera: *Culicidae*). J Med Entomol 44:60–64.

Diabaté A, Dabire RK, Kengne P, et al (2006) Mixed swarms of the molecular M and S forms of *Anopheles*

gambiae (Diptera: *Culicidae*) in sympatric area from Burkina Faso. J Med Entomol 43:480–483.

Diabaté A, Dao A, Yaro AS, et al (2009) Spatial swarm segregation and reproductive isolation between the

molecular forms of *Anopheles gambiae*. Proc R Soc B Biol Sci 276:4215–4222.

Diabaté A, Yaro AS, Dao A, et al (2011) Spatial distribution and male mating success of *Anopheles gambiae*

swarms. BMC Evol Biol 11:184.

Dormont L, Bessière J-M, Cohuet A (2013a) Human Skin Volatiles: A Review. J Chem Ecol 39:569–578.

Dormont L, Bessiere JM, McKey D, Cohuet A (2013b) New methods for field collection of human skin volatiles

and perspectives for their application in the chemical ecology of human-pathogen-vector interactions. J

Exp Biol 216:2783–2788.

Downes JA (1969) The swarming and mating flight of Diptera. Annu Rev Entomol 14:271–298.

Facchinelli L, Valerio L, Lees RS, et al (2015) Stimulating *Anopheles gambiae* swarms in the laboratory:

application for behavioural and fitness studies. Malar J 14:271.

Feugère L, Gibson G, Manoukis NC, Roux O (2021) Mosquito sound communication: are male swarms loud

enough to attract females? J R Soc Interface 18:20210121.

Filipiak W, Filipiak A, Sponring A, et al (2014) Comparative analyses of volatile organic compounds (VOCs)

from patients, tumors and transformed cell lines for the validation of lung cancer-derived breath markers. J

Breath Res 8:027111.

Gibson G (1985) Swarming behaviour of the mosquito *Culex pipiens quinquefasciatus*: a quantitative analysis.

Physiol Entomol 10:283–296.

Gibson G, Warren B, Russell IJ (2010) Humming in tune: sex and species recognition by mosquitoes on the

wing. JARO - J Assoc Res Otolaryngol 11:527–540.

Gibson NHE (1945) On the mating swarms of certain *Chironomidae* (Diptera). Trans R Entomol Soc London

95:263–294.

Gomulski L (1988) Aspects of mosquito mating behaviour. University of London.

Guidobaldi F, May-Concha IJ, Guerenstein PG (2014) Morphology and physiology of the olfactory system of

blood-feeding insects. J. Physiol. Paris 108:96–111.

Hahn MW, White BJ, Muir CD, Besansky NJ (2012) No evidence for biased co-transmission of speciation

Islands in *Anopheles gambiae*. Philos Trans R Soc B Biol Sci 367:374–384.

Howell PI, Knols BG (2009) Male mating biology. Malar J 8:S8.

Iatrou K, Biessmann H (2008) Sex-biased expression of odorant receptors in antennae and palps of the African

malaria vector *Anopheles gambiae*. Insect Biochem Mol Biol 38:268–274.

Kelly DW, Dye C (1997) Pheromones, kairomones and the aggregation dynamics of the sandfly *Lutzomyia*

longipalpis. Anim Behav 53:721–731.

Kostiainen R (1995) Volatile organic compounds in the indoor air of normal and sick houses. Atmos Environ

29:693–702.

Kruza M, Lewis AC, Morrison CG, Carslaw N (2017) Impact of surface ozone interactions on indoor air

chemistry: A modeling study. Indoor Air 27:1001–1011.

Lefèvre T, Gouagna LC, Dabire KR, et al (2009) Evolutionary lability of odour-mediated host preference by the

malaria vector *Anopheles gambiae*. Trop Med Int Heal 14:228–236.

Lefèvre T, Gouagna LC, Dabiré KR, et al (2010) Beer consumption increases human attractiveness to malaria

mosquitoes. PLoS One 5:e9546.

Lehmann T, Diabaté A (2008) The molecular forms of *Anopheles gambiae*: a phenotypic perspective. Infect

Genet Evol 8:737–746.

Levi-Zada A, Fefer D, David M, et al (2014) Diel periodicity of pheromone release by females of *Planococcus*

citri and *Planococcus ficus* and the temporal flight activity of their conspecific males. Naturwissenschaften

101:671–678.

Lu T, Qiu YT, Wang G, et al (2007) Odor coding in the maxillary palp of the malaria vector mosquito *Anopheles*

gambiae. Curr Biol 17:1533–1544.

Maïga H, Dabiré RK, Lehmann T, et al (2012) Variation in energy reserves and role of body size in the mating

system of *Anopheles gambiae*. J Vector Ecol 37:289–297.

Maïga H, Niang A, Sawadogo SP, et al (2014) Role of nutritional reserves and body size in *Anopheles gambiae*

males mating success. Acta Trop 132S:S102–S107.

Marchand RP (1984) Field observations on swarming and mating in *Anopheles gambiae* mosquitoes in Tanzania.

Netherlands J Zool 34:367–387.

McBride CS, Baier F, Omondi AB, et al (2014) Evolution of mosquito preference for humans linked to an

odorant receptor. Nature 515:222–227.

Merlin C, Lucas P, Rochat D, et al (2007) An antennal circadian clock and circadian rhythms in peripheral

pheromone reception in the moth *Spodoptera littoralis*. J Biol Rhythms 22:502–514.

Mosqueira B, Soma DD, Namountougou M, et al (2015) Pilot study on the combination of an organophosphate-

based insecticide paint and pyrethroid-treated long lasting nets against pyrethroid resistant malaria vectors

in Burkina Faso. Acta Trop 148:162–169.

Mozūraitis R, Hajkazemian M, Zawada JW, et al (2020) Male swarming aggregation pheromones increase female attraction and mating success among multiple African malaria vector mosquito species. Nat Ecol Evol 4:1395–1401.

Nguyen PL, Vantaux A, Hien DFS, et al (2017) No evidence for manipulation of *Anopheles gambiae*, *An. coluzzii* and *An. arabiensis* host preference by *Plasmodium falciparum*. Sci Rep 7:9415.

Niang A, Nignan C, Poda BS, et al (2019) Semi-field and indoor setups to study malaria mosquito swarming behavior. Parasit Vectors 12:446.

Nyasembe VO, Tchouassi DP, Kirwa HK, et al (2014) Development and assessment of plant-based synthetic odor baits for surveillance and control of malaria vectors. PLoS Negl Trop Dis 9:e89818.

Pandey SK, Kim K (2011) Human body-odor components and their determination. Trends Anal Chem 30:784–796.

Pennetier C, Warren B, Dabiré KR, et al (2010) "Singing on the wing" as a mechanism for species recognition in the malarial mosquito *Anopheles gambiae*. Curr Biol 20:131–136.

Persiani A, Dideco MA, Petrangeli G (1986) Osservzioni di laboratorio su polimorfismi da inversione originati da incroci tra popolazioni diverse di *Anopheles gambiae s.s.* Ann Dell'Istituto Super Di Sanita 22:221–224.

Pitts RJ, Rinker DC, Jones PL, et al (2011) Transcriptome profiling of chemosensory appendages in the malaria vector *Anopheles gambiae* reveals tissue- and sex-specific signatures of odor coding. BMC Genomics 12:271.

Poda SB, Buatois B, Lapeyre B, et al (2020) No evidence for long-range male sex pheromones in two malaria mosquitoes. bioRxiv Prepr.

Poda SB, Nignan C, Gnankiné O, et al (2019) Sex aggregation and species segregation cues in swarming mosquitoes: Role of ground visual markers. Parasit Vectors 12:589.

Poda SB, Soma DD, Hien A, et al (2018) Targeted application of an organophosphate-based paint applied on windows and doors against *Anopheles coluzzii* resistant to pyrethroids under real life conditions in Vallée du Kou, Burkina Faso (West Africa). Malar J 17:136.

Poli D, Goldoni M, Corradi M, et al (2010) Determination of aldehydes in exhaled breath of patients with lung cancer by means of on-fiber-derivatisation SPME-GC/MS. J Chromatogr B 878:2643–2651.

Pombi M, Kengne P, Gimonneau G, et al (2017) Dissecting functional components of reproductive isolation among closely related sympatric species of the *Anopheles gambiae* complex. Evol Appl 00:1–19.

Robledo N, Arzuffi R (2012) Influence of host fruit and conspecifics on the release of the sex pheromone by

Toxotrypana curvicauda males (Diptera: *Tephritidae*). Environ Entomol 41:387–391.

Rund SSC, Bonar NA, Champion MM, et al (2013) Daily rhythms in antennal protein and olfactory sensitivity in

the malaria mosquito *Anopheles gambiae*. Sci Rep 3:2494.

Santolamazza F, Mancini E, Simard F, et al (2008) Insertion polymorphisms of SINE200 retrotransposons within

speciation islands of *Anopheles gambiae* molecular forms. Malar J 7:163.

Savolainen E (1978) Swarming in *Ephemeroptera*: the mechanism of swarming and the effects of illumination

and weather. Ann Zool Fennici 15:17–52.

Sawadogo PS, Costantini C, Pennetier C, et al (2013) Differences in timing of mating swarms in sympatric

populations of *Anopheles coluzzii* and *Anopheles gambiae s.s.* (formerly *An. gambiae* M and S molecular

forms) in Burkina Faso, West Africa. Parasit Vectors 6:275.

Sawadogo PS, Namountougou M, Toé KH, et al (2014) Swarming behaviour in natural populations of *Anopheles*

gambiae and *An. coluzzii*: review of 4 years survey in rural areas of sympatry, Burkina Faso (West Africa).

Acta Trop 130:24–34.

Shelly TE, Whittier TS (1997) Lek behavior of insects. In: Choe JC, Crespi BJ (eds) The evolution of mating

systems in insects and Arachnids. Cambridge University Press, New York, pp 273–293.

Sivinski JM, Petersson E (1997) Mate choice and species isolation in swarming insects. In: Choe JC, Crespi BJ

(eds) The evolution of mating systems in insects and Arachnids. Cambridge University Press, New York,

pp 294–309.

Souto-Vilarós D, Proffit M, Buatois B, et al (2018) Pollination along an elevational gradient mediated both by

floral scent and pollinator compatibility in the fig and fig-wasp mutualism. J Ecol 106:2256–2273.

Suh E, Choe D, Saveer AM, Zwiebel LJ (2016) Suboptimal larval habitats modulate oviposition of the malaria

vector mosquito *Anopheles coluzzii*. PLoS One 11:e0149800.

Tchouassi DP, Sang R, Sole CL, et al (2013) Common host-derived chemicals increase catches of disease-

transmitting mosquitoes and can improve early warning systems for rift valley fever virus. PLoS Negl

Trop Dis 7:e2007.

Tienpont B, David F, Bicchi C, Sandra P (2000) High capacity headspace sorptive extraction. J Microcolumn

Sep 12:577–584.

Tripet F, Dolo G, Traoré S, Lanzaro GC (2004) The "wingbeat hypothesis" of reproductive isolation between

members of the *Anopheles gambiae complex* (Diptera: *Culicidae*) does not fly. J Med Entomol 41:375–

384.

Tripet F, Touré YT, Taylor CE, et al (2001) DNA analysis of transferred sperm reveals significant levels of gene flow between molecular forms of *Anopheles gambiae*. Mol Ecol 10:1725–1732.

Vantaux A, Hien DFS, Yameogo B, et al (2015) Host-seeking behaviors of mosquitoes experimentally infected with sympatric field isolates of the human malaria parasite *Plasmodium falciparum*: no evidence for host manipulation. Front Ecol Evol 3:86.

Verhulst NO, Andriessen R, Groenhagen U, et al (2010) Differential attraction of malaria mosquitoes to volatile blends produced by human skin bacteria. PLoS One 5:e15829.

Wang G, Diabaté A, Liu J, et al (2021) Clock genes and environmental cues coordinate *Anopheles* pheromone synthesis, swarming, and mating. Science 371(6527):411–415.

Wondwosen B, Birgersson G, Seyoum E, et al (2016) Rice volatiles lure gravid malaria mosquitoes, *Anopheles arabiensis*. Sci Rep 6:37930.

Wondwosen B, Birgersson G, Tekie H, et al (2018) Sweet attraction: sugarcane pollen-associated volatiles attract gravid *Anopheles arabiensis*. Malar J 17:90.

Zellner B d'Acampora, Bicchi C, Dugo P, et al (2008) Linear retention indices in gas chromatographic analysis: a review. Flavour Fragr J 23:297–314.

Figure legends

Fig. 1 Schematic of the olfactometer set up. S: swarm; M: marker. Not at scale.

Fig. 2 Mosquito activation rate, expressed as the proportion of females caught in both collecting boxes out of the

total number released females for each of the four tested combinations. Different letters indicate difference at

$P<0.05$.

Fig. 3 Mosquito choice, expressed as the proportion of females caught in one or the other collecting box out of

the total number of "activated" females for each test.

Fig. 4 Quantities (integration units) of acetoin, sulcatone, octanal, nonanal and decanal contained in the

headspace of the empty bottles, the breath of the manipulator and flying males (A) and females (B) of *Anopheles*

gambiae Kisumu and collected with SPME fibers. The intensity values correspond to the counts related to the

abundance of the specific ions representative of each molecule formed in the mass spectrometer and correspond

to the amount of compound analyzed. The box plots indicate the median (wide horizontal bars), the 25th and

75th percentiles (squares), and the minimum and maximum values (whiskers). The black dots represent outliers

and grey dots raw data. NS=not-significant; * =$P < 0.05$; ** =$P < 0.01$; *** =$P < 0.001$.

Fig. 5 Quantities (integration units) of sulcatone, octanal, nonanal and decanal contained in a natural swarm

sampled with twisters (A); in the headspace of the empty box, swarming *Anopheles coluzzii* males and *An.*

gambiae males sampled with Tenax/Carbotrap tubes (B); or collected with Porapack tubes (C). The intensity

values correspond to the counts related to the abundance of the specific ions representative of each molecule

formed in the mass spectrometer and correspond to the amount of compound analyzed. The box plots indicate

the median (wide horizontal bars), the 25th and 75th percentiles (squares), and the minimum and maximum

values (whiskers). The black dots represent outliers and grey dots raw data. NS=not-significant; * =$P < 0.05$; **

=$P < 0.01$; *** =$P < 0.001$. NA: The analytical procedure for these analyses did not allow detection of acetoin.

Fig. 6 Mean electrophysiological responses of female *An. coluzzii* antennae to male swarm solvent extracts,

recorded by coupled gaz chromatography – electroantennography detection (GC-EAD). Dashed lines indicate

retention times (and related LRI) for which the most significant EAG responses were recorded. Only 2 females

in 8 tested produced this pattern.

Table 1: Air inlet and outlet rates, and VOC extraction duration according to the type of trap used.

	Air flow rate (L/min)		VOCs extraction duration (hour)
	Air inlet	Air outlet	
Tenax/Carbotrap tube	2.5	2.0	2.0
Porapak-Q VCT tube	1.5	1.0	2.5
Micro-trap	0.9	0.4	5.0

Fig. S1 Test for olfactometer symmetry. Female choice, expressed as the proportion of female mosquitoes

caught in one or the other collecting box out of the total number retrieved from both collecting boxes for the test

with empty boxes (control *vs*. control combination).

Fig. S2 A. Chromatograms obtained with eluted Porapak extracts from headspaces of the control, an *An. coluzzii*

swarm and an *An. gambiae* swarm (linear retention index (LRI) 800 to 1700). **B.** Enlarged view of the retention

time (RT) interval containing the LRI at which *An. coluzzii* antennal responses were detected in GC-EAD. Doted

white lines correspond to the theoretical RT of Sulcatone, octanal, nonanal and decanal. No difference was

detected between the control and the swarm extracts. All the peaks correspond to emissions born from the

material.

Fig. S3 A. Chromatograms obtained with the desorption of twisters obtained with the static sorptive extraction

methods under natural conditions in the field (linear retention index (LRI) 800 to 1700). The twister was either

located within a non-enclosed swarm of *An. coluzzii* or 3 m away upwind (control). **B.** Enlarged view of the

retention time (RT) interval containing the LRI at which *An. coluzzii* antennal responses were detected in GC-

EAD. Doted white lines correspond to the theoretical RT of sulcatone, octanal, nonanal and decanal. When in

italic, the compound was absent. No difference was detected between the control and the swarm extracts. All the

peaks correspond to environmental emissions.

Fig. S4 Quantities (integration units) of acetoin, sulcatone, octanal, nonanal and decanal contained in the headspace of the empty bottles, the breath of the manipulator and the breath of the manipulator after blowing for the second time (breath x2) sampled with SPME fibers. The intensity values correspond to the counts related to the abundance of the specific ions representative of each molecule formed in the mass spectrometer and correspond to the amount of compound analyzed. The box plots indicate the median (wide horizontal bars), the 25th and 75th percentiles (squares), and the minimum and maximum values (whiskers). The black dots represent outliers and grey dots raw data. NS=not-significant; * =$P < 0.05$; ** =$P < 0.01$.

An. coluzzii females

Control	An. gambiae swarm	ns
Control	An. coluzzii swarm	ns
An. gambiae swarm	An. coluzzii swarm	ns

An. gambiae females

Control	An. gambiae swarm	ns
Control	An. coluzzii swarm	ns
An. coluzzii swarm	An. gambiae swarm	ns

Treatment combinations

Female choice (95%CI)