3 # The ER membrane chaperone Shr3 acts in a progressive manner to

4 assist the folding of related plasma membrane transport proteins

Ioanna Myronidi[§], Andreas Ring[§] and Per O. Ljungdahl*

Department of Molecular Biosciences, The Wenner-Gren Institute, SciLifeLab, Stockholm

University, SE-106 91, Sweden

Running title: Membrane-localized chaperone substrate interactions

Keywords: Shr3; membrane-localized chaperone; amino acid permease; polytopic membrane

protein folding; endoplasmic reticulum; *Saccharomyces cerevisiae*

[§] Contributed equally, the names are listed in alphabetic order.

* Corresponding author. Mailing address: SciLifeLab, Box 1031, SE- 171 21 Solna, Sweden.

Phone: 46 8 16 41 01. E-mail: per.ljungdahl@scilifelab.se.

Abstract

Proteins with multiple membrane-spanning segments (MS) co-translationally insert into the endoplasmic reticulum (ER) membrane of eukaryotic cells. In *Saccharomyces cerevisiae*, Shr3 is an ER membrane-localized chaperone that is specifically required for the functional expression of amino acid permeases (AAP), a family of eighteen transporters comprised of 12 MS. Here, comprehensive scanning mutagenesis and deletion analysis of Shr3, combined with a modified split-ubiquitin approach, were used to probe chaperone-substrate interactions with seven different AAP *in vivo*. Our findings indicate that Shr3 specifically recognizes AAP substrates, largely independent of sequence-specific interactions involving membrane and luminally oriented domains. Shr3 selectively and robustly interacts with nested C-terminal AAP truncations in marked contrast to similar truncations of non-Shr3 substrate polytopic sugar transporters. Strikingly, Shr3-AAP interactions initiate with the first 4 MS of AAP and successively strengthen, but abruptly weaken when all 12 MS are present. The data are consistent with Shr3 acting in a temporal manner as a scaffold preventing AAP translation intermediates from engaging in non-productive interactions.

Introduction

The integration and concomitant folding of membrane proteins in the lipid bilayer of the endoplasmic reticulum (ER) are critical steps in the biogenesis of transport proteins destined to function at the plasma membrane (PM). Most eukaryotic membrane proteins are co-translationally inserted into the ER membrane via the Sec61-complex, also known as the translocon. The translocon forms a protein-conducting channel that mediates protein translocation and the co-translational partitioning of membrane-spanning segments (MS) (Johnson and van Waes, 1999; Rapoport et al., 2017; Seinen and Driessen, 2019). During the synthesis of complex polytopic membrane proteins comprised of multiple MS, each MS sequentially exits the channel and partitions into the ER membrane via the lateral gate of the translocon. Although the central channel of the translocon is too small to accommodate multiple MS, the translocon appears to have a limited capacity to promote the folding of membrane proteins comprised of up to three MS; two separate extra-channel MS-binding sites have been reported to act in a chaperone-like manner to delay the release of N-terminal MS until translation is completed (Hou et al., 2012). However, as the complexity of membrane proteins grows beyond three MS, the challenge of preventing inappropriate interactions between incompletely translated nascent chains apparently exceeds the chaperone-like activity of the translocon. Specifically, during the translation of complex polytopic membrane proteins, e.g., amino acid permeases with 12 MS, the N-terminal MS partition into the membrane prior to the synthesis and partitioning of C-terminal MS. To prevent the MS of translation intermediates from entering non-productive folding pathways, discrete and highly specialized ER resident

membrane proteins have been described in fungi that prevent misfolding of specific families of
polytopic membrane proteins (Kota and Ljungdahl, 2005; Lau et al., 2000; Ljungdahl et al.,
1992; Luo et al., 2002; Martínez and Ljungdahl, 2000; 2004; Sherwood and Carlson, 1999;
Shurtleff et al., 2018).

Shr3, the most comprehensively studied of these specialized ER components, was
identified as an integral membrane protein required for the functional expression of the
conserved family of amino acid permeases (AAP) (Ljungdahl *et al.*, 1992). The AAP family in
Saccharomyces cerevisiae, belonging to the Amino acid-Polyamine-Organocation (APC)
super-family of transporters (Gilstring and Ljungdahl, 2000; Jack et al., 2000; Saier, 2000;
Wong et al., 2012), is comprised of 18 genetically distinct but structurally similar proteins with
12 MS. Shr3 is composed of 210 amino acids organized into two functional domains; a N-
terminal membrane domain comprised of four hydrophobic α-helices connected by three
hydrophilic loops and a hydrophilic cytoplasmically oriented C-terminal domain. Initially, Shr3
was recognized as an essential factor facilitating the packaging of AAP into ER-derived
secretory vesicles (Kuehn et al., 1996), and it was found to be important for the proper
presentation of ER-exit motifs, located within the hydrophilic C-terminal tails of AAP, to the
inner COPII coatomer subunit Sec24 (Kuehn et al., 1998; Malkus et al., 2002; Miller et al.,
2002; Miller et al., 2003). Additional data regarding the packaging activity of Shr3 included
genetic and physical interactions of Shr3 with components of the COPII-coated vesicles
(Gilstring et al., 1999). Also, it was shown that Shr3 facilitated ER-vesicle formation in close
proximity to fully integrated and folded AAP (Gilstring and Ljungdahl, 2000; Gilstring *et al.*,
1999) and hence Shr3 was designated a packaging chaperone. The ability of Shr3 to interact
with COPII components is primarily linked to its hydrophilic C-terminal tail. Consistently, Shr3
was found to associate with newly synthesized Gap1 in a transient manner, the interaction was
reported to exhibit a half-life of approximately 15 min. The observed association with Gap1
was initially thought to occur post-translationally since Gap1 fully integrates into the ER-
membrane with each MS correctly oriented independently of Shr3 (Gilstring and Ljungdahl,
2000). Also, the AAP that accumulate in the ER of *shr3Δ* strains do not activate the unfolded
protein stress response (UPR) (Gilstring *et al.*, 1999).

However, subsequent studies revealed that Shr3 has an important function that is separate
from and precedes its packaging function. Consequently, the view of Shr3 evolved from being
a packaging into a specialized membrane-localized chaperone that interacts early with substrate
AAP during their co-translational insertion into the ER membrane (Kota et al., 2007; Kota and
Ljungdahl, 2005). In these studies, Shr3 was found to prevent the aggregation of AAP in the
ER-membrane, a function associated with its N-terminal membrane domain (Kota and
Ljungdahl, 2005). Critical evidence demonstrating the importance of Shr3 in facilitating the
folding of AAP includes the finding that co-expressed split N- and C-terminal portions of Gap1
assemble into a functional permease in a Shr3-dependent manner (Kota *et al.*, 2007). The

membrane domain of Shr3 is required and suffices to prevent aggregation of the first five MS
of Gap1 enabling productive folding interactions with the C-terminal portions of Gap1. Similar
to full-length Gap1, the N-terminal fragment displays an increased propensity to aggregate in
membranes isolated from cells lacking Shr3, and importantly, its aggregation appeared not to
be affected by the presence or absence of the C-terminal fragment. In marked contrast, the
aggregation status of the C-terminal fragment was dependent on the presence of both Shr3 and
the N-terminal fragment. Since the N- and C-terminal fragments individually insert into the
membrane, Shr3 apparently can maintain the N-terminal fragment in a conformation that
enables the C-terminal fragment to interact and assemble with it. Although direct physical
interactions between Shr3 and substrate AAP have not been demonstrated, all available data is
consistent with Shr3 interacting early with N-terminal MS as they co-translationally partition
into the ER membrane.

During translocation, exclusively hydrophobic MS partition readily into the lipid phase of
the membrane, whereas less hydrophobic MS containing (Heinrich et al., 2000) charged or
polar residues partition into the membrane less readily and are retained in proximity to the
translocon or to translocon associated proteins, e.g., TRAM (Heinrich and Rapoport, 2003). In
analogy to TRAM, we posited that Shr3 facilitates the partitioning of MS of AAP containing
charged or polar amino acid residues as they emerge from the translocon. According to this
hypothesis, Shr3 may physically shield charged or polar residues within MS, thereby preventing
these thermodynamically challenging segments from engaging in nonproductive interactions
(Kota *et al.*, 2007). More recent findings in yeast regarding the conserved ER membrane protein
complex (EMC) have been interpreted in a similar manner (Miller-Vedam et al., 2020; Shurtleff
et al., 2018). However, a striking difference between Shr3 and EMC function is that null alleles
of *SHR3* do not activate the UPR (Gilstring *et al.*, 1999).

Despite the clear requirement of Shr3 in AAP biogenesis, we currently lack critical
information regarding the mechanisms underlying Shr3 function. Here we have focused on the
membrane domain of Shr3 and employed a comprehensive scanning mutagenesis approach to
define amino acid residues involved in recognizing AAP substrates. Further, we have exploited
a split-ubiquitin approach to directly probe and characterize interactions with seven different
AAP substrates *in vivo*. The data support Shr3 acting as a MLC providing a scaffold-like
structure to help nascent chains of partially translated AAP maintain a structure required to
enter and follow a productive folding pathway as translation proceeds to completion.

Results

Systematic scanning mutagenesis of MS within the membrane domain of Shr3

We have previously shown that the membrane domain of Shr3 is required and sufficient for
facilitating the folding of AAPs (Kota and Ljungdahl, 2005). Here, a systematic scanning
mutagenesis approach was used to identify residues within the Shr3 membrane domain required
for function. Intramembrane residues were mutated to leucine; the length of consecutive
substitution mutations varied, ranging from 2 to 13 residues. The extramembrane residues
within ER lumenal loops L1 and L3 and cytoplasmic oriented NT and loop L2 where mutated
to alanine; the length of consecutive alanine replacements ranged from 2 to 3.

The biological activity of the mutant proteins was initially assessed using growth-based
assays on YPD supplemented with metsulfuron-methyl (MM), which provides a sensitive
measure of Shr3 function. MM targets and inhibits branched-chain amino acid synthesis and
growth is strictly dependent on the combined activity of multiple SPS-sensor regulated AAP
that facilitate high-affinity isoleucine, leucine and valine uptake (Andréasson and Ljungdahl,
2002; Jørgensen et al., 1998). Serial dilutions of cell suspensions from strain JKY2 (*shr3Δ*)
carrying vector control (VC), *SHR3* or the *shr3* mutant alleles were spotted on YPD and
YPD+MM plates (Supplementary Material Fig. S1 – S10). Only three mutant alleles, *shr3-35*,
shr3-50 and *shr3-76*, failed to support growth (Fig. 1A and B). The steady state levels of the
three mutant proteins were similar to wildtype Shr3 (Fig. 1B), suggesting that the mutant
proteins were not grossly misfolded, and consequently, not prematurely targeted for ER-
associated degradation.

The *shr3-35* allele encodes a protein with residues 17 through 19 (serine-alanine-threonine)
in MS I replaced by leucine (Fig. 1A). To more precisely define the critical residues, we
constructed additional mutant alleles with paired leucine substitutions at residues 17-18 (LLT),
18-19 (SLL), and 17 and 19 (LAL). Expression of *SHR3-36* (LLT) and *SHR3-38* (LAL) alleles
complemented *shr3Δ* and supported wildtype growth on YPD + MM (Fig. 1C, dilutions 4 and
6). By contrast, cells expressing *shr3-37* (SLL) did not complement, exhibiting a phenotype
similar to the *shr3-35* (LLL) mutant (Fig. 1C, dilutions 3 and 5).

The *shr3-76* allele carries leucine replacements at residues 139 through 142 (serine-
asparagine-isoleucine-isoleucine) in MS IV (Fig. 1A). Again, the importance of the affected
residues was tested by creating alleles with paired leucine substitutions at positions 139-140
(LLII) and 141-142 (SNLL). Cells expressing *shr3-77* (LLII) allele grew poorly, although
slightly better than cells expressing the *shr3-76* allele (Fig. 1C, dilutions 15 and 16). Cells
expressing *SHR3-78* (SNLL) grew as wildtype *SHR3* (Fig. 1C, dilutions 14 and 17).

The third non-functional allele, *shr3-50*, encodes a mutant protein with alanine
substitutions at residues 51 through 53 (leucine-arginine-histidine) located within the ER
lumenal loop L1 (Fig. 1A). The importance of these residues was tested by paired alanine
substitutions of the positions 51-52 (AAH), 52-53 (LAA), and 51 and 53 (ARA). Expression of
these alleles demonstrated that *SHR3-51* (AAH) and *SHR3-52* (LAA) complemented *shr3Δ*

similar to wildtype *SHR3* (Fig.1C, dilutions 8, 10 and 11). The *shr3-53* (ARA) allele exhibited

reduced growth (Fig. 1C, dilution 12), indicative of compromised function. The finding that

mutations affecting residues 53-55 in the ER lumenal loop L1 abolish function suggests that

extramembrane sequences are important for guiding the folding of AAP sequences destined to

be oriented towards the extracellular side of the PM.

Deletion analysis of ER-lumen oriented loops

The possibility that Shr3 engages and interacts with its substrate AAP through contacts with

extramembrane sequences prompted us to specifically test the functional significance of loops

L1 and L3 (Fig. 2A). The residues 44-57 within L1 are predicted to fold into an amphipathic α-

helix (Fig. 2B). We constructed four internal deletions in L1 that affect this secondary structure

motif to varying extent: *shr3Δ90* (Δ34-48); *shr3Δ91* (Δ39-47); *shr3Δ92* (Δ44-54); and *shr3Δ93*

(Δ55-60). Also, an internal deletion in L3 was constructed: *SHR3Δ94* (Δ121-127). The five

deletion alleles directed the expression of mutant proteins at levels comparable to *SHR3* (Fig.

2C); the deletions do not decrease the steady state levels of protein. The function of these

deletion alleles was assessed as before using YPD + MM. We also extended the analysis with

more nuanced growth-based assays capable of monitoring amino acid uptake catalyzed

predominantly by a single or a couple of AAP. This was accomplished by examining growth

on minimal media individually supplemented with toxic amino acid analogues D-histidine, L-

canavanine, and azetidine-2-carboxylate (AzC), which are taken up by Gap1 (Gresham et al.,

2010), Can1 (Ono et al., 1983), Agp1/Gnp1 (Andréasson et al., 2004), respectively. The

expression of functional alleles of *SHR3* results in impaired growth in the presence of these

toxic analogues. Serial dilutions of cell suspensions from strain JKY2 (*shr3Δ*) carrying vector

control (VC), *SHR3*, *shr3Δ90*, *shr3Δ91*, *shr3Δ92*, *shr3Δ93* or *SHR3Δ94* were spotted on SAD

containing D-histidine, SD + L-canavanine, SD + AzC, and YPD + MM plates. The four

internal deletion alleles affecting L1 failed to complement *shr3Δ*; the strains grew similar as

the VC (Fig. 2C, dilutions 1, 3, 4, 5 and 6). By contrast, the strain expressing the internal

deletion in L3 showed a more complex pattern of growth. On SAD + D-histidine and SD + L-

canavanine, *SHR3Δ94* appeared to express a non-functional protein, the strain grew in the

presence of these toxic amino acid analogues (Fig. 2C, compare dilution 1 with 7). However,

on SD + AzC and YPD + MM, the *SHR3Δ94* allele exhibited growth similar to wildtype *SHR3*

(Fig. 2C, compare dilution 2 with 7). Note that on YPD + MM the strain carrying *SHR3Δ94*

exhibited enhanced growth compared to wildtype, and consequently, on YPD + MM we

designated the phenotype WT[+].

Lumenal loop L3 influences substrate specificity

The finding that *SHR3Δ94* exhibited a range of phenotypes, i.e., from null to apparently
enhanced functionality, prompted us to reexamine the growth characteristics of the 44 leucine-
and alanine-scanning mutant alleles using the more nuanced growth-based assays. Serial
dilutions of cell suspensions from strain JKY2 (*shr3Δ*) carrying vector control (VC), *SHR3* or
one of the individual mutant alleles were spotted on SAD + D-histidine, SD + L-canavanine,
SD + AzC and YPD+MM (Supplementary Material Fig. S1-S10). The growth characteristics
were evaluated and the results, including the internal loop deletions, are summarized in an
ordered heat-map (Fig. 3A). As was found in the initial evaluation of growth on YPD+MM,
most of the mutant alleles were judged to encode functional proteins; the strains grew similarly
as the strain carrying the *SHR3* wildtype control. Reevaluation of the three non-functional
alleles, *shr3-35*, *shr3-50* and *shr3-76*, confirmed that the mutations exhibit major defects on all
of the selective media (Fig. 3A). However, in some instances, several of the mutations, similar
to *SHR3Δ94*, conferred robust growth on YPD+MM but did not complement *shr3Δ* on the other
selective media. Strikingly, *SHR3-45*, *-63*, *-65*, *-68*, *-71*, *-74*, and *-75*, supported more robust
growth on YPD + MM than *SHR3*, but exhibited a null phenotype on media containing toxic
amino acid analogues. In summary, growth in the presence of D-histidine was found to be the
most sensitive monitor of mutations in *SHR3*, perhaps due to the fact that D-amino acids are
taken up by a single AAP, Gap1 (Grenson et al., 1970; Rytka, 1975). We note that mutations
localized to the MS III and IV and the ER lumenal oriented loop L3 exhibited the most
pleiotropic affects, suggesting that these regions of Shr3 facilitate interactions with discrete
AAP, and potentially, comprise substrate specific determinants.

We performed multiple sequence alignments of the membrane domain of Shr3 and orthologs
from two *Saccharomyces sensu stricto* strains, *S. paradoxus* and *S. mikatae*, and from three
divergent *lato* fungal strains, *Candida albicans* (Csh3), *Schizosaccharomyces pombe* (Psh3)
and *Aspergillus nidulans* (ShrA) and obtained a consensus identity plot (Fig. 3B) (Madeira et
al., 2019). The Shr3 orthologs of *C. albicans, S. pombe* and *A. nidulans* have been shown to
function analogously and are required for proper amino acid uptake. Heterologous expression
of *CSH3* complements *shr3Δ* (Martínez and Ljungdahl, 2004), whereas heterologous
expression of *PSH3* or *SHRA* only partially complement *shr3Δ*, merely facilitating the
functional expression of limited subset of permeases (Erpapazoglou et al., 2006; Martínez and
Ljungdahl, 2000). The Shr3 sequence is well-conserved in the *Saccharomyces sensu stricto*
strains, exhibiting almost absolute identity. Several positions throughout the membrane domain
of Shr3 are conserved between the full set of selected sequences. Interestingly, threonine 19,
which is the single critical amino acid residue in MS I, is conserved in all orthologs (Fig. 3B,
T19 is highlighted in dark blue). The requirement for a polar amino acid in MS IV of Shr3 is
conserved as well (Fig. 3B, S139 highlighted in dark blue). A higher sequence divergence is
evident in the luminal loop L3, with the extreme case of the *A. nidulans* orthologue that contains
an extra sequence of twelve amino acid residues. The limited sequence identity in loop L3

aligns with the observation that mutations in L3 of Shr3 exhibit the most pleiotropic effects,
and is consistent with the notion that L3 affects substrate specificity.

The finding that the Shr3Δ94 mutation supported robust growth on YPD+MM, suggested
that the mutant protein retained the capacity to facilitate Ssy1 folding. In contrast to the other
members of the AAP transporter family, Ssy1 functions as the primary receptor of extracellular
amino acids in the context of the plasma membrane-localized SPS sensor (Didion et al., 1998;
Iraqui et al., 1999; Klasson et al., 1999). In response to extracellular amino acids, Ssy1 initiates
signaling events leading to the proteolytic activation of Stp1, which in turn induces the
expression of several AAP genes including *AGP1* and *GNP1* and multiple permeases
facilitating branched amino acid uptake. As an indirect measure of Shr3-Ssy1 interactions, we
examined the proteolytic cleavage of the transcription factor Stp1 (Fig. 3C). Consistent with
the growth assays, leucine induction led to Stp1 processing in strain FGY135 (*shr3Δ*)
expressing *SHR3* or *SHR3Δ94* (Fig. 3C, lanes 4 and 8), but not the non-functional alleles *shr3-*
35, *shr3-50* or *shr3-76* (Fig. 3C, lanes 6, 10 and 12).

Shr3-AAP substrate interactions

To test if the observed growth phenotypes of the mutated *SHR3*/*shr3* alleles correlated with the
ability of mutant Shr3/shr3 proteins to interact with specific AAP and facilitate their functional
expression we exploited a split-ubiquitin approach to monitor Shr3-AAP interactions *in vivo*
(Fig. 4A and 4B). A sequence encoding the N-terminal fragment of ubiquitin carrying the I13A
mutation (NubA), which reduces the propensity of non-specific interactions (Johnsson, 2002;
Johnsson and Varshavsky, 1994), was fused at the C-terminal end of *SHR3*, *shr3-35*, and
SHR3Δ94, creating the Shr3-NubA constructs schematically depicted in Fig. 4A. Next, we
created a *GAP1* allele encoding the C-terminal fragment of ubiquitin (Cub) tagged with GST-
6xHA (Cub-GST) (Fig. 4A). The resulting *GAP1-Cub-GST* was placed under the control of the
GAL1-promoter. When co-expressed, productive interactions between Gap1 and Shr3 enable
the split NubA and Cub domains to assemble a functional ubiquitin moiety that is recognized
by ubiquitin-specific proteases, resulting in the release of the GST-6xHA reporter (Fig. 4B).
The functional attributes of the NubA fusion constructs were tested by their ability to
complement *shr3Δ* (Fig. 4C, dilutions 3-5); strains carrying *SHR3-NubA* or *SHR3Δ94-NubA*
grew as well as *SHR3* without NubA (Fig. 4C, compare dilution 2 with 3 and 5), whereas the
shr3-35-NubA allele did not (dilution 4). The *GAP1-Cub-GST* allele encodes a functional Gap1
protein that facilitates citrulline uptake as well as wildtype Gap1 (Fig. 4C, compare dilution 7
with 8). The functionality of Gap1-Cub-GST is dependent on its ability to exit the ER; a
construct lacking the ER exit motif in the hydrophilic C-terminal domain of Gap1 is not
functional (Fig. 4C dilution 9), presumably due its retention in the ER.

To test *in vivo* interactions, we analyzed protein extracts from strain FGY135 (*shr3Δ gap1Δ*)
carrying plasmids *GAP1-Cub-GST* and *SHR3-NubA*, *shr3-35-NubA* or *SHR3Δ94-NubA* using
anti-HA immunoblot analysis. In cells expressing *SHR3-NubA* or *SHR3Δ94-NubA* and *GAP1-*
Cub-GST-6xHA, two bands were detected, corresponding to full-length Gap1-Cub-GST-6xHA
and the cleaved GST-6xHA (Fig. 4D, lane 1). We calculated the fraction of split-ubiquitin
cleavage in the *SHR3-NubA* strain to be ≈ 20% by dividing the intensities of the cleaved band
with the intensities from full-length plus cleaved species; whereas in the *SHR3Δ94-NubA* strain
the cleavage was 5% (Fig. 4D); By contrast, only a single band, full-length Gap1-Cub-GST-
6xHA, was detected in extracts from the strain expressing the non-functional *shr3-35-NubA*
and *GAP1-Cub-GST-6xHA* constructs (Fig. 4D, lane 2). Although expressed at similar levels,
the ER retained gap1-ERX$_{AAA}$-Cub-GST protein did not exhibit an enhanced propensity to
interact with Shr3, suggesting the split-ubiquitin assay primarily monitors transient interactions
during AAP biogenesis (Fig. S11).

Based on the success of the split ubiquitin approach to analyze Shr3-Gap1 interactions, we
created Cub-GST-6xHA tagged constructs with five additional AAP, i.e., Agp1, Gnp1, Bap2,
Can1 and Lyp1, the non-transporting but Shr3-dependent AAP homologue Ssy1 (Klasson *et*
al., 1999), and two Shr3-independent sugar transporters (HXT), i.e., the low-affinity glucose
transporter Hxt1 and the galactose transporter Gal2 (Fig. 5A). Extracts from strain FGY135
(*shr3Δ gap1Δ*) carrying *SHR3-NubA*, *shr3-35-NubA* or *SHR3Δ94-NubA* and a single *AAP-Cub-*
GST-6xHA or *HXT-Cub-GST-6xHA* construct were prepared and the levels of the GST-6xHA
reporter were determined (Fig. 5B-E). Consistent with the general requirement of Shr3 for AAP
folding, robust interactions were detected between Agp1-, Gnp1-, Bap2- and Ssy1-Cub
constructs and wildtype Shr3-NubA (Fig. 5B, C). Although Shr3 is required for their functional
expression, Can1- and Lyp1-Cub constructs exhibited low levels of reporter cleavage, similar
to the Shr3-independent sugar transporters (Fig. 5D, E). These latter observations are consistent
with Shr3 functioning in a transient manner, primarily interacting with AAP substrates during
early stages of AAP folding. In the context of full-length AAP, the split-ubiquitin signal may
reflect weaker post-folding interactions.

AAP-Cub interactions with shr3-35-NubA were weak or absent, which is precisely aligned
with it being non-functional on all selective media tested (Fig. 1, 3). Unexpectedly, the split-
Ub interactions with unrelated sugar transporters (HXT) were also weaker compared to the
wildtype Shr3-NubA (Fig. 5E). This prompted us to test the trivial explanation that the NubA
domain of Shr3-35-NubA is incorrectly oriented. The NubA is correctly oriented to the
cytoplasm and thus in a context capable of supporting potential interactions (Supplementary
Material, Fig. 12A). Together these findings indicate that the shr3-35 protein, although
expressed at similar levels as Shr3, is incapable of engaging in both specific and non-specific
secretory substrate interactions.

Interestingly, the Shr3Δ94-NubA interacted with all AAP-Cub constructs, but at
significantly reduced levels, and consequently, the data did not explain the enhanced growth
conferred by the *SHR3Δ94* allele on YPD + MM (Fig. 2 and 3). In marked contrast, Shr3Δ94-
NubA interacted with Ssy1-Cub-GST-6xHA at levels comparable to Shr3-NubA (Fig.5 C),
which provided the mechanistic explanation for the enhanced growth phenotype. Ssy1 is a
unique member of the AAP family that strictly requires Shr3 for folding (Klasson *et al.*, 1999)
and constitutes the integral membrane component of the PM-localized SPS sensor that induces
the expression of AAP genes in response to extracellular amino acids (Ljungdahl and Daignan-
Fornier, 2012). Apparently, the Shr3Δ94 mutant retains ability to assist the folding of Ssy1,
restoring the transcriptional circuits abrogated by *shr3Δ*. Consistent with this notion, the
Shr3Δ94 allele supports Stp1 processing (Fig. 3C). The induced expression of multiple AAP
that facilitate branched-chain amino acid uptake correlates well with the growth-based
phenotypes. Together, the results indicate that the *in vivo* interactions monitored by the split
ubiquitin cleavage provide a nuanced assessment of Shr3 function.

Shr3 interacts with substrates in a progressive manner

As a proxy to investigate the temporal aspects of Shr3-facilitated AAP folding, we constructed
a series of truncated *gap1-Cub-GST, hxt1-Cub-GST* and *gal2-Cub-GST* alleles capable of
encoding 2, 4, 6, 8, 10 and 12 MS (Fig. 6A). Strain FGY135 (*shr3Δ gap1Δ*) carrying plasmid
SHR3-NubA and a truncated *gap1-Cub-GST, hxt1-Cub-GST* or *gal2-Cub-GST* allele was
employed and potential interactions were monitored by immunoblot. Shr3-NubA did not
interact with gap1-2TM, even though the Cub domain is presented in the context of proper
membrane topology oriented towards the cytoplasm (Supplementary Material, Fig. S12 B). The
presence of two additional MS of Gap1 (gap1-4TM) supported an interaction with Shr3 (Fig. 6
B left panel, C in black). The intensity of the interactions increased and eventually plateaued in
the strains carrying the *gap1-6TM/-8TM/-10TM* alleles, respectively (Fig. 6 B left panel, C in
black). Strikingly, the gap1-12TM construct interacted only weakly with the functional Shr3-
NubA (Fig. 6 B left panel, C in black). In marked contrast to the interaction pattern of the gap1-
Cub-GST truncations with Shr3-NubA, we could not detect the GST-6xHA reporter in extracts
from the strain expressing *SHR3-NubA* and any of the truncated *hxt1-Cub-GST* or *gal2-Cub-
GST* alleles (Fig. 6B, center and left panel, C in grey and white). These findings support the
notion that our split ubiquitin approach is suitable to monitor specific Shr3-AAP interactions.
As a critical test, we examined if robust interactions could be detected between Shr3-NubA and
two truncated Can1 constructs with eight and ten MS (Fig. S13). The rational being that growth-
based and biochemical assays have clearly defined Can1 as a *bona fide* substrate of Shr3.
However, interactions between full-length Can1 and Shr3 are weak, similar to that
corresponding to the non-Shr3 substrates Hxt1 and Gal2 (Fig. 5D and E). We posited that if

truncations of AAP are indeed proxies of translation intermediates, then truncations of Can1
would readily interact with Shr3. Interactions between the can1-8TM and -10TM Cub
constructs with Shr3-NubA were readily detected (Fig. S13 B, C), findings clearly consistent
with Shr3 functioning at early stages of AAP biogenesis.

To more fully understand Shr3-substrate interactions, we created a series of truncated Agp1
and Ssy1 split-ubiquitin constructs (Fig. 7A). Interestingly, in contrast to gap1-2TM, the agp1-
2TM construct clearly interacted with Shr3-NubA (Fig. 7A; black bars). Aside from this
difference, the pattern of interactions with the remaining Agp1 constructs was strikingly similar
to that observed with Gap1 truncations; the intensity of the GST-6xHA reporter increased
successively as the number of MS increased from 4 to 10, and greatly reduced when all 12 TM
were present (Fig. 7A, black bars). The interactions of the Shr3Δ94-NubA with the agp1-Cub
constructs followed a similar pattern to that of the wildtype Shr3-NubA, albeit of lower intensity
(Fig. 7A, white bars). Notedly, the interaction pattern with the Ssy1 constructs was quite
different (Fig. 7B). Interestingly, as did agp1-2TM, the ssy1-2TM interacted with Shr3-NubA,
strongly suggesting that Shr3 engages early during the biogenesis of Ssy1. The ssy1-4TM and
-6TM constructs exhibited weaker interactions, however, the ssy1-8TM, -10TM and -12TM
constructs exhibited robust interactions. As anticipated from growth-based assays, Shr3Δ94-
NubA exhibited an interaction pattern very similar to the wildtype Shr3-NubA construct.

Discussion

The biogenesis of AAP can be divided into three interconnected but discrete functional steps:
1) co-translational partitioning and integration into the ER membrane; 2) folding into native
structures; and 3) packaging into ER-derived COPII-coated transport vesicles. Shr3 is not
essential for integration (Gilstring and Ljungdahl, 2000), but is required for folding and
packaging into COPII-coated vesicles (Gilstring *et al.*, 1999; Kuehn *et al.*, 1998; Kuehn *et al.*,
1996); Kota and Ljungdahl, 2005;(Kota *et al.*, 2007; Kota and Ljungdahl, 2005). Here, our
studies were aimed at further elucidating the role of Shr3 in facilitating AAP folding, i.e. its
membrane-localized chaperone (MLC) function.

Saturation scanning mutagenesis of the N-terminal membrane domain of Shr3 allowed us
to define amino acid residues that are critical for the recognition of AAP as folding substrates
(Supplementary Material Fig. S1-S10; Fig. 3A). Strikingly, mutations affecting a few amino
acids residues at three discrete sites resulted in a complete loss of function, suggesting that Shr3
generally recognizes its folding substrates independently of sequence-specific interactions, but
rather based on the presence of structural determinants shared by the AAP. The critical residues
identified in MSI and MSIV shared the common feature of being polar (Fig. 1C). Threonine19
is conserved among closely and more distant fungal species (Fig. 3B), and it is of interest to
note that one of the original spontaneous mutations that led to the identification of *SHR3* is a

T19R mutation (Ljungdahl *et al.*, 1992). Strikingly, two mutant proteins carrying 10
consecutive leucine residues (aa 62-71, Shr3-57, MSII) or 9 (aa 145-153, Shr3-79, MSIV)
support functional expression of AAP in manner indistinguishable from wildtype Shr3, and the
Shr3-58 mutant with 13 consecutive leucine residues (aa 69-81, MSII) functions well for all
AAP except Gap1 (Fig. 3A, Fig. S6). These findings suggest that hydrophobic interactions
between MS of Shr3 and AAP can develop even at the expense of larger sections of specific
Shr3 sequence, presumably provided that its overall membrane structure is retained. The data
are consistent with Shr3 acting as a scaffold for AAP folding, a function that primarily depends
on hydrophobic interactions but with the capacity to shield energetically unfavorable polar
residues of AAP MS. Importantly, exposed polar residues may be recognized by a hydrophilic
pocket identified in the structure of the ERAD-associated E3 ubiquitin ligase Hrd1 (Schoebel
et al., 2017), which participates in the degradation of misfolded AAP in cells lacking Shr3 (Kota
et al., 2007).

The finding that one of the three loss-of-function mutations disrupting Shr3 function
resides in the lumen-oriented loop L1 (Fig. 1) suggests that the role of Shr3 as a MLC is not
restricted to MS-mediated hydrophobic interactions. Apparently, Shr3 facilitates the folding of
AAP in a manner dependent on extramembrane lumen-oriented sequences. Consistent with this,
all internal deletions in L1 of Shr3 strongly affected its function (Fig. 2). Interestingly, L1
contains a predicted α-helical structure with amphipathic characteristics (aa 44-57) that is
disrupted in each of the internal deletion mutant proteins.

By contrast to mutations affecting L1, the deletion affecting L3 (*SHR3Δ94*) exhibited a
variable effect on amino acid uptake (Fig. 2), a phenotype that we could trace to differential
interactions with distinct AAP substrates. Consequently, the data implicate L3 as an important
determinant that influences substrate interactions. This region appears to be critical for
interactions with Gap1 and Can1, but less important for interactions with Agp1/Gnp1, and
clearly dispensable for interactions with Ssy1 (Fig. 2, Fig. 4, Fig. 5). Consistently, some
mutations in *SHR3* result in WT+ phenotypes, i.e., exhibiting more robust growth than wildtype
(Fig. 3A). This latter gain-of-function phenotype is presumably due to proper Ssy1 folding;
cells that carry these mutations remain capable of inducing AAP gene expression in a manner
that is augmented by the derepression of nitrogen regulation (Ljungdahl and Daignan-Fornier,
2012). Thus, under the specific growth conditions used, cells carrying L3 mutations express
enhanced levels of AAP leading to WT+ growth phenotype.

The observation that Ssy1, the only non-transporting AAP, exhibits a lax requirement for
L3 to fold, suggests that transporting AAP may have an enhanced requirement for the chaperone
function of Shr3. This notion is consistent with recent evidence showing that extracellular loops
of AAP are not merely MS-connecting sequences but rather have important roles affecting
intracellular trafficking and transport function (van't Klooster et al., 2020). Some of the

hydrophilic loops of AAP are of considerable length, especially extracellular loops 3 and 4
connecting MS V-VI and VII-VIII, respectively. Secondary structure predictions suggest the
extracellular loop 4 of the lysine permease Lyp1 (van't Klooster *et al.*, 2020) and Gap1
(Ghaddar et al., 2014a) possesses α-helical regions that appear to influence the substrate
specificity of amino acid transport (Risinger et al., 2006). Consistent with a requirement for the
lumen-oriented loops of Shr3, the loop regions of AAP that face the extracellular milieu are
lumen-oriented during biogenesis. Interactions between the lumen-oriented loops of Shr3 may
prevent precocious folding of the extracellular loops, maintaining them in a more flexible state
required for subsequent folding events, e.g., involving MS that fold in a context with more
distal C-terminal MS. In analogy, proper trafficking and functional expression of the closest
AAP homologues in mammals, the L-type amino acid transporters (LAT) (SLC7 family)
depend on extramembrane region-mediated recognition by the 4F2hc, or rBAT, members of
the SLC3 protein family (Fotiadis et al., 2013). The extracellular orientation of the interacting
regions upon plasma-membrane localization of the SLC7-SLC3 holo-LAT transporters in in
line with the concept of extramembrane regions containing motifs for substrate recognition and
specificity exhibited by amino acid transporter biogenesis factors (Rosell et al., 2014).

Although there is no crystal structure of an AAP, Bap2 (Usami et al., 2014), Can1 (Ghaddar
et al., 2014a), Gap1 (Ghaddar et al., 2014b) and Tat2 (Kanda and Abe, 2013) have been
successfully modeled onto the *E. coli* arginine/agmatine antiporter AdiC (Gao et al., 2010).
AdiC has 12 MS and belongs to the Amino Acid-Polyamine-Organocation (APC) super-family
of transporters. The 12 MS are arranged in a 5+5 inverted repeat fold that form the transporter
core, with MS XI and XII appearing to hold the two halves together. MS I, III, VI, VIII and X
shape the binding pocket. To directly probe the *in vivo* interactions between Shr3 and AAP we
adapted and applied a split-ubiquitin approach that is independent of a transcriptional readout.

In the context of full-length AAP, we found that the split-ubiquitin signal was relatively
weak, specifically in comparison to the signals with truncated substrates. Also, we detected
low levels of interactions with full length non-Shr3 substrates that do not rely on Shr3
chaperone function for folding, indicating that the split-ubiquitin approach is sensitive and
accurately reflects biologically relevant interactions. Perhaps, Shr3 can interact with many
secretory substrates but some interactions become stronger and specific when structural
characteristics distinct from the general ones are involved. In support of this notion, the pattern
of interactions between Shr3 and truncated Gap1, Agp1, Can1 and Ssy1 is striking, particularly
in contrast to truncated HXT (Hxt1 and Gal2) exhibiting essentially no interactions (Fig. 6, 7,
and S13). The specific interactions with AAP were detected when only the first 2 to 4 MS were
present. As more MS were added the interactions increased and plateaued until MS XI and XII
were present, at which point the interactions significantly lessen. By contrast, full-length (Fig.
5) and 12MS truncated (Fig. 7) forms of Ssy1 exhibited more persistent interactions with Shr3
and Shr3Δ94, perhaps attributable to Ssy1 being substantially larger than transporting AAP.

Together, the data we acquired in this study provide further support to our previously
described model (Kota *et al.*, 2007; Kota and Ljungdahl, 2005) whereby Shr3 transiently
interacts with AAP early as their MS partition into the ER membrane, acting as an assembly
site for MS helices. This activity is required to prevent AAP translation intermediates from
engaging in nonproductive interactions, shielding polar residues, until all MS are available and
presented in a context of the long-range intramolecular interactions inherent to the native 3D-
structure of AAP (Fig.8). Our model for the Shr3 chaperone function in the biogenesis of AAP
is analogous to that of the bacterial insertase/chaperone YidC (Beck et al., 2001; (Dalbey and
Kuhn, 2014) acting as an assembly site for alpha helices in the folding of LacY whereby
hydrophobic interactions mediate shielding of LacY to provide a protective chamber that
reduces energetically unfavorable contacts in the non-native structure during translation
(Nagamori et al., 2004; Serdiuk et al., 2016; Serdiuk et al., 2019; Wagner et al., 2008; Zhu et
al., 2013). Similar transient MLC-substrate interactions have been reported in the case of the
mammalian PAT intramembrane chaperone complex, comprised of four MS, three contributed
by Asterix and one by CCDC47 (Chitwood and Hegde, 2020). Although in these studies an *in
vitro* translation system was employed, in contrast to our *in vivo* split-ubiquitin approach, the
PAT complex interaction was found to be selective for truncated, immature β1-adrenergic
receptor constructs compared to the full-length substrate, which presumably is capable of helix
packing and polar residue shielding.

The chaperone-like capacity of the conserved eukaryotic ER membrane protein complex
(EMC) insertase has recently been explored (Bai et al., 2020; O'Donnell et al., 2020; Pleiner et
al., 2020; Volkmar and Christianson, 2020). The results point to EMC acting in close proximity
with nascent polytopic membrane proteins typically enriched for MS containing polar or
charged residues, shielding them from degradation during folding (Miller-Vedam *et al.*, 2020;
Shurtleff *et al.*, 2018). Importantly, the EMC can associate with a number of ER-integral
substrate-specific chaperones such as Sop4 (Luo *et al.*, 2002), Gsf2 (Kota and Ljungdahl, 2005;
Sherwood and Carlson, 1999) and Ilm1 (Shurtleff *et al.*, 2018). Shr3, although an abundant ER
membrane protein, has not been identified as an interacting partner of EMC components
(Shurtleff *et al.*, 2018). Potentially, Shr3 and the EMC act in proximity to distinct ribosome
populations, functioning in a parallel manner, enabling the Sec61 translocon to pair with diverse
and distinct sets of partners, and thereby facilitate the efficient biogenesis of more challenging
versus canonical substrates of the secretory pathway (O'Keefe and High, 2020).

In the case of Shr3, the early interactions with nascent AAP being inserted in the lipid
bilayer together with the ability of Shr3 to interact with COPII components via its C-terminal
cytoplasmic tail (Gilstring *et al.*, 1999), converge to function as a nexus between AAP folding
and packaging into COPII-coated vesicles. The potential network of dynamic interactions in
the ER remains to be explored for an integral substrate-specific chaperone as well as what
structural determinants in the substrates dictate a remarkable degree of substrate specificity. To

this end, the broad collection of mutations that we have acquired in combination to our extended
growth-based phenotype analysis and in vivo interaction studies lay the ground for future
structural studies that are required to gain insights into the mechanistic details that underlie this
substrate-specific MLC function.

Materials and methods

Yeast strains and plasmids

515 Yeast strains and plasmid used are listed in Supplementary Material Tables S1 and S2,
516 respectively.

Media

Standard media, YPD (yeast extract, peptone, dextrose), SD (synthetic defined with ammonium
as nitrogen source and glucose as carbon source) were prepared as previously described (Burke
et al., 2000). Ammonia-based synthetic complete dextrose (SC) drop-out medium, were
prepared as described (Andréasson and Ljungdahl, 2002) and SAD (synthetic minimal
dextrose, with allantoin as sole nitrogen source) was prepared as previously described. Media
were made solid with 2% (wt/vol) bacto Agar (Difco), 2% (wt/vol) washed bacto Agar (Difco)
or 2% (wt/vol) washed pure Agar where indicated. Sensitivity to 200 µg/ml MM (2-{[({[(4-
methoxy-6-methyl)-1,3,5-triazin-2-yl]-amino}carbonyl) amino-]-sulfonyl}-benzoic acid) was
tested on YPD as described previously (Jørgensen et al., 1998). Sensitivity to 1 mM AzC
(azetidine-2-carboxylate), 10 µg/ml DL-ethionine, 50 µg/ml p-Fluoro-DL-phenylalanine and 1
µg/ml L-canavanine was tested on SD. Sensitivity to 0,5% (wt/vol) D-histidine was tested on
SAD media made solid with washed pure Agar. Cells were grown over night in SC-uracil
medium, cells were then resuspended in water to OD=1, 10-fold dilutions were prepared in
water and then spotted on the indicated medium. Plates were then incubated at 30°C for 2–3 d
and photographed. Gap1-dependent citrulline uptake was monitored on minimal medium
containing 2 % galactose as carbon source, 1 mM L-citrulline as sole nitrogen source and uracil.
Media were made solid with washed bacto Agar. Plates were incubated at 30 °C for 7 d and
photographed.

Immunoblot analysis

Whole-cell extracts were prepared under denaturing conditions using NaOH and trichloroacetic
acid as described previously (Silve et al., 1991). Proteins were separated using SDS-PAGE and
blotted onto Amersham Protran 0.45 µm nitrocellulose membrane (GE Healthcare). The
primary antibodies and dilutions were, mouse anti-Dpm1 5C5A7 (Abcam), 1:2500; rat anti-

HA-HRP 3F10 (Roche Applied Science), 1:2500-1:5000; mouse anti-Pgk1 22C5D8 (Thermo
Fisher Scientific), 1:10000 and rabbit anti-Shr3, 1:9000. Secondary antibodies and dilutions
used were, goat anti-mouse-poly-HRP (Thermo Fisher Scientific), 1:5000 and goat anti-rabbit-
poly-HRP (Thermo Fisher Scientific), 1:5000. Immunoreactive bands were visualized by
chemiluminescence using (SuperSignal West Dura Extended-Duration Substrate; Thermo
Fisher Scientific) as substrate in a ChemiDoc imaging system (Biorad).

Split-ubiquitin assay

Cells were pre-grown in SD+R (synthetic defined with ammonium as nitrogen source and 2 %
raffinose and 0.1 % glucose as carbon source) to logarithmic phase. Approximately 10 OD of
logarithmically cells were induced in 5 ml of SD+G (synthetic defined with ammonium as
nitrogen source and 2 % galactose as carbon source) for 1 hour. Cells were collected and washed
once in ddH$_2$0. Cells were resuspended in 150 µl lysis buffer (0.8 M sorbitol; 10 mM MOPS,
pH 7.2; 2 mM EDTA; 1 mM PMSF; 1X cØmplete, mini, EDTA-free protease inhibitor cocktail,
Roche). Cells were lysed by bead beating with 0.5 mm glass beads for 3x20s at 6.5 m/s in a
benchtop homogenizer (Fastprep-24, MP Biomedical). The cell lysates were centrifuged at
500g for 10 min and 25 µl of the resulting supernatant was diluted 1:1 with 2x sample buffer.
Proteins were separated using SDS-PAGE and blotted onto Amersham Protran 0.45 µm
nitrocellulose membrane (GE Healthcare). The primary antibody and dilution used was rat anti-
HA-HRP 3F10 (Roche Applied Science), 1:2500-1:5000. Immunoreactive bands were
visualized by chemiluminescence using (SuperSignal West Dura Extended-Duration Substrate;
Thermo Fisher Scientific) as substrate in a ChemiDoc imaging system (Biorad).

Protease protection assay

Cells were pre-grown in SD+R (synthetic defined with ammonium as nitrogen source and 2 %
raffinose and 0.1 % glucose as carbon source) to logarithmic phase. Approximately 5 OD of
logarithmically cells were induced in 5 ml of SD+G (synthetic defined with ammonium as
nitrogen source and 2 % galactose as carbon source) for 1 hour. Cells were collected and washed
once in ddH$_2$0. Cells were resuspended in 150 µl lysis buffer (0.8 M sorbitol; 10 mM MOPS,
pH 7.2; 2 mM EDTA; 1 mM PMSF; 1X cØmplete, mini, EDTA-free protease inhibitor cocktail,
Roche). Cells were lysed by bead beating with 0.5 mm glass beads for 3x20s at 6.5 m/s in a
benchtop homogenizer (Fastprep-24, MP Biomedical). The cell lysates were centrifuged at
500g for 10 min and 100 µl of the resulting supernatant was centrifuged at 100 000 g for 30
minutes. The membrane pellet was resuspended in 50 µl lysis buffer (0.8 M sorbitol; 10 mM
MOPS, pH 7.2; 2 mM EDTA; 5 mM CaCl$_2$). The resulting membrane preparations were
digested with 20 µg Proteinase K (Thermo Fisher Scientific) on ice with 0.2 % NP-40 as

indicated. Time points where taken at 0 and 2 h. Proteins were precipitated using trichloroacetic

acid as described previously (Silve *et al.*, 1991). Proteins were separated using SDS-PAGE and

blotted onto Amersham Protran 0.45 μm nitrocellulose membrane (GE Healthcare). The

primary antibodies and dilutions used were, rat anti-HA-HRP 3F10 (Roche Applied Science),

1:2500-1:5000 and rabbit anti-Kar2, 1:5000. Secondary antibody and dilution used was goat

anti-rabbit-poly-HRP (Thermo Fisher Scientific), 1:5000. Immunoreactive bands were

visualized by chemiluminescence using (SuperSignal West Dura Extended-Duration Substrate;

Thermo Fisher Scientific) as substrate in a ChemiDoc imaging system (Biorad).

Acknowledgements

589 We thank the members of the Ljungdahl laboratory and Claes Andréasson for constructive

590 comments throughout the course of this work. In particular we acknowledge Nina Horwege,

591 and Carlos Sacristán for early contributions in creating plasmid constructs. This research was

592 supported by funding from Swedish Research Council (P.O.L.), Grant/Award numbers: 2011-

593 5925 and 2015-04202.

Competing interests

596 The authors have no conflicts of interest to report.

References

Andréasson, C., and Ljungdahl, P.O. (2002). Receptor-mediated endoproteolytic activation

of two transcription factors in yeast. Genes Dev. *16*, 3158-3172.

Andréasson, C., Neve, E.P.A., and Ljungdahl, P.O. (2004). Four permeases import proline

and the toxic proline analogue azetidine-2-carboxylate into yeast. Yeast *21*, 193-199.

Bai, L., You, Q., Feng, X., Kovach, A., and Li, H. (2020). Structure of the ER membrane

complex, a transmembrane-domain insertase. Nature *584*, 475-478. 10.1038/s41586-

020-2389-3.

Chitwood, P.J., and Hegde, R.S. (2020). An intramembrane chaperone complex facilitates

membrane protein biogenesis. Nature *584*, 630-634. 10.1038/s41586-020-2624-y.

Dalbey, R.E., and Kuhn, A. (2014). How YidC inserts and folds proteins across a membrane.

Nat Struct Mol Biol *21*, 435-436. 10.1038/nsmb.2823.

Didion, T., Regenberg, B., Jørgensen, M.U., Kielland-Brandt, M.C., and Andersen, H.A.

(1998). The permease homologue Ssy1p controls the expression of amino acid and

peptide transporter genes in *Saccharomyces cerevisiae*. Mol. Microbiol. *27*, 643-650.

Drozdetskiy, A., Cole, C., Procter, J., and Barton, G.J. (2015). JPred4: a protein secondary

structure prediction server. Nucleic Acids Res *43*, W389-394. 10.1093/nar/gkv332.

Erpapazoglou, Z., Kafasla, P., and Sophianopoulou, V. (2006). The product of the SHR3
orthologue of Aspergillus nidulans has restricted range of amino acid transporter targets.
Fungal Genet Biol *43*, 222-233. 10.1016/j.fgb.2005.11.006.

Fotiadis, D., Kanai, Y., and Palacin, M. (2013). The SLC3 and SLC7 families of amino acid
transporters. Mol Aspects Med *34*, 139-158. 10.1016/j.mam.2012.10.007.

Gao, X., Zhou, L., Jiao, X., Lu, F., Yan, C., Zeng, X., Wang, J., and Shi, Y. (2010). Mechanism of
substrate recognition and transport by an amino acid antiporter. Nature *463*, 828-832.
10.1038/nature08741.

Ghaddar, K., Krammer, E.M., Mihajlovic, N., Brohee, S., Andre, B., and Prevost, M. (2014a).
Converting the yeast arginine can1 permease to a lysine permease. J Biol Chem *289*, 7232-
7246. 10.1074/jbc.M113.525915.

Ghaddar, K., Merhi, A., Saliba, E., Krammer, E.M., Prevost, M., and Andre, B. (2014b).
Substrate-induced ubiquitylation and endocytosis of yeast amino acid permeases. Mol
Cell Biol *34*, 4447-4463. 10.1128/MCB.00699-14.

Gilstring, C.F., and Ljungdahl, P.O. (2000). A method for determining the in vivo topology
of yeast polytopic membrane proteins demonstrates that Gap1p fully integrates into the
membrane independently of Shr3p. J. Biol. Chem. *275*, 31488-31495.

Gilstring, C.F., Melin-Larsson, M., and Ljungdahl, P.O. (1999). Shr3p mediates specific
COPII coatomer-cargo interactions required for the packaging of amino acid permeases
into ER-derived transport vesicles. Mol. Biol. Cell *10*, 3549-3565.

Grenson, M., Hou, C., and Crabeel, M. (1970). Multiplicity of the amino acid permeases in
Saccharomyces cerevisiae. IV. Evidence for a general amino acid permease. J Bacteriol
103, 770-777.

Gresham, D., Usaite, R., Germann, S.M., Lisby, M., Botstein, D., and Regenberg, B. (2010).
Adaptation to diverse nitrogen-limited environments by deletion or extrachromosomal
element formation of the GAP1 locus. Proc Natl Acad Sci U S A *107*, 18551-18556.
10.1073/pnas.1014023107.

Heinrich, S.U., Mothes, W., Brunner, J., and Rapoport, T.A. (2000). The Sec61 complex
mediates the integration of a membrane protein by allowing lipid partitioning of the
transmembrane domain. Cell *102*, 233-244.

Heinrich, S.U., and Rapoport, T.A. (2003). Cooperation of transmembrane segments
during the integration of a double-spanning protein into the ER membrane. Embo J *22*,
3654-3663.

Hou, B., Lin, P.J., and Johnson, A.E. (2012). Membrane protein TM segments are retained
at the translocon during integration until the nascent chain cues FRET-detected release
into bulk lipid. Mol Cell *48*, 398-408. 10.1016/j.molcel.2012.08.023.

Iraqui, I., Vissers, S., Bernard, F., de Craene, J.O., Boles, E., Urrestarazu, A., and André, B.
(1999). Amino acid signaling in *Saccharomyces cerevisiae*: a permease-like sensor of
external amino acids and F-Box protein Grr1p are required for transcriptional induction

of the *AGP1* gene, which encodes a broad-specificity amino acid permease. Mol. Cell. Biol.
19, 989-1001.

Jack, D.L., Paulsen, I.T., and Saier, M.H. (2000). The amino acid/polyamine/organocation
(APC) superfamily of transporters specific for amino acids, polyamines and
organocations. Microbiology *146 (Pt 8)*, 1797-1814. 10.1099/00221287-146-8-1797.

Johnson, A.E., and van Waes, M.A. (1999). The translocon: a dynamic gateway at the ER
membrane. Annu Rev Cell Dev Biol *15*, 799-842. 10.1146/annurev.cellbio.15.1.799.

Johnsson, N. (2002). A split-ubiquitin-based assay detects the influence of mutations on
the conformational stability of the p53 DNA binding domain in vivo. FEBS Lett *531*, 259-
264. S0014579302035330 [pii].

Johnsson, N., and Varshavsky, A. (1994). Split ubiquitin as a sensor of protein interactions
in vivo. Proc Natl Acad Sci U S A *91*, 10340-10344.

Jørgensen, M.U., Bruun, M.B., Didion, T., and Kielland-Brandt, M.C. (1998). Mutations in
five loci affecting GAP1-independent uptake of neutral amino acids in yeast. Yeast *14*, 103-
114.

Kanda, N., and Abe, F. (2013). Structural and functional implications of the yeast high-
affinity tryptophan permease Tat2. Biochemistry *52*, 4296-4307. 10.1021/bi4004638.

Klasson, H., Fink, G.R., and Ljungdahl, P.O. (1999). Ssy1p and Ptr3p are plasma membrane
components of a yeast system that senses extracellular amino acids. Mol Cell Biol *19*,
5405-5416.

Kota, J., Gilstring, C.F., and Ljungdahl, P.O. (2007). Membrane chaperone Shr3 assists in
folding amino acid permeases preventing precocious ERAD. J Cell Biol *176*, 617-628.
10.1083/jcb.200612100.

Kota, J., and Ljungdahl, P.O. (2005). Specialized membrane-localized chaperones prevent
aggregation of polytopic proteins in the ER. J Cell Biol *168*, 79-88.

Kuehn, M.J., Herrmann, J.M., and Schekman, R. (1998). COPII-cargo interactions direct
protein sorting into ER-derived transport vesicles. Nature *391*, 187-190.

Kuehn, M.J., Schekman, R., and Ljungdahl, P.O. (1996). Amino acid permeases require
COPII components and the ER resident membrane protein Shr3p for packaging into
transport vesicles in vitro. J. Cell Biol. *135*, 585-595.

Lau, W.T., Howson, R.W., Malkus, P., Schekman, R., and O'Shea, E.K. (2000). Pho86p, an
endoplasmic reticulum (ER) resident protein in *Saccharomyces cerevisiae*, is required for
ER exit of the high-affinity phosphate transporter Pho84p. Proc. Natl. Acad. Sci. USA *97*,
1107-1112.

Ljungdahl, P.O., and Daignan-Fornier, B. (2012). Regulation of amino acid, nucleotide, and
phosphate metabolism in Saccharomyces cerevisiae. Genetics *190*, 885-929.
10.1534/genetics.111.133306.

Ljungdahl, P.O., Gimeno, C.J., Styles, C.A., and Fink, G.R. (1992). SHR3: A novel component
of the secretory pathway specifically required for the localization of amino acid
permeases in yeast. Cell *71*, 463-478.

Luo, W.J., Gong, X.H., and Chang, A. (2002). An ER membrane protein, Sop4, facilitates ER
export of the yeast plasma membrane [H+]ATPase, Pma1. Traffic *3*, 730-739.
10.1034/j.1600-0854.2002.31005.x.

Madeira, F., Park, Y.M., Lee, J., Buso, N., Gur, T., Madhusoodanan, N., Basutkar, P., Tivey,
A.R.N., Potter, S.C., Finn, R.D., and Lopez, R. (2019). The EMBL-EBI search and sequence
analysis tools APIs in 2019. Nucleic Acids Research *47*, W636-W641.
https://doi.org/10.1093/nar/gkz268.

Malkus, P., Jiang, F., and Schekman, R. (2002). Concentrative sorting of secretory cargo
proteins into COPII-coated vesicles. J. Cell Biol. *159*, 915-921.

Martínez, P., and Ljungdahl, P.O. (2000). The SHR3 homologue from *S. pombe*
demonstrates a conserved function of ER packaging chaperones. J. Cell Sci. *113*, 4351-
4362.

Martínez, P., and Ljungdahl, P.O. (2004). An ER packaging chaperone determines the
amino acid uptake capacity and virulence of *Candida albicans*. Mol Microbiol *51*, 371-384.

Miller, E., Antonny, B., Hamamoto, S., and Schekman, R. (2002). Cargo selection into COPII
vesicles is driven by the Sec24p subunit. EMBO J. *21*, 6105-6113.

Miller, E.A., Beilharz, T.H., Malkus, P.N., Lee, M.C., Hamamoto, S., Orci, L., and Schekman, R.
(2003). Multiple cargo binding sites on the COPII subunit Sec24p ensure capture of
diverse membrane proteins into transport vesicles. Cell *114*, 497-509.

Miller-Vedam, L.E., Brauning, B., Popova, K.D., Schirle Oakdale, N.T., Bonnar, J.L., Prabu,
J.R., Boydston, E.A., Sevillano, N., Shurtleff, M.J., Stroud, R.M., et al. (2020). Structural and
mechanistic basis of the EMC-dependent biogenesis of distinct transmembrane clients.
Elife *9*. 10.7554/eLife.62611.

Nagamori, S., Smirnova, I.N., and Kaback, H.R. (2004). Role of YidC in folding of polytopic
membrane proteins. J. Cell Biol. *165*, 53-62.

O'Donnell, J.P., Phillips, B.P., Yagita, Y., Juszkiewicz, S., Wagner, A., Malinverni, D., Keenan,
R.J., Miller, E.A., and Hegde, R.S. (2020). The architecture of EMC reveals a path for
membrane protein insertion. Elife *9*. 10.7554/eLife.57887.

O'Keefe, S., and High, S. (2020). Membrane translocation at the ER: with a little help from
my friends. FEBS J. 10.1111/febs.15309.

Ono, B.I., Ishino, Y., and Shinoda, S. (1983). Nonsense mutations in the can1 locus of
Saccharomyces cerevisiae. J Bacteriol *154*, 1476-1479.

Pleiner, T., Tomaleri, G.P., Januszyk, K., Inglis, A.J., Hazu, M., and Voorhees, R.M. (2020).
Structural basis for membrane insertion by the human ER membrane protein complex.
Science *369*, 433-436. 10.1126/science.abb5008.

Rapoport, T.A., Li, L., and Park, E. (2017). Structural and Mechanistic Insights into Protein
Translocation. Annu Rev Cell Dev Biol *33*, 369-390. 10.1146/annurev-cellbio-100616-
060439.

Risinger, A.L., Cain, N.E., Chen, E.J., and Kaiser, C.A. (2006). Activity-dependent reversible
inactivation of the general amino acid permease. Mol Biol Cell *17*, 4411-4419.
10.1091/mbc.e06-06-0506.

Rosell, A., Meury, M., Alvarez-Marimon, E., Costa, M., Perez-Cano, L., Zorzano, A.,
Fernandez-Recio, J., Palacin, M., and Fotiadis, D. (2014). Structural bases for the
interaction and stabilization of the human amino acid transporter LAT2 with its ancillary
protein 4F2hc. Proc Natl Acad Sci U S A *111*, 2966-2971. 10.1073/pnas.1323779111.

Rytka, J. (1975). Positive selection of general amino acid permease mutants in
Saccharomyces cerevisiae. J Bacteriol *121*, 562-570.

Saier, M.H., Jr. (2000). Families of transmembrane transporters selective for amino acids
and their derivatives. Microbiology *146 (Pt 8)*, 1775-1795. 10.1099/00221287-146-8-
1775.

Schoebel, S., Mi, W., Stein, A., Ovchinnikov, S., Pavlovicz, R., DiMaio, F., Baker, D., Chambers,
M.G., Su, H., Li, D., et al. (2017). Cryo-EM structure of the protein-conducting ERAD channel
Hrd1 in complex with Hrd3. Nature *548*, 352-355. 10.1038/nature23314.

Seinen, A.B., and Driessen, A.J.M. (2019). Single-Molecule Studies on the Protein
Translocon. Annu Rev Biophys *48*, 185-207. 10.1146/annurev-biophys-052118-115352.

Serdiuk, T., Balasubramaniam, D., Sugihara, J., Mari, S.A., Kaback, H.R., and Muller, D.J.
(2016). YidC assists the stepwise and stochastic folding of membrane proteins. Nat Chem
Biol *12*, 911-917. 10.1038/nchembio.2169.

Serdiuk, T., Steudle, A., Mari, S.A., Manioglu, S., Kaback, H.R., Kuhn, A., and Muller, D.J.
(2019). Insertion and folding pathways of single membrane proteins guided by
translocases and insertases. Sci Adv *5*, eaau6824. 10.1126/sciadv.aau6824.

Sherwood, P.W., and Carlson, M. (1999). Efficient export of the glucose transporter Hxt1p
from the endoplasmic reticulum requires Gsf2p. Proc. Natl. Acad. Sci. USA *96*, 7415-7420.

Shurtleff, M.J., Itzhak, D.N., Hussmann, J.A., Schirle Oakdale, N.T., Costa, E.A., Jonikas, M.,
Weibezahn, J., Popova, K.D., Jan, C.H., Sinitcyn, P., et al. (2018). The ER membrane protein
complex interacts cotranslationally to enable biogenesis of multipass membrane proteins.
Elife *7*. 10.7554/eLife.37018.

Silve, S., Volland, C., Garnier, C., Jund, R., Chevallier, M.R., and Haguenauer-Tsapis, R.
(1991). Membrane insertion of uracil permease, a polytopic yeast plasma membrane
protein. Mol. Cell. Biol. *11*, 1114-1124.

Usami, Y., Uemura, S., Mochizuki, T., Morita, A., Shishido, F., Inokuchi, J., and Abe, F. (2014).
Functional mapping and implications of substrate specificity of the yeast high-affinity
leucine permease Bap2. Biochim Biophys Acta *1838*, 1719-1729.
10.1016/j.bbamem.2014.03.018.

van't Klooster, J.S., Bianchi, F., Doorn, R.B., Lorenzon, M., Lusseveld, J.H., Punter, C.M., and
Poolman, B. (2020). Extracellular loops matter - subcellular location and function of the
lysine transporter Lyp1 from Saccharomyces cerevisiae. FEBS J. 10.1111/febs.15262.

Volkmar, N., and Christianson, J.C. (2020). Squaring the EMC - how promoting membrane
protein biogenesis impacts cellular functions and organismal homeostasis. J Cell Sci *133*.
10.1242/jcs.243519.

Wagner, S., Pop, O.I., Haan, G.J., Baars, L., Koningstein, G., Klepsch, M.M., Genevaux, P.,
Luirink, J., and de Gier, J.W. (2008). Biogenesis of MalF and the MalFGK(2) maltose
transport complex in Escherichia coli requires YidC. J Biol Chem *283*, 17881-17890.
10.1074/jbc.M801481200.

Waterhouse, A.M., Procter, J.B., Martin, D.M., Clamp, M., and Barton, G.J. (2009). Jalview
Version 2--a multiple sequence alignment editor and analysis workbench. Bioinformatics
25, 1189-1191. 10.1093/bioinformatics/btp033.

Wong, F.H., Chen, J.S., Reddy, V., Day, J.L., Shlykov, M.A., Wakabayashi, S.T., and Saier, M.H.,
Jr. (2012). The amino acid-polyamine-organocation superfamily. J Mol Microbiol
Biotechnol *22*, 105-113. 10.1159/000338542.

Zhu, L., Kaback, H.R., and Dalbey, R.E. (2013). YidC protein, a molecular chaperone for
LacY protein folding via the SecYEG protein machinery. J Biol Chem *288*, 28180-28194.
10.1074/jbc.M113.491613.

Figure 1. Scanning mutagenesis of the Shr3 membrane domain

(A) Graphical representation of Shr3 topology and position of residues resulting in a non-functional protein. (B) Top: Serial dilutions of cell suspensions from strain JKY2 (*shr3Δ*) carrying pRS316 (VC), pPL210 (*SHR3*), pAR4 (*shr3-35*), pAR18 (*shr3-50*) or pPL1349 (*shr3-76*) spotted on YPD and YPD+MM. The plates were incubated at 30 °C for 2 d and photographed. Bottom: Immunoblot analysis of Shr3 proteins in extracts prepared from the strains; the levels of Pgk1 were used as loading controls. The blots were developed using α-Shr3 and α-Pgk1 antibodies. The signal intensities of the immunoreactive forms of Shr3 and Pgk1 were quantified, and the Shr3 signals were normalized with respect to Pgk1; the mean values are plotted, error bars show standard deviation (n=3). (C) Serial dilutions of cell suspensions from strain JKY2 (*shr3Δ*) carrying pRS316 (VC), pPL210 (*SHR3*), pAR4 (*shr3-35*), pPL1330 (*SHR3-36*), pAR47 (*shr3-37*), pAR37 (*SHR3-38*), pAR18 (*shr3-50*), pAR51 (*SHR3-51*), pAR52 (*SHR3-52*), pAR50 (*shr3-53*), pPL1349 (*shr3-76*), pAR48 (*shr3-77*) or pAR49 (*SHR3-78*) spotted on YPD and YPD+MM plates. Plates were incubated at 30 °C for 2 d and photographed.

Figure 2. Deletion analysis of ER-lumen oriented loops

(A) Graphical representation of Shr3 topology and the positions of the internal deletions in loops L1 and L3. (B) Based on structural predictions (Drozdetskiy et al., 2015), amino acid residues 44-57 in L1 are predicted to fold into an α-helix with amphipathic character. Helical wheel projection of the L1 α-helix with non-polar (yellow), polar (grey), negatively- (blue) and positively-charged (red) residues indicated. (C)   Serial dilutions of cell suspensions from strain JKY2 (*shr3Δ*) carrying pRS316 (VC), pPL210 (*SHR3*), pAR41 (*shr3Δ90*), pAR42 (*shr3Δ91*), pAR43 (*shr3Δ92*), pAR44 (*shr3Δ93*) or pAR45 (*shr3Δ94*) spotted on SAD containing D-histidine (D-his), SD + L-canavanine (L-Can), SD + AzC, and YPD + MM plates. Plates were incubated at 30 °C for 2 d and photographed. Bottom: Immunoblot analysis of Shr3 proteins in extracts prepared from the strains; the levels of Dpm1 were used as loading controls. The blots were developed using α-Shr3 and α-Dpm1 antibodies. The signal intensities of the immunoreactive forms of Shr3 and Dpm1 were quantified, and the Shr3 signals were normalized with respect to Dpm1; the mean values are plotted, error bars show standard deviation (n=3).

Figure 3. Mutational analysis of Shr3 function and substrate specificity

(A) Summary of growth characteristics of JKY2 (*shr3Δ*) individually expressing 44 Shr3-mutant proteins. Cells were spotted on media containing toxic amino analogues and nitrogen sources as follows: D-his, D-histidine (0.5% w/v), allantoin; L-can, L-canavanine (1 μg/ml), ammonium; AzC, azetidine-2-carboxylate (1 mM), ammonium; MM (200 μg/ml), yeast extract and peptone. Growth was scored after 2 - 3 d of incubation at 30 °C (Supplementary Materials Fig. S1-S10). Colors reflect Shr3 function relative to wildtype activity: red, no function (-); orange, weak but detectable function (+/-); yellow, intermediate function but less than wildtype (+); light blue, wildtype function (WT); dark blue, enhanced function (WT+). (B) Clustal O (Madeira *et al.*, 2019) comparison of Shr3 sequences, corresponding to aa residues 1-159 of *S. cerevisiae*, and orthologs of members from the *Saccharomyces sensu stricto* group (*S. paradoxus*, *S. mikatae*) and orthologs from *sensu lato* fungi (*S. pombe, A. nidulans,* and *C. albicans*). The consensus plot (identity; (Waterhouse et al., 2009) and detailed multiple sequence alignments are presented for the regions with mutations giving rise to major growth defects on selective media; identical residues in three (light blue), four (blue), and five or six homologs (dark blue) are highlighted. (C) Shr3-dependent Ssy1 folding and function assessed by Stp1 processing. Immunoblot analysis of extracts from FGY135 (*shr3Δ*) carrying pCA204 (*STP1-13xMYC*) and pRS316 (VC), pPL210 (*SHR3*), pAR004 (*shr3-35*), pAR45 (*SHR3Δ94*), pAR018 (*shr3-50*) or pPL1351 (*shr3-76*). Cells were grown in SD and induced 30 min with 1.3 mM leucine (+) as indicated.

Figure 4. Assessing Shr3-Gap1 interactions using split ubiquitin

(A) Schematic diagram of the split ubiquitin Shr3-NubA, shr3-35-NubA, Shr3Δ94-NubA and Gap1-Cub-GST constructs. (B) Overview of the split-ubiquitin assay and expected outcomes. (C) Left panels: serial dilutions of cell suspensions from strain JKY2 (*shr3Δ*) carrying pRS316 (VC), pPL210 (*SHR3*), pPL1262 (*SHR3-NubA*), pAR67 (*shr3-35-NubA*) or pAR76 (*SHR3Δ94-NubA*) spotted on YPD and YPD+MM plates. Plates were incubated at 30 °C for 2 d and photographed. Right panel: serial dilutions of cell suspensions from strain FGY15 (*gap1Δ*) carrying pRS317 (VC), pJK92 (*GAP1*), pPL1257 (*GAP1-Cub-GST*) or pIM28 (*gap1-ERX_{AAA}-Cub-GST*) were spotted on minimal medium with 2 % galactose as carbon source and 1 mM L-citrulline as sole nitrogen source. Plates were incubated for 7 d and photographed. (D) Strain FGY135 (*gap1Δ shr3Δ*) expressing *SHR3-NubA* (pPL1262), *shr3-35-NubA* (pAR67) or *SHR3Δ94-NubA* (pAR76) and carrying pPL1257 (*GAP1-Cub-GST*) were induced with 2% galactose for 1 h. Proteins extracts were prepared, separated by SDS-PAGE and analyzed by immunoblotting using α-HA antibody. The signal intensities of the immunoreactive forms of full-length and cleaved Gap1 were quantified. The fraction of split-ubiquitin cleavage was determined; the mean values plotted with error bars showing standard deviation (n=3).

Figure 5. Monitoring Shr3-AAP interactions *in vivo*

(A) Schematic diagram of the split ubiquitin constructs used to evaluate Shr3-AAP and Shr3-HXT interactions.
(B) Shr3-Agp1, Shr3-Gnp1 and Shr3-Bap2 interactions. (C) Shr3-Ssy1 interactions. (D) Shr3-Can1 and Shr3-
Lyp1 interactions. (E) Shr3-Hxt1 and Shr3-Gal2 interactions. Strain FGY135 (*gap1Δ shr3Δ*) expressing *SHR3-*
NubA (pPL1262), *shr3-35-NubA* (pAR67) or *SHR3Δ94-NubA* (pAR76) and carrying (B) pIM6 (*AGP1-Cub-GST*),
pIM17 (*GNP1-Cub-GST*), or pIM7 (*BAP2-Cub-GST*) or (C) pIM19 (*SSY1-Cub-GST*) or (D) pIM8 (*CAN1-Cub-*
GST) or pIM18 (*LYP1-Cub-GST*), or (E) pIM32 (*HXT1-Cub-GST*) or pIM33 (*GAL2-Cub-GST*) were induced with
2% galactose for 1 h. Proteins extracts were prepared, separated by SDS-PAGE and analyzed by immunoblotting
using α-HA antibody. The signal intensities of the immunoreactive forms of full-length and cleaved Agp1, Gnp1,
Bap2, Ssy1, Can1, Lyp1, Hxt1 and Gal2 constructs were quantified. The fraction of split-ubiquitin cleavage was
determined; the mean values plotted with error bars showing standard deviation (n=3).

Figure 6. Progressivity of Shr3-Gap1 chaperone-substrate interactions

(A) Schematic diagram of split ubiquitin constructs including the gap1-Cub-GST, hxt1-Cub-GST and gal2-Cub-GST truncation constructs. (B) Strain FGY135 (*gap1Δ shr3Δ*) carrying pPL1262 (*SHR3-NubA*) and pIM1 (*gap1-2TM-Cub-GST*), pIM2 (*gap1-4TM-Cub-GST*), pIM3 (*gap1-6TM-Cub-GST*), pIM4 (*gap1-8TM-Cub-GST*), pIM5 (*gap1-10TM-Cub-GST*) or pIM16 (*gap1-12TM-Cub-GST*) (left panel), or pIM34 (*hxt1-2TM-Cub-GST*), pIM35 (*hxt1-4TM-Cub-GST*), pIM36 (*hxt1-6TM-Cub-GST*), pIM37 (*hxt1-8TM-Cub-GST*), pIM38 (*hxt1-10TM-Cub-GST*) or pIM39 (*hxt1-12TM-Cub-GST*) (center panel) or pIM40 (*gal2-2TM-Cub-GST*), pIM41 (*gal2-4TM-Cub-GST*), pIM42 (*gal2-6TM-Cub-GST*), pIM43 (*gal2-8TM-Cub-GST*), pIM44 (*gal2-10TM-Cub-GST*) or pIM45 (*gal2-12TM-Cub-GST*) (right panel) were induced with 2% galactose for 1 h. Extracts were prepared, separated by SDS-PAGE and analyzed by immunoblotting using α-HA antibody. (C) The signal intensities of the immunoreactive forms of uncleaved Cub constructs and cleaved interaction marker (GST-6xHA) were quantified; the mean values of the fraction of split ubiquitin cleavage is plotted with error bars showing standard deviation (n=3).

Figure 7. Progressive Shr3-Agp1 and Shr3-Ssy1 chaperone-substrate interactions

(A) Schematic diagram of agp1-Cub-GST truncation constructs. Strain FGY135 (*gap1Δ shr3Δ*) expressing *SHR3-NubA* (pPL1262) or *SHR3Δ94-NubA* (pAR76) and carrying pIM9 (*agp1-2TM-Cub-GST*), pIM10 (*agp1-4TM-Cub-GST*), pIM11 (*agp1-6TM-Cub-GST*), pIM12 (*agp1-8TM-Cub-GST*), pIM13 (*agp1-10TM-Cub-GST*) or pIM26 (*agp1-12TM-Cub-GST*) were induced with 2% galactose for 1 h. Extracts were prepared, separated by SDS-PAGE and analyzed by immunoblotting using α-HA antibody. The signal intensities of the immunoreactive forms of uncleaved Cub constructs and cleaved interaction marker (GST-6xHA) were quantified; the mean values of the fraction of split ubiquitin cleavage is plotted with error bars showing standard deviation (n=3). (B) Schematic diagram of ssy1-Cub-GST truncation constructs. Strain FGY135 (*gap1Δ shr3Δ*) expressing *SHR3-NubA* (pPL1262) or *SHR3Δ94-NubA* (pAR76) and carrying pIM20 (*ssy1-2TM-Cub-GST*), pIM21 (*ssy1-4TM-Cub-GST*), pIM22 (*ssy1-6TM-Cub-GST*), pIM23 (*ssy1-8TM-Cub-GST*), pIM24 (*ssy1-10TM-Cub-GST*) or pIM25 (*ssy1-12TM-Cub-GST*) were induced with 2% galactose for 1 h. Extracts were prepared, and analyzed as in (A) and the mean values of the fraction of split ubiquitin cleavage is plotted with error bars showing standard deviation (n=3).

Figure 8. Model of Shr3 facilitated AAP folding

Shr3 interacts transiently with AAP as they are co-translationally inserted into the ER membrane. Interactions start early, when 2-4MS have partitioned into the lipid bilayer, and continue until all MS are inserted. When AAP have fully integrated into the membrane and attain native conformations, the interactions with Shr3 diminish. The co-translational Shr3 function is specifically required for AAP folding and ER-exit. The chaperone activity depends on relatively few residues of the Shr3 sequence, suggesting that it functions as folding template. In the absence of Shr3, AAP are specifically retained in the ER and form high-molecular weight aggregates that are recognized as ERAD substrates.

895   **Supplementary Material**