Multilevel community assembly of the tadpole gut microbiome

Decio T. Correa[1,2*], David Rodriguez[3], Carine Emer[4], Daniel Saenz[5], Cory K. Adams[5], Luis

Schiesari[6], Mikhail Matz[1], Mathew A. Leibold[7]

1. Department of Integrative Biology, The University of Texas at Austin, Austin, TX, USA

2. Current affiliation: IBM Corporation, Chief Analytics Office, Armonk, NY, USA

3. Department of Biology, Texas State University, San Marcos, TX, USA

4. Chico Mendes Institute for Biodiversity Conservation, National Centre for Wild Birds

Research and Conservation, João Pessoa, PB, Brazil

5. Southern Research Station, US Forest Service, Nacogdoches, TX, USA

6. School of Arts, Sciences and Humanities, University of São Paulo, São Paulo, SP, Brazil

7. Department of Biology, University of Florida, Gainesville, FL, USA

* Corresponding author, decio.tadeu@gmail.com

Abstract

The assembly of local communities is likely to reflect the effects of local environmental

factors associated with filters that act at larger spatial scales. Dissecting these multiscale

effects remains a timely challenge that is particularly important for host-associated

microbiomes. We investigated the relative roles of local selection (due to host species

identity) and regional effects (due to water body identity) on the community structure

of bacteria in the gut of tadpoles from three biogeographic areas and used graph theory

and metanetwork approaches to explore and illustrate the distribution of bacteria

across different ponds. The pond of origin, which represents a regional species pool of

bacteria, was in general more important in shaping the gut microbiome of tadpoles than

host species identity. The resulting metanetworks are modular and indicate relatively

few species of bacteria occurring in more than one pond. Thus, each pond represents a

relatively distinct species pool of bacteria available for community assembly of the

tadpole microbiomes. Our findings indicate that microbiome community assembly in

amphibian larvae, as in many other communities, is a multiscale process with important

regional effects that constrain how local (i.e. host-dependent) filters act to influence

microbiome community composition.

Introduction

The origin and assembly of host-associated microbial communities

(microbiomes) is a complex process of fundamental importance in helping us

understand the symbiotic relationship between hosts and their microbes [1, 2]. Various

factors can play a role in the community assembly of microbiomes at multiple levels.

Perhaps the most obvious is the effect of host diet on the gut microbiome, which has

been shown in many organisms from insects to humans [3–6]. Other local factors

include host sex, health state, and genetics [7–11]. These effects can be thought of as

local 'filters' to microbial community assembly. Here we evaluate these effects as they

are determined by host species identity only, recognizing that there are likely to be

other more subtle effects involving inter-individual factors. However, there can also be

processes that constrain which microbes are available to colonize any given host. Here,

we focus on factors that determine the microbial species pool available to colonize the

host and evaluate these by the identity of the site of the host community.

We view community assembly of microbiomes as being driven by

metacommunity processes [12]. This recognizes that the assembly of a local community

depends strongly on the species available to colonize that community, known as the

regional species pool [13, 14]. From that pool, if a species is able to disperse to a given

community, its persistence in that community will depend on local demographic

processes and on the ability of the species to tolerate the local biotic and abiotic

conditions, which is known as the local selection component of community assembly

[15]. Therefore, communities with identical environmental conditions could have a

completely different species composition if colonized by different species pools [16].

While most work in metacommunity ecology has focused on macroorganisms, these

ideas should apply equally well to microbes [1, 2, 12, 17, 18].

Host-associated microbial communities appear to be influenced by effects

related to the individual host or individual species such as diet, physiology, and

genotype [10, 19, 20], as well as regional and historical processes, as in the case of

exposure to potentially different species pools of microorganisms [21–26], or vertical

and horizontal transmission of the microbiome [27, 28]. Considering the importance

that the microbiome can have to host health and well-being, understanding the relative

importance of effects acting at different levels is fundamental, but studies explicitly

tackling that question are still lacking [29].

We analyzed variation in tadpole microbiomes to evaluate the effects of local

environmental filters by focusing on host species identity, and regional effects by

focusing on the locality of the water body. We assume that different microbes would be

differentially favored by selection associated with the physiology and diet of different

tadpole species, and different microbes may be present at different sites due to either

site-specific habitat variables or to isolation and other differences among water bodies.

Our studies were conducted across three different geographic locations to test if they

show consistent patterns. We also used graph theory and metanetwork approaches to

explore the degree to which each pond consists of a unique species pool, and how they

are connected to each other in terms of shared bacterial species.

The gut microbiome of tadpoles is critically important to their digestive function

[30], and some evidence shows gut microbiomes from the same host species are more

similar to each other than from different species [31]. Further, there are often multiple

species of tadpole per water body, which represent different hosts with different

characteristics that are exposed to the same pool of bacteria. Many of these tadpole

species can occupy a variety of different water bodies, each with potentially distinct

bacterial species pools, which makes them a suitable system to test questions regarding

the effects of environment and host identity on microbiome assembly [32]. We evaluate

two hypotheses: i) if the bacterial community of each pond is different across the

landscape, then the pond of origin would be more important in determining the gut

microbiome of tadpoles, i.e., the microbiomes of different species within each pond

would be more similar to each other than the microbiomes of the same species across

distinct ponds, and the metanetworks would be structured into modules according to

each pond; ii) if ponds have very similar bacterial communities across the landscape,

then the species of tadpole would be more important in determining their gut

microbiome, i.e., tadpoles of the same species across ponds would have more similar

microbiomes than tadpoles of different species from the same pond, and the

metanetworks would be structured into modules according to each species.

Methods

    Study area

    We sampled tadpoles from 22 water bodies located in three different localities: i) six water bodies in the Boracéia Biological Station, located in the Atlantic Forest east of the state of São Paulo, Brazil, hereafter BR; ii) nine water bodies in the David Crockett and Angelina National Forests, located in the pine forest ecosystem in eastern Texas, United States, hereafter ET; and iii) seven water bodies in the Edwards Plateau, a mix of grassland and juniper/oak woodlands in the central part of Texas, United States, hereafter CT. We classified water bodies as either lentic or lotic. For ET ponds, we did a more thorough environmental characterization quantifying in each pond the dissolved oxygen (mg/L), pH, conductivity (uS/cm), chlorophyll a (ug/L), and water temperature (°C) using a Eureka multiparameter sonde, and the absence or presence of piscivorous (Bass) and/or insectivorous fish (Green Sunfish). Even though three water bodies in BR were lotic, we will refer to all of them as *ponds* for convenience.

    Tadpole and microbiome sampling

    In each pond we collected tadpoles with a dipnet that was swept through the pond. We then placed the larvae in a sterile plastic bag (Whirl Pak®) filled with water from their place of origin. Environmental microorganisms from water from CT and ET were sampled by filtering approximately 50 ml of water from three different places through a sterile syringe filter with a 0.45 μm cellulose acetate membrane (VWR). From

BR, approximately 2L of water was collected in a sterile plastic bag (Whirl Pak®), mixed,

and 100 mL was vacuum-filtered through a 0.45 μm cellulose acetate membrane. The

sediment in all ponds was sampled in three locations in each pond by removing the first

three centimeters with a sterile plastic straw of 1.3 cm of diameter and placed in a

sterile tube.

In the lab, tadpoles were euthanized by overdose in a solution of Milli-Q® filtered

water and tricaine methanesulfonate (MS 222). After the death was confirmed, each

individual was dissected using instruments that were cleaned in 100% ethanol and

flame-sterilized. The entire tadpole gut was removed and placed directly in a MoBio

PowerSoil® bead tube. The sediment from each pond was homogenized, centrifuged to

remove the excess water, and an aliquot of approximately 0.2 g was used for DNA

extraction. DNA from the water sample was extracted from the entire filter membrane.

We performed DNA extraction using MoBio PowerSoil® DNA isolation kit following the

manufacturer's recommendation. The same procedures without a real sample were

repeated to count as negative control samples, which did not have DNA when quantified

using Qubit™ dsDNA high sensitivity assay.

We amplified the V4 hypervariable region of the 16s rRNA gene (515F, 806F). CT

samples were amplified using the version F: GTGYCAGCMGCCGCGGTA / R:

GGACTACHVGGGTWTCTAAT and ET and BR the version F: GTGYCAGCMGCCGCGGTAA /

R: GGACTACNVGGGTWTCTAAT. The library preparation and sequencing (Illumina MiSeq

2x250bp) of the CT samples was realized at the Genomic Sequencing and Analysis

Facility (wikis.utexas.edu/display/GSAF) at The University of Texas at Austin following

their protocols. The library preparation and sequencing (Illumina MiSeq 2x250bp) of the

ET samples was realized at the Argonne National Laboratory (www.anl.gov) following

their protocols. For BR samples, we prepared the libraries following the Earth

Microbiome Project protocol (www.earthmicrobiome.org), with the exception of using

20uL PCR reactions and dual-index barcodes instead. Samples were sequenced in an

Illumina MiSeq platform (2x300 bp paired-end sequences).

The sequences were processed using the dada2 pipeline [33] in the software

Microsoft R Open v 3.5.1 [34]. Sequences from each site were processed separately.

Briefly, primers, adapters, and barcodes were removed and the quality profile of the

reads were visually inspected using the function plotQualityProfile aggregating over all

the samples from a site. BR, CT, and ET reads were truncated at 200 and 120, 200 and

180, and 200 and 160 forward and reverse reads, respectively. Sequencing errors were

calculated, sequences were clustered as Amplicon Sequence Variants based on the

DADA2 algorithm [33] and paired ends were merged. Chimeras were removed using the

consensus method in the function removeBimeraDenovo. Taxonomy was assigned

based on the Silva database v128 [35]. ASV sequences were exported to Qiime2 2018.4

[36], aligned using the MAFFT algorithm [37], and a phylogenetic tree was constructed

using FastTree [38]. We discarded all ASVs that were classified as Archaea, chloroplast,

mitochondria, or that were not at least assigned to bacteria. Hereafter, we will refer to

ASVs as species of bacteria.

Data analysis

    Diversity

    For each sample we constructed a rarefaction curve and compared richness and Faith's phylogenetic diversity [39]. To avoid sequencing depth effects, each sample was rarefied 100 times to the number of reads of the sample with the lowest coverage using the R package BAT [40]. For comparison, we considered the median richness from all rarefactions from a sample as our richness measure.

    Species and pond effects

    We analyzed our data as compositional data (termed CODA) [41]. It requires a centered log-ratio transformation of the read counts and therefore cannot have zeros, so we added a pseudocount of one to all ASVs in all samples. We transformed the data using the codaSeq.clr function from the R package CoDaSeq [42, 43]. We also analyzed data that takes into account phylogenetic relatedness of microbes by using the PHILR transformation from the R package philr [44]. This metric takes into account the phylogenetic relationship between the ASVs, which is equivalent to the Unifrac metric [45], but it considers the compositional nature of the data [44]. We then ran a Principal Component Analysis (PCA) on the Euclidean distance matrix of the CODA- and PHILR-transformed data to visualize the relationship between samples and tested if samples from tadpoles, water, and sediment are different from each other using a Permutational

Multivariate Analysis of Variance (perMANOVA). If a difference was detected we then

used additional perMANOVAs for pairwise comparisons with Bonferroni correction.

To first test if either the identity of the species of tadpole or the pond of origin

are more related to the gut microbiome composition we used a perMANOVA to fit a

model with both pond and species identity to the CODA- and PHILR-transformed data

using the function adonis2 from the R package vegan [46]. The significance of the

marginal effects was tested based on 999 permutations. In addition, to test for the

unique effects of the species of tadpole or the pond of origin as well their shared effects

on the gut microbiome we performed variance partitioning and Redundancy Analysis

(RDA) [47] using the function varpart in the R package vegan [46]. The significance of

each unique component in the variation partitioning was tested using the vegan

function anova.cca with 1000 permutations. We repeated perMANOVA and variation

partitioning analyses to the data grouping the species of bacteria at Genus and Family

level.

Inter-pond variation

We also investigated if there was evidence for dispersal limitation in the bacteria

from water and sediment using multiple regression on distance matrices using the

function MRM from the R package ecodist [48]. We used the Euclidean distance matrix

based on the PHILR and CODA transformations as the dependent variable and a

geographical distance matrix between sites as the independent variable. In addition, to

account for environmental effects, we also include in the model a distance matrix based

on the type of the pond (lentic or lotic) for BR, and a distance matrix based on the

environmental variables measured for ET. The latter was a Gower distance matrix [49]

created by the function dist.ktab from the R package ade4 [50]. Quantitative variables

were Gower-standardized (divided by maximum minus minimum). All CT ponds were

lentic and there were no other variables besides spatial distance to include in the

model. For models where distance was significant in the MRMs, we applied a Partial

Mantel Correlogram approach using the function mpmcorrelogram from the R package

with the same name [51]. That tests, at several distance classes, in our case estimated

by Sturge's rule [52], the relationship between water and sediment microbiome with

geographical distance while controlling for environmental effects.

Metanetworks

We further explored the structure of three different metanetworks within each

locality (BR, CT, and ET) to understand the distribution of bacteria across ponds. Shared

species (here bacteria) connect different ponds in a metacommunity perspective [53,

54]. We built three bacteria metanetworks for each locality: (i) the tadpole microbiome

metanetwork, composed by the bacteria living in the gut of all tadpoles within a pond,

independently of the tadpole species, (ii) the water metanetwork, with bacteria found in

the water, and (iii) the sediment metanetwork, composed by bacteria found in the

sediment of each pond. Then, we built an a_{mn} adjacency matrix for each metanetwork in

which *m* corresponds to a single pond (i.e., considering bacteria from water, sediment

or gut of tadpoles), and *n* to each bacterium. The *mn* element represents the

presence/absence of species of bacteria *n* in a pond *m,* and is represented by a link in

the graphical representation. Therefore, each node in our metanetwork is either a

bacterium or a pond; when the same bacterium is found in more than one pond, there

is a link connecting them, forming the edges of the metanetwork.

To test whether ponds consist of different bacterial communities, we calculated

the modularity of the metanetworks using the FastGreedy algorithm [55, 56] in the

software Modular v 0.1 [57]. We tested the significance of the observed modularity

against two different null models with 100 permutations each: the Erdős-Rényi [58], in

which pond-bacteria interactions are connected randomly, and the 'null model 2' [59],

in which the probability of a pond-bacteria interaction is proportional to their number of

links in the observed matrix. To identify the bacteria with higher potential to connect

the metanetwork through shared occurrence in a higher number of ponds, we

calculated the degree (k) for each metanetwork using the software Pajek v. 4.10 [60].

Degree measures how many links each node establishes in its correspondent network.

For the bacteria, it represents the number of ponds in which it occurs; for the pond, it

represents the number of bacteria it has (from tadpoles, water or sediment). For each

metanetwork we also calculated the proportion of bacteria found uniquely in p_1, p_2, …,

p_n, where n is the total number of ponds in one locality.

Results

Diversity

We registered 19 species of tadpole in all ponds: three in CT, six in ET, and

eleven in BR (Table S1). After the filtering steps, there were a total of 533 777, 1 289

662, and 2 064 298 16s rRNA gene fragment reads for CT, ET, and BR, respectively. The

rarefaction curve shows that all samples reached an asymptote, even the ones with the

lowest number of reads (Figure S1). For richness and phylogenetic diversity measures of

bacteria, CT, ET, and BR samples were rarefied to 3 950, 1 791, and 3 916 reads,

respectively. There was no significant difference among CT samples in terms of richness,

but there was in terms of phylogenetic diversity, with sediment samples having the

highest diversity and *Rana berlandieri* tadpoles the lowest (Figure S2). Sediment

samples had the highest richness and phylogenetic diversity in ET and *Hylodes phyllodes*

the highest in BR (Figures S3 and S4). A great part of the tadpole microbiome was

composed by Fusobacteria, Firmicutes, Proteobacteria, and Bacteroidetes, while

sediment and water samples were mostly composed by Proteobacteria (Figures S5-S7).

Species and pond effects

The composition of the tadpole gut microbiome is different from both water and

sediment samples in all localities (Figure 1, Figure S8, Table S2). Also, in most cases,

tadpole gut microbiome samples cluster more with other samples from the same pond

than with samples from the same species from a different pond (Figure 1, Figures S8-

S10). However, within a pond, there can be separation by species and, in fact, the

perMANOVA shows that both species identity of tadpole and pond of origin are

significant predictors of the tadpole gut microbiome in all localities when using either

PHILR- or CODA-transformed data, except for CT CODA, in which species identity was

only marginally significant (Table 1). In CT and ET, pond of origin was a stronger

predictor of the tadpole gut microbiome than tadpole species identity. The overall

results are consistent over all the taxonomic levels considered (Tables S3-S6).

For CT and ET, the pond of origin explained considerably more of the variance in

tadpole gut microbiome than the species of tadpole (Figure 2). For BR, the tadpole

species explained a slightly higher proportion of the variance in their gut microbiome.

However, most of the variation in that environment is explained by the joint effects of

tadpole species and pond of origin, i.e., the variation that could not be separated in

unique components (Figure 2). Again, the overall results are comparable across all the

taxonomic levels considered (Tables S7-S8).

Inter-pond variation

Only the bacterial communities of water from CT were significantly related to the

spatial distance between ponds (Table S9). However, that was the only dataset where

environmental data is not available and could not be accounted for, allowing for the

possibility that such distance effects are due to spatially structured environmental

factors. Partial Mantel Correlograms showed that, for both CODA- (p_{adj} = 0.044) and

PHILR-transformed data (p_{adj} = 0.022), there is a spatial signal only for the first distance

range, from 174 to 6 223 m. ET water and sediment communities were significantly

related to environmental variables for PHILR-transformed data and BR water

communities were significantly related to both PHILR- and CODA-transformed data

(Table S9).

Metanetworks

All metanetworks are significantly modular according to the two null models

used ($p < 0.001$ for all null models; Figure 3, Figure S11). In most cases, each module

corresponds to a single pond (Figure 3, Figure S11 and Table S10). Metanetworks of

water and sediment are more modular than those of tadpoles for all localities (Figure 3,

Figure S11). In all cases, the vast majority of bacteria occur in a single pond (k = 1),

varying from 63.2% in East Texas to 91.3% in Brazil (Figure 4, Figures S12-S16). Only 0.1

to 1.1% of bacteria were present in all ponds (Figure 4, Figure S12-S16). From those, for

BR and ET they were mostly from the phyla Bacteroidetes and Proteobacteria and

Orders Bacteroidales and Desulfovibrionales. CT had only three bacteria present across

all ponds. From the ponds' perspective, their degree, which corresponds to the number

of unique links they have in the metanetwork, varied from 118 to 615 for water, 90 to

478 for sediment, and 241 to 2 751 for bacteria from tadpoles (Figures S17-S19).

Discussion

We found that microbial community assembly of tadpole microbiomes was

structured by processes happening at different levels and broadly resemble findings

from macroorganisms [1, 15, 61–63]. At higher levels, there are inter-pond differences

in microbial communities due to site-specific variables, isolation, and/or stochasticity. It

was not our goal to understand the drivers of inter-pond differences, which is ultimately

responsible for the different bacterial species pool tadpoles are exposed to. However, in

some cases, for the ponds where we had environmental data there is a relationship

between environmental variables and the bacterial community of water and/or

substrate. In these cases, each pond could act as a regional scale environmental filter for

bacteria. For CT, we found some evidence for a spatial correlation in bacterial

communities among nearby ponds, which could indicate dispersal limitation structuring

the pond communities, but that was the only dataset for which we did not have

environmental variables. Nevertheless, independently of the causes of the differences

among ponds, each one of them constitutes a unique species pool of colonizers for the

microbiomes of tadpoles. As tadpoles develop, the community assembly of their

microbiome is restricted by that species pool of microbes. Each tadpole comprises a

smaller-scale filter to a subset of the microbes within the species pool. Therefore, even

if similar habitats (same species of tadpole in our case) are exposed to different species

pools, they will not end up with the same microbial community.

The composition and diversity of the gut microbiome of tadpoles can be variable

across species. The microbiome of tadpoles from this study is slightly different than

previous studies. Proteobacteria and Firmicutes dominated in other studies [31, 64–66]

but here we found that Fusobacteria were also important. Alpha and phylogenetic

diversity were also mostly higher in our study than in previous ones [31, 64–66]. The

knowledge about the gut microbiome of tadpoles is still limited to a few species and

quantitative comparisons of diversity across studies is problematic because of

differences in sampling procedures, sample processing, data cleaning, and the

algorithms and databases. However, overall differences in microbiome composition

could be due to the exposure of tadpoles to different pools of colonist microbes.

The pond of origin of the tadpoles was more important in determining their gut

microbiome than the species of the tadpole in both of the Texas locations. We assume

the same species of tadpole has similar requirements in terms of microbes and/or

constitutes a similar patch for colonization by microbes. There were only a few bacteria

in common across all ponds in all three localities, therefore each pond constitutes a

unique bacterial species pool from which the tadpoles can be colonized. In Brazil, the

pond of origin and species of tadpole showed similar relationships with the gut

microbiome of tadpoles. However, most of variation in Brazil was due to variation that

could not be separately attributed to either because of the uniquely distinct tadpole

faunas of ponds.

The limited and unique pool of bacteria tadpoles are exposed to and the higher

similarity within a pond can be visualized on the metanetworks of bacteria from water,

sediment, and tadpole gut. The metanetworks are modular and each pond constitutes a

unique community of bacteria. The vast majority of bacteria occur only in a single pond,

and just a few are present in all of them. In fact, for water and sediment, the maximum

number of bacteria found in all ponds was only eight. That number would probably

increase if more sediment or water was sampled. On the other hand, we likely sampled

all species of tadpoles from each water body and the proportion of bacteria from their

gut found in all water bodies is still very low (0.1-1.1%) despite similar phyla found in all

of them. The major bacteria in tadpole guts belong to orders Bacteroidales and

Desulfovibrionales, which do not overlap with the most widespread taxa found in water

or sediment. It is common to not find much overlap between bacteria from environment

and gut of aquatic organisms [67, 68]. Likewise, we show that the gut microbiome of

tadpoles is distinct from both water and sediment from the ponds suggesting that the

gut constitutes a strong environmental filter that favors certain taxa. Desulfovibrionales,

for example, is a group of sulfur-reducing bacteria that is mostly anaerobic [69].

Therefore, at least certain parts of the digestive tract of tadpoles might be anaerobic

[31, 70]. Pryor and Bjorndal [30] detected significant levels of fermentation in the

hindgut of bullfrog tadpoles. It is unknown to which extent other species of tadpoles are

hindgut fermenters. For those that are, even though they might not have a core

microbiome, at least not at fine-scale taxonomy levels, there may be a functional core

gut microbiome.

In terrestrial and aquatic environments, the importance of the regional species

pool in shaping the microbial communities can be noticed even when not tested

directly. In many species, from insects to humans, individuals likely exposed to the same

regional species pool, such as being from the same environment, have more similar

microbiomes [21–26]. There are many difficulties in quantifying and testing how

regional effects affect community assembly [29, 62], but comparing results with

predictions from theory can lead to important insights. If hosts from different

environments have similar microbiomes it could indicate that (i) there is not much

difference in the regional species pools between environments, (ii) the hosts move

between species pools, or (iii) there are other methods for microbiome acquisition such

as transmission from conspecifics. For example, Kueneman *et al*. [71] and McKenzie *et*

al. [32] studying the skin microbiome of tadpoles found that the identity of the species

was the strongest predictor of the skin microbiome, with the water body of origin

explaining additional [71] or no variance at all [32]. Vertical transmission of the

microbiome is unlikely in amphibians [72], and they probably acquire their microbiome

from the environment at every generation [73]. Therefore, the results from Kueneman

et al. [71] and McKenzie *et al*. [32] indicate there is at least a partially shared species

pool of microbes across their sampling sites.

The enormous intra and inter-specific variation in the microbiomes of different

species of animals and plants can be at least partially explained by differences in

regional species pool during the community assembly of the microbiome. The gut

microbiome of animals can be stable over time [63, 74] and can also be resistant to

invasions and colonization by other bacteria [75–77]. Therefore, if the community

assembly of individuals of the same species occurred while they were exposed to

different species pools, historical effects can cause lasting effects in the development of

the microbiome [78]. Moreover, recent efforts to manipulate the microbiome of plants

and animals to improve health and/or increase productivity might not be as effective if

the manipulation is done only at the level of the individual host, i.e., at the local scale.

As stated by Chase [16] "if local and regional processes determine the community

composition, then both processes need to be restored to achieve the desired

community". Our results indicate that the same logic is valid for microbes.

In summary, we showed that local assembly of host-associated bacterial

communities are affected by regional scale processes, more specifically changes in the

regional species pool of colonizers. The inter-pond variation in bacterial community can

be due to stochastic, historical, spatial, and/or environmental processes or even the

result of local evolution [79]. Nevertheless, microbiome assembly, as any other

community, is a multiscale process. Microbial and microbiome ecology could be better

linked to ecological theory by considering the multiscale dynamics of community

assembly [12, 17, 80, 81]. Because it is much easier to quantify effects at the scale of the

individual host or host species, i.e., local effects [29, 62], the role of regional effects is

probably underestimated. Regional effects, however, appear to be a fundamental piece

of the community assembly puzzle that can be used to understand variation in

microbiomes.

Acknowledgements

We thank Ananda Brito de Assis and Carlos Arturo Navas for laboratory access

during field work in Brazil and Christopher Schalk and James Childress for help in the

field in East Texas. We thank the Zoology Museum of University of São Paulo (MZUSP),

The National Forests and Grasslands of Texas, and several private landowners for

granting permission to access sampling sites and ICMBio (47263-1), Texas Parks and

Wildlife (SPR-0414-067), and IACUC (AUP-2014-00286) for collecting and animal care

permits. Funding was provided by Conselho Nacional de Desenvolvimento Científico e

Tecnológico (CNPq, Brasil, grant number 458796/2014-0) and Texas Ecolab. DTC

received a Science Without Borders scholarship during the research (CAPES 1176/13-7).

Finally, we thank Daniel Bolnick, Michel Ryan, Ulrich Mueller, and members of Leibold

and Matz labs for comments throughout the project.

References

1. Costello EK, Stagaman K, Dethlefsen L, Bohannan BJM, Relman DA. The
application of ecological theory toward an understanding of the human
microbiome. *Science* 2012; **336**: 1255–62.
2. Christian N, Whitaker BK, Clay K. Microbiomes: unifying animal and plant systems
through the lens of community ecology theory. *Front Microbiol* 2015; **6**: 1–15.
3. Broderick NA, Raffa KF, Goodman RM, Handelsman J. Census of the Bacterial
Community of the Gypsy Moth Larval Midgut by Using Culturing and Culture-
Independent Methods. *Appl Environ Microbiol* 2004; **70**: 293–300.
4. Turnbaugh PJ, Ridaura VK, Faith JJ, Rey FE, Knight R, Gordon JI. The Effect of Diet
on the Human Gut Microbiome: A Metagenomic Analysis in Humanized
Gnotobiotic Mice. *Sci Transl Med* 2009; **1**: 6ra14-6ra14.
5. Bolnick DI, Snowberg LK, Hirsch PE, Lauber CL, Knight R, Caporaso JG, et al.
Individuals' diet diversity influences gut microbial diversity in two freshwater fish
(threespine stickleback and Eurasian perch). *Ecol Lett* 2014; **17**: 979–87.
6. Gomez A, Rothman JM, Petrzelkova K, Yeoman CJ, Vlckova K, Umaña JD, et al.
Temporal variation selects for diet–microbe co-metabolic traits in the gut of
Gorilla spp. *ISME J* 2016; **10**: 514–526.
7. Fierer N, Hamady M, Lauber CL, Knight R. The influence of sex, handedness, and
washing on the diversity of hand surface bacteria. *Proc Natl Acad Sci U S A* 2008;
105: 17994–9.
8. Pflughoeft KJ, Versalovic J. Human Microbiome in Health and Disease. *Annu Rev
Pathol Mech Dis* 2012; **7**: 99–122.
9. Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. Diversity, stability
and resilience of the human gut microbiota. *Nature* 2012; **489**: 220–30.
10. Goodrich JK, Waters JL, Poole AC, Sutter JL, Koren O, Blekhman R, et al. Human
genetics shape the gut microbiome. *Cell* 2014; **159**: 789–799.
11. Bolnick DI, Snowberg LK, Hirsch PE, Lauber CL, Org E, Parks B, et al. Individual diet
has sex-dependent effects on vertebrate gut microbiota. *Nat Commun* 2014; **5**:
4500.
12. Miller ET, Svanbäck R, Bohannan BJM. Microbiomes as Metacommunities:
Understanding Host-Associated Microbes through Metacommunity Ecology.
Trends Ecol Evol 2018; **33**: 926–935.
13. Ricklefs RE. Community Diversity: Relative Roles of Local and Regional Processes.
Science (80-) 1987; **235**: 167–171.
14. Leibold MA, Holyoak M, Mouquet N, Amarasekare P, Chase JM, Hoopes MF, et al.
The metacommunity concept: a framework for multi-scale community ecology.
Ecol Lett 2004; **7**: 601–613.
15. Vellend M. Conceptual synthesis in community ecology. *Q Rev Biol* 2010; **85**: 183–
206.

16. Chase JM. Community assembly: when should history matter? *Oecologia* 2003;
136: 489–98.
17. Prosser JI, Bohannan BJM, Curtis TP, Ellis RJ, Firestone MK, Freckleton RP, et al.
The role of ecological theory in microbial ecology. *Nat Rev Microbiol* 2007; **5**:
384–392.
18. Fierer N, Ferrenberg S, Flores GE, González A, Kueneman J, Legg T, et al. From
Animalcules to an Ecosystem: Application of Ecological Concepts to the Human
Microbiome. *Annu Rev Ecol Evol Syst* 2012; **43**: 137–155.
19. Phillips CD, Phelan G, Dowd SE, McDonough MM, Ferguson AW, Delton Hanson J,
et al. Microbiome analysis among bats describes influences of host phylogeny, life
history, physiology and geography. *Mol Ecol* 2012; **21**: 2617–2627.
20. Fitzpatrick BM, Allison AL. Similarity and differentiation between bacteria
associated with skin of salamanders (*Plethodon jordani*) and free-living
assemblages. *FEMS Microbiol Ecol* 2014; **88**: 482–94.
21. Staubach F, Baines JF, Künzel S, Bik EM, Petrov DA. Host Species and
Environmental Effects on Bacterial Communities Associated with Drosophila in
the Laboratory and in the Natural Environment. *PLoS One* 2013; **8**: e70749.
22. Schloss PD, Iverson KD, Petrosino JF, Schloss SJ. The dynamics of a family's gut
microbiota reveal variations on a theme. *Microbiome* 2014; **2**: 25.
23. McCord AI, Chapman CA, Weny G, Tumukunde A, Hyeroba D, Klotz K, et al. Fecal
microbiomes of non-human primates in Western Uganda reveal species-specific
communities largely resistant to habitat perturbation. *Am J Primatol* 2014; **76**:
347–354.
24. Eichmiller JJ, Hamilton MJ, Staley C, Sadowsky MJ, Sorensen PW. Environment
shapes the fecal microbiome of invasive carp species. *Microbiome* 2016; **4**: 44.
25. Hernández-Gómez O, Hoverman JT, Williams RN. Cutaneous Microbial
Community Variation across Populations of Eastern Hellbenders (*Cryptobranchus
alleganiensis alleganiensis*). *Front Microbiol* 2017; **8**: 1–16.
26. Muletz Wolz CR, Yarwood SA, Campbell Grant EH, Fleischer RC, Lips KR. Effects of
host species and environment on the skin microbiome of Plethodontid
salamanders. *J Anim Ecol* 2018; **87**: 341–353.
27. Sanders JG, Powell S, Kronauer DJC, Vasconcelos HL, Frederickson ME, Pierce NE.
Stability and phylogenetic correlation in gut microbiota: Lessons from ants and
apes. *Mol Ecol* 2014; **23**: 1268–1283.
28. Moeller AH, Suzuki TA, Phifer-Rixey M, Nachman MW. Transmission modes of the
mammalian gut microbiota. *Science (80-)* 2018; **362**: 453–457.
29. Lindström ES, Langenheder S. Local and regional factors influencing bacterial
community assembly. *Environ Microbiol Rep* 2012; **4**: 1–9.
30. Pryor GS, Bjorndal KA. Symbiotic fermentation, digesta passage, and
gastrointestinal morphology in bullfrog tadpoles (*Rana catesbeiana*). *Physiol
Biochem Zool* 2005; **78**: 201–215.

31. Vences M, Lyra ML, Kueneman JG, Bletz MC, Archer HM, Canitz J, et al. Gut
bacterial communities across tadpole ecomorphs in two diverse tropical anuran
faunas. *Sci Nat* 2016; **103**: 25.
32. McKenzie VJ, Bowers RM, Fierer N, Knight R, Lauber CL. Co-habiting amphibian
species harbor unique skin bacterial communities in wild populations. *ISME J*
2012; **6**: 588–96.
33. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
High-resolution sample inference from Illumina amplicon data. *Nat Methods*
2016; **13**: 581–583.
34. Microsoft and R Core Team. Microsoft R Open. 2018. Redmond, Washington.
35. Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, et al. The SILVA and
"All-species Living Tree Project (LTP)" taxonomic frameworks. *Nucleic Acids Res*
2014; **42**: D643–D648.
36. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet C, Al-Ghalith GA, et al. QIIME
2: Reproducible, interactive, scalable, and extensible microbiome data science.
PeerJ Prepr 2018; **6**: e27295v2.
37. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. *Mol Biol Evol* 2013; **30**: 772–780.
38. Price MN, Dehal PS, Arkin AP. FastTree 2 - Approximately maximum-likelihood
trees for large alignments. *PLoS One* 2010; **5**.
39. Faith DP. Conservation evaluation and phylogenetic diversity. *Biol Conserv* 1992;
61: 1–10.
40. Cardoso P, Rigal F, Carvalho JC. BAT - Biodiversity Assessment Tools, an R package
for the measurement and estimation of alpha and beta taxon, phylogenetic and
functional diversity. *Methods Ecol Evol* 2015; **6**: 232–236.
41. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome Datasets
Are Compositional: And This Is Not Optional. *Front Microbiol* 2017; **8**: 2224.
42. Gloor GB, Reid G. Compositional analysis: a valid approach to analyze microbiome
high-throughput sequencing data. *Can J Microbiol* 2016; **62**: 692–703.
43. Gloor GB, Wu JR, Pawlowsky-Glahn V, Egozcue JJ. It's all relative: analyzing
microbiome data as compositions. *Ann Epidemiol* 2016; **26**: 322–329.
44. Silverman JD, Washburne AD, Mukherjee S, David LA. A phylogenetic transform
enhances analysis of compositional microbiota data. *Elife* 2017; **6**: 1–20.
45. Lozupone C, Knight R. UniFrac: a new phylogenetic method for comparing
microbial communities. *Appl Environ Microbiol* 2005; **71**: 8228–35.
46. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:
Community Ecology Package. 2018.
47. Legendre P, Legendre L. Numerical ecology, 3rd ed. 2012. Elsevier.
48. Goslee SC, Urban DL. The ecodist Package for Dissimilarity-based Analysis of
Ecological Data. *J Stat Softw* 2007; **22**.
49. Pavoine S, Vallet J, Dufour A-B, Gachet S, Daniel H. On the challenge of treating

various types of variables: application for improving the measurement of
functional diversity. *Oikos* 2009; **118**: 391–402.

50.   Dray S, Dufour A. The ade4 package: implementing the duality diagram for
ecologists. *J Stat Softw* 2007; **22**: 1–20.

51.   Matesanz S, Gimeno TE, de la Cruz M, Escudero A, Valladares F. Competition may
explain the fine-scale spatial patterns and genetic structure of two co-occurring
plant congeners. *J Ecol* 2011; **99**: 838–848.

52.   Borcard D, Gillet F, Legendre P. Numerical Ecology with R. 2011. Springer New
York, New York, NY.

53.   Hagen M, Kissling WD, Rasmussen C, De Aguiar MAM, Brown LE, Carstensen DW,
et al. Biodiversity, Species Interactions and Ecological Networks in a Fragmented
World. *Advances in Ecological Research*, 1st ed. 2012. Elsevier Ltd., pp 89–210.

54.   Emer C, Galetti M, Pizo MA, Guimarães PR, Moraes S, Piratelli A, et al. Seed-
dispersal interactions in fragmented landscapes – a metanetwork approach. *Ecol*
Lett 2018; **21**: 484–493.

55.   Clauset A, Newman MEJ, Moore C. Finding community structure in very large
networks. *Phys Rev E* 2004; **70**: 066111.

56.   Wakita K, Tsurumi T. Finding Community Structure in Mega-scale Social Networks.
WWW. 2007. pp 1275–1276.

57.   Marquitti FMD, Guimarães PR, Pires MM, Bittencourt LF. MODULAR: Software for
the autonomous computation of modularity in large network sets. *Ecography*
(Cop) 2014; **37**: 221–224.

58.   Erdős P, Rényi A. On random graphs I. *Publ Math* 1959; **6**: 290–297.

59.   Bascompte J, Jordano P, Melian CJ, Olesen JM. The nested assembly of plant-
animal mutualistic networks. *Proc Natl Acad Sci* 2003; **100**: 9383–9387.

60.   Batagelj V, Mrvar A. Pajek - Analysis and Visualization of Large Network. In:
Juenger M, Mutzel P (eds). *Graph Drawing Software*. 2003. Springer, Berlin, pp
77–103.

61.   Emerson BC, Gillespie RG. Phylogenetic analysis of community assembly and
structure over space and time. *Trends Ecol Evol* 2008; **23**: 619–630.

62.   Cornell H V., Harrison SP. What Are Species Pools and When Are They Important?
Annu Rev Ecol Evol Syst 2014; **45**: 45–67.

63.   Coyte KZ, Schluter J, Foster KR. The ecology of the microbiome: Networks,
competition, and stability. *Science (80-)* 2015; **350**: 663–666.

64.   Kohl KD, Cary TL, Karasov WH, Dearing MD. Restructuring of the amphibian gut
microbiota through metamorphosis. *Environ Microbiol Rep* 2013; **5**: 899–903.

65.   Chai L, Dong Z, Chen A, Wang H. Changes in intestinal microbiota of *Bufo*
gargarizans and its association with body weight during metamorphosis. *Arch*
Microbiol 2018; **0**: 1–13.

66.   Lyra ML, Bletz MC, Haddad CFB, Vences M. The Intestinal Microbiota of Tadpoles
Differs from Those of Syntopic Aquatic Invertebrates. *Microb Ecol* 2018; **76**: 121–

124.

67.   Sullam KE, Essinger SD, Lozupone CA, O'Connor MP, Rose GL, Knight R, et al.
Environmental and ecological factors that shape the gut bacterial communities of
fish: a meta-analysis. *Mol Ecol* 2012; **21**: 3363–3378.

68.   Parris DJ, Brooker RM, Morgan MA, Dixson DL, Stewart FJ. Whole gut microbiome
composition of damselfish and cardinalfish before and after reef settlement.
PeerJ 2016; **4**: e2412.

69.   Madigan MT, Martinko JM, Bender KS, Buckley DH, Stahl DA. Brock Biology of
Microorganisms. *International Microbiology* . 2015. Pearson.

70.   Pryor G. Anaerobic bacteria isolated from the gastrointestinal tracts of bullfrog
tadpoles (*Rana catesbeiana*). *Herpetol Conserv Biol* 2008; **3**: 176–181.

71.   Kueneman JG, Parfrey LW, Woodhams DC, Archer HM, Knight R, McKenzie VJ. The
amphibian skin-associated microbiome across species, space and life history
stages. *Mol Ecol* 2014; **23**: 1238–1250.

72.   Hughey MC, Delia J, Belden LK. Diversity and stability of egg-bacterial
assemblages: The role of paternal care in the glassfrog *Hyalinobatrachium*
colymbiphyllum. *Biotropica* 2017; **49**: 792–802.

73.   Walke JB, Becker MH, Loftus SC, House LL, Cormier G, Jensen R V, et al.
Amphibian skin may select for rare environmental microbes. *ISME J* 2014; **8**:
2207–2217.

74.   Faith JJ, Guruge JL, Charbonneau M, Subramanian S, Seedorf H, Goodman AL, et
al. The Long-Term Stability of the Human Gut Microbiota. *Science (80-)* 2013;
341: 1237439–1237439.

75.   Britton RA, Young VB. Interaction between the intestinal microbiota and host in
Clostridium difficile colonization resistance. *Trends Microbiol* 2012; **20**: 313–319.

76.   Risely A, Waite D, Ujvari B, Klaassen M, Hoye B. Gut microbiota of a long-distance
migrant demonstrates resistance against environmental microbe incursions. *Mol*
Ecol 2017; **26**: 5842–5854.

77.   Zmora N, Zilberman-Schapira G, Suez J, Mor U, Dori-Bachash M, Bashiardes S, et
al. Personalized Gut Mucosal Colonization Resistance to Empiric Probiotics Is
Associated with Unique Host and Microbiome Features. *Cell* 2018; **174**: 1388-
1405.e21.

78.   Fukami T. Historical Contingency in Community Assembly: Integrating Niches,
Species Pools, and Priority Effects. *Annu Rev Ecol Evol Syst* 2015; **46**: 1–23.

79.   Urban MC, De Meester L. Community monopolization: local adaptation enhances
priority effects in an evolving metacommunity. *Proc Biol Sci* 2009; **276**: 4129–38.

80.   Mihaljevic JR. Linking metacommunity theory and symbiont evolutionary ecology.
Trends Ecol Evol 2012; **27**: 323–9.

81.   Shade A, Dunn RR, Blowes SA, Keil P, Bohannan BJM, Herrmann M, et al.
Macroecology to Unite All Life, Large and Small. *Trends Ecol Evol* 2018; **33**: 731–
744.

Tables

Table 1. Results of a Permutational Analysis of Variance (perMANOVA) testing if the
species of tadpole or the pond of origin are related to the gut microbiome composition
of tadpoles using the bacteria as Amplicon Sequence Variants (ASVs). Models were
fitted to CODA and PHILR transformed data. The significance of the marginal effects was
tested based on 999 permutations. BR = Brazil, CT = Central Texas, ET = East Texas.
635

		PHILR			CODA		
		R^2	F	p	R^2	F	p
BR	Water body$_{(4,80)}$	0.08	3	0.005	0.1	2.7	0.001
	Species$_{(9,80)}$	0.15	2.5	0.002	0.16	2.1	0.001
CT	Water body$_{(5,25)}$	0.48	6	0.001	0.41	3.6	0.001
	Species$_{(1,25)}$	0.05	2.8	0.010	0.04	1.8	0.056
ET	Water body$_{(8,76)}$	0.27	4.7	0.001	0.24	3.2	0.001
	Species$_{(5,76)}$	0.18	5.1	0.001	0.12	2.6	0.001

Figures

Figure 1. Plot of the first two axes of a Principal Component Analysis on CODA
transformed microbiome data (amplicon sequence variants) from tadpole samples from
Brazil (BR), Central Texas (CT), and Eastern Texas (ET). Panels on the left show sediment,
water, and tadpole samples. Panels on the right show samples from tadpoles only. Left
and right panels are results from separate analyses. Shapes represent different species
of tadpoles and colors represent water body of origin. See Table S1 for abbreviations.

Figure 2. Results of a variation partitioning analysis testing the unique effects of the
species of tadpole or pond of origin as well their shared effects on the tadpole gut
microbiome considering the bacteria as Amplicon Sequence Variants (ASVs) for CODA-
and PHILR-transformed data. The effects of pond of origin and species of tadpole were
all significant ($p < 0.05$) based on 1000 permutations, except for the shared component,
which cannot be tested.

Figure 3. Microbiome metanetworks of the shared bacteria (amplicon sequence variant -
ASV) found in the gut of tadpoles. Larger nodes represent each water body for each
locality, *i.e.*, Brazil (BR), Central Texas (CT), and East Texas (ET). Smaller nodes represent
a unique bacterium that can be found in a single water body (link between smaller and
larger nodes) or shared among ponds thus connecting them. Numbers on the bottom-
right are the modularity values for each metanetwork. Distinct colors represent
different modules which, with a few exceptions, correspond to a single water body
(Table S10).

Figure 4. Number of bacteria from the gut of tadpoles for each degree (k) on the left and Order of bacteria present in the highest degree on the right. Each degree represents one pond. Bacteria found in only one pond are represented in the degree category 1. Likewise, bacteria present in tadpoles across all ponds are represented in the highest category of each histogram. The Order of bacteria present in tadpoles across all ponds from each locality is shown on the barplot on the right. Same color across barplots represent the same Order. See Figures S11-S16 for water and sediment metanetworks. BR = Brazil, CT = Central Texas, ET = East Texas.