MDA5 is an essential vita-PAMP sensor necessary for host resistance against *Aspergillus fumigatus*[1]

Xi Wang[*,§], Alayna K. Caffrey-Carr[*,†,2, §], Ko-wei Liu[*], Vanessa Espinosa[‡], Walburga Croteau[*], Sourabh Dhingra[*,3], Amariliz Rivera[‡], Robert A. Cramer[*], Joshua J. Obar[*,#]

[*]Geisel School of Medicine at Dartmouth, Department of Microbiology & Immunology, Lebanon, NH 03756, United States

[†]Montana State University, Department of Microbiology & Immunology, Bozeman, MT 59718, United States

[‡]Center for Immunity & Inflammation, Rutgers - New Jersey Medical School, Newark, NJ, 07103, United States

[§]These authors contributed equally to this work

Conflict of Interests: The authors have declared that no conflict of interest exists.

[#]Corresponding author:
Joshua J. Obar
Geisel School of Medicine at Dartmouth
Department of Microbiology & Immunology
1 Medical Center Drive
Lebanon, NH 03756
Fax: +1 (603) 650-6223
Telephone: +1 (603) 650-6858
Email: joshua.j.obar@dartmouth.edu

<u>ABSTRACT</u>

RIG-I like receptors (RLR) are cytosolic RNA sensors that signal through the MAVS adaptor to

activate interferon responses against viruses. Whether the RLR family has broader effects on

host immunity against other pathogen families remains to be fully explored. Herein we

demonstrate that MDA5/MAVS signaling was essential for host resistance against pulmonary

Aspergillus fumigatus challenge through the regulation of antifungal leukocyte responses.

Activation of MDA5/MAVS signaling was driven by dsRNA from live *A. fumigatus* serving as a

key vitality-sensing pattern-recognition receptor. Interestingly, induction of type I interferons

after *A. fumigatus* challenge was only partially dependent on MDA5/MAVS signaling, whereas

type III interferon expression was entirely dependent on MDA5/MAVS signaling. Ultimately,

type I and III interferon signaling drove the expression of CXCL10, which is critical in resistance

against invasive aspergillosis in transplant patients. Furthermore, the MDA5/MAVS-dependent

interferon response was critical for the induction of optimal antifungal neutrophil killing of *A.*

fumigatus spores. In conclusion, our data broaden the role of the RLR family to include a role in

regulating antifungal immunity against *A. fumigatus*.

INTRODUCTION

Aspergillus fumigatus is a ubiquitous environmental mold that humans inhale on a daily basis. Multiple environmental surveys have demonstrated a range of several hundred to thousands of *A. fumigatus* conidia can be inhaled daily (1-3). Individuals with normal immune systems readily clear *A. fumigatus* conidia from their airways without problems. However, immunocompromised individuals are at significantly greater risk of developing invasive pulmonary aspergillosis (IPA)[4]. IPA is a life-threatening disease with over 300,000 cases globally and a high mortality rate of 30% - 50% (4). Patients at increased risk of developing IPA include those receiving chemotherapy treatments for cancer and patients receiving immunosuppressive regimens to prevent graft-versus-host disease (GVHD) following hematopoietic stem cell transplants or solid organ transplants (5-9). Furthermore, epidemiology studies predict a steady increase of IPA cases in the future due to the projected continual increase of immune compromised patients (10). Additionally, individuals with single nucleotide polymorphisms (SNPs) or primary immunodeficiencies in key antifungal signaling or effector pathways, such as CARD9 (11), pentraxin 3 (12, 13), or NADPH oxidase (14, 15) are highly susceptible to IPA. Thus, these clinical observations demonstrate that host innate immune responses are imperative in deactivating viable fungal conidia to avoid fungal establishment in the lung, and failure of such processes result in fungal germination and fungal growth and dissemination, ultimately culminating in IPA.

If the physical barriers of the lungs are bypassed by the inhaled *A. fumigatus* conidia, airway epithelial cells and lung-resident macrophages comprise the first line of defense against inhaled conidia. It is well established that lung resident macrophages, including alveolar macrophages (16, 17) and CCR2$^+$ monocytes (18, 19) are critical for initiating the early

inflammatory milieu necessary for the recruitment of innate immune cells to the respiratory tract to mediate host resistance against invasive disease. These innate immune effector cells include neutrophils (20, 21), inflammatory monocytes (18, 19), NKT cells (22), and plasmacytoid dendritic cells (23-25). The cytokine/chemokine signaling networks necessary to regulate the recruitment of inflammatory cells to the respiratory tract are rapidly emerging, but still much remains to be learnt.

Recently, it has become well appreciated that the host inflammatory response induced by *A. fumigatus* is tightly tuned to the virulence of the individual strain under study (26-30). Being able to recognize pathogenic traits of microbes is key to appropriately responding to infection (31), one such trait is being able to resist killing and remain viable (32). Two crucial inflammatory pathways in signaling bacterial vitality are inflammasome-dependent secretion of IL-1β and induction of the interferon response, which both can be driven by bacterial RNA (33, 34). We and others have previously demonstrated that inflammasomes are essential for IL-1β production after *A. fumigatus* challenge (35-38). Interestingly, Kanneganti and colleagues have shown to the NLRP3 and AIM2 inflammasomes could be activated by fungal RNA and DNA, respectively (35). Moreover, both type I and type III interferons have been shown to be essential for host resistance against pulmonary *A. fumigatus* challenge (39), but the fungal PAMP and host pattern-recognition pathways leading to type I and type III interferon expression following *A. fumigatus* challenge have not been well elucidated.

Type I and type III interferon responses are best understood in the context of virus infections. Following virus infections, Toll-like receptors (including TLR3, TLR7/8, TLR9), cGAS/STING, and RIG-I-like receptors (RIG-I and MDA5) are known to be critical in the induction of type I and type III interferons (review by (40-42)). However, during fungal

infections much less is currently known. Both TLR3 and TLR9 are known to be involved in the antifungal defenses against *A. fumigatus* (43-46), but only their role in the induction of type I interferons has been explored (45, 47). Following *Candida albicans* infection TLR7 is known to be involved in the induction of type I interferons (48, 49). Moreover, a Dectin-1/Syk/Irf5 signaling network has been shown to be critical in the induction of type I interferons following *C. albicans* challenge (50). Finally, Dectin-1 has been shown to be partially involved in the induction of type I and III interferons following respiratory *A. fumigatus* challenge (Dutta *et al,* in press). However, the role of cytosolic sensing pattern-recognition receptors systems in the antifungal interferon response has not been well studied.

Our data demonstrate that MDA5/MAVS-dependent signaling is necessary for host resistance against *A. fumigatus* infection through the accumulation and activation of antifungal effector functions of neutrophils in the fungal challenged airways. MAVS signaling drove expression of both type I and type III interferon, which ultimately drove the expression of the interferon inducible chemokines CXCL9 and CXCL10. Overall, our study reveals a critical role for MDA5 in host anti-fungal immunity, which supports a broader role of MDA5 in monitoring the health of the host cytosol.

MATERIALS AND METHODS

Mice. C57BL/6J (Jackson Laboratory, Stock #000664), *Mavs*$^{-/-}$ (Jackson Laboratory, Stock #008634), and *Ifih1*$^{-/-}$ (Jackson Laboratory, Stock # 015812) were all originally purchased from Jackson Laboratories and sequentially bred in-house at Geisel School of Medicine at Dartmouth. *Ifnar*$^{-/-}$, *Ifnlr*$^{-/-}$, and *Ifnar*$^{-/-}$ *Ifnlr1*$^{-/-}$ DKO were kindly provided by Dr. Sergei Kotenko and Dr. Joan Durbin (51) and bred in-house at Rutgers – New Jersey Medical School. Wild-type B6.129F2 (Jackson Laboratory, Stock #101045) mice, used as controls for the *Mavs*$^{-/-}$ mice, were purchased directly from Jackson Laboratories. All mice were 8-16 weeks of age at the time of challenge. All animal experiments were approved by either the Rutgers – New Jersey Medical School Institutional Animal Care and Use Committee or Dartmouth College Institutional Animal Care and Use Committee.

Preparation of Aspergillus fumigatus conidia. *A. fumigatus* CEA10 and Af293 strains were used for this study. Each strain was grown on glucose minimal media (GMM) agar plates for 3 days at 37°C. Conidia were harvested by adding 0.01% Tween 80 to plates and gently scraping conidia from the plates using a cell scraper. Conidia were then filtered through sterile Miracloth, were washed, and resuspended in phosphate buffered saline (PBS), and counted on a hemocytometer.

Preparation of swollen conidia and heat-killing. *A. fumigatus* CEA10 conidia were collected, as described above. To generate a homogenous population of swollen conidia, conidia were resuspended at 5×10^6 conidia per ml in RPMI containing 0.5 μg/ml of voriconazole (Sigma-Aldrich) and incubated at 37°C for 15 h, as previously done (52). Conidia were then washed twice and adjusted to 4×10^8 conidia per ml in PBS-Tween 80. To heat-kill the swollen conidia,

the conidial suspension was incubated in a 100°C water bath for 30 minutes. The conidial

concentration and efficiency of heat-killing was verified by plating on GMM agar.

Aspergillus fumigatus pulmonary challenge model. Mice were challenged with *A. fumigatus*

conidia by the intratracheal (i.t.) route. Mice were anesthetized by inhalation of isoflurane;

subsequently, mice were challenged i.t. with ~4×10^7 *A. fumigatus* conidia in a volume of 100 µl

PBS. At the indicated time after *A. fumigatus* challenge, mice were euthanized using a lethal

overdose of pentobarbital. Bronchoalveolar lavage fluid (BALF) was collected by washing the

lungs with 2 ml of PBS containing 0.05*M* EDTA. BALF was clarified by centrifugation and

stored at -20°C until analysis. After centrifugation, the cellular component of the BALF was

resuspended in 200 µl of PBS and total BAL cells were determined by hemocytometer count.

BALF cells were subsequently spun onto glass slides using a Cytospin4 cytocentrifuge (Thermo

Scientific) and stained with Diff-Quik (Siemens) or Hema 3™ Stat Pack (Fisher Scientific) stain

sets for differential counting. For histological analysis lungs were filled with and stored in 10%

buffered formalin phosphate for at least 24 hours. Lungs were then embedded in paraffin and

sectioned into 5-micron sections. Sections were stained with Grocott-Gomori methenamine

silver (GMS) using standard histological techniques to assess lung inflammatory infiltrates and

fungal germination, respectively. Representative pictures of lung sections were taken using an

Olympus BX50WI microscope with a QImaging Retiga 2000R camera.

Luminex assay for cytokine and chemokine secretion from experimental murine models of

invasive aspergillosis. BALF and lung homogenates from B6/129F2 mice and *Mavs$^{-/-}$* mice

challenged with *A. fumigatus* 12 or 48 h prior were analyzed for cytokines and chemokines using

Milliplex Mouse Cytokine & Chemokine 32-plex (Millipore). Plates were read using a BioPlex

200 (Bio-Rad) in the Immune Monitoring and Flow Cytometry Core Facility at Dartmouth

College.

Quantitative RT-PCR analysis. Total RNA from lungs was extracted with Trizol (Invitrogen).

One microgram of total RNA was reverse transcribed using High Capacity cDNA Reverse

Transcription Kit (Applied Biosystems). TaqMan Fast Universal Master Mix (2X) No AmpErase

UNG and TaqMan probes (Applied Biosystems, Catalog #4331182) for *Ifna2*

(Mm00833961_s1), *Ifnb1* (Mm00439552_s1), *Ifnl2/3* (Mm0404158_gH), *Ifng*

(Mm01168134_m1), *Cxcl9* (Mm00434946_m1), and *Cxcl10* (Mm00445235_m1) were used and

normalized to *Gapdh* (Mm99999915_g1). Gene expression was calculated using $\Delta\Delta$CT method

relative to naïve sample.

Western blot analysis. Total protein from whole lungs was extracted using RIPA buffer. Total

protein was quantified using a DC™ Protein Assay (Bio-Rad). For Western Blot analysis, 40μg

of total protein was loaded per well. Gels were run at 100V for ~1 hour at room temperature and

then transferred to PVDF membranes at 100V for ~1 hour at 4°C. After transfer, the membranes

were blocked with TBST + 5% milk for 1 hour at room temperature. Following blocking, blots

were rinsed twice with TBST buffer. Staining with primary antibodies was done in TBST with

5% BSA at 4°C for overnight on a roller. Primary antibodies used for these studies were rabbit

anit-STAT1 (Cell Signaling, #9172S, 1/1000), rabbit anti-pSTAT1 (Cell Signaling, #7649P,

1/1000 dilution), and rabbit anti-β-actin (Abcam, ab75186, 1/40,000). After primary antibody

staining blots were washed four times with TBST. Blots were then stained with a donkey anti-

rabbit IgG conjugated to HRP (Cell Signaling, #7074P2, 1/4000 dilution) in TBST + 5% milk for

1 hour at room temperature with gentle shaking. After secondary antibody staining blots were

washed four times with TBST. Blots were developed using ECL Clarity (Bio-Rad) for 5 minutes,

then analyzed using an Alpha Innotech FluorChem Q imager.

Generation of a Fluorescent Aspergillus fumigatus reporter (FLARE) strain in the CEA10

background. CEA10:H2A.X$^{A.\ nidulans}$::_ptrA_ was constructed in two steps. First, _gpdA_(p) was

amplified from DNA isolated from strain _A. nidulans_ A4 (Source: FGSC) using primers

RAC2888 and RAC2799. Histone variant H2A.X (AN3468) was amplified (without stop codon)

from A4 DNA using primers RAC4582 and RAC4583. mRFP fragment was amplified from

plasmid pXDRFP4 (Source: FGSC) using primers RAC2600 and RAC4575 and terminator for

A. nidulans trpC gene was amplified using primers RAC2536 and RAC2537. The four fragments

were then fused together with primers RAC1981 and RAC4134 using fusion PCR resulting in

H2A.X first round fragment as described earlier (53). All primers are listed in Supplemental

Table 1.

Secondly, we targeted integration of the H2A.X::rfp to the intergenic locus between

AFUB_023460 and AFUB_023470. For this, left homology arm was amplified from CEA10

genomic DNA using primers RAC3873 and RAC3874. Right homology arm was amplified with

primers RAC3875 and RAC3876. Dominant selection marker gene _ptrA_ conferring resistance to

pyrithiamine hydrobromide was amplified from plasmid pSD51.1 with primers RAC2055 and

RAC2056. The four fragments were then fused together with primers RAC3877 and RAC3878

using fusion PCR as described earlier (53). After the construct generation, polyethylene glycol

mediated transformation of protoplast was performed as described earlier (53). Transformants

were screened by PCR (data not shown) and confirmed with southern blot analysis as described

earlier (54). mRFP fluorescence was confirmed with FACS (Fluorescence activated cell sorting)

analysis.

Fluorescent Aspergillus fumigatus reporter (FLARE) assay. To measure both conidial uptake

and viability in distinct immune cell populations we used fluorescent *Aspergillus* reporter

(FLARE). The CEA10-based FLARE was labelled with AlexaFlour 633 membrane labeling, as

described elsewhere (55). Bronchoalveolar lavage (BAL) and lung cell suspensions were

prepared as described elsewhere (55) and stained with anti-Ly6G (1A8), anti-CD11b (M1/70),

anti-CD11c (HL3), anti-CD45 (30-F11), anti–CD206 (C068C2) and a fixable viability dye

(eBioscience; Catalog #65-0865-14). Neutrophils were identified as $CD45^+CD11b^+Ly6G^+$,

alveolar macrophages as $CD45^+CD11b^+Ly6G^-CD206^{dim}$, interstitial macrophages as

$CD45^+CD11b^+Ly6G^-CD206^-$, and monocytes as $CD45^+CD11b^{dim}Ly6G^-CD206^+$ cells, as

previously defined (56, 57). Flow cytometric data were collected on a CytoFLEX (Beckman

Coulter). All data was analyzed using FlowJo software.

Primary murine fibroblast cultures. Ears from 6-12 week-old C57BL/6J and *Ifih1*$^{-/-}$ mice were

collected, washed in 70% EtOH, then PBS, and finally minced using scissors. For each ear, 500

μl of 1000 U/ml collagenase (Gibco, REF 17101-015) in HBSS was used to digest them at 37°C

for 25 minutes. The tissues were then centrifuged at 500 x g for 3 minutes and then washed once

with HBSS. Ears were subsequently digested with 500 µl of 0.05% trypsin in HBSS for a further

20 minutes in 37°C. The tissues were then centrifuged again at 500 x g for 3 minutes, and the

trypsin was discarded and replaced by 0.5ml fibroblast media [10% FCS, 1% MEM Non-

essential amino acids, 1% pencillin-streptomycin in DMEM]. The tissues were then pushed

through a sterile 40 µm cell strainer via the plunger of a 3ml syringe to obtain single cell

suspensions. For every ear, 25ml of fibroblast media was used to initiate the murine fibroblast

culture in tissue culture flasks and a media change was conducted every two days. The

fibroblasts were usually confluent and harvested on day 6 by 0.25% trypsin digest for 5min in

37°C and then collected with fresh fibroblast media.

Isolation of total RNA from Aspergillus fumigatus. *A. fumigatus* conidia was harvested from

GMM plates and inoculated at 1 x 10^6 per ml in liquid GMM. Cultures were grown for 24h in

37°C with shaking at 250 rpm in an orbital shaker. Mycelia were then filtered and wrapped in

sterile Miracloth, dried by repeated paper towel absorption, and weighed. For every 50 mg of

mycelium, 1 ml of Trisure (Bioline) was added and mixed. The Trisure-mycelium mixture was

frozen by liquid nitrogen to release cellular content and then homogenized using a mortar and

pestle. For every 1 ml of Trisure used, 0.2 ml chloroform was added and vortexed, and the

solution was centrifuged for 15 min at 12000 x g at 4°C. The aqueous phase was then transferred

to a new tube and 0.5 ml of isopropanol was added, vortexed, and centrifuged for 10 min at

12000 x g at 4°C. The supernatant was then removed, and the pellet was washed by 1 ml 75%

ethanol via vortexing followed with centrifuging for 5 min at 7500 x g at 4°C. The 75% ethanol

was then removed to allow a complete air dry of the RNA, followed by dissolving in molecular

biology grade water and stored in -80°C.

Stimulation of murine fibroblasts with fungal RNA. Freshly harvested murine fibroblasts were

dosed at 5 x10^4/ml in fibroblast media and transferred to 24-well tissue culture treated plates

with 0.5 ml per well. The plate was incubated for one day to allow the attachment of murine

fibroblasts. Fibroblasts were then transfected via LyoVec™ (Invivogen)/RNA complexes. For

every 200 μl of LyoVec™, 2 μg of either poly-IC (Invivogen), 5'-ppp RNA (Invivogen), or *A.*

fumigatus RNA was diluted in 80 μl molecular biology grade water, mixed, and incubated in

room temperature for 30 minutes to assemble the liposome. The assembled liposome complexes

were then added to 5 ml of fibroblast medium and 0.5 ml of the above transfection media was

transferred to each well of 24-well plates after removal of the original medium. Twenty-four

hours after transfection the supernatant was collected for subsequent ELISA analysis.

ELISA analysis for cytokine and chemokine secretion. Cell culture supernatants from fibroblast

stimulated with *A. fumigatus* RNA were analyzed by ELISA for CXCL10 (R&D Systems, Cat.

DY466) and Interferon alpha (Invitrogen, Cat. BMS6027TWO). Plates were read using an Epoch

BioTek Gen5 microplate reader at 450nm, and the background was subtracted at 570nm.

Statistical analysis. Statistical significance for *in vitro* and *ex vivo* data was determined by a

Mann-Whitney U test, one-way ANOVA using a Tukey's or Dunn's post-test, or two-way

ANOVA with Tukey's post-test through the GraphPad Prism 7 software as outlined in the figure

legends. Mouse survival data were analyzed with the Mantel-Cox log rank test using GraphPad

Prism.

RESULTS

Induction of an optimal interferon response by *Aspergillus fumigatus* is dependent on viable conidia.

The polysaccharide-rich cell wall is a major driver of the inflammatory response against *A. fumigatus* and other fungi. Following *C. albicans* challenge the Dectin-1 (*Clec7a*) receptor is critical in the induction of type I interferons following (50). Dectin-1 is only partially responsible for induction of the type I and type III interferon response induced by *A. fumigatus* (Dutta *et al*, in press). To test whether the fungal cell wall was essential for driving the type I and type III interferon response, we stimulated C57BL/6J mice with 4×10^7 live or heat-killed swollen conidia of the CEA10 strain of *A. fumigatus*. Forty hours post-challenge with live swollen conidia there was an induction of IL-28 (IFN-λ), CXCL10, and TNFα secretion (Figure 1). In contrast, when challenged with heat-killed swollen conidia secretion of both IL-28 (IFN-λ) and CXCL10 was markedly reduced, while TNFα was still induced (Figure 1). These finding demonstrate that while cell carbohydrates in heat-killed swollen conidia still drove robust inflammatory cytokines (such as TNFα), as has previously been demonstrated (52, 58), there is an alternative pattern-recognition receptor pathway which is largely responsible for the interferon response induced by live *A. fumigatus* conidia.

dsRNA from *Aspergillus fumigatus* drives an MDA5-dependent interferon response.

Vitality sensing during bacterial infection through recognition of bacterial RNA has been shown to be critical for the induction of protective immune responses through the secretion of both IL-1β and type I interferons (33). Thus, we next wanted to test whether *A. fumigatus* RNA could drive interferon release. To test this hypothesis, primary murine fibroblasts from C57BL/6J

mice were stimulated with naked or liposome packaged total RNA from the Af293 strain of *A.*

fumigatus. As expected, untreated fibroblast or empty liposome treated did not drive significant

secretion of IFNα or CXCL10, while liposome packaged polyI:C drove significant secretion of

both (Figure 2A). Liposome-packaged Af293 RNA drove secretion of both IFNα and CXCL10,

while naked Af293 could not drive their production (Figure 2A). Taken together, these data

demonstrate that intracellular sensing of *A. fumigatus* RNA can drive an interferon-dependent

inflammatory response.

Intracellular sensing of RNA is mediated by the RIG-I-like receptor family, composed of

RIG-I and MDA5, for the induction of interferons. Both RIG-I and MDA5 recognize distinct

structures in foreign RNA (59, 60). To test whether ssRNA or dsRNA from *A. fumigatus* was

required to drive type I IFN secretion, we extended these *in vitro* fibroblast stimulation studies

by pre-treating the *A. fumigatus* RNA pool with either RNase S1 or RNase III, which will

selectively cleave ssRNA and dsRNA, respectively, prior to liposome packaging. We also treated

the fungal RNA pool with DNase I as a control. Similar to our previous result, total *A. fumigatus*

RNA packaged in liposomes was able to induce IFNα secretion from murine fibroblast (Figure

2B). Pre-treatment of the *A. fumigatus* RNA with either RNase S1 or DNase I had no impact on

the secretion of IFNα (Figure 2B). In contrast, pre-treatment of the *A. fumigatus* RNA with

RNase III completely ablated the secretion of IFNα from primary murine fibroblasts (Figure 2B).

These data suggest that recognition of double-stranded fungal RNA in the cytosol is necessary

for the interferon response induced by *A. fumigatus.*

To date the most well characterized intracellular receptor for dsRNA is MDA5 (59). To

test whether *A. fumigatus* RNA initiates an interferon response in a MDA5-dependent manner we

treated primary murine fibroblasts isolated from C57BL/6J or *Ifih1*[-/-] mice liposome packaged

fungal RNA. As positive controls, we utilized high molecular weight polyI:C, a known MDA5

ligand, and 5'ppp-RNA, a known RIG-I ligand. Eighteen hours are stimulation cell culture

supernatants were collected for ELISA analysis to assess IFNα secretion. Unstimulated or

LyoVec™ only stimulated fibroblast did not secrete any IFNα (Figure 2B). As positive controls,

5'ppp-RNA stimulated equivalent secretion of IFNα from both wild-type and *Ifih1*-deficient

fibroblasts; while high molecular weight polyI:C stimulated robust IFNα secretion from wild-

type fibroblast, which was significantly reduced in *Ifih1*-deficient fibroblasts (Figure 2B). The

induction of IFNα secretion by *A. fumigatus* RNA was partially dependent on MDA5, like what

we also observed with high molecular weight polyI:C (Figure 2B). Interestingly, RNase III

treatment ablated both the MDA5-dependent and -independent IFN-α response (Figure 2B),

which suggests there is another dsRNA receptor that contributions to the IFN-α response in

response to encapsulated fungal RNA.

MAVS-dependent signaling is required for interferon expression after *Aspergillus*

***fumigatus* challenge**

During viral infections RLR family members are critical in initiating both the type I and

type III interferon responses (61). Moreover, stimulation of *Ifih1*-deficient murine macrophages

with *C. albicans* resulted in a slight decrease in *Ifnb* mRNA expression, which did not reach

significance (62). However, the specific role of MDA5/MAVS in initiating the induction of a

broad range of interferons to other fungal pathogens has not been explored. To test the role of

MAVS signaling in the induction of interferons after respiratory challenge with *A. fumigatus*, we

challenged B6.129F2 mice and *Mavs*$^{-/-}$ mice with 4×10^7 conidia of CEA10. We collected lungs

from the *A. fumigatus* challenged mice at 3 and 48 h post-inoculation for quantitative RT-PCR

analysis to assess interferon expression patterns. Similar to previous observations with C57BL/6J

mice (39), wild-type B6.129F2 mice expressed high levels of type I interferons, *Ifna4* (Figure

3A) and *Ifnb* (Figure 3B), at 3 h post-inoculation that waned with time. Both type II interferon

(*Ifng*) (Figure 3D) and type III interferon (*Ifnl2/3*) (Figure 3C) were expressed at higher levels at

48 h post-inoculation in wild-type B6.129F2 mice. Interestingly, in the *Mavs*-deficient mice type

I interferon levels were only decreased by approximately 50% (Figure 3A-B), whereas the

expression of type II and type III interferons were nearly completely ablated in the absence of

MAVS (Figure 3C-D). Thus, MAVS-dependent signaling is essential for the late expression of

type III interferon and only partially responsible for early expression of type I interferons.

To confirm that the expression of interferons resulted in down-stream IFN signaling, we

next examined the phosphorylation of STAT1. STAT1 phosphorylation is essential for the

signaling of all interferon classes. Lungs from C57BL/6J mice challenged 6, 24, or 48 h prior

with either the CEA10 or Af293 strains were collected, and total proteins were extracted for

Western blot analysis. Both the CEA10 and Af293 strains induced a bi-phasic induction of

STAT1 phosphorylation (Figure 4A). Specifically, significant STAT1 phosphorylation was

observed at 6 and 48 h after challenge, but minimal phosphorylation was observed at 24 h

(Figure 4A). These data fit with our earlier observation that type I interferons are produced by 3

h post-challenge, while type II and type III interferons are highly expressed at later times (Figure

3). We next examined the phosphorylation of STAT1 in the lungs of B6.129F2 and *Mavs*$^{-/-}$ mice

challenged with the CEA10 stain of *A. fumigatus* at 6 and 48 h after challenge, times when we

saw high levels of STAT1 phosphorylation. Like C57BL/6J mice, wild-type B6.129F2 mice

induced robust phosphorylation of STAT1 at both 6 and 48 h (Figure 4B). Interestingly, at 6 h

after challenge with *A. fumigatus* the phosphorylation of STAT1 in the *Mavs*$^{-/-}$ mice was not

significantly different, although there was a slight but reproducible reduction (Figure 4B-C),

which fits with our earlier observation that type I interferon mRNA levels were only mildly

reduced at 3 h post-challenge with *A. fumigatus* (Figure 3A-B). In contrast, phosphorylation of

STAT1 in the *Mavs*[-/-] mice was significantly decreased at 48 h post-challenge with *A. fumigatus*

(Figure 4B-C), which corresponds with the significant decrease in type III interferon mRNA

levels we observed at this time (Figure 3C). Taken together, these results strongly support the

conclusion that MAVS signaling is critical for the late expression of type III interferons in

response to *A. fumigatus* instillation and initiation of the interferon response that is necessary for

host resistance against *A. fumigatus* infection.

Alteration of the airway inflammatory milieu after *Aspergillus fumigatus* challenge in the

absence of MAVS-dependent signaling

While our data demonstrates that all the classes of interferons are decreased to vary

degrees after pulmonary *A. fumigatus* challenged, we next wanted to more broadly understand

the cytokine and chemokine response in the airways of *Mavs*[-/-] mice. To achieve this, we

challenged B6.129F2 and *Mavs*[-/-] mice with $4x10^7$ conidia of *A. fumigatus*. At 12 and 48 h after

challenge with *A. fumigatus*, the BALF was analyzed using a 32-plex Milliplex cytokine assay.

Numerous cytokines and chemokines--TNFα, IL-17A, CCL2, and CCL4--were expressed in a

manner that was completely independent of MAVS signaling (Figure 5). Interestingly, IL-1α, IL-

1β, CXCL1, and CXCL2 secretion were significantly greater in the absence of MAVS signaling

(Figure 5). In contrast, IL-12p70, CXCL9, and CXCL10 secretion was significantly impaired in

the absence of MAVS signaling, particularly at 48 h post-challenge despite the large increase in

fungal burden (Figure 5). These data suggest that MAVS-dependent signaling is critical for

induction and/or maintenance of these cytokines.

***Cxcl9* and *Cxcl10* mRNA expression after pulmonary *Aspergillus fumigatus* challenge**

requires both type I and type III interferons

The marked decrease in CXCL9 and CXCL10 expression in the absence of MAVS

following *A. fumigatus* challenge was quite striking, because both CXCL9 and CXCL10 are

interferon-inducible chemokines (63). To test the role of type I and type III interferons in the

induction of CXCL9 and CXCL10 expression, we challenged C57BL/6, *Ifnar*$^{-/-}$, *Ifnlr*$^{-/-}$, and

Ifnar$^{-/-}$ *Ifnlr*$^{-/-}$ (DKO) mice with 4×10^7 conidia of *A. fumigatus*. Forty-eight hours later, lungs

were collected for quantitative RT-PCR analysis to assess *Cxcl9* and *Cxcl10* mRNA expression

patterns. Both *Ifnar-* and *Ifnlr*-deficient mice had a significant decrease in their expression of

both *Cxcl9* (Figure 6A) and *Cxcl10* (Figure 6B). In general, the defect in *Cxcl9* and *Cxcl10*

mRNA expression was greater in the absence of type III interferon signaling, particularly

regarding *Cxcl9* expression. Importantly, the *Ifnar*$^{-/-}$ *Ifnlr*$^{-/-}$ (DKO) mice had the most severe

defect in *Cxcl9* and *Cxcl10* expression (Figure 6). Overall, these data demonstrate that type I and

type III interferon signaling are essential for the expression of both *Cxcl9* and *Cxcl10*.

MDA5/MAVS-dependent signaling is required for host resistance against *Aspergillus*

***fumigatus* infection**

These previous data suggest that MDA5 can sense *A. fumigatus* RNA to drive an

interferon response, which could be critical in vitality sensing following an *in vivo* challenge

with *A. fumigatus*. Thus, we wanted to test the role of MDA5/MAVS signaling in host resistance

against respiratory challenge with the human fungal pathogen, *A. fumigatus*. To globally assess

the role of the RLR family in antifungal immunity, we initially challenged B6.129F2 mice and

Mavs[-/-] mice with 4×10^7 conidia of the CEA10 strain, since MAVS is the central signaling

adaptor for both RIG-I and MDA5-mediated responses (64-67). We monitored the survival of the

A. fumigatus challenged mice over the next two weeks. *Mavs*-deficient mice were more

susceptible to pulmonary challenge with *A. fumigatus* than wild-type, B6/129F2 mice (Figure

7A; Mantel-Cox log rank test, p = 0.0001). Additionally, susceptibility of the *Mavs*[-/-] mice to *A.*

fumigatus challenge is not dependent on the strain of *A. fumigatus* used because *Mavs*[-/-] mice

challenged with the Af293 strain were also highly susceptible to infection (Supplemental Figure

1; Mantel-Cox log rank test, p < 0.0001).

It is widely acknowledged that neutrophils and macrophages are critical antifungal

effector cells for clearing *A. fumigatus* from the lungs. Thus, we assessed inflammatory cell

accumulation in the airways via differential microscopic counting of cytospins stained with Diff-

Quik from the BALF. The increased susceptibility of *Mavs*-deficient mice to *A. fumigatus*

challenge was associated with decreased accumulation of macrophages and neutrophils in the

airways compared with B6.129F2 mice at 40 h post-challenge (Figure 7B). Next, we assessed

fungal growth and tissue invasion in the lung by histological analysis at 40 h after conidial

instillation. Strikingly, GMS staining of lung tissue from *Mavs*-deficient mice revealed the

presence of germinating *A. fumigatus* conidia at 40 h that was not observed to the same extent in

B6.129F2 mice (Figure 7C). When the presence of germinating *A. fumigatus* conidia was

quantified, wild-type B6.129F2 mice displayed low levels of fungal germination (17.6% ± 2.9)

compared with *Mavs*[-/-] mice in which most of the fungal material had germinated (73.8% ± 7.6)

by 40 h post-challenge (Figure 7C). B6.129F2 mice had very few areas of fungal material

throughout their lungs, but it was associated with robust cellular infiltration and tissue

destruction (Figure 7C).

To specifically address which RLR sensor is involved in regulating the innate anti-fungal

immune response against *A. fumigatus* we chose to assess the role of MDA5 because of its role

in the *in vitro* sensing of *A. fumigatus* RNA (Figure 2). Specifically, we challenged C57BL/6J

and *Ifih1*$^{-/-}$ mice with 4×10^{7} conidia of *A. fumigatus*. We monitored the survival of the *A.*

fumigatus challenged mice over the next ten days. Like the *Mavs*-deficient mice, *Ifih1*-deficient

mice were more susceptible to pulmonary challenge with *A. fumigatus* than wild-type C57BL/6J

mice (Figure 7D; Mantel-Cox log rank test, p = 0.0004). Analogous to what we observed in the

Mavs-deficient mice, the increased susceptibility of *Ifih1*-deficient mice to *A. fumigatus*

challenge was associated with decreased accumulation of neutrophils in the airways compared

with C57BL/6J mice at 40 h post-challenge (Figure 7E). GMS staining of lung tissue from *Ifih1*-

deficient mice revealed the presence of germinating *A. fumigatus* conidia at 42 h that was not

observed to the same extent in C57BL/6J mice (Figure 7F) similar to our observation in the

Mavs-deficient mice.

While immune competent animal models have provided important insights into the host

immune pathways necessary for antifungal control (11, 13, 55, 68-70), we wanted to assess the

importance of MDA5/MAVS signaling in a clinically relevant immune compromised model of

IPA. For this we utilized the triamcinolone model of IPA (71). Briefly, C57BL/6J and *Ifih1*$^{-/-}$

mice were immunosuppressed with triamcinolone and then subjected to challenge with 10^{6}

conidia of the Af293 strain of *A. fumigatus*. At 72 h post-inoculation lungs were collected for

histological analysis for fungal growth. Interestingly, *Ifih1*$^{-/-}$ mice had a significantly greater

number of fungal lesions per lung section than C57BL/6J mice (Supplemental Figure 2A).

Moreover, the fungal lesions appear to be larger and contain more fungi (Supplemental Figure

2A). Additionally, the BALF from *Ifih1*$^{-/-}$ challenged mice had higher levels of LDH activity

(Supplemental Figure 2B) and albumin (Supplemental Figure 2C), which are indicators of lung

damage and leakage, respectively, than C57BL/6J challenged mice. Taken together, these results

strongly support the conclusion that MDA5/MAVS signaling is critical for host resistance

against *A. fumigatus* in an immune competent murine model and critical for disease progression

in a corticosteroid immune compromised murine model.

Neutrophil antifungal effector functions are decreased in the absence of MDA5.

To determine whether MDA5/MAVS signaling regulates antifungal effector cell

functions, we used FLARE conidia challenge of C57BL/6 or *Ifih1*$^{-/-}$ mice to assess neutrophil-

mediated conidial uptake and killing. FLARE conidia encode a fungal viability indicator

(DsRed) and contain a tracer fluorophore (Alexa Fluor 633) (55). For this study we generated a

novel FLARE strain in the CEA10 strain background (Supplemental Figure 3). FLARE conidia

emit two fluorescence signals (DsRed and Alexa Fluor 633) when the conidia are alive, but only

emit a single fluorescence signal (Alexa Fluor 633) when the conidia are dead. This approach

allows us to determine the frequency of conidia-engaged immune cells that contain either live or

dead fungal cells in the BAL and lungs. The frequency of conidia-engaged neutrophils was

similar between the C57BL/6 and *Ifih1*$^{-/-}$ mice (Figure 8A). However, the frequency of conidia-

engaged neutrophils that contain live conidia was increased among *Ifhi1*$^{-/-}$ mice compared with

C57BL/6J mice, indicating a defect in conidial killing (Figure 8B). Specifically, the frequency of

neutrophils that contain live conidia was 1.9-fold and 2.1-fold higher for BAL and lung fluid of

Ifih1$^{-/-}$ mice compared with their C57BL/6J counterparts (Figure 8B). Thus, these data indicate

that neutrophil-mediated antifungal killing of *A. fumigatus* conidia requires MDA5/MAVS

activation and signaling.

DISCUSSION

It is now well appreciated that the host inflammatory response induced by *A. fumigatus* is tightly tuned to the virulence of the individual strain under study (26-30). In filamentous fungi one key determinant for sensing the threat posed by the fungi is the changes in that occur in the fungal cell wall composition. Specifically, detection of β-1,3-glucan linked polysaccharides by Dectin-1 (*Clec7a*) occurs only upon conidial swelling and germling formation, which are the earliest steps of fungal growth (52, 58). Interestingly, our data demonstrate that while heat-killed swollen conidia of *A. fumigatus* can induce the secretion of pro-inflammatory cytokines, such as TNFα, they do not drive the secretion of IFNα and CXCL10 (Figure 1). These data suggest that fungal factors other than cell wall polysaccharides are responsible for driving the interferon response following respiratory *A. fumigatus* challenge. Consistent with this observation, individual polysaccharides commonly found in the *A. fumigatus* cell wall are not sufficient to induce an interferon response (72).

During bacterial infections one key virulence trait sensed by the host immune system to tune the inflammatory response is the ability of the bacteria to resist killing and remaining viable. Two crucial inflammatory pathways in signaling bacterial vitality are inflammasome secretion of IL-1β and interferon response through TRIF and cGAS/STING signaling (33, 34, 73). Following respiratory challenge with *A. fumigatus* both IL-1β and type I/III interferons have been shown to be critical for antifungal immunity (35, 39). While we and others have previously demonstrated that inflammasomes are essential for the IL-1β response after respiratory challenge with *A. fumigatus* (35-38), the innate pattern-recognition receptor(s) leading to type I/III interferon expression have not been elucidated.

In bacteria one critical PAMP associated with vitality (vita-PAMP) is RNA. Using an *E.*

coli infection model, bacterial RNA was shown to be sufficient to drive both inflammasome-

dependent IL-1β secretion and TRIF-dependent IFNβ secretion (33). Heat-killing of *E. coli*

drove the loss of RNA and its subsequent inflammatory response (33). Several innate RNA

sensors have been identified in mammalian hosts, including TLR3 (74), TLR7 (75), MDA5 (76),

and RIG-I (77). TLR3 and TLR7 signal through TRIF (78), while MDA5 and RIG-I signal

through MAVS (64-67) to drive the interferon response. Our data using RNase III treatment of

total RNA isolated from *A. fumigatus* demonstrates that dsRNA from *A. fumigatus* is sufficient to

drive an interferon response (Figure 2B). TLR3 and MDA5 are both receptors for dsRNA (59,

74, 77). Our data demonstrate that mice lacking *Mavs* displayed a markedly reduced type III

interferon (IL-28/IFNλ), CXCL9, and CXCL10 secretion, but only a moderate defect in type I

interferons (Figure 4). In contrast, the expression of *Ifna1* and *Ifnb1* are regulated by both TRIF

(46) and TLR3 (45). Interestingly, the role of TRIF appears to be restricted to the non-

hematopoietic compartment (46) likely epithelial cells (79). Taken together these data suggest

the type I interferon response following *A. fumigatus* challenge is primarily driven by

TLR3/TRIF-dependent signaling, while the type III interferon response is primarily drive by

MDA5/MAVS-dependent signaling.

The purpose of vitality sensing is to drive robust inflammation that is in line with the

threat posed by the invading pathogen. Neutrophils are well known to be critical antifungal

effector cells needed for the prevention of fungal germination and clearance of *A. fumigatus* from

the lungs (20). Our results demonstrate that MDA5/MAVS-dependent signaling is necessary for

both the optimal accumulation (Figure 7) and activation of neutrophils in the *Aspergillus*

challenged airways (Figure 8). Following pulmonary challenge with *A. fumigatus* it has been

established that both type I and type III interferons are essential for host resistance against

through enhancing neutrophil ROS production and antifungal effector functions (39). Our

FLARE data support as neutrophils in both the lungs and airways of *Ifih1*-deficient mice are

significantly impaired in their antifungal killing capacity at 48 h after challenge (Figure 8),

which corresponds with a time when type III interferons are nearly absent mice lacking

MDA5/MAVS signaling. Decreased accumulation of neutrophils in the *Mavs*- and *Ifih1*-deficient

mice did not correlate with classical neutrophil chemoattractants, such as CXCR2 ligands, that

are required to maintain host resistance against *A. fumigatus* (36, 69, 80). Rather, in the absence

of *Mavs* and *Ifih1* there was a marked decrease in CXCL9 and CXCL10, which are both ligands

for CXCR3. Recently, CXCR3 has been suggested to be a potential chemoattractant for

neutrophils in certain systems (81, 82). However, *Cxcr3*-deficient mice appear to recruit normal

numbers of neutrophils following *A. fumigatus* challenge (25). Thus, why *Mavs*- and *Ifih1*-

deficient mice accumulate few neutrophils in their lungs following challenge warrants further

research.

While the role of RLRs in host immunity was originally associated with antiviral innate

immune responses (reviewed by (42)), recent studies have begun to expand the role of RLRs in

immunity to other pathogens, including *C. albicans, Listeria monocytogenes*, *Mycobacterium*

tuberculosis, and *Plasmodium falciparum* (40, 62, 83-87). Our study adds *A. fumigatus* to the list

of non-viral pathogens which can activate the RLR family. Thus, cytosolic sensing of pathogens

by RLRs, both RIG-I and MDA5, may have a broader role in initiating innate immune responses

in several other clinically relevant non-viral pathogens. Additionally, MDA5/MAVS signaling

appears to be crucial in sensing vita-PAMPs for the enhancing the inflammatory response to

invading pathogens.

AUTHOR CONTRIBUTIONS

Conceived and designed the experiments: XW, AKC, AR, RC, JJO. Performed the experiments: XW, AKC, VE, KL, WC, SD. Analyzed the data: XW, AKC, VE, KL, WC, AR, JJO. Wrote the paper: XW, JJO.

XW and AKC are co-first authors: AKC initiated these studies before graduating, while XW picked up and completed these studies and helped write the manuscript, which warranted listing XW first.

ACKNOWLEDGEMENTS

Thank you to Dr. Brent Berwin (Geisel School of Medicine at Dartmouth), Dr. David Leib (Geisel School of Medicine at Dartmouth), and Dr. Tobias Hohl (Memorial Sloan Kettering Cancer Center) for helpful discussion on this project and manuscript. Thank you to Stacy Ceron for assistance in generation of the primary murine fibroblast cultures.

REFERENCES

1. Chazalet, V., J.P. Debeaupuis, J. Sarfati, J. Lortholary, P. Ribaud, P. Shah, M. Cornet, H. Vu Thien, E. Gluckman, G. Brucker, and J.P. Latge. 1998. Molecular typing of environmental and patient isolates of Aspergillus fumigatus from various hospital settings. *J Clin Microbiol* 36: 1494-1500.

2. Goodley, J.M., Y.M. Clayton, and R.J. Hay. 1994. Environmental sampling for aspergilli during building construction on a hospital site. *J Hosp Infect* 26: 27-35.

3. Hospenthal, D.R., K.J. Kwon-Chung, and J.E. Bennett. 1998. Concentrations of airborne Aspergillus compared to the incidence of invasive aspergillosis: lack of correlation. *Med Mycol* 36: 165-168.

4. Brown, G.D., D.W. Denning, N.A. Gow, S.M. Levitz, M.G. Netea, and T.C. White. 2012. Hidden killers: human fungal infections. *Sci Transl Med* 4: 165rv113.

5. Baddley, J.W., D.R. Andes, K.A. Marr, D.P. Kontoyiannis, B.D. Alexander, C.A. Kauffman, R.A. Oster, E.J. Anaissie, T.J. Walsh, M.G. Schuster, J.R. Wingard, T.F. Patterson, J.I. Ito, O.D. Williams, T. Chiller, and P.G. Pappas. 2010. Factors associated with mortality in transplant patients with invasive aspergillosis. *Clin Infect Dis* 50: 1559-1567.

6. Steinbach, W.J., K.A. Marr, E.J. Anaissie, N. Azie, S.P. Quan, H.U. Meier-Kriesche, S. Apewokin, and D.L. Horn. 2012. Clinical epidemiology of 960 patients with invasive aspergillosis from the PATH Alliance registry. *J Infect* 65: 453-464.

7. Garcia-Vidal, C., A. Upton, K.A. Kirby, and K.A. Marr. 2008. Epidemiology of invasive mold infections in allogeneic stem cell transplant recipients: biological risk factors for infection according to time after transplantation. *Clin Infect Dis* 47: 1041-1050.

8. Upton, A., K.A. Kirby, P. Carpenter, M. Boeckh, and K.A. Marr. 2007. Invasive aspergillosis following hematopoietic cell transplantation: outcomes and prognostic factors associated with mortality. *Clin Infect Dis* 44: 531-540.

9. Thompson, G.R., 3rd and T.F. Patterson. 2008. Pulmonary aspergillosis. *Semin Respir Crit Care Med* 29: 103-110.

10. Bongomin, F., S. Gago, R.O. Oladele, and D.W. Denning. 2017. Global and Multi-National Prevalence of Fungal Diseases-Estimate Precision. *J Fungi (Basel)* 3: 67.

11. Rieber, N., R.P. Gazendam, A.F. Freeman, A.P. Hsu, A.L. Collar, J.A. Sugui, R.A.
Drummond, C. Rongkavilit, K. Hoffman, C. Henderson, L. Clark, M. Mezger, M.
Swamydas, M. Engeholm, R. Schule, B. Neumayer, F. Ebel, C.M. Mikelis, S. Pittaluga,
V.K. Prasad, A. Singh, J.D. Milner, K.W. Williams, J.K. Lim, K.J. Kwon-Chung, S.M.
Holland, D. Hartl, T.W. Kuijpers, and M.S. Lionakis. 2016. Extrapulmonary Aspergillus
infection in patients with CARD9 deficiency. *JCI Insight* 1: e89890.

12. Fisher, C.E., T.M. Hohl, W. Fan, B.E. Storer, D.M. Levine, L.P. Zhao, P.J. Martin, E.H.
Warren, M. Boeckh, and J.A. Hansen. 2017. Validation of single nucleotide
polymorphisms in invasive aspergillosis following hematopoietic cell transplantation.
Blood 129: 2693-2701.

13. Cunha, C., F. Aversa, J.F. Lacerda, A. Busca, O. Kurzai, M. Grube, J. Loffler, J.A.
Maertens, A.S. Bell, A. Inforzato, E. Barbati, B. Almeida, P. Santos e Sousa, A. Barbui,
L. Potenza, M. Caira, F. Rodrigues, G. Salvatori, L. Pagano, M. Luppi, A. Mantovani, A.
Velardi, L. Romani, and A. Carvalho. 2014. Genetic PTX3 deficiency and aspergillosis in
stem-cell transplantation. *N Engl J Med* 370: 421-432.

14. Morgenstern, D.E., M.A. Gifford, L.L. Li, C.M. Doerschuk, and M.C. Dinauer. 1997.
Absence of respiratory burst in X-linked chronic granulomatous disease mice leads to
abnormalities in both host defense and inflammatory response to Aspergillus fumigatus. *J*
Exp Med 185: 207-218.

15. Pollock, J.D., D.A. Williams, M.A. Gifford, L.L. Li, X. Du, J. Fisherman, S.H. Orkin,
C.M. Doerschuk, and M.C. Dinauer. 1995. Mouse model of X-linked chronic
granulomatous disease, an inherited defect in phagocyte superoxide production. *Nat*
Genet 9: 202-209.

16. Grimm, M.J., R.R. Vethanayagam, N.G. Almyroudis, C.G. Dennis, A.N. Khan, A.C.
D'Auria, K.L. Singel, B.A. Davidson, P.R. Knight, T.S. Blackwell, T.M. Hohl, M.K.
Mansour, J.M. Vyas, M. Rohm, C.F. Urban, T. Kelkka, R. Holmdahl, and B.H. Segal.
2013. Monocyte- and macrophage-targeted NADPH oxidase mediates antifungal host
defense and regulation of acute inflammation in mice. *J Immunol* 190: 4175-4184.

17. Bhatia, S., M. Fei, M. Yarlagadda, Z. Qi, S. Akira, S. Saijo, Y. Iwakura, N. van Rooijen,
G.A. Gibson, C.M. St Croix, A. Ray, and P. Ray. 2011. Rapid host defense against
Aspergillus fumigatus involves alveolar macrophages with a predominance of
alternatively activated phenotype. *PLoS One* 6: e15943.

18.   Espinosa, V., A. Jhingran, O. Dutta, S. Kasahara, R. Donnelly, P. Du, J. Rosenfeld, I.
623   Leiner, C.C. Chen, Y. Ron, T.M. Hohl, and A. Rivera. 2014. Inflammatory monocytes
624   orchestrate innate antifungal immunity in the lung. *PLoS Pathog* 10: e1003940.

625   19.   Hohl, T.M., A. Rivera, L. Lipuma, A. Gallegos, C. Shi, M. Mack, and E.G. Pamer. 2009.
626   Inflammatory monocytes facilitate adaptive CD4 T cell responses during respiratory
627   fungal infection. *Cell Host Microbe* 6: 470-481.

628   20.   Mircescu, M.M., L. Lipuma, N. van Rooijen, E.G. Pamer, and T.M. Hohl. 2009. Essential
629   role for neutrophils but not alveolar macrophages at early time points following
630   Aspergillus fumigatus infection. *J Infect Dis* 200: 647-656.

631   21.   Bonnett, C.R., E.J. Cornish, A.G. Harmsen, and J.B. Burritt. 2006. Early neutrophil
632   recruitment and aggregation in the murine lung inhibit germination of Aspergillus
633   fumigatus Conidia. *Infect Immun* 74: 6528-6539.

634   22.   Cohen, N.R., R.V. Tatituri, A. Rivera, G.F. Watts, E.Y. Kim, A. Chiba, B.B. Fuchs, E.
635   Mylonakis, G.S. Besra, S.M. Levitz, M. Brigl, and M.B. Brenner. 2011. Innate
636   recognition of cell wall beta-glucans drives invariant natural killer T cell responses
637   against fungi. *Cell Host Microbe* 10: 437-450.

638   23.   Loures, F.V., M. Rohm, C.K. Lee, E. Santos, J.P. Wang, C.A. Specht, V.L. Calich, C.F.
639   Urban, and S.M. Levitz. 2015. Recognition of Aspergillus fumigatus hyphae by human
640   plasmacytoid dendritic cells is mediated by dectin-2 and results in formation of
641   extracellular traps. *PLoS Pathog* 11: e1004643.

642   24.   Ramirez-Ortiz, Z.G., C.K. Lee, J.P. Wang, L. Boon, C.A. Specht, and S.M. Levitz. 2011.
643   A nonredundant role for plasmacytoid dendritic cells in host defense against the human
644   fungal pathogen Aspergillus fumigatus. *Cell Host Microbe* 9: 415-424.

645   25.   Guo, Y., S. Kasahara, A. Jhingran, N.L. Tosini, B. Zhai, M.A. Aufiero, K.A.M. Mills, M.
646   Gjonbalaj, V. Espinosa, A. Rivera, A.D. Luster, and T.M. Hohl. 2020. During
647   Aspergillus Infection, Monocyte-Derived DCs, Neutrophils, and Plasmacytoid DCs
648   Enhance Innate Immune Defense through CXCR3-Dependent Crosstalk. *Cell Host*
649   *Microbe* doi: 10.1016/j.chom.2020.1005.1002.

650   26.   Caffrey-Carr, A.K., C.H. Kowalski, S.R. Beattie, N.A. Blaseg, C.R. Upshaw, A.
651   Thammahong, H.E. Lust, Y.W. Tang, T.M. Hohl, R.A. Cramer, and J.J. Obar. 2017. IL-
652   1alpha is Critical for Resistance Against Highly Virulent Aspergillus fumigatus Isolates.
653   *Infect Immun* 85: e00661-00617.

654   27.   Caffrey-Carr, A.K., K.M. Hilmer, C.H. Kowalski, K.M. Shepardson, R.M. Temple, R.A.
655   Cramer, and J.J. Obar. 2017. Host-Derived Leukotriene B4 Is Critical for Resistance
656   against Invasive Pulmonary Aspergillosis. *Front Immunol* 8: 1984.

657   28.   Caffrey, A.K. and J.J. Obar. 2016. Alarmin(g) the innate immune system to invasive
658   fungal infections. *Curr Opin Microbiol* 32: 135-143.

659   29.   Rizzetto, L., G. Giovannini, M. Bromley, P. Bowyer, L. Romani, and D. Cavalieri. 2013.
660   Strain dependent variation of immune responses to A. fumigatus: definition of pathogenic
661   species. *PLoS One* 8: e56651.

662   30.   Rosowski, E.E., N. Raffa, B.P. Knox, N. Golenberg, N.P. Keller, and A. Huttenlocher.
663   2018. Macrophages inhibit Aspergillus fumigatus germination and neutrophil-mediated
664   fungal killing. *PLoS Pathog* 14: e1007229.

665   31.   Vance, R.E., R.R. Isberg, and D.A. Portnoy. 2009. Patterns of pathogenesis:
666   discrimination of pathogenic and nonpathogenic microbes by the innate immune system.
667   *Cell Host Microbe* 6: 10-21.

668   32.   Blander, J.M. and L.E. Sander. 2012. Beyond pattern recognition: five immune
669   checkpoints for scaling the microbial threat. *Nat Rev Immunol* 12: 215-225.

670   33.   Sander, L.E., M.J. Davis, M.V. Boekschoten, D. Amsen, C.C. Dascher, B. Ryffel, J.A.
671   Swanson, M. Muller, and J.M. Blander. 2011. Detection of prokaryotic mRNA signifies
672   microbial viability and promotes immunity. *Nature* 474: 385-389.

673   34.   Barbet, G., L.E. Sander, M. Geswell, I. Leonardi, A. Cerutti, I. Iliev, and J.M. Blander.
674   2018. Sensing Microbial Viability through Bacterial RNA Augments T Follicular Helper
675   Cell and Antibody Responses. *Immunity* 48: 584-598.

676   35.   Karki, R., S.M. Man, R.K. Malireddi, P. Gurung, P. Vogel, M. Lamkanfi, and T.D.
677   Kanneganti. 2015. Concerted activation of the AIM2 and NLRP3 inflammasomes
678   orchestrates host protection against Aspergillus infection. *Cell Host Microbe* 17: 357-
679   368.

680   36.   Caffrey, A.K., M.M. Lehmann, J.M. Zickovich, V. Espinosa, K.M. Shepardson, C.P.
681   Watschke, K.M. Hilmer, A. Thammahong, B.M. Barker, A. Rivera, R.A. Cramer, and J.J.
682   Obar. 2015. IL-1alpha signaling is critical for leukocyte recruitment after pulmonary
683   Aspergillus fumigatus challenge. *PLoS Pathog* 11: e1004625.

37. Said-Sadier, N., E. Padilla, G. Langsley, and D.M. Ojcius. 2010. Aspergillus fumigatus
stimulates the NLRP3 inflammasome through a pathway requiring ROS production and
the Syk tyrosine kinase. *PLoS One* 5: e10008.

38. Moretti, S., S. Bozza, V. Oikonomou, G. Renga, A. Casagrande, R.G. Iannitti, M.
Puccetti, C. Garlanda, S. Kim, S. Li, F.L. van de Veerdonk, C.A. Dinarello, and L.
Romani. 2014. IL-37 inhibits inflammasome activation and disease severity in murine
aspergillosis. *PLoS Pathog* 10: e1004462.

39. Espinosa, V., O. Dutta, C. McElrath, P. Du, Y.J. Chang, B. Cicciarelli, A. Pitler, I.
Whitehead, J.J. Obar, J.E. Durbin, S.V. Kotenko, and A. Rivera. 2017. Type III interferon
is a critical regulator of innate antifungal immunity *Science Immunology* 2: eaan5357.

40. Wu, J., L. Tian, X. Yu, S. Pattaradilokrat, J. Li, M. Wang, W. Yu, Y. Qi, A.E. Zeituni,
S.C. Nair, S.P. Crampton, M.S. Orandle, S.M. Bolland, C.F. Qi, C.A. Long, T.G. Myers,
J.E. Coligan, R. Wang, and X.Z. Su. 2014. Strain-specific innate immune signaling
pathways determine malaria parasitemia dynamics and host mortality. *Proc Natl Acad Sci*
U S A 111: E511-520.

41. Lazear, H.M., J.W. Schoggins, and M.S. Diamond. 2019. Shared and Distinct Functions
of Type I and Type III Interferons. *Immunity* 50: 907-923.

42. Dixit, E. and J.C. Kagan. 2013. Intracellular pathogen detection by RIG-I-like receptors.
Adv Immunol 117: 99-125.

43. Herbst, S., A. Shah, M. Mazon Moya, V. Marzola, B. Jensen, A. Reed, M.A. Birrell, S.
Saijo, S. Mostowy, S. Shaunak, and D. Armstrong-James. 2015. Phagocytosis-dependent
activation of a TLR9-BTK-calcineurin-NFAT pathway co-ordinates innate immunity to
Aspergillus fumigatus. *EMBO Mol Med* 7: 240-258.

44. Ramaprakash, H., T. Ito, T.J. Standiford, S.L. Kunkel, and C.M. Hogaboam. 2009. Toll-
like receptor 9 modulates immune responses to Aspergillus fumigatus conidia in
immunodeficient and allergic mice. *Infect Immun* 77: 108-119.

45. Carvalho, A., A. De Luca, S. Bozza, C. Cunha, C. D'Angelo, S. Moretti, K. Perruccio,
R.G. Iannitti, F. Fallarino, A. Pierini, J.P. Latge, A. Velardi, F. Aversa, and L. Romani.
2012. TLR3 essentially promotes protective class I-restricted memory CD8(+) T-cell
responses to Aspergillus fumigatus in hematopoietic transplanted patients. *Blood* 119:
967-977.

46. de Luca, A., S. Bozza, T. Zelante, S. Zagarella, C. D'Angelo, K. Perruccio, C. Vacca, A.
Carvalho, C. Cunha, F. Aversa, and L. Romani. 2010. Non-hematopoietic cells contribute
to protective tolerance to Aspergillus fumigatus via a TRIF pathway converging on IDO.
Cell Mol Immunol 7: 459-470.

47. Khan, N.S., P.V. Kasperkovitz, A.K. Timmons, M.K. Mansour, J.M. Tam, M.W. Seward,
J.L. Reedy, S. Puranam, M. Feliu, and J.M. Vyas. 2016. Dectin-1 Controls TLR9
Trafficking to Phagosomes Containing beta-1,3 Glucan. *J Immunol* 196: 2249-2261.

48. Biondo, C., A. Malara, A. Costa, G. Signorino, F. Cardile, A. Midiri, R. Galbo, S.
Papasergi, M. Domina, M. Pugliese, G. Teti, G. Mancuso, and C. Beninati. 2012.
Recognition of fungal RNA by TLR7 has a nonredundant role in host defense against
experimental candidiasis. *Eur J Immunol* 42: 2632-2643.

49. Biondo, C., G. Signorino, A. Costa, A. Midiri, E. Gerace, R. Galbo, A. Bellantoni, A.
Malara, C. Beninati, G. Teti, and G. Mancuso. 2011. Recognition of yeast nucleic acids
triggers a host-protective type I interferon response. *Eur J Immunol* 41: 1969-1979.

50. del Fresno, C., D. Soulat, S. Roth, K. Blazek, I. Udalova, D. Sancho, J. Ruland, and C.
Ardavin. 2013. Interferon-beta production via Dectin-1-Syk-IRF5 signaling in dendritic
cells is crucial for immunity to C. albicans. *Immunity* 38: 1176-1186.

51. Lin, J.D., N. Feng, A. Sen, M. Balan, H.C. Tseng, C. McElrath, S.V. Smirnov, J. Peng,
L.L. Yasukawa, R.K. Durbin, J.E. Durbin, H.B. Greenberg, and S.V. Kotenko. 2016.
Distinct Roles of Type I and Type III Interferons in Intestinal Immunity to Homologous
and Heterologous Rotavirus Infections. *PLoS Pathog* 12: e1005600.

52. Hohl, T.M., H.L. Van Epps, A. Rivera, L.A. Morgan, P.L. Chen, M. Feldmesser, and
E.G. Pamer. 2005. Aspergillus fumigatus triggers inflammatory responses by stage-
specific beta-glucan display. *PLoS Pathog* 1: e30.

53. Szewczyk, E., T. Nayak, C.E. Oakley, H. Edgerton, Y. Xiong, N. Taheri-Talesh, S.A.
Osmani, and B.R. Oakley. 2006. Fusion PCR and gene targeting in Aspergillus nidulans.
Nat Protoc 1: 3111-3120.

54. Willger, S.D., S. Puttikamonkul, K.H. Kim, J.B. Burritt, N. Grahl, L.J. Metzler, R.
Barbuch, M. Bard, C.B. Lawrence, and R.A. Cramer, Jr. 2008. A sterol-regulatory
element binding protein is required for cell polarity, hypoxia adaptation, azole drug
resistance, and virulence in Aspergillus fumigatus. *PLoS Pathog* 4: e1000200.

55.   Jhingran, A., K.B. Mar, D.K. Kumasaka, S.E. Knoblaugh, L.Y. Ngo, B.H. Segal, Y.
Iwakura, C.A. Lowell, J.A. Hamerman, X. Lin, and T.M. Hohl. 2012. Tracing conidial
fate and measuring host cell antifungal activity using a reporter of microbial viability in
the lung. *Cell Rep* 2: 1762-1773.

56.   Schyns, J., F. Bureau, and T. Marichal. 2018. Lung Interstitial Macrophages: Past,
Present, and Future. *J Immunol Res* 2018: 5160794.

57.   Misharin, A.V., L. Morales-Nebreda, G.M. Mutlu, G.R. Budinger, and H. Perlman. 2013.
Flow cytometric analysis of macrophages and dendritic cell subsets in the mouse lung.
Am J Respir Cell Mol Biol 49: 503-510.

58.   Steele, C., R.R. Rapaka, A. Metz, S.M. Pop, D.L. Williams, S. Gordon, J.K. Kolls, and
G.D. Brown. 2005. The beta-glucan receptor dectin-1 recognizes specific morphologies
of Aspergillus fumigatus. *PLoS Pathog* 1: e42.

59.   Pichlmair, A., O. Schulz, C.P. Tan, J. Rehwinkel, H. Kato, O. Takeuchi, S. Akira, M.
Way, G. Schiavo, and C. Reis e Sousa. 2009. Activation of MDA5 requires higher-order
RNA structures generated during virus infection. *J Virol* 83: 10761-10769.

60.   Pichlmair, A., O. Schulz, C.P. Tan, T.I. Naslund, P. Liljestrom, F. Weber, and C. Reis e
Sousa. 2006. RIG-I-mediated antiviral responses to single-stranded RNA bearing 5'-
phosphates. *Science* 314: 997-1001.

61.   Odendall, C., E. Dixit, F. Stavru, H. Bierne, K.M. Franz, A.F. Durbin, S. Boulant, L.
Gehrke, P. Cossart, and J.C. Kagan. 2014. Diverse intracellular pathogens activate type
III interferon expression from peroxisomes. *Nat Immunol* 15: 717-726.

62.   Jaeger, M., R. van der Lee, S.C. Cheng, M.D. Johnson, V. Kumar, A. Ng, T.S. Plantinga,
S.P. Smeekens, M. Oosting, X. Wang, W. Barchet, K. Fitzgerald, L.A. Joosten, J.R.
Perfect, C. Wijmenga, F.L. van de Veerdonk, M.A. Huynen, R.J. Xavier, B.J. Kullberg,
and M.G. Netea. 2015. The RIG-I-like helicase receptor MDA5 (IFIH1) is involved in
the host defense against Candida infections. *Eur J Clin Microbiol Infect Dis* 34: 963-974.

63.   Groom, J.R. and A.D. Luster. 2011. CXCR3 ligands: redundant, collaborative and
antagonistic functions. *Immunol Cell Biol* 89: 207-215.

64.   Seth, R.B., L. Sun, C.K. Ea, and Z.J. Chen. 2005. Identification and characterization of
MAVS, a mitochondrial antiviral signaling protein that activates NF-kappaB and IRF 3.
Cell 122: 669-682.

65.   Kawai, T., K. Takahashi, S. Sato, C. Coban, H. Kumar, H. Kato, K.J. Ishii, O. Takeuchi,
778   and S. Akira. 2005. IPS-1, an adaptor triggering RIG-I- and Mda5-mediated type I
779   interferon induction. *Nat Immunol* 6: 981-988.

780   66.   Xu, L.G., Y.Y. Wang, K.J. Han, L.Y. Li, Z. Zhai, and H.B. Shu. 2005. VISA is an
781   adapter protein required for virus-triggered IFN-beta signaling. *Mol Cell* 19: 727-740.

782   67.   Meylan, E., J. Curran, K. Hofmann, D. Moradpour, M. Binder, R. Bartenschlager, and J.
783   Tschopp. 2005. Cardif is an adaptor protein in the RIG-I antiviral pathway and is targeted
784   by hepatitis C virus. *Nature* 437: 1167-1172.

785   68.   Garlanda, C., E. Hirsch, S. Bozza, A. Salustri, M. De Acetis, R. Nota, A. Maccagno, F.
786   Riva, B. Bottazzi, G. Peri, A. Doni, L. Vago, M. Botto, R. De Santis, P. Carminati, G.
787   Siracusa, F. Altruda, A. Vecchi, L. Romani, and A. Mantovani. 2002. Non-redundant role
788   of the long pentraxin PTX3 in anti-fungal innate immune response. *Nature* 420: 182-186.

789   69.   Jhingran, A., S. Kasahara, K.M. Shepardson, B.A. Junecko, L.J. Heung, D.K. Kumasaka,
790   S.E. Knoblaugh, X. Lin, B.I. Kazmierczak, T.A. Reinhart, R.A. Cramer, and T.M. Hohl.
791   2015. Compartment-specific and sequential role of MyD88 and CARD9 in chemokine
792   induction and innate defense during respiratory fungal infection. *PLoS Pathog* 11:
793   e1004589.

794   70.   Drummond, R.A., A.L. Collar, M. Swamydas, C.A. Rodriguez, J.K. Lim, L.M. Mendez,
795   D.L. Fink, A.P. Hsu, B. Zhai, H. Karauzum, C.M. Mikelis, S.R. Rose, E.M. Ferre, L.
796   Yockey, K. Lemberg, H.S. Kuehn, S.D. Rosenzweig, X. Lin, P. Chittiboina, S.K. Datta,
797   T.H. Belhorn, E.T. Weimer, M.L. Hernandez, T.M. Hohl, D.B. Kuhns, and M.S.
798   Lionakis. 2015. CARD9-Dependent Neutrophil Recruitment Protects against Fungal
799   Invasion of the Central Nervous System. *PLoS Pathog* 11: e1005293.

800   71.   Kowalski, C.H., S.R. Beattie, K.K. Fuller, E.A. McGurk, Y.W. Tang, T.M. Hohl, J.J.
801   Obar, and R.A. Cramer, Jr. 2016. Heterogeneity among Isolates Reveals that Fitness in
802   Low Oxygen Correlates with Aspergillus fumigatus Virulence. *mBio* 7: :e01515-01516.

803   72.   Fermaintt, C.S., K. Sano, Z. Liu, N. Ishii, J. Seino, N. Dobbs, T. Suzuki, Y.X. Fu, M.A.
804   Lehrman, I. Matsuo, and N. Yan. 2019. A bioactive mammalian disaccharide associated
805   with autoimmunity activates STING-TBK1-dependent immune response. *Nat Commun*
806   10: 2377.

807   73.   Moretti, J., S. Roy, D. Bozec, J. Martinez, J.R. Chapman, B. Ueberheide, D.W.
808   Lamming, Z.J. Chen, T. Horng, G. Yeretssian, D.R. Green, and J.M. Blander. 2017.

STING Senses Microbial Viability to Orchestrate Stress-Mediated Autophagy of the
Endoplasmic Reticulum. *Cell* 171: 809-823.

74.   Alexopoulou, L., A.C. Holt, R. Medzhitov, and R.A. Flavell. 2001. Recognition of
double-stranded RNA and activation of NF-kappaB by Toll-like receptor 3. *Nature* 413:
732-738.

75.   Lund, J.M., L. Alexopoulou, A. Sato, M. Karow, N.C. Adams, N.W. Gale, A. Iwasaki,
and R.A. Flavell. 2004. Recognition of single-stranded RNA viruses by Toll-like receptor
7. *Proc Natl Acad Sci U S A* 101: 5598-5603.

76.   Gitlin, L., W. Barchet, S. Gilfillan, M. Cella, B. Beutler, R.A. Flavell, M.S. Diamond,
and M. Colonna. 2006. Essential role of mda-5 in type I IFN responses to
polyriboinosinic:polyribocytidylic acid and encephalomyocarditis picornavirus. *Proc
Natl Acad Sci U S A* 103: 8459-8464.

77.   Kato, H., O. Takeuchi, S. Sato, M. Yoneyama, M. Yamamoto, K. Matsui, S. Uematsu, A.
Jung, T. Kawai, K.J. Ishii, O. Yamaguchi, K. Otsu, T. Tsujimura, C.S. Koh, C. Reis e
Sousa, Y. Matsuura, T. Fujita, and S. Akira. 2006. Differential roles of MDA5 and RIG-I
helicases in the recognition of RNA viruses. *Nature* 441: 101-105.

78.   Yamamoto, M., S. Sato, K. Mori, K. Hoshino, O. Takeuchi, K. Takeda, and S. Akira.
2002. Cutting edge: a novel Toll/IL-1 receptor domain-containing adapter that
preferentially activates the IFN-beta promoter in the Toll-like receptor signaling. *J
Immunol* 169: 6668-6672.

79.   Beisswenger, C., C. Hess, and R. Bals. 2012. Aspergillus fumigatus conidia induce
interferon-beta signalling in respiratory epithelial cells. *Eur Respir J* 39: 411-418.

80.   Mehrad, B., R.M. Strieter, T.A. Moore, W.C. Tsai, S.A. Lira, and T.J. Standiford. 1999.
CXC chemokine receptor-2 ligands are necessary components of neutrophil-mediated
host defense in invasive pulmonary aspergillosis. *J Immunol* 163: 6086-6094.

81.   Lang, S., L. Li, X. Wang, J. Sun, X. Xue, Y. Xiao, M. Zhang, T. Ao, and J. Wang. 2017.
CXCL10/IP-10 Neutralization Can Ameliorate Lipopolysaccharide-Induced Acute
Respiratory Distress Syndrome in Rats. *PLoS One* 12: e0169100.

82.   Ichikawa, A., K. Kuba, M. Morita, S. Chida, H. Tezuka, H. Hara, T. Sasaki, T. Ohteki,
V.M. Ranieri, C.C. dos Santos, Y. Kawaoka, S. Akira, A.D. Luster, B. Lu, J.M.
Penninger, S. Uhlig, A.S. Slutsky, and Y. Imai. 2013. CXCL10-CXCR3 enhances the

development of neutrophil-mediated fulminant lung injury of viral and nonviral origin.
Am J Respir Crit Care Med 187: 65-77.

83.   Hagmann, C.A., A.M. Herzner, Z. Abdullah, T. Zillinger, C. Jakobs, C. Schuberth, C.
Coch, P.G. Higgins, H. Wisplinghoff, W. Barchet, V. Hornung, G. Hartmann, and M.
Schlee. 2013. RIG-I detects triphosphorylated RNA of Listeria monocytogenes during
infection in non-immune cells. *PLoS One* 8: e62872.

84.   Abdullah, Z., M. Schlee, S. Roth, M.A. Mraheil, W. Barchet, J. Bottcher, T. Hain, S.
Geiger, Y. Hayakawa, J.H. Fritz, F. Civril, K.P. Hopfner, C. Kurts, J. Ruland, G.
Hartmann, T. Chakraborty, and P.A. Knolle. 2012. RIG-I detects infection with live
Listeria by sensing secreted bacterial nucleic acids. *EMBO J* 31: 4153-4164.

85.   Liehl, P., V. Zuzarte-Luis, J. Chan, T. Zillinger, F. Baptista, D. Carapau, M. Konert, K.K.
Hanson, C. Carret, C. Lassnig, M. Muller, U. Kalinke, M. Saeed, A.F. Chora, D.T.
Golenbock, B. Strobl, M. Prudencio, L.P. Coelho, S.H. Kappe, G. Superti-Furga, A.
Pichlmair, A.M. Vigario, C.M. Rice, K.A. Fitzgerald, W. Barchet, and M.M. Mota. 2014.
Host-cell sensors for Plasmodium activate innate immunity against liver-stage infection.
Nat Med 20: 47-53.

86.   Cheng, Y. and J.S. Schorey. 2019. Extracellular vesicles deliver Mycobacterium RNA to
promote host immunity and bacterial killing. *EMBO Rep* 20:

87.   Cheng, Y. and J.S. Schorey. 2018. Mycobacterium tuberculosis-induced IFN-beta
production requires cytosolic DNA and RNA sensing pathways. *J Exp Med* 215: 2919-
2935.

FOOTNOTES:

[1]Research in this study was supported in part by institutional startup funds to JJO in part through the Dartmouth Lung Biology Center for Molecular, Cellular, and Translational Research grant P30 GM106394 (PI: Bruce A. Stanton) and Center for Molecular, Cellular and Translational Immunological Research grant P30 GM103415 (PI: William R. Green). JJO was partially supported by a Munck-Pfefferkorn Award from Dartmouth College and NIH R01 AI139133 grant. Work by AR was partially funded by NIH R01 AI114647 grant. AR and RAC are Investigators in the Pathogenesis of Infectious Diseases supported by the Burroughs Wellcome Fund. The funders had no role in the preparation or publication of the manuscript.

[2]Current Address: Loxo Oncology, Boulder, CO 80301

[3]Current Address: Clemson University, Biological Sciences Department, Clemson, SC, 29634

[4]Abbreviations: RLR, RIG-I like receptors; IPA, invasive pulmonary aspergillosis; GVHD, graft-versus-host disease; SNP, single nucleotide polymorphism; GMM, glucose minimal medium; PBS, phosphate buffered saline; i.t., intratracheal; BALF, bronchoalveolar lavage fluid; GMS, Grocott-Gomori methenamine silver; FLARE, fluorescent *Aspergillus* reporter; vita-PAMP, vitality pathogen-associated molecular pattern

FIGURE LEGENDS

Figure 1. Heat-killed swollen conidia have a decreased ability to induce CXCL10. C57BL/6J mice were challenged i.t. with $4x10^7$ CEA10 live or heat-killed swollen conidia. Forty hours later BALF was collected and IL-28/IFN-λ **(A)**, CXCL10 **(B)**, and TNFα **(C)** levels were determined by ELISA. Data are pooled from 2 independent experiments with 10-15 total mice per group. Statistical significance was determined using a one-way ANOVA with Dunn's post-test (*$p<0.05$; **$p < 0.01$).

Figure 2. *Aspergillus fumigatus* RNA induces an interferon response in a partially MDA5-dependent manner. **(A)** Primary murine fibroblasts from C57BL/6J mice were stimulated with LyoVec™ encapsulated polyI:C (pIC) or total RNA isolated from Af293 for 18 h. After stimulation cell supernatants were collected and analyzed for IFNα (left) and CXCL10 (right) by ELISA. Data from 2 independent experiments with 3 samples per group. **(B)** Primary murine fibroblasts from C57BL/6J and *Ifih1*$^{-/-}$ mice were stimulated with LyoVec™ encapsulated polyI:C (pIC) (MDA5 agonist), 5'-ppp (RIG-I agonist), or total RNA isolated from Af293 for 18 h. The total RNA pool isolated from Af293 was also treated with either RNase S1, RNase III, or DNase to degrade ssRNA, dsRNA, or DNA, respectively prior to encapsulation in LyoVec™. After stimulation cell supernatants were collected and analyzed for IFNα by ELISA. Data from 2 independent experiments with 3 samples per group. Statistical significance was determined by a two-way ANOVA with a Tukey's post-test (τττ $p<0.001$ – LyoVec™ only vs. experimental group; *** $p<0.001$ – B6 vs. *Ifih1*$^{-/-}$; ^^^ $p<0.001$ – enzyme treated vs. Af293).

<u>Figure 3.</u> *Mavs*[-/-] **mice have decreased interferon mRNA levels after** *Aspergillus fumigatus*

challenge particularly at later times. B6.129F2 and *Mavs*[-/-] mice were challenged with $4x10^7$

conidia of CEA10**.** Lungs were collected 3 or 48h post-inoculation. Total RNA was extracted

from whole lungs. Gene expression as determined by quantitative reverse transcription

polymerase chain reaction (qRT-PCR) using TaqMan probes for *Ifna* **(A)**, *Ifnb* **(B)**, *Ifnl2/3* **(C)**,

and *Ifng* **(D)**, which were normalized to *Gapdh* expression. Bars represent data means ± SEM

with each dot representing individual mice. Data are representative of results from at least 2

independent experiments with at least 3 mice per group. Data were analyzed using a Mann-

Whitney U-test (** $p < 0.01$; *** $p < 0.0001$).

<u>Figure 4.</u> *Mavs*[-/-] **mice have decreased STAT1 phosphorylation after** *Aspergillus fumigatus*

challenge at later times. **(A)** C57BL/6J mice were challenged with $4x10^7$ conidia of CEA10 or

Af293. Lungs were collected at 6, 24, and 48 hours post-inoculation and total protein was

extracted from whole lungs. Protein expression and phosphorylation was determined by Western

blot analysis. **(B)** B6.129F2 and *Mavs*[-/-] mice were challenged with $4x10^7$ conidia of CEA10**.**

Lungs were collected at 6- or 48-hours post-inoculation and total protein was extracted from

whole lungs. Protein expression and phosphorylation was determined by Western blot analysis.

(C) Western blots from panel B were quantified by densitometry. Bars represent data means ±

SEM with each dot representing individual mice. Data are representative of results from at least

2 independent experiments with at least 3 mice per group. Data were analyzed using a Mann-

Whitney U-test.

Figure 5. *Mavs*$^{-/-}$ **mice have an altered inflammatory milieu after** *Aspergillus fumigatus*

challenge in the airways. B6.129F2 and *Mavs*$^{-/-}$ mice were challenged with 4×10^{7} conidia of

CEA10. Bronchoalveolar lavage fluid (BALF) was collected 12 or 48h post-inoculation.

Cytokine and chemokine levels were determined using a Milliplex Mouse Cytokine &

Chemokine 32-plex (Millipore). Bars represent data means ± SEM. Data are representative of

results 2 independent experiments with 4-6 mice per group. Data were analyzed using a Mann-

Whitney U-test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$).

Figure 6. Expression of *Cxcl9* and *Cxcl10* are dependent on IFN signaling. C57BL/6J, *Ifnar*$^{-/-}$

, *Ifnlr*$^{-/-}$, and *Ifnar*$^{-/-}$ *Ifnlr*$^{-/-}$ (DKO) mice were challenged with 4×10^{7} conidia of CEA10. Lungs

were collected 48h post-inoculation and total RNA was extracted. Gene expression as

determined by quantitative reverse transcription polymerase chain reaction (qRT-PCR) using

TaqMan probes for *Cxcl9* **(A)** and *Cxcl10*, which were normalized to *Gapdh* expression. Bars

represent data means ± SEM with each symbol representing individual mice. Data are

representative of results from at least 2 independent experiments with at least 4 mice per infected

group. Data were analyzed using a Mann-Whitney U-test (n.s. = not significant; **** $p <$

0.0001).

Figure 7. Both *Mavs*$^{-/-}$ and *Ifih1*$^{-/-}$ mice are highly susceptible to *Aspergillus fumigatus*. (A-

C) B6.129F2 and *Mavs*$^{-/-}$ mice or **(D-F)** C57BL/6J and *Ifih1*$^{-/-}$ mice were challenged with 4×10^{7}

conidia of CEA10. **(A & D)** Survival analysis in immune-competent wild-type and knock-out

mice were tracked over the first 8-15 days. Statistical significance was assessed using a Mantel-

Cox log rank test (**(A)** $p = 0.0001$ and **(D)** $p = 0.0004$). **(B & E)** At 36 hpi, mice were euthanized

and BALF collected for quantification of macrophage and neutrophil recruitment to the airways. Data from 2 independent experiments for a total of 7-10 mice per group. Data are presented as box-and-whisker plots with Tukey whiskers and outliers displayed as dots. Statistical significance was determined using a Mann-Whitney U test. **(C & F)** At 40-42 hpi mice were euthanized and lungs saved for histological analysis. Formalin-fixed lungs were paraffin embedded, sectioned, and stained with GMS for analysis by microscopy. Representative lung sections from are shown to the right using the 40× objective. *A. fumigatus* germination rates were determined by microscopically counting both the number of conidia and number of germlings in GMS-stained sections. Data from 2 independent experiments with 4-6 mice per group. Statistical significance was determined using a Mann-Whitney U test (*p < 0.05; **p < 0.01).

Figure 8. _Ifih1$^{-/-}$_ mice have decrease antifungal killing by neutrophils _in vivo._ C57BL/6J and *Ifih1$^{-/-}$* mice were challenged with 3x10^7 conidia of CEA10 fluorescent *Aspergillus* reporter (FLARE). Bronchoalveolar lavage (BAL) fluid and lungs were harvested at 48 hours after infection and fungal uptake and viability were analyzed with flow cytometry. Fungal uptake **(A)** and viability **(B)** in neutrophils were measured. Data pooled from 2 independent experiments with 7-9 mice per group. Statistical significance was determined using a Mann-Whitney U test (***p < 0.01).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8