Title

Prime editing efficiently generates W542L and S621I double mutations in two ALS genes of maize

Yuan-Yuan **Jiang**[1†], Yi-Ping **Chai**[1†], Min-Hui **Lu**[2†], Xiu-Li **Han**[2], Yu **Zhang**[1], Qiang **Zhang**[1], Yun **Zhou**[3],

Xue-Chen **Wang**[1], Qi-Jun **Chen**[1,2*]

[1]State Key Laboratory of Plant Physiology and Biochemistry, College of Biological Sciences, China

Agricultural University, Beijing 100193, China

[2]Center for Crop Functional Genomics and Molecular Breeding, China Agricultural University, Beijing

100193, China

[3]Collaborative Innovation Center of Crop Stress Biology, Henan Province. Institute of Plant Stress

Biology, School of Life Science, Henan University, Kaifeng 475004, China

[†] These authors contributed equally to this work

[*] Correspondence (email: qjchen@cau.edu.cn)

Email addresses:

Yuan-Yuan Jiang: jiangyyuan111@163.com

Yi-Ping Chai: chaiyping@163.com

Min-Hui Lu: luminhui@cau.edu.cn

Xiu-Li Han: hanxiuli120@cau.edu.cn

Yu Zhang: yu_zhang68@163.com

Qiang Zhang: 1012224278@qq.com

Yun Zhou: zhoumouyun@henu.edu.cn

Xue-Chen Wang: 15201377438@163.com

Abstract

A novel and universal CRISPR/Cas-derived precision genome editing technology named prime editing was developed. However, low prime editing efficiency was shown in transgenic rice lines. We reasoned that enhancing pegRNA expression would be able to improve prime editing efficiency. In this report, we used two strategies to enhance pegRNA expression and constructed a prime editing vector harboring two pegRNA variants for W542L and S621I double mutations in *ZmALS1* and *ZmALS2*. As compared with previous reports in rice, we achieved much higher prime editing efficiency in maize. Our results are inspiring and provide a direction for optimization of plant prime editors.

Keywords: prime editing, *PE2*, pegRNA, CRISPR/Cas9, maize, *ALS*, herbicide-resistance

Background

Despite rapid advances in genome editing technologies, precision genome editing remains challenging [1]. Recently, a novel and universal precision genome editing technology named prime editing was developed and tested in mammalian cells [1]. More recently, seven groups reported applications of prime editors (PE) in rice and wheat, and six groups achieved prime-edited rice lines [2-8]. Excluding the data based on enriching strategies or unsuitable for comparison, total 164 prime-edited rice lines transformed with 39 prime editors targeting 11 endogenous rice genes were generated in these previous reports [2, 3, 5, 6, 8]. The highest editing efficiencies achieved by the five groups are from 2.22% to 31.3% [2, 3, 5, 6, 8]. Out of the total 164 prime-edited rice lines, only one line achieved by Xu et al. [6] harbors homozygous mutations whereas all the other lines mainly harbor chimeric mutations. These results indicate that the reported prime editing tools require significant optimization before they are adopted for use by plant researchers. In this paper, in comparison with the previous reports in rice, we achieved much higher prime editing efficiency in maize possibly owing to the optimized pegRNA expression, clean pegRNA sequences trimmed by tRNA and ribozyme processing systems, or both.

Results and discussion

We selected two maize acetolactate synthase (*ALS*) genes to test prime editing efficiency in maize and attempted to generate maize herbicide-resistance lines harboring P165S mutation or W542L/S621I double mutations [9] in *ZmALS1* and *ZmALS2* by prime editing (Fig. 1a). We used maize *Ubi1* promoter to drive the expression of maize codon-optimized *PE2* and used *OsU3* and *TaU3* promoters to drive the expression of pegRNA and sgRNA, respectively. In this way, we constructed two pGreen3 binary vectors [10, 11], pZ1PE3 and pZ1PE3b, for the same P165S mutation but based on two different strategies, PE3 and PE3b [1], respectively (Fig. 1a,b). To generate mutant maize lines harboring W542L/S621I double mutations, we based the W542L and S621I mutations on PE3b and PE3 strategies, respectively, and assembled two pegRNA and two nicking sgRNA variants into the same binary vectors (Fig. 1a,b). According to the structure of pegRNAs, a hairpin can potentially form between the PBS and the protospacer of a pegRNA (Fig. 1a), and thus will possibly weaken the pegRNA activity. Based on this possibility, we reasoned that enhancing pegRNA expression will be able to improve editing efficiency. We used two strategies to enhance pegRNA expression: doubling pegRNA expression cassettes and using two promoter systems to drive pegRNA expression, together with use of the tRNA, ribozyme, and Csy4 RNA processing systems [12, 13] when appropriate (Fig. 1b). In this way, we generated two prime editors named pZ1WS-Csy4 and pZ1WS (Fig. 1b). We also generate two additional prime editors including p4xZ1PE3-Csy4 and p4xZ1PE3b-Csy4 for comparison with pZ1PE3 and pZ1PE3b, respectively. The p4xZ1PE3-Csy4 and p4xZ1PE3b-Csy4 editors have similar structure to pZ1WS-Csy4 but harbor redoubled pegRNA and sgRNA expression cassettes for the P165S mutation. The pZ1WS vector was constructed as a control of pZ1WS-Csy4 for comparison of Csy4 and non-Csy4 systems. We transformed maize with these six PE vectors via the Agrobacterium-mediated method. We obtained no transgenic lines transformed with the three Csy4 vectors. Almost at the same time, we found that transformations with 13 additional pGreen3 or pCambia binary vectors harboring the Csy4-P2A-ZsGreen fusion gene, including 1 maize and 12 Arabidopsis transformations, were unsuccessful, suggesting Csy4 protein severely affects Agrobacterium-mediated transformation of at least these two plant species.

To detect targeted mutations, we amplified fragments spanning the targets from genomic DNA of

the transgenic lines and detected mutations by direct sequencing of PCR fragments. We observed

no editing for the 71 and 156 lines transformed with pZ1PE3 and pZ1PE3b, respectively. Fortunately,

we observed that 7 out of 16 lines transformed with pZ1WS harbor S621I edits (Fig. 1c,d).

Interestingly, one line displayed homozygous mutations in both *ZmALS1* and *ZmALS2*.

We cloned the PCR products from the 6 lines harboring chimeric or heterozygous mutations and

sequenced ~86 clones for each PCR fragment (Additional file 1: Table S1). Interestingly, we observed

the W542L edits in some cloned fragments, suggesting that low-frequency mutations were not able

to be displayed in sequencing chromatograms (Additional file 1: Table S1). To further analyze the

low-frequency W542L and/or S621I mutations, we cloned the PCR fragments from the line (#4)

harboring homozygous mutations and an additional line (#3) harboring no mutations according to

direct sequencing of PCR fragments. We observed the S621I edits in some cloned fragments from

#3, further demonstrating that low-frequency mutations could not be reflected by direct

sequencing of PCR products (Additional file 1: Table S1). Three cloned fragments harbor W542L and

S621I double mutations, suggesting that PE can simultaneously edit two non-allelic targets in a cell.

We also cloned 6 PCR fragments from three pZ1PE3 and three pZ1PE3b transgenic lines, sequenced

the cloned PCR fragments, but found no mutations in total 213 *ZmALS1* and 196 *ZmALS2* clones

(Additional file 1: Table S1).

We observed two types of undesired byproducts from the sequencing results of cloned PCR

fragments. One type is derived from the pegRNA scaffold, which acts as extended RT templates and

thus introduces unwanted edits; the other is only involved in editing of multiple nucleotides and is

derived from unbiased and well-balanced double-strand repair (rather than single-strand-biased

repair) of multiple mismatches in heteroduplex DNA, which leads to incomplete editing of multiple

nucleotides (Fig. 2a). In addition, the two mechanisms of forming these two types of byproducts

were sometimes overlapped (Fig. 2a). We observed high frequencies of byproducts at the S621

target: 23.8% and 7.2% derived from the pegRNA scaffold and the double-strand even DNA repair,

respectively (Fig. 2b). For the W542 target, frequencies of both types of byproducts remained to be

investigated because of limited number of cloned PCR fragments harboring W542L edits. Since the

T0 line #4 harbors homozygous mutations in the two *ALS* and the T0 line #15 harbors obvious

pegRNA scaffold-derived byproducts (Fig. 1d), we excluded these two lines for possibly unbiased

analysis of the two types of byproducts. The recalculated frequencies of the byproducts for the

S621I edits are 18.1% and 15.3% for the pegRNA scaffold-derived and double-strand even DNA

repair-derived byproducts, respectively (Additional file 1: Table S1), indicating that frequencies of

byproducts excluding the two lines are still high. Surprisingly, we observed no indel-type

byproducts, suggesting that indels induced by PE are infrequent in maize.

To further figure out the low-frequency mutations and byproducts, we analyzed all the 243 lines

transformed with the three PE vectors by next-generation sequencing (NGS) of PCR amplicons. Out

of the 227 T0 lines transformed with pZ1PE3 or pZ1PE3b, we revealed only one line transformed

with pZ1PE3 harboring the low frequency (0.07%, 21/30306) P165S edits. On the contrary, we found

that 12 out of the 16 T0 lines transformed with pZ1WS harbor the S621I edits, and 6 out of the 16

lines also harbor the W542L edits (Additional file 1: Table S2). For the S621 target, average

frequencies of pegRNA scaffold-derived and double-strand even DNA repair-derived byproducts

from NGS are 17.5% and 8.5%, respectively; those excluding the #4 and #15 lines are 12.1% and

16.2%, respectively (Additional file 1: Table S2). We observed no byproducts for the W542L edits

from NGS possibly owing to the low frequency of desired edits, which affected analysis of

byproducts with lower frequency.

We repeatedly transformed maize with pZ1WS after we observed that our first transformation

could not generate many transgenic lines. In the second transformation, we obtained 46 transgenic

lines. We detected targeted mutations induced by prime editing and the results demonstrated again

that the prime editing efficiency of pZ1WS was much higher than previous reports (Fig. 2c and

Additional file 1: Fig. S1). Interestingly, this time we observed the signals of the W542L edits in the

sequencing chromatograms (Fig. 2d and Additional file 1: Fig. S1). These results suggest that we will

be able to achieve homozygous mutants harboring W542L and S621I double mutations in both

ZmALS1 and *ZmALS2* in the progeny of some lines, such as #38 (Additional file 1: Fig. S1). In

addition, we achieved two additional lines harboring the S621I homozygous mutations in the two

ALS genes (Additional file 1: Fig. S2), and these two lines, together with the above described line,

can be used to generate homozygous double mutants harboring W542L and S621I double mutations

in the progeny. Not surprisingly, we again observed pegRNA scaffold-derived byproducts in the

sequencing chromatograms (Additional file 1: Fig. S1). We cloned PCR products from the 4 lines

harboring strong W542L edit signals in chromatograms (Additional file 1: Fig. S1) and sequenced

~106 clones for each PCR fragment (Fig. 2e,f and Additional file 1: Table S3). The results indicated that 11.5% cloned fragments from the line #41 harbor S621I edits although direct sequencing of PCR fragment from this line displayed weak S621I edit signals in chromatograms (Additional file 1: Fig. S1 and Table S3). We detected no pegRNA scaffold-derived byproducts from the W542L edits (Fig. 2e,f and Additional file 1: Table S3), suggesting low frequency of pegRNA scaffold-derived byproducts was produced at this target and frequency of byproducts depends on targets. Out of 242 cloned PCR fragments harboring edits, 17.8% (43/242) harbor W542L and S621I double mutations in *ZmALS1* or *ZmALS2* (Fig. 2f), demonstrating again that PE can simultaneously edit two non-allelic targets in a cell. For the W542L edits, the protospacer of the sgRNA nicking non-edited strand fully matches the edited *ZmALS1* target but has a mismatch adjacent to PAM of the edited *ZmALS2* target (Figure 1a), therefore actually PE2 was used for editing *ZmALS2* although PE3b was done for *ZmALS1*. Editing frequencies at the W542 target of *ZmALS1* and *ZmALS2* are 7.1% and 39.7%, respectively (Additional file 1: Table S3), suggesting that PE2 has much higher editing efficiencies than PE3b.

Collectively, we achieved sufficiently high prime editing efficiency in maize possibly owing to the optimized pegRNA expression, clean pegRNA sequences trimmed by tRNA and ribozyme processing systems, or both. In addition, we observed high frequency the pegRNA scaffold-derived byproducts and to reduce them we propose a rule of termination to design pegRNAs: one to a few nucleotides (C, GC, or TGC, et al.) of the pegRNA scaffold adjoining RT template can be used as termination signals at genomic DNA matching RT templates.

Although we achieved sufficient high prime editing efficiency, a few questions remain to be answered. Is the pegRNA sequence, the optimized pegRNA expression, or both the decisive factor for the high efficiency of the S621I edits? Are the Pol2-Pol3 fusion promoters (such as 35S-CmYLCV-U6) or only Pol2 promoters (such as 35S-CmYLCV), together with the tRNA and ribozyme RNA processing systems, better systems to drive pegRNA expression than the commonly used Pol3 promoters (such as OsU3 and TaU3)? Is doubling or redoubling pegRNA expression cassettes crucial for high prime editing efficiency? Is clean pegRNA sequences trimmed by tRNA and ribozyme processing systems, or both vital for high prime editing efficiency? Why does Csy4 system affect maize transformation? To facilitate to answer these questions, we are developing additional user-friendly tools for a variety of plant species and promise to share them with researchers as soon as

possible. Application of these tools to test more pegRNA variants in amenable plants, such as rice,

will help to clarify decisive factors for high prime editing efficiency in plants and thus facilitate the

development of more powerful prime editors for high throuput precise genome editing in plants.

Conclusions

As compared with previous reports in rice, we achieved much higher prime editing efficiency in

maize: 53.2% (33/62) and 6.5% (4/62) transgenic lines harbor S621I and W542L mutations,

respectively, in *ZmALS1* and/or *ZmALS2*; 4.8% (3/62) and 4.8% (3/62) lines harbor homozygous

S621I mutations and S621I/W542L double mutations, respectively, in the two *ALS* genes. We also

observed high frequency of the pegRNA scaffold-derived byproducts and to potentially overcome

this defect, we propose a rule to design pegRNAs. Our results are inspiring and provide a new

reference standard for future optimization of prime editors. In addition, the pZ1WS prime editor

generated in this study is useful for breeding transgene-free maize lines harboring W542L and S621I

double mutations in the two ALS genes, which confer resistance to multiple herbicides functioning

as ALS inhibitors.

Methods

All primers used in this study are listed in Additional file 1: Table S4 and sequences of PE2 and

pegRNA expression cassettes are listed in Additional file 1: Supplemental material. Vectors

described in this study together with their annotated sequences are available from Addgene and/or

MolecularCloud (GenScript).

Vector construction

We replaced the *Xba*I-*Sbf*I fragment of pUC57-dCas9 [14] with a synthetic fragment, resulting in

the generation of pUC57-PE2P1. We replaced the *Mlu*I and *Sac*I fragment of pUC57-PE2P1 with

another synthetic fragment, resulting in the generation of pUC57-PE2. We modified the maize Ubi1

promoter in pG3R23-U3Ub [11] so that the first of the two *Pst*I sites, the first of the four *Xba*I sites

(the last 3 *Xba*I sites are blocked by Dam methylation), the *Nco*I site, and the *Eco*RI site in the

promoter were disrupted. We replaced the *Xba*I and *Sac*I fragment of the modified pG3R23-U3Ub

with the *Xba*I and *Sac*I fragment of pUC57-PE2, resulting in the generation of pG3R23-PE2U3A. We

replaced the *Eco*RI-*Sac*II fragment of pG3R23-PE2U3A with the *Eco*RI-*Sac*II fragment of pG3GB411

199 [11], resulting in the generation of pG3GB-PE2U3A. To generate the final PE binary vectors

200 harboring a single pegRNA, we conducted the Golden Gate reaction to insert a synthetic pegRNA-

201 OsU3t-TaU3p-sgRNA fragment into the two *Bsa*I sites of pG3GB-PE2U3A. In this way, we generated

202 pZ1PE3 and pZ1PE3b.

203 We replaced the *Asc*I-*Pac*I fragment of pL21-iCre [11] with a synthetic fragment of 35S-CmYLCV-

204 Csy4-P2A-ZsGreen-polyA-Csy4-BB-HDV-HSPt, resulting in the generation of pL2L1-35Csy4. We

205 replaced the *Xba*I-*Eco*RI fragment of pL2L1-35Csy4 with a synthetic fragment of U6-tGly-BB-HDV-

206 HSPt, resulting in the generation of pL2L1-35C-U6. We replaced the *Hin*dIII-*Eco*RI fragment of pR14-

207 BWM [11] with an insert prepared by annealing two oligos oHEASmE-F/R, resulting in the

208 generation of pR1R4-HEAS. We replaced the *Eco*RI-*Aar*I fragment of pR1R4-HEAS with the *Eco*RI-

209 *Spe*I fragment of synthetic OsAct1p-CP4EPSPS-NOSt, resulting in the generation of pR1R4-Gly. We

210 amplified the U3p-BB-sgRNA cassette from pG3R23-PE2U3A with primers OsU3p-AsF and TaU3t-

211 EcR, purified the PCR products, digested them with *Asc*I and *Eco*RI, and allowed them to ligate with

212 *Asc*I and *Eco*RI-digested pR1R4-Gly and pL43-U3-Bar, resulting in the generation of pR1R4-U3G.0

213 and pL4L3-U3B.0, respectively.

214 We generated the two binary vectors harboring four pegRNA and four sgRNA cassettes in two

215 steps. First, we set up the Golden Gate reaction to clone synthetic fragments harboring pegRNAs

216 and/or sgRNAs into the destination vector pG3R23-PE2U3A and the three entry vectors including

217 pL2L1-35Csy4 or pL2L1-35C-U6, pR1R4-U3G.0, and pL4L3-U3B.0 (Additional file 1: Supplemental

218 material). Second, we set up the MultiSite Gateway reaction to assemble the final PE vectors. In this

219 way, we generated pZ1WS-Csy4 and pZ1WS. We constructed p4xZ1PE3-Csy4 and p4xZ1PE3b-Csy4

220 harboring redoubled pegRNA and sgRNA expression cassettes for the P165S mutation in a similar

221 way to the generation of pZ1WS-Csy4.

222 **Maize transformation and analysis of prime editing**

223 We separately transformed the six PE vectors, pZ1PE3, pZ1PE3b, p4xZ1PE3-Csy4, p4xZ1PE3b-

224 Csy4, pZ1WS-Csy4, and pZ1WS into the engineered Agrobacterium strain LBA4404/pVS1-VIR2 to

225 generate strains harboring the ternary vector system [11]. We separately used these strains to

226 transform maize ND73, a B73-derived inbred line, according to the published protocols [11].

227 To analyze the P165S mutations in *ZmALS1* and *ZmALS2*, we simultaneously amplified two

228 fragments spanning the target site in the two *ALS* genes from genomic DNA of transgenic lines by

PCR using primers ALS1&2P-F/R. We then submitted purified PCR products for direct sequencing

with the primer ALS1&2P-F. To analyze the W542L and S621I mutations in *ZmALS1* and *ZmALS2*, we

simultaneously amplified two fragments spanning the two target sites in the two *ALS* genes by PCR

using primers ALS1&2WS-F/R and submitted purified PCR products for direct sequencing with

primer ALS1&2WS-F2. Since we designed the primers according to homologous sequences of

ZmALS1 and *ZmALS2*, the PCR fragments contain both *ZmALS1* and *ZmALS2* sequences. Since

ZmALS1 and *ZmALS2* are highly homologous, the co-existence of two PCR fragments can be

determined according to the double peaks of the sequencing chromatograms. Similarly, chimeric

and heterozygous mutations can also be determined according to the double peaks. For sequencing

cloned PCR fragments, we picked ~86 or ~106 colonies from each transformation and submitted

them for Sanger sequencing. We also used Hi-TOM assay [15] with 0.5% threshold to analyze the

mutations in T0 transgenic plants. To analyze the P165S, W542L, and S621I mutations in *ZmALS1*

and *ZmALS2*, we simultaneously amplified two fragments spanning the target site in the two *ALS*

genes from genomic DNA of transgenic lines by PCR using primers ALS1&2P-NGSF/R, ALS1&2W-

NGSF/R, and ALS1&2S-NGSF/R, respectively.

Supplementary information

Additional file 1: **Figure S1.** Sequencing chromatograms from 7 prime-edited lines harboring W542L

edits. **Figure S2.** Sequencing chromatograms from 2 prime-edited lines harboring homozygous

S621I edits. **Table S1.** Edits and byproducts revealed from cloned PCR fragments. **Table S2.** Analysis

of mutations in T0 transgenic plants by NGS with 0.5% threshold. **Table S3.** Edits and byproducts

from the 4 additional lines. **Table S4.** Sequences of primers, targets, and rtT-PBS of pegRNAs.

Supplemental material. Sequences of PE2 and pegRNA expression cassettes.

Acknowledgments

The transgenic maize lines were created by the Maize Functional Genomic Platform of China

Agricultural University. We thank our colleagues from the platform for help in maize transformation.

256   Authors' contributions

257   QJC, YZhou, and XCW conceived and designed the research. YYJ, YPC, MHL, XLH, YZ, and QZ

258   conducted experiments and analyzed the data. QJC, YZhou, and XCW wrote the paper. All authors

259   read and approved the final version.

260   Funding

261   This work was supported by grants from the National Transgenic Research Project (grant nos.

262   2019ZX08010003-001-011 and 2016ZX08009002), the National Crop Breeding Fund (grant no.

263   2016YFD0101804), the National Natural Science Foundation of China (grant nos. 31872678 and

264   31670371).

265   Availability of data and materials

266   Vectors described in this study together with their annotated sequences are available from Addgene

267   and/or MolecularCloud (GenScript).

268   Ethics approval and consent to participate

269   Not applicable.

270   Consent for publication

271   Not applicable.

272   Competing interests

273   The authors declare that they have no competing interests.

274   Author details

275   Not applicable.

276   References

1.   Anzalone AV, Randolph PB, Davis JR, Sousa AA, Koblan LW, Levy JM, Chen PJ, Wilson C, Newby GA, Raguram

A, Liu DR: Search-and-replace genome editing without double-strand breaks or donor DNA. *Nature* 2019,

576:149-157.

2.   Xu R, Li J, Liu X, Shan T, Qin R, Wei P: Development of a plant prime editing system for precise editing in the

rice genome. *Plant Communications* 2020:100043.

3.   Hua K, Jiang Y, Tao X, Zhu JK: Precision genome engineering in rice using prime editing system. *Plant*

Biotechnol J 2020.

4.   Butt H, Rao GS, Sedeek K, Aman R, Kamel R, Mahfouz M: Engineering herbicide resistance via prime editing
in rice. *Plant Biotechnol J* 2020.

5.   Lin Q, Zong Y, Xue C, Wang S, Jin S, Zhu Z, Wang Y, Anzalone AV, Raguram A, Doman JL, et al: Prime genome
editing in rice and wheat. *Nature Biotechnology* 2020.

6.   Xu W, Zhang C, Yang Y, Zhao S, Kang G, He X, Song J, Yang J: Versatile Nucleotides Substitution in Plant Using
an Improved Prime Editing System. *Molecular Plant* 2020.

7.   Tang X, Sretenovic S, Ren Q, Jia X, Li M, Fan T, Yin D, Xiang S, Guo Y, Liu L, et al: Plant Prime Editors Enable
Precise Gene Editing in Rice Cells. *Molecular Plant* 2020.

8.   Li H, Li J, Chen J, Yan L, Xia L: Precise Modifications of Both Exogenous and Endogenous Genes in Rice by
Prime Editing. *Molecular Plant* 2020.

9.   Kawai K, Kaku K, Izawa N, Shimizu T, Fukuda A, Tanaka Y: A novel mutant acetolactate synthase gene from
rice cells, which confers resistance to ALS-inhibiting herbicides. *Journal of Pesticide Science* 2007, 32:89-98.

10.   Zhang Y, Zhang Q, Chen QJ: Agrobacterium-mediated delivery of CRISPR/Cas reagents for genome editing in
plants enters an era of ternary vector systems. *Sci China Life Sci* 2020.

11.   Zhang Q, Zhang Y, Lu MH, Chai YP, Jiang YY, Zhou Y, Wang XC, Chen QJ: A Novel Ternary Vector System United
with Morphogenic Genes Enhances CRISPR/Cas Delivery in Maize. *Plant Physiol* 2019, 181:1441-1448.

12.   Zhang Q, Xing HL, Wang ZP, Zhang HY, Yang F, Wang XC, Chen QJ: Potential high-frequency off-target
mutagenesis induced by CRISPR/Cas9 in Arabidopsis and its prevention. *Plant Mol Biol* 2018, 96:445-456.

13.   Cermak T, Curtin SJ, Gil-Humanes J, Cegan R, Kono TJY, Konecna E, Belanto JJ, Starker CG, Mathre JW,
Greenstein RL, Voytas DF: A Multipurpose toolkit to enable advanced genome engineering in plants. *Plant
Cell* 2017, 29:1196-1217.

14.   Xing HL, Dong L, Wang ZP, Zhang HY, Han CY, Liu B, Wang XC, Chen QJ: A CRISPR/Cas9 toolkit for multiplex
genome editing in plants. *BMC Plant Biol* 2014, 14:327.

15.   Liu Q, Wang C, Jiao X, Zhang H, Song L, Li Y, Gao C, Wang K: Hi-TOM: a platform for high-throughput tracking
of mutations induced by CRISPR/Cas systems. *Sci China Life Sci* 2019, 62:1-7.

Figure legends for Fig. 1

Fig. 1 Efficiency of prime editing in maize. **a** Sequences and their relationship of the two *ALS* genes, the pegRNA targets, the pegRNA RT templates, the pegRNA primer binding sites (PBS), and the sgRNA targets for nicking non-edited strands used for generation of 4 maize prime editing vectors. The PAMs, nicking sites, RT template and PBS lengths, and introduced mutations are indicated. **b** T-DNA structures of 4 PE vectors. The pZ1WS and pZ1WS-Csy4 vectors are designed for editing double-targets. Each double-PE vector includes two expression cassettes for each of two pegRNAs or sgRNAs. Two strategies are designed in the peg1&2: one is based on Csy4 RNA processing system and the other is based on tRNA and HDV ribozyme-based RNA processing systems integrated with double-drives of polymerase II (35S enhancer-CmYLCV) and III (shortened U6-26) promoters. The structure of pegRNA and sgRNA expression cassettes in pegR1-sgR1 and pegR2-sgR2 is the same as that in pZ1PE3/3b. The expressions of two sgRNAs in sgR1-sgR2 are driven by OsU3p and TaU3p, respectively. **c** Prime editing efficiencies based on direct sequencing of PCR products. The transgenic lines transformed with pZ1WS-Csy4 were not able to be achieved. **d** The sequencing chromatograms of PCR fragments from the prime-edited plants transformed with pZ1WS. Double peaks represent heterozygous or chimeric mutations and an asterisk indicates a mutation induced by PE. Note that the first mutation in #15 is the unwanted mutation introduced by pegRNA scaffold.

Figure legends for Fig. 2

Fig. 2 Desired edits and unwanted byproducts in the two transformations with pZ1WS. **a** Desired

edits and unwanted byproducts at the two target sites in the first transformation. Only partial

sgRNA sequence is shown. The homologous sgRNA-rtT sequences shared by all the aligned

sequences are shaded with yellow and mutated nucleotides are indicated with red letters. Number

of cloned PCR fragments harboring the same edits is indicated in parentheses and a vertical line

indicates the same type of byproducts. For convenience, the byproducts derived from the two

mechanisms were assorted into the pegRNA scaffold-derived byproducts. **b** Summary of the desired

edits and byproducts in the first transformation according to sequences of cloned PCR fragments. **c**

Prime editing efficiency achieved in the second transformation. Strong and Weak, T0 lines harboring

strong and weak peak signals for edits, respectively, in the sequencing chromatograms. Both, total

No. of lines harboring strong and weak edits. **d** Sequencing chromatograms from a prime-edited line

harboring W542L edits and obtained in the second transformation. Double peaks represent

heterozygous or chimeric mutations and an asterisk indicates a mutation induced by PE. Note that

the first mutation in the S621 target is the pegRNA scaffold-derived byproducts. **e** Desired edits and

unwanted byproducts at the two target sites in the second transformation. **f** Summary of the

desired edits and byproducts in the second transformation.

Figure 1

a

P165S mutation based on PE3b
CCG (Pro) > AGT (Ser)

P165S mutation based on PE3
CCG (Pro) > AGT (Ser)

W542L mutation based on PE3b
TGG (Trp) > TTA (Leu)

S621I mutation based on PE3
AGT (Ser) > ATT (Ile)

T1, prime editing target; T2, target for nicking non-edited strand; rtT, RT template; RP/PR, rtT-PBS or PBS-rtT of pegRNA
ALS, ZmALS1&2; S, C (*ZmALS1*) or G (*ZmALS2*); Y, T (*ZmALS1*) or C (*ZmALS2*)

b

pZ1PE3 or pZ1PE3b for the P165S mutation

RB ▷ OsU3p:pegRNA ▷ TaU3p:sgRNA ▷ Ubi1p:PE2-E9t ▷ Gly ▷ Bar ▷ LB

pegR1/2, pegRNA1/2
sgR1/2, sgRNA1/2
pegR1&sgR1: the W542L mutation
pegR2&sgR2: the S621I mutation

pZ1WS-Csy4 or pZ1WS for the W542L and S621I mutations

RB ▷ pegR1-sgR1 ▷ Ubi1p:PE2-E9t ▷ pegR1&2 ▷ sgR1&2 ▷ Gly ▷ pegR2-sgR2 ▷ Bar ▷ LB

pZ1WS-Csy4
CaMV 35S enhancer | CmYLCV ▷ Csy4ZsG // pA | Csy4 pegR1 HDV | Csy4 pegR2 HDV | Csy4 | HSPt

pZ1WS
CaMV 35S enhancer | CmYLCV ▷ U6 ▷ tGly | pegR1 HDV | tMet pegR2 HDV | polyT-HSPt

Csy4ZsG, Csy4-P2A-ZsGreen; pA, polyA; tGly/Met, tRNA(Gly/Met)

c

Vector	Mutation	No. of double mutants		No. of single mutants		Total	Ratio of mutants
		Ho	He/Ch	Ho	He/Ch		
pZ1PE3b	P165S	0	0	0	0	156	0
pZ1PE3	P165S	0	0	0	0	71	0
pZ1WS	W542L	0	0	0	0	16	0
	S621I	1	6	0	0	16	43.8%

Ho, homozygous; He/Ch, heterozygous or chimeric; Double mutants, T0 lines harboring mutations in *ZmALS1* and *ZmALS2*; Single mutants, T0 lines harboring mutations in *ZmALS1* or *ZmALS2*; Total, total No. of T0 lines.

d

#4 A C C T A T   #8 C C C T A G   #15 T T G C C T A T G A T A C C T A T

(The sequencing chromatograms of #5, #6, #14, and #16 are similar to that of #8)

Figure 2

a

S621I edits
```
CACCAGGAGCATGTGTTGCCTATGATCCCTAGTGGTGGGGCTTTC Ref 1
GGCACCGAGTCGGTGCTGCCTATGATACCTATTGGTGGGGCTTTC Ref 2
CACCAGGAGCATGTGTTGCCTATGATACCTATTGGTGGGGCTTTC Seq (×154)
CACCAGGAGCATGTGTTGCCTATGATACCTAGTGGTGGGGCTTTC Seq (×10)
CACCAGGAGCATGTGTTGCCTATGATCCCTATTGGTGGGGCTTTC Seq (×6)
CACCAGGAGCATGTGCTGCCTATGATACCTATTGGTGGGGCTTTC Seq (×41)
CACCAGGAGCATGTGCTGCCTATGATCCCTAGTGGTGGGGCTTTC Seq (×10)
CACCAGGAGCATGTG.TGCCTATGATACCTATTGGTGGGGCTTTC Seq (×1)
CACACCGAGTCGGTGCTGCCTATGATCCCTAGTGGTGGGGCTTTC Seq (×1)
```

W542L edits
```
ATGGTGGTGCAGTGGGAGGACAGGTTCTATAAGGC Ref 1
ATGGTGGTGCAGTTAGAAGACAGGTTCTAGCACCG Ref 3
ATGGTGGTGCAGTTAGAAGACAGGTTCTATAAGGC Seq (×7)
ATGGTGGTGCAGTGGGAAGACAGGTTCTATAAGGC Seq (×1)
ATGGTGGTGCAGTTAGAAGACAGGTTCTAGCAGGC Seq (×1)
```

Ref 1/2/3, reference sequence 1/2/3; Ref 1, ZmALS1 & 2
Ref 2, sgRNA-rtT-PBS; Ref 3, PBS-rtT-sgRNA
Seq, sequence of cloned PCR fragments

b

Mutation	De	Sc	Re	To	De/To	Sc/To	Re/To
S621I	154	53	16	223	69.1%	23.8%	7.2%
W542L	7	1	1	9	77.8%	11.1%	11.1%

De, Sc, or Re, No. of clones harboring specified mutations
De, desired edits
Sc, pegRNA scaffold-derived byproducts
Re, double-strand even DNA repair-derived byproducts
To, total No. of cloned fragments harboring mutations

c

Mutation	No. of prime-edited lines			Total No. of	Ratio of prime-edited lines		
	Strong	Weak	Both	T0 lines	Strong	Weak	Both
W542L	4	3	7	46	8.7%	6.5%	15.2%
S621I	26	7	33	46	56.5%	15.2%	71.7%
W542L & S621I	3	4	7	46	6.5%	8.7%	15.2%

d

e

S621I edits
```
CACCAGGAGCATGTGTTGCCTATGATCCCTAGTGGTGGGGCTTTC Ref 1
GGCACCGAGTCGGTGCTGCCTATGATACCTATTGGTGGGGCTTTC Ref 2
CACCAGGAGCATGTGTTGCCTATGATACCTATTGGTGGGGCTTTC Seq (x117)
CACCAGGAGCATGTGTTGCCTATGATCCCTATTGGTGGGGCTTTC Seq (x15)
CACCAGGAGCATGTGTTGCCTATGATACCTAGTGGTGGGGCTTTC Seq (x11)
CACCAGGAGCATGTGCTGCCTATGATACCTATTGGTGGGGCTTTC Seq (x34)
CACCAGGAGCATGTGCTGCCTATGATCCCTAGTGGTGGGGCTTTC Seq (x4)
CACCAGGAGCATGTGCTGCCTATGATCCCTATTGGTGGGGCTTTC Seq (x2)
CACCAGGAGCATGTGCTGCCTATGATACCTAGTGGTGGGGCTTTC Seq (x1)
CACCAGGAGTCGGTGCTGCCTATGATACCTATTGGTGGGGCTTTC Seq (x1)
```

W542L edits
```
ATGGTGGTGCAGTGGGAGGACAGGTTCTATAAGGC Ref 1
ATGGTGGTGCAGTTAGAAGACAGGTTCTAGCACCG Ref 3
ATGGTGGTGCAGTTAGAAGACAGGTTCTATAAGGC Seq (×93)
ATGGTGGTGCAGTTAGAGGACAGGTTCTATAAGGC Seq (×4)
ATGGTGGTGCAGTGGGAGACAGGTTCTATAAGGC Seq (×3)
```

Ref 1/2/3, reference sequence 1/2/3; Ref 1, ZmALS1 & 2
Ref 2, sgRNA-rtT-PBS; Ref 3, PBS-rtT-sgRNA
Seq, sequence of cloned PCR fragments

f

Mutation	De	Sc	Re	To	De/To	Sc/To	Re/To
W542L	93	0	7	100	93.0%	0.0%	7.0%
S621I	117	42	26	185	63.2%	22.7%	14.1%
WS	27		16	43	62.8%		37.2%

De, Sc, or Re, No. of clones harboring specified mutations
De, desired edits
Sc, pegRNA scaffold-derived byproducts
Re, double-strand even DNA repair-derived byproducts
To, total No. of cloned fragments harboring mutations
WS, W542L and S621I double mutations