1

# Ghat: An R package for identifying adaptive polygenic traits

**Medhat Mahmoud**[*,‡]**, Mila Tost**[*,‡]**, Ngoc-Thuy Ha** [†]**, Henner Simianer** [†,‡]**, and Timothy**

**Beissinger**[*,‡,1]

[*]Department of Crop Science, University of Goettingen, Goettingen, Germany

[†]Department of Animal Sciences, University of Goettingen, Goettingen, Germany

[‡]Center for Integrated Breeding Research, University of Goettingen, Goettingen, Germany

[1] Corresponding author

Timothy Beissinger (beissinger@gwdg.de)

19
20

**Abstract**

Identifying selection on polygenic complex traits in crops and livestock is important for understanding evolution and helps prioritize important characteristics for breeding. The QTL that contribute to polygenic trait variation often exhibit small or infinitesimal effects. This hinders the ability to detect QTL controlling polygenic traits because enormously high statistical power is needed for their detection. Recently, we circumvented this challenge by introducing a method to identify selection on complex traits by evaluating the relationship between genome-wide changes in allele frequency and estimates of effect-size. The approach involves calculating a composite-statistic across all markers that captures this relationship, followed by implementing a linkage disequilibrium-aware permutation test to evaluate if the observed pattern differs from that expected due to drift during evolution and population stratification. In this manuscript, we describe "Ghat", an R package developed to implement this method to test for selection on polygenic traits. We demonstrate the package by applying it to test for polygenic selection on 15 published European wheat traits including yield, biomass, quality, morphological characteristics, and disease resistance traits. Moreover, we applied Ghat to different simulated populations with different breeding history and genetic architecture. The results highlight the power of Ghat to identify selection on complex traits. The Ghat package is accessible on CRAN, the Comprehensive R Archival Network, and on GitHub.

## Introduction

Many important traits in plants, animals, and humans are polygenic and depend on the cumulative effects of many loci, each contributing a small proportion of the total genetic variation [1]. Studying the evolution and selection history of such traits can be challenging due to the small effects of the quantitative trait loci (QTL) that contribute to genetic variation. However, regardless of effect size, the theory of natural selection tells us that alleles with positive effects on fitness will tend to increase in frequency over successive generations [2]. Several statistical methods have been developed to identify loci carrying such beneficial alleles. These methods were reviewed in [3], and include approaches relying on polymorphism variability, haplotype frequency, linkage disequilibrium, allele frequency change over time, and techniques that combine some or all of these signals. However, in the case of highly polygenic traits, it is difficult or impossible to identify regions carrying individually selected alleles/haplotypes due to their small effects, regardless of which sophisticated methodologies are leveraged. For instance, it has been shown that selected loci are difficult to identify when per-locus selection intensity is low [3], as is the case with low heritability traits and those controlled by many alleles of small effect [4]. This imposes serious limitations to the previous methods when identifying selection on polygenic traits (e.g. [5]).

Recently, large sample sizes, especially as utilized in human genetic studies, have allowed a more powerful understanding of the genetic architecture of polygenic traits using genome-wide association studies (GWAS) [6–8]. After conceptual work solidified the importance of polygenic adaptation [9,10], researchers began placing more of an emphasis on identifying this phenomenon using an infinitesimal approach [11–14]. When polygenic adaptation is studied with an infinitesimal approach, the estimates of individual allelic effects on a phenotype are often calculated in a GWAS[12,15,16]. The "GWAS hits" are then used to test for selection [12,15,16]Studies in human genetics usually use summary statistics from meta-analysis GWAS

like the GIANT data set, where many individual GWAS are combined [12,15–19]. These studies observed strong signals for polygenic selection [12,15–19]]. The observed signals did not replicate when the estimates of individual allelic effects were calculated based on GWAS conducted in the unstructured UK Biobank data set, especially when the data set was filtered for individuals with a common ancestry [15,16]. Apparently estimates of allelic effects derived from GWAS are affected by population confounding effects, and when they are combined in polygenic scores, they are highly prone to cumulative bias due to their systematic nature [15,16,19]. Furthermore, Berg et al. (2019) [16] observed that methods like principal component analysis are not properly accounting for population confounding effects.

Zeng et al. (2018) [20] recently developed a Bayesian mixed linear model (BayesS) based on the relationship between SNP effect sizes and minor allele frequency. This approach uses genomic data and fits all SNP effects together as random effects [20]. The identified signals of polygenic selection from Zeng et al. (2018) [20] conflict with the results from Berg et al. (2019) [16], even though both studies worked with the unstructured UK Biobank data set. Berg et al. (2019) [16] filtered the samples based on British ancestry, whereas Zeng et al. (2018) [20] included samples of all European ancestry, which might suggest that residual European population structure continues to be confound, even when advanced methods for corrections for population structure were applied [16]. On the other hand, BayesS used by Zeng et al. (2018) [20] is based on the relationship between SNP effect sizes and minor allele frequency, whereas minor allele frequency and SNP effects might correlate due to estimation error during genomic prediction [4].

The previously mentioned approaches were developed for human genetics, and most of these approaches are based on summary statistics from massive GWAS which are not always feasible for other species, e.g. due to budgetary limitations. Josephs et al. (2019) [21] developed a statistic, referred as conditional $Q_{PC}$, which detects adaptive divergence while controlling for

population substructure. Conditional $Q_{PC}$ detects adaptive divergence based on the excess of

variation observed among populations versus variation observed within populations [21]. The

underlying assumption of this test is that adaptive divergence among populations increases the

trait variance explained by the first principal components (PCs) relative to the later PCs [21].

The PCs are computed based on the relatedness matrix, therefore the conditional $Q_{PC}$ detects

substructures which are associated with trait divergence and based on these observations,

conclusions about polygenic adaptation regarding the trait are drawn [21].

Another test, denoted $\hat{G}$ (or Ghat), has been developed to identify selection on polygenic traits

using whole-genome data [4]. Ghat combines information from all loci simultaneously to

evaluate whether there is a trend for loci contributing to a phenotype to collectively show

evidence of selection [4,22]. Thereby, signals that are individually insignificant for selection

(at the locus level) may become highly significant when evaluated at the genome-wide scale (at

the level of finding selection on a trait). Furthermore, because Ghat uses pre- and post-selection

data, it can identify polygenic selection within shorter sub-periods of time and compare these

to each other [4]. Another advantage of Ghat is that it is based on the relationship between SNP

effects and allele frequency change over time at every locus, which should not correlate unless

there has been on-going selection in comparison to minor allele frequency and SNP effects

[4,20].

We have developed and released an R [22,23] package called "Ghat" to implement the $\hat{G}$ test.

The package allows users to test for selection and evolution in quantitative traits, and as shown

by Beissinger et al. (2018) [4], it is particularly powerful when testing for selection on highly

polygenic traits. To evaluate and demonstrate the package in this manuscript, we applied the

method and corresponding Ghat package to 15 different traits measured in a winter wheat

collection of 191 cultivars, registered mainly in Western Europe between 1966 and 2013, which

were published by Voss-Fels et al. (2019) [24]. We then applied Ghat package on a simulated

cattle population to validate the package and explore different situations where it is most and

least powerful.

## Implementation of the Ghat Package

**Overview**

Ghat identifies selection by leveraging all genotyped loci simultaneously [4]. This makes it possible to identify selection in complex traits that are controlled by many genes with small effects [4]. First, all allelic effects are estimated using a genomic prediction approach. Next, allele frequencies are estimated in two or more different generations (e.g. generation 0 and 10) [4]. Third, the rate of linkage disequilibrium decay is estimated to approximate the number of independent genome segments in the study population [4]. Finally, an LD-aware permutation test is implemented to evaluate if there is evidence of significant selection between generations. The direction of selection is calculated based on the relationship between the change in allele frequency between the generations and the estimated additive effect [4]. Complete theoretical and methodological details are provided in [4].

**Installation**

The Ghat package is available on the Comprehensive R Archival Network (CRAN) [23] and can be installed within the R terminal (Fig1, step-1). The package requests the following optional R-package dependencies:

- Parallel-Package {parallel}: Support for parallel computation, to speed the computation time in large data analysis. [25]
- rrBLUP-Package {rrBLUP}: Ridge regression for estimating marker effects (RR-BLUP). [26]

These maybe installed and loaded before running Ghat by typing:

>install.packages(c("parallel", "rrBLUP")) and then typing >library(parallel) and >library(rrBLUP) (Fig1, step-1).

## Input Data format

For its simplest implementation, Ghat requires only two vectors of input. The first of these is a vector of allele substitution effects, and the second is a vector containing the allele frequency change between the two generations (Table 1) [4]. Required parameters are, "method": to specify how the Ghat permutation test should account for LD between markers ("scale" is usually the most appropriate method to implement because it accounts for linkage disequilibrium between markers, as described below), "perms": for the number of permutations to perform, "plot": to specify whether or not simple Ghat plots should be output, "blocksize": for setting the size of blocks that should be used for LD-trimming (only required when method = "trim"), and "num_eff": for setting the effective number of independent segments of the genome (only required when method = "scale"). All parameter details are described in Table 1.

## Establishing the number of effective markers

Treating each marker as an independent observation can lead to overestimated test statistics. This is because allele frequencies and effect estimates from markers that are in LD are not independent [4] . Any level of LD between markers leads to fewer independent observations than the total number of markers [4]. To avoid such inflated signal of polygenic adaptation, we include an option in Ghat package called "scale", which scales the variance of the permutation test statistic according to the realized extent of LD [4]. More details on the statistical background of scaling method were published by Beissinger et al. (2018) [4] . In the Ghat package, we provide a function called "ld_decay" that can estimate the LD decay across the entire genome and uses this information to approximate the total number of independent genome segments, or "genotype blocks" [4]. The following command runs the ld_decay function >Ld <- ld_decay(gen, map, max_win_snp, max.chr, cores=1, max_r2) (Fig 1, Step-

3)[4]. The output information is used by Ghat to scale the variance of the permutation test

statistic according to the actual number of independent markers (Fig 1, Step-4) [4].

**Running the test**

To run Ghat, estimates of allele substitution effects and allele frequency changes between two

generations (or populations) are required [4]. Next, to perform the Ghat test with this

information, only one line of code is required: >Ghat_res <- Ghat(effects, change, method,

perms, plot, num_eff) (Fig 1, Step-4) [4]. All of the functions, options and data sets available

in Ghat are provided in Table 1

**Fig 1. Four steps to go from raw individual marker data to results from the Ghat test for**

**selection.** An illustrated example using the "Maize_wqs" data available in the Ghat package.

**Step-1**: install the package, load the dependencies and "Maize_wqs" data and estimate allele

substitution effects (e.g., by using rrBLUP [26]). **Step-2**: calculate the differences in allele

frequency between two different generations (cycle 1 and cycle 3). **Step-3**: estimate the decay

of linkage disequilibrium (LD) to calculate the effective number of independent markers. **Step-**

**4**: calculate the Ghat value and p-value from the permutation test.

191    **Table 1**. Objects, data, functions, and options found in Ghat package.

| Function | | Description |
|---|---|---|
| Ghat (Function) | effects | Input vector of allele substitutional effects. |
| | change | Input vector of changes in allele frequency between two different generations (could be positive, negative or zero). |
| | method | One of "vanilla" (assumes complete linkage equilibrium between markers), "trim" (excludes markers to approximate linkage equilibrium some of the extreme values), or "scale" (scales results to reflect underlying levels of linkage LD). |
| | perms | Input number to set the number of permutations to test for significance. |
| | plot | Input option to select which plot you need to be returned: "Ghat", "Cor", or "Both". |
| | blockSize | Input number to set the size of the window on each chromosome for trimming, only required if method = "trim". |
| | num_eff | Input number to set the effective number of independent markers, only required if method = " scale". Can be calculated using "ld_decay" function above. |
| ld_decay (Function) | gen | Input matrix of genotype data. Individuals in rows, genotypes (0, 1, 2) in columns. |
| | map | Input data frame including the name for each marker with a corresponding chromosome number and physical position. |
| | max_win_snp | Input number to set the maximum number of markers allowed per window within a chromosome before estimating the LD. The default value is 2000. |
| | max.chr | Input number to select to which chromosome you want to run the analysis, chromosomes above this number will be excluded from the analysis. |
| | cores | Input number to set the optimal number of cores for using parallelized calculation, since calculation of LD require too much memory. The default value is 1. |
| | max_r2 | Input number to set the threshold of correlation between the neighbourhood variant ($r^2$) to calculate the effective number of independent variants. |
| Maize_wqs (Data object) | Maize_wqs[[1]] | Data frame, including all SNP names (SNP Names), effects on Acid detergent fiber (adf) (effects. adf), and changes in allele frequencies (change) between cycle 1 and cycle 3 of selection. |
| | Maize_wqs[[2]] | Data frame, including plant ID's (taxa), estimated breeding values for Acid detergent fiber (adf), and generation when the phenotyped strains were grown (year). |
| | Maize_wqs[[3]] | Data frame, map file, each line of the Map file describes a single marker and contain three columns. 1: Name (SNP id); 2: Chromosome (Chromosome number); 3: SNP position (in base-pairs). |
| | Maize_wqs[[4]] | Data frame, Genotype (Illumina MaizeSNP50 BeadChip); an Infinium HD assay (Illumina, Inc. San Diego, CA). 10,017 SNP markers (0, 1 and 2) after filtration, distributed across the maize genome [27]. |

**Part I: Testing for selection during 50 years of wheat breeding**

To demonstrate the power and flexibility of the Ghat R package, we used it to test for selection

during the past 50 years of wheat breeding in Western Europe. To achieve this, we leveraged a

large dataset representing 50 years of commercial winter wheat varieties that was previously

published by Voss-Fels and colleagues [24]. Our analysis methods and results are described

below.

**Methods**

**Genotypes and phenotypes.** A data set of 191 cultivars, registered mainly in Western Europe

between 1966 and 2013, was previously published by Voss-Fels and his group[24]. In total, 19

traits were considered and used to evaluate and quantify the phenotypic progress of wheat

breeding during the last five decades. The cultivars in the data set are mainly registered and

bred for the German market [24].

After filtering for data quality, 15 traits remained in the data set. Spikes per sqm, heading,

sedimentation, falling number, and green canopy duration were removed from the analysis

because of missing phenotypic information (>70% of total phenotypes were missing). All traits

were tested in three different environments by Voss-Fels et al. (2019) [24]. The environments

were: 1: A high-intensity nitrogen supply along with best-practices fungicide, insecticide, and

growth regulator applications (HiN/HiF treatment), 2: A high level of nitrogen fertilization with

a fungicide-free treatment (HiN/NoF), and 3: A low level of nitrogen fertilization with a

fungicide-free treatment (LoN/NoF). The evaluation was based on five main groups of traits:

**1- Yield parameters traits**, including grain yield, harvest index, radiation use efficiency and

radiation interception efficiency; **2- Biomass parameter traits**, including above-ground

biomass, kernels per sqm, and thousand-kernel weight. **3- Quality parameter traits**, including

crude protein, protein yield and nitrogen use efficiency. **4- Morphological traits**, including

plant height, green canopy duration and kernel per spike. **5- Disease resistances traits**,

including resistance to powdery mildew and resistance to stripe rust. All cultivars were

genotyped with a 15K SNP Illumina Infinium iSelect genotyping array [28] by Voss-Fels et al.

(2019) [24]. After filtering, 8,710 SNPs passed the genotype quality control criteria [24] and

were included in their subsequent analyses. In our analysis, we re-analyzed the exact set of

SNPs that was published by Voss-Fels et al. (2019) [24], as described above. All estimates of

genetic correlation between the same traits measured in different environments were published

[24].

**Phenotypic progress.** For establish a baseline for progress that we could use to compare against

Ghat results, we re-estimated the phenotypic progress of the 15 wheat traits over 50 years of

selection and adaptation using simple *Pearson* correlation between phenotype and the year of

release [29].

**Ghat.** Allelic effects at every marker locus for the 15 traits were estimated using rrBLUP [26],

and changes in allele frequencies were calculated with R [23,30]. Ghat is calculated as the

summation of the estimated allele frequency change of every SNP multiplied by its effect size,

according to

$$\hat{G} = \sum_{j=1}^{m} \Delta_j \alpha_j \ , \qquad (1)$$

where $\Delta_j$ is the frequency change in locus $j$ between two time-points and $\alpha_j$ is the additive effect

of locus $j$. We implemented two different analyses with Ghat to test for selection, which enabled

us to measure the efficiency of the test in different situations. In the first analysis, which we

call "**All Phenotypes**", allelic effects were estimated using phenotypic information from all

available cultivars in the Voss-Fels et al. (2019)[24] study, which represent the last 50 years of

wheat breeding. This represents the best-case Ghat analysis, in which all phenotypic

information is available. In the second analysis, which we call **"Modern Phenotypes",** allelic

effects were estimated using only modern phenotypic information (from 2010 to 2013), while

changes in allele frequency were still calculated based on genotypes from opposite ends of the

50-year dataset. This represents a realistic use-case for Ghat, when all genotypic information is

available, but phenotypic information is only available for a subset of the individuals (e.g. from

modern germplasm). In other words, this mimics the situation where reliably measured

historical phenotypes do not exist, but historical tissues or seeds are available and can be used

for DNA collection.

**PCA:** Principal component analysis (PCA) was performed with an R [23,31] package called

"adegenet", which was developed for multivariate analysis of genetic markers [31]. The PCA

was conducted with all 191 cultivars with 8,710 markers to detect the presence of population

structure. For this analysis missing data were imputed according to the mean allele frequency

[32].

## Results

### Testing for selection during 50 years of wheat breeding

The main theory behind this test for polygenic selection is that alleles at every locus, distributed across the entire genome, may contribute a small proportion of the total variance for each trait .Therefore, previous studies either required large sample sizes [5] or were unsuccessful in identifying the signals of selection on a genetic level [5]. To demonstrate the Ghat package, we used it to test for selection on the 15 wheat traits, enabling us to establish how much power is lost when phenotypic data are only available from modern lines.

**Phenotypic progress.** After estimating the phenotypic trend (correlation between phenotypic value and the registration year) (Fig 2). We can classify the 15 traits into three groups. The first group is showing comprised of traits that increased in value during breeding. These were above-ground plant biomass, grain yield, green canopy duration, harvest index, kernels per spike, kernels per square meter, nitrogen use efficiency, protein yield, radiation interception efficiency, radiation use efficiency, resistance to powdery mildew, and resistance to stripe rust. Next, we identified a group of traits with a decreasing phenotypic trend, which was comprised of plant height and crude protein. The last group contained traits without a strong overarching phenotypic trend. Only thousand-kernel weight was in this group. The correlation for thousand-kernel weight was positive, but relatively weak in comparison correlations observed in the first group (Fig 2).

**Ghat test for selection.** Fig 2 shows the results of the Ghat test when testing for selection between 1966 and 2013 in winter wheat breeding programs from mainly Western Europe. First, we investigated the precision and accuracy of the Ghat test by estimating the significance, and direction of selection and comparing the results to phenotypic progress over a time frame of 50 years. Overall, the selection size and direction on the 15 traits using all phenotypes were in the

strong agreement with the phenotypic trend. Next, we evaluated the results from Ghat using

phenotypic information from only the most-recent three years of the study (but all genotypic

data were still included). Notably, this was also in a strong agreement with the phenotypic trend.

Ghat successfully detected the signals of selection from the past in each trait, even when only

modern phenotypes were utilized. For example, the results from the Ghat test indicated strong,

directional selection for plant height and crude protein traits, which agrees with their phenotypic

trends during the last 50 years[24]. Ghat showed very weak selection for thousand-kernel

weight, which again aligned with this trait's observed phenotypic trend [24]. The rest of the

traits were under high to moderate directional selection. To validate the results of Ghat on wheat

data, *spearman* correlations between the Ghat results from 3 and 50 years of phenotypes were

estimated. A significant (r=0.57; P<0.001) correlation was found between Ghat values from 3

and 50 years of phenotypes, indicating that even in the absence of historical phenotypic

information, Ghat can be used to understand historical patterns of selection.

**Fig 2. The power of Ghat test of selection in European winter wheat under three agrochemical treatments**. 1: A high-intensity nitrogen supply along with best-practices fungicide, insecticide, and growth regulator applications (HiN/HiF), 2: A high level of nitrogen fertilization with a fungicide-free treatment (HiN/NoF), and 3: A low level of nitrogen fertilization with a fungicide-free treatment (LoN/NoF). **Left**: the phenotypic trend over the last 50 years. **Right**: The Ghat test of selection based on two analyses. "All Phenotypes" refers to estimating allelic effects using all available phenotypic information over the 50 years of wheat breeding; "Modern Phenotypes" refers to estimating allelic effects using modern phenotypic information only (2010 to 2013). Blue, positive selection; red, negative selection. Large light ellipses correspond to weaker selection intensities; thin dark ellipse correspond to more selection intensity. P-values printed on each cell on the left side of the figure (phenotypic trend section) correspond to the significance of the phenotypic trend. P-values printed on each cell on the right side of the figure (Ghat test section) correspond to implementing the Ghat permutation test.

**Assessing population stratification.** Our PCA showed little evidence for population

stratification: no major groupings could be identified (Fig 3). Voss-Fels et al. (2019) [24]

observed the same, and the authors explain that this observation may be caused by a large

germplasm exchange between breeding companies. In the PCA, the first three principal

components (PCs) explain together 16.05% of the variance, which do not correspond to their

geographic location (Fig 3). This analysis was performed because excessive population

stratification can cause erroneous signals of selection to be identified [16]. The minimal

stratification observed in this PCA (Fig 3) and by Voss-Fels et al. (2019) [24] suggests this is

not the case (Fig 3), which could be also explained by the strong exchange between the breeding

programs [24].

**Fig 3. The results from the principal component analysis (PCA)**. A- PC1 plotted against PC2. B-

PC1 plotted against PC3. C- PC2 plotted against PC3. D-The eigenvalues of all the detected PCs.

**Adaptation history of Western European winter wheat.**

To dissect the history of selection in wheat more precisely, we divided the 50 years of breeding into 10 five-year time periods and applied the Ghat test successively for each period. We excluded the green canopy duration trait from this analysis since it was only measured under one environmental condition. For the remaining 14 traits, we calculated the average phenotypic value across the three different environmental conditions. For example, during the period from 1966 to 1975, significantly selected traits according to the Ghat test included grain yield, harvest index (Fig 4-A), above-ground plant biomass, thousand-kernel weight (Fig 4-B), crude protein, nitrogen use efficiency (Fig 4-C), kernel per spike and resistance to stripe rust (Fig 4-D). However, during the same period, selection operated significantly in the negative direction against radiation use efficiency (Fig 4-A), thousand-kernel weight (Fig 4-B), plant height and resistance to powdery mildew (Fig 4-D). Therefore, by considering the single time periods of selection within the past 50 years (Fig 4), and regarding the traits which were selected within the different time periods, we can deduce information about the focus of the breeders during that time period or the environmental conditions during the time period. The number of cultivars available for the 10 adaptation periods are as follows: 18 cultivars from 1966 to 1975; 19 cultivars from 1976 to 1985; 25 cultivars from 1986 to 1995; 48 cultivars from 1996 to 2005; 81 cultivars from 2006 to 2015.

**Fig 4. Ghat correlation estimates of the selection size and direction for 50 years of wheat breeding.**
Results are for 14 productivity traits in winter wheat from Western Europe measured during the past 50
years. Points above zero level (red dashed lines) indicate positive selection, while points under zero
indicate negative selection; bold lines represent 99% confidence interval. A- Adaptation history for the
**Yield parameters traits**. B- Adaptation history for the **Biomass parameters traits**. C- Adaptation
history for **Quality parameters traits**. D- Adaptation history for **Morphological and disease
resistance traits**.

**Part II: The power of Ghat to identify selection in a divergent of simulated dairy cattle population**

To complement the above wheat analysis, we additionally generated a simulated cattle data set with divergent selection to further test and demonstrate the efficacy of Ghat. This allowed us to dissect the Ghat package and how it performs under different genetic architectures and experimental parameters, which are never perfectly known using real data. In this analysis, we assessed the impact of trait heritability ($h^2$), sample size ($n$), trait polygenicity (number of QTL ($nQTL$)), and marker density ($MD$) on the performance of Ghat.

**Data Simulation**

We simulated 30 bovine chromosomes of a length of 100 cM using QMsim [33] . Our simulations started with 100 historical generations of drift-only, and then we sampled two distinct populations to undergo selection. Population-A (unselected population) is a population that underwent complete random mating (i.e. no selection) for 20 generations; Population-B (selected population) is a population that underwent strong phenotypic selection for 20 generations (Fig 5). Aside from selection, Population-A and Population-B were identical and were treated identically during the simulated experiment. After simulating the populations, we applied Ghat to test for selection during the most recent 5 generations (generation #20 – generations #15) in both populations (unselected population and selected populations), creating a realistic situation in dairy cattle breeding. We estimated the allele effects using the genotypes of all 20 generations for more accurate estimation.

381
382 **Fig 5.** QMSim simulation scheme.
383
384
385

Four groups of datasets were simulated for studying the performance of Ghat (Table 2).

Each group involved the creation of different datasets simulated with three constants parameters

and one varying parameter (i.e. the parameter under investigation). Group 1 was simulated to

investigate the effects of $n$ in the Ghat test; we simulated ten datasets with a constant $h^2$, $nQTL$,

and $MD$, but with a different level of $n$ (from 50 to 25600 animals) in each dataset. Then we

measured the P values coming from using these different datasets with different $n$. Group 2 was

simulated to investigate the effect of $h^2$ on Ghat performance. For this, we simulated ten datasets

with a constant $n$, $nQTL$, and $MD$, but with a different $h^2$ (from 0.1 to 0.95) in each dataset.

Group 3 investigated the effect of $nQTl$ on Ghat performance. For group 3, we simulated ten

datasets with a constant $n$, $h2$, and $MD$, but with a different level of nQTL (from 30 to 15360

QTL) in each dataset. Group 4 was simulated to investigate the effect of $MD$ on Ghat

performance. In this group we simulated ten datasets with a constant $n$, $h2$, and $nQTL$ but with

a different level of MD (from 30 to ~ 1 million markers) in each dataset.

**Table 2:** Presentation of all datasets used in the Ghat test.

| Dataset | To investigate | Sample size | Heritability | QTL number | Marker density |
|---------|----------------|-------------|--------------|------------|----------------|
| Group 1 | Sample size | 50 : 25600[a] | 0.5 | 150 | 10020 |
| Group 2 | Heritability | 400 | 0.1 : 0.95[b] | 150 | 10020 |
| Group 3 | QTL number | 400 | 0.5 | 30 : 15360[c] | 10020 |
| Group 4 | Marker density | 400 | 0.5 | 150 | 30 : 1 million[d] |

401    [a]Sample size (n) : 50, 100, 200, 400, 800, 1600, 3200, 6400, 12800 and 25600 animals.
402    [b] Heritability ($h^2$) : 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 and 0.95 of $h^2$.
403    [c] QTL number (nQTL) : 30, 60, 120, 240, 480, 960, 1920, 3840, 7680 and 15360 QTL.
404    [d] Marker density (MD) : 30, 300, 3k, 30k, 60k, 120k, 240k, 480k, 960k and ~1 million markers.
405

**Method and implementation**

Genomic prediction methods like BayesR, BayesB and BayesC assume that the allele

substitution effects follow a distribution in which many effects are zero and some have

moderate effects, subsequently these methods result in higher accuracy than the

GBLUP/rrBLUP methods [34]. Since most of the bovine traits were simulated with a large

number of QTL of small effect, allele substitution effects at every marker locus were estimated

using BayesC [35]. The effective number of independent marker genome segments were

estimated using SimpleM algorithm in R [23,36], and changes in allele frequencies were

calculated with R [22,23]. All R-codes are publicity available in GitHub

(https://github.com/Medhat86/Selection-and-adaptation-on-simulated-bovine-genome).    Ghat

is calculated as the summation of the estimated effect of every SNP scored multiplied by its

effect size, according to model 1. We implemented identical analyses on selected and

unselected populations across 1000 simulation replications for four different tests ($n$, $h^2$, $nQTL$

*and MD*) to evaluate the detection accuracy by evaluating true discovery rates and false

positives. The confidence intervals were obtained from p-values of the same traits from the

Ghat permutation test using the formula from Altman and Bland [37].

**The effect of sample size on Ghat**. Based on 1000 replications, Fig 6 shows the impact

of *n* on the performance of Ghat through quantifying the P values of the Ghat test for selection

(Fig 6). When we test the unselected population with a constant $h^2 = 0.5$, $nQTL = 150$ and

10,020 markers, 99% of replicated traits were not significantly (p-value < 0.01) under selection.

Conversely, in the selected population 99.3% of replicated traits found highly significant under

selection (p-value < 0.01). The p-value also shows a decreasing trend with increasing *n* (from

50 to 25,600 animals) in the selected population. Moreover, the confidence interval decreases

with increasing *n*.

**Fig 6. The effect of *n* on Ghat.** Results are for 1000 replicated traits for testing the effect of *n* on

Ghat-P.values, light blue for unselected population and dark blue for selected population.

**The effect of heritability on Ghat**. Based on 1000 replications, Fig 7 shows the impact

of $h^2$ on the power of Ghat (Fig 7). Higher $h^2$ led to a more powerful test. When we tested the

unselected population with a constant $n = 400$, $nQTL = 150$ and 10,020 markers, 100% of

replicates were not significantly (p-value < 0.01) under selection. In the selected populations

100% of replicates were significant for selection (p-value < 0.01). The p-value shows a

consistence decrease with increasing trait $h^2$ (from 0.1 to 0.95 of $h^2$).

**Fig 7. The effect of $h^2$ on Ghat.** Results depict the effect of $h^2$ on the power of Ghat. Light green

indicates the range of 1000 unselected population replicates and dark green indicates the range of 1000

selected population replicates.

**The effect of the number of QTL controlling a trait *nQTL* on Ghat**. Based on 1000

replicates, Fig 8 shows the impact of *nQTL* on the power of Ghat. When we tested the unselected

population with a constant $n = 400$, $h^2 = 0.5$ and 10,020 markers, 99.7% of replicated traits were

not significantly (p-value $< 0.01$) under selection. In selected population 100% of replicated

traits found highly significant selection (p-value $< 0.01$). The power of the test increased with

increasing the *nQTL* in selected population. This can be restated to emphasize that, all else

being equal, Ghat is more able to detect selection on traits that are more complex (ie controlled

by large number of QTL).

**Fig 8. The effect of *nQTL* in the Ghat-test for selection.** Results are for 1000 replicated traits for

testing the effect of *nQTL* on Ghat-P.values, light red for unselected population and dark red for the

selected population.

**The effect of marker density on Ghat**. Based on 1000 replications, Fig 9 shows the

impact of *MD* on the performance of Ghat through quantifying the p-values of the Ghat-test for

selection (Fig 9). When we tested the unselected population with a constant $n = 400$, $h^2 = 0.5$

and 150 *QTL*, 90% of replicated traits were not significantly (p-value < 0.01) under selection.

While, in selected population 100% of replicated traits were highly significant for selection (p-

value < 0.01). The p-value also shows an inconsistent increase of significance along with an

increasing MD in the selected population (from 30 to ~ 1 million MD). This may be caused by

multicollinearity in the data when there are a high number of markers (>120,000 MD) and

relatively small n (400 animals).

**Fig 9. The effect of *MD* in Ghat-test for selection.** Results are for 1000 replicated simulations testing

the effect of *MD* on the power of Ghat. Gray indicates no-selection simulations and black indicates

selection simulations.

**Testing and system requirements.** We have released three versions of Ghat package

in CRAN (https://cran.r-project.org/package=Ghat) for the three major operating systems: MS

Windows, Linux/Unix, and Mac OS. We have also released the source code on GitHub

(https://github.com/timbeissinger/ComplexSelection) so that users can compile for specific

platforms or modify the code as needed.

For users who have already computed allele effect and allele frequencies changes and

are applying the Ghat test to these data, the computational cost of Ghat is unnoticeable.

Practically, estimating Ghat for maize data [5] with 1000 permutations takes one second per

one phenotype on a 2019 Dell XPS 23 with 2.4 Intel Core i7 processor.

## Future directions

As shown herein, the Ghat package can be powerfully used to detect selection on polygenic traits. Ghat identified polygenic selection on several candidate traits which were actually selected in winter wheat breeding programs. However, the wheat dataset included a relatively small number of markers (8,710) relative to current high-density marker panels and/or whole-genome sequencing approaches, so this analysis did not represent all the challenges associated with implementing Ghat in these settings.

To address this, we used a simulated SNP panel to further explore the utility of Ghat when high LD between markers is present. We found that Bayesian methods, such as BayesC [34,38], as implemented here, can be useful because of its ability to employ feature selection for marker number reduction [38–40]. BayesC does this by assuming that some fraction of the markers have zero effect on the trait of interest and forcing all other effects to equal zero. While Ghat proved to be powerful when paired with BayesC, Ghat is flexible and able to use marker effect estimates extracted from any prediction method. Further research should be conducted to determine the ideal feature-selection approaches to pair with Ghat.

Unlike methods for detecting selected loci, Ghat can quantify the direction of selection even in the absence of phenotypes from a pre-selection population. For breeders and breeding organizations, this represents a practical solution for tracing selection and adaptation history based on phenotypes measured from modern germplasm coupled with DNA samples saved from the past. From economical perspective, this may allow a retrospective evaluation of traits that were inadvertently selected in the past, and therefore may productively serve as the targets of intentional selection in the future. Advances in phenotyping technologies make measuring novel traits increasingly possible [41,42] , and Ghat allows researchers to understand selection on traits that could not be measured previously for practical or technological reasons.

The Ghat test and implementation via the package described herein does not require

immense sample size, individual genotypes and/or genome-wide significant SNPs, as

demonstrated by our implementation with a relatively modest sample size of 191 wheat

individuals genotyped for 8,710 SNPs. Notably, this approach is more powerful when traits are

controlled by large number of QTLs [4] . These advantages allow the Ghat package to be used

for identifying selection for various complex traits in other species of plants, animals, and

humans.

## 524    Data Availability

525    The Ghat R package described in this manuscript is open source and available on CRAN

526    (https://cran.r-project.org/package=Ghat). All analysis and simulation scripts used in this

527    manuscript are available on GitHub (https://github.com/timbeissinger/Maize-Teo-Scripts). The

528    wheat dataset we utilized was previously published and released by [24].

## 530    Acknowledgements

This project was supported with funds from the University of Göttingen, Faculty of Agriculture,

the Center for Integrated Breeding Research, and the Division of Plant Breeding Methodology.

We are indebted to the authors of [24] for making their data publicly available. Two anonymous

reviewers provided suggestions that substantially improved this manuscript and we are grateful

for their input. The authors gratefully acknowledge the hardware support from the High-

Performance Computing team (HPC) at "Gesellschaft für wissenschaftliche Datenverarbeitung

mbH Göttingen" (GWDG).

# References

1.  Fisher RA. XV.—The Correlation between Relatives on the Supposition of Mendelian Inheritance. Earth Environ Sci Trans R Soc Edinb. 1919;52: 399–433. doi:10.1017/S0080456800012163

2.  Wright S. Evolution in Mendelian Populations. Genetics. 1931;16: 97–159.

3.  Ma Y, Ding X, Qanbari S, Weigend S, Zhang Q, Simianer H. Properties of different selection signature statistics and a new strategy for combining them. Heredity. 2015;115: 426–436. doi:10.1038/hdy.2015.42

4.  Beissinger T, Kruppa J, Cavero D, Ha N-T, Erbe M, Simianer H. A Simple Test Identifies Selection on Complex Traits. Genetics. 2018;209: 321–333. doi:10.1534/genetics.118.300857

5.  Lorenz AJ, Beissinger TM, Silva RR, de Leon N. Selection for Silage Yield and Composition Did Not Affect Genomic Diversity Within the Wisconsin Quality Synthetic Maize Population. G3 GenesGenomesGenetics. 2015;5: 541–549. doi:10.1534/g3.114.015263

6.  Wood AR, Esko T, Yang J, Vedantam S, Pers TH, Gustafsson S, et al. Defining the role of common variation in the genomic and biological architecture of adult human height. Nat Genet. 2014;46: 1173–1186. doi:10.1038/ng.3097

7.  Visscher PM, Hemani G, Vinkhuyzen AAE, Chen G-B, Lee SH, Wray NR, et al. Statistical Power to Detect Genetic (Co)Variance of Complex Traits Using SNP Data in Unrelated Samples. PLOS Genet. 2014;10: e1004269. doi:10.1371/journal.pgen.1004269

8.  Visscher PM, Wray NR, Zhang Q, Sklar P, McCarthy MI, Brown MA, et al. 10 Years of GWAS Discovery: Biology, Function, and Translation. Am J Hum Genet. 2017;101: 5–22. doi:10.1016/j.ajhg.2017.06.005

9.  Pritchard JK, Di Rienzo A. Adaptation – not by sweeps alone. Nat Rev Genet. 2010;11: 665–667. doi:10.1038/nrg2880

10. Pritchard JK, Pickrell JK, Coop G. The Genetics of Human Adaptation: Hard Sweeps, Soft Sweeps, and Polygenic Adaptation. Curr Biol. 2010;20: R208–R215. doi:10.1016/j.cub.2009.11.055

11. Hancock AM, Alkorta-Aranburu G, Witonsky DB, Di Rienzo A. Adaptations to new environments in humans: the role of subtle allele frequency shifts. Philos Trans R Soc B Biol Sci. 2010;365: 2459–2468. doi:10.1098/rstb.2010.0032

12. Berg JJ, Coop G. A Population Genetic Signal of Polygenic Adaptation. W. Feldman M, editor. PLoS Genet. 2014;10: e1004412. doi:10.1371/journal.pgen.1004412

13. Field Y, Boyle EA, Telis N, Gao Z, Gaulton KJ, Golan D, et al. Detection of human adaptation during the past 2000 years. Science. 2016;354: 760–764. doi:10.1126/science.aag0776

14. Csilléry K, Rodríguez-Verdugo A, Rellstab C, Guillaume F. Detecting the genomic
signal of polygenic adaptation and the role of epistasis in evolution. Mol Ecol. 2018;27:
606–612. doi:10.1111/mec.14499

15. Sohail M, Maier RM, Ganna A, Bloemendal A, Martin AR, Turchin MC, et al.
Polygenic adaptation on height is overestimated due to uncorrected stratification in
genome-wide association studies. eLife. 2019;8: e39702. doi:10.7554/eLife.39702

16. Berg JJ, Harpak A, Sinnott-Armstrong N, Joergensen AM, Mostafavi H, Field Y, et al.
Reduced signal for polygenic adaptation of height in UK Biobank. eLife. 2019;8:
e39725. doi:10.7554/eLife.39725

17. Lloyd-Jones LR, Zeng J, Sidorenko J, Yengo L, Moser G, Kemper KE, et al. Improved
polygenic prediction by Bayesian multiple regression on summary statistics. Nat
Commun. 2019;10: 5086. doi:10.1038/s41467-019-12653-0

18. Turchin MC, Chiang CW, Palmer CD, Sankararaman S, Reich D, Hirschhorn JN.
Evidence of widespread selection on standing variation in Europe at height-associated
SNPs. Nat Genet. 2012;44: 1015–1019. doi:10.1038/ng.2368

19. Racimo F, Berg JJ, Pickrell JK. Detecting Polygenic Adaptation in Admixture Graphs.
Genetics. 2018;208: 1565–1584. doi:10.1534/genetics.117.300489

20. Zeng J, de Vlaming R, Wu Y, Robinson MR, Lloyd-Jones LR, Yengo L, et al.
Signatures of negative selection in the genetic architecture of human complex traits. Nat
Genet. 2018;50: 746–753. doi:10.1038/s41588-018-0101-4

21. Josephs EB, Berg JJ, Ross-Ibarra J, Coop G. Detecting Adaptive Differentiation in
Structured Populations with Genomic Data and Common Gardens. Genetics. 2019;211:
989–1004. doi:10.1534/genetics.118.301786

22. Beissinger, Tim. ComplexSelection. 2019. Available:
https://github.com/timbeissinger/ComplexSelection

23. R: The R Project for Statistical Computing. [cited 2 Jan 2020]. Available: https://www.r-
project.org/

24. Voss-Fels KP, Stahl A, Wittkop B, Lichthardt C, Nagler S, Rose T, et al. Breeding
improves wheat productivity under contrasting agrochemical input levels. Nat Plants.
2019;5: 706–714. doi:10.1038/s41477-019-0445-5

25. Ooi H, Corporation M, Weston S, Tenenbaum D. doParallel: Foreach Parallel Adaptor
for the "parallel" Package. 2019. Available: https://CRAN.R-
project.org/package=doParallel

26. Endelman JB. Ridge Regression and Other Kernels for Genomic Selection with R
Package rrBLUP. Plant Genome. 2011;4: 250–255.
doi:10.3835/plantgenome2011.08.0024

27. Ganal MW, Durstewitz G, Polley A, Bérard A, Buckler ES, Charcosset A, et al. A Large
Maize (Zea mays L.) SNP Genotyping Array: Development and Germplasm
Genotyping, and Genetic Mapping to Compare with the B73 Reference Genome. Lukens
L, editor. PLoS ONE. 2011;6: e28334. doi:10.1371/journal.pone.0028334

28. Muqaddasi QH, Brassac J, Börner A, Pillen K, Röder MS. Genetic Architecture of Anther Extrusion in Spring and Winter Wheat. Front Plant Sci. 2017;8: 754. doi:10.3389/fpls.2017.00754

29. Benesty J, Chen J, Huang Y, Cohen I. Pearson Correlation Coefficient. Noise Reduction in Speech Processing. Berlin, Heidelberg: Springer Berlin Heidelberg; 2009. pp. 1–4. doi:10.1007/978-3-642-00296-0_5

31. Jombart T. adegenet: a R package for the multivariate analysis of genetic markers. Bioinformatics. 2008;24: 1403–1405. doi:10.1093/bioinformatics/btn129

32. Jombart T, Devillard S, Balloux F. Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. BMC Genet. 2010;11: 94. doi:10.1186/1471-2156-11-94

33. Sargolzaei M, Schenkel FS. QMSim: a large-scale genome simulator for livestock. Bioinformatics. 2009;25: 680–681. doi:10.1093/bioinformatics/btp045

34. Meuwissen T, Hayes B, Goddard M. Accelerating Improvement of Livestock with Genomic Selection. Annu Rev Anim Biosci. 2013;1: 221–237. doi:10.1146/annurev-animal-031412-103705

35. Pérez P, de los Campos G. Genome-wide regression and prediction with the BGLR statistical package. Genetics. 2014;198: 483–495. doi:10.1534/genetics.114.164442

36. Gao X, Starmer J, Martin ER. A multiple testing correction method for genetic association studies using correlated single nucleotide polymorphisms. Genet Epidemiol. 2008;32: 361–369. doi:10.1002/gepi.20310

37. Bland MJ, Altman DG. STATISTICAL METHODS FOR ASSESSING AGREEMENT BETWEEN TWO METHODS OF CLINICAL MEASUREMENT. The Lancet. 1986;327: 307–310. doi:10.1016/S0140-6736(86)90837-8

38. Habier D, Fernando RL, Kizilkaya K, Garrick DJ. Extension of the bayesian alphabet for genomic selection. BMC Bioinformfatics. 2011;12: 186. doi:10.1186/1471-2105-12-186

39. Habier D, Fernando RL, Dekkers JCM. The Impact of Genetic Relationship Information on Genome-Assisted Breeding Values. Genetics. 2007;177: 2389–2397. doi:10.1534/genetics.107.081190

40. Habier D, Tetens J, Seefried F-R, Lichtner P, Thaller G. The impact of genetic relationship information on genomic breeding values in German Holstein cattle. Genet Sel Evol. 2010;42: 5. doi:10.1186/1297-9686-42-5

41. Cobb JN, DeClerck G, Greenberg A, Clark R, McCouch S. Next-generation phenotyping: requirements and strategies for enhancing our understanding of genotype–phenotype relationships and its relevance to crop improvement. Theor Appl Genet. 2013;126: 867–887. doi:10.1007/s00122-013-2066-0

42. Mahmoud M, Yin T, Brügemann K, König S. Phenotypic, genetic, and single nucleotide polymorphism marker associations between calf diseases and subsequent performance and disease occurrences of first-lactation German Holstein cows. J Dairy Sci. 2017;100: 2017–2031. doi:10.3168/jds.2016-11767

659