Babesia bovis Rad51 ortholog influences switching of *ves* genes but is not essential for segmental gene conversion in antigenic variation

Erin A. Mack[1]#¶, Massimiliano S. Tagliamonte[2,3]#, Yu-Ping Xiao[1], Samantha Quesada[1], and David R. Allred[1,2,3,4]*

[1]Department of Infectious Diseases and Immunology, University of Florida;

[2]Department of Pathology, Immunology, and Laboratory Medicine, University of Florida; [3]Emerging Pathogens Institute, University of Florida; [4]Genetics Institute, University of Florida, Gainesville, Florida, United States of America

*Corresponding author
email: allredd@ufl.edu (DRA)

#These individuals contributed equally to this project.

¶Current address: Department of Biological Sciences, University of Idaho, Moscow, Idaho, United States of America.

short title: Rad51 and antigenic variation in *Babesia*

Abstract

The tick-borne apicomplexan parasite, *Babesia bovis*, a highly persistent bovine pathogen, expresses VESA1 proteins on the infected erythrocyte surface to mediate cytoadhesion. The cytoadhesion ligand, VESA1, which protects the parasite from splenic passage, is itself protected from a host immune response by rapid antigenic variation. *B. bovis* relies upon segmental gene conversion (SGC) as a major mechanism to vary VESA1 structure. Gene conversion has been considered a form of homologous recombination (HR), a process for which Rad51 proteins are considered pivotal components. This makes BbRad51 a choice target for development of inhibitors that could both interfere with parasite genome integrity and disrupt HR-dependent antigenic variation. Previously, we knocked out the Bb*rad51* gene from the *B. bovis* haploid genome, resulting in a phenotype of sensitivity to methylmethane sulfonate (MMS) and apparent loss of HR-dependent integration of exogenous DNA. In a further characterization of BbRad51, we demonstrate here a failure to upregulate the Bbrad51 gene in response to DNA damage. Moreover, we demonstrate that ΔBb*rad51* parasites are not more sensitive than wild-type to DNA damage induced by γ-irradiation, and repair their genome with similar kinetics. To assess the need for BbRad51 in SGC, RT-PCR was used to observe alterations to a highly variant region of *ves1*α transcripts over time. Mapping of these amplicons to the genome revealed a significant reduction of in situ transcriptional switching (isTS) among *ves* loci, but not cessation. By combining existing pipelines for analysis of the amplicons, we demonstrate that SGC continues unabated in ΔBb*rad51* parasites, albeit at an overall reduced rate, and a reduction in SGC tract lengths was observed. By contrast, no differences were observed in the lengths of homologous sequences at which recombination occurred. These results indicate that, whereas BbRad51 is not essential to babesial antigenic variation, it influences epigenetic control of *ves* loci, and its absence significantly reduces successful variation. These results necessitate a reconsideration of the likely enzymatic mechanism(s) underlying SGC and suggest the existence of additional targets for development of small molecule inhibitors.

Author summary

B. bovis establishes highly persistent infections in cattle, in part by using cytoadhesion

to avoid passage through the spleen. While protective, a host antibody response targeting the cytoadhesion ligand is quickly rendered ineffective by antigenic variation. In *B. bovis*, antigenic variation relies heavily upon segmental gene conversion (SGC), presumed to be a form of homologous recombination (HR), to generate variants. As Rad51 is generally considered essential to HR, we investigated its contribution to SGC. While diminishing the parasite's capacity for HR-dependent integration of exogenous DNA, the loss of BbRad51 did not affect the parasite's sensitivity to ionizing radiation, overall genome stability, or competence for SGC. Instead, loss of BbRad51 diminished the extent of in situ transcriptional switching (isTS) among *ves* gene loci, the accumulation of SGC recombinants, and the mean lengths of SGC sequence tracts. Given the overall reductions in VESA1 variability, compromise of the parasite's capacity for in vivo persistence is predicted.

Introduction

The apicomplexan parasite, *Babesia bovis*, is a quick-change artist with the ability to rapidly alter proteins it expresses on the infected erythrocyte surface. This ability is needed because, during asexual reproduction *B. bovis* cytoadheres to the capillary and post-capillary venous endothelium within its bovine mammalian host [1-3]. It is thought that this behavior allows this microaerophilic parasite to avoid splenic clearance and to complete asexual development under hypoxic conditions, analogous to the human malarial parasite, *Plasmodium falciparum* [4]. Cytoadhesion is mediated by heterodimeric variant erythrocyte surface antigen-1 (VESA1) proteins, which are exported by the parasites to the erythrocyte where they integrate into the erythrocyte membrane [5, 6]. Cytoadhesion is compromised by a host antibody response targeting VESA1, preventing and potentially reversing cytoadhesion [6]. However, *B. bovis* has evolved the ability to rapidly vary the structure and antigenicity of VESA1 polypeptides, abrogating recognition by existing antibodies [7-10]. Antigenic variation in *B. bovis* involves the *ves* multigene family encoding VESA1a and 1b polypeptides [9, 11, 12], and possibly the *smorf* multigene family [13, 14]. *B. bovis* intraerythrocytic stages reproduce asexually, with a haploid genome of only 8-8.5 Mbp [13]. Despite its small genome size, approximately 135 genes comprise the *B. bovis ves* multigene family, amounting to approximately 4.7% of all coding sequences [10, 13]. Transcription of *ves* genes is monoparalogous, arising from a single *ves* locus at any one time (but typically involves transcription of both a *ves*1α and *ves*1β gene from the same locus to encode both subunits), whereas the remainder of the family remains transcriptionally inactive [15]. In situ transcriptional switching (isTS) from one *ves* locus to another over time has been implicated in *B. bovis* antigenic variation [10, 15, 16], although segmental gene conversion (SGC) is the only mechanism of variation critically demonstrated in this parasite to date [12]. Progressive replacement of short sequence patches within the actively transcribed *ves* genes by SGC, yields *ves* genes (and VESA1 polypeptides) that are mosaics comprised of sequences from many *ves* loci [11, 12]. The short lengths of the SGC conversion tracts, ability to acquire sequences from any chromosome, and involvement of two similarly variant subunits may enable this gene family to provide practically unlimited diversity in epitope structure [5, 7, 9, 11, 12].

Canonical gene conversion is a form of homologous recombination (HR)-mediated DNA

repair. In this process a damaged sequence is repaired by incorporating duplicated

homologous sequences from an undamaged allele or paralog to replace the damaged

sequences. At least three models have been proposed to explain this process (reviewed

in [17]). Common to all models is the assembly of a repair complex at the site of damage.

Activities of the repair complex include 5' to 3' resection of the broken ends of the

damaged molecule, and stabilization of the single-stranded 3' ends by assembly of RPA

ssDNA-binding proteins onto the strands (reviewed in [17, 18]). Rad52 may stabilize the

RPA and maintain spatial proximity of the broken ends [19]. The Rad52-RPA complexes

then are replaced by Rad51 protein, which forms helical filaments on the ssDNA

strands. The Rad51-ssDNA complexes, together with Rad54, mediate both a search for

homologous sequences elsewhere in the genome and strand invasion when such

sequences are found [20-22]. Once found, single-stranded invasion of the identified

sequence allows sequence acquisition by extension of the 3' end of the invading strand.

Depending upon the model, the inter-chromosomal entanglement involves either one or

two Holliday junction structures that are resolved to yield two independent

chromosomes again. Sequence acquisition may occur with or without crossover,

depending upon how the junction structures are resolved [23]. Like canonical gene

conversion, we hypothesized SGC to occur via HR, a possibility consistent with the

stretches of homologous sequence flanking SGC tracts that are shared between donor

and recipient. However, other factors call this into question. For example, the reasons

for consistently short conversion tracts are unknown, and crossover appears to be a rare

outcome in *B. bovis* [10]. These traits suggest that minimal end resection, acquisition of

only short tracts of differing sequence, and rapid resolution of intermolecular junction

structures all may define this process mechanistically. Among the existing models of

HR, those most consistent with these traits are synthesis-dependent strand annealing,

either with rapid disentanglement of the invading strand(s), or double-strand break

repair but with convergent branch migration and junction dissolution. By contrast,

resolution of distal Holliday junctions would be expected to result in frequent crossover

events [17]. Alternatively, a more exotic explanation may hold, such as the involvement

of a template-switching repair polymerase, but no direct evidence yet supports this

possibility.

Rad51 is considered essential to HR and the gene conversion process in eukaryotes.

Organisms with defective Rad51 consistently suffer reduced viability and enhanced

sensitivity to environmental insult [24-29]. For example, knockout of the *Rad51* gene in

mice resulted in embryonic lethality [30], whereas loss of the Tb*rad51* gene in the

kinetoplastid parasite, *Trypanosoma brucei*, yielded parasites that were compromised

in growth and hypersensitive to methyl methanesulfonate (MMS) [28]. Interestingly,

Tb*rad51* knockout parasites continued to undergo variation of their variant surface

glycoprotein (*VSG*) genes, both by isTS and gene conversion mechanisms, but the rate at

which variation occurred was slowed dramatically. This observation suggested a role for

TbRad51 in facilitating or regulating trypanosomal antigenic variation, but not an

essential role in catalysis. Moreover, TbRad51-independent mechanisms may act in

trypanosomal antigenic variation. Recently, we knocked out the Bb*rad51* gene of *B.*

bovis. Unlike higher eukaryotes, there was no apparent effect on parasite viability or

growth. However, parasites were made hypersensitive to MMS and failed to integrate

exogenous DNA, suggesting defects in HR [31]. Given the importance of SGC to *B. bovis*

antigenic variation and survival, and of Rad51 to gene conversion and HR, we

investigated further the interplay between DNA repair and antigenic variation. We

provide evidence that overall DNA repair remains highly robust in the absence of

BbRad51. SGC also continues, albeit at an overall reduced rate, concomitant with a

significant reduction in isTS. We hypothesize that these results reflect unique roles for

BbRad51 in antigenic variation, and suggest that alternative enzymes catalyze

recombination during SGC.

Results

Bbrad51 knockout parasites behave similarly to wild-type in response to

DNA damage. Previously, we knocked out the Bb*rad51* gene, resulting in a parasite

phenotype of MMS-sensitivity and apparent loss of the ability to integrate exogenous

DNA by HR, but not significant difference in rates of growth [31]. Cells that suffer

environmental insult may up-regulate DNA repair proteins, including Rad51 [32]. To

test this in *B. bovis* we needed to create a reporter parasite line with which we could

easily quantify BbRad51 protein levels. This was done by integrating the linearized

plasmid, pBRNHLg, into the genome by homologous double-crossover, resulting in

Bb*rad51* coding sequences fused in-frame with NanoLuciferase and dual-HA tag

sequences. Immediately following the stop codon, the *E. coli glmS* ribozyme element, a

self-cleaving autoregulatory sequence controlling glucosamine-6-phosphate synthase

expression [33], was also incorporated. Upon double crossover integration, a *gfp-bsd*

selectable marker cassette also becomes incorporated downstream of the gene in this

approach [34]. This combination allowed us to conveniently measure levels of BbRad51

protein via the activity of the NanoLuciferase fusion partner, and to inducibly knock

down expression via the *glmS* ribozyme element by the addition of the D-glucosamine

cofactor [33], an approach used previously in *Plasmodium* [35]. As a negative control

for induction, samples of *B. bovis* CE11 parasites were transiently transfected with the

plasmid pHggb, in which NanoLuciferase-2xHA expression is driven by the *Babesia*

divergens EF1α-B promoter. This promoter was chosen because it would be unlikely to

respond to upregulation of DNA repair proteins. Structures of these plasmids are shown

in S1 Figure, and sequences are provided in S2_File and S3_File. Reporter parasites

were exposed to 250 μM MMS, a level that reproducibly inhibits growth by

approximately 10% after 48 hours in wild-type *B. bovis* CE11 parasites [31], presumably

sufficient to induce Bb*rad51* expression in response but low enough not to significantly

reduce viability. Under these conditions, no evidence of an alteration in BbRad51

expression was observed after either 3 hours (Figure 1A) or 16 hours of post-treatment

recovery (Figure 1B). Treatment with glcN resulted in significant knockdown of

NanoLuciferase levels, on the order of 80-95%. No upregulation of BbRad51 was

observed if knockdown of existing BbRad51 via the *glmS* element was performed. This

was done to allow more sensitive detection in the case of post-translational stabilization,

or to prevent any possible negative feedback (Figure 1). This result indicates that

BbRad51 levels are not regulated in response to repair of alkylated DNA following acute,

sublethal exposure to MMS.

A second measure of phenotype commonly used in studies of DNA repair is sensitivity to

γ-irradiation. Sensitivity of the knockout parasites to DNA damage was assessed by

exposure to γ-irradiation provided by a Cs[137] source. Wild type CE11 parasites first were

titrated for sensitivity, over a range from 10 - 1230 gray (Gy). A dosage of 100 Gy

resulted in approximately 80-90% lethality by 24 hours, but allowed some parasite
survival at 48 hours and beyond, whereas at 200 Gy no parasites were observed to
survive 48 hours post-irradiation (S2 Figure). Radiation sensitivities of three
independently-derived Bb*rad51* knockout lines (all on a CE11 genetic background)
therefore was compared with CE11 wild-type parasites over a range from 0-200 Gy.
There were no reproducible differences among any of the three knockout lines and CE11
wild type parasites in growth assays (Figure 2).
202
Because no differential survival phenotype was apparent following DNA damage by γ-
irradiation, we asked whether there were any detectable differences in overall rates of
DNA repair. To assess this, pulsed-field gel electrophoresis (PFGE) was used to monitor
the disintegration of chromosomes and their subsequent reassembly. Parasites were
subjected to 100 Gy γ-irradiation, allowed to recover for up to 24 hours, and then were
processed for PFGE analysis. In growth experiments, 100 Gy had resulted in killing of
80-90% of the parasites. Consistent with this level of killing, 100 Gy γ-irradiation
severely damaged *B. bovis* chromosomes, virtually eliminating full-length chromosomes
3 and 4, and greatly diminishing the proportion of intact chromosomes 1 and 2.
Remarkably, by 24 hours post-irradiation both wild-type and knockout parasites had
reassembled their genomes into full-length chromosomes of apparently normal size,
recovering approximately 50% of non-irradiated control values (Figure 3). Given the 10-
20% viability of parasites receiving this dosage (S2 Figure) it is likely that a large
proportion is not viable in the longer-term, but remains metabolically active long
enough to reassemble chromosomes. These data, when considered together,
demonstrate that the loss of BbRad51 has little, if any, effect on the extensive DNA
repair required to recover from such damage and suggest that BbRad51-dependent HR
plays little role in this type of repair.
221

Bbrad51 knockout did not prevent in situ transcriptional switching.
Although overall DNA repair following damage from ionizing radiation was not
measurably impaired by the loss of BbRad51, sensitivity to MMS and loss of ability to
integrate selectable plasmids via long sequence tracts suggested that BbRad51 plays a
role in aspects of HR [31], and perhaps in repair of stalled replication forks [36].

Previously, it was demonstrated that SGC is a major mechanism of antigenic variation in

B. bovis [12]. Given the seemingly conflicting outcomes obtained with MMS and

ionizing radiation, we wished to determine whether BbRad51 plays any role in SGC. In

order to assess the nature of any changes occurring in transcribed *ves1α* genes of wild

type and Bb*rad51* knockout parasites, we adapted a previously published assay in which

the highly variant cysteine-lysine-rich domain (CKRD) region of *ves1α* transcripts

(Figure 4A) is amplified by RT-PCR, and the amplicons undergo deep sequencing [15].

Three immediate subclones of CE11 wild-type and three independent knockout clonal

lines were studied; their origin is described in [31]. Total RNAs were collected from each

clonal line at one month and five months post-cloning. These two timepoints were used

to observe for increases in the numbers of unique recombinants over time, and for some

analyses were pooled to minimize the loss of unique variants from the population over

time. The forward primer, *ves*UniF2, was selected because it represents a sequence

almost universally conserved among *ves1α* genes, and in combination with the highly-

conserved primer PD1R was anticipated to generate amplicons of approximately 340-

460 bp. By constructing bar-coded paired-end libraries from the amplicons and

generating 250 bp reads, reads could be merged with high-confidence overlaps of 55 -

185 bp. Merged, full-length sequences were obtained from 69.8 - 84.5% of amplicon

reads. Following the removal of ambiguous and low-quality reads, adaptor and primer

sequences, and sequences found likely to be PCR chimeras, individual libraries ranged

from a minimum of 853,688 to a maximum of 1,738,649 merged reads. Mean merged

read lengths ranged from 326.2 ± 60.3 to 334.7 ± 52.9 bp. To determine their probable

loci of origin, reads were mapped (using non-global settings) onto the *B. bovis* C9.1 line

genomic sequence (available at Wellcome Trust Sanger Institute;

ftp://ftp.sanger.ac.uk/pub/pathogens/Babesia/). The C9.1 line genome was used

because we do not currently have a high quality genome for the CE11 line. However, as

these are closely-related clonal sibling lines [12] this allowed us to easily identify the

probable locus of origin for nearly all reads. In all six knockout and wild-type lines, the

earlier time-point *ves1α* transcripts mapped predominantly, sometimes almost solely, to

a single locus (Figure 4B), consistent with prior observations of monoparalogous *ves*

gene transcription in the C9.1 clonal line [15]. At the latter time-point, lines ko1[H5] and

CE11[B8] continued to transcribe almost solely from the original locus, and all lines still

transcribed most heavily from the original locus. Minor but significant subpopulations

were detected in all lines that had switched to transcription from alternative loci.

Without immune pressure there is no obvious selection for parasites expressing specific

VESA1a isoforms to predominate. Regardless, detectable transcription occurred from

more alternative loci in wild type CE11 subclones B8, C2, and C5 (ranging from 28.7 ±

4.0 at 1 month to 33.3 ± 1.5 loci at 5 months) than were observed for the three knockout

lines, which ranged from 22.0 ± 3.5 at 1 month to 23.3 ± 1.5 at 5 months (Figure 4C).

While transcribing most heavily from a single locus, the CE11 C5 subclone (as a

population) also transcribed significantly from several alternative loci at the early time-

point, but by the 5-month time-point transcription levels had been reduced from all but

the single, major locus. Interestingly, the same alternative loci seemed to dominate as

sites to which switching occurred (S3 Figure), suggesting a hierarchy in locus

transcription. However, nothing can be inferred from these data regarding an order in

switching like that documented for *P. falciparum var* genes [37].

Bbrad51 knockout failed to prevent segmental gene conversion. The ideal

situation for accurate identification and characterization of SGC tracts would be to

directly map amplicon sequences against a reference genome. Although the CE11 and

C9.1 lines are closely related sibling clonal progeny of the MO7 clonal line, they have

quite different histories [12]. Given the nature and rapidity of SGC many loci would have

been extensively modified. The ideal would be to map to a high quality genome from

each subclone, so that unique variants in each line would be known, but this was not

feasible. As a more feasible alternative, we chose to observe for recombination among

loci represented by the *ves1α* transcript amplicons, considering only sequences unique

to a given line, on a line by line basis. By taking this approach, we could assess for

recombination among sequences that were definitively present in each line at the time

the experiment was performed. Using an RT-PCR strategy employed previously to

characterize variation in *ves1α* transcripts [15], we identified all unique transcribed

sequences by cluster analysis, then identified representative reads for each cluster.

Among those, we then identified sequences for which there was very strong statistical

support for them being the result of a true recombination event between two other

unique sequences, based upon a consensus of 4 out of 7 statistical analyses (see Methods

for details). For comparative analysis, we included only those recombinant sequences
found in a single clonal line, on the assumption that any sequences found in more than
one line was present prior to the act of parasite cloning and not a result of post-cloning
recombination. Information on all identified SGC tracts are provided in S1_File.zip. We
propose that most of the unique sequences were in fact recombinant, as a plot of all
unique sequences against all statistically-supported recombinant sequences (both
normalized per million reads) a regression of R^2 = 0.9228 was obtained. Thus, the use of
a statistical consensus approach was highly conservative and likely underestimates the
true number of recombinant sequences (S4 Figure), but allows for rigorous comparative
results.
301
Variants with strong statistical support as true recombinant SGC tracts were observed
among the transcripts of all six clonal lines, but clear distinctions are seen between
knockout and wild type. The mean lengths of conversion tracts differed between groups,
decreasing from means of 109.01 ± 39.39 in wild type parasites to 91.77 ± 40.78 in
knockouts (p <0.001; Figure 5). The distributions of SGC tract lengths, plotted as
cumulative proportions of all SGC tracts from that population, resulted in wild type and
knockout medians of 107.0 and 81.0 (S5 Figure; p <0.001). Interestingly, the difference
between the two populations arose primarily at tract lengths <150 bp. The two
populations were not distinct above that length, suggesting the possibility that more
than one mechanism gave rise to the SGC tracts. The number of unique SGC tracts
arising per active *ves* locus was not significantly different, ranging from 0.35- 1.20 SGC
tracts per locus (S6 Figure; p = 0.22 among all group comparisons). Although there was
a rise in the frequency of recombinants per locus in CE11 wt parasites over time this was
not statistically significant (p = 0.18). However, given the differences in the numbers of
transcriptionally active *ves* loci in wild type and knockout parasites, this led to
significantly larger total numbers of unique SGC tracts per million reads among
members of the wild type population (Figure 6; at 5 months, p <0.02). Thus, while the
frequency of SGC alterations that may be observed at any given transcriptionally active
ves locus is approximately constant, the numbers of *ves* loci that are activated is
significantly higher in the presence of BbRad51. The number of unique SGC tracts per
million reads may be considered a surrogate measure of overall levels of variability in

antigen structure presented by the population of parasites. Taken together, these data

demonstrate clearly that neither SGC nor apparent isTS to alternative loci is abrogated

by knockout of the Bb*rad51* gene, although statistically significant quantitative effects

on the lengths of SGC tracts and on the frequency and extent of *ves* locus switching were

observed.

Involvement of homologous sequences flanking SGC breakpoints. We

wished to assess whether there is anything shared at SGC breakpoint sites, or unique

where SGC tracts differed from the active locus of *ves* transcription (LAT). To do this,

alignments were made of each pair of sequences identified as having given rise to a

unique recombinant, and the recombinant sequence itself. Regions of homology

between all three sequences were then identified manually that represented a region in

which transition occurred from one parent locus providing the sequence to the other (S7

Figure).. No significant differences were observed among wild type or Bb*rad51* knockout

genotype parasites in the lengths of homology patches (Figure 7). While this is clearly

not an exhaustive analysis of all SGC tracts, the results of this subsampling indicate that

there is no apparent difference between wild type and Bb*rad51* knockout parasites with

regard to the lengths of the homologous patches possibly involved in recombination.

Importantly, for SGC to occur patches of homology between donor and recipient may

not be required, as patches of as little as 2 bp were observed. In a few instances no

transition region was present, and in still others there was a brief patch in which no

match existed between the three sequences at the site of transition, suggesting a

sometimes chaotic process (S7 Figure).

Discussion

SGC is a major mechanism of antigenic variation in *B. bovis*, and to date the only one

that has been demonstrated critically [12]. This phenomenon, in which short DNA

patches are duplicated from a donor to a recipient gene, typically occurs without

modification of the donor and at least superficially resembles HR-mediated DNA repair.

DNA repair in apicomplexan and other protozoal parasites overall is not well

understood [38, 39]. It is even difficult to predict from the parasites' proteomes which

repair pathways may be active. Orthologs are recognizable for only a fraction of the

proteins known to be important in higher eukaryotes, and many orthologs are simply not present. For example, the proteins RPA, Rad51, Rad52, Rad54, and ATM are considered key participants in DNA repair pathways, including HR and gene conversion. Yet, in *B. bovis* orthologs may be identified only for RPA, Rad51, and Rad54, and many similar examples of "missing" proteins hold [13, 31], suggesting the merging of functions. In this study, we wished to understand the contributions of BbRad51 to SGC because proteins of this family are considered essential to HR and gene conversion in other systems, including other apicomplexans [29, 40]. Our prior identification of BbRad51 as the true Rad51 ortholog was based upon several criteria, including sequence and structural similarities to established Rad51 proteins, a greatly reduced or abrogated ability to achieve HR-dependent integration of exogenous sequences in the absence of BbRad51, and enhanced sensitivity to MMS in knockouts that could be complemented by Bb*rad51* coding sequences [31].

In contrast with our prior work, we present evidence herein that the absence of BbRad51 does not influence the parasite's survival or extent and rate of general repair of dsDNA breaks engendered by acute exposure to γ-irradiation, and that BbRad51 expression fails to up-regulate in response to MMS-induced damage. The apparent insignificance of BbRad51 to repair and survival of IR-induced DSBs may be attributable to the haploid nature of the asexual stages studied here. A large proportion of IR-caused DSBs would occur in unique regions of the genome, with no intact second copy of the damaged sequence available to support true gene conversion, except briefly during mitosis. Thus, in the absence of available sequence donors for HR, BbRad51 may be largely superfluous to surviving heavy dosages of IR, yielding similar ionizing radiation survival outcomes in wild-type and knockout parasites. *B. bovis* may depend instead upon error-prone end-joining reactions for survival of such significant damage. Unlike most eukaryotes, including *Toxoplasma* [41], *B. bovis* lacks genes for key players in canonical non-homologous end-joining repair, such as Ku70/80 and DNA ligase 4 ([13]; this study). This parasite instead may depend upon a synthesis-dependent microhomology-mediated end-joining mechanism like that demonstrated in *P. falciparum* [42], but this remains to be determined. In contrast with ionizing radiation, the absence of BbRad51 does render *B. bovis* sensitive to alkylation damage caused by acute exposure to MMS

[31]. In diploid organisms recovery from either type of insult is typically compromised
by loss of Rad51 [43]. However, in diploids such repair makes frequent use of the second
allele for repair through gene conversion. Unlike IR, MMS does not directly cause DSBs,
but rather alkylates adenosine and guanidine bases which must be removed and
replaced [36]. DSBs still may result when abasic sites or single-strand breaks created
during base excision repair of the methylated bases stall or cause the collapse of mitotic
replication forks [44]. In diploid cells HR is the major mechanism used in repairing
such DSBs during replication, and in at least one mechanism makes use of Rad51 [43,
45]. In TbRad51-intact *T. brucei* (a diploid parasite), the repair of DSBs created within
VSG bloodstream-stage expression sites resulted in a massive increase in the rate of
VSG switching through gene conversion, presumably due to the ready availability of
related sequences [46]. This also may be accomplished in haploid organisms, if
appropriate sequences have already been replicated on the opposite strand, but would
limit the timing of gene conversion only to S phase, prior to separation of sister
chromatids.
402
The distribution and structure of *ves* genes might suggest that they have evolved for
efficient application of the SGC mechanism. In most microorganisms that undergo
antigenic variation the variant multigene families involved typically are arranged in
large, subtelomeric clusters [47]. By contrast, *ves* genes are found in numerous small
clusters scattered throughout the genome, and often are interspersed with *smorf* genes
[10, 13]. Despite extremely high overall variability, *ves* genes possess periodic tracts of
highly conserved sequence, and *ves* genes on the same or different chromosomes may
provide targets for strand invasion via such conserved tracts [12]. A small-scale
chromatin conformation capture (3C) query of sequences in proximity to the transcribed
ves genes of the LAT suggested close proximity primarily to other *ves* genes [16]. In that
organization, the ability to find short patches of homologous sequence in which to
initiate strand invasion might occur via those conserved patches. However, when the
sequences flanking the breakpoint sites were compared, patches of homologous
sequence unrelated to the highly conserved sequence regions were observed at the SGC
tract breakpoints between the LAT and donor sequences. These stretches of homologous
sequence ranged from 2 bp to 35 bp in length (Figure 7), and in some sequences no

identifiable homologous sequences could be identified. These data suggest that, for SGC

to proceed only local microhomology is required, and perhaps no homology. While these

data do not specifically identify the process responsible they are consistent with SGC not

relying upon classical HR mechanisms. The gene family- and subfamily-specific

conserved tracts found in most *ves* genes instead likely serve other functions, either in

the chromatin or in the VESA1 polypeptides they encode. Interestingly, very similar

patterns were reported in *P. falciparum* of recombination between *var2CSA* genes

associated with adhesion of infected erythrocytes to chondroitin sulfate in placental

malaria [48].

When initiating this study we hypothesized that, as Rad51 proteins are considered

essential to HR and SGC is thought to be a form of HR, BbRad51 should be essential to

this process and its loss should abrogate SGC. Our results disprove that hypothesis. The

analysis of sequences surrounding SGC breakpoints supports error-free recombination

via homologous sequences in most, but not all, instances. However, unlike Rad51-

mediated HR where tracts of homologous sequence at least 8 bp long are needed for

synapse formation [49, 50], only local sequence microhomology of at least 2 bp, or

perhaps no homology, appears to be needed (Figure 7). It is possible that this is an

artifactual result arising from misidentification of recombining sequences during

analysis. Alternatively, this may be the actual biological result, due to mismatch repair

or deletion subsequent to sequence acquisition or replication. Whereas these results

reflect the mechanism(s) responsible, they do not identify the cause. This result is

consistent with loss of the ability to incorporate exogenous plasmid sequences into the

genome via HR in the absence of BbRad51 [31], whereas only the rate and product

lengths of SGC are affected. The reason for the apparent shortening of SGC tracts

observed in knockouts is not clear. One possibility is that it may reflect the loss of some

recombinants where longer unique tracts were involved, which might require BbRad51

for successful disengagement of the invading and donor strands, repair of breaks

created during the disengagement process, or stabilization of a longer crossover region

during exchange. Regardless of any secondary effects, BbRad51 is clearly not essential to

the SGC process.

The overall reduction in isTS observed among Bb*rad51* knockouts, based upon the

numbers of alternative loci to which transcripts mapped in each line, is particularly

intriguing. At the 5-month time point, in the knockout lines transcripts appear to have

arisen from a mean of only 23.3 *ves* loci, whereas in wild-type parasites a mean of 33.3

loci had been activated within the population. Although statistically significant (p=

0.014), the biological significance of this difference is not as clear. The reason is because

these data are derived from *ves1α* genes only. Also, the maximal number of *ves* loci that

are competent to be activated is not known and the ability to modify a single locus is

extensive. About half of the *ves* family is organized in divergent (head-to-head) pairs

(approximately 33 pairs) of *ves1α/ves1β* or *ves1α/ves1α* genes that flank quasi-

palindromic, bidirectional promoter regions [12, 16]. The remainder are present as

individual *ves* genes with potentially unidirectional promoters. Among the *ves* loci

putatively activated in this study, the ratio of loci with divergent/ unidirectional

promoters ranged from 1.07 - 1.60 (mean 1.28 ± 0.24) in CE11, and 1.25 - 2.00 (mean

1.61 ± 0.29) in knockouts. Although wild type parasites, on average, appeared to activate

a higher proportion of unidirectional promoters than did knockouts, this did not reach

statistical significance (p = 0.059; S1 Table). The functionality of several bidirectional

ves promoters has been demonstrated experimentally [16], but function has not yet been

similarly tested for promoters preceding individual *ves* genes. This result clearly

indicates that a significant proportion of *ves* loci have the potential to be activated,

including individual *ves* genes, consistent with a study on the transcriptomes of pairs of

virulent *B. bovis* lines and attenuated lines derived from them. In that study, virulent

lines transcribed from a significantly wider variety of *ves* loci that included both

divergent and non-divergent loci [51]. Interestingly, among attenuated parasites *ves*

transcription was upregulated only from loci that are not divergently-oriented. The

difference in *ves* transcriptional behavior observed here and in the attenuation study

suggests that in vivo attenuation of *B. bovis* is unlikely to be related to BbRad51

expression or function. This conclusion is supported by the unperturbed Bb*rad51*

transcription observed in attenuated parasites [51].

The basis for a connection between the SGC and isTS mechanisms is not clear. Indirect

evidence allows us to propose at least three feasible explanations, each with varying

levels of support. (i) First, of the various *ves* loci represented among the transcripts, a

subset of the sequences may reflect the complete replacement of the observed region

within the original locus of active *ves* transcription (LAT) by much longer conversion

tracts, rather than by isTS. In this case, such a long replacement sequence would cause

the read to map artifactually to the donor locus rather than to the locus from which the

full *ves* gene was actually being transcribed. Given the inability to integrate exogenous

DNAs into the genome of Bb*rad51* knockout parasites [31] it is anticipated that this

would be a rare event in knockouts. From our data, we cannot rule out this possibility.

(ii) The sites failing to activate in Bb*rad51* knockouts may be unusually sensitive to

recombination, with most such events leading to lethality. However, comparison of all

the putatively activated *ves* loci in wild-type and knockout parasites reveals that there is

essentially complete overlap in the *ves* gene clusters that can be activated (S3 Figure),

directly arguing against this explanation. Rather, the knockouts appear to achieve

comparable inter-locus switches, but with a lower frequency than wild-type. (iii) A

mechanistically distinct possibility is that Bb*rad51* knockouts have a diminished

capacity to activate *ves* loci epigenetically. As a part of DNA repair, chromatin first must

be remodeled to make it accessible to the repair machinery. This occurs in part by local

chromatin decondensation through histone acetylation [52-54]. Rad51 has been

proposed to assist in the assembly of the histone acetylation machinery during repair of

dsDNA breaks and stalled replication forks [55]. In the presence of BbRad51, double-

stranded breaks in silenced *ves* loci may be successfully acetylated and repaired, but

may not always be remodeled again for silencing. With transcription of a single *ves* locus

being the default state, a choice would have to be made between the existing LAT (the

single active locus of *ves* transcription [15]) and the newly repaired/acetylated *ves* locus.

Silencing of the existing LAT would lead to isTS and establishment of a new LAT. Thus,

BbRad51 may be epigenetically effecting isTS of *ves* genes as an unintended side-effect

of the DNA repair process. If true, then in Bb*rad51* knockouts the absence of BbRad51

would be anticipated to result in frequent failure to assemble the full repertoire of repair

machinery, including epigenetic modifiers. Accordingly, isTS would be a less common

event. While explanation (i) is consistent with well-established Rad51 protein functions,

possibility (iii) is neither implausible nor inconsistent with less well-characterized

functions, and even could provide a potential mechanism for the stochastic switch

events of isTS. If substantiated experimentally, this could provide a direct link between

DNA repair and antigenic variation via isTS as well as recombination. Our currently

available evidence does not distinguish these two possibilities, and this question

warrants further investigation.

The limitations of our study include the lack of genomic data for our culture

populations. High coverage whole genome sequencing, and good quality assemblies

might improve the assessment of SGC and recombination. Still, our results demonstrate

that BbRad51 is not necessary for survival of asexual *B. bovis* in vitro or for overall

genome stability in the absence of environmental insult. Moreover, this protein is

dispensable to SGC-based antigenic variation in *B. bovis*, although it influences the rates

of SGC antigenic variation and isTS. It is not clear whether its absence would be

similarly benign during in vivo infection, where there is strong selection by host immune

responses. Evolutionary retention of BbRad51 and its involvement in recovery from

alkylation damage indicates clearly that it plays some role(s) in parasite DNA repair,

including in asexual developmental stages. The clear implication of this work is that

some component besides BbRad51 provides for the recombination observed in SGC.

Whether this is from a more distantly related member of the RecA/RadA/Rad51

superfamily proteins encoded by the *B. bovis* genome, or another enzyme class

altogether, awaits experimental evidence.

Materials and Methods

Parasite culture, transfection, and selection. This project used the clonal *B.*

bovis CE11 parasite line as starting material [6]. In vitro parasite cultures were

maintained as microaerophilous stationary phase cultures under 90% N_2/ 5% O_2/ 5%

CO_2 (v/v), essentially as described [5, 56]. Cloning of parasites was conducted by two

sequential rounds of limiting dilution cloning as described previously [7]. Parasites were

transfected with DNAs purified from *E. coli* DH5α, using EndoFree Plasmid Maxi kits

(Qiagen; Valencia, CA). Both parasitized erythrocytes and DNAs were suspended in

cytomix (120 mM KCl, 0.15 mM $CaCl_2$, 10 mM K_2HPO_4/KH_2PO_4 pH 7.6, 25 mM HEPES

pH 7.6, 2 mM EGTA, 5 mM $MgCl_2$, pH 7.6) [57] prior to electroporation.

Electroporation was performed in 2 mm-gap cuvettes, using 5 pmol linearized DNA at

1.25 kV, 25 µF, and 200 Ω, as described by Wang et al. [16], with plating at 1.25% packed

cell volume. After 24h recovery in culture, selection was initiated by addition of

blasticidin-s hydrochloride (TOKU-E; Bellingham, WA) to a final concentration of 15 µg

ml⁻¹ (32.7 µM) [58]. Every three days medium was removed, and replaced with a 2.5%

packed cell volume of uninfected erythrocytes in medium plus blasticidin-s. Once viable

parasites emerged, usually after approximately two weeks, they were maintained under

drug selection and were immediately cloned.

Validation of Bb*rad51* knock-out. Validation of Bb*rad51* gene knock-out was

performed by diagnostic PCR, Southern blotting, RT-PCR, and sequencing of the

Bb*rad51* locus. These data are presented in [31].

Phenotypic analyses.

Parasite growth assays. Parasite growth was assayed by counting Giemsa-stained

smears, with samples collected at 0, 24, and 48h growth (approximately 0, 3 and 6 cell

cycles [59]). Alternatively, in some experiments a DNA-based SYBR Green I method was

performed, essentially as described [60, 61], on parasites grown in bovine erythrocytes

depleted of leukocytes [62]. For experiments involving γ-irradiation, parasites were

exposed to a calibrated [Cs¹³⁷] source (Gammacell GC-10 gamma irradiator), on ice.

Control cells were maintained on ice for the duration of the treatment time. Samples

were immediately diluted into fresh medium containing 10% packed cell volume

uninfected erythrocytes, and placed into culture. For experiments involving MMS

exposure, parasites were exposed to MMS (diluted in complete medium) for 90 minutes,

followed by washout as described [31].

Creation of a Bbrad51-NanoLuciferase-tagged parasite line. The Bb*rad51* locus from

nucleotide 1,858,332- 1,860,393 of the *B. bovis* C9.1 line genome, already fused with 2x

HA tag and a glmS element in plasmid pBbRad51HAglmS, was opened by inverse PCR

with primers DA350 and DA351 [63]. NanoLuciferase coding sequences were amplified

from plasmid p2xHAglmS-gfp-bsd (pHggb) with primers DA348 and DA349, and

inserted into the opened plasmid with NEBuilder reagents (New England Biolabs;

Berverley, MA) to create plasmid pBbRad51NLHAglmS (pBRNLHg; S2_File). This

plasmid was linearized and transfected into *B. bovis* CE11 parasites [16]. Transfectants

were selected for resistance to blasticidin-S. Recovered transformants had experienced

fusion of the cassette to the 3' end of Bb*rad51* coding sequences by double crossover

homologous recombination, as described [31].

Chromosome reassembly. Parasites grown in leukocyte-depleted erythrocytes as

described above. Cultures, at 2.5% parasitized erythrocytes, were given 100 Gy exposure

to [^{137}Cs] on ice to fragment chromosomes. Irradiated cells were placed back into culture

to recover for designated times, then were processed for pulsed-field gel analysis [64].

Plugs were embedded into 1% SeaKem Gold agarose in 0.5x TBE buffer, and

electrophoresed for 23.5 hours at 180V, with a 50-165 second ramped switch time [65].

Gels were stained with SYBR Gold Nucleic Acid Stain (Invitrogen) for DNA

visualization, and photographed. Integrated pixel intensities were plotted for each

chromosome and the "smear" of DNA below chromosome 1 using ImageJ v. 1.52 "Gels"

and "Measure" algorithms. Corresponding blank gel regions were used for background

correction.

SGC-mediated antigenic variation assay. This experiment was performed with three

biological replicates per genotype, comprised of one clone each from three independent

Bb*rad51* knockout lines (CE11Δrad51^{ko1} H5, CE11Δrad51^{ko2} E8, and CE11Δrad51^{ko3} A5;

referred to as ko1^{H5}, ko2^{E8}, and ko3^{A5}), and three subclones of wild type CE11 line

parasites (CE11 B8, CE11 C2, and CE11 C5). Bb*rad51* knockout and wild type parasites

were cloned by limiting dilution [7]. RNAs were isolated one and five months after

parasite cloning, using Ribozol (Amresco). RNAs were treated two times with

TurboDNase (Ambion) supplemented with 1 mM MnCl$_2$ [66], followed by inactivation

with DNase Inactivation Reagent (Ambion). M-MuLV reverse transcriptase (New

England Biolabs) and oligo-d(T) primers were used to make cDNAs. A hypervariable

segment containing most of the CKRD domain of *ves*1α transcripts was amplified by RT-

PCR, using "universal" primers vesUniF2 (TGGCACAGGTACTCAGTG) and PD1R

(TACAANAACACTTGCAGCA) as described [15].

Sequencing and recombination analysis. Four independent amplifications of each

cDNA were pooled in stoichiometrically equal amounts to maximize detection of rare

variants and minimize single-sample PCR artifacts during sequencing. Paired-end

amplicon libraries were generated with NEBNext reagents (New England Biolabs) by

the University of Florida NextGen Sequencing Core Laboratory, incorporating Illumina

TruSeq index sequences. Libraries, spiked with 8% PhiX genomic library as internal

control, were sequenced on the Illumina MiSeq platform, using the Illumina Pipeline

1.8. Fastq reads were analyzed using Qiime2 pipeline [67]. Quality and adapter

trimming were performed using CutAdapt [68, 69]. Further de-noising and amplicon

sequence variant (ASV)-calling were performed using DADA2 [70], truncating the reads

at 230 nt and allowing a maximum of 5 expected errors per read. In order to remove

potential contaminating sequences, ASVs were aligned with the bovine genome, using

BLAST [71]. In order to identify recombinant sequences, ASVs were first aligned using

CLC Main Workbench, version 6.9.2. Recombination analyses were then performed on

the alignments with RDP, GENECONV, Bootscan, Maxchi, Chimaera, SiSscan, and

3Seq, as implemented in RDP4 [72]. Only recombinant events identified by a minimum

of 4 out of seven tests (at $p \leq 0.05$) were considered statistically supported and included

in downstream analyses. Note that no genome is currently available for the CE11 line.

Therefore, the C9.1 line genome was used, as the C9.1 and CE11 lines are sibling clonal

lines derived from the MO7 clonal line [6, 7, 12]. Recombinant results, including

extraction of SGC tracts, were summarized using custom scripts written with R version

3.6.3 [73] through the RStudio shell. Mean lengths of SGC tracts were compared by one-

way ANOVA, whereas tract length distributions were compared by the Mann-Whitney

Rank Sum test, without expectation of a normal distribution of variance, using

SigmaPlot version 11.0 (Systat Software, Inc.; San Jose, CA).

Data availability. Raw sequence reads are publicly available for download from

NCBI, as BioProject #PRJNA357248 (accession numbers SRR5110992-SRR5111003).

All recombinant transfection constructs are available upon request.

Acknowledgments

This work was supported by funds from National Institutes of Health grants R01

AI0055864 and T32 AI0007110, USDA Animal Health project FLA-VME-005011, and

U.F. College of Veterinary Medicine. The authors thank Allison Vansickle for assistance

with animal care and handling, and Kevin Brown, Linda Bloom, Salvador Castaneda

Barba, Eva Top, and Olivia Kosterlitz for helpful comments.

Author contributions

Conceptualization: DRA, EAM

Data curation: DRA

Experimental design: DRA, EAM, Y-PX

Formal analysis: DRA, EAM, MST

Funding acquisition: DRA

Investigation: DRA, EAM, SQ, Y-PX

Methodology: DRA, EAM, MST, Y-PX

Project administration: DRA

Resources: DRA

Supervision: DRA

Writing- original draft: DRA, EAM

Writing- review and editing: DRA, EAM, MST

Figure Captions

Figure 1. BbRad51 levels are not upregulated in response to DNA damage. *B. bovis* CE11 parasites were given acute exposure to 250 μM MMS to induce DNA alkylation, then placed back into culture to allow upregulation of DNA repair proteins. Cells were harvested and assayed for expression levels of NanoLuciferase. 1-4 samples are transiently transfected parasites expressing NanoLuciferase from episomal plasmid pHggb; 5-8 samples are expressing BbRad51-NanoLuciferase-2xHA fusion protein from the endogenous Bb*rad51* locus and under control of the Bb*rad51* promoter. Because expression was at quite different levels depending upon the source, values have been normalized as a proportion of the signal produced by the untreated control in each set (samples 1). "1" and "5", mock-treated, grown in complete medium (CM); "2" and "6", MMS-treated, CM; "3" and "7", mock-treated, grown in CM containing 400 μM glcN; "4" and "8", MMS-treated, 400 μM glcN. Parasites were allowed to recover from MMS treatment for 3 hours **(A)** or for 16h **(B)**. Asterisks indicate statistically significance difference from "1" control samples of each series (p < 0.001). No other differences were statistically significant.

Figure 2. Survival and growth of Bb*rad51* knockouts following γ-irradiation is not different from wild-type. A. Parasites were exposed to varying dosages of γ-irradiation, placed back into culture, sampled at 24 and 48h post-irradiation, and percent parasitized erythrocytes determined microscopically. In this individual experiment a significant difference was observed at 48h in the non-irradiated control samples only. This was not a reproducible difference. This assay shows that there is no increase in percent parasitized erythrocytes over time after 200 Gy exposure. **B.** Comparison of CE11 wild-type with knockout clonal lines ko2[E8] and ko3[A5], determined by a Sybr Green DNA-detection assay [31]. Sybr Green data are plotted as the proportion of the signal exhibited by the non-irradiated control samples (± 1 s.d.). No reproducible significant differences were observed among the parasite lines at these dosages. Note that the ko1[H5] line was derived approximately one year prior to ko2[E8] and ko3[A5] lines and was assessed differently, by microscopic observation of percent parasitized erythrocytes. In each case, this experiment was repeated three times with four replicates per sample. No reproducible differences were observed between *B. bovis*

CE11 wild-type and any of the knockout lines.

Figure 3. Chromosomes of γ-irradiated parasites reassembled with the same

kinetics. *B. bovis* CE11 wild type and ko1^{H5} clonal line were exposed to 100 Gy γ-

irradiation, then allowed allowed to recover for 0, 8, 16, or 24h. **A.** Reassembly of

chromosomes was assessed by PFGE (CE11 wild type is shown as an example). **B.** Plot of

relative signal intensities of *B. bovis* CE11 wild type and ko1^{H5} DNAs within individual

chromosomes, and in the "smear" below chromosome 1. The plot includes the means (±

1 s.d.) of four independent experiments. Statistical analyses using Student's T-test

revealed no significant differences between wild type and knockout parasites at any time

point. NC, non-irradiated control.

Figure 4. Variation arose over time in the sequences of *ves*1α transcripts.

RT-PCR amplification of the highly variable CKRD domain of *ves*1α transcripts and

deep sequencing of amplicons was used to assess variation arising over time. Three

freshly cloned lines of *B. bovis* CE11 wild type (lines CE11^{B8}, CE11^{C2}, and CE11^{C5}), and

one clonal line from each of three independent CE11Δ*rad51* knockout lines were

observed 1 and 5 months post-cloning. **A.** Typical structure of *ves*1α genes. Positions of

primers vesUniF2 and PD1R used in PCR reactions are shown, delineating the region

amplified (modified from [15]). **B.** All merged amplicon sequences were mapped to the

B. bovis C9.1 genome, using CLC Genomics Workbench. Mapping tracks are shown for

each sample. The X axis represents the 8 Mbp *B. bovis* C9.1 genome as a single linear

sequence, the Y axis represents the proportion of sequences aligning to a given *ves* locus

within the genome, and the alternating blue and white bands represent data from 1

month (blue) and 5 month (white) time-points for each parasite line. The area of each

peak is proportional to the numbers of reads mapping to that locus relative to the total.

Transcripts from many activated loci are present in too low abundance to be observable

in this plot. These data confirm earlier results suggesting that *ves* transcription is

monoparalogous [15], but also show that isTS appears to occur over time. **C.** Numbers of

transcriptionally active *ves* loci detected at each time point (data is pooled for knockouts

and for CE11 samples). Slotted crossbars indicate means, solid crossbars indicate

medians, and box boundaries represent 25% and 75% confidence intervals. Statistical

significance of differences among samples, based upon one-way ANOVA, are indicated.

Figure 5. SGC tracts in knockout parasite populations were shorter than in

wild-type. The lengths of SGC tracts identified in amplicons with full statistical

support were compared for each of the six clonal lines under study. Data from 1 mo. and

5 mo. samples were pooled for each sample. A statistically significant difference (p <

0.001) was observed between wild-type and knockout parasites SGC tracts.

Figure 6. The numbers of unique SGC tracts created over time by knockouts

is reduced. The numbers of unique recombinant SGC tracts with full statistical

support were plotted, normalized per million merged reads. This value provides a

surrogate measure of the total variation presented by a parasite population. Slotted

crossbars indicate means, solid crossbars indicate medians, and the box boundaries

represent 25% and 75% confidence intervals. Asterisk: a statistically significant

difference of p = 0.035 was observed between wt and knockout parasites when

comparing the total populations of unique recombinants (Student's t-test). wt, pooled

CE11 wild-type samples; ko, pooled knockout parasite samples.

Figure 7. Lengths of homology patches at SGC breakpoints did not differ.

Patches of homologous sequence were observed at the breakpoints between the

sequences of transcripts representing loci that had served as sequence donors and those

of the recipient loci during SGC modifications of actively transcribed ves genes. These

tracts of homology were identified through alignments of donor and recipient sequences

(S7 Figure), and plotted for a minimum of 15 recombinant sequences from each sample.

Slotted crossbars indicate means, solid crossbars indicate medians, and the box

boundaries represent 25% and 75% confidence intervals. No statistically significant

differences were observed between any of the samples (p= 0.133; Kruskal-Wallis one

way analysis of variance on ranks).

Supporting Information Captions

S1 Figure. Structures of plasmids pBbRad51-NanoLuciferase-2xHA-glmS and p2xHA-glmS-gfp-bsd plasmids. Complete sequences of these plasmids are provided in S2_File.fa and S3_File.fa, in fasta format. Prior to transfection, pBbRad51-NanoLuciferase-2xHA-glmS was linearized at the NotI site to facilitate double crossover integration as described in Results.

S2 Figure. Titration of γ-irradiation dosage on *B. bovis* CE11 (wild type). Parasites were exposed to 0- 1000 Gy irradiation from a calibrated Cs[137] source, then placed back into culture and the percent parasitized erythrocytes determined from Giemsa-stained smears made at 0, 24, and 48h growth. For experiments requiring survival of a proportion of the parasites the 100 Gy dosage was chosen.

S3 Figure. Locations of *ves* loci from which transcripts and SGC tracts arose. The genomic locations of *ves* loci to which amplicons mapped (plotting the full 8 Mbp genome as a single linear element), and from which they were presumed to be transcribed, is plotted relative to the numbers of parasite lines transcribing from that locus. No *ves* gene clusters within the genome were transcribed by CE11 parasites that could not also be transcribed by knockout parasites, and vice versa. For this plot, data were pooled from both time-points of all three wt or all three knockout lines (i.e., maximally 6 possible per *ves* locus).

S4 Figure. Statistically supported recombinants *vs*. total unique amplicons. A regression plot was made of the frequencies of merged amplicons with full statistical support for identification as true recombinants against the total numbers of unique amplicons observed for each population, with each type of value normalized per million reads (R^2= 0.9228).

S5 Figure. Cumulative distribution of unique SGC tract lengths in wild-type and knockout parasites. The length distribution of unique SGC tracts was plotted as the cumulative proportion of total SGC tracts against tract lengths. Samples were pooled for all six samples for wild-type and knockout parasites. These pooled data are from the

same CE11 wild-type and knockout lines shown individually in Figure 4B. The vertical

index lines indicates the median values of each population. Differences among the

distributions in each sample type were determined by the Mann-Whitney Rank Sum test

(p = <0.001).

S6 Figure. Unique SGC tracts per active *ves* locus. The numbers of unique SGC

tracts with full statistical support per transcriptionally active *ves* locus is shown for *B.*

bovis CE11 wild-type and knockout populations at 1 and 5 months growth post-cloning.

The data were pooled from all three lines of each population type. Slotted crossbars

indicate means, solid crossbars indicate medians, and the box boundaries represent 25%

and 75% confidence intervals. Values did not vary significantly among samples (p=

0.986, based upon one-way ANOVA).

S7 Figure. Examples of homology patches observed in recombinant

amplicons. In each case, the top sequence is the minor parent, the bottom the major

parent, and the middle sequence is the recombinant sequence. The gray boxes identify

the regions of homology within which recombination presumably occurred. Illustrated

are patches of (A) 2 bp, (B) 9 bp, and (C) a region wherein no identifiable patch of

homology may be found.

S1 Table. Organizational nature of the *ves* loci to which reads mapped. Loci

to which amplicons mapped are listed for all 12 samples (i.e., 1-month and 5-month

RNAs from all three wild-type and knockout lines), along with mean lengths and the

nature of each locus. Both divergent, bidirectional loci and unidirectional loci were

active in transcription at some level in all samples.

S1_File.zip. This .zip file contains individual files of SGC tracts identified for all six

clonal lines, in .csv format.

S2_File.fa. This file contains the full sequence of the pBbRad51-NanoLuciferase-

2xHA-glmS plasmid (pBRNLHg), in fasta format.

S3_File.fa. This file contains the full sequence of the p2xHA-glmS-gfp-bsd plasmid

(pHggb), in fasta format.

References

1.   Wright IG. An electron microscopic study of intravascular agglutination in the
cerebral cortex due to *Babesia argentina* infection. Int J Parasitol. 1972;2:209-15.

2.   Wright IG. Ultrastructural changes in *Babesia argentina*-infected erythrocytes in
kidney capillaries. J Parasitol. 1973;59:735-6.

3.   Wright IG, Goodger BV, McKenna RV, Mahoney DF. Acute *Babesia bovis*
infection: A study of the vascular lesions in kidney and lung. Z Parasitenkd. 1979;60:19-
27.

4.   Allred DR, Al-Khedery B. Antigenic variation and cytoadhesion in *Babesia bovis*
and *Plasmodium falciparum*: different logics achieve the same goal. Mol Biochem
Parasitol. 2004;134:27-35.

5.   Allred DR, Hines SA, Ahrens KP. Isolate-specific parasite antigens of the *Babesia*
bovis-infected erythrocyte surface. Mol Biochem Parasitol. 1993;60:121-32.

6.   O'Connor RM, Allred DR. Selection of *Babesia bovis*-infected erythrocytes for
adhesion to endothelial cells co-selects for altered variant erythrocyte surface antigen
isoforms. J Immunol. 2000;164:2037-45.

7.   Allred DR, Cinque RM, Lane TJ, Ahrens KP. Antigenic variation of parasite-
derived antigens on the surface of *Babesia bovis*-infected erythrocytes. Infect Immun.
1994;62:91-8.

8.   O'Connor RM, Lane TJ, Stroup SE, Allred DR. Characterization of a variant
erythrocyte surface antigen (VESA1) expressed by *Babesia bovis* during antigenic
variation. Mol Biochem Parasitol. 1997;89:259-70.

9.   Xiao Y-P, Al-Khedery B, Allred DR. The *Babesia bovis* VESA1 virulence factor
subunit 1b is encoded by the 1β branch of the *ves* multigene family. Mol Biochem
Parasitol. 2010;171:81-8.

10.   Jackson AP, Otto TD, Darby A, Ramaprasad A, Xia D, Echaide IE, et al. The
evolutionary dynamics of variant antigen genes in *Babesia* reveal a history of genomic
innovation underlying host-parasite interaction. Nucleic Acids Res. 2014;42:7113-31.

11.   Allred DR, Carlton JM-R, Satcher RL, Long JA, Brown WC, Patterson PE, et al.
The *ves* multigene family of *B. bovis* encodes components of rapid antigenic variation at
the infected erythrocyte surface. Mol Cell. 2000;5:153-62.

12.   Al-Khedery B, Allred DR. Antigenic variation in *Babesia bovis* occurs through
segmental gene conversion of the *ves* multigene family, within a bidirectional site of
transcription. Mol Microbiol. 2006;59:402-14.

13.   Brayton KA, Lau AOT, Herndon DR, Hannick L, Kappmeyer LS, Berens SJ, et al.
Genome sequence of *Babesia bovis* and comparative analysis of Apicomplexan
hemoprotozoa. PLoS Pathog. 2007;3:e148.

14.   Ferreri LM, Brayton KA, Sondgeroth KS, Lau AOT, Suarez CE, McElwain TF.
Expression and strain variation of the novel "*small open reading frame*" (*smorf*)
multigene family in *Babesia bovis*. Int J Parasitol. 2012;42:131-8.

15.   Żupańska AK, Drummond PB, Swetnam DM, Al-Khedery B, Allred DR. Universal
primers suitable to assess population dynamics reveal apparent mutually exclusive
transcription of the *Babesia bovis ves*1α gene. Mol Biochem Parasitol. 2009;166:47-53.

16.   Wang X, Bouchut A, Xiao Y-P, Al-Khedery B, Allred DR. Characterization of the
unusual bidirectional *ves* promoters driving VESA1 expression and associated with
antigenic variation in *Babesia bovis*. Eukaryotic Cell. 2012;11:260-9.

17.   Symington LS, Rothestein R, Lisby M. Mechanisms and regulation of mitotic
recombination in *Saccharomyces cerevisiae*. Genetics. 2014;198:795-835.

18.   Wyman C, Kanaar R. DNA double-strand break repair: All's well that ends well.
Annu Rev Genet. 2006;40:363-83.

19.   Sugiyama T, Kantake N. Dynamic regulatory interactions of Rad51, Rad52, and
Replicaton Protein-A in recombination intermediates. J Mol Biol. 2009;390:45-55.

20.   Petukhova G, Stratton S, Sung P. Catalysis of homologous DNA pairing by yeast
Rad51 and Rad54 proteins. Nature. 1998;393:91-4.

21.   Inbar O, Kupiec M. Homology search and choice of homologous partner during
mitotic recombination. Mol Cell Biol. 1999;19(6):4134-42.

22.   Bugreev DV, Mazina OM, Mazin AV. Rad54 protein promotes branch migration
of Holliday junctions. Nature. 2006;442:590-3.

23.   Mortimer RK, Fogel S. Genetical interference and gene conversion. In: Grell RF,
editor. Mechanisms in Recombination. New York: Plenum Publishing Corporation;
1974. p. 263-75.

24.   Anderson DG, McKay LL. Isolation of a recombination-deficient mutant of
Streptococcus lactis ML3. J Bacteriol. 1983;155(2):930-2.

25.   Morita T, Yoshimura Y, Yamamoto A, Murata K, Mori M, Yamamoto H, et al. A
mouse homolog of the *Escherichia coli recA* and *Saccharomyces cerevisiae RAD51*
genes. Proc Natl Acad Sci (USA). 1993;90(14):6577-80.

26.   Siede W, Friedl AA, Dianova I, Eckardt-Schupp F, Friedberg EC. The
Saccharomyces cerevisiae Ku autoantigen homologue affects radiosensitivity only in the
absence of homologous recombination. Genetics. 1996;142(1):91-102.

27.   Essers J, Hendriks RW, Swagemakers SM, Troelstra C, de Wit J, Bootsma D, et
al. Disruption of mouse RAD54 reduces ionizing radiation resistance and homologous
recombination. Cell. 1997;89(2):195-204.

28.   McCulloch R, Barry JD. A role for RAD51 and homologous recombination in
Trypanosoma brucei antigenic variation. Genes Dev. 1999;13:2875-88.

29.   Roy N, Bhattacharyya S, Chakrabarty S, Laskar S, Babu SM, Bhattacharyya MK.
Dominant negative mutant of *Plasmodium* Rad51 causes reduced parasite burden in
host by abrogating DNA double strand break repair. Mol Microbiol. 2014;94(2):353-66.

30.   Tsuzuki T, Fukii Y, Sakumi K, Tominaga Y, Nakao K, Sekiguchi M, et al. Targeted
disruption of the *Rad51* gene leads to lethality in embryonic mice. Proc Natl Acad Sci
(USA). 1996;93:6236-40.

31.   Mack EA, Xiao Y-P, Allred DR. Knockout of *Babesia bovis rad51* ortholog and its
complementation by expression from the BbACc3 artificial chromosome platform. PLoS
One. 2019;14(8):e0215882. doi: https://doi.org/10.1371/journal.pone.0215882.

32.   Cohen Y, Dardalhon M, Averbeck D. Homologous recombination is essential for
RAD51 up-regulation in *Saccharomyces cerevisiae* following DNA crosslinking damage.
Nucleic Acids Res. 2002;30(5):1224-32.

33.   Collins JA, Irnov I, Baker S, Winkler WC. Mechanism of mRNA destabilization by
the *glmS* ribozyme. Genes Dev. 2007;21:3356-68.

34.   Suarez CE, McElwain TF. Stable expression of a GFP-BSD fusion protein in
Babesia bovis merozoites. Int J Parasitol. 2008;39:289-97.

35.   Prommana P, Uthaipibull C, Wongsombat C, Kamchonwongpaisan S, Yuthavong
Y, Knuepfer E, et al. Inducible knockdown of *Plasmodium* gene expression using the
glmS ribozyme. PLoS One. 2013;8(8):e73783.

36.   Lundin C, North M, Erixon K, Walters K, Jenssen D, Goldman AS, et al. Methyl
methanesulfonate (MMS) produces heat-labile DNA damage but no detectable in vivo
DNA double-strand breaks. Nucleic Acids Res. 2005;33(12):3799-811.

37.   Horrocks P, Pinches R, Christodoulou Z, Kyes SA, Newbold CI. Variable *var*
transition rates underlie antigenic variation in malaria. Proc Natl Acad Sci (USA).
2004;101:11129-34.

38.   Kelso AA, Waldvogel SM, Luthman AJ, Sehorn MG. Homologous recombination
in protozoan parasites and recombinase inhibitors. Front Microbiol. 2017;8:1716.

39.   Bhattacharyya MK, Norris DE, Kumar N. Molecular players of homologous
recombination in protozoan parasites: implication for generating antigenic variation.
Infect Genet Evol. 2004;4:91-8.

40.   Achanta SS, Varunan SM, Bhattacharyya S, Bhattacharyya MK. Characterization
of Rad51 from apicomplexan parasite *Toxoplasma gondii*: an implication for inefficient
gene targeting. PLoS One. 2012;7(7):e41925.

41.   Fox BA, Falla A, Rommereim LM, Tomita T, Gigley JP, Mercier C, et al. Type II
Toxoplasma gondii KU80 knockout strains enable functional analysis of genes required
for cyst development and latent infection. Eukaryotic Cell. 2011;10:1193-206.

42.   Kirkman LA, Lawrence EA, Deitsch KW. Malaria parasites utilize both
homologous recombination and alternative end joining pathways to maintain genome
integrity. Nucleic Acids Res. 2014.

43.   Krogh BO, Symington LS. Recombination proteins in yeast. Annu Rev Genet.
2004;38:233-71.

44.   Ensminger M, Iloff L, Ebel C, Nikolova T, Kaina B, Löbrich M. DNA breaks and
chromosomal aberrations arise when replication meets base excision repair. J Cell Biol.
2014;206(1):29-43.

45.   Yeeles JTP, Poli J, Marians KJ, Pasero P. Rescuing stalled or damaged replication
forks. Cold Spring Harb Perspect Biol. 2013;5:a012815.

46.   Boothroyd CE, Dreesen O, Leonova T, Ly KI, Figueiredo LM, Cross GAM, et al. A
yeast-endonuclease-generated DNA break induces antigenic switching in *Trypanosoma
brucei*. Nature. 2009;459:278-81. doi: doi:10.1038/nature07982.

47.   Deitsch KW, Moxon ER, Wellems TE. Shared themes of antigenic variation and
virulence in bacterial, protozoal, and fungal infections. Microbiol Mol Biol Rev.
1997;61:281-93.

48.   Talundzic E, Shah S, Fawole O, Owino S, Moore JM, Peterson DS. Sequence
polymorphism, segmental recombination and toggling amino acid residues within the
DBL3X domain of the VAR2CSA placental malaria antigen. PLoS One.
2012;7(2):e31565. doi: doi:10.1371/journal.pone.0031565.

49.   Hsieh P, Camerini-Otero CS, Camerini-Otero RD. The synapsis event in the
homologous pairing of DNAs: RecA recognizes and pairs less than one helical repeat of
DNA. Proc Natl Acad Sci (USA). 1992;89:6492-6.

50.   Lee JY, Terakawa T, Zi Z, Steinfeld JB, Redding S, Kwon Y, et al. Base triplet
stepping by the Rad51/RecA family of recombinases. Science. 2015;349(6251):977-81.

51.   Pedroni MJ, Sondgeroth KS, Gallego-Lopez GM, Echaide I, Lau AOT.
Comparative transcriptome analysis of geographically distinct virulent and attenuated
Babesia bovis strains reveals similar gene expression changes through attenuation.
BMC Genomics. 2013;14:763.1-13.

52.   Sivanand S, Rhoades S, Jiang Q, Lee JV, Benci J, Zhang J, et al. Nuclear acetyl-
CoA production by ACLY promotes homologous recombination. Mol Cell. 2017;67:252-
65.

53.   Gong F, Miller KM. Mammalian DNA repair: HATs and HDACs make their mark
through histone acetylation. Mutat Res. 2013;750:23-30.

54.   Price BD, D'Andrea AD. Chromatin remodeling at DNA double-strand breaks.
Cell. 2013;152:1344-54.

55.   Rowlands H, Dhavarasa P, Cheng A, Yankulov K. Forks on the run: can the
stalling of DNA replication promote epigenetic changes? Frontiers in Genetics.
2017;8:86.

56.   Levy MG, Ristic M. *Babesia bovis*: continuous cultivation in a microaerophilous
stationary phase culture. Science. 1980;207:1218-20.

57.   van den Hoff MJB, Moorman AFM, Lamers WH. Electroporation in 'intracellular'
buffer increases cell survival. Nucleic Acids Res. 1992;20:2902.

58.   Mack E. Unexpected effects of Rad51 deletion on viability, chromosome repair,
and antigenic variation in *Babesia bovis* [Ph.D. Dissertation]: University of Florida;
2014.

59.   Franssen FFJ, Gaffar FR, Yatsuda AP, de Vries E. Characterisation of erythrocyte
invasion by *Babesia bovis* merozoites efficiently released from their host cell after high-
voltage pulsing. Microbes Inf. 2003;5:365-72.

60.   Smilkstein M, Sriwilaijaroen N, Kelly JX, Wilairat P, Riscoe M. Simple and
inexpensive fluorescence-baseed technique for high-throughput antimalarial drug
screening. Antimicrob Agents Chemother. 2004;48(5):1803-6.

61.   Bennett TN, Paguio M, Gligorijevic B, Seudieu C, Kosar AD, Davidson E, et al.
Novel, rapid, and inexpensive cell-based quantification of antimalarial drug efficacy.
Antimicrob Agents Chemother. 2004;48(5):1807-10.

62.   Ambrosio RE, Potgieter FT, Nel N. A column purification procedure for the
removal of leucocytes from parasite-infected bovine blood. Onderstepoort J Vet Res.
1986;53:179-80.

63.   Triglia T, Peterson MG, Kemp DJ. A procedure for in vitro amplification of DNA
segments that lie outside the boundaries of known sequences. Nucleic Acids Res.
1988;16:8186.

64.   Ray BK, Bailey CW, Jensen JB, Carson CA. Chromosomes of *Babesia bovis* and
Babesia bigemina. Mol Biochem Parasitol. 1992;52:123-6.

65.   Alleman AR, Kamper SM, Viseshakul N, Barbet AF. Analysis of the *Anaplasma*
marginale genome by pulsed-field electrophoresis. J Gen Microbiol. 1993;139:2439-44.

66.   Bauer P, Rolfs A, Regitz-Zagrosek V, Hildebrandt A, Fleck E. Use of manganese
in RT-PCR eliminates PCR artifacts resulting from DNase I digestion. Biotechniques.
1997;22:1128-32.

67.   Bolyen E, Riedeout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.
Reproducible, interactive, scalable and extensible microbiome data science using QIIME
2. Nat Biotechnol. 2019;37:852-7.

68.   Bokulich NA, Subramanian S, Faith JJ, Gevers D, Gordon JI, Knight R, et al.
Quality-filtering vastly improves diversity estimates from Illumina amplicon
sequencing. Nature Methods. 2013;10(1):57-9. doi: DOI:10.1038/NMETH.2276.

69.   Martin M. Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnetJournal. 2011;17(1):1-3.

70.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP.
DADA2: high-resolution sample inference from Illumina amplicon data. Nature
Methods. 2016;13(7):581-3. doi: DOI:10.1038/NMETH.3869.

71.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment
search tool. J Mol Biol. 1990;215:403-10.

72.   Martin DP, Murrell B, golden M, Khoosal A, Muhire B. RDP4: detection and
analysis of recombination patterns in virus genomes. Virus Evolution. 2015;1(1):vev003.
doi: doi: 10.1093/ve/vev003.

73.   R_Core_Team. R: A language and environment for statistical computing Vienna,
Austria: R Foundation for Statistical Computing; 2018. Available from: https://www.r-
project.org/.
1020

Figure 1

Figure 4

Figure 5

Figure 6

Figure 7

Figure 2

Figure 3