Insights into gene expression changes under conditions that facilitate horizontal gene transfer (mating) of a model Archaeon

Authors: Andrea Makkay[1,$], Artemis S. Louyakis[1,$], Nikhil Ram-Mohan[2], Uri Gophna[3], J. Peter Gogarten[1,4], R. Thane Papke[1*]

Affiliations:
[1]Department of Molecular and Cell Biology, University of Connecticut, Storrs, CT, USA
[2]Department of Emergency Medicine, Stanford University School of Medicine, Stanford, CA, USA
[3]School of Molecular Cell Biology and Biotechnology, Tel Aviv University, P.O. Box 39040, Tel Aviv 6997801, Israel
[4] Institute for Systems Genomics, University of Connecticut, Storrs, Connecticut, USA

*Corresponding author: thane@uconn.edu
$These authors contributed equally

Keywords: Transcriptome, Archaea, Haloarchaea, horizontal gene transfer, mating

Funding: This work was supported through grants from the Binational Science Foundation (BSF 2013061); the National Science Foundation (NSF/MCB 1716046) within the BSF-NSF joint research program; and NASA exobiology (NNX15AM09G, and 80NSSC18K1533).

Abstract: Horizontal gene transfer is a means by which bacteria, archaea, and eukaryotes are able to trade DNA within and between species. While there are a variety of mechanisms through which this genetic exchange can take place, one means prevalent in the archaeon *Haloferax volcanii* involves the transient formation of cytoplasmic bridges between cells and is referred to as mating. This process can result in the exchange of very large fragments of DNA between the participating cells. Genes governing the process of mating, including triggers to initiate mating, mechanisms of cell fusion, and DNA exchange, have yet to be characterized. We used a transcriptomic approach to gain a more detailed knowledge of how mating might transpire. By examining the differential expression of genes expressed in cells harvested from mating conditions on a filter over time and comparing them to those expressed in a shaking culture, we were able to identify genes and pathways potentially associated with mating. These analyses provide new insights into both the mechanisms and barriers of mating in *Hfx. volcanii*.

---

1. Introduction

In eukaryotic sexual reproduction, the recombination of DNA is intertwined with reproduction through the processes of meiosis and syngamy, but in prokaryotes the two processes are separated; they reproduce asexually and transfer fragments of DNA between cells in a donor-recipient fashion, a process called horizontal or lateral gene transfer (HGT or LGT) [1, 2] . There is a diversity of known mechanisms by which DNA can move between cells. The best-characterized means of HGT include transduction, the movement of DNA between cells via a phage or virus; transformation, the uptake of naked environmental DNA (eDNA); and conjugation, the movement of genetic material, usually plasmids (selfish genetic elements), via a narrow channel that transfers single-stranded DNA [3].

Very little is known about HGT mechanisms in archaeal species. Natural transformation has been detected in four archaeal species [4-7]. However, the molecular machinery by which DNA is transported across the membrane remains elusive. For instance, the required pore protein gene *comEC* and its homologs have not been detected in Archaea [8, 9]. A cell-cell contact mechanism, the "crenarchaeal system for exchange of DNA" (Ced) [10] found in *Sulfolobus acidocaldarius* imports DNA from the donor, rather than export it like bacterial conjugation requires protein families that have only been identified in crenarchaeal organisms to date [10]. The oldest report for HGT in an archaeon is the cell-cell contact dependent mechanism called "mating" found in the hypersaline adapted euryarchaeon *Haloferax volcanii* [11-13]. The cell-cell contact mechanism exhibited by *Hfx. volcanii* involves fusion of (at least) two cells that results in a heteroploid state [14, 15]. Analysis of mating during the formation of a biofilm demonstrated approximately the same efficiency as seen under filter-mediated laboratory experiments, suggesting that biofilms are a natural condition under which *Hfx. volcanii* undergoes HGT, and that biofilm formation and HGT could be intertwined physiological pathways [16].

The process and genes involved in this archaeal mating system have been interrogated, though many of the details remain unknown. Mating cells require cell-cell contact before they can recombine DNA and there is a

limited mating range [17], so it is reasonable to assume the presence of a self/non-self recognition system. Laboratory experiments have confirmed this, showing that when cell fusion events were compared, higher efficiency was observed within than between closely related *Haloferax* species by a factor of about 8 [15]. This self/other distinction may be the result of differential glycosylation of S-layer proteins. S-layer glycosylation is required for mating [18, 19] and their pathways can be highly diverse within haloarchaeal species. *Hfx. volcanii* expresses different S-layer glycosylation when required to grow at lower-than-optimal salinity [20], and these pathways are constructed uniquely amongst individuals in a population found on the rocks of Tel Aviv that are identical for their rpoB gene [21].

Though little is known about which additional genes contribute to mating, or how the molecular "machinery" works together for the exchange of DNA, it is recognized that the impact of HGT on Haloarchaea is perhaps greater than for other organisms. For instance, while a greater than one-thousand-fold reduction in recombination is also observed in Haloarchaea, the drop off is more gradual with phylogenetic distance, indicating *Hfx. volcanii* has lower barriers to interspecies gene flow and that interspecies recombination potentially plays a more important evolutionary role than in *E. coli*, for instance [15]. Phylogenetic analysis of 21 haloarchaeal genomes revealed that homologous recombination declines log-linearly as genetic relatedness decreases, as seen between *E. coli* and *Salmonella* [22]. Notably however, Haloarchaea have acquired several hundreds of genes and gene fragments from Bacteria [23-25].

To uncover more details regarding how the HGT mating mechanism might work in *Hfx. volcanii*, we undertook a transcriptome approach to determine which genes are differentially expressed when moving a population of cells from a shaking culture environment to being immobilized under filter-mating conditions, over a 24-hour period. Here we examine the changes in the expression of some groups of genes we feel are relevant to the overall course of mating in *Hfx. volcanii*. While further work is warranted for the intricacies of the mating process to be unraveled, these data provide a foundation upon which to direct additional research and can provide some insight into barriers of mating in halophilic archaea.

2. Methods

Haloferax volcanii strains H53 and H98, auxotrophs for tryptophan and thymidine respectively, were grown at 42°C with shaking in casamino acids (CA) medium [26]. CA mediums contains per liter: 144 g of NaCl, 21 g of $MgSO_4 \cdot 7H_2O$, 18 g of $MgCl_2 \cdot 6H_2O$, 4.2 g of KCl, 5.0 g casamino acids, and 12 mM Tris HCl (pH 7.5). As needed the medium was supplemented with 50 µg/mL uracil (ura), 50 µg/mL tryptophan (trp), and/or 40 µg/mL both thymidine and hypoxanthine (thy).

Liquid cultures of *Hfx. volcanii* were grown until late exponential phase; the cells were then pelleted and resuspended in higher volumes of fresh CA medium and allowed to grow overnight to ensure cells were in mid-log phase when mating was commenced. Cells were diluted in CA medium to $OD_{600}=0.25$ and equal volumes of each cell line were mixed together per mating replicate. In the case of suspended (non-mating) cells, the cell cultures were mixed, pelleted and resuspended directly in either 1 mL Trizol reagent (for RNA extraction) or unsupplemented CA medium (for plating). In the case of mated replicates, 25 mL cell suspension were pushed with a syringe on to 25 mm diameter, 0.45 µm nitrocellulose filters. The filters were subsequently laid, cells up, on plates of solid CA medium (+ ura, trp, thy) for the duration of mating. Each time point (0, 2, 4, 8, and 24 hours) was prepared in triplicate. At the time of harvesting, cells from one filter were resuspended in 1 mL unsupplemented CA medium by incubating at 42°C with agitation (for plating), and the remaining two filters were resuspended in 1 mL Trizol reagent (for RNA extraction). The overall mating was done in twice in triplicate each time and a graphical outline was created with BioRender.com (Fig. 1A; Table S1).

Mating counts were determined by plating cells on selective media. Whole cells counts were determined by plating 10-fold dilutions of cells on solid CA medium supplemented with ura, trp, and thy, and mated cell counts were determined by plating dilutions on unsupplemented CA agar. Dilutions were plated in triplicate, and the averages of colony counts were used. Mating percentage was determined by comparing the cell counts from unsupplemented agar to those from supplemented agar.

RNA was isolated from cells in Trizol reagent as per manufacturer's instructions. The RNA from two filters was pooled for each sample. Isolated RNA was subjected to two rounds of DNAse digestion using Turbo DNA-free (Thermo Fisher) as per manufacturer's directions. DNAse-treated RNA was finally purified using Qiagen RNeasy columns, as per manufacturer's directions. RNA was assessed for DNA contamination using PCR, and RNA quality was determined using a 2100 Bioanalyzer (Agilent genomics). cDNA libraries for sequencing were constructed using a ScriptSeq Complete kit (Illumina), including depletion of rRNA using the Ribo-Zero rRNA removal beads for bacteria. cDNA was sequenced using a 150-cycle MiSeq reagent kit v3. Libraries were pooled and sequenced at 10 pM with a 15% 12 pM PhiX spike-in.

Bioinformatics. All commands with specific parameters can be found at Gogarten-Lab github repository [27]. Quality control on the sequences included Scythe v0.991 [28] adapter trimming against Illumina universal adapter sets followed by Sickle v1.33 [29] read trimming. Trimmed reads were aligned to the *Haloferax volcanii* DS2

genome (NCBI accession GCF_000025685.1_ASM2568v1; 4.0129 Mb length; 65.46 %GC; 4,023 genes) and quantified using Salmon v0.9.1 [30] with Variational Bayesian Estimation Method (VBEM). The *Haloferax volcanii* DS2 genome annotations were cross referenced against the UniProt database [31] and a complete table of the annotations and abundance estimates is available (Table S3). Additionally, alignments were completed using bowtie2 v2.3.1 [32-34] Counts were TMM normalized and differential expression analysis was completed using NOISeq v2.26.1 [35, 36] in R by running noiseqbio function with default parameters. Differential expression analysis was completed for all comparisons [27] and comparisons between planktonic state and each time point are presented here. Interactive volcano plots and Manhattan plots were created in R using manhattanly v0.2.0. [37]. Pearson and Spearman correlation coefficients were calculated using the cor function in R stats package v3.5.2. Principle component analysis was completed in Noiseq (Euclidian distance) and using r stats prcomp function with linear regression models calculated for PC1, PC2, and PC3. Bar charts and violin plots were constructed using ggplot2 v3.1.0 [38-40]. Phylogeny for HVO_RS15650 (conjugal transfer protein) was constructed using NCBI; briefly, protein sequence was BLASTed against the non-redundant database, top 100 hits were BLAST pairwise aligned, using Fast Minimum Evolution tree method, and Grishin distance with maximum sequence distance of 0.85 [41-43]. Genes were clustered using normalized expression counts via DESeq2 v1.24.0 [44] with a likelihood ratio test and an adjusted p-value cutoff of 0.05. Differential expression using NOISeq v2.26.1 was completed for cluster three and volcano plots were produced with Manhattanly v0.2.0. Additional data preparations and organizing were completed in R using tidyr [45], dplyr [46], and reshape2 [47]. Reads were deposited to NCBI under BioProject PRJNA632928 and SRA accession numbers can be found in Table S2.

3. Results and Discussion

Mating and mating efficiency

Cells were mated by mixing the H53 and H98 auxotrophs in equal volumes and transferring the cells onto filters. H53 cells are auxotrophic for tryptophan, whereas H98 cells are auxotrophic for thymidine. The genes *trpA* and *hdrB* [26] were thus used as markers for mating. Cells able to grow on CA plates which were supplemented with neither tryptophan nor thymidine were considered as representing mated cells and the number of colony-forming units (cfus) on unsupplemented plates compared with total counts of cfus (Allers et al., 2004) on plates supplemented with both thymidine and tryptophan yielded a measure of mating efficiency at each time point. We note that this measure excludes the majority of mating events: all mating between two H53 and between two H98 go undetected, and mating between H53 and H98 are only detected in case the heteroploid state is resolved after a recombination event in a way that results in a hybrid chromosome with both functional genes. Nevertheless, this measure is expected to be proportional to the mating frequency and is a standard means of gauging mating efficiency [15, 18, 19, 48]. Mating for planktonic cells over time is not shown, because under our experimental conditions we saw no mating in planktonic cells. Even though mating in planktonic cells has once been reported, rates were 10^{-4} times less than for cells on filters, and thus mating in planktonic cells can be considered to be negligible under our experimental conditions [49].

Mating was allowed to proceed by incubating the filters on solid medium and cells were harvested for RNA isolation and plating at various time points (see Materials and Methods and Fig. 1A). The intention of this paper was to observe changes in gene expression between non-mated and mated cells over time, but likewise some changes might also reflect the cells' reaction to moving from a suspended to sessile milieu. Additionally, the total cfu count did increase over time, suggesting the cells were dividing (Fig. 1B) with a doubling time below 4 hours. While this is in agreement with some sources [50], others consider this rapid [51]. Given the rather loose adherence of the typical *Hfx. volcanii* biofilm [16, 52], the increase in cells could be representative of a combination of cell growth as well as more of the cells sloughing off filters at later time points. Regardless, some gene expression level changes could be in response to growth and changes in local nutrient availability.

Transcriptome data were taken from cells exposed to conditions that encourage mating, but there was no means of specifically selecting out mating cells from this group. Nonetheless, we feel that this data is a means of surveying for genes that may have a significant role in the cellular processes surrounding mating. Mating was observed to start rapidly, as even those cells harvested immediately were shown to produce colonies able to grow on unsupplemented CA medium (Fig. 1C). It is unlikely that mating is completed so rapidly, but the mating at 0 hours could be indicative of the onset of mating. In other words, the cells were able to adhere to one-another to initiate mating immediately, and this adherence was robust enough that the mating was able to continue to completion as the cells were washed from the filters where mating was initiated. Over the course of 24 hours, mating efficiency rose from approximately 3×10^{-4} to 4×10^{-3} which is comparable to that of previous work [15, 19]. The main increase in cfus occurred during the first 2 hours. While the mating efficiency for both replicates is in good agreement for most time points, those for 4 hours of mating are divergent. Principle component analysis of the expression data (Fig. 1D), however, shows excellent agreement between the replicates in all time points.

The transcripts were aligned to 3,941 genes in the *Hfx. volcanii* genome and expression was identified in 3,582 genes after filtering any genes with a total expression estimate of less than ten transcripts. Significant differential expression for each time point against the planktonic state transcriptome ranged from 109 genes (p-value <0.01) in the zero hour time point up to 1,092 genes at the four hour time point (Table 1). Many of those genes are related to cell growth and moving from a planktonic to sessile state, while many are associated with processes related to mating conditions. Mating occurs through stages including cell contact, cell surface fusion, a heteroploid state, and separation into hybrid cells [53]. There are genes which may contribute to or limit mating at each of these steps, so we focused on the genes, pathways, and systems we hypothesize are involved in each of these processes, and on genes that show a similar expression pattern to those involved in mating.

Glycosylation and low salt glycosylation genes

While little is known about which genes are involved with mating of *Haloferax* spp. cells, it was previously noted that mating efficiencies were greater within than between species, though inter-species mating still occurs and the barriers to this are relatively lax. More recent work was done to determine if recognition of potential mating partners might be mediated in part by the glycosylation of cell-surface glycoproteins. Briefly, it was shown that disruption of normal glycosylation by either gene knock-out or culturing cells in low-salt conditions can reduce mating efficiencies [19]. It is for this reason that glycosylation and low-salt glycosylation gene expression levels were examined.

Overall, for the majority of both glycosylation and low-salt glycosylation genes expression increased when compared to suspended cells (Fig. 2). Many of the low-salt glycosylation genes do not appear to change in expression in any manner that reflects time of mating, but the glycosylation genes show a small increase at 0 hours of mating followed by a higher increase in expression at 2 hours of mating and some tapering thereafter. This pattern, however, is by no means consistent. The two-hour time point had significantly higher expression in eleven glycosylation genes with the highest expression for the major cell surface glycoprotein, HVO_RS14660, (fold change > 16). Of the glycosylation genes that were previously investigated in relation to mating [19], *aglD* (HVO_RS08530) shows an increase in expression after 0 h of mating, and *aglB* (HVO_RS12050) shows a small decrease in expression levels after 0 hours of mating, although in the former case only the changes in gene expression after 2 and 4 hour mating are significant (p<0.05) , and in the latter, none of the changes are significant. The low-salt glycosylation gene *agl*15 (HVO_RS20240) showed a significant increase in expression levels in comparison to that of planktonic cells in all time points after 0 hours of mating (p<0.05).

In the work done by Shalev et al., the deletion of both the *aglB* and *agl15* genes showed a large negative impact on mating efficiency, with deletion of both genes together reducing mating to almost undetectable levels [19]. The expression data agrees with this observation showing an immediate increase and tapering over time. Deletion of the *aglD* gene had a very small impact on mating efficiency with even a small increase shown when both mating partners had the gene deleted [19]. In the transcriptome data, however, the expression of this gene shows a comparable pattern to *aglB* and *agl*15, but from 2 hours of mating onward. Other genes known to have an impact on cell-surface glycosylation (e.g., *aglF*, *aglG*, *aglI*; HVO_RS12035, HVO_RS12045, and HVO_RS12040, respectively), but with as-yet undescribed impact on mating, show significant increases in expression in mated cells compared with unmated cells.

These data overall suggest that while expression levels already existing in cells when they are subjected to mating conditions may have an impact, the mating itself may also have an impact on the expression of these genes. This notion, however, can be argued against as mating appears to begin immediately upon cells moving into proximity with one-another and it would not then be necessary for the cells to increase the expression of genes related to glycosylation. The changes observed thus may merely represent fluctuations resulting from the cells' growth or, most likely, a change from planktonic to sessile cells. The fact that genes whose deletions have a positive impact on mating, and those who have a negative impact on mating both show increases in expression post-mating implies that increased glycosylation is a physiological response to growth on a surface that probably affects multiple processes including mating.

Community Development and Biofilm Formation

Hfx. volcanii is known to form biofilms [16, 52] and incubating cells after filtration will form a biofilm over time. A lack of observable mating in planktonic cells (Table S1), combined with prior observation of an increase in mating in cells which are part of a biofilm [16], suggests that the majority of mating in the natural environment will take place amongst *Hfx. volcanii* that are part of a biofilm. This may be mediated in part by molecules involved in cell adhesion, for example the glycosylation genes already discussed, as well as biofilm formation and quorum sensing genes. As such, we examined which transcripts predicted to be involved in those processes increased or decreased in comparison to planktonic cells (Fig. 3).

Biofilm formation begins with adhesion of cells to a surface. Archaeal flagella, also referred to as archaella [54, 55], although sharing a role comparable to that of bacterial flagella, are assembled in a manner more similar to bacterial type IV pili [56-58]. Archaella and chemotaxis proteins modulate mobility of cells to find

alternative conditions in their aqueous environment and allow cells to move to surfaces [49, 52]. The *Hfx. volcanii* archaella are also important for adhesion [49, 52, 59, 60] and their chemotaxis proteins have been implicated in mediating social mobility within a biofilm [16]. Archaella assembly proteins had non-monotonous transcriptional changes (Fig. 3). The biogenesis components, *pilC3* and *pilB3* (HVO_RS09660, and HVO_RS09665, respectively) [61], had significantly lower expression with trending decreases in fold-change (Fig. 3). The use of the archaella for movement in the planktonic condition is supported by the decreased expression over time. However, some of the genes associated with motility, such as *pilA4*, *pilA5*, and *flaCE* (HOV_RS16515, HVO_RS04680, and HVO_RS10530, respectively), showed increased expression, but the change was not significant. While there is no notable expression pattern across all chemotaxis genes, many time points do show significant increases and decreases in expression. For example, *cheW* (HVO_RS10590), coding a purine-binding chemotaxis protein that stimulates rotation of the archaella [62], has a significant increase at all time points except 2 hours, while *cheB* (HVO_RS10585), which responds to stimuli and modulates chemotaxis [63], is significantly decreased at 2 hours and moderately decreased at all other timepoints (Fig. 3). Genes used for motility specifically would be expected to decrease in expression once cells have adhered to a surface, while those used for adherence would be expected to increase temporarily until the biofilm structure is formed. This speculation is supported by observations in the formation of bacterial biofilms [64]. When *Hfx. volcanii* archaellin genes *flgA1* and *flgA2* (HVO_RS10515 and HVO_RS10520, respectively) and the prepilin peptidase gene *pibD* (HVO_RS19185) were examined for their roles, deletion of the archaellin genes showed no impact on surface adhesion or on mating efficiency, but deletion of *pibD* did reduce adhesion to surfaces, but again mating efficiency was unaffected [49]. Neither of *flgA*1 and *flgA*2, however, had any significant changes, suggesting the mating process is not mediated by the same structures that *Haloferax* uses for surface attachment (Fig. 3). The *pibD* gene transcription levels increase in response to transfer to mating conditions with a significant increase at 8 hours (Fig. 3). This initial decrease in fold change at 0-hours is minor and does not contradict a role in adhesion with the remaining timepoint supporting this role.

Biofilm adhesion is also facilitated by the presence of extracellular polymeric substances (EPSs) made up of DNA, amyloids, and glycoproteins, which makes up the biofilm structure [52, 65-67]. EPS is produced by the cell and transported utilizing secretion system machinery. Archaeosortase genes, for example, mediate the attachment of proteins to the cell surface and have been shown to impact not only the cell growth and motility but also the amount of mating between cells [18]. Deletion of the *artA* archaeosortase gene (HVO_RS09085) decreases mating [18], an observation that correlates well with the transcriptome data: *artA* increases at all timepoints with peak expression at 4 hours and significance from 2-8 hours (Fig. 3). HVO_RS15650, which has been annotated as both a type IV secretion system DNA-binding domain-containing protein (TraD) and as a "conjugal transfer protein", had highly significant increases in expression, also with peak expression at 4 hours (Fig. 3, Fig. S1). In bacteria, type IV secretion systems (T4SS) are responsible for conjugation in some species, as well as a variety of other roles, including delivery of effector molecules to eukaryotic hosts, biofilm formation, and killing other bacteria [68]. The phylogeny of HVO_RS15650 and its homologs shows those belonging to archaeal species are in a clade separate from the bacterial homologs (Fig. S2), thus one cannot assume a conjugation-related role for this protein with certainty. While it will be interesting to see if this gene has a further role in the mating process in Archaea, the observed expression pattern could be due to a role in biofilm formation. Many putative secretion system genes appear to have lower expression levels in the filter-bound cells and many expression patterns are variable with rapid increases and decreases consistent with biofilm maturation (Fig. 3, Fig. S1). Overall, archaeosortase and type IV secretion system genes are consistent with our hypothesis for expression changes associated with a mating environment.

DedA/SNARE genes

Fusion processes have long been studied in eukaryotic systems in intracellular trafficking and vesicle maturation, viral entry, and gamete fusion [69-71]. Similar mechanisms appear to be at play in *Hfx. volcanii* mating which involves fusion and cleavage of membranes of neighbouring cells [14, 15, 53]. SNARE proteins have been implicated in eukaryotic vesicle fusion in mammalian neuronal cells and similar machinery and processes have also been described in yeast [72]. The SNARE-associated Tvp38 family of proteins in yeast have homologs in other single- and multi-cellular eukaryotes and are homologous to the DedA protein family in bacteria and archaea [73-75]. While the Tvp38 proteins do not appear essential in yeast, the DedA family of proteins in bacteria has been implicated in division defects causing long chains of incomplete cells to form [76]. While the proteins implicated in cell division are initially appropriately localized at a site of division, they disassemble again when division fails (Sikdar and Doerrler, 2010). Of archaeal genomes examined, 27% lack genes with significant sequence similarity to DedA, but all Halobacteriales thus far examined contain a DedA homolog [70]. Given the proposed manner of cell mating (partial or whole cell fusion followed by cleavage), it is reasonable to predict that DedA- or SNARE-type proteins will have a role in this process.

Two DedA family proteins found in *Hfx. volcanii* show an appreciable change in transcription (Fig. 4). HVO_RS01005 shows an immediate increase in transcription at the 0-hour time point, lower increases at 2 and

4 hours, and higher increases at 8 and 24 hours of mating, but none of the differential expression is statistically significant ($p > 0.05$). Another DedA homologue, HVO_RS09165, shows an immediate drop in expression ($p = 0.059$) followed by significantly lower expression at 2, 4, 8, and 24 hours (p-values: 0.003, 0.006, 0.010, and 0.005 respectively). HVO_RS09165 is considered to be a member of the TVP family of proteins, which are yeast SNARE proteins conserved in numerous eukaryotes [75]. It is intriguing that the expression of the TVP family member protein expression is in the opposite direction expected for genes involved in mating, and it could indicate that the transient fusion of cells during mating is mediated by the balance of DedA proteins present before cells are transferred to mating conditions.

CetZ/FtsZ

As can be seen in Fig. 1A, cells that have been incubated on filters for mating also undergo cell division over the course of 24 hours. It is probable that there is an overlap in machinery used for cell division and mating, in particular, those genes responsible for DNA replication and the cytoskeletal proteins. In mating, *Haloferax* are known to exchange large segments of both genomic and plasmid DNA [15], and passive diffusion of DNA through a cell is limited by size [77], so one can postulate that the movement of such large molecules requires the involvement of the cytoskeletal matrix. Research in eukaryotes has shown the requirement of the cytoskeleton for trafficking of DNA to the nucleus in transfected cells [78], and it is conceivable that an analogous system may be at play in the transport of DNA between mating *Haloferax* cells. Additionally, the CetZ proteins appear to mediate the shape of *Haloferax* cells [79], and thus might be responsible for the formation of cytoplasmic bridges. For these reasons, the expression levels of the CetZ and FtsZ genes were investigated.

CetZ1 has been implicated in cell motility in *Hfx. volcanii* [79]. Surprisingly, transcriptome data showed that expression of CetZ1 (HVO_RS15305) was increased in sessile cells as compared with planktonic cells (Fig. 5). While Duggin et al. (2015) showed that CetZ1 converted cells to the rod-shape required for motility, the observed increase in expression may indicate that the protein is also required for mating. Indeed, many of the CetZ genes show increased expression levels under mating conditions, with the exception of CetZ2 (HVO_RS08275) and CetZ3 (HVO_RS10050). While it is tempting to speculate that the rigidity of the cell shape may play a role in mating, further understanding as to the function of these proteins is essential.

In addition to the CetZ genes, *Hfx. volcanii* also has the genes FtsZ1 and FtsZ2 (HVO_RS08145 and HVO_RS07495, respectively). These genes have been implicated in cell division: the encoded proteins form a ring structure that effectively pinches off dividing cells from one-another [80, 81]. The FtsZ1 gene is also well-conserved amongst many Haloarchaea [82-84]. In the case of this mating transcriptome, FtsZ1 shows higher rates of transcription in the mated cells as opposed to the unmated, but FtsZ2 shows little change overall, with transcription rates being slightly lower at 0- and 2-hour time points in mating and a small increase in later time points (Fig. 5). While it is conceivable that the same mechanism of separation of dividing cells is used in the separation of mating cells, the growth and division of cells after plating may also account for the increase in transcription of FtsZ1. As yet, our understanding of the roles of the CetZ and FtsZ proteins in *Haloferax* is limited, and thus further research into their roles in mating and cell functions are warranted.

Selfish Genes

Hfx. volcanii is known to harbor selfish genes, as well as innate and adaptive means of defense against such invasion [85, 86]. Studies in natural populations have shown a larger amount of HGT within than between haloarchaeal phylogroups (Papke et al., 2007; Fullmer et al., 2014), which could, to some degree, be attributable to barriers to mating which are active after cell fusion. Although the *Hfx. volcanii* cells were mated to members of the same species in this experiment, it is possible that the act of mating might trigger expression of selfish genes and those associated with defense against invasion of selfish genetic elements.

The *Hfx. volcanii* transcriptome contains expressed transposases from at least nine insertion sequence (IS) families on the main chromosome and on plasmids pHV1, pHV3, and pHV4 (Fig. 6A). An overwhelming majority show increased expression at all or most time points compared to the shaking control; forty-two have significantly higher expression, while only ten have significantly lower expression (Fig. 6A, Fig. S3), supporting the hypothesis that selfish genes entering a mating condition will be activated. The IS5 and IS6 families' expression, for example, are almost entirely increased after mating (Fig. S3). While most of the insertion sequences have higher expression than the shaking control, there are still many expressed during shaking and a few with expression decreased in a sessile state (Fig. 6B, Fig. S3). One family, the IS110 transposases, all have higher expression during shaking (Fig. S3). The largest family of transposases (IS4) has both increased and decreased fold changes, but is mostly expressed after plating, even at the zero-time point.

Although it has been previously observed that the expression of CRISPR-associated genes is low in *Haloferax* species under normal growth conditions [87], it has been shown that not only are CRISPR spacers acquired during cross-species mating, but that such spacer acquisition will reduce the gene transfer during cross-species mating [48]. All the CRISPR-associated genes in *Hfx. volcanii* are located on pHV4 (Fig. S4A) [88]. All *cas* genes were expressed at all time points (Fig. S4B), but changes in expression levels varied. Two genes that

encode components of the interference machinery, *cas8b* and *cas7b* (HVO_RS02760 and HVO_RS02765, respectively), had significantly increased expression from 4-8 hours after plating. Two genes, *cas6* and *cas2* (HVO_RS02755 and HVO_RS2790, respectively), the former required for processing of CRISPR RNAs, and the latter involved in acquiring new spacers, were expressed significantly less than in planktonic cells starting at the 2-hour time point to 24-hours and 8-hours, respectively. *cas4* (HVO_RS02780), also assumed to be a part of the spacer acquisition module, had a significant increase in expression immediately at 0-hour turning to lower expression for the rest of the experiment. Acquisitions of spacers were shown to be induced by mating with different *Hfx.* species, e.g. *Hfx mediterranei,* but to much lower levels than within-species mating [48]. It therefore appears that while CRISPR-surveillance against threats (i.e. interference module) is maintained or even increased during mating conditions, spacer acquisition remains generally low, potentially preventing acquisition of DNA spacers from closely related cells during mating, which could result in CRISPR autoimmunity.

Another potential limiting factor on genetic exchange in *Haloferax* during mating is the activity of restriction modification (RM) system enzymes, provided the mating pair possess different sets of RM systems. RM systems also constitute addiction cassettes, and therefore can be considered selfish genetic elements (see Fullmer et al., 2019 for discussion). *Hfx. volcanii* contains a Type I RM system and a number of DNA and RNA methylases [89-91]. The strains of the *Hfx. volcanii* auxotophs used in this mating study, H53 and H98, were constructed lacking most of the original RM system genes in prior experiments [89, 90]. Studies by Ouelette (2019) suggest that frequency of recombination during mating was lower for strains with incompatible RM systems than those with compatible RM systems, although the effect could have been accounted for by the presence of an RM motif close to one of the marker genes [92]. In contrast, mating a strain harboring an intein with a homing endonuclease and a strain without the intein, resulted in increased recombination rates [93], revealing that DNA double strand breaks can lead to increased recombination during mating. All RM genes show increased transcription during mating conditions in comparison to planktonic cells with the only exception being an initial decrease at the 0-hour time point for the gene in the *rmeRMS* operon (consisting of the genes HVO_RS15605, HVO_RS20260, and HVO_RS20265) encoding the specificity unit (HVO_RS20265) (Fig. 6C). The transcription of the genes HVO_RS01890 and HVO_RS02905, predicted methyltransferases, are increased in response to mating, although HVO_RS02905 sees an increase to a lesser degree than that of HVO_RS01890. It has been shown computationally that these two genes were at one point likely a single RM system, but had been split due to genome rearrangements [89]. A latter study showed that all methylation within *Hfx. volcanii* is mediated by the gene HVO_RS08510 and the operon *rmeRMS*. HVO_RS08510 is an orphan methyltransferase and is responsible for the methylation of a CTAG motif [90], which is widely distributed amongst the Haloarchaea [91]. This gene shows some inconsistent variation of transcription in mating cells as compared to planktonic cells, but none of the changes observed are significant and expression values are low (Table S3). The *rmeRMS* operon was shown to be responsible for all the adenine methylation in *Hfx. volcanii* [90], and it encodes a type I restriction-methylation system. The transcription of all three genes showed an increase after the 0-hour time point of mating. The genes HVO_RS02230 and HVO_RS20385, were identified as methyltransferase homologs, but were not functionally active [90]. Both these genes also showed an increase in transcription in mated cells as compared to planktonic cells, although in the case of HVO_RS02230, this increase only appeared in the 4- and 8-hour time points. Overall, these results coupled with the study showing increased mating between *Haloferax* cells with compatible RM systems suggests that RM systems may play a role in limiting recombination after cell fusion.

Clustering

Beyond our hypothesized or known genes for mating, we were interested in identifying additional unknown genes that could also play a role in mating. To that end we applied a clustering analysis to group genes based on expression changes across the time points resulting in clusters of genes with shared expression patterns. Clustering analysis based on expression patterns resulted in twenty-two clusters with a range of 3 to 289 genes per cluster (Fig. S5). Those genes that increased from shaking or time 0 rapidly to the two-hour time-point and then decreased in expression after two hours show a pattern consistent with the timeline for mating (Fig. S5; Table S4). Clusters with continuous increases to twenty-four hours, such as the pattern seen in clusters 2 and 16, are less likely to be associated with a mating and more likely associated with cell growth. This is because cells are in log phase through the twenty-four hour time-point and mating is initiated predominately within the first two hours of plating (Fig. 1b,c). Cluster three contained the most genes of all clusters and also presented the putative mating pattern of gene expression.

Differential expression analysis on cluster three between the shaking control and time points 2-24 hours revealed a number of significant genes (Fig. S7). Most of the highly significant genes in cluster three in the differential expression analysis comparing shaking or t0 with the two- and four-hour time-points were ribosomal proteins or hypothetical proteins (Table S4). Cluster three also contained a number of operon genes. Notably, co-located genes linked to glycosylation were identified – HVO_RS20145, HVO_RS20160, and HVO_RS20165 (Fig. S8). These genes are *aglP*, a SAM-dependent methyltransferase, *aglR*, a MATE family

membrane protein, and *aglS/arnT*, a glycosylation protein, respectively. Cluster twelve, which has a similar expression pattern to cluster three but peaking at four-hours instead of two, contained additional genes in this location, HVO_RS20150 (*aglQ*, a glycosol hydrolase) and HVO_RS20155 (*aglI*, a glycosyl transferase).

This analysis suggests a large number of potential targets for further examination. In addition to laboratory based knockout experiments for genes of known or presumed function, there is a gap in functional annotation that would benefit from computational examination.

4. Summary and conclusions

This work represents an overview of the gene expression changes between unmated *Hfx. volcanii* and cells grown under mating conditions. It has raised many avenues for further work into genes controlling mating behavior, and likewise has suggested genes which may have a role in biofilm formation. One of the more remarkable observations is that the process of mating begins rapidly in *Hfx. volcanii*. While prior published mating experiments on filters include incubation times of 16 to 48 hours to allow for mating [15, 18, 19, 48, 49], here we show that cells initiate mating immediately after contact. Mating efficiency appears to peak between 4 to 8 hours after initiation, with very little difference between those two time points. Future experiments involving mating may be better streamlined in light of this.

Glycosylation has been previously examined for its connection to mating and is likely driving mating through s-layer decorating that serves to attract cells to one another. Disruption of glycosylation genes has resulted in reduction of mating [19], but no other genes have been distinctly linked to mating to date. Glycosylation genes were examined and followed a pattern consistent with the mating efficiency over time. Cells cultured in low-salt were also found to decrease mating efficiency [19] and the patterns of expression of many low-salt glycosylation genes were consistent with mating, though some were more erratically expressed. Multiple glycosylation genes were identified in this study for future examination on their impact on mating, including *aglF*, *aglG*, and *aglI*.

Glycosylation of the cell wall influences mating as the cells move closer together, a condition most often seen in biofilm formation and community development. Indeed, cells are observed to mate in biofilms at a significantly higher rate than in the planktonic state [16]. The genes involved in construction of the *Hfx. volcanii* archaella and chemotaxis did not show a uniform pattern of expression, but that may indicate that the cells already have their archaella from the planktonic state and it may be used to move cells closer in preparation for mating during the early sessile conditions, as seen in bacterial biofilm formation [94]. Nevertheless, one previous study showed that disruption of the archaellin had no effect on mating efficiency [49]. Additionally, genes associated with biofilm formation, namely adhesion, showed no effect on mating efficiency. Previously published gene deletion data in conjunction with our findings suggests that the genes associated with biofilm formation and community development are not driving mating.

The physiology of mating in *Hfx. volcanii* involves cellular fusion via the formation of a cytoplasmic bridge followed by bidirectional transfer of DNA and ends with cell cleavage. SNARE/DedA family proteins in Haloarchaea do not have a clear transcriptional pattern to connect them to the mating process, but the mating pattern is not in conflict with that hypothesis. The balance of DedA and SNARE homologs may be mediating the fusion required for mating and are considered a primary target for future work. Additionally, genes implicated in mediating cell shape (the CetZ genes) may potentially be involved in formation of the cytoplasmic bridge and many of those genes have a pattern of expression consistent with mating warranting further examination.

This work also identified significantly increased expression under a mating condition for selfish genetic elements and host defense genes. The increased expression of insertion sequence transposases indicates that mating conditions may be activating transposition. Simultaneously, expression of the host defense, CRISPR-Cas and RM Systems, are also activated under mating conditions, likely limiting recombination and defending against "unwanted", invading DNA, and suggesting post-fusion barriers against mating are present in *Hfx. volcanii*.

This study has identified a number of genes in addition to those discussed that follow an overall pattern of expression consistent with mating efficiency observations. Many of those genes are of unknown function and would benefit from further computational and experimental analysis to identify potential links to mating or to other important aspects of *Haloferax volcanii* molecular processes. Work on this intriguing method of DNA transfer remains ongoing.

Acknowledgements

The Computational Biology Core, Institute for Systems Genomics, University of Connecticut provided computational resources.

This work was supported through grants from the Binational Science Foundation (BSF 2013061); the National Science Foundation (NSF/MCB 1716046) within the BSF-NSF joint research program and NASA Exobiology (NNX15AM09G, and 80NSSC18K1533).

Author contributions

RTP and AMM designed experiments and JPG and ASL designed computational analysis. AMM carried out experiments. RTP, JPG, AMM, ASL, NRM, and UG interpreted the analytical output. AMM, ASL, RTP, and JPG wrote the manuscript. ASL prepared the figures. RTP, JPG, AMM, ASL, NRM, and UG reviewed and approved the manuscript.

Competing Interests

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Figure 1: (A) Experimental overview (repeated for two biological replicates): auxotrophic cells were combined and filter plated for mating in casamino acids (CA) medium with uracil (ura), tryptophan (trp), and both thymidine and hypoxanthine (thy); two filters were used for sequencing and one for dilution plating on selective and non-selective media to calculate mean colony forming units (cfu) and mating efficiency; (B) Growth curve of *Haloferax volcanii* plotting mean cfu on $\log_{10}$ scale (standard error bars based on triplicate plates for two replicates); (C) mating efficiency based on triplicate plates for two replicates excluding shaking and time 0, linear model regression fit calculated for time 2 to 24 h and comparison tested with ANOVA; and (D) principle component analysis of mating transcriptome time-points identified by replicate pairs and ANOVA tested on PC1 for time.

Figure 2: Bar plot depiction of differential expression compared to planktonic state for (A) glycosylation genes and (B) low-salt glycosylation genes; gene id on the x-axis and log$_2$ fold change on the y-axis, *p-value <0.05 and **p-value<0.01.

Figure 3. Bar plot depiction of differential expression compared to planktonic state for genes involved in community development and biofilm formation; gene id on the y-axis and log2 fold change on the x-axis, *p-value <0.05 and **p-value<0.01.

Figure 4. Bar plot depiction of differential expression compared to planktonic state for SNARE/*dedA* motif genes; gene id on the x-axis and log₂ fold change on the y-axis, *p-value <0.05 and **p-value<0.01.

Figure 5: Bar plot depiction of differential expression compared to planktonic state for FitZ and CetZ genes; gene id on the x-axis and log$_2$ fold change on the y-axis, *p-value <0.05 and **p-value<0.01.

Figure 6: (A) Manhattan plot with chromosome number on the x-axis and -log10 p-value of differential expression for planktonic cells vs plated timepoints on the y-axis, all genes are plotted in grey and genes for insertion sequences are highlighted by color according to the key; (B) Violin plot showing $\log_{10}$ of average normalized expression values for each treatment on the y-axis; (C) Bar plot depiction of differential expression compared to planktonic state for restriction modification system genes; gene id on the x-axis and $\log_2$ fold change on the y-axis, *p-value <0.05 and **p-value<0.01.

Figure S1: Bar plot depiction of differential expression compared to planktonic state for type IV secretion system genes; gene id on the x-axis and log$_2$ fold change on the y-axis, *p-value <0.05 and **p-value<0.01.

Figure S2: Phylogeny of homologs to HVO_RS15650, conjugal transfer protein, across both Archaeal and Bacterial domains.

Figure S3. Bar plot depiction of differential expression compared to planktonic state for insertion sequence gene families; gene id on the y-axis and log₂ fold change on the x-axis.

Figure S4. (A) Manhattan plot with chromosome number on the x-axis and -log10 p-value of differential expression for planktonic cells vs plated timepoints on the y-axis, all genes are plotted in grey and genes for CRISPR-Cas are highlighted by color according to the key; (B) Bar plot depiction of differential expression compared to planktonic state for CRISPR-Cas genes; gene id on the x-axis and log2 fold change on the y-axis, *p-value <0.05 and **p-value<0.01.

Figure S5. Clustering of all expressed *Haloferax volcanii* genes for identification of similar expression patterns; time is on the x-axis and expression values scaled to the mean are on the y-axis.

Figure S6. Bar plot depiction of differential expression compared to planktonic state for top and bottom ten genes for each time-point; gene id on the x-axis, $\log_2$ fold change on the y-axis, and titles indicating the time-point and top or bottom; *p-value <0.05 and **p-value<0.01.

Figure S7. Volcano plots of differential expression for the genes in cluster three comparing shaking to each time-point as labeled; x-axis is $\log_2$ fold change, y-axis is $-\log_{10}$ p-value, orange lines mark $\log_2$ fold change of -1 and 1, and green line marks a p-value of 0.01.

Figure S8. Gene map of co-located glycosylation genes.

Table 1: Sample information including experimental parameters, sequencing and assembly information

Sample ID	Treatment[a]	Raw Pairs[b]	Trimmed Pairs[c]	%remaining	Bowtie2 alignment rate[d]	No. D.E. genes[e]
shaking	planktonic	3,169,158	2,918,605	92.09%	91.50%	n/a
t00	0 hours	3,256,394	2,805,158	86.14%	91.61%	3,320 (109)
t02	2 hours	3,036,028	2,789,144	91.87%	94.17%	3,434 (941)
t04	4 hours	3,155,477	2,900,100	91.91%	93.50%	3,480 (1092)
t08	8 hours	3,054,785	2,756,910	90.25%	94.08%	3,472 (757)
t24	24 hours	3,136,767	2,873,436	91.61%	93.27%	3,462 (643)

[a]Shaking represents cells suspended and not mating; each hour following is time on filter before sampling (see methods for details)

[b]Paired end sequencing (2x75) using Illumina MiSeq summed for two replicates

[c]Trimmed using Sickle v1.33 summed for two replicates

[d]Trimmed reads were aligned back to the *Haloferax volcanii* DS2 genome (NCBI assembly: ASM2568v1); average across two replicates

[e]Total number of differentially expressed genes for each time point compared to shaking control and genes with p-value < 0.01 in parenthesis

Time	Replicate[a]	Average non-selective cfu	Selective media	Mating efficiency
		Table S1. Growth and mating efficiency colony counts.		
shaking	pelleted A	3.540E+08	0	0
shaking	pelleted A	3.540E+08	0	0
shaking	pelleted A	3.540E+08	0	0
shaking	pelleted B	3.437E+08	0	0
shaking	pelleted B	3.437E+08	0	0
shaking	pelleted B	3.437E+08	0	0
0h	filtered A	1.557E+07	6.400E+01	4.111E-06
0h	filtered A	1.557E+07	5.100E+01	3.276E-06
0h	filtered A	1.557E+07	7.500E+01	4.818E-06
0h	filtered B	1.250E+07	1.900E+01	1.520E-06
0h	filtered B	1.250E+07	2.800E+01	2.240E-06
0h	filtered B	1.250E+07	1.100E+01	8.800E-07
2h	filtered A	9.267E+06	1.950E+02	2.104E-05
2h	filtered A	9.267E+06	1.510E+02	1.629E-05
2h	filtered A	9.267E+06	1.500E+02	1.619E-05
2h	filtered B	4.933E+06	8.200E+01	1.662E-05
2h	filtered B	4.933E+06	1.010E+02	2.047E-05
2h	filtered B	4.933E+06	9.700E+01	1.966E-05
4h	filtered A	2.643E+07	3.600E+03	1.362E-04
4h	filtered A	2.643E+07	5.100E+03	1.929E-04
4h	filtered A	2.643E+07	3.600E+03	1.362E-04
4h	filtered B	2.240E+07	1.000E+01	4.464E-07
4h	filtered B	2.240E+07	2.000E+01	8.929E-07
4h	filtered B	2.240E+07	3.000E+01	1.339E-06
8h	filtered A	1.313E+07	3.000E+02	2.284E-05
8h	filtered A	1.313E+07	5.000E+02	3.807E-05
8h	filtered A	1.313E+07	7.000E+02	5.330E-05
8h	filtered B	5.300E+07	1.500E+03	2.830E-05
8h	filtered B	5.300E+07	8.000E+02	1.509E-05
8h	filtered B	5.300E+07	5.000E+02	9.434E-06
24h	filtered A	8.067E+08	3.900E+04	4.835E-05
24h	filtered A	8.067E+08	2.700E+04	3.347E-05
24h	filtered A	8.067E+08	4.800E+04	5.950E-05
24h	filtered B	9.433E+08	2.600E+04	2.756E-05
24h	filtered B	9.433E+08	2.800E+04	2.968E-05
24h	filtered B	9.433E+08	3.700E+04	3.922E-05

[a]Filtered indicates cells that have been applied to a filter and allowed to mate prior to being washed off the filter and plated. The step of washing cells off the filter results in incomplete recovery of cells and is reflected in the decrease of cfu/mL for these cells.

Table S2: Sample information including experimental parameters, sequencing and assembly information

Sample ID	Treatment[a]	Replicate	Raw Pairs[b]	trimmed_pairs[c]	%remaining	Bowtie2 alignment rate[d]	fwd_seq_name	rev_seq_name	SRA Accession[e]
Ap0	shaking	1	1,500,539	1,417,634	94.47%	91.86%	Ap0h_S1_L001_R1_001.fastq	Ap0h_S1_L001_R2_001.fastq	SRR11797459
A00	0 hours	1	1,584,563	1,442,125	91.01%	94.11%	A0h_S2_L001_R1_001.fastq	A0h_S2_L001_R2_001.fastq	SRR11797455
A02	2 hours	1	1,411,487	1,332,354	94.39%	94.92%	A2h_S3_L001_R1_001.fastq	A2h_S3_L001_R2_001.fastq	SRR11797453
A04	4 hours	1	1,637,874	1,483,767	90.59%	93.03%	A4h_S4_L001_R1_001.fastq	A4h_S4_L001_R2_001.fastq	SRR11797451
A08	8 hours	1	1,670,817	1,532,585	91.73%	94.06%	A8h_S5_L001_R1_001.fastq	A8h_S5_L001_R2_001.fastq	SRR11797449
A24	24 hours	1	1,654,754	1,507,569	91.11%	92.81%	A24h_S6_L001_R1_001.fastq	A24h_S6_L001_R2_001.fastq	SRR11797457
Bp0	shaking	2	1,668,619	1,500,971	89.95%	91.14%	Bp0h_S7_L001_R1_001.fastq	Bp0h_S7_L001_R2_001.fastq	SRR11797458
B00	0 hours	2	1,671,831	1,363,033	81.53%	89.10%	B0h_S8_L001_R1_001.fastq	B0h_S8_L001_R2_001.fastq	SRR11797454
B02	2 hours	2	1,624,541	1,456,790	89.67%	93.42%	B2h_S9_L001_R1_001.fastq	B2h_S9_L001_R2_001.fastq	SRR11797452
B04	4 hours	2	1,517,603	1,416,333	93.33%	93.96%	B4h_S10_L001_R1_001.fastq	B4h_S10_L001_R2_001.fastq	SRR11797450
B08	8 hours	2	1,383,968	1,224,325	88.46%	94.10%	B8h_S11_L001_R1_001.fastq	B8h_S11_L001_R2_001.fastq	SRR11797448
B24	24 hours	2	1,482,013	1,365,867	92.16%	93.72%	B24h_S12_L001_R1_001.fastq	B24h_S12_L001_R2_001.fastq	SRR11797456

[a]Shaking represents cells suspended and not mating; each hour following is time on filter before sampling (see methods for details)

[b]Paired end sequencing (2x75) using Illumina MiSeq

[c]Trimmed using Sickle v1.33

[d]Trimmed reads were aligned back to the *Haloferax volcanii* DS2 genome (NCBI assembly: ASM2568v1)

[e]All sequences available on NCBI under BioProject PRJNA632928

Tables S3 and S4 too large to insert – see attached supplemental

References

1. Thomas, C. M. & Nielsen, K. M. Mechanisms of, and barriers to, horizontal gene transfer between bacteria. *Nature reviews.Microbiology* **3**, 711 (2005).

2. Goodenough, U. & Heitman, J. Origins of eukaryotic sexual reproduction. *Cold Spring Harbor perspectives in biology* **6**, a016154 (2014).

3. Soucy, S. M., Huang, J. & Gogarten, J. P. Horizontal gene transfer: building the web of life *Nat. Rev. Genet.* **16**, 472-482 (2015).

4. Bertani, G. & Baresi, L. Genetic transformation in the methanogen Methanococcus voltae PS *J. Bacteriol.* **169**, 2730-2738 (1987).

5. Worrell, V. E., Nagle, D. P., McCarthy, D. & Eisenbraun, A. Genetic transformation system in the archaebacterium Methanobacterium thermoautotrophicum Marburg *J. Bacteriol.* **170**, 653-656 (1988).

6. Sato, T., Fukui, T., Atomi, H. & Imanaka, T. Improved and versatile transformation system allowing multiple genetic manipulations of the hyperthermophilic archaeon Thermococcus kodakaraensis *Appl. Environ. Microbiol.* **71**, 3889-3899 (2005).

7. Waege, I., Schmid, G., Thumann, S., Thomm, M. & Hausner, W. Shuttle vector-based transformation system for Pyrococcus furiosus *Appl. Environ. Microbiol.* **76**, 3308-3313 (2010).

8. Johnsborg, O., Eldholm, V. & Håvarstein, L. S. Natural genetic transformation: prevalence, mechanisms and function *Res. Microbiol.* **158**, 767-778 (2007).

9. Johnston, C., Martin, B., Fichant, G., Polard, P. & Claverys, J. P. Bacterial transformation: distribution, shared mechanisms and divergent control *Nat. Rev. Microbiol.* **12**, 181-196 (2014).

10. van Wolferen, M., Wagner, A., van der Does, C. & Albers, S. V. The archaeal Ced system imports DNA *Proc. Natl. Acad. Sci. U. S. A.* **113**, 2496-2501 (2016).

11. Mevarech, M. & Werczberger, R. Genetic transfer in Halobacterium volcanii *J. Bacteriol.* **162**, 461-462 (1985).

12. Torreblanca, M. *et al.* Classification of Non-alkaliphilic Halobacteria Based on Numerical Taxonomy and Polar Lipid Composition, and Description of *Haloarcula* gen. nov. and *Haloferax* gen. nov. *Systematic and Applied Microbiology* **8**, 89-99 (1986).

13. Rosenshine, I., Tchelet, R. & Mevarech, M. The mechanism of DNA transfer in the mating system of an archaebacterium *Science* **245**, 1387-1389 (1989).

14. Ortenberg, R., Tchelet, R. & Mevarech, M. in *Microbiology and biogeochemistry of hypersaline environments* (ed Oren, A.) 331-338 (CRC Press, Boca Raton, FL, 1998).

15. Naor, A., Lapierre, P., Mevarech, M., Papke, R. T. & Gophna, U. Low species barriers in halophilic archaea and the formation of recombinant hybrids *Curr. Biol.* **22**, 1444-1448 (2012).

16. Chimileski, S., Franklin, M. J. & Papke, R. T. Biofilms formed by the archaeon Haloferax volcanii exhibit cellular differentiation and social motility, and facilitate horizontal gene transfer *BMC Biol.* **12**, 65-014 (2014).

17. Tchelet, R. & Mevarech, M. Interspecies genetic transfer in halophilic archaebacteria. *Systematic and Applied Microbiology* **16**, 578-581 (1993).

18. Abdul Halim, M. F. *et al.* Haloferax volcanii archaeosortase is required for motility, mating, and C-terminal processing of the S-layer glycoprotein *Mol. Microbiol.* **88**, 1164-1175 (2013).

19. Shalev, Y., Turgeman-Grott, I., Tamir, A., Eichler, J. & Gophna, U. Cell Surface Glycosylation Is Required for Efficient Mating of Haloferax volcanii *Front. Microbiol.* **8**, 1253 (2017).

20. Guan, Z., Naparstek, S., Calo, D. & Eichler, J. Protein glycosylation as an adaptive response in Archaea: growth at different salt concentrations leads to alterations in Haloferax volcanii S-layer glycoprote in N-glycosylation. *Environmental microbiology* **14**, 743-753 (2012).

21. Shalev, Y. *et al.* Comparative Analysis of Surface Layer Glycoproteins and Genes Involved in Protein Glycosylation in the Genus Haloferax *Genes (Basel)* **9**, 10.3390/genes9030172 (2018).

22. Williams, D., Gogarten, J. P. & Papke, R. T. Quantifying homologous replacement of loci between haloarchaeal species *Genome Biol. Evol.* **4**, 1223-1244 (2012).

23. Ng, W. V. *et al*. Genome sequence of Halobacterium species NRC-1 *Proc. Natl. Acad. Sci. U. S. A.* **97**, 12176-12181 (2000).

24. Koonin, E. V., Makarova, K. S. & Aravind, L. Horizantal gene transfer in porkaryotes: quantification and classification. *Annu.Rev.Microbiol.* **55**, 709-742 (2001).

25. Meheust, R. *et al*. Hundreds of novel composite genes and chimeric genes with bacterial origins contributed to haloarchaeal evolution *Genome Biol.* **19**, 75-018 (2018).

26. Allers, T., Ngo, H. P., Mevarech, M. & Lloyd, R. G. Development of additional selectable markers for the halophilic archaeon Haloferax volcanii based on the leuB and trpA genes *Appl. Environ. Microbiol.* **70**, 943-953 (2004).

27. https://github.com/Gogarten-Lab/hvo_mating.

28. https://github.com/ucdavis-bioinformatics/scythe.

29. Joshi, N. A. & Fass, J. N. No title. *Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files (Version 1.33)[Software]* (2011).

30. Patro, R., Duggal, G. & Kingsford, C. Salmon: accurate, versatile and ultrafast quantification from RNA-seq data using lightweight-alignment. *BioRxiv*, 021592 (2015).

31. UniProt Consortium. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res.* **47**, D506-D515 (2019).

32. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome *Genome Biol.* **10**, R25 (2009).

33. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2 *Nat. Methods* **9**, 357-359 (2012).

34. Langmead, B., Wilks, C., Antonescu, V. & Charles, R. Scaling read aligners to hundreds of threads on general-purpose processors *Bioinformatics* **35**, 421-432 (2019).

35. Tarazona, S., García, F., Ferrer, A., Dopazo, J. & Conesa, A. NOIseq: a RNA-seq differential expression method robust for sequencing depth biases. *EMBnet.journal* **17**, 18-19 (2011).

36. Tarazona, S. *et al*. Data quality aware analysis of differential expression in RNA-seq with NOISeq R/Bioc package. *Nucleic Acids Res.* **43**, e140 (2015).

37. https://rdrr.io/cran/manhattanly/.

38. Wickham, H. & Chang, W. ggplot2: An implementation of the Grammar of Graphics. *R package version 0.7, URL: http://CRAN.R-project.org/package=* ggplot2 **3** (2008).

39. Wickham, H. ggplot2. *Wiley Interdisciplinary Reviews: Computational Statistics* **3**, 180-185 (2011).

40. Wickham, H., Chang, W. & Wickham, M. H. Package 'ggplot2'. *Create Elegant Data Visualisations Using the Grammar of Graphics.Version* **2**, 1-189 (2016).

41. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403-410 (1990).

42. Grishin, N. V. Estimation of the number of amino acid substitutions per site when the substitution rate varies among sites. *Journal of molecular evolution* **41**, 675-679 (1995).

43. Desper, R. & Gascuel, O. Theoretical foundation of the balanced minimum evolution method of phylogenetic inference and its relationship to weighted least-squares tree fitting. *Moleclar Biology and Evolution* **21**, 587-598 (2004).

44. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2 *Genome Biol.* **15**, 550 (2014).

45. Wickham, H. & Henry, L. Tidyr: Easily tidy data with'spread ()'and'gather ()'functions. *R package version 0.6* **1** (2017).

46. Wickham, H., Francois, R., Henry, L. & Müller, K. dplyr: A grammar of data manipulation. *R package version 0.4* **3** (2015).

47. Wickham, H. reshape2: Flexibly reshape data: a reboot of the reshape package. *R package version* **1** (2012).

48. Turgeman-Grott, I. *et al.* Pervasive acquisition of CRISPR memory driven by inter-species mating of archaea can limit gene transfer and influence speciation *Nat. Microbiol.* **4**, 177-186 (2019).

49. Tripepi, M., Imam, S. & Pohlschroder, M. Haloferax volcanii flagella are required for motility but are not involved in PibD-dependent surface adhesion *J. Bacteriol.* **192**, 3093-3102 (2010).

50. Robinson, J. L. *et al.* Growth kinetics of extremely halophilic archaea (family halobacteriaceae) as revealed by arrhenius plots *J. Bacteriol.* **187**, 923-929 (2005).

51. Dyall-Smith, M. "The halohandbook." *Protocols for haloarchaeal genetics. .* (2009).

52. Frols, S., Dyall-Smith, M. & Pfeifer, F. Biofilm formation by haloarchaea *Environ. Microbiol.* **14**, 3159-3174 (2012).

53. Naor, A. & Gophna, U. Cell fusion and hybrids in Archaea: prospects for genome shuffling and accelerated strain development for biotechnology *Bioengineered* **4**, 126-129 (2013).

54. Jarrell, K. F. & Albers, S. V. The archaellum: an old motility structure with a new name *Trends Microbiol.* **20**, 307-312 (2012).

55. Albers, S. V. & Jarrell, K. F. The Archaellum: An Update on the Unique Archaeal Motility Structure *Trends Microbiol.* **26**, 351-362 (2018).

56. Thomas, N. A., Bardy, S. L. & Jarrell, K. F. The archaeal flagellum: a different kind of prokaryotic motility structure. *FEMS Microbiol. Rev.* **25**, 147-174 (2001).

57. Ng, S. Y., Chaban, B. & Jarrell, K. F. Archaeal flagella, bacterial flagella and type IV pili: a comparison of genes and posttranslational modifications *J. Mol. Microbiol. Biotechnol.* **11**, 167-191 (2006).

58. Trachtenberg, S. & Cohen-Krausz, S. The archaeabacterial flagellar filament: a bacterial propeller with a pilus-like structure *J. Mol. Microbiol. Biotechnol.* **11**, 208-220 (2006).

59. Näther, D. J., Rachel, R., Wanner, G. & Wirth, R. Flagella of Pyrococcus furiosus: multifunctional organelles, made for swimming, adhesion to various surfaces, and cell-cell contacts *J. Bacteriol.* **188**, 6915-6923 (2006).

60. Zolghadr, B. *et al.* Appendage-mediated surface adherence of Sulfolobus solfataricus *J. Bacteriol.* **192**, 104-110 (2010).

61. Albers, S. V. & Pohlschroder, M. Diversity of archaeal type IV pilin-like structures. *Extremophiles* **13**, 403-410 (2009).

62. Liu, J. D. & Parkinson, J. S. Role of CheW protein in coupling membrane receptors to the intracellular signallingsystem of bacterial chemotaxis. *Proc.Natl.Acad.Sci.U.S.A.* **86**, 8703-8707 (1989).

63. Yonekawa, H., Hayashi, H. & Parkinson, J. S. Rewuirement of the cheB function for sensory adaptation in *Escherichia coli. Journal of Bacteriology* **156**, 1228-1235 (1983).

64. Guttenplan, S. B. & Kearns, D. B. Regulation of flagellar motility during biofilm formation. *FEMS Microbiol.Rev.* **37**, 849-871 (2013).

65. Liao, Y. *et al.* Morphological and proteomic analysis of biofilms from the Antarctic archaeon, Halorubrum lacusprofundi *Sci. Rep.* **6**, 37454 (2016).

66. Taglialegna, A., Lasa, I. & Valle, J. Amyloid structures as biofilm matrix scaffolds. *J. Bacteriol.* **198**, 2579-2588 (2016).

67. van Wolferen, M., Orell, A. & Albers, S. V. Archaeal biofilm formation *Nat. Rev. Microbiol.* **16**, 699-713 (2018).

68. Grohmann, E., Christie, P. J., Waksman, G. & Backert, S. Type IV secretion in Gram-negative and Gram-positive bacteria *Mol. Microbiol.* **107**, 455-471 (2018).

69. Jahn, R. & Scheller, R. H. SNAREs--engines for membrane fusion *Nat. Rev. Mol. Cell Biol.* **7**, 631-643 (2006).

70. Doerrler, W. T., Sikdar, R., Kumar, S. & Boughner, L. A. New functions for the ancient DedA membrane protein family *J. Bacteriol.* **195**, 3-11 (2013).

71. Yoon, T. Y. & Munson, M. SNARE complex assembly and disassembly *Curr. Biol.* **28**, R397-R401 (2018).

72. Bennett, M. K. & Scheller, R. H. The molecular machinery for secretion is conserved from yeast to neurons *Proc. Natl. Acad. Sci. U. S. A.* **90**, 2559-2563 (1993).

73. Inadome, H., Noda, Y., Kamimura, Y., Adachi, H. & Yoda, K. Tvp38, Tvp23, Tvp18 and Tvp15: novel membrane proteins in the Tlg2-containing Golgi/endosome compartments of Saccharomyces cerevisiae *Exp. Cell Res.* **313**, 688-697 (2007).

74. Liang, F. T. *et al.* BB0250 of Borrelia burgdorferi is a conserved and essential inner membrane protein required for cell division *J. Bacteriol.* **192**, 6105-6115 (2010).

75. Keller, R. & Schneider, D. Homologs of the yeast Tvp38 vesicle-associated protein are conserved in chloroplasts and cyanobacteria *Front. Plant. Sci.* **4**, 467 (2013).

76. Sikdar, R. & Doerrler, W. T. Inefficient Tat-dependent export of periplasmic amidases in an Escherichia coli strain with mutations in two DedA family genes *J. Bacteriol.* **192**, 807-818 (2010).

77. Lukacs, G. L. *et al.* Size-dependent DNA mobility in cytoplasm and nucleus *J. Biol. Chem.* **275**, 1625-1629 (2000).

78. Bai, H., Lester, G. M. S., Petishnok, L. C. & Dean, D. A. Cytoplasmic transport and nuclear import of plasmid DNA *Biosci. Rep.* **37**, BSR20160616 (2017).

79. Duggin, I. G. *et al.* CetZ tubulin-like proteins control archaeal cell shape *Nature* **519**, 362-365 (2015).

80. Bi, E. F. & Lutkenhaus, J. FtsZ ring structure associated with division in Escherichia coli *Nature* **354**, 161-164 (1991).

81. Wang, X. & Lutkenhaus, J. FtsZ ring: the eubacterial division apparatus conserved in archaebacteria *Mol. Microbiol.* **21**, 313-319 (1996).

82. Kazumichi, O., Yatsunami, R. & Nakamura, S. Cloning and sequencing of ftsZ homolog from extremely halophilic archaeon Haloarcula japonica strain TR-1 *Nucleic Acids Symp. Ser.* **(44)**, 155-156 (2000).

83. Poplawski, A., Gullbrand, B. & Bernander, R. The ftsZ gene of Haloferax mediterranei: sequence, conserved gene order, and visualization of the FtsZ ring *Gene* **242**, 357-367 (2000).

84. Aylett, C. H. S. & Duggin, I. G. The Tubulin Superfamily in Archaea *Subcell. Biochem.* **84**, 393-417 (2017).

85. Hartman, A. L. *et al.* The complete genome sequence of Haloferax volcanii DS2, a model archaeon *PLoS One* **5**, e9605 (2010).

86. Naor, A., Lazary, R., Barzel, A., Papke, R. T. & Gophna, U. In vivo characterization of the homing endonuclease within the polB gene in the halophilic archaeon Haloferax volcanii *PLoS One* **6**, e15833 (2011).

87. Artieri, C. G. *et al.* Cis-regulatory evolution in prokaryotes revealed by interspecific archaeal hybrids *Sci. Rep.* **7**, 3986-017 (2017).

88. Maier, L. -. *et al.* The nuts and bolts of the Haloferax CRISPR-Cas system I-B *RNA Biol.* **16**, 469-480 (2019).

89. Ouellette, M., Jackson, L., Chimileski, S. & Papke, R. T. Genome-wide DNA methylation analysis of Haloferax volcanii H26 and identification of DNA methyltransferase related PD-(D/E)XK nuclease family protein HVO_A0006 *Front. Microbiol.* **6**, 251 (2015).

90. Ouellette, M., Gogarten, J. P., Lajoie, J., Makkay, A. M. & Papke, R. T. Characterizing the DNA Methyltransferases of Haloferax volcanii via Bioinformatics, Gene Deletion, and SMRT Sequencing *Genes (Basel)* **9**, 129 (2018).

91. Fullmer, M. S., Ouellette, M., Louyakis, A. S., Papke, R. T. & Gogarten, J. P. The Patchy Distribution of Restriction(-)Modification System Genes and the Conservation of Orphan Methyltransferases in Halobacteria *Genes (Basel)* **10**, 233 (2019).

92. Ouellette, M. Characterizing Restriction-Modification Systems and DNA Methyltransferases in the Halobacteria. *Doctoral Dissertations*. 2280. https://opencommons.uconn.edu/dissertations/2280. (2019).

93. Naor, A. *et al.* Impact of a homing intein on recombination frequency and organismal fitness *Proc. Natl. Acad. Sci. U. S. A.* **113**, E4654-61 (2016).

94. O'Toole, G., Kaplan, H. B. & Kolter, R. Biofilm formation as microbial development. *Annu.Rev.Microbiol.* **54**, 49-79 (2000).