Neutrophil extracellular traps impair regeneration.

Authors:

Eric Wier[1], Mayumi Asada[1], Gaofeng Wang[1,2], Martin P. Alphonse[1], Ang Li[1], Chase Hintelmann[1],

Christine Youn[1], Brittany Pielstick[3], Roger Ortines[1], Lloyd S. Miller[1,4], Nathan K. Archer[1], Luis A.

Garza[1]

Affiliations:

[1]Department of Dermatology, Johns Hopkins University School of Medicine, Baltimore,

Maryland, USA.

[2]Department of Plastic and Aesthetic Surgery, Nanfang Hospital of Southern Medical University,

Guangzhou, Guangdong Province, China.

[3]Department of Molecular Biology and Genetics, Johns Hopkins University, Baltimore,

Maryland, USA.

[4]Immunology, Janssen Research and Development, 1400 McKean Road, Spring House, PA,

19477, USA (Current affiliation. All work performed at prior affiliation[1])

* Correspondence to: LAG@jhmi.edu

Abstract

Fibrosis is a major health burden across diseases and organs. To remedy this, we study wound

induced hair follicle regeneration (WIHN) as a model of non-fibrotic healing that recapitulates

embryogenesis for de novo hair follicle morphogenesis after wounding. We have previously

demonstrated that TLR3 promotes WIHN through binding dsRNA, but the source of which is still

unclear. Here, we demonstrate that multiple distinct contexts of high WIHN all show a strong

neutrophil signature, and given the likelihood of nuclear dsRNA release during the production

of neutrophil extracellular trap (NETs), we hypothesized that mature neutrophils and NETs

might promote WIHN. Consistent with this, in addition to the presence of mature neutrophils

shortly after wounding, neutrophils remain within the wound after the barrier is reestablished,

where they produce extracellular traps (NETs) that likely release spliceosomal U1 dsRNA.

Contrary to our hypothesis, genetic models of neutrophil depletion show enhanced WIHN.

Pad4 null mice that are defective in NET production also augment WIHN. Finally, using single-

cell RNA sequencing, we identified a dramatic increase in mature neutrophils in the wound

beds of low regenerating Tlr3-/- mice. Taken together, these results demonstrate that although

mature neutrophils are stimulated by a common pro-regenerative cue, their presence and NETs

hinder WIHN.

Introduction

After suffering a wound, the body initiates a well-coordinated physiological process to

restore homeostasis and reestablish the barrier. This spontaneous process comprises four

discrete phases: hemostasis, inflammation, proliferation, and remodeling[1]. Although all of these
phases have been extensively studied, the molecular details as to why repair tends to result in
fibrotic scar tissue rather than complete regeneration have not been fully elucidated. This
tendency of wound repair to lead to fibrosis, and in some cases, hypertrophic scars, contributes
an enormous burden on human health[1-3]. Inflammation and different components of the
immune system have been shown to contribute to regeneration in salamanders and zebrafish,
as well as to promote healing and maintain barrier function in mammals upon mucosal injury[4-6].
Although the cellular effects of macrophages and T cells are well studied[7-10] , the role of
neutrophils in modulating regeneration remains elusive.
Unlike urodele salamanders, complete skin regeneration after wounding in mammals is
rare[11]. It occurs via de novo hair follicle generation in mice and rabbits through a process that
mimics skin embryogenesis[12-15]. This process of de novo follicle neogenesis (Wound Induced
Hair Neogenesis; WIHN) was first fully characterized in mice after receiving full thickness
wounds[16]. These regenerated follicles establish a distinct stem cell population, express
characteristic differentiation markers, produce functional hair shafts, and can complete the hair
cycle. In addition to the hair follicles, sebaceous glands, specialized vascular and nerve
supports, and surrounding fat cells are regenerated[16,17].
Immediately after wounding, a robust inflammatory phase occurs, which allows the
ingress of keratinocytes and fibroblasts to proceed afterward. The early stage of wound healing
is defined by the dramatic recruitment of mature neutrophils, which are instrumental in
providing defense against microbial pathogens[18-22]. This is followed by an influx of macrophages
(Mϕ's) that continue the phagocytic processes begun by neutrophils and aid in the transition to
the proliferative phase of wound healing[23-25]. While much is known about the importance of
this inflammatory phase for preventing and abrogating infections, less is known about how it
influences regenerative capacity or WIHN. Increasingly, macrophages have been shown to be
essential for WIHN via TNF-induced AKT/ β-catenin signaling[8-10]. However, the role of
neutrophils in modeling regeneration or WIHN is unclear.
Toll-like receptors (TLRs) are highly conserved single-pass membrane-spanning
receptors that recognize structurally conserved molecular components of invading microbes
and activate a cascade of inflammatory signaling pathways[26]. Rather than simply recognizing
pathogen associated molecules, they can also initiate "sterile" inflammation upon recognizing
damage-associated molecular patterns (DAMPs), which are critical to recruit immune cells and
initiate wound healing[27]. TLR3 is activated by dsRNA and has primarily been studied in the
context of viral infection[28]. Mounting evidence shows that TLR3 also plays an important role in
wound repair[29-37]. Synthetic double strand RNA (dsRNA) polyriboinosinic-polyribocytidylic acid
(poly(I:C)) treatment dramatically increases WIHN in mice. Furthermore, wound-released
dsRNA activates TLR3 to promote hair follicle regeneration[34]. The identity and source of the
physiological RNA remains an open and important question in the field. Notably, the dsRNA U1
spliceosomal small nuclear RNA (snRNA), may be an important endogenous RNA sensed via
TLR3[32,33,38,39]. Specifically, UV damage releases U1 snRNA that stimulates cytokine production in
keratinocytes and increases barrier gene transcription[32,33].
In addition to phagocytosis and degranulation, mature neutrophils can produce
extracellular traps (ET), large extracellular web-like structures composed of decondensed
chromatin bound to various cytosolic and granule proteins[21,22,40-42]. While originally recognized

as a defense mechanism against pathogens[21,40,41], they have also been found to mediate sterile
inflammatory processes[43,44]. In the absence of infection, ETs can be stimulated in sterile tissue
environments through various cytokines[41,45-47] and by activated platelets[43,44]. Interestingly, ETs
are found within sterile wounds of mice and delay wound healing[48]. Mechanistically, ETs are
formed by the rapid decondensation of the cellular chromatin, followed by the fragmentation
of the nuclear membrane and mixing of the nuclear and cytoplasmic compartments, before
being expelled from the cell. The ability of neutrophils to rapidly migrate to the wound site and
produce ETs, coupled with the nuclear localization of some dsRNA, made us question whether
neutrophils were a source of the dsRNA critical for WIHN. Interestingly, while there are
extensive studies on the DNA components released during ET formation[22,42,49], the RNA
components are poorly understood.
To probe how mature neutrophils influence wound regeneration and WIHN, we
analyzed multiple microarrays from distinct contexts of high regenerating mice and found a
common neutrophil signature. Using immunofluorescence and flow cytometry we found that
neutrophils remain in the wound bed, albeit at low levels, after the acute inflammatory phase,
where they produce NETs that contain the nuclear U1 dsRNA. To define how this influences
regeneration, we used antibody mediated neutrophil depletion to eliminate neutrophils from
the wound bed at select time points, but discovered that the technique is ineffective in the
context of large full-thickness wounds, which are necessary to initiate WIHN. However, using a
neutrophil-specific diphtheria toxin ablation model, we were able to deplete mature
neutrophils in the wound bed and found that—contrary to our initial hypothesis-- the absence
of mature neutrophils enhances WIHN. Eliminating neutrophil's ability to produce NETs by
knocking out Pad4 also boosted WIHN, confirming the negative influence of neutrophils on
regeneration. Finally, we used single-cell RNA sequencing to characterized WIHN deficient Tlr3-
/- mice and found that they have a dramatically increased population of mature neutrophils in
the re-epithelized wound bed, compared to wild-type mice, likely contributing to their
diminished regenerative capacity. These results indicate that, while important for preventing
infection, mature neutrophils and their NETs negatively impact regeneration and WIHN.
Although a common pro-regenerative signal might increase neutrophil infiltration, mature
neutrophils instead likely contribute to fibrosis.
115
116   **Results**
117
118   **Neutrophil signature present during skin regeneration**
119
To begin to characterize the role of neutrophils in WIHN, we performed bioinformatic
analysis on previous microarrays of multiple distinct high regenerating mouse models, probing
for innate immune and neutrophil signatures. First, we analyzed the proteome comparing the
center of the re-epithelialized wound bed (high WIHN) to the surrounding periphery/edge (low
WIHN) (Fig. 1a)[37]. Gene ontology analysis revealed that, in the area of high WIHN, neutrophil
aggregation and other defense pathways against bacterium are enriched, characterized by an
abundance of antimicrobial and granular proteins, such as Neutrophil Elastase (Elane),
Cathelicidin (Camp), and Myeloperoxidase (Mpo) (Fig. 1b-c). In a second model system, high
WIHN Rnasel-/- mice (manuscript in review), at the time of scab detachment, are also enriched

in genes associated with neutrophils. Neutrophil chemotaxis is the most significant upregulated
gene ontology category, with other chemotactic and inflammatory pathways up as well (Fig.
1d). Finally, we analyzed the gene expression changes between wounded specific pathogen free
(SPF) mice, which have increased regeneration and WIHN, when compared to germ free (GF)
mice (manuscript in review), at the time of scab detachment. Like the other two high
regeneration models, when compared to GF, SPF mice had elevated neutrophil chemotaxis and
immune response transcripts (Fig. 1e). Together, these disparate experimental contexts
demonstrate that neutrophil chemotaxis can correlate with high regeneration and WIHN.
137
138 **Neutrophils persist in the wound bed after the acute inflammatory phase, producing**
139 **extracellular traps**
140
Given the above correlations of high neutrophil infiltration to high WIHN, we first
characterized mature neutrophil infiltration in the wound beds of C57BL/6J mice after a large
square wound cut into the center of their backs, to the depth of fascia. As anticipated,
neutrophils are abundant in the acute phase of the wound healing process (1-3 days post
wounding), as seen by H&E staining (Fig. 2a) and immunofluorescence of Myeloperoxidase
(MPO), a major neutrophil granule protein (Fig. 2b). While neutrophils predominate at early
time points, they are still present in the wound bed as late as wound day 11, well after re-
epithelization has completed (Fig. 2c). In contrast, macrophages only accumulate starting at
wound day 3 (Fig. 2b) but remain a major component of the wound even after re-epithelization
(Fig. S1). These results show the dynamic changes of immune cell infiltration, but with an
underappreciated persistence of some neutrophils late in wound healing, during the time of
morphogenesis.
Given the importance of dsRNA to promote WIHN, and the association of neutrophils
with high WIHN, we hypothesized that neutrophil extracellular traps (NETs) release of
neutrophil nuclear content might release dsRNA, in addition to dsDNA, from the nucleus to
promote WIHN. We visualized citrullinated histone H3 (H3Cit) to identify NETS as early as
wound day 3 (Fig. 2d). The production of NETs is mediated by Pad4, an enzyme that modifies
the arginine residues on histones to citrulline, which changes their charge, leading to massive
chromatin decondensation[50,51]. Consistent with this, NETs are virtually absent in mice lacking
Pad4, while abundant in the wound bed after the acute phase of wound healing in WT mice
(Fig. 2e). Although NETs are characterized by their extruded DNA, which forms a web-like
scaffold containing cytosolic and granular proteins, little is known about the RNA content within
them and how that might influence wound healing. Given that U1 small nuclear (sn) RNA is
proposed as a TLR3 agonist damage-associated molecular pattern important for skin barrier
repair[32,33], we visualized it by FISH. U1 snRNA is present throughout the wound beds of
C57BL/6J mice (Fig. 2f). Rather than the baseline nuclear appearance of unwounded tissue, U1
snRNA in the wound bed is markedly cytoplasmic, suggesting that its cellular localizations shifts
from the nucleus as a potential step towards cellular release. Consistent with this, we noted a
fine haze of signal extracellularly. Taken together, these data suggest a potential model where
neutrophils persist to the morphogenesis stage of wound healing and release extracellular traps
that contain U1 dsRNA to perhaps modulate WIHN.
172

Large full-thickness wounds dramatically reduce antibody-mediated neutrophil depletion

To functionally assess how neutrophils modulate wound induced hair neogenesis (WIHN), we sought to deplete neutrophils throughout the wound bed at select times. To accomplish this, we adapted a widely used antibody mediated depletion scheme, where we injected neutrophil specific antibodies intraperitoneally (Fig. 3a)[52]. Most experiments were completed using a Ly6G positive monoclonal antibody (1A8 clone), which recognizes a 21-25kD glycosylphosphatidylinositol (GPI)-linked differentiation antigen that is expressed by myeloid-derived cells. While monocytes transiently express Ly6G during bone marrow development, Ly6G expression in peripheral neutrophils directly correlates its level of differentiation and maturation. The less specific anti-granulocyte receptor-1 (Gr-1) antibody (RB6-8C5 clone), which recognizes both Ly6G and Ly6C and thereby also targets dendritic cells and subpopulations of lymphocytes and monocytes, was also used in select experiments. To prevent confounding flow cytometry measurements caused by epitope masking from the initial Ly6G antibody injected during the antibody mediated neutrophil depletion experiments[53], a distinct Ly6G antibody clone was used for analysis via flow cytometry. Injections were done one day before and one day after large full-thickness wounds on the back of mice. The depletion scheme focused on the early time points of the wound healing process because those were shown to be most critical in dsRNA-Tlr3 enhanced WIHN[34].

Unexpectedly, on wound day 3, after receiving two IP injections of either Ly6G or Gr-1 antibodies, using our careful detection strategy, there was no apparent neutrophil depletion in the wound bed, as measured via flow cytometry (Fig. 3b). Depletion was also ineffective in other tissues after wounding, including the spleen and liver (Fig. S3). Looking one day earlier (wound day 2), neutrophil depletion was still ineffective in the wound bed, but there was a statistically significant depletion in the blood of the same mice (Fig. 3c). Although significant, the neutrophil depletion was less than 50%, far less efficient than the previous reports[52,54,55]. An important distinction between ours and previous work is that we employed depletion in the context of large wounds to fascia, prompting us to question whether large area wounding globally modifies neutrophil blood trafficking dynamics to result in depletion failure. Indeed, neutrophil depletion was very efficient in the blood of unwounded mice who had received an IP injection of the Ly6G (1A8 clone) antibody 24 hours earlier (Fig. 3d), but this depletion was abolished after wounding the mice (Fig. 3e). To explore this further, we repeated the injection scheme in Fig. 3a. With and without wounding mice, we examined the neutrophil depletion in the blood and wound bed at wound day 2. Consistent with our previous experiments, in the context of wounding, the Ly6G (1A8) neutralizing antibody achieved less than 50 percent neutrophil depletion in the blood. In contrast, mice that were not wounded, achieved substantially more efficient depletion in the blood (Fig. 3f). Unwounded skin does not typically contain neutrophils, so the Ly6G (1A8 clone) antibody injection had no effect (Fig. 3g). Wounding caused dramatic recruitment of neutrophils to the wound bed, as shown earlier (Fig 3b-c, e). Ly6G antibody injection could not overcome this to deplete neutrophils from the wound bed (Fig. 3g).

A previous report had success with antibody-mediated neutrophil depletion in a wounding model where small 2cm excisions were made[55], leading us to question whether Ly6G

(1A8 clone) antibody injection was more effective in smaller, full-thickness wounds. To address this, mice were injected with Ly6G (1A8 clone) antibody and then wounded with large wounds (1.25x1.25cm2), like the previous experiments, or smaller wounds (0.6x0.6cm2) (Fig. 3h). Neutrophil depletion efficacy was then assessed at wound day two in the blood and wound bed. Mice that had been given small wounds had significant neutrophil depletion in both the blood and wound bed, while those with the large wounds did not (Fig 3i-j). Taken together, these data suggest that antibody mediated neutrophil depletion, while efficient in certain non-wounding contexts, is not effective after the large full-thickness wounds necessary to induce WIHN—perhaps because of the unusually robust stimuli of neutrophil recruitment in this context.

Mature Neutrophils inhibit wound induced hair neogenesis

Despite the minimal inhibition of neutrophils using conventional antibody depletion with large wounds, we sought to begin testing how neutrophils might modulate WIHN. We used the same antibody depletion scheme described (Fig 3a) but allowed the mice to recover for 21 days before quantifying WIHN via confocal scanning laser microscopy (CSLM) (Fig 4a)[56]. Antibody injection did not affect wound closure speed (Fig. S4). Despite minimal neutrophil depletion within the wound beds of mice post Ly6G (1A8 clone) injection (Fig 3b, c, e, g), the number of regenerated hair follicles was increased, suggesting that neutrophils may hinder WIHN (fold = 3.94) (Fig 4b).

To surmount the difficulties with meager neutrophil depletion using Ly6G antibodies, we generated a transgenic mouse model for selective and inducible ablation of neutrophils upon injection of diphtheria toxin (DT)[57]. MRP8-Cre mice expressing Cre recombinase under the control of the neutrophil-associated human MRP8 promoter[58-60] were crossed with ROSA-iDTR[KI] mice, which have a Cre-inducible simian DT receptor (DTR)[61]. This generated mice with DTR expression restricted to mature neutrophils that suffer cell death selectively after the injection of DT. Two injections, one before wounding and one a day after, had no effect on control mice (PMN[WT] mice: MRP8-Cre−, ROSA-iDTR[KI]) but substantially reduced neutrophil numbers in the blood (fold = -4.33) and wound beds (fold = -4.27) of PMN[DTR] (MRP8-Cre+, ROSA-iDTR[KI]) mice at wound day 2 (Fig. 4c-d). We next followed the mice after neutrophil ablation in the early phase of wounding to find WIHN was substantially elevated (fold = 3.23) (Fig. 4e). The ablation of neutrophils later in the wound healing process yielded the same results (Wound days 6, 8, and 10) (Fig. S5). Antibody mediated depletion (1A8 clone) at these later time points had no effect on WIHN, suggesting that although effective in the blood, it is not able to target neutrophils already present within the wound bed (Fig. S6). After overcoming the technical hurdles above, these data suggest that neutrophils have a detrimental effect on the regeneration of hair follicles.

To further explore this possibility, we sought to try a gain-of-function approach, rather than the loss-of-function approach above. We injected purified neutrophils from the bone marrow of mice into the wound beds of mice during the re-epithelialization process. Although not significant, likely due to the high variability of WIHN, the addition of 200,000 neutrophils reduced the number of regenerated hair follicles (fold = -0.73) (Fig. 4f).

Finally, given the presence of NETs late in wound healing (Fig 2e), we sought to test the
role of NETs in hair follicle neogenesis directly. We, therefore, tested WIHN in NET-deficient
Pad4 KO mice, as employed in Figure 2e. In the absence of Pad4 and with dramatically reduced
NETs, WIHN is enhanced (fold = 2.47, p = 0.026) (Fig. 4g). This suggests that NETs play a role in
reducing the regenerative capacity of mice during the wound healing process. This correlates
with the capacity of NETs to damage tissue in diseases such as small vessel vasculitis[62], systemic
lupus erythematosus[63-67], rheumatoid arthritis[68], and psoriasis[69].
267

Single-cell RNAseq correlation of mature neutrophils with poor WIHN

Double-stranded RNA sensing, mediated by Tlr3 and downstream effector pathways Il-
6/ Stat3, has been shown to be critical for WIHN, with Tlr3-/- mice having substantially less
regenerated hair follicles than their wild-type controls[34]. Given that TLR3 dsRNA sensing has
been shown to be critical for neutrophil recruitment and NET production in a model for acute
lung injury (ALI) and glomerulonephritis (GN), we wondered if Tlr3-/- mice paradoxically have
increased neutrophil levels, contributing to lower WIHN[70,71]. As anticipated, Tlr3-/- mice had
substantially less WIHN than their WT controls (fold = -3.75, p = 0.0218) (Fig. 4h, Fig. S7a). We
performed single-cell RNA sequencing on wildtype and Tlr3-/- re-epithelialized wound beds of
mice 10 days post wounding and prior to morphogenesis. Approximately 8014 sequenced cells
met standard quality control metrics and were further analyzed in the Seurat R package[72].
Unsupervised clustering and UMAP non-linear dimensional reduction identified 18 cell clusters.
Seurat generated conserved and differentially expressed genes to assign cluster identities. We
found the mature neutrophil cluster to be substantially increased in the Tlr3-/- mice (Fig. 4i, Fig.
S7b). This disparity is evident when looking at select differentially expressed neutrophil
associated genes that are all elevated in Tlr3 -/- mice (Fig. 4j). Together, these data suggest that
elevated mature neutrophil levels within Tlr3-/- mice correlate with their deficiency in WIHN.

Discussion

The wound healing process is a careful balance of interconnected steps that must weigh
the benefits of quick barrier repair, which leads to fibrous scarring, and more complete
regeneration, that restores function and appearance. The role inflammation plays in
regeneration and scarring is still being elucidated, but increasing evidence suggests that neither
excess nor lack of inflammation supports regeneration. Fgf9-producing γ-δ T cells are critical for
WIHN, infiltrating into wound bed immediately before re-epithelialization and onset of hair
follicle regeneration[7]. Macrophages have also been shown to be important in the process, with
their ablation eliminating WIHN[8-10]. Additionally, the injection of the dsRNA mimic poly(I:C), as
early as 3 days after wound, dramatically enhances WIHN[34]. Intriguingly, spiny mice (*Acomys*)
have a dramatically reduced inflammatory response post wounding, with less cytokines and
virtually no macrophages until late in the process, when compared to laboratory mice (*Mus*),
despite having substantially improved regeneration[73,74]. Consistent with this idea of a complex
network of inflammatory cells influencing regeneration, we show that neutrophils persist in the

wound bed after the acute inflammatory phase and—despite multiple correlations to the
contrary-- have a detrimental effect on regeneration.

Using histological and flow cytometry techniques, we show that while abundant
immediately after wounding, mature neutrophils remain in low levels within the wound bed
after re-epithelization. These late stage neutrophils produce NETs, which are eliminated in mice
that are deficient in Pad4, an enzyme critical for chromatin decondensation and NET
formation[50,51]. Critically, Pad4-/- mice have increased WIHN, when compared with WT mice.
Furthermore, antibody mediated and selective genetic neutrophil ablation dramatically boost
WIHN. Single-cell RNA sequencing also revealed that mice deficient in dsRNA sensing Tlr3, who
have severely reduced WIHN, have substantially more mature neutrophils present in the re-
epithelized wound bed, immediately preceding regeneration. All these data suggest a model
where neutrophils play an important role in defense against bacterial pathogens, but if they
persist within the wound bed too long after barrier repair, regeneration is hindered. In the
future, it will be interesting to see if selectively targeting NETs (e.g., PAD4 inhibitors, DNase I, N-
acetylcysteine) enhance regenerative wound repair.

When attempting to deplete neutrophils to assess their effects on WIHN, we also
discovered that the commonly used antibody depletion methods were not effective. Two
neutrophil-specific antibodies, which recognize different epitopes of the GPI-linked
differentiation antigen Ly6G, are commonly injected to rapidly deplete circulating neutrophils.
The most widely used clone (1A8) specifically recognizes Ly6G, while the anti-granulocyte
receptor-1 (Gr-1) antibody (RB6-8C5 clone) recognizes a heterodimer of Ly6G and Ly6C, making
it partially target monocytes. Both antibodies efficiently depleted neutrophils in unwounded
animals. When large full-thickness wounds ($1.25x1.25cm^2$) were used, which are necessary to
initiate WIHN, depletion levels in the blood were drastically reduced, while having negligible
effects on neutrophils within the wound bed. Depletion efficacy was restored if smaller wounds
were used ($0.6x0.6cm^2$), albeit at reduced levels when compared to unwounded mice. These
results suggest that wounding has a dramatic effect on recruiting neutrophil precursors for
mobilization and release, overwhelming the standard ability of depletion antibodies to work
effectively. This highlights the importance of rigorously testing antibody mediated depletion
methods in mouse models, to ensure they are achieving appreciable depletion. Furthermore, it
is important to carefully design neutrophil detection methods to assess depletion efficacy,
particularly with flow cytometry. Robust depletion can be erroneously detected if similar Ly6G
clones are used for both depletion and detection, due to epitope masking[53]. This can occur
since identical monoclonal clones are sometimes given different clone numbers. Antibody
manufacturers are increasingly manipulating antibodies to reduce undesirable features like
non-specific binding, without affecting the epitope-binding variable region, yet renaming the
clones, necessitating substantial effort to ensure non-overlapping epitope binding by both
depletion and analytical flow antibodies.

Our work generates important areas of future investigation. One question is unraveling
the paradox of why neutrophil infiltration signatures correlate with high WIHN, but mature
neutrophils inhibit WIHN. One possible model is that a common upstream cue or factor both
promotes WIHN and promotes neutrophil infiltration, but the latter serves to limit WIHN in
favor of decreasing infection risk. Defining this common upstream signal will be important for

future work. It will also be interesting to define the function of neutrophils within the re-epithelized wound since the barrier has been restored. Another important question is whether nuclear RNAs released in NETs have any function besides the general theorized one for released DNA. Though NETs inhibit WIHN, nuclear RNA release in NETs might have functions in separate physiological processes that will yield important insights. A final question is the identity of the neutrophil mobilization factor whose strength correlates to wound size. Potential signals include products of complement activation (C5a) or a number of small molecular weight C-X-C chemokines, which are recognized by Cxcr1 and Cxcr2[75,76].

In summary, we here demonstrate a novel role for NETs and mature neutrophils to inhibit regeneration. Future studies will be important to further understand the biology of regeneration and test the capacity for neutrophil inhibition to promote regenerative healing.

Acknowledgments: The authors also thank Conover Talbot Jr. (JHMI Deep Sequencing and Microarray Core) for assistance with microarray analysis.

Competing financial interests: L.S.M. is a full-time employee of Janssen Pharmaceuticals and may hold Johnson & Johnson stock and stock options. L.S.M. performed all work at his prior affiliation at Johns Hopkins University School of Medicine and he has received prior grant support from AstraZeneca, Pfizer, Boehringer Ingelheim, Regeneron Pharmaceuticals, and Moderna Therapeutics, was a paid consultant for Armirall and Janssen Research and Development, was on the scientific advisory board of Integrated Biotherapeutics and is a shareholder of Noveome Biotherapeutics, which are all developing therapeutics against infections and/or inflammatory conditions.

Funding: Research reported in this publication was supported by the National Institute of Arthritis, Musculoskeletal, and Skin Diseases, part of the National Institutes of Health, under 1F32AR074865-01 to EW and R01AR064297/AR068280 to LAG.

Methods

Mouse Lines

All wild-type and control mice used for in vivo experiments were on the C57BL/6J background. All mice were age-matched and co-housed until 3-weeks of age. *Pad4* knockout mice were purchased from the Jackson Laboratory (B6.Cg-Padi4[tm1.1Kmow]/J, 030315). The diphtheria toxin (DT) mediated neutrophil ablative mice were generated by crossing ROSA26iDTR (C57BL/6-Gt(ROSA)26Sor[tm1(HBEGF)Awai]/J , 007900) and MRP8-Cre-ires/GFP (B6.Cg-Tg(S100A8-cre,-EGFP)1Ilw/J, 021614) from the Jackson Laboratory to get heterogeneous mice and genotyped according to their specifications. Mice who genotyped positive for Cre (MRP8-Cre+; ROSA-iDTR[KI]) were considered PMN[DTR] mice, while those that were negative for Cre (MRP8- Cre−; ROSA-iDTR[KI]) were PMN[WT] littermate controls[57]. Tlr3 knockout mice (B6N.129S1-Tlr3[tm1Flv]/J,

009675) and C57BL/6NJ controls (005304) were purchased from the Jackson Laboratory. All mice were bred and housed at an American Association for the Accreditation of Laboratory Animal Care (AAALAC)-compliant facility, and all experimental procedures were reviewed and approved by the Johns Hopkins University Institutional Animal Care and Use Committee (IACUC).

Wound Induced Hair Neogenesis (WIHN) Assay
All in vivo experimental surgical procedures were performed as previously characterized[7,16,34,37,77]. Briefly, after exposure to anesthesia (Baxter, Isoflurane), the dorsal side of 3-week-old (21 days) male and female mice were shaved. Surgical scissors were used to excise 1.25x1.25cm^2 of skin, creating wounds deep into the fascia. Approximately 3 weeks after wounding (~21 days), neogenic hair follicles in the re-epithelialized skin tissue were quantified using reflectance confocal scanning laser microscopy (CSLM) as previously published[34,37].

Neutrophil Depletion
C57BL/6J mice were depleted with 100 or 500µg of anti-Ly6G (Bio-X-Cell, 1A8 clone, BP0075-1) via intraperitoneal injection (IP) injection one day prior to and one day after wounding mice[52,54]. Select experiments also used 200µg of anti-Gr1 (Bio-X-Cell, RB6-8C5 clone, BP0075). LgG2a (Bio-X-Cell, BE0089) and IgG2b (Bio-X-Cell, BP0090) isotype controls were used for Ly6G or Gr1 experiments, respectively. DT depletion was done with PMN^{DTR} and PMN^{WT} littermate control mice that were IP injected with 250ng DT (Sigma-Aldrich). The injections were primarily done one day before and after wounding, or at wound days 6, 8, and 10.

Flow Cytometry
Flow cytometry was used to access neutrophil depletion. Blood was collected via retro-orbital sinus bleeds, and red blood cells were lysed RBC lysis buffer (BioLegend, 420301). Wound beds were surgically removed, and cell suspensions were prepared by digesting the tissue in a cocktail consisting of Liberase TL (Roche, 5401020001) and DNase I (Sigma, DN25) in RPMI 1640 (Gibco, 11875093). Cells were washed and then Fc blocked (BioLegend, 101320), before staining with an antibody cocktail (Extended Table 1). Finally, cells were washed and resuspended in FACs buffer containing Propidium Iodide (Miltenyi, 130-093-233). All flow cytometry experiments were performed on a BD LSR II, and downstream analysis of data was performed using FlowJo.

Neutrophil Extracellular Trap measurement
Wound beds were surgically removed at wound days 2 and 7, using a 6mm biopsy punch to remove excess tissue. Cell suspensions were prepared by digesting the tissue in a cocktail consisting of Liberase TL (Roche, 5401020001) and DNase I (Sigma, DN25) in RPMI 1640 (Gibco, 11875093). Cells were washed, and then Fc blocked (BioLegend, 101320), before staining with an antibody cocktail containing anti-MPO (Abcam, ab208670, 1:500) or respective isotype control (Abcam, ab172730, 1:500). Cells were then stained with a secondary Alexa Fluor 647 antibody (Abcam, ab150083, 1:2000). On the final wash, SYTOX green was added (ThermoFisher, S7020, 1:1000). This was performed on a BD LSR II, and downstream analysis of data was performed using FlowJo.

425
3'-end single-cell RNA-sequencing:
The re-epithelialized wounds beds (wound day 10) of a Tlr3-/- and a C57BL/6NJ control mouse
were excised, and cell suspensions were prepared by digesting the mouse skin tissue in a
cocktail consisting of Liberase TL (Roche, 5401020001) and DNase I (Sigma, DN25) in RPMI 1640
(Gibco, 11875093). Propidium iodide and DAPI positive dead cells were removed via cell sorting
with a BD FACSAria II. Single-cell libraries were prepared via a 10× Genomics Chromium Single-
Cell Platform, followed by sequencing using Illumina NovaSeq 6000. The results were run
through Cell Ranger pipeline software for sequence alignment and basic filtering. GEM
generation, barcoding, cDNA amplification, library preparation, quality control, and sequencing
were performed at the Genomics High Throughput Sequencing facility at Johns Hopkins School
of Medicine.
437
Downstream analysis, after the Cell Ranger pipeline, was done using the Seurat R package. A
standard pre-processing workflow was done, removing low quality cells or doublets, filtering
unique feature count over 3750 and below 200, as well as filtering out cells with higher than 5%
mitochondrial counts. This resulted in 4150 WT and 5648 Tlr3-/- cells for downstream
bioinformatics. Expression matrices then underwent normalization, scaling, principal
components analysis, and subsequent t-SNE analysis using Seurat packages. Seurat was then
used to generate conserved genes, differentially expressed genes, feature plots, dot plots, and
ridge plots. Cell clusters were then defined querying conserved genes and differentially
expressed genes against the Immgen gene expression database (www.immgen.org) using the
interactive tool "My Gene Set."
448
Histology
Biopsies from mouse skin tissue were removed and fixed in 4% paraformaldehyde overnight
and then transferred to 70% ethanol. Samples were then submitted to the Johns Hopkins
Oncology Tissue Services Core facility where they were embedded in paraffin. Tissue sections
were obtained at 4µm thickness and mounted onto glass slides, followed by hematoxylin and
eosin (H&E) staining.
455
Immunofluorescence and immunohistochemistry
Immunofluorescence microscopy was performed on de-paraffinized tissue sections that
received heat-induced antigen retrieval using Target Retrieval Solution (Agilent Dako, S169984-
2). After washing and permeabilization in TBS-T universal buffer (0.2% Triton X-100 in tris-
buffered saline), sections were blocked at room temperature in 5% goat, donkey, or fetal
bovine serum with 1% bovine serum albumin. Tissue sections were then incubated overnight at
4°C with primary antibodies (Extended Table 2) in Antibody Diluent (Agilent Dako, S080983-2).
Following a wash step, sections were incubated in fluorescent-dye conjugated secondary
antibodies diluted in antibody diluent for 1 hour at room temperature. After final washing,
sections were mounted with VECTASHIELD® Hardset™ Antifade Mounting Medium with DAPI
(Vector Laboratories, H-1500) for nuclear staining. All imaging was done on a DFC365FX (Leica)
at 20x and 40x magnifications.
468

U1 in situ hybridization

U1 in situ probes were designed and ordered in the Stellaris Probe Designer (Biosearch Technologies) (Extended Table 3). Tissue sections were de-parafinized and stained following Biosearch Technologies Stellaris RNA FISH protocol for Formalin-Fixed Paraffin-Embedded Tissue. Briefly, tissue sections were washed in Wash Buffer A (Biosearch Technologies, SMF-WA1-60), before adding 200µl hybridization buffer (Biosearch Technologies, SMF-HB1-10) containing the U1 probe and covering the tissue with a glass coverslip. The slides were then incubated overnight in a humid box at 37°C. Slides were then immersed in Wash Buffer A in the dark at 37°C for 30 minutes, allowing the coverslips to float off. Slides were then washed for 5 minutes with Wash Buffer B (Biosearch Technologies, SMF-WB1-20), before sections were mounted with VECTASHIELD® Hardset™ Antifade Mounting Medium with DAPI (Vector Laboratories, H-1500) for nuclear staining. All imaging was done on a DFC365FX (Leica) at 63x magnifications.

Microarray, RNA-seq and proteomic analysis

Proteins from the wound center and wound edges were analyzed by proteomics, as previously described[37]. Briefly, we used Orbitrap Fusion Tribrid mass spectrometer (Thermo Scientific) for protein profiling and obtained the protein expression level by the MS Amanda algorithm. Standard bioinformatics procedures were performed, including standardization of gene expression, the definition of differentially expressed genes, and GO enrichment analysis. For both Rnasel-/- and WT mice, total RNA was isolated from mouse tissue at the time of scab detachment from the wound (~10 days post-wounding). RNA was submitted to the JHMI Deep Sequencing & Microarray core facility and profiled using the Affymetrix Clariom™ S mouse array platform, according to the manufacturer's protocols. Gene chips were scanned, generating CEL pixel intensity files, which were processed and analyzed using Partek® Genomics Suite™ software, and the Robust Multichip Analysis (RMA) algorithm was used for normalization. For SPF and GF analysis, total RNA from early wound bed skin (~WD12) was submitted to the JHMI Transcriptomics and Deep Sequencing Core. The 1.0ST exon sequencing of mouse RNA was performed according to the manufacturer's standard protocol. The raw affymetrix CEL data was standardized using Robust Multichip Analysis (RMA) algorithm for comparison.

Quantification and Statistical Analysis

All in vivo and in vitro experiments were performed in at least individual instances. Univariate statistical analysis was performed using Student's t-test, and multivariate analysis was performed using ANOVA. All statistical analyses and graphical representations were generated using GraphPad Prism software. Statistical significance is defined as p-values <0.05 derived from the standard error of mean calculations.

Extended Table 1

Name	Host	Fluorophore	Manufacturer/Product #

MHCII (IA/IE)	Rat	BV421/Pacific Blue	BioLegend/107631
CD3	Rat	BV510/AmCyan	BD/740147
Ly6C	Rat	FITC	BD/553104
Ly6G	Rat	PE	BioLegend/127607
CD45	Rat	PE-Cy5.5	Invitrogen/35-0451-82
CD115	Rat	PE-Vio770	BioLegend/135523
CD11c	Hamster	APC	BioLegend/117310
CD11b	Rat	APC-Cy7	BD/557657

511 **Extended Table 2**

Name	Host	Dilution	Company/ Product #
MPO	Goat	1:200	Abcam/ab208670
F4/80	Rat	1:200	Abcam/ab6640
Ly6G	Rat	1:200	BioXCell/BP0075-1
H3Cit	Rabbit	1:500	Abcam/ab5103
Alexa Fluor® 488 Anti-Goat IgG (H+L)	Rabbit	1:1000	Invitrogen/A27012
Alexa Fluor® 488 Anti-Rabbit Ig (H+L)	Goat	1:1000	Invitrogen/A-11008
Alexa Fluor® 594 Anti-Rabbit IgG (H+L)	Goat	1:1000	Invitrogen/A-11037

Alexa Fluor® 488 Anti-Rat Ig (H+L)	Goat	1:1000	Invitrogen/A-11006
Alexa Fluor® 594 Anti-Rat Ig (H+L)	Donkey	1:1000	Invitrogen/ A-21209

512
513 **Extended Table 3**

Sequence Name	Sequence
U1 snRNA_1	cccctgccaggtaagtat
U1 snRNA_2	caccttcgtgatcatggt
U1 snRNA_3	aagcctcgccctgggaaa
U1 snRNA_4	acatccggagtgcaatgg
U1 snRNA_5	gggaaatcgcaggggtca
U1 snRNA_6	cagtcgagtttcccacat
U1 snRNA_7	ccccactaccacaaatta
U1 snRNA_8	aggggaaagcgcgaacgc

514
515
516 **Figure Legends**
517
Figure 1: Neutrophil signature correlates with high skin regeneration in multiple models
a. Schematic of hair neogenesis preferential localization to wound center (high WIHN) rather
than wound edge (low WIHN). **b.** Proteomic gene ontology (GO) analysis of the top 100 genes
wound center versus wound edge in wild type mice (at scab detachment) shows a
predominance in innate immune response pathways and neutrophil signatures. **c.** Abundance
ratios of genes from select GO terms highlighted in b. show enrichment of antimicrobial and
granular proteins, labeled in red. **d.** GO analysis of the top 200 genes from high WIHN Rnasel[-/-]
versus wild-type mice at scab detachment shows a predominance of neutrophil and innate
immune cell chemotaxis pathways. Inset graphs show the gene fold expression changes for
genes present in that category. **e.** Neutrophil chemotaxis and innate immune categories are
included in Gene Ontology (GO) enrichment analysis of the top versus bottom 500 differentially
expressed genes between Specific pathogen-free (SPF; high WIHN) and germ-free (GF; low
WIHN) mice on ~WD12 (scab detachment) wound beds. (n=3 independent animals per group).
Inset graphs show the gene fold expression changes for genes present in that category.
532
533

Figure 2: Neutrophils persist in wound bed after the acute inflammatory phase, producing extracellular traps

a. Neutrophils are present in the wound beds of C57BL/6J mice at early time points, visible in representative hematoxylin and eosin [H&E] staining. Red arrows show select neutrophils, evident by their multilobular nuclei. The black dashed line signifies the migrating epithelial front. Black scale bar = 50 μm. **b.** Neutrophils predominate throughout the wound beds of C57BL/6J mice on wound days 1 and 3, visible in prominent MPO immunofluorescence (green). Few macrophages are present (Red, F4/80). Red arrows show select neutrophils. The white dashed line signifies the dorsal edge of the wound bed. White scale bar = 200 μm. **c.** Percent neutrophil (Ly6G+ cells from total CD45+ cells) levels are consistent in the blood throughout the wound time course, but drop in the wound bed at wound day 11, as measured by FACS. ****$p < 0.0001$, as calculated by two-way ANOVA. n.s., not significant. N = 2 vs 4. Results are representative of at least two independent experiments. **d.** Citrullinated histone H3 (H3Cit, red) co-localized with Ly6G+ neutrophils (green), beginning at wound day 3 in the wound beds of immunofluorescence stained C57BL/6J mice, indicating the formation of extracellular traps. Red arrows show select neutrophils. The white dashed line signifies the dorsal edge of the wound bed. White scale bar = 200 μm. **e.** Neutrophil extracellular trap positive cells (MPO+, SYTOX green +) are present at late wound time points, but are absent in the wound beds of PAD4-/- mice, as measured by FACs ****$p < 0.0001$, as calculated by two-way ANOVA. n.s., not significant. N = 7 vs 4. Results are representative of at least two independent experiments. **f.** Cytoplasmic U1 snRNA is present in the wound bed of C57BL/6J mice, while it localized exclusively in the nuclei of unwounded controls, as visualized by representative FISH. White arrows show select cells with nuclear or cytoplasmic U1 snRNA. The white dashed line signifies the dorsal edge of the wound bed. The solid white line delineates a hair follicle. White scale bar = 80 μm

Figure 3: Large full-thickness wounds dramatically reduce antibody-mediated neutrophil depletion

a. Schematic of neutrophil depletion via intraperitoneal injection (IP) of Ly6G antibody (1A8 clone, 500μg), Gr1 (Ly6G/Ly6C, RB6-8C5 clone, 200ng), or isotype control. IP injections are done one day before and one day after wounding C57BL/J6 mice with 1.25cm^2 square wounds. Mice were then sacrificed wound day 2 or 2 for FACs analysis of blood or wound bed (WB). **b.** Antibody mediated depletion is not successful in the wound bed. Representative flow plots from the wound beds (wound day 3) of mice treated as in **a.** The percent neutrophils (CD11b+, Ly6c+) present in these samples are presented to the right. The difference between samples is statistically insignificant, as calculated by the two-tailed Student's t-test. N = 2. Results are representative of at least two independent experiments. **c.** Antibody mediated. Neutrophil depletion is marginally successful in the blood, but not the wound bed, two days after wounding. The percent neutrophils (CD11b+, Ly6c+), at wound day 2, present in the blood and wound beds of mice treated as in **a.** (Ly6G, 1A8). **$p < 0.0011$, as calculated by two-way ANOVA. n.s., not significant. N = 4. Results are representative of at least two independent experiments. **d.** Neutrophil depletion from the blood is successful 1-day post injection (Ly6G, 1A8), in the absence of wounding, via FACs. The neutrophil depletion is statistically significant.

****p < 0.0001, as calculated by two-tailed Student's t-test. N = 4 vs 6. **e.** The efficacy of
neutrophil depletion from the blood diminishes after 1.25X1.25cm^2 wounds are inflicted. The
percent neutrophils (Gr1+, Ly6c+) in the blood at wound day 2, via FACs. The neutrophil
depletion is statistically insignificant. n.s., not significant. N = 2 vs 3. Results are representative
of at least three independent experiments. **f.** Neutrophil depletion from the blood is more
pronounced in unwounded mice. The percent neutrophils (Gr1+, Ly6c+) present in the blood at
wound day 2 of mice IP injected with Ly6G antibody (1A8, 500 μg), with or without wounding,
via FACs. ****p < 0.0001 vs **p=0.0038 as calculated by two-way ANOVA. N = 6 vs 7. **g.**
Neutrophil depletion from the skin is ineffective, regardless of wound status. The percent
neutrophils (Gr1+, Ly6c+) present in the wound bed or skin at wound day 2 of mice IP injected
with Ly6G antibody (1A8, 500ug), with or without wounding. The neutrophil depletion is
insignificant in both cases, as calculated by two-way ANOVA. n.s., not significant. N = 6 vs 7. **h.**
C57BL/6J mice were given full thickness wounds on the center of their backs. The normal large
wound (LW) is 1.25x1.25cm^2, while the smaller wound (SW) is 0.6x0.6cm^2. **i.** Neutrophil
depletion from the blood is successful on small, but not large wounded mice. Mice wounded
with small or large wounds (**h.**) were IP injected with Ly6G antibody (1A8, 500μg) one day
before and after wounding. Blood was extracted on wound day 2 and analyzed by flow
cytometry for percent neutrophils (Gr1+, Ly6c+). *p = 0.0362 as calculated by two-way ANOVA.
n.s., not significant. N = 2 vs 3 and 2 vs 2. Results are representative of at least two independent
experiments. **j.** Neutrophil depletion from the wound bed is successful on small, but not large
wounded mice. Mice wounded with small or large wounds (**h.**) were IP injected with Ly6G
antibody (1A8, 500μg) one day before and after wounding. Wound beds were collected on
wound day 2 and analyzed by flow cytometry for percent neutrophils (Gr1+, Ly6c+). Depletion
was significant in the small wound, but not the large wound setting. *p = 0.0458 as calculated
by two-way ANOVA. n.s., not significant. N = 2 vs 3 and 2 vs 2. Results are representative of at
least two independent experiments.
604
605
Figure 4: Mature neutrophils inhibit wound induced hair neogenesis
a. Schematic of neutrophil depletion via IP of Ly6G antibody (100μg) or isotype control. IP
injections are done one day before and one day after wounding C57BL/J6 mice with
1.25x1.25cm^2 square wounds. Wound induced hair neogenesis (WIHN) was measured 21 days
after wounding via confocal scanning laser microscopy (CSLM). **b.** Neutrophil targeted Ly6G
antibody injected mice exhibit increased WIHN (CSLM, images; fold = 3.94, p = 0.011, N = 10 vs
9). In each image, the dash red box indicates the area of hair follicle regeneration. **c.** Diphtheria
toxin (DT, 250ng) injection in heterozygous ROSA26iDTR/MRP8-Cre-ires mice (PMN^{DTR} and
PMN^{WT}), following the injection scheme in **a.**, successfully depletes neutrophils from the blood
in mice with large wounds**.** Fold = -4.33. **p = 0.0098. N = 2 vs. 3. Results are representative of
at least two independent experiments. **d.** Mice treated as in **c.** are depleted of neutrophils in
their wound beds at wound day 2. Fold = -4.27. **p = 0.0069. N = 2 vs. 3. Results are
representative of at least two independent experiments. **e.** PMN^{DTR} mice IP injected with
diphtheria toxin (DT, 250ng) on wound days -1 and 1 exhibit increased WIHN (CSLM, images;
fold = 3.23, ***p = 0.0010, N = 14 vs 6). In each image, the dash red box indicates the area of
hair follicle regeneration. **f.** Regenerated hair follicles after the injection of 50k-200k purified

neutrophils underneath the scab at WD7-8. (CSLM, images; n.s., not significant, N = 9 vs 3 vs 6).
In each image, the dash red box indicates the area of hair follicle regeneration. **g.** Pad4$^{-/-}$ mice
defective in extracellular traps exhibit increased WIHN (CSLM, images; fold = 2.47, p = 0.026, N
= 10 vs 19). In each image, the dash red box indicates the area of hair follicle regeneration. **h.**
Tlr3$^{-/-}$ mice exhibit decreased WIHN (fold = -3.75 p = 0.0218, N = 8 vs 10). **i.** The presence of
increased neutrophils correlates with decreased WIHN in TLR3$^{-/-}$ mice. scRNA-seq t-SNE plot
shows differences between WT (red, 4150 cells) and Tlr3-/- (blue, 5648 cells) wound beds at
wound day 10. The plots were generated via Seurat. The neutrophil cluster is circled in red.
Percent of neutrophils are graphed to the right. **j.** Neutrophil associated gene expression is
more pronounced within the neutrophils of Tlr3-/- mice, compared to WT. Generated in Seurat
with RidgePlot function.
633
634   **Supplementary Figure Legends**
635
636   **S1:** Macrophage (F4/80) levels are largely absent from the blood and low in the wound bed
637   during the early phase of healing, but increase dramatically at wound day 11, as measured by
638   FACS. ***p < 0.004, as calculated by two-way ANOVA. n.s., not significant. N = 2 vs 4. Results
639   are representative of at least two independent experiments.
640
S2: Ly6G 1A8 antibody clone (Bio-X-Cell) masks the Ly6G epitope, preventing detection by the
Ly6G REA526 clone (Miltenyi, engineered 1A8 clone) and partially preventing detection by the
Gr1 RB6-8C5 clone (Bio-X-Cell) antibodies. Blood was extracted from a single C57BL/6J and
stained sequentially with combinations of two antibodies, to test Ly6G epitope masking, before
detecting by flow cytometry. If only stained with the Ly6G (REA526) or Gr1 (RB6-8C5), the
neutrophil population is detected normally (13.7% or 13.2%, respectively). If you first stain with
Ly6G (1A8 clone), followed by Ly6G (REA526) or Gr1 (RB6-8C5), you get varying degrees of Ly6G
antibody masking. Being a derivative of the 1A8 clone, the REA526 Ly6G binding site is almost
completely blocked by the prior incubation with the Ly6G 1A8 clone (1.10%), making the pair
unusable for neutrophil depletion experiments. Prior incubation with the Ly6G 1A8 clone
followed by Gr1 (recognized an epitope of Ly6G/Ly6C) shifts the neutrophil population to the
left, but they remain distinct from the negatively stained cells (12.7%). With careful gating, this
makes depletion with the Ly6G (1A8 clone) antibody and detection with the Gr1 (RB6-8C5)
antibody possible. The neutrophil population is boxed in red.
655
S3: Antibody mediated Neutrophil depletion is not efficient in the blood, spleen, liver, or wound
bed, two days after wounding. The percent neutrophils (CD11b+, Ly6c+), at wound day 2,
present in the blood, spleen, liver, and wound beds of mice IP injected with Ly6G antibody
(500μ, 1A8) one day before and one day after given 1.25x1.25cm^2 full thickness wounds. The
neutrophil depletion is statistically insignificant, as calculated by two-way ANOVA. n.s., not
significant. N = 4 vs 4, except for the Blood and Wound bed samples, which were 7 vs. 7.
662
663   **S4:** The wound closure rate was not affected by neutrophil depletion via IP of Ly6G antibody
664   (100μg) or isotype control. IP injections were done one day before and one day after wounding

665 C57BL/J6 mice with 1.25x1.25cm^2 square wounds. N = 4 vs 4. Results are representative of
666 three independent experiments.
667
668 **S5.** PMN^{DTR} mice IP injected with diphtheria toxin (DT, 250ng) on wound days 6, 8, and 10
669 exhibit increased WIHN (CSLM, images; fold = 3.28, **p = 0.0085, N = 13 vs 5).
670
S6: a. Schematic of neutrophil depletion via IP of Ly6G antibody (500µg) or isotype control. IP
injections are done on wound day 8 and 10, after wounding C57BL/J6 mice with 1.25x1.25cm^2
square wounds. Wound induced hair neogenesis (WIHN) was measured 21 days after wounding
via confocal scanning laser microscopy (CSLM). **b.** Neutrophil depletion in the blood is
successful on wound day 9, one day after antibody injection (Ly6G, 1A8), via FACs. The
neutrophil depletion is statistically significant. ****p < 0.0001, as calculated by two-tailed
Student's t-test. Fold = -87.9. N = 2 vs 3. **c.** Late stage Ly6G antibody injected mice exhibit
normal WIHN (CSLM, images; n.s., not significant, N = 3 vs 3). In each image, the dash red box
indicates the area of hair follicle regeneration.
680
681 **S7: a.** Tlr3$^{-/-}$ mice exhibit decreased WIHN (representative CSLM images, N = 8 vs. 10). In each
682 image, the dash red box indicates the area of hair follicle regeneration. **b.** Neutrophil associated
683 genes used to identify the neutrophil cluster in UMAP non-linear dimensional reduction, via
684 Seurat R package, of WT and Tlr3-/- mice.
685
686
687
688 **Works Cited:**
689
690

1. Diegelmann, R.F. & Evans, M.C. Wound healing: an overview of acute, fibrotic and
delayed healing. *Frontiers in bioscience : a journal and virtual library* **9**, 283-289 (2004).
2. O'Connell, J.B. The economic burden of heart failure. *Clinical cardiology* **23**, III6-10
(2000).
3. Chung, F.*, et al.* Assessing the burden of respiratory disease in the UK. *Respiratory*
medicine **96**, 963-975 (2002).
4. Godwin, J.W., Pinto, A.R. & Rosenthal, N.A. Macrophages are required for adult
salamander limb regeneration. *Proceedings of the National Academy of Sciences of the*
United States of America **110**, 9415-9420 (2013).
5. Kyritsis, N.*, et al.* Acute inflammation initiates the regenerative response in the adult
zebrafish brain. *Science* **338**, 1353-1356 (2012).
6. Taniguchi, K.*, et al.* A gp130-Src-YAP module links inflammation to epithelial
regeneration. *Nature* **519**, 57-62 (2015).
7. Gay, D.*, et al.* Fgf9 from dermal gammadelta T cells induces hair follicle neogenesis after
wounding. *Nature medicine* **19**, 916-923 (2013).

8. Kasuya, A., Ito, T. & Tokura, Y. M2 macrophages promote wound-induced hair
neogenesis. *Journal of dermatological science* **91**, 250-255 (2018).
9. Rahmani, W.*, et al.* Macrophages Promote Wound-Induced Hair Follicle Regeneration in
a CX3CR1- and TGF-beta1-Dependent Manner. *The Journal of investigative dermatology*
138, 2111-2122 (2018).
10. Wang, X.*, et al.* Macrophages induce AKT/beta-catenin-dependent Lgr5(+) stem cell
activation and hair follicle regeneration through TNF. *Nature communications* **8**, 14091
(2017).
11. Seifert, A.W.*, et al.* Skin shedding and tissue regeneration in African spiny mice
(Acomys). *Nature* **489**, 561-565 (2012).
12. Billingham, R.E. & Russell, P.S. Incomplete wound contracture and the phenomenon of
hair neogenesis in rabbits' skin. *Nature* **177**, 791-792 (1956).
13. Breedis, C. Regeneration of hair follicles and sebaceous glands from the epithelium of
scars in the rabbit. *Cancer research* **14**, 575-579 (1954).
14. Kligman, A.M. & Strauss, J.S. The formation of vellus hair follicles from human adult
epidermis. *The Journal of investigative dermatology* **27**, 19-23 (1956).
15. Lacassagne, A. & Latarjet, R. Action of methylcholanthrene on certain scars of the skin in
mice. *Cancer research* **6**, 183-188 (1946).
16. Ito, M.*, et al.* Wnt-dependent de novo hair follicle regeneration in adult mouse skin after
wounding. *Nature* **447**, 316-320 (2007).
17. Plikus, M.V.*, et al.* Regeneration of fat cells from myofibroblasts during wound healing.
Science **355**, 748-752 (2017).
18. Kim, M.H.*, et al.* Dynamics of neutrophil infiltration during cutaneous wound healing
and infection using fluorescence imaging. *The Journal of investigative dermatology* **128**,
1812-1820 (2008).
19. Guo, S. & Dipietro, L.A. Factors affecting wound healing. *Journal of dental research* **89**,
219-229 (2010).
20. Segal, A.W. How neutrophils kill microbes. *Annual review of immunology* **23**, 197-223
(2005).
21. Brinkmann, V.*, et al.* Neutrophil extracellular traps kill bacteria. *Science* **303**, 1532-1535
(2004).
22. Papayannopoulos, V. Neutrophil extracellular traps in immunity and disease. *Nature
reviews. Immunology* **18**, 134-147 (2018).
23. Acosta, J.B.*, et al.* The pro-inflammatory environment in recalcitrant diabetic foot
wounds. *International wound journal* **5**, 530-539 (2008).
24. Singer, A.J. & Clark, R.A. Cutaneous wound healing. *The New England journal of
medicine* **341**, 738-746 (1999).
25. DiPietro, L.A. Wound healing: the role of the macrophage and other immune cells. *Shock*
4, 233-240 (1995).
26. Takeuchi, O. & Akira, S. Pattern recognition receptors and inflammation. *Cell* **140**, 805-
820 (2010).
27. Huebener, P. & Schwabe, R.F. Regulation of wound healing and organ fibrosis by toll-like
receptors. *Biochimica et biophysica acta* **1832**, 1005-1017 (2013).

28. Kawai, T. & Akira, S. Toll-like receptor and RIG-I-like receptor signaling. *Annals of the*
New York Academy of Sciences **1143**, 1-20 (2008).
29. Cavassani, K.A.*, et al.* TLR3 is an endogenous sensor of tissue necrosis during acute
inflammatory events. *The Journal of experimental medicine* **205**, 2609-2621 (2008).
30. Lin, Q.*, et al.* Impaired wound healing with defective expression of chemokines and
recruitment of myeloid cells in TLR3-deficient mice. *Journal of immunology* **186**, 3710-
3717 (2011).
31. Lin, Q.*, et al.* Toll-like receptor 3 ligand polyinosinic:polycytidylic acid promotes wound
healing in human and murine skin. *The Journal of investigative dermatology* **132**, 2085-
2092 (2012).
32. Bernard, J.J.*, et al.* Ultraviolet radiation damages self noncoding RNA and is detected by
TLR3. *Nature medicine* **18**, 1286-1290 (2012).
33. Borkowski, A.W.*, et al.* Toll-like receptor 3 activation is required for normal skin barrier
repair following UV damage. *The Journal of investigative dermatology* **135**, 569-578
(2015).
34. Nelson, A.M.*, et al.* dsRNA Released by Tissue Damage Activates TLR3 to Drive Skin
Regeneration. *Cell stem cell* **17**, 139-151 (2015).
35. Zhu, A.S., Li, A., Ratliff, T.S., Melsom, M. & Garza, L.A. After Skin Wounding, Noncoding
dsRNA Coordinates Prostaglandins and Wnts to Promote Regeneration. *The Journal of*
investigative dermatology **137**, 1562-1568 (2017).
36. Lai, Y.*, et al.* Commensal bacteria regulate Toll-like receptor 3-dependent inflammation
after skin injury. *Nature medicine* **15**, 1377-1382 (2009).
37. Kim, D.*, et al.* Noncoding dsRNA induces retinoic acid synthesis to stimulate hair follicle
regeneration via TLR3. *Nature communications* **10**, 2811 (2019).
38. Liu, Y.*, et al.* Tumor Exosomal RNAs Promote Lung Pre-metastatic Niche Formation by
Activating Alveolar Epithelial TLR3 to Recruit Neutrophils. *Cancer cell* **30**, 243-256
(2016).
39. Zhang, L.J.*, et al.* Antimicrobial Peptide LL37 and MAVS Signaling Drive Interferon-beta
Production by Epidermal Keratinocytes during Skin Injury. *Immunity* **45**, 119-130 (2016).
40. Chow, O.A.*, et al.* Statins enhance formation of phagocyte extracellular traps. *Cell host &*
microbe **8**, 445-454 (2010).
41. Mohanan, S., Horibata, S., McElwee, J.L., Dannenberg, A.J. & Coonrod, S.A. Identification
of macrophage extracellular trap-like structures in mammary gland adipose tissue: a
preliminary study. *Frontiers in immunology* **4**, 67 (2013).
42. Boe, D.M., Curtis, B.J., Chen, M.M., Ippolito, J.A. & Kovacs, E.J. Extracellular traps and
macrophages: new roles for the versatile phagocyte. *Journal of leukocyte biology* **97**,
1023-1035 (2015).
43. Rossaint, J.*, et al.* Synchronized integrin engagement and chemokine activation is crucial
in neutrophil extracellular trap-mediated sterile inflammation. *Blood* **123**, 2573-2584
(2014).
44. Maugeri, N.*, et al.* Activated platelets present high mobility group box 1 to neutrophils,
inducing autophagy and promoting the extrusion of neutrophil extracellular traps.
Journal of thrombosis and haemostasis : JTH **12**, 2074-2088 (2014).

45.   Keshari, R.S., *et al.* Cytokines induced neutrophil extracellular traps formation:
793   implication for the inflammatory disease condition. *PloS one* **7**, e48111 (2012).
794   46.   Gupta, A.K., Giaglis, S., Hasler, P. & Hahn, S. Efficient neutrophil extracellular trap
795   induction requires mobilization of both intracellular and extracellular calcium pools and
796   is modulated by cyclosporine A. *PloS one* **9**, e97088 (2014).
797   47.   Wong, K.W. & Jacobs, W.R., Jr. Mycobacterium tuberculosis exploits human interferon
798   gamma to stimulate macrophage extracellular trap formation and necrosis. *The Journal*
799   *of infectious diseases* **208**, 109-119 (2013).
800   48.   Wong, S.L., *et al.* Diabetes primes neutrophils to undergo NETosis, which impairs wound
801   healing. *Nature medicine* **21**, 815-819 (2015).
802   49.   Delgado-Rizo, V., *et al.* Neutrophil Extracellular Traps and Its Implications in
803   Inflammation: An Overview. *Frontiers in immunology* **8**, 81 (2017).
804   50.   Wang, Y., *et al.* Histone hypercitrullination mediates chromatin decondensation and
805   neutrophil extracellular trap formation. *The Journal of cell biology* **184**, 205-213 (2009).
806   51.   Wang, Y., *et al.* Human PAD4 regulates histone arginine methylation levels via
807   demethylimination. *Science* **306**, 279-283 (2004).
808   52.   Daley, J.M., Thomay, A.A., Connolly, M.D., Reichner, J.S. & Albina, J.E. Use of Ly6G-
809   specific monoclonal antibody to deplete neutrophils in mice. *Journal of leukocyte*
810   *biology* **83**, 64-70 (2008).
811   53.   Ribechini, E., Leenen, P.J. & Lutz, M.B. Gr-1 antibody induces STAT signaling,
812   macrophage marker expression and abrogation of myeloid-derived suppressor cell
813   activity in BM cells. *European journal of immunology* **39**, 3538-3551 (2009).
814   54.   Carr, K.D., *et al.* Specific depletion reveals a novel role for neutrophil-mediated
815   protection in the liver during Listeria monocytogenes infection. *European journal of*
816   *immunology* **41**, 2666-2676 (2011).
817   55.   Dovi, J.V., He, L.K. & DiPietro, L.A. Accelerated wound closure in neutrophil-depleted
818   mice. *Journal of leukocyte biology* **73**, 448-455 (2003).
819   56.   Fan, C., *et al.* Characterization and quantification of wound-induced hair follicle
820   neogenesis using in vivo confocal scanning laser microscopy. *Skin research and*
821   *technology : official journal of International Society for Bioengineering and the Skin* **17**,
822   387-397 (2011).
823   57.   Reber, L.L., *et al.* Neutrophil myeloperoxidase diminishes the toxic effects and mortality
824   induced by lipopolysaccharide. *The Journal of experimental medicine* **214**, 1249-1258
825   (2017).
826   58.   Passegue, E., Wagner, E.F. & Weissman, I.L. JunB deficiency leads to a myeloproliferative
827   disorder arising from hematopoietic stem cells. *Cell* **119**, 431-443 (2004).
828   59.   Elliott, E.R., *et al.* Deletion of Syk in neutrophils prevents immune complex arthritis.
829   *Journal of immunology* **187**, 4319-4330 (2011).
830   60.   Abram, C.L., Roberge, G.L., Pao, L.I., Neel, B.G. & Lowell, C.A. Distinct roles for
831   neutrophils and dendritic cells in inflammation and autoimmunity in motheaten mice.
832   *Immunity* **38**, 489-501 (2013).
833   61.   Buch, T., *et al.* A Cre-inducible diphtheria toxin receptor mediates cell lineage ablation
834   after toxin administration. *Nature methods* **2**, 419-426 (2005).

62.   Kessenbrock, K.*, et al.* Netting neutrophils in autoimmune small-vessel vasculitis. *Nature*
medicine **15**, 623-625 (2009).

63.   Hakkim, A.*, et al.* Impairment of neutrophil extracellular trap degradation is associated
with lupus nephritis. *Proceedings of the National Academy of Sciences of the United*
States of America **107**, 9813-9818 (2010).

64.   Lande, R.*, et al.* Neutrophils activate plasmacytoid dendritic cells by releasing self-DNA-
peptide complexes in systemic lupus erythematosus. *Science translational medicine* **3**,
73ra19 (2011).

65.   Garcia-Romo, G.S.*, et al.* Netting neutrophils are major inducers of type I IFN production
in pediatric systemic lupus erythematosus. *Science translational medicine* **3**, 73ra20
(2011).

66.   Villanueva, E.*, et al.* Netting neutrophils induce endothelial damage, infiltrate tissues,
and expose immunostimulatory molecules in systemic lupus erythematosus. *Journal of*
immunology **187**, 538-552 (2011).

67.   Leffler, J.*, et al.* Neutrophil extracellular traps that are not degraded in systemic lupus
erythematosus activate complement exacerbating the disease. *Journal of immunology*
188, 3522-3531 (2012).

68.   Khandpur, R.*, et al.* NETs are a source of citrullinated autoantigens and stimulate
inflammatory responses in rheumatoid arthritis. *Science translational medicine* **5**,
178ra140 (2013).

69.   Hu, S.C.*, et al.* Neutrophil extracellular trap formation is increased in psoriasis and
induces human beta-defensin-2 production in epidermal keratinocytes. *Scientific reports*
6, 31119 (2016).

70.   Gan, T.*, et al.* TLR3 Regulated Poly I:C-Induced Neutrophil Extracellular Traps and Acute
Lung Injury Partly Through p38 MAP Kinase. *Frontiers in microbiology* **9**, 3174 (2018).

71.   Liu, Q.*, et al.* Toll-Like Receptor 3 Signaling Contributes to Regional Neutrophil
Recruitment in Cultured Human Glomerular Endothelial Cells. *Nephron* **139**, 349-358
(2018).

72.   Satija, R., Farrell, J.A., Gennert, D., Schier, A.F. & Regev, A. Spatial reconstruction of
single-cell gene expression data. *Nature biotechnology* **33**, 495-502 (2015).

73.   Jiang, T.X., Harn, H.I., Ou, K.L., Lei, M. & Chuong, C.M. Comparative regenerative biology
of spiny (Acomys cahirinus) and laboratory (Mus musculus) mouse skin. *Exp Dermatol*
28, 442-449 (2019).

74.   Brant, J.O., Yoon, J.H., Polvadore, T., Barbazuk, W.B. & Maden, M. Cellular events during
scar-free skin regeneration in the spiny mouse, Acomys. *Wound Repair Regen* **24**, 75-88
(2016).

75.   Trujillo, G.*, et al.* Neutrophil recruitment to the lung in both C5a- and CXCL1-induced
alveolitis is impaired in vitamin D-binding protein-deficient mice. *Journal of immunology*
191, 848-856 (2013).

76.   Metzemaekers, M., Gouwy, M. & Proost, P. Neutrophil chemoattractant receptors in
health and disease: double-edged swords. *Cell Mol Immunol* **17**, 433-450 (2020).

77.   Lim, C.H.*, et al.* Hedgehog stimulates hair follicle neogenesis by creating inductive
dermis during murine skin wound healing. *Nature communications* **9**, 4903 (2018).

78.   Kim, S.*, et al.* Simple cell culture media expansion of primary mouse keratinocytes.
879   *Journal of dermatological science* **93**, 135-138 (2019).
880

Figure 1: Neutrophil signature correlates with high skin regeneration in multiple models

Figure 2: Neutrophils persist in wound bed after the acute inflammatory phase, producing extracellular traps

Figure 3: Large full-thickness wounds dramatically reduce antibody-mediated neutrophil depletion

Figure 4: Neutrophils inhibit wound induced hair neogenesis