The genetic architecture of host response reveals the importance of

arbuscular mycorrhizae to maize cultivation

M. Rosario Ramírez-Flores[1], Sergio Perez-Limón[2], Meng Li[3], Benjamin Barrales-Gamez[2],

Víctor Olalde-Portugal[1] and Ruairidh J. H. Sawers[3*]

[1] Departamento de Biotecnología y Bioquímica, Centro de Investigación y de Estudios

Avanzados (CINVESTAV-IPN), Irapuato, C.P. 36821, Guanajuato, México.

[2] Laboratorio Nacional de Genómica para la Biodiversidad/Unidad de Genómica Avanzada,

Centro de Investigación y de Estudios Avanzados, Instituto Politécnico Nacional (CINVESTAV-

IPN), Irapuato, C.P. 36821, Guanajuato, México

[3] Department of Plant Science, The Pennsylvania State University, State College, PA, USA.

11
12 * Author for correspondence: rjs6686@psu.edu

15 Main text word count: 3, 500

16 Number of Figures: 4

17 Number of Tables: 1

19
20
21
22

ABSTRACT

Arbuscular mycorrhizal fungi (AMF) are ubiquitous in cultivated soils, forming symbiotic

relationships with the roots of major crop species. Although studies in controlled conditions have

demonstrated the potential of the symbiosis to enhance host plant nutrition and alleviate

environmental stress, practical difficulties make it hard to estimate the actual benefit in cultivated

fields, not least because of the lack of availability of suitable AMF-free controls. Furthermore,

the response can vary depending on the plant variety in a manner which is not fully understood.

Here, we implemented a novel strategy based on the selective incorporation of AMF-resistance

into a genetic mapping population to evaluate maize response in the field. We found AMF to

account for about one third of the grain production in a rain-fed medium input field, as well as to

impact the relative performance of plant varieties. Characterization of the genetic architecture of

host response allowed us to distinguish mycorrhizal benefit from dependence and indicated a

trade-off between mycorrhizal and non-mycorrhizal performance, both at the level of individual

QTL and genomewide. This approach is applicable to other crop species, permits further

mechanistic analysis and is scalable to full yield trials.

INTRODUCTION

There is a pressing need to develop sustainable agricultural systems that improve productivity whilst minimizing adverse environmental impacts (Lynch, 2019). In the face of this challenge, there is great interest in the contribution of the arbuscular mycorrhizal (AM) symbiosis and other microbial mutualisms (Sawers et al., 2008; Fester and Sawers, 2011; Bender et al., 2016). AM fungi (AMF) occur broadly in cultivated soils, forming symbiotic relationships with the roots of major crop species and, indeed, many commercial "biofertilizers" are already on the market (Faye et al., 2013). Nonetheless, the value of AM symbiosis in agriculture remains a matter of debate (Ryan and Graham, 2018; Rillig et al., 2019). Experimental studies under controlled conditions have demonstrated that AMF have much to offer host plants, including enhanced uptake of immobile soil nutrients (especially phosphorus) and water, and improved tolerance to a range of abiotic and biotic stresses (Borowicz 2001; Chandrasekaran et al., 2014; Augé et al., 2015; Chiu and Paszkowski, 2019). AMF also contribute to important ecosystem services (van der Heijden et al., 2015), including promoting soil aggregation (Rillig and Mummey, 2006) and reducing nutrient loss through leaching and emission (Bender et al., 2014; Cavagnaro et al., 2015). However, efforts to integrate AMF into agricultural practice are hampered by the difficulty of evaluating host response in the field and the lack of appropriate AMF free controls: when exogenous fungal inoculum is applied to a field, it is difficult to gauge its effectiveness with respect to the native fungal community; when looking to evaluate the impact of the native AMF community under any given agronomic scenario, there is no easy way to estimate what the baseline performance of plants would be if the AMF were not present. Although experimental designs using chemical treatments and/or mechanical barriers might be implemented at small scale they are limited and not practical for use in larger genetic studies or trials. Here, to address

these difficulties, we propose the use of genetic variants conferring resistance to AMF in the

design of experiments to study AM symbiosis in the field.

AM symbiosis dates back over 450 million years (Strullu-Derrien et al., 2018), In

contrast, agriculture is a recent development. Modern breeding efforts have focused on

maximizing yield under homogeneous environments with large quantities of synthetic inputs,

and it is not clear how the breeding process has impacted the AM symbiosis (discussed in

(Sawers et al., 2008; Sawers et al., 2018). Nonetheless, modern varieties of our major crops do

retain the capacity to interact with AMF and the molecular machinery necessary for symbiotic

establishment (*e.g.* Gutjahr et al., 2008). Elegant molecular genetic analyses, largely conducted

in *Medicago truncatula* and *Lotus japonicus,* have identified a series of plant-encoded receptor

and signal-transduction components necessary for AM symbiosis and, in relevant host plants, for

interaction with nodule-forming rhizobia. Fungal signals are perceived by the host SYMRK

(DMI2) receptor , triggering cellular calcium spiking events through the action of the nuclear

ion-channels CASTOR and POLLUX (DMI1) and activate downstream events via CCamK

(DMI3) and CYCLOPS (IPD3) transduction components (Parniske, 2008; Zipfel and Oldroyd,

2017). Notwithstanding the great antiquity of AM symbiosis, there is little redundancy in

symbiotic signalling and single gene mutations in the host can be sufficient to generate a

complete early block in symbiotic interaction (*e.g.* Charpentier et al., 2008; Gutjahr et al., 2008).

In non-nodulating hosts, mutations in symbiotic signaling pathway genes appear to specifically

impact AM symbiosis and do not condition obvious pleiotropic effects on plant development. To

date, of the major cereal crops, mutations in the symbiotic signalling pathway have only been

reported in rice ((Charpentier et al., 2008; Gutjahr et al., 2008)), although the genes themselves

can be unambiguously identified in the genomes of plants such as maize, wheat and sorghum.

While the symbiotic pathway is essential for plant interaction with AMF, there is no

evidence that it plays a role in regulating the progression of the symbiosis or that variation in the

pathway underlies the range of host response that is observed among different plant varieties

(Kaeppler et al., 2000; Mascher et al., 2014; Sawers et al., 2017; Watts-Williams et al., 2019).

Indeed, defining variation in plant response to AMF can in itself be challenging (discussed in

Janos, 2007; Sawers et al., 2010). Here, we define *response* as the difference between

mycorrhizal and non-mycorrhizal plants, applied to any phenotypic measurement. In addition,

we define *dependence* as the capacity of a given variety to perform in the absence of AMF

(Janos, 2007), and *benefit* as the degree to which a plant host profits from the association.

Crucially, variation in host response confounds differences in both dependence and benefit. The

optimization of AM symbiosis in crop varieties is concerned with benefit: there is no interest in

elevating host response by increasing dependence (Sawers et al., 2008). However, these

somewhat abstract concepts can be difficult to distinguish, especially when comparing a small

number of varieties at the whole genotype level: if one variety shows a greater proportional

increase in growth when inoculated with AMF than another, does this reflect a superior ability to

profit from the symbiotic interaction, or poorer performance in the absence of symbiosis? This

question has relevance both to hypotheses concerning the impact of past selection of AM

symbiosis and to any future breeding effort to optimize the interaction. Mechanistic

understanding can provide useful insight: *e.g.* response variation linked to differences in root

development might be reasonably assigned to dependence (Paszkowski and Boller, 2002),

response variation linked to differences in the growth of root-external fungal hyphae to benefit

(Sawers et al., 2017). However, it is not practical to characterize all the physiological differences

among a larger collection of varieties and far from trivial to use these data to interpret variation

in host response to AMF. Here, we propose to distinguish dependence and benefit using the

concepts of genotype x environment interaction (Des Marais and Juenger, 2010), where the

mycorrhizal and non-mycorrhizal conditions are considered as two distinct "environments". In

conjunction with the ability to perform large scale mapping experiments, the GEI framework has

the potential to define dependence and response at the level of the variety and at the level of the

action of individual genomic regions.

In this study, we identify a loss-of-function allele of the maize *CASTOR* ortholog,

confirm it confers resistance to AMF and selectively incorporate it into a biparental population

for quantitative trait locus (QTL) mapping. The resulting population contained both AM

susceptible and resistant families, allowing us to map QTL for agronomically important yield

components in mycorrhizal and non-mycorrhizal "environments". Our results indicate AM

symbiosis tp makes a substantial contribution to maize performance in the field. Furthermore, we

present an empirical demonstration of the dependence and benefit variation, as well as evidence

of a trade-off between the two.

RESULTS

Mutation in a maize common symbiosis gene demonstrates the importance of arbuscular

mycorrhizae in the field

To generate AMF-resistant maize varieties for use as a control in field experiments, we identified

maize orthologs of the common-symbiosis genes *CASTOR* and *POLLUX* (*DMI1*) (Parniske,

2008). *CASTOR* and *POLLUX* were found to be single copy genes in maize, and designated

Hunahpu (*Hun1*; GRMZM2G099160; Zm00001d012863) and *Xbalanque* (*Xba1*;

GRMZM2G110897; Zm00001d042694) after the hero twins of Mayan mythology (Ramírez-

Flores, 2015). We searched publicly available genetic resources (McCarty et al., 2005) and

identified a transposon-insertion allele of the *Hun1* gene (*hun1-2::mu,* hereafter, *hun1*) that we

found to confer resistance to mycorrhizal colonization (Ramírez-Flores, 2015). We then crossed

the *hun1* mutant in its original temperate W22 background to the subtropical inbred line

CML312 and advanced material to generate a mapping population of 73 wild-type (susceptible;

AMF-S) and 64 *hun1* (resistant; *AMF-R*) $F_{2:3}$ families (Fig. 1A, S2). We confirmed that AM

resistance was stable in the field, colonization being absent or greatly reduced in *AMF-R* families

in a preliminary trial (Fig. S3). In a second preliminary experiment, we grew *AMF-S* and *AMF-R*

families under highly managed nutrient-replete field conditions and found there to be no major

difference in performance, indicating that *hun1* does not condition a general defect in plant

growth or show marked pleiotropic effects beyond the block in mycorrhizal colonization (Fig.

S4, S5).

To estimate the importance of AM symbiosis, we evaluated *AMF-S* and *AMF-R* families

in a replicated trial under rain-fed, medium-input, conditions in Ameca, Mexico (Fig. 1A, S4, S6;

triplicated three-row plots, ~1,200 rows of 15 individuals for a total of ~ 20, 000 plants). We

observed *AMF-R* families to generally exhibit mild-chlorosis (Fig. 1B, S7) and to produce

visibly poorer ears than *AMF-S* families (Fig. 1C-D). We collected fifteen phenological and

morphological traits and yield-components (Fig. 1E-H, S8; Table S1), eleven of which showed a

significant marginal difference between *AMF-S* and *AMF-R* families, equating to host response

ranging from 3 to 51 % (Table 1). *AMF-R* families were reduced in stature and delayed in silking

(female flowering), resulting in an extension of anthesis-silking interval (ASI; Fig 1E, F; Table

S1) - a classic symptom of abiotic stress in maize. Ear size was reduced in *AMF-R* families,

along with ear and total kernel weight (Table 1; Fig. S9). Individual kernels were not

significantly different in weight or size between *AMF-S* and *AMF-R* families (Fig. 1G; Table 1).

The total number of kernels per ear, however, was greatly reduced in *AMF-R* families (Fig. 1H;

Table 1), indicative of poor seed set, a possible consequence of increased ASI.

QTL x AMF effects underlie variation in mycorrhiza response

To characterize the genetic architecture of host response, we performed a Quantitative Trait

Locus (QTL) analysis using the trait estimates obtained for the $F_{2:3}$ families and the genotypes of

their respective F_2 parents (Fig. S10, S11). We combined *AMF-S* and *AMF-R* families in a single

analysis, treating the *Hun* genotype as an interactive covariate. Under this model, the *Hun*

additive effect (hereafter, AMF effect) estimated the marginal host response across all families,

QTL additive effects captured genetic differences between the parents, and QTL x AMF effects

indicated underlying heritable differences in the response of CML312 and W22 parents. We

identified a total of 24 QTLs, 17 of which also showed AMF x QTL interaction (Fig. 2A, S12;

Table 1). We combined significant QTL, AMF and AMF x QTL terms, into a single multiple-

QTL model for each trait and calculated the percentage of phenotypic variation explained by

AMF, by additive QTL and by AMF x QTL terms (Fig. 2B; Table S2). For plant height, ear

weight and total kernel weight, the combination of AMF and AMF x QTL effects explained

more than half of the total genetic variation (based on estimation of H^2), and over a quarter of the

total phenotypic variance (Fig. 2B). The identification of significant AMF x QTL effects reveals

genetic variation for host response between the CML312 and W22 parents.

The genetic architecture of response variation distinguishes dependence and benefit

Variation in host response variation confounds differences in dependence and benefit (Fig. 3A,

B, S13; Janos, 2007; Sawers et al., 2010) - the former being the capacity of a given variety to

perform in the absence of AMF, the latter the degree to which a plant host profits from the

association. Having identified significant AMF x QTL interaction, we distinguished dependence

and benefit by estimating the size of the QTL effect separately for *AMF-S* and *AMF-R* families

(Fig. S14). For 12 of the 17 QTLs showing significant interaction, the effect was *conditional -*

i.e. expressed in either the *AMF-S* or *AMF-R* families, but not in both (Fig. 2A, 3; Table 1; Des

Marais and Juenger, 2010). We considered conditional QTLs expressed specifically in the *AMF-*

R families to represent variation in dependence. Conversely, we considered conditional QTLs

expressed specifically in the *AMF-S* families to be variation in benefit (Fig. 2, 3; Table 1). Two

regions of the genome were associated with multiple QTLs, suggesting a shared mechanistic

basis: a region on the long arm of chromosome (chr) 1 linked to ear-traits and a region on chr 4

linked to plant height and flowering time (Fig. 2A). Combining traits using Principal

Components (PCs) further refined these two QTL hot-spots and clearly demonstrated the

difference between dependence and benefit (Fig. 3D, S15. Table 1).

Antagonistic QTL effects suggest a trade-off between mycorrhizal and non-mycorrhizal

growth

Several QTLs showed a more extreme form of QTL x AMF in which the effect was expressed in

both *AMF-S* and *AMF-R* populations, but with a change of sign (*i.e.* a "swap" in the relative

performance of the parental alleles) - a condition described as *antagonistic pleiotropy* (Fig. 4A,

S13, S14; Des Marais and Juenger, 2010). Under such genetic architecture, the superior allele for

mycorrhizal plants is detrimental to performance in the absence of mycorrhiza, and *vice versa*,

providing evidence for a trade-off at the single-locus level. Significantly, several QTLs

associated with key yield components showed antagonistic pleiotropy (Fig. 2, 4B, 4C, S14;

Table 1). Analogous "rank changing" effects were also seen at the level of the whole genotype.

Although the mapping design did not permit direct evaluation of the same genotype as both

AMF-S and *AMF-R*, we could fit QTL models to all genotypes across both levels of AMF. The

resulting estimates showed evidence of rank change with respect to performance with or without

AMF (Fig. 4D). To further explore response at the genotype level, we used whole genome

models to predict mycorrhizal and non-mycorrhizal trait values across the 137 families.

Specifically, we used the *AMF-S* families to train a model for mycorrhizal performance that was

then applied across the population. Similarly, we used the *AMF-R* families to train a model for

non-mycorrhizal performance. Whole genome predictions aligned well with the observed values

of their training sets, although they did not capture the more extreme observed values (Fig. 4E).

Comparison of the two models indicated that genotypes associated with the highest values of the

major yield components in one condition were unexceptional or poor in the other (Fig. 4F),

indicating trade-off between dependence and benefit.

DISCUSSION

We have presented evidence that AM symbiosis makes a significant contribution to maize

performance. Extrapolating our observations to a cultivated field planted at 80,000 plants/ha

(typical in the locality of the trial site), we estimate that AM colonization would contribute ~ 2

tonnes/ha to a total yield of ~ 5.5 tonnes/ha. Although a rough figure to be treated with caution,

this estimate serves to provide a striking indication of the importance of AMF, and is consistent

with the previous evaluation of an inbred mutant line that linked loss of mycorrhizal phosphate

uptake with a 15% reduction in ear-weight (Willmann et al., 2013). Although maize varieties

have consistently been shown to benefit from AM symbiosis (*e.g.* Kaeppler et al., 2000; Sawers

et al., 2017), the impact of AMF in other plant species may vary along a continuum from

mutualism to parasitism (Bergmann et al., 2020; Johnson et al., 1997). In wild species, host

response is centered around neutral commensalistic interactions, with equal occurrence of

parasitism and mutualism (Klironomos, 2003). A similar range of host response has also been

reported in crop plants (Wen et al., 2019). In sorghum, a close relative of maize, host response

varies from positive to negative depending on both the plant variety and the AMF species

(Ramírez-Flores *et al*., unpublished observations; Watts-Williams et al., 2019). However, host

response has typically been measured in young plants, grown under controlled conditions, and

the values obtained may under-estimate the true importance of AM symbiosis in the field.

We found evidence for substantial variation in host response among plant genotypes.

Based on the range of fitted values obtained from the total grain weight multiple-QTL model, we

predict a range of yield response from 0.4 to 3.7 tonnes/ha at 80,000 plants/ha. In terms of

variance explained, AMF x QTL effects were more important than the QTL main effects for total

kernel weight, total kernel number and fifty-kernel weight. While we saw no evidence that our

W22 and CML312 parents, as "modern" inbred lines, have lost the capacity to profit from AM

symbiosis (*e.g.* see Koide et al., 1988; Hetrick et al., 1992), we did observe QTL linked to

differences in benefit, implying that it would be possible to select improve the capacity of plants

to profit from the symbiosis. Further study will be required to understand the mechanistic basis

of the differences we saw in host response. Although the level of AM colonization has been

reported to differ among varieties in a number of plant species, it is not necessarily an indicator

of greater host response (Kaeppler et al., 2000; Sawers et al., 2017; Watts-Williams et al., 2019;

Plouznikoff et al., 2019; Pawlowski et al., 2020). Indeed, the capacity of the host to limit fungal

colonization under certain conditions may maximize plant benefit, and it is well established that

colonization is reduced under nutrient replete conditions (Karlo et al., 2020; Nouri et al., 2014;

Sawers et al., 2017). As much as root-internal fungal growth, the extent of the root-external

hyphal network impacts host response (Yao et al., 2001; Munkvold et al., 2004; Schnepf et al.,

2008) and may be a factor in benefit variation(Sawers et al., 2017). Recently, it has been shown

that certain bacterial taxa exert beneficial and synergistic effects on AM symbiosis, acting as so-

called "mycorrhiza helpers" (Frey-Klett et al., 2007; Heijden et al., 2016; Ferreira et al., 2020).

Characterization of the root microbiome of *AMF-S* and *AMF-R* families will provide insight into

the interplay between AMF, the broader microbial community and host response.

We predicted trade-offs between host dependence and benefit: the genotypes predicted to

be the best in AM susceptible families were predicted to be unremarkable or poor in the context

of AM resistance, and *vice versa*. At the level of individual loci, examples of antagonistic

pleiotropy (Des Marais and Juenger, 2010) were seen for QTL associated with days-to-silking,

ear weight, total kernel weight, number of kernels per row and fifty kernel weight. Our

identification of intraspecific genetic trade-offs is analogous to patterns of plant nutritional

strategy seen at higher taxonomic levels where the differing demands placed on plant anatomy

and physiology by direct foraging and engagement in AM symbiosis appear to prevent co-

optimization for both (Lambers et al., 2008; Wen et al., 2019). Detailed characterization of root

anatomy has suggested that such trade-offs will exist: for example, increasing the proportion of

root cortical aerenchyma (root air space) reduces the carbon demand of the root system,

promoting foraging efficiency (Postma and Lynch, 2011), but, potentially, at the cost of limiting

AMF colonization and host response (Strock et al., 2019). It was somewhat surprising to see a

QTL for a morphological trait such as tassel branch number to be conditional on AM

susceptibility; such observations may indicate the action of pleiotropic systemic signals. Indeed,

the phytohormones so important to the regulation of plant development (including patterning of

tassel architecture) are also implicated in the establishment and regulation of AM symbiosis

(Gutjahr, 2014; Bonfante and Genre, 2015). For example, disruption of ethylene signalling has

been shown to modulate the level of (negative) host response to AMF in *Nicotiana attenuata*

(Riedel et al., 2008); a DELLA repressor in the gibberellic acid signalling pathway is essential

for AM colonization in rice (Yu et al., 2014); a feedback-loop involving strigolactone modulates

the extent of AMF colonization in *Medicago truncatula* (Müller et al., 2019). It may be

significant that many of these same signals have been the targets of domestication and plant

improvement (Sawers et al., 2018).

QTL x AMF effects, including those associated with "trade-off", were primarily driven

by plasticity of the temperate W22 alleles. It was interesting that CML312 alleles, derived from

the parent better adapted to the location of our trial, were more stable with respect to the

presence or absence of AMF. It has been hypothesised that mutualisms will be strengthened

under stressful conditions and that they might play an important role as non-adapted genotypes

move into new environments (O'Brien et al., 2018, 2019). Beyond an impact on yield in any

given site, the genetic architecture described here implies a role for the AMF in influencing yield

stability. In the face of variation and unpredictability in the AM community from site to site, or

year to year, "optimization" of the plant host might reasonably place a greater emphasis on

consistency over maximum benefit. Although we have found no evidence of a selective pressure

to minimize the AM symbiosis in maize breeding, the trade-offs we have described suggest that

efforts to optimize plants for better performance either with or without the contribution of

microorganisms may not be easily aligned. Ultimately, we may need to make decisions about the

types of agroecosystems we want to create and develop our crop varieties accordingly.

MATERIALS AND METHODS

Plant material and generation of the $F_{2:3}$ mapping population

An $F_{2:3}$ population was developed from the cross between a W22 stock homozygous for the

hun1-2 allele (*Mutator* insertion mu::1045305 in the gene

GRMZM2G099160/Zm00001d012863. www.maizegdb.org. Ramírez-Flores MR. M.Sc. thesis,

2015) and the subtropical CIMMYT inbred line CML312. The F_1 was self-pollinated to generate

an F_2 segregating *hun1-2* and W22/CML312 genome content. F_2 plants were genotyped by PCR

to identify homozygous wild-type and mutant individuals, and these self-pollinated to generate

susceptible (*AMF-S*) and resistant (*AMF-R*) $F_{2:3}$ families, respectively. $F_{2:3}$ families were

increased by sibling-mating. Material was generated in Valle de Banderas, Nayarit, Mexico and

Irapuato, Guanajuato, Mexico from 2016 to 2019. Small amounts of seed are available for

distribution through direct contact with the authors.

DNA extraction and PCR genotyping of hun1-2

DNA extraction was performed as described in (Fulton et al., 1995). The genotype at the *Hun*

locus was determined by running wild-type and mutant PCR reactions. To amplify the wild-type

fragment, a pair of gene-specific primers were used (HUNF01-

CGCGAAGAAACGCAGACATTCC and HUNR04-TAACCTGGAGCGAACAGAATCCAC),

generating a product of 606 bp. To amplify the mutant fragment a combination of *Mutator*

primer and gene-specific primer were used (HUNF03-CTTGGGCGCATTGGAAATTCATCG

and RS183-CGCCTCCATTTCGTCGAATCCSCTT), generating a fragment of 839 bp. PCR

conditions were: 1 cycle of initial incubation at 94ºC for 3 min, 32 cycles of 94ºC for 30 sec,

60ºC for 30 sec and 72ºC for 1 min, and 1 cycle of a final extension at 72ºC for 5 min. PCR used

the Kapa Taq PCR kit from Kapa Biosystems (Wilmington, Massachusetts, United States)

following the manufacturer's instructions. Products were visualized on 1 % agarose gel.

Whole genome genotyping and genetic map construction

Approximately 200 F_2 parents were analyzed using the Illumina MaizeLD BeadChip

(https://www.illumina.com/products/by-type/microarray-kits/maize-ld.html). Over 3,000 single

nucleotide polymorphisms (SNPs) were detected, with a call rate of> 96% for all samples. The

SNP set was processed using TASSEL 5 (Bradbury et al., 2007). SNPs were transformed to

ABH format (A: CML312; B: W22) and filtered to remove sites matching any of the following

criteria: > 30% missing data; monomorphic sites outside of the *Hun* locus; SNPs outside the *Hun*

locus showing segregation distortion (X^2 test, p < 0.05); redundant sites. The markers

"5_2269274", "5_3096229" and "5_4270584" (physical position on chromosome 5 at 2.67, 3.1

and 4.27 Mb, respectively) were used to confirm the expected genotype at the *Hun* locus. The

genetic map was built in R (R core team 2020) using R/ASMAP (Taylor and Butler, 2017) and

R/qtl (Broman et al., 2003); (Broman and Sen, 2009a).

Field Evaluation and phenotypic analysis

The $F_{2:3}$ population was evaluated in the summer of 2019 at the UNISEM experimental station in

Ameca, Jalisco, Mexico (20.78, -105.243). Three complete blocks of 73 *AMF-S* and 64 *AMF-R*

families were evaluated. Within each block, *AMF-S* and *AMF-R* families were alternated. The

order of the families within the *AMF-S* or *AMF-R* sub-populations was randomized within each

block. For each replicate, 45 seeds of each family were sown in a plot of three 2 m long rows.

Only the second row of each plot was considered for evaluation, the first and third rows

providing a buffer between adjacent families. Phenotypic data were collected at flowering and

after harvest as follows: stand count (STD), number of plants per row; days to anthesis (DTA),

number of days from planting until anthers visible on the main spike of half of the plants in the

row; days to silking (DTS), number of days from planting until silks visible on half of the plants

in the row; anthesis-silking interval (ASI), difference in days between anthesis and silking; plant

height (PH), distance in cm from the ground to the flag leaf; tassel branch number (TBN),

number of primary lateral branches originating from the main spike; ear weight (EW), the weight

in g of the ear; ear length (EL), distance in centimeters from the base to the tip of the ear; ear

diameter (ED), the diameter in cm at the middle part of the ear; kernel row number (KRN),

number of rows in the middle part of the ear; kernels per row (KPR), number of kernels in a

single row from the base to the tip of the ear; cob diameter (CD), diameter in cm at the middle

part of the cob; fifty kernel weight (FKW), weight in g of 50 randomly selected kernels; total

kernel weight (TKW), weight in g of all kernels on the ear; total kernel number (TKN), estimated

from TKW/FKW * 50.

Image analysis

Image-based phenotyping method was used to quantify the size and shape of maize ear, cob, and

kernel. Ears and cobs (1-3) from each plot, and approximately 50 kernels from associated ears,

were scanned using an Epson Perfection V600 scanner at 1200 dots per inch. Scanned images

were uploaded to Cyverse (Merchant et al., 2016) and analyzed using an automatic pipeline to

quantify ear, cob, and kernel attributes (Miller et al., 2017). Width profiles of ear and cob and

contour data of kernels were extracted in R to compute their shapes.

Principal Component Analysis

Principal Component Analysis (PCA) was conducted to analyze major sources of plant

phenotypic variance and to visualize the relationships among all measured phenotypic traits

across two maize families. PCA was performed using the prcomp function in R.

QTL mapping and whole genome prediction

QTL mapping was performed using the R/qtl package (Broman et al., 2003) on a population

composed of 73 AMF-S and 64 AMF-R F2:3 families. Single-QTL standard interval mapping

was run using the scanone function with Haley-Knott regression, with the genotype at *Hun* (wt

=1; *Hun* = 0) treated as a covariate (hereafter, AMF). To assess QTL x AMF interaction

(Broman and Sen, 2009), QTL analysis was performed under four different models: separate

analyses for AMF-S (H_{0wt}) and AMF-R (H_{0hun}) subpopulations; considering AMF as an additive

covariate (H_a) and considering AMF as an interactive covariate (H_f). The LOD significance

threshold for each trait and model was established with a permutation test ($\alpha = 0.05$, 1000

permutations). Evidence of QTL x AMF interaction was obtained by comparing H_f and H_a

models, where the difference $LOD_i = LOD_f - LOD_a$ was considered with reference to the

threshold difference $LOD\_thr_i = LOD\_thr_f - LOD\_thr_a$ for evidence of possible interaction.

Individual QTL were combined into multiple-QTL models on a per trait basis. Where evidence

was found of QTL x AMF interaction in the single scan, the interaction was also included in the

multiple-QTL model. Multiple-QTL models were evaluated using the fitqtl function and non-

significant (p < 0.1) terms removed according to the drop-one table.

To compare the potential of a given genotype in the presence or absence of AM

colonization, we estimated the performance of the 137 families for both levels of AMF using two

approaches. In the first approach, we generated fitted values from the multiple-QTL models,

under both levels of the AMF covariate. In the second approach, we performed whole genome

prediction using the R/rrBLUP package (Endelman, 2011). Missing genotypes were imputed

with the A.mat function and marker effects and BLUE values were obtained with the

mixed.solve function. The observed values for AMF-S families were used to train a mycorrhizal

model, that was then applied to the genotypes of the AMF-R families; the observed values from

the AMF-R families were used to train a non-mycorrhizal model, that was then applied to the

genotypes of the AMF-S families. For interpretation of the results, predicted values were

compared for all genotypes in both levels of AMF, avoiding possible bias in extreme observed

values.

ACKNOWLEDGEMENTS

ACKNOWLEDGEMENTS

We thank Beda Angehrn and Mario Rivera (UNISEM) for assistance with field evaluation, Cruz

Robledo (PV Winter Nurseries) for support with generating material, and Jessica Carcaño-

Macias for technical support and seed stock management. We acknowledge Karina Picazarri-

Delgado for assistance with preliminary characterization of the *hun* mutants and Nathan Miller

for his help with image analysis. This study was funded by the Mexican National Commission

for the Study and Use of Biodiversity (CONABIO) project *Impact of native arbuscular*

mycorrhizal fungi on maize performance (Nº 62, 2016-2018). M. Rosario Ramírez-Flores was

supported by a Ph.D. scholarship from the Mexican National Council of Science and Technology

(CONACYT).

REFERENCES

Augé RM, Toler HD, Saxton AM. 2015. Arbuscular mycorrhizal symbiosis alters stomatal conductance of host plants more under drought than under amply watered conditions: a meta-analysis. *Mycorrhiza* **25**:13–24.

Bender SF, Plantenga F, Neftel A, Jocher M, Oberholzer H-R, Köhl L, Giles M, Daniell TJ, van der Heijden MG. 2014. Symbiotic relationships between soil fungi and plants reduce N2O emissions from soil. *ISME J* **8**:1336–1345.

Bender SF, Wagg C, van der Heijden MGA. 2016. An Underground Revolution: Biodiversity and Soil Ecological Engineering for Agricultural Sustainability. *Trends Ecol Evol* **31**:440–452.

Bergmann J, Weigelt A, van der Plas F, Laughlin DC, Kuyper TW, Guerrero-Ramirez N, Valverde-Barrantes OJ, Bruelheide H, Freschet GT, Iversen CM, Kattge J, McCormack ML, Meier IC, Rillig MC, Roumet C, Semchenko M, Sweeney CJ, van Ruijven J, York LM, Mommer L. 2020. The fungal collaboration gradient dominates the root economics space in plants. *Sci Adv* **6**:eaba3756.

Bonfante P, Genre A. 2015. Arbuscular mycorrhizal dialogues: do you speak "plantish" or "fungish"? *Trends in Plant Science*. doi:10.1016/j.tplants.2014.12.002

Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. 2007. TASSEL: software for association mapping of complex traits in diverse samples. *Bioinformatics*. doi:10.1093/bioinformatics/btm308

Broman KW, Sen S. 2009. A Guide to QTL Mapping with R/qtl. *Statistics for Biology and Health*. doi:10.1007/978-0-387-92125-9

Broman KW, Wu H, Sen S, Churchill GA. 2003. R/qtl: QTL mapping in experimental crosses. *Bioinformatics* **19**:889–890.

Cavagnaro TR, Bender SF, Asghari HR, Heijden MGA van der. 2015. The role of arbuscular mycorrhizas in reducing soil nutrient loss. *Trends Plant Sci* **20**:283–290.

Chandrasekaran M, Boughattas S, Hu S, Oh S-H, Sa T. 2014. A meta-analysis of arbuscular mycorrhizal effects on plants grown under salt stress. *Mycorrhiza* **24**:611–625.

Charpentier M, Bredemeier R, Wanner G, Takeda N, Schleiff E, Parniske M. 2008. Lotus japonicus CASTOR and POLLUX are ion channels essential for perinuclear calcium spiking in legume root endosymbiosis. *Plant Cell* **20**:3467–3479.

Chiu CH, Paszkowski U. 2019. Mechanisms and Impact of Symbiotic Phosphate Acquisition. *Cold Spring Harb Perspect Biol* **11**. doi:10.1101/cshperspect.a034603

Des Marais DL, Juenger TE. 2010. Pleiotropy, plasticity, and the evolution of plant abiotic stress tolerance. *Ann N Y Acad Sci* **1206**:56–79.

Faye A, Dalpé Y, Ndung'u-Magiroi K, Jefwa J, Ndoye I, Diouf M, Lesueur D. 2013. Evaluation of commercial arbuscular mycorrhizal inoculants. *Can J Plant Sci* **93**:1201–1208.

Ferreira DA, da Silva TF, Pylro VS, Salles JF, Andreote FD, Dini-Andreote F. 2020. Soil Microbial Diversity Affects the Plant-Root Colonization by Arbuscular Mycorrhizal Fungi. *Microb Ecol*. doi:10.1007/s00248-020-01502-z

Fester T, Sawers R. 2011. Progress and Challenges in Agricultural Applications of Arbuscular Mycorrhizal Fungi. *Critical Reviews in Plant Sciences*. doi:10.1080/07352689.2011.605741

Frey-Klett P, Garbaye J, Tarkka M. 2007. The mycorrhiza helper bacteria revisited. *New Phytol* **176**:22–36.

Fulton TM, Chunwongse J, Tanksley SD. 1995. Microprep protocol for extraction of DNA from

tomato and other herbaceous plants. *Plant Molecular Biology Reporter*. doi:10.1007/bf02670897

Gutjahr C. 2014. Phytohormone signaling in arbuscular mycorhiza development. *Curr Opin Plant Biol* **20**:26–34.

Gutjahr C, Banba M, Croset V, An K, Miyao A, An G, Hirochika H, Imaizumi-Anraku H, Paszkowski U. 2008. Arbuscular mycorrhiza-specific signaling in rice transcends the common symbiosis signaling pathway. *Plant Cell* **20**:2989–3005.

Heijden MGA van der, van der Heijden MGA, de Bruin S, Luckerhoff L, van Logtestijn RSP, Schlaeppi K. 2016. A widespread plant-fungal-bacterial symbiosis promotes plant biodiversity, plant nutrition and seedling recruitment. *The ISME Journal*. doi:10.1038/ismej.2015.120

Hetrick BAD, Wilson GWT, Cox TS. 1992. Mycorrhizal dependence of modern wheat varieties, landraces, and ancestors. *Canadian Journal of Botany*. doi:10.1139/b92-253

Janos DP. 2007. Plant responsiveness to mycorrhizas differs from dependence upon mycorrhizas. *Mycorrhiza*. doi:10.1007/s00572-006-0094-1

Johnson NC, Graham JH, Smith FA. 1997. Functioning of mycorrhizal associations along the mutualism-parasitism continuum. *New Phytologist*. doi:10.1046/j.1469-8137.1997.00729.x

Kaeppler SM, Parke JL, Mueller SM, Senior L, Stuber C, Tracy WF. 2000. Variation among Maize Inbred Lines and Detection of Quantitative Trait Loci for Growth at Low Phosphorus and Responsiveness to Arbuscular Mycorrhizal Fungi. *Crop Sci* **40**:358.

Karlo M, Boschiero C, Landerslev KG, Blanco GS, Wen J, Mysore KS, Dai X, Zhao PX, de Bang TC. 2020. The CLE53-SUNN genetic pathway negatively regulates arbuscular mycorrhiza root colonization in Medicago truncatula. *J Exp Bot*. doi:10.1093/jxb/eraa193

Klironomos JN. 2003. VARIATION IN PLANT RESPONSE TO NATIVE AND EXOTIC ARBUSCULAR MYCORRHIZAL FUNGI. *Ecology*. doi:10.1890/02-0413

Koide R, Li M, Lewis J, Irby C. 1988. Role of mycorrhizal infection in the growth and reproduction of wild vs. cultivated plants. *Oecologia*. doi:10.1007/bf00377271

Lambers H, Raven JA, Shaver GR, Smith SE. 2008. Plant nutrient-acquisition strategies change with soil age. *Trends Ecol Evol* **23**:95–103.

Lynch JP. 2019. Root phenotypes for improved nutrient capture: an underexploited opportunity for global agriculture. *New Phytol* **223**:548–564.

Mascher M, Gerlach N, Gahrtz M, Bucher M, Scholz U, Dresselhaus T. 2014a. Sequence and ionomic analysis of divergent strains of maize inbred line B73 with an altered growth phenotype. *PLoS One* **9**:e96782.

Mascher M, Gerlach N, Gahrtz M, Bucher M, Scholz U, Dresselhaus T. 2014b. Sequence and ionomic analysis of divergent strains of maize inbred line B73 with an altered growth phenotype. *PLoS One* **9**:e96782.

McCarty DR, Settles AM, Suzuki M, Tan BC, Latshaw S, Porch T, Robin K, Baier J, Avigne W, Lai J, Messing J, Koch KE, Hannah LC. 2005. Steady-state transposon mutagenesis in inbred maize. *Plant J* **44**:52–61.

Merchant N, Lyons E, Goff S, Vaughn M, Ware D, Micklos D, Antin P. 2016. The iPlant Collaborative: Cyberinfrastructure for Enabling Data to Discovery for the Life Sciences. *PLoS Biol* **14**:e1002342.

Miller ND, Haase NJ, Lee J, Kaeppler SM, de Leon N, Spalding EP. 2017. A robust, high-throughput method for computing maize ear, cob, and kernel attributes automatically from images. *Plant J* **89**:169–178.

Müller LM, Flokova K, Schnabel E, Sun X, Fei Z, Frugoli J, Bouwmeester HJ, Harrison MJ.
2019. A CLE-SUNN module regulates strigolactone content and fungal colonization in
arbuscular mycorrhiza. *Nat Plants* **5**:933–939.
Munkvold L, Kjøller R, Vestberg M, Rosendahl S, Jakobsen I. 2004. High functional diversity
within species of arbuscular mycorrhizal fungi. *New Phytologist*. doi:10.1111/j.1469-
8137.2004.01169.x
Nouri E, Breuillin-Sessoms F, Feller U, Reinhardt D. 2014. Phosphorus and nitrogen regulate
arbuscular mycorrhizal symbiosis in Petunia hybrida. *PLoS One* **9**:e90841.
O'Brien AM, Sawers RJH, Ross-Ibarra J, Strauss SY. 2018. Evolutionary Responses to
Conditionality in Species Interactions across Environmental Gradients. *The American
Naturalist*. doi:10.1086/700118
O'Brien AM, Sawers RJH, Strauss SY, Ross-Ibarra J. 2019. Adaptive phenotypic divergence in
an annual grass differs across biotic contexts. doi:10.1101/382770
Parniske M. 2008. Arbuscular mycorrhiza: the mother of plant root endosymbioses. *Nat Rev
Microbiol* **6**:763–775.
Paszkowski U, Boller T. 2002. The growth defect of lrt1 , a maize mutant lacking lateral roots,
can be complemented by symbiotic fungi or high phosphate nutrition. *Planta*.
doi:10.1007/s004250100642
Pawlowski ML, Vuong TD, Valliyodan B, Nguyen HT, Hartman GL. 2020. Whole-genome
resequencing identifies quantitative trait loci associated with mycorrhizal colonization of
soybean. *Theor Appl Genet* **133**:409–417.
Plouznikoff K, Asins MJ, de Boulois HD, Carbonell EA, Declerck S. 2019. Genetic analysis of
tomato root colonization by arbuscular mycorrhizal fungi. *Ann Bot* **124**:933–946.
Postma JA, Lynch JP. 2011. Root cortical aerenchyma enhances the growth of maize on soils
with suboptimal availability of nitrogen, phosphorus, and potassium. *Plant Physiol*
156:1190–1201.
Ramírez-Flores MR. 2015. Caracterización genética de mutantes de HUN e IXBA, ortólogos de
maíz de los canales de potasio CASTOR y POLLUX. Centro de Investigación y de Estudios
Avanzados del Instituto Politécnico Nacional.
Riedel T, Groten K, Baldwin IT. 2008. Symbiosis between Nicotiana attenuata and Glomus
intraradices: ethylene plays a role, jasmonic acid does not. *Plant Cell Environ* **31**:1203–
1213.
Rillig MC, Aguilar-Trigueros CA, Camenzind T, Cavagnaro TR, Degrune F, Hohmann P,
Lammel DR, Mansour I, Roy J, van der Heijden MGA, Yang G. 2019. Why farmers should
manage the arbuscular mycorrhizal symbiosis. *New Phytol*.
Rillig MC, Mummey DL. 2006. Mycorrhizas and soil structure. *New Phytol* **171**:41–53.
Ryan MH, Graham JH. 2018. Little evidence that farmers should consider abundance or diversity
of arbuscular mycorrhizal fungi when managing crops. *New Phytol* **220**:1092–1107.
Sawers RJH, Gebreselassie MN, Janos DP, Paszkowski U. 2010. Characterizing variation in
mycorrhiza effect among diverse plant varieties. *Theor Appl Genet* **120**:1029–1039.
Sawers RJH, Gutjahr C, Paszkowski U. 2008. Cereal mycorrhiza: an ancient symbiosis in
modern agriculture. *Trends Plant Sci* **13**:93–97.
Sawers RJH, Ramírez-Flores MR, Olalde-Portugal V, Paszkowski U. 2018. The impact of
domestication and crop improvement on arbuscular mycorrhizal symbiosis in cereals:
insights from genetics and genomics. *New Phytol* **220**:1135–1140.
Sawers RJH, Svane SF, Quan C, Grønlund M, Wozniak B, Gebreselassie M-N, González-Muñoz

E, Chávez Montes RA, Baxter I, Goudet J, Jakobsen I, Paszkowski U. 2017. Phosphorus acquisition efficiency in arbuscular mycorrhizal maize is correlated with the abundance of root-external hyphae and the accumulation of transcripts encoding PHT1 phosphate transporters. *New Phytol* **214**:632–643.

Schnepf A, Roose T, Schweiger P. 2008. Impact of growth and uptake patterns of arbuscular mycorrhizal fungi on plant phosphorus uptake—a modelling study. *Plant and Soil*. doi:10.1007/s11104-008-9749-3

Strock CF, Schneider HM, Galindo-Castañeda T, Hall BT, Van Gansbeke B, Mather DE, Roth MG, Chilvers MI, Guo X, Brown K, Lynch JP. 2019. Laser ablation tomography for visualization of root colonization by edaphic organisms. *J Exp Bot* **70**:5327–5342.

Strullu-Derrien C, Selosse M-A, Kenrick P, Martin FM. 2018. The origin and evolution of mycorrhizal symbioses: from palaeomycology to phylogenomics. *New Phytol* **220**:1012–1030.

Taylor J, Butler D. 2017. R Package ASMap: Efficient Genetic Linkage Map Construction and Diagnosis. *Journal of Statistical Software*. doi:10.18637/jss.v079.i06

van der Heijden MGA, Martin FM, Selosse M-A, Sanders IR. 2015. Mycorrhizal ecology and evolution: the past, the present, and the future. *New Phytol* **205**:1406–1423.

Watts-Williams SJ, Emmett BD, Levesque-Tremblay V, MacLean AM, Sun X, Satterlee JW, Fei Z, Harrison MJ. 2019. Diverse Sorghum bicolor accessions show marked variation in growth and transcriptional responses to arbuscular mycorrhizal fungi. *Plant Cell Environ* **42**:1758–1774.

Wen Z, Li H, Shen Q, Tang X, Xiong C, Li H, Pang J, Ryan MH, Lambers H, Shen J. 2019. Tradeoffs among root morphology, exudation and mycorrhizal symbioses for phosphorus-acquisition strategies of 16 crop species. *New Phytol* **223**:882–895.

Willmann M, Gerlach N, Buer B, Polatajko A, Nagy R, Koebke E, Jansa J, Flisch R, Bucher M. 2013. Mycorrhizal phosphate uptake pathway in maize: vital for growth and cob development on nutrient poor agricultural and greenhouse soils. *Front Plant Sci* **4**:533.

Yao Q, Li X, Feng G, Christie P. 2001. Influence of extramatrical hyphae on mycorrhizal dependency of wheat genotypes. *Communications in Soil Science and Plant Analysis*. doi:10.1081/css-120001122

Yu N, Luo D, Zhang X, Liu J, Wang W, Jin Y, Dong W, Liu J, Liu H, Yang W, Zeng L, Li Q, He Z, Oldroyd GED, Wang E. 2014. A DELLA protein complex controls the arbuscular mycorrhizal symbiosis in plants. *Cell Res* **24**:130–133.

Zipfel C, Oldroyd GED. 2017. Plant signalling in symbiosis and immunity. *Nature* **543**:328–336.

Figure 1. Mycorrhizal symbiosis impacts maize growth under cultivation. A, Susceptible (*AMF-S*) and resistant (*AMF-R*) families segregate genomic content from two founder parents, shown as red and blue bars. *AMF-R* families are homozygous for the *hun1-2* mutation, blocking AM symbiosis (black arrows). B, Border between representative *AMF-S* and *AMF-R* plots, Jalisco, Mexico, 2019. C, Representative AMF-S ears. D, Representative AMF-R ears. E, Plant height (PH) of 73 *AMF-S* and 64 *AMF-R* families. F, Kernel shape (KS). G, Anthesis-silking interval (ASI). H, Total kernel number (TKN). The violin plots in **E, F** and **H** are a hybrid of boxplot and density plot. The box represents the interquartile range with the horizontal line representing the median and whiskers representing 1.5 times the interquartile ranges. The shape of the violin plot represents probability density of data at different values along the y-axis.

599
600

Figure 2. QTL x AMF effects contribute significantly to variation in host response. A)
Genomic position of QTL identified in this study. Boxes indicate the position of the peak
marker. Bars represent the drop-1 LOD interval. Colors denote patterns of AMF x QTL
interaction as described in the text. **B)** The contribution of different components to phenotypic
variance in different plant traits among *AMF-S* and *AMF-R* families. Total genetic variance (H^2)
was estimated based on differences between experimental blocks. Variance explained by the
QTL model for each trait is partitioned into that explained by the additive effect of AMF, the
additional variation explained by interactive QTL and QTL x AMF interaction (QTL x AMF)

610   and the additional variation explained by non-additive QTL. Traits codes as in Table 1.

611
Figure 3. Mycorrhiza response confounds benefit and dependence. A, host response (R) is
the difference in trait value between mycorrhizal (M; green) and non-mycorrhizal (NM, yellow)
plants. Increased response can result from either greater dependence (D; poorer performance of
NM plants) or greater benefit (B; greater performance of M plants). **B**, QTL x AMF effects
underlying variation in response reveal the balance of benefit and dependence. The difference in
trait value between plants carrying the CML312 (red) or W22 (blue) allele. In this theoretical
example, the effect is conditional on mycorrhizal colonization, reflecting a difference in benefit
more than dependence. **C, D,** effect plots for major QTL associated with PC1 and PC2,
respectively. The PC1 QTL is conditional on *AMF-R*, indicating a difference in dependence. The
PC2 QTL is conditional on *AMF-S*, indicating a difference in benefit.
622
623

Figure 4. The genetic architecture of mycorrhiza response implies a trade-off between dependence and benefit. A, QTL x AMF effects underlying variation in response reveal the balance of benefit and dependence. The difference in trait value between plants carrying the CML312 (red) or W22 (blue) allele. In this theoretical example, the effect is conditional on mycorrhizal colonization, reflecting a difference in benefit more than dependence. **B**, QTL effects for a total kernel weight (TKW) QTL located on Chromosome 1. TKW of families homozygous for the CML312 allele at this locus (red) is stable across susceptible (*AMF-S*) and resistant (*AMF-R*) families. In contrast, families homozygous for the W22 allele (blue) or segregating at this locus (brown), show superior TKW if *AMF-S* but lower TKW if *AMF-R*. In this extreme example of QTL x AMF, the QTL is linked to differences in both dependence and benefit, an example of antagonistic pleiotropy or "trade-off". **C**, Theoretical trade-off between the performance of non-mycorrhizal (NM) and mycorrhizal (M) plants. For any position on this plot, the vertical distance above the diagonal indicates the magnitude of host response. Trade-off prevents occupancy of the upper-right of the plot, defining a so-called "Pareto front" (solid arc). Here, an increase in the NM performance of variety X is necessarily accompanied by a reduction in M performance, as described by movement along the arc to condition Y. Biological constraint prevents the path towards condition Z which would represent NM improvement without any negative impact on M. **D**, Fitted ear weight values for all genotypes from the QTL model (2 QTL and their interactions with AMF) under both levels of AMF (*AMF-S* and *AMF-R*). Line segments

645 connect the same QTL genotype under the two AMF levels. **E**, Observed ear weight against
646 whole-genome prediction for 73 susceptible (*AMF-S*) and 64 resistant (*AMF-R*) families. Best fit
647 regression line in dashed blue. **F**, Predicted ear weights for all 137 genotypes based on *AMF-S*
648 (M) and *AMF-R* (NC) whole-genome models. Dashed red line shows *AMF-S* = *AMF-R*, *i.e.* no
649 host response.
650

651 **Table 1. Summary of host response and QTLs**

Trait	Mean AMF-S	Mean AMF-R	Adj. P-value[1]	HR[2]	HR (%)[3]	QTLs[4]	AMFxQTL[5]	Type[6]
STD	11	10	NS	1	10			
DTA	56	57	NS	-1	-1	1	1	S
DTS	59	60	***	-1	-3	1	1	AP
ASI	2	4	***	-2	-34	1	1	R
PH	196.8	171.8	***	25	15	2	2	R, S
TBN	13	12	NS	1	10	2	1	C, S
GC				0		1	0	C
EW	86.6	60.6	***	26	43	2	2	AP, S
EL	13.9	12.4	***	1.5	13	1	1	R
ED	40.3	37.2	***	3.1	8	2	2	R, R
KRN	16	15	*	1	6	1	0	C
KPR	25	18	***	7	37	2	2	R, AP
KC				0		1	0	C
CD	26.6	26.1	NS	0.5	2			
FKW	10.4	9.4	NS	1	10	2	2	R, AP
TKW	67.7	44.8	***	22.9	51	1	1	AP
TKN	330	230	***	100	44	1	1	R
PC1	-1.23	1.7	***			2		
PC2	0.126	-0.182	NS			1		
PC3	0.251	-0.288	*			1		
PC4	-0.033	0.037	NS			2		
PC5	-0.245	0.282	***			1		

[1]Wilcoxon tests with Bonferonni adjusted P-values. Note: *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$; NS:
not significant. [2]Host response calculated as mean *AMF-S - AMF-R*. [3]Host response expressed as

percentage of *AMF-R* mean. [4]Number of QTL associated with a given trait. [5]Number of QTL showing
significant QTL x AMF effect. [6]Pattern of QTL x AMF effect: R: conditional on *AMF-R*; S, conditional
on *AMF-S*; AP, antagonistic pleiotropy; C, constitutive. Trait codes: STD, stand count; DTA, days to
anthesis; DTS, days to silking; ASI, anthesis-silking interval; PH, plant height; TBN, tassel branch
number; EW, ear weight; EL, ear length; ED, ear diameter; KRN, kernel row number; KPR, kernels per
row; CD, cob diameter; FKW, fifty kernel weight; TKW, total kernel weight; TKN, total kernel number.
660

661
662