Differential spatial processing in ventral and lateral face-selective regions is scaffolded by

structural connections

Authors

Dawn Finzi[1], Jesse Gomez[2,3], Marisa Nordt[1], Alex A. Rezai[1], Sonia Poltoratski[1], and Kalanit Grill-

Spector[1,2,4]

[1] Department of Psychology, Stanford University, Stanford, CA

[2] Neurosciences Program, Stanford University, Stanford, CA

[3] Princeton Neuroscience Institute, Princeton University, NJ

[4] Wu Tsai Neurosciences Institute, Stanford University, Stanford, CA

Abstract

Face-processing occurs across ventral and lateral visual streams, which are involved in static and

dynamic face perception, respectively. However, the nature of spatial computations across

streams is unknown. Using functional MRI and novel population receptive field (pRF) mapping,

we measured pRFs in face-selective regions. Results reveal that spatial computations by pRFs in

ventral face-selective regions are concentrated around the center of gaze (fovea), but spatial

computations in lateral face-selective regions extend peripherally. Diffusion MRI reveals that

these differences are mirrored by a preponderance of white matter connections between ventral

face-selective regions and foveal early visual cortex (EVC), while connections with lateral regions

are distributed more uniformly across EVC eccentricities. These findings suggest a rethinking of

spatial computations in face-selective regions showing that they vary across ventral and lateral

streams, and further propose that spatial processing in high-level regions is scaffolded by the

fine-grain pattern of white matter connections from EVC.

Introduction

Face perception is crucial for every day social interactions and relies on brain computations across a series of face-selective regions in occipital and temporal cortex. In humans, face-selective regions span two processing streams[1–4]: the ventral stream and the lateral stream[2]. The ventral face processing stream is thought to be involved in face categorization, recognition, and identification[5–9]. Cortically, it begins in early visual cortex (EVC: union of V1, V2, V3), continues to the inferior aspects of occipital and temporal cortex, and contains several face-selective regions: one on the inferior occipital gyrus (IOG-faces, also referred to as the occipital face area, OFA[10]), and two on the fusiform gyrus—one on the posterior fusiform (pFus-faces[11]) and one in the mid fusiform (mFus-faces[11])—which are collectively referred to as the fusiform face area (FFA[5]). These regions project[12] to an anterior temporal face patch[3]. The lateral face processing stream is thought to be involved in perception of dynamic[3,13,14], transient[15], and changeable aspects[7] of faces including motion[13,14,16,17], expression[7,8], and gaze[18,19]. This stream also emerges in early visual cortex and then continues to superior temporal cortex. It consists of a region in the posterior superior temporal sulcus (pSTS-faces[1,11]) and a region on the main branch of the STS (mSTS-faces[1,11]), then projecting to a region in the anterior STS (aSTS-faces[11,12,20]). Interestingly, pSTS-faces neighbors hMT+[21], an area involved in processing motion, and mSTS-faces neighbors regions that process biological motion[22,23]. As the ventral and lateral face processing stream are optimized for different tasks, a central open question is: what are the basic computations separating these streams?

The basic computation in the visual system is spatial processing performed by the receptive field[24] (RF), which is the region in visual space that is processed by a neuron. As neurons with overlapping RFs are clustered, we can measure the population receptive field (pRF)[25], i.e., the region in visual space processed by the collection of neurons in an fMRI voxel. Additionally, how a region spatially processes a stimulus depends on the way pRFs in the region tile the visual field[26]. This tiling is referred to as the visual field coverage (VFC) of the region. Therefore, we asked: what are the properties of pRFs and VFC in face-selective regions of the human ventral and lateral processing streams?

One possibility is that face-selective regions in both lateral and ventral streams perform

similar spatial computations, and thus have similar pRF properties and VFC. As face perception

requires the high visual acuity of central (foveal) vision, people tend to fixate on faces[27–30]. This

habitual fixation on faces has led researchers to suggest that face-selective regions mainly

process foveal information[31,32]. This hypothesis (Eccentricity Bias Theory[31]) is supported by

findings that ventral face-selective regions respond more strongly to central than peripheral

visual stimuli[31–33], and have a foveal bias, that is, denser coverage of the central than peripheral

visual field[26,34]. In contrast, visual information related to other categories such as places, which

in the real world occupy the entire visual field, extends to the periphery of the visual field

irrespective of where people fixate. Supporting Eccentricity Bias Theory, researchers found that

place-selective regions are peripherally-biased[32,33]. Thus, Eccentricity Bias Theory predicts that

due to habitual fixation on faces, spatial processing in all face-selective regions, across ventral

and lateral processing streams, will be foveally-biased.

An alternative hypothesis predicts that spatial computations may differ across face-

selective regions in the ventral and lateral streams because these streams are optimized for

different tasks with different computational demands. Specifically, face recognition, associated

with the ventral stream, requires the fine spatial acuity afforded by central vision. This predicts

that ventral face-selective regions that are involved in face recognition will have foveally-biased

spatial computations[31,35]. In contrast, dynamic information in faces, which is associated with

processing in lateral face-selective regions, requires fast temporal computations of transient

visual information. As processing of visual information in the periphery is faster than in the

fovea[15,36,37], spatial computations in lateral face-selective regions may extend to the periphery.

To test these hypotheses, we designed a novel pRF mapping experiment optimized to

map pRFs in high-level visual regions (**Figure 1a**). Using these data, we estimated pRFs in each

voxel and then compared pRFs and VFC across face-selective regions in the ventral and lateral

streams.

We also considered an important, related question: how do spatial processing

characteristics and eccentricity biases emerge in high-level visual regions in the first place? A

prevalent view suggests that the hierarchical organization of visual processing streams, as well

as pooling operations from one stage to the next, generate successively larger pRFs in higher-

level stages of the processing hierarchy[38,39]. Indeed, large pRFs in the end stages of the processing

hierarchy may be optimized to process whole stimuli such as faces[40]. This hierarchical

organization may be supported by sequential connections along regions constituting the

processing steam[12,40,41]. However, other evidence shows that the human visual stream is not

strictly hierarchical[41–43] as there are skip connections[40] between areas that are not consecutive

in the hierarchy. In fact, recent studies report the existence direct white matter connections

between EVC and face-selective regions of both lateral and ventral processing streams[41,42], which

may enable fast visual processing[44].

93   However, the nature of direct connections from early stages of the visual hierarchy to

later stages and how they relate to spatial processing in downstream regions remains a mystery.

As EVC has a topographic representation of visual space, we examined the pattern of white

matter connections between specific eccentricity bands in EVC and face-selective regions. We

reasoned that to support large pRFs in high-level regions, there should be connections between

a range of EVC eccentricities and face-selective regions. However, we hypothesized that one

source of emergent spatial biases in face-selective regions could be an uneven distribution of

connections emerging from central and peripheral eccentricities in EVC. That is, we predicted

that regions that are foveally-biased will have a preponderance of white matter tracts emerging

from central compared to peripheral eccentricities of EVC. In contrast, regions in which spatial

processing extends to the periphery will have a more uniform distribution of connections across

EVC eccentricities. We tested these predictions in a second experiment, in which we used

diffusion MRI (dMRI) and fMRI in the same participants to determine the distribution of white

matter tracts between EVC eccentricity bands and each of the face-selective regions.

Results

"Toonotopy" (retinotopy with cartoons) drives responses in face-selective cortex more than

standard checkerboard retinotopy

To measure pRFs, 21 adults (10 female) participated in a wide-field pRF mapping experiment

which used colorful cartoon images ("toonotopy", **Figure 1A**). In the experiment, subjects fixated

on a central dot, and pressed a button when it changed color. As they performed this task, they

viewed a bar extending 40° in length that systematically swept the visual field (**Figure 1a**). Unlike

standard pRF mapping[25,34] that uses flickering checkerboards, our bars contained random images

from cartoons including colorful faces, objects, scenes, and text presented at a rate of 8Hz.

Additionally, subjects participated in a 10-category localizer experiment[45] to independently

define their face-selective regions.

Figure 1. **Toonotopy experiment and regions of interest.** (A) Experimental design. A bar containing colored cartoon images changing at 8 Hz is swept across a gray background in 4 orientations (0°, 45°, 90°, 135°), each in 2 directions orthogonal to the bar. Each sweep takes 24 seconds, with blanks after six bar steps (12 seconds) for each diagonal direction. Participants were instructed to fixate and indicate when the fixation dot changed color. (B) Example polar angle (left) and eccentricity (right) maps in a representative participant's inflated right hemisphere. *Inset:* zoomed medial view. Maps are thresholded at 10% variance explained,

voxel level. *Solid lines:* boundary of retinotopic areas. *Dashed contour:* early visual cortex: union of V1, V2 and V3.
(C) Proportion of voxels in each right hemisphere region in which pRF model explains >10% of their variance. Data
averaged across five participants who underwent both checkerboard retinotopy and toonotopy. *Error bars*: ±SE
(standard error of the mean). Asterisks indicate significant differences between toonotopy vs. checkerboard
retinotopy; post-hoc Tukey t-tests (** *p* < .01) (D) Face-selective regions defined from the localizer experiment in a
representative participant's inflated right hemisphere using contrast: faces vs. all other categories, *t*>3, voxel level.
Warm colors: ventral face-selective regions; *Cold colors*: lateral face-selective regions. *Acronyms*: *IOG:* inferior
occipital gyrus; *pFus:* posterior fusiform. *mFus:* mid fusiform; *OTS:* occipito-temporal sulcus; *ITS:* inferior temporal
sulcus; *STS:* superior temporal gyrus.

From the localizer experiment, we defined three ventral face-selective regions (IOG, pFus,

mFus-faces) and two lateral face-selective regions (pSTS, mSTS-faces) in each subject's brain

(**Figure 1D**, Methods). Additionally, we defined in each participant their ventral place-selective

region (CoS-places), which we include to replicate prior findings and as a control in certain

analyses. All regions were found bilaterally except for mSTS-faces, which was lateralized to the

right hemisphere in all but six participants. Due to the lack of mSTS-faces in the left hemisphere,

as well as the vast literature reporting lateralization of face-selectivity[3,46], we analyze data from

each hemisphere separately throughout this study, unless otherwise specified.

In the toonotopy pRF mapping experiment, we used colorful cartoon stimuli and fast

presentation rates in an attempt to maximally drive responses in high-level visual cortex. To

quantify the success of this approach, in five participants we compared toonotopy to a standard

checkerboard pRF mapping experiment, which was identical to the toonotopy experiment except

the bars contained black and white checkerboards that flickered at a rate of 2Hz. In each

participant and experiment, we fit a pRF for each voxel and identified the voxels for which the

pRF model explained at least 10% of their variance.

Qualitatively, toonotopy pRF mapping yielded the typical retinotopic polar angle and

eccentricity maps (**Figure 1B**). Along the calcarine we found a hemifield representation,

corresponding to V1, followed by mirror reversals of the polar angle (**Figure 1B**-left). Likewise,

we found the standard eccentricity map of the occipital lobe, with foveal representations close

to the occipital pole, and systematically more peripheral eccentricities proceeding from posterior

to anterior along the calcarine sulcus (**Figure 1B**-right). Using the polar angle map, we defined in

each subject V1, V2, V3 and their union (referred to as early visual cortex: EVC, **Figure 1B**-right).

Quantitatively, we compared the amount of retinotopically-modulated voxels across

experiments. Differences between the ability of mapping stimuli to drive high-level face-selective

regions are striking. Nearly 90% of mFus-faces voxels in the right hemisphere were driven in the

toonotopy experiment, compared to less than 25% in the checkboard experiment (**Figures 1C,**

S1-left hemisphere). As linear mixed models (LMMs) tolerate missing values[47] (e.g. when there

are missing regions of interest (ROIs)), here and in subsequent analyses, we applied LMMs to

analyze the data. The significance of fixed effects in these models were evaluated using analyses-

of-variance with Satterthwaite approximations for degrees of freedom[48] (referred to as LMM

ANOVA). A 2-way repeated measures LMM ANOVA on the proportion of retinotopically-driven

voxels with factors of experiment (toonotopy/checkerboards) and ROI

(V1/IOG/pFus/mFus/pSTS/mSTS) revealed a significant ROI x experiment interaction in both the

right ($F(5, 44)=5.0$, *p*=.0010) and left hemispheres ($F(5, 40)=3.9$, *p*=.0057) as the type of mapping

experiment had a pronounced effect on driving responses in face-selective regions, but not V1

(**Figures 1C, S1**).

pRF centers in ventral face-selective ROIs have a foveal bias while pRFs in lateral face-selective

regions extend to the periphery

Next, we compared pRF characteristics across ventral and lateral face-selective ROIs. Visualizing

pRF centers across participants for each ROI (**Figure 2A**) reveals differences across ventral and

lateral regions. pRF centers of ventral face-selective regions, IOG, pFus and mFus, are largely

confined within the central 10° (**Figures 2, S2**). In contrast, pRF centers of lateral face-selective

regions, pSTS and mSTS, extend to the periphery, even past 30° from fixation. Notably, the

distribution of pRF centers of pSTS-faces is more similar to CoS-places, a ventral place-selective

region with a peripheral bias[33], than to ventral face-selective regions.

We quantified these observations by calculating the proportion of pRF centers within

each of four eccentricity bands (0–5°/5–10°/10–20°/20–40°) for each participant and ROI (**Figure

2B**). These eccentricities bands were chosen as they approximately occupy a similar cortical

expanse due to cortical magnification[25,49]. As illustrated in **Figure 2B**, the foveal bias is particularly

striking in the right hemisphere, where ~70% of pRF centers in right ventral face-selective regions

are located within the central 5° (right IOG-faces: mean±SE=0.71±0.07; pFus-faces: 0.69±0.06;

mFus-faces: 0.76±0.06), with very few pRF centers located outside the central 10°.

Lateral face-selective ROIs show the opposite distribution of pRF centers across

eccentricity bands. In these regions, the proportion of centers increases from central to

peripheral eccentricity bands. In stark contrast to ventral face-selective ROIs, less than 20% of

centers were located within the central 5° eccentricity band for lateral face-selective regions

(right pSTS-faces: 0.15±0.05; right mSTS-faces: 0.16±0.11). Here, the majority of pRF centers are

outside the central 10°.

Figure 2. pRF centers in ventral face-selective ROIs have a foveal bias, while pRF centers in lateral face-selective regions (and CoS-places) extend to the periphery. (A) Distribution of pRF centers; Each dot is a pRF center. *Dark colors:* Right hemisphere. Left mSTS-faces is not shown as it could only be localized in three participants. (B) *Bars:* mean proportion of pRF centers of face-selective ROIs and CoS-places across eccentricity bands (0°–5°, 5°–10°, 10°–20°, and 20°–40°) averaged across participants. *Error bars*: ±SE. *Dots:* individual participants.

A 2-way repeated measures LMM ANOVA on the proportion of centers with eccentricity

band (0–5°/5–10°/10–20°/20–40°) and stream (ventral: IOG/pFus/mFus and lateral: pSTS/mSTS;

mSTS, right hemisphere only) as factors revealed a significant eccentricity band x stream

interaction in both hemispheres (right: $F_{(3, 304)}=80.8$, $p<2.2x10^{-16}$; left: $F_{(3,256)}=30.1$, $p=2.2x10^{-}$

[16]). Post-hoc Tukey's tests establish that this is driven by a significantly higher proportion of

centers in the most foveal 0–5° eccentricity band in ventral vs. lateral face-selective regions

(proportion higher in ventral than lateral 0–5°: right: 0.56±0.05 , $t(304)=12.0$, $p<.0001$; left:

0.43±0.07, $t(256)=5.9$, $p<.0001$), as well as a significantly lower proportion of centers for ventral

vs. lateral regions in the two most peripheral eccentricity bands (proportion lower in ventral than

lateral, 10–20° right: 0.27±0.05, $t(304)=-5.7$, $p<.0001$; left: 0.29±0.07, $t(256)=-3.9$, $p=.0001$; 20–

40° right: 0.38±0.05, $t(304)=-8.0$, $p<.0001$; left: 0.40±0.07, $t(256)=-5.4$, $p<.0001$). Additionally,

differences in pRF center distributions are not a general characteristic of ventral vs. lateral

regions, as CoS-places, which is a ventral region, has pRFs that extend into the far periphery

(**Figure 2**-green).

There were also significant differences in the distribution of pRF centers between

hemispheres in ventral face-selective ROIs, revealed by a significant hemisphere by band by ROI

interaction ($F(6, 400)=5.0$, $p<6.2x10^{-5}$, 3-way repeated measures LMM ANOVA with factors of

hemisphere, eccentricity band, and ROI) and a significant hemisphere by band interaction ($F(3,$

$400)=13.7$, $p<1.6x10^{-8}$). As evident in **Figure 2B,** pRF centers were concentrated more foveally in

the right hemisphere than in the left, particularly in pFus-faces and mFus-faces. For both pFus-

faces and mFus-faces, there were significantly more pRF centers in the 0°–5° band in the right

than left hemisphere (proportion of pRFs in the 0°–5°: pFus-faces - right: mean±SE=0.69±0.06;

left: 0.34±0.07;   post-hoc   Tukey   test $t(400)=-5.2$, $p<.0001$;   mFus-faces   -   right:

mean±SE=0.76±0.06; left: 0.51±0.08; post-hoc Tukey test $t(400)=-3.7$, $p=.0002$).

Together this analysis reveals (i) differences in distribution of pRF centers across face-

selective regions of the lateral and ventral streams, and (ii) a higher foveal bias in right than left

ventral face-selective regions.

pRFs in lateral face-selective ROIs are larger than pRFs in ventral face-selective regions

In addition to differences in pRF locations, pRF sizes differ between lateral and ventral face-

selective regions. Specifically, median pRF sizes are larger in lateral than ventral face-selective

regions (**Figure 3A**). In the right hemisphere, for example, the average across participants of

median pRF sizes in pSTS faces was 21.7°±1.3°, while in pFus-faces (the ventral region with the

largest pRFs) the average was 9.4°±1.2°. To determine if these differences were significant, we

calculated in each participant the median pRF size across ROIs in each stream (ventral:

IOG/pFus/mFus and lateral: pSTS/mSTS; mSTS, right hemisphere only) and then compared values

across streams. Results show that in both hemispheres pRFs were significantly larger in lateral

than ventral face-selective regions (paired t-tests; right: $t(16)=-5.5$, $p=4.8x10^{-5}$; left: $t(13)=-5.8$,

$p=5.8x10^{-5}$). Differences between ROIs were significant (1-way repeated measures LMM ANOVAs

on median pRF size, right ROIs: IOG/pFus/mFus/pSTS/mSTS, $F(4,60)=25.1$, $p=3.6x10^{-12}$; left ROIs:

IOG/pFus/mFus/pSTS, $F(3,46)=27.9$, $p=2.1x10^{-10}$), and were driven by significant differences

between pSTS-faces and each of the ventral face-selective regions (post-hoc Tukey tests, all

ts>5.9, ps<.0001, **Table S1**).

Figure 3. pRFs in lateral face-selective regions are larger than in ventral face-selective regions. (A) Median pRF size $\left(\frac{\sigma}{\sqrt{n}}\right)$ in face-selective regions and CoS-places. *Box:* median and 25%, 75% percentiles; *lines:* ±1.5 times interquartile range; *Dots:* individual participant median pRF size. (B) Average median pRF size across participants for each eccentricity band of right hemisphere regions. Left hemisphere ROIs were not included as we could only identify in 5 participants any voxels whose pRF eccentricity was less than 5º in any of the lateral ROIs. *Dots:* Individual participant median pRF size. *Error bars:* ±SE.

As pRF size increases with eccentricity in retinotopic regions[25], we next tested if

differences in pRF sizes across pSTS and ventral regions were simply driven by the higher number

of peripheral pRFs in the former than the latter. We reasoned that if that were the case, then

comparison of pRF sizes within the same eccentricity band should reveal no difference across

ROIs. However, comparison of pRF sizes within the same eccentricity band revealed significant

differences between median pRF across ROIs (**Figure 3B,** 1-way LMM ANOVAs with factor of ROI,

right hemisphere, 0–5° band: $F_{(4,62)}=6.8$, p=.00013; 5–10° band: $F_{(4,53)}=8.7$, $p=1.9x10^{-5}$).

Moreover, across both the 0–5° and 5–10° bands, pRFs were significantly larger in right pSTS-

faces than in any other region except mSTS-faces (all ts>4.1, ps<=.0012, **Table S2**). We note that

pRF size in pSTS faces did not increase much with eccentricity, consistent with prior studies[50].

Ventral face-selective ROIs have dense coverage of the central visual field, while lateral face-

selective ROIs have coverage that extends to the periphery

How do differences in pRF locations and sizes across face-selective ROIs affect their visual field

coverage (VFC)? VFC was calculated as the proportion of pRFs covering each point in the visual

field, for each participant and ROI, then averaged across subjects. Results reveal three main

findings illustrated in **Figure 4A**: (1) In all ROIs, VFC is concentrated in the contralateral visual field

(see also **Figure S3**), (2) ventral face-selective ROIs have dense coverage concentrated around

the center of the visual field, and (3) lateral face-selective ROIs have more diffuse coverage that

extends further into the periphery than in ventral face-selective ROIs. These differences are not

a general characteristic of ventral vs. lateral regions, as ventral CoS-places exhibits VFC that

extends into the periphery.

To quantify differences in VFC, we calculated average pRF density as a function of

eccentricity for the contralateral visual field of each ROI. Ventral face-selective regions displayed

high pRF density close to the fovea that decreased sharply beyond ~10°, while pRF density in

lateral face-selective regions decreased more moderately with increasing eccentricity (**Figure**

4B). We summarized the relationship between pRF density and eccentricity by fitting a line for

each participant's pRF density curve per ROI. Negative slopes indicate higher pRF density near

the fovea than periphery, and slopes close to 0 indicate similar pRF densities across eccentricities.

Results reveal (i) significant differences between the average slopes of ventral and lateral face-

selective regions (paired t-tests; right: $t(16)=-9.4$, $p=6.9x10^{-8}$; left: $t(13)=-5.2$, p=.00016), whereby

slopes for ventral face-selective ROIs were more negative than for lateral face-selective ROIs and

(ii) significant differences between the average slopes of individual face-selective ROIs (right:

$F(4,58)=56.8$, $p<2.2x10^{-16}$; left: $F(362)=16.3$, $p=6.1x10^{-8}$, 1-way repeated measures LMM ANOVAs

on the slopes with factor ROI). Specifically, slopes in lateral face-selective regions—pSTS-faces

and mSTS-faces—were significantly closer to zero than any of the ventral face-selective regions

(all $ts>3.7$, $ps<=.0022$, post-hoc Tukey tests, **Table S3**). Additionally, bilateral pFus-faces and right

mFus-faces had significantly more negative slopes than IOG-faces (all $ts<-3.1$, $ps<=.017$, **Table**

S3), indicating that the former ROIs have a larger foveal bias than the latter.

Figure 4. Dense coverage of the central visual field in ventral face-selective regions, but coverage extends to the
periphery in lateral face-selective regions (and CoS-places). (A) Average visual field coverage of each ROI across
participants (number indicated above each plot). Visual field coverage is computed as the proportion of pRFs
covering each point in the visual field for each participant and then averaged across participants. *Asterisk:* average
location of the center of mass of all pRF centers in each ROI; (B) Average pRF density as a function of eccentricity for
the contralateral visual field of each ROI. Data were calculated per participant and then averaged across participants.
*Shaded area: s*tandard error of the mean.

Face-selective regions have direct white matter connections to early visual cortex

Given our findings of differences in pRF properties and visual field coverage across ventral vs.

lateral face-selective regions, we asked how white matter connections from early visual cortex

(EVC) contribute to this differentiation. We reasoned that the foveal bias in ventral face-selective

regions may arise in part from a preponderance of connections from foveal eccentricities of EVC.

In contrast, the more uniform tiling of the visual field by pRFs in lateral face-selective regions may

rise from a more uniform pattern of white matter connections from EVC eccentricities to these

face-selective ROIs along the STS.

Figure 5. Functionally-defined white matter tracts (fWMT) between functional ROIs and early visual cortex (DVC).
We combined dMRI and fMRI in each participant to determine fWMT between each functional ROI and EVC. *From*
left to right: fMRI: from the localizer experiment, we defined in each participant their face-selective ROIs; from the
toonotopy experiment, we defined in each subject their EVC and eccentricity map. *dMRI:* using MRTrix3 with ACT
we defined the whole brain connectome of each participant, which was culled to remove false alarm tracts with
Linear Fascicle Evaluation{ref}. *fWMT of each ROI:* all tracts that intersect with both the functional ROI and EVC.
Endpoints in EVC: We projected the end points of each fWMT in EVC and related it to the eccentricity map. All
analyses were done in each participant's native brain space.

To test these predictions, we acquired an additional diffusion-weighted MRI (dMRI) scan

from 16 of our participants (see Methods). We combined dMRI data with the fMRI data from

both localizer and toonotopy experiments to identify the functionally-defined white matter tracts

(fWMT) that connect each face-selective region with EVC (**Figure 5**). We used anatomically-

constrained tractography (ACT)[51], which uses the gray-white matter-interface to seed the

tractography. This guarantees that the resulting fiber tracts reach the gray matter, which is crucial

for testing our hypothesis. As a control, we also defined the fWMT that connect CoS-places with

EVC. We asked: (1) Are there white matter tracts that connect each of the face-selective ROIs and

EVC? (2) If so, how are they distributed across EVC eccentricities?

We found direct white matter connections between each face- and place-selective region

and EVC in all participants where we could localize the region (**Figure 6A**, representative subjects;

all subjects, **Supplementary Figure S4**). Qualitatively, fWMT between face-selective regions and

EVC were largely consistent across participants, although the proportion of tracts varied

substantially by region (**Figure 7B**). IOG-faces had the highest percentage of its tracts connecting

to EVC (right: mean±SE=37.7±2.9%, left: 39.6±3.1%), followed by pFus-faces (right: 13.2±1.7%,

left: 14.7±2.8%). mSTS-faces had the lowest percentage, with less than 3% of all mSTS-faces tracts

connecting to EVC (right: 2.1±0.6%; left: 2.6±0.9%). However, in both hemispheres, ventral mFus-

faces and lateral pSTS-faces had similar percentages of tracts to EVC (right mFus-faces: 9.5±1.6%,

left mFus-faces: 8.4±0.9%; right pSTS-faces: 10.8±1.7%, left pSTS-faces: 9.0±1.6%), suggesting

that differences in fWMT to EVC do not stem from differences in our ability to identify tracts

across ventral and lateral streams. Additionally, despite their adjacent locations, ventral regions

mFus-faces and CoS-places have different percentages of connections to EVC. Specifically, CoS-

places has twice the percentage of connections to EVC as mFus-faces (right: 22.0±2.9%; left:

17.4±3.1%). These data further underscore that factors other than anatomical location

determine the percentage of tracts between functional ROIs and EVC.

Figure 6. White matter tracts connecting EVC and face-selective regions (and CoS-places). (A) *Top:* right
hemisphere, Subject 1; *Bottom:* left hemisphere, Subject 2. (B) Mean percentage of fWMT between each functional
ROI and EVC. 100% indicates that all tracts starting at an ROI connect to EVC and 0 indicates that no tracts connect
to EVC. Percentage is calculated for each participant (circles) and then averaged across participants (bars). *Lighter*
bars: left hemisphere; *Darker bars:* right hemisphere.

Ventral and lateral face-selective regions differentially connect to EVC eccentricities

To test whether ventral and lateral face-selective regions are differentially connected to EVC eccentricities, we evaluated to which eccentricity band in EVC these tracts connect. Eccentricity values were determined from each participant's toonotopy data. We then quantified what proportion of tracts from an ROI to EVC connected to each of four EVC eccentricity bands: 0–5°, 5–10°, 10–20°, 20–40° (**Figure 7**).

Figure 7. Tracts from lateral and ventral face-selective regions have a different distribution across early visual cortex (EVC) eccentricities. (A) Mean endpoint density of fiber tracts connecting each ROI with EVC shown on the zoomed medial inflated cortical surface of the FreeSurfer average brain; *Top:* right hemisphere; *Bottom:* left hemisphere. *Color map:* average density across participants. Eccentricity bands (white) were derived from the average of all participants' retinotopies. (B) Average proportion of tract endpoints from each ROI that terminate in each of four EVC eccentricity bands (0°–5°; 5°–10°; 10°–20º°; 20°–40°). *Bars:* mean across participants; Each dot is a participant; *Error bars*: ±SE. Analyses done in native subject brain space

For ventral face-selective regions in both hemispheres, the highest proportion of white

matter endpoints were located within the central 5° of EVC, while the fewest endpoints were

within the most peripheral EVC eccentricity band, mirroring the results for pRF centers (**Figure**

7A, IOG, pFus, mFus). In fact, the majority (>66%) of the tracts connecting EVC to ventral face-

selective ROIs originate within the central 10°. This overrepresentation of tract endpoints within

the central EVC is not the case for lateral face-selective regions. In these regions, tract endpoints

are instead more evenly distributed across EVC eccentricity bands, with at least 20% of endpoints

in the most peripheral eccentricity band (>40°, **Figure 7A**-pSTS, mSTS). Importantly, this

difference is not due to a general bias that favors connections from foveal eccentricities in EVC

to VTC, as the tracts that connect to ventral CoS-places to EVC are more uniformly distributed

across EVC eccentricities. In fact, nearly half of the tracts from EVC to CoS-places (right: 48%, left:

43%) originate in eccentricities ≥20° .

To test the significance of endpoint distributions in EVC between streams, we ran a 2-way

repeated-measures LMM ANOVA on the proportion of tract endpoints in EVC with factors of

eccentricity band (0–5°/5–10°/10–20°/20–40°) and stream (ventral: IOG/pFus/mFus and lateral:

pSTS/mSTS; mSTS right hemisphere only). We found a significant eccentricity band x stream

interaction in both hemispheres (right: $F(3, 268)=10.3$, $p=2.0x10^{-6}$; left: $F(3,200)=19.5$, $p=3.8x10^{-11}$).

As with the proportion of pRF centers, post-hoc Tukey's tests establish that stream

differences are driven by a significantly higher proportion of tract endpoints in the most central

eccentricity band (0–5°) in ventral vs. lateral face-selective regions (proportion higher in ventral

than lateral, right: 0.17±0.04, $t(268)=4.4$, $p<.0001$; left: 0.20 ± 0.04, $t(200)=5.0$ $p<.0001$), as well

as a significantly lower proportion of centers for ventral vs. lateral regions in the two most

peripheral eccentricity bands (proportion lower in ventral than lateral, 10–20°, right: 0.09±0.04,

$t(268)=-2.2$, $p=.0253$; left: 0.13±0.04, $t(200)=-3.4$, $p=.0009$; 20–40°, right: 0.09±0.04, $t(268)=-2.5$,

$p=.0148$; left: 0.16±0.04, $t(200)=-4.0$, $p=.0001$). To control for any effects of ROI size, we repeated

this analysis with 5 mm disk ROIs, created at the center of each functionally defined region

(**Supplementary Figure S5**). As in the main analysis, a 2-way repeated-measures ANOVA on the

proportion of fiber endpoints in EVC confirms significant eccentricity band x stream interactions

in both hemispheres (right: $F(3, 256) = 8.7$, $p = 1.65 \times 10^{-5}$; left: $F(3,200) = 17.3$, $p = 5.248 \times 10^{-10}$).

Together, we find that the ventral and lateral face-selective regions differentially connect

to EVC. Ventral face-selective regions connect more to central eccentricity bands in EVC, while

lateral face-selective regions show connections that are distributed across eccentricity bands and

extend into the far periphery. White matter connectivity differences between ventral and lateral

face-selective ROIs echo the different distributions of pRF centers, suggesting that white matter

connections may provide a scaffolding for differential spatial processing between streams.

Discussion

Using a multimodal approach, we tested how different face-selective regions of visual cortex

represent visual space and whether this can be traced to white matter connections with early

visual cortex. We find that (i) contrary to the prevailing view, not all face-selective regions are

foveally biased, as lateral face-selective regions show visual field coverage extending into the far

periphery, and that (ii) these differences are reflected in differential patterns of white matter

connections with EVC, as lateral face-selective regions exhibit white matter connections that are

more evenly distributed across eccentricities than ventral face-selective regions.

Consistent with prior research illustrating a foveal bias in face-selective regions[32,33,52], our

data show that pRF properties and VFC of ventral occipital and temporal face-selective regions

process information around the center of gaze. Surprisingly, we find that this is not a general

property of face-selective regions: pRFs of face-selective regions of the lateral stream are both

more peripheral and larger than those of the ventral stream. This finding requires a rethinking of

spatial computations in face perception.

A substantial body of research[1,3,4] has highlighted differences in the computational goals

of ventral and lateral streams: ventral regions are thought to be involved in static aspects of face

perception such as face identification, while lateral regions are thought to be involved in

dynamic[3,13,14] and transient[15] aspects of face perception such as expression[7,8] and gaze[18,19]. As

such, the foveal bias of ventral face-selective regions has been attributed to the high visual acuity

necessary for face identification, which is achieved by foveal vision[31]. Indeed, people tend to

fixate on the center of the face during recognition[30,31,34], putting the VFC of ventral face selective

regions on the internal face features that carry information about identity[34,53]. This raises the

question: what is the benefit of peripheral processing in the lateral stream? We propose a new

hypothesis: peripheral processing in lateral face-selective regions and direct white matter

connections between these regions to peripheral EVC may facilitate rapid processing of face

information. In-line with this prediction, behavioral studies show that (i) tasks associated with

the lateral stream such as face emotion judgements[27,53] or person recognition from dynamic face

and body stimuli[54] use peripheral information, and (ii) the speed of visual processing is faster in

the periphery[36]. Additionally, not only do lateral face-selective regions contain large and

peripheral pRFs, but they are also mainly driven by transient visual information[15]. Future studies

can test these hypotheses by measuring spatio-temporal pRFs, as well as behavioral performance

in the periphery for tasks related to the lateral stream.

Though dMRI is not capable of resolving whether the connections we find are

feedforward or feedback, the majority of connections in cortex are likely reciprocal[55]. Thus, we

conjecture that included in the white-matter connections we discovered are tracts that originate

in EVC and project to face-selective regions. Additionally, we do not intend to suggest that all

differences in spatial processing between ventral and lateral face-selective regions can be

attributed to direct connections with EVC. It is likely that hierarchical connections throughout the

ventral stream also contribute to shaping pRFs in face-selective regions, though we are not able

to resolve topographic information for these shorter connections with the present dMRI

resolution. Nonetheless, together with hierarchical connections, small differences in the

distribution of connections from EVC eccentricities to face-selective regions may result in

pronounced differences in spatial processing by pRFs across processing streams.

Our finding that visual field properties in face-selective regions are reflected in differential

white matter connections with EVC supports the idea that structural connections serve a

functional role. Our data suggest that it is not just that face-selective regions in different streams

use the input that they are receiving from early in the visual hierarchy differently, but that they

are actually receiving different input to begin with. Further, we hypothesize that white matter

connections between EVC and face-selective regions may explain higher functional connectivity

between ventral face-selective regions and central than peripheral eccentricities in EVC[56–58], a

prediction that can be tested in future research. As these functional connections are present

early in infancy[56], another important question for future research is whether these structural

connections between EVC and downstream regions are present at birth and constrain where

face-selective regions emerge during development[59].

Our results not only explicate the nature of connections between EVC and face-selective

regions[41,42], but also substantially advance the precision with which we can understand

connections within the human brain. In contrast to many dMRI studies that examine white matter

connections of entire fascicles or between brain regions, to our knowledge this is the first study

that identifies white matter connections associated with a topographic map in individual human

brains. As receptive fields, topographic maps, and processing streams are key computational

characteristics of sensory (e.g., audition[60]) and cognitive (e.g., numerosity[61]) brain systems and

across species[52], our innovative approach can be applied to study how topographic computations

emerge and are scaffolded by white matter connections in many brain systems and species.

In sum, we discovered differential spatial processing across face-selective regions in the ventral

and lateral processing streams, which is mirrored by differences in white matter connections with

eccentricity bands in EVC. These findings suggest a rethinking of computations in face-selective

regions, and further propose that spatial processing in high-level regions may be scaffolded by

the pattern of white matter connections from EVC.

Methods

Participants

Twenty-one participants (10 female, ages 22 to 45, mean age 27.1 ± 5.8 years) from Stanford University

participated in the study, sixteen of whom returned for diffusion MRI. All participants had normal or

corrected-to-normal vision. Participants gave informed consent and all procedures were approved by the

Stanford Internal Review Board on Human Subjects Research.

MRI data acquisition

Participants were scanned using a General Electric Discovery MR750 3T scanner located in the Center for

Cognitive and Neurobiological Imaging (CNI) at Stanford University. A phase-array 32-channel head coil

was used for all data acquisition except for the retinotopic mapping experiment (16-channel head coil).

Anatomical Scans

We obtained for each of 15 participants a whole-brain, anatomical volume using a T1-weighted BRAVO

pulse sequence (resolution: 1 mm x 1 mm x 1 mm, TI=450 ms, flip angle: 12°, 1 NEX, FoV: 240 mm). In 6

participants we used quantitative MRI measurements using the protocols from[62] to generate a 1 mm x 1

mm x 1 mm, T1-weighted anatomical image of their brain. Anatomical images of each participant's brain

were used for gray/white matter tissue segmentation, which was then used for cortical surface

reconstruction and visualization of data on the cortical surface.

fMRI category localizer experiment: A category localizer was run to identify voxels that responded more

strongly to one category vs. other categories of stimuli in order to localize face- and place-selective regions

of interest (ROIs). In the experiment, participants were presented with grayscale images of stimuli from

five domains, each with two categories (faces: child, adult; bodies: whole, limbs; places: corridors, houses;

characters: pseudowords, numbers; objects: cars, guitars) as described previously[45]. Images from each

category were presented in 4s trials at a rate of 2 Hz, intermixed with 4s blank trials. Each category was

shown eight times per run in counterbalanced order. Participants were instructed to fixate on a central

point and perform an oddball detection task, identifying 0-2 randomly presented phase scrambled images

within a trial. Runs were 318s long and each participant completed 3 runs with different images and

category order. Data were collected with a simultaneous multi-slice EPI sequence with a multiplexing

factor of 3 to acquire near whole-brain (48 slices) volumes at TR=1s, TE=30 ms at a resolution of 2.4 mm

isotropic. Data were acquired with a one-shot T2*-sensitive gradient echo sequence and slices were

aligned parallel to the parieto-occipital sulcus.

pRF mapping experiment: Each participant completed four runs of a wide-field pRF mapping experiment

to model the region of the visual field that elicits a response from each voxel (i.e. each voxel's pRF). In

each run, participants fixated on a central stimulus and were required to press a button whenever the

central fixation dot changed color as bars were swept across the screen (mean accuracy=$95.2 \pm 6.9\%$).

Due to our interest in higher-level category selective regions, bars of width $5.7°$ swept across high-

contrast, colorful, cartoon stimuli that spanned a wide-field circular aperture ($40°$ x $40°$ of visual angle).

Images randomly changed at a rate of 8 Hz. In each run, bars of four orientations ($0°$, $45°$, $90°$, $135°$) swept

the visual field in eight directions (2 opposite directions orthogonal to the bar orientation) with blanks

(mean luminance gray background with a fixation) interspersed at regular intervals. Each run lasted 3 min

and 24 s. Eye-tracking was not conducted in the scanner but visual field coverage of early visual field maps

(V1-V3) show the expected hemifields and quarterfields (**Figure 1B** for an example), indicating that

participants maintained stable fixation.

Diffusion magnetic resonance imaging (dMRI): In 16 of the participants, we acquired whole-brain

diffusion-weighted, dual-spin echo sequences (60 slices, TE=96.8 ms, TR=8,000 ms, 96 diffusion directions,

b0=2,000s/mm^2) at a resolution of 2.0 mm x 2.0 mm x 2.0 mm. Ten non-diffusion-weighted images were

collected at the beginning of each scan.

Preprocessing and data analysis

Anatomical data: T1-weighted images were automatically segmented using FreeSurfer 5.3

(https://surfer.nmr.mgh.harvard.edu/) and then manually validated and corrected using ITKGray

(http://web.stanford.edu/group/vista/cgi-bin/wiki/index.php/ItkGray). Using these segmentations, we

generated cortical surfaces in FreeSurfer.

fMRI data analysis: fMRI data analysis was performed in MATLAB (www.mathworks.com) and using the

mrVista analysis software (http://github.com/vistalab) developed at Stanford University. All data were

analyzed within the individual participants' native brain anatomy space without spatial smoothing. Data

were motion corrected within and between scans using mrVista motion correction algorithms, and then

manually aligned to the anatomical volume. The manual alignment was optimized using robust
multiresolution alignment.

Definition of V1-V3 (EVC): V1, V2, and V3 were defined in all participants using their individual data from
the pRF mapping experiment. Maps of pRF phase and eccentricity were projected onto an inflated cortical
surface reconstruction for each participant and borders between the retinotopic maps were drawn on
this cortical surface. The boundary was defined as the center of polar angle reversal occurring at the
vertical or horizontal meridian[34]. V2 and V3 were drawn as quarterfields separated by V1 and were later
combined to produce a hemifield representation. The early visual cortex (EVC) ROI was then defined as
the union of V1, V2, and V3.

Definition of face-selective and place-selective functional regions of interest: The functional ROIs were
defined in individual participants from the category localizer experiment using anatomical and functional
criteria[45]. Statistical contrasts of faces > all other stimuli (for face-selective ROIs) and places > all other
stimuli (for place-selective ROIs) were thresholded at t-values >3, voxel level, in each participant, as in
previous work[34,45]. Face-selective ROIs were defined in the inferior occipital gyrus (IOG-faces, right
hemisphere: N=20; left hemisphere: N=18), posterior fusiform gyrus (pFus-faces, right: N=18; left: N=18),
mid fusiform gyrus (mFus-faces; right: N=17; left: N=19), posterior superior temporal sulcus (pSTS-faces,
right: N=19; left: N=17), and medial superior temporal sulcus (mSTS-faces, right: N=18; left: N=9). An
additional place-selective ROI was defined along the collateral sulcus (CoS-places, right: N=21; left: N=21),
corresponding to the parahippocampal place area[63].

Estimating pRFs: We modeled the population receptive field (pRF) in each voxel using the compressive
spatial summation (CSS) model[64]. Time course data was transformed from the functional slices to the T1-
weighted anatomy using trilinear interpolation. We estimated pRFs within each gray matter voxel. The
pRF model fits a 2-dimensional Gaussian for each voxel, with parameters x and y describing the location
of the center of the pRF, and σ, the standard deviation of the Gaussian in degrees of visual angle. An
additional compressive summation exponent (n) is fit for each voxel to capture the nonlinear summation
properties of pRFs in later stages of the visual hierarchy[26,64]. We define the size of the pRF as $\sigma/\sqrt{n}$[64].
Candidate pRFs are estimated and a predicted time-series is produced by convolving the product of the
binarized stimulus sequence and the candidate pRF with an HRF. The model parameters x, y, and σ, are
then iteratively adjusted to minimize the sum of squared errors between the predicted time-series and

the observed time-series. Phase (atan(y/x)) and eccentricity (sqrt($x^2 + y^2$) of each voxel are derived from

the x and y coordinates of the pRF center to generate phase and eccentricity maps respectively. Voxels

were included for subsequent analysis if the variance explained by the pRF model was greater than 10%.

Additionally, to ensure accurate pRF fits by the optimization procedure, voxels whose σ remained at the

initial minimum value (0.21°) were excluded from further analysis. For analyses of pRFs in each ROI, we

include data from participants for whom the ROI was identified and at least 25 voxels (corresponding to

a volume of $\geq$ 25 mm^3) of the ROI were retinotopically modulated according to the above criteria

(variance explained $\geq$10%, pRF size > 0.21°). The number of participants per ROI is as follows: IOG-faces,

right hemisphere N=19, left N=17; pFus-faces, right N=18, left N=18; mFus-faces, right N=17, left N=17;

pSTS-faces, right N=16, left N=13; mSTS-faces, right N=6, left N=2; CoS-places, right N=21, left N=21. Note

that left mSTS-faces in not included in further analyses as we were only left with two participants.

Visual field coverage (VFC): is defined as the portion of the visual field that is processed by the set of pRFs

spanning the ROI. To calculate the VFC for a given ROI and participant, all voxels in an ROI that passed the

above inclusion criteria are included. Each pRF is modeled with a binary circular mask centered at the pRF

center and with a diameter of 2σ/$\sqrt{n}$. For each location in the visual field, the VFC value is the pRF density

at that location. Group average VFC maps are created by averaging the individual VFC maps for that ROI.

Thus, the group VFC depicts the mean pRF density at each location averaged across subjects. To assess

differences in VFC across ROIs, we fitted a line relating pRF density to eccentricity in each ROI and

participant, and then compared the average slopes (across participants) of different ROIs in each stream.

Linear fits were chosen over more complicated models as the linear model fit (R^2) was high for all ROIs

(right: mean R^2s > 0.71, left: mean R^2s > 0.66)

For each functional ROI, we also calculated indices of contra- vs. ipsi-laterality and upper vs. lower

visual field bias of the VFC (**Supplementary Figure S3**). Laterality of VFC was calculated for each participant

and ROI as the mean coverage in the contralateral visual field minus the mean coverage in the ipsilateral

visual field, divided by the sum of the two ($\frac{contra - ipsi}{contra + ipsi}$). Similarly, the upper field bias for each participant

and ROI was quantified as the average coverage in the contralateral upper visual field minus the average

coverage in the contralateral lower visual field, divided by the sum of the two ($\frac{upper - lower}{upper + lower}$).

dMRI data analysis: dMRI data were preprocessed using a combination of tools from MRtrix3[65], ANTs, and

FSL, as in[66,67]. We denoised the data using: (i) a principal component analysis, (ii) Rician based denoising,

and (iii) Gibbs ringing corrections[68]. We also corrected for eddy currents and motion using FSL and
performed bias correction using ANTs. The dMRI data were then registered to the average of the non-
diffusion-weighted images and aligned to the corresponding anatomical brain volume of the participant
using rigid body transformation. We used constrained spherical deconvolution[65] with up to eight spherical
harmonics (lmax=8) to calculate the voxel-wise tract orientation distributions (FOD). These FODs were
then used for tractography.

Tractography: Ensemble tractography[69] with linear fascicle evaluation (LiFE[70]) was performed using the
preprocessed dMRI data and consisted of 3 main steps:

1) We used MRtrix3[65] to generate five candidate connectomes which varied in the maximum angle
(2.9°, 5.7°, 11.5°, 23.1°, and 47.2°). Each candidate connectome was seeded on the gray white
matter interface (GWMI) using anatomically constrained tractography (ACT[51]) and consisted of
500,000 streamlines. This approach allows us to generate multiple connectomes with different
degrees of curvature, as opposed to restricting our estimations to a single connectome with one
particular set of parameters[69]. Each candidate connectome was generated using probabilistic
tract tracking with the following parameters: algorithm: IFOD2, step size: 1 mm, minimum length:
4 mm, maximum length: 200 mm. ACT[51] uses the GWMI from the FreeSurfer segmentation of
each participant's brain to seed the tracking algorithm. This enabled us to focus only on those
fiber tracts that reach the gray matter.
2) The five candidate connectomes were concatenated into one ensemble connectome containing
a total of 2,500,000 streamlines.
3) We used linear fascicle evaluation (LiFE[70]) to validate the ensemble connectome. LiFE was used to
determine which fiber estimates make a significant contribution to predicting the dMRI data and
removes false alarm tracts that did not significantly contribute to predicting the dMRI data.

Functionally defined white matter tracts (FDWT): To determine the white matter tracts that are associated
with each functional ROI, we intersected the tracts with the GWMI directly adjacent to each functional
ROI for each participant. This yielded functionally defined white matter tracts (FDWT) for each face and
place-selective ROI.
EVC-fROI connections: To identify tracts connecting these functional ROIs and early visual cortex, we
intersected the FDWM of the fROI with the GWMI underneath EVC (union of V1, V2, V3), and then used
these pairwise tracts for individual subject visualization and endpoint density estimation.

Quantification of endpoint density

We used MRtrix3[65] and track-density imaging (TDI) to create tract maps of the streamline endpoints of

the FDWT for each functional ROI. These endpoint density maps were transformed to the cortical surface

using FreeSurfer. We examined the portion of the endpoint density maps that intersected EVC and

determined the corresponding eccentricity values. Eccentricity was derived from the pRF mapping

experiment and each participant's individual retinotopic maps. We used the track-density values to weight

each eccentricity value based on the density of endpoints connecting to that eccentricity. We then

quantified the proportion of endpoints in each of four eccentricity bands (0° to 5°, 5° to 10°, 10° to 20°,

20° to 40°) for each fROI and participant separately.

Size control: As functional ROIs vary in size across participants and regions, we did a control analysis to

evaluate the endpoint density using constant-size spherical ROIs (radius=5mm) centered on each

participant's functional ROIs. The goal of this analysis was to test whether differences across ROIs could

be attributed to differences in ROI size. Results of these analyses are presented in the Supplementary

Materials (**Supplementary Figure S5**).

Statistical analyses

Statistical analyses were computed using the *lmerTest*[48] and *lsmeans* packages in R (https://cran.r-

project.org/web/packages/lmerTest/, https://cran.r-project.org/web/packages/lsmeans/). Linear mixed-

effects models (LMMs) were used for statistical analyses because our data includes missing data points

(not all participants have all ROIs). LMMs were constructed separately for each hemisphere, with one

exception: to test the effect of hemisphere on the proportion of pRF centers by eccentricity band in the

ventral face-selective ROIs (i.e. to test the lateralization of the foveal bias), we constructed an LMM which

can be expressed as: *proportion of pRF centers ~ eccentricity band + ROI + hemisphere + (eccentricity*

band*ROI) + (ROI*hemisphere) + (eccentricity band*hemisphere) + (eccentricity band*ROI*hemisphere) +

(1|subject). The significance of LMM model terms was evaluated using repeated-measures analyses-of-

variance (LMM ANOVAs; type III), with Satterthwaite's method of correction for degrees of freedom[47].

Post-hoc tests were done by computing Tukey's Honest Significant Difference (HSD) where the family-

wise confidence level is set at 0.95. p-values reported from these post-hoc tests are thus corrected for

multiple comparisons. All statistical tests conducted in the manuscript followed this procedure other than

two paired t-tests. One paired t-test was conducted to compare the average pRF size across streams

(average pRF size across ventral ROIs vs. average pRF size across lateral ROIs) and one was conducted to

compare slopes of the pRF density by eccentricity lines across streams (average slope across ventral ROIs

vs. average slope across lateral ROIs).

Data and code availability

fMRI data were analyzed using the open source mrVista software package (available on GitHub:

http://github.com/vistalab). T1-weighted anatomicals for six subjects were generated using the mrQ

software package (https://github.com/mezera/mrQ). dMRI data were analyzed using mrTrix3

(http://www.mrtrix.org/). Custom code (all available on GitHub) was used for dMRI preprocessing

(https://github.com/vistalab/RTP-preproc; https://github.com/vistalab/RTP-pipeline) and processing of

the pRF experiment (https://github.com/VPNL/toonotopy). Code for all further analysis, including

reproducing all figures and statistics, can be found at https://github.com/VPNL/fibeRFs. Processed data is

additionally available at https://github.com/VPNL/fibeRFs while the individual nifti files are available upon

request.

References

1. Haxby, J. V, Hoffman, E. a & Gobbini, M. I. The distributed human neural system for face perception [Record Supplied By Publisher]. *Trends Cogn Sci* (2000). doi:10.1016/S1364-6613(00)01482-0

2. Weiner, K. S. & Grill-Spector, K. Neural representations of faces and limbs neighbor in human high-level visual cortex: Evidence for a new organization principle. *Psychol. Res.* **77**, 74–97 (2013).

3. Duchaine, B. & Yovel, G. A Revised Neural Framework for Face Processing. *Annu. Rev. Vis. Sci.* (2015). doi:10.1146/annurev-vision-082114-035518

4. Freiwald, W., Duchaine, B. & Yovel, G. Face Processing Systems: From Neurons to Real-World Social Perception. *Annu. Rev. Neurosci.* (2016). doi:10.1146/annurev-neuro-070815-013934

5. Kanwisher, N., McDermott, J. & Chun, M. M. The fusiform face area: a module in human extrastriate cortex specialized for face perception. *J. Neurosci.* **17**, 4302–4311 (1997).

6. Grill-Spector, K., Knouf, N. & Kanwisher, N. The fusiform face area subserves face perception, not generic within-category identification. *Nat. Neurosci.* **7**, 555–562 (2004).

7. Andrews, T. J. & Ewbank, M. P. Distinct representations for facial identity and changeable aspects of faces in the human temporal lobe. *Neuroimage* (2004). doi:10.1016/j.neuroimage.2004.07.060

8. Winston, J. S., Henson, R. N. A. A., Fine-Goulden, M. R. & Dolan, R. J. fMRI-adaptation reveals dissociable neural representations of identity and expression in face perception. *J. Neurophysiol.* (2004). doi:10.1152/jn.00155.2004

9. Pitcher, D., Walsh, V., Yovel, G. & Duchaine, B. TMS Evidence for the Involvement of the Right Occipital Face Area in Early Face Processing. *Curr. Biol.* **17**, 1568–1573 (2007).

10. Gauthier, I., Skudlarski, P., Gore, J. C. & Anderson, A. W. Expertise for cars and birds recruits brain areas involved in face recognition. *Nat. Neurosci.* **3**, 191–197 (2000).

11. Weiner, K. S. & Grill-Spector, K. *The evolution of face processing networks*. *Trends in Cognitive Sciences* (2015). doi:10.1016/j.tics.2015.03.010

12. Pyles, J. A., Verstynen, T. D., Schneider, W. & Tarr, M. J. Explicating the Face Perception

Network with White Matter Connectivity. *PLoS One* **8**, e61611 (2013).

13.   Fox, C. J., Iaria, G. & Barton, J. J. S. Defining the face processing network: Optimization of

the functional localizer in fMRI. *Hum. Brain Mapp.* (2009). doi:10.1002/hbm.20630

14.   Pitcher, D., Dilks, D. D., Saxe, R. R., Triantafyllou, C. & Kanwisher, N. Differential

selectivity for dynamic versus static information in face-selective cortical regions.

Neuroimage **56**, 2356–2363 (2011).

15.   Stigliani, A., Jeska, B. & Grill-Spector, K. Differential sustained and transient temporal

processing across visual streams. *PLoS Comput. Biol.* (2019).

doi:10.1371/journal.pcbi.1007011

16.   Puce, A., Allison, T., Bentin, S., Gore, J. C. & McCarthy, G. Temporal cortex activation in

humans viewing eye and mouth movements. *J. Neurosci.* **18**, 2188–99 (1998).

17.   Pitcher, D., Duchaine, B. & Walsh, V. Combined TMS and fMRI reveal dissociable cortical

pathways for dynamic and static face perception. *Curr. Biol.* **24**, 2066–2070 (2014).

18.   Calder, A. J. *et al.* Separate Coding of Different Gaze Directions in the Superior Temporal

Sulcus and Inferior Parietal Lobule. *Curr. Biol.* (2007). doi:10.1016/j.cub.2006.10.052

19.   Carlin, J. D. & Calder, A. J. The neural basis of eye gaze processing. *Current Opinion in*

Neurobiology (2013). doi:10.1016/j.conb.2012.11.014

20.   Pitcher, D., Walsh, V. & Duchaine, B. The role of the occipital face area in the cortical face

perception network. *Exp. Brain Res.* (2011). doi:10.1007/s00221-011-2579-1

21.   Huk, A. C., Dougherty, R. F. & Heeger, D. J. Retinotopy and functional subdivision of

human areas MT and MST. *J. Neurosci.* (2002). doi:20026661

22.   Grossman, E. D. & Blake, R. Brain Areas Active during Visual Perception of Biological

Motion. *Neuron* **35**, 1167–1175 (2002).

23.   Beauchamp, M. S., Lee, K. E., Haxby, J. V. & Martin, A. Parallel visual motion processing

streams for manipulable objects and human movements. *Neuron* (2002).

doi:10.1016/S0896-6273(02)00642-6

24.   Hubel, D. H. & Wiesel, T. N. Receptive fields and functional architecture in two nonstriate

visual areas (18 And 19) of the cat. *J Neurophysiol* **28**, 229–289 (1965).

25.   Dumoulin, S. O. & Wandell, B. A. Population receptive field estimates in human visual

cortex. *Neuroimage* **39**, 647–660 (2008).

26.   Kay, K. N., Weiner, K. S. & Grill-Spector, K. Attention reduces spatial uncertainty in

human ventral temporal cortex. *Curr. Biol.* **25**, 595–600 (2015).

27.   Pelphrey, K. A. *et al.* Visual Scanning of Faces in Autism. *J. Autism Dev. Disord.* (2002).

doi:10.1023/A:1016374617369

28.   Hsiao, J. H. W. & Cottrell, G. Two fixations suffice in face recognition. *Psychol. Sci.* (2008).

doi:10.1111/j.1467-9280.2008.02191.x

29.   van Belle, G., Ramon, M., Lefèvre, P. & Rossion, B. Fixation patterns during recognition of

personally familiar and unfamiliar faces. *Front. Psychol.* (2010).

doi:10.3389/fpsyg.2010.00020

30.   Peterson, M. F. & Eckstein, M. P. Looking just below the eyes is optimal across face

recognition tasks. *Proc. Natl. Acad. Sci. U. S. A.* (2012). doi:10.1073/pnas.1214269109

31.   Malach, R., Levy, I. & Hasson, U. The topography of high-order human object areas.

Trends Cogn Sci **6**, 176–184 (2002).

32.   Hasson, U., Levy, I., Behrmann, M., Hendler, T. & Malach, R. Eccentricity bias as an

organizing principle for human high-order object areas. *Neuron* **34**, 479–490 (2002).

33.   Levy, I. *et al.* Center-periphery organization of human object areas. *Nat. Neurosci.* (2001).

doi:10.1038/87490

34.   Gomez, J., Natu, V., Jeska, B., Barnett, M. & Grill-Spector, K. Development differentially

sculpts receptive fields across early and high-level human visual cortex. *Nat. Commun.* **9**,

788 (2018).

35.   Behrmann, M. & Plaut, D. C. A vision of graded hemispheric specialization. *Ann. N. Y.*

Acad. Sci. **1359**, 30–46 (2015).

36.   Carrasco, M., McElreel, B., Denisova, K. & Giordano, A. M. Speed of visual processing

increases with eccentricity. *Nat. Neurosci.* (2003). doi:10.1038/nn1079

37.   Azzopardi, P. & Cowey, A. Preferential representation of the fovea in the primary visual

cortex. *Nature* (1993). doi:10.1038/361719a0

38.   Jiang, X. *et al.* Evaluation of a Shape-Based Model of Human Face Discrimination Using

fMRI and Behavioral Techniques. *Neuron* (2006). doi:10.1016/j.neuron.2006.03.012

39. Yamins, D. L. K. *et al.* Performance-optimized hierarchical models predict neural

responses in higher visual cortex. *Proc. Natl. Acad. Sci.* (2014).

doi:10.1073/pnas.1403112111

40. Grill-Spector, K., Weiner, K. S., Kay, K. N. & Gomez, J. The Functional Neuroanatomy of

Human Face Perception. *Annu. Rev. Vis. Sci.* **3**, 167–196 (2017).

41. Gschwind, M., Pourtois, G., Schwartz, S., Van De Ville, D. & Vuilleumier, P. White-matter

connectivity between face-responsive regions in the human brain. *Cereb. Cortex* **22**,

1564–1576 (2012).

42. Weiner, K. S. *et al.* The face-processing network is resilient to focal resection of human

visual cortex. *J. Neurosci.* **36**, 8425–8440 (2016).

43. Rossion, B., Dricot, L., Goebel, R. & Busigny, T. Holistic face categorization in higher order

visual areas of the normal and prosopagnosic brain: Toward a non-hierarchical view of

face perception. *Front. Hum. Neurosci.* (2011). doi:10.3389/fnhum.2010.00225

44. Stigliani, A., Jeska, B. & Grill-Spector, K. Encoding model of temporal processing in human

visual cortex. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E11047-E11056; (2017).

45. Stigliani, A., Weiner, K. S. & Grill-Spector, K. Temporal processing capacity in high-level

visual cortex is domain specific. *J. Neurosci.* **35**, 12412–12424 (2015).

46. Song, S. *et al.* Local but not long-range microstructural differences of the ventral

temporal cortex in developmental prosopagnosia. *Neuropsychologia* (2015).

doi:10.1016/j.neuropsychologia.2015.10.010

47. Luke, S. G. Evaluating significance in linear mixed-effects models in R. *Behav. Res.*

Methods (2017). doi:10.3758/s13428-016-0809-y

48. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in Linear

Mixed Effects Models . *J. Stat. Softw.* (2017). doi:10.18637/jss.v082.i13

49. Wandell, B. A. & Winawer, J. Computational neuroimaging and population receptive

fields. **19**, 349–357 (2015).

50. Barton, B. & Brewer, A. A. Visual field map clusters in high-order visual processing:

Organization of V3a/V3B and a new cloverleaf cluster in the posterior superior temporal

sulcus. *Front. Integr. Neurosci.* (2017). doi:10.3389/fnint.2017.00004

51.   Smith, R. E., Tournier, J.-D., Calamante, F. & Connelly, A. Anatomically-constrained

tractography: Improved diffusion MRI streamlines tractography through effective use of

anatomical information. *Neuroimage* **62**, 1924–1938 (2012).

52.   Janssens, T., Zhu, Q. Q., Popivanov, I. D. & Vanduffel, W. Probabilistic and Single-Subject

Retinotopic Maps Reveal the Topographic Organization of Face Patches in the Macaque

Cortex. *J. Neurosci.* (2014). doi:10.1523/JNEUROSCI.2914-13.2013

53.   Gosselin, F. & Schyns, P. G. Bubbles: A technique to reveal the use of information in

recognition tasks. *Vision Res.* (2001). doi:10.1016/S0042-6989(01)00097-9

54.   O'Toole, A. J. *et al.* Recognizing people from dynamic and static faces and bodies:

Dissecting identity with a fusion approach. *Vision Res.* (2011).

doi:10.1016/j.visres.2010.09.035

55.   Felleman, D. J. & Van Essen, D. C. Distributed hierarchical processing in the primate

cerebral cortex. *Cereb. Cortex* (1991). doi:10.1093/cercor/1.1.1

56.   Arcaro, M. J. & Livingstone, M. S. A hierarchical, retinotopic proto-organization of the

primate visual system at birth. *Elife* **6**, (2017).

57.   Butt, O. H., Benson, N. C., Datta, R. & Aguirre, G. K. The Fine-Scale Functional Correlation

of Striate Cortex in Sighted and Blind People. *J. Neurosci.* (2013).

doi:10.1523/JNEUROSCI.0363-13.2013

58.   Arcaro, M. J., Honey, C. J., Mruczek, R. E. E. B., Kastner, S. & Hasson, U. Widespread

correlation patterns of fMRI signal across visual cortex reflect eccentricity organization.

Elife **4**, (2015).

59.   Saygin, Z. M. *et al.* Anatomical connectivity patterns predict face selectivity in the

fusiform gyrus. *Nat. Neurosci.* **15**, 321–327 (2012).

60.   Formisano, E. *et al.* Mirror-symmetric tonotopic maps in human primary auditory cortex.

Neuron (2003). doi:10.1016/S0896-6273(03)00669-X

61.   Harvey, B. M., Klein, B. P., Petridou, N. & Dumoulin, S. O. Topographic representation of

numerosity in the human parietal cortex. *Science (80-.).* (2013).

doi:10.1126/science.1239052

62.   Mezer, A. *et al.* Quantifying the local tissue volume and composition in individual brains

with magnetic resonance imaging. *Nat. Med.* **19**, (2013).

63. Epstein, R. & Kanwisher, N. A cortical representation the local visual environment.
Nature (1998). doi:10.1038/33402

64. Kay, K. N., Winawer, J., Mezer, A. & Wandell, B. A. Compressive spatial summation in
human visual cortex. *J. Neurophysiol.* **110**, 481–494 (2013).

65. Tournier, J. D. *et al.* MRtrix3: A fast, flexible and open software framework for medical
image processing and visualisation. *NeuroImage* (2019).
doi:10.1016/j.neuroimage.2019.116137

66. Lerma-Usabiaga, G., Mukherjee, P., Ren, Z., Perry, M. L. & Wandell, B. A. Replication and
generalization in applied neuroimaging. *Neuroimage* (2019).
doi:10.1016/j.neuroimage.2019.116048

67. Grotheer, M., Zhen, Z., Lerma-Usabiaga, G. & Grill-Spector, K. Separate lanes for adding
and reading in the white matter highways of the human brain. *Nat. Commun.* (2019).
doi:10.1038/s41467-019-11424-1

68. Veraart, J. *et al.* Denoising of diffusion MRI using random matrix theory. *Neuroimage*
(2016). doi:10.1016/j.neuroimage.2016.08.016

69. Takemura, H., Caiafa, C. F., Wandell, B. A. & Pestilli, F. Ensemble Tractography. *PLoS*
Comput. Biol. **12**, e1004692 (2016).

70. Pestilli, F., Yeatman, J. D., Rokem, A., Kay, K. N. & Wandell, B. A. Evaluation and statistical
inference for human connectomes. *Nat. Methods* **11**, 1058–1063 (2014).

Acknowledgements

This work was supported by a William R. and Sara Hart Kimball Stanford Graduate Fellowship

awarded to DF; NIH grants (grant numbers 1RO1EY02231801, 1RO1EY02391501 to KGS); an NIH

training grant (grant number 5T32EY020485 to VN); Ruth L. Kirschstein National Research Service

Award (grant number F31EY027201 to JG) and the NSF Graduate Research Development Program

(grant number DGE-114747 to JG).

Author contributions

D.F. designed the experiment, collected data, analyzed the data, and wrote the manuscript.

J.G. collected the data, contributed to data analysis and contributed to the manuscript.

M.N., and A.A.R., collected the data, and contributed to the manuscript.

S.P. contributed to data analysis and contributed to the manuscript.

K.G-S. designed the experiment, contributed to the data analysis, and wrote the manuscript.