Fungal microbiomes are determined by host phylogeny and exhibit widespread associations with the bacterial microbiome

Xavier A. Harrison[1] [#], Allan D. McDevitt[2], Jenny C. Dunn[3], Sarah Griffiths[4], Chiara Benvenuto[2],

Richard Birtles[2], Jean P. Boubli[2], Kevin Bown[2], Calum Bridson[4,5], Darren Brooks[2], Samuel S.

Browett[2], Ruth F. Carden[6,7], Julian Chantrey[8], Friederike Clever[4,9], Ilaria Coscia[2], Katie L. Edwards[10],

Natalie Ferry[2], Ian Goodhead[2], Andrew Highlands[2], Jane Hopper[11], Joseph Jackson[2], Robert Jehle[2],

Mariane da Cruz Kaizer[2], Tony King[11,12], Jessica M. D. Lea[5], Jessica L. Lenka[2], Alexandra

McCubbin[13], Jack McKenzie[2], Bárbara Lins Caldas de Moraes[14], Denise B. O'Meara[15], Poppy

Pescod[2], Richard F. Preziosi[4], Jennifer K. Rowntree[4], Susanne Shultz[5], Matthew J. Silk[1], Jennifer E.

Stockdale[13,16], William O. C. Symondson[13], Mariana Villalba de la Pena[5], Susan L. Walker[10], Michael

D. Wood[2], Rachael E. Antwis[2] [#]

[#] Corresponding authors:

Dr Rachael Antwis, University of Salford, r.e.antwis@salford.ac.uk

Dr Xavier Harrison, University of Exeter, x.harrison@exeter.ac.uk

1. School of Biosciences, University of Exeter, UK. 2. School of Science, Engineering and Environment, University of Salford,
UK. 3. School of Life Sciences, Joseph Banks Laboratories, University of Lincoln, UK 4. Ecology and Environment Research
Centre, Department of Natural Sciences, Manchester Metropolitan University, UK. 5. School of Earth and Environmental
Sciences, University of Manchester, UK. 6. School of Archaeology, University College Dublin, Ireland. 7. Wildlife Ecological &
Osteological Consultancy, Wicklow, Ireland. 8. Institute of Veterinary Science, University of Liverpool, UK. 9. Smithsonian
Tropical Research Institute, Ancon, Republic of Panama. 10. North of England Zoological Society, Chester Zoo, Upton-by-
Chester, UK. 11. The Aspinall Foundation, Port Lympne Reserve, Hythe, Kent, UK. 12. School of Anthropology and
Conservation, University of Kent. 13. School of Biosciences, University of Cardiff, UK. 14. Department of Zoology, Federal
University of Pernambuco, Recife, Brazil. 15. School of Science & Computing, Waterford Institute of Technology, Ireland. 16.
School of Life Sciences, University of Nottingham, UK

RUNNING TITLE: Cross-kingdom microbiome phylosymbiosis and associations

KEYWORDS: amplicon sequencing, bacteria, co-occurrence, diet, diversity, fungi, host-microbe

interactions, mycobiome, phylosymbiosis,

ABSTRACT

Interactions between hosts and their resident microbial communities are a fundamental component of fitness for both agents. Though recent research has highlighted the importance of interactions between animals and their bacterial communities, comparative evidence for fungi is lacking, especially in natural populations. Using data from 49 species, we present novel evidence of strong covariation between fungal and bacterial communities across the host phylogeny, indicative of recruitment by hosts for specific suites of microbes. Using co-occurrence networks, we demonstrate that fungi form critical components of putative microbial interaction networks, where the strength and frequency of interactions varies with host taxonomy. Host phylogeny drives differences in overall richness of bacterial and fungal communities, but the effect of diet on richness was only evident in mammals and for the bacterial microbiome. Collectively these data indicate fungal microbiomes may play a key role in host fitness and suggest an urgent need to study multiple agents of the animal microbiome to accurately determine the strength and ecological significance of host-microbe interactions.

SIGNIFICANCE STATEMENT

Microbes perform vital metabolic functions that shape the physiology of their hosts. However, almost all research to date in wild animals has focused exclusively on the bacterial microbiota, to the exclusion of other microbial groups. Although likely to be critical components of the host microbiome, we have limited knowledge of the drivers of fungal composition across host species. Here we show that fungal community composition is determined by host species identity and phylogeny, and that fungi form extensive interaction networks with bacteria in the microbiome of a diverse range of animal species. This highlights the importance of microbial interactions as mediators of microbiome-health relationships in the wild.

INTRODUCTION

Multicellular organisms support diverse microbial communities critical for physiological functioning, immunity, development, evolution and behaviour (1–3). Variability in host-associated microbiome composition may explain asymmetries among hosts in key traits including susceptibility to disease (4, 5), fecundity (6), and resilience to environmental change (7). Although the microbiota is a complex assemblage of bacteria, fungi, archaea, viruses and protozoa, the overwhelming majority of research has focused solely on the bacterial component (8, 9). Although relatively well documented in soils and plants (10–13), relatively few studies have examined the dynamics of non-bacterial components of the microbiome in animal hosts (but see (14–16)), especially in non-model or wild systems. As such, our current understanding of host-microbe interactions is skewed by a bacteria-centric view of the microbiome. Although not well understood, there is growing evidence that the fungal microbiota, termed the 'mycobiome', may drive diverse functions such as fat, carbon and nitrogen metabolism (17, 18), degradation of cellulose and other carbohydrates (19), pathogen resistance (20), initiation of immune pathways and regulation of inflammatory responses (9, 21), and even host dispersal (22).

Host phylogeny has repeatedly been shown to be an important predictor of bacterial microbiome structure in multiple vertebrate clades, a phenomenon known as 'phylosymbiosis' (23–27). This phenomenon often reflects phylogenetic patterns in life history traits, such as diet, physiology or spatial distribution (23–27). However, evidence of phylosymbiosis, and its drivers, in other microbial kingdoms or domains is lacking. Addressing this major gap in our knowledge is crucial as we likely underestimate the strength and importance of coevolution between animal hosts and their resident communities, particularly in the context of cross-kingdom interactions within the microbiome (28).

Here we used ITS and 16S rRNA gene amplicon sequencing to characterise fungal and bacterial communities of primarily gut and faecal samples from 49 host species across eight classes, including both vertebrates and invertebrates (Table S1). We predicted that both fungal and bacterial microbiomes demonstrated strong signals of phylosymbiosis across the broad host taxonomic range tested. Specifically, we predicted that patterns of phylosymbiosis within microbial kingdoms will also drive significant positive covariance in patterns of microbial community structure between microbial kingdoms within individual hosts, suggestive of evolutionary constraints that favour co-selection of

86   specific bacterial and fungal communities in tandem. We also used network analysis to identify key

87   bacteria-fungi interactions whilst quantifying variation in the frequency and strength of bacteria-fungi

88   interaction networks across host taxonomic groups. Finally, we tested the prediction that cross-

89   kingdom phylosymbiosis may be partially driven by similarity in host dietary niche across the 32 bird

90   and mammal species sampled.

91
RESULTS

Fungal and Bacterial Microbiome Diversity Varies with Host Phylogeny

Our data revealed consistent patterns in fungal and bacterial alpha diversity across host taxonomic

groups. Bacterial community alpha-diversity was generally greater than, or similar to, fungal

community alpha-diversity at the host species level (Fig. 1A), although two species exhibited greater

fungal diversity than bacterial (great tit, tsetse fly; Fig. 1A). Comparisons between microbial richness

values within individuals (i.e., *relative* richness) using a binomial GLMM supported these patterns,

indicating that bacterial richness was higher on average than fungal in 80% of individuals [95%

credible interval (CI) 0.55 - 0.95]. When conditioning on Class, samples from both Mammalia and

Insecta were more likely to have higher bacterial diversity than fungal diversity (credible intervals not

crossing zero on the link scale). Mammalia were more likely to have higher bacterial relative to fungal

diversity than Aves in our study organisms (mean difference in probability 22.9% [1.6 - 45.7%]).

Variation among species in this model explained 19.5% [7.3 - 31.2%] of the variation in relative

microbial richness. Using a bivariate model with both fungal and bacterial diversity as response

variables to examine patterns of absolute microbial richness across host taxonomy, only Mammalia

exhibited bacterial diversity that was consistently higher than fungal diversity when controlling for

variation among species (mean difference in index 5.16; [3.33 - 6.96]). There was no evidence of

positive covariance between fungal and bacterial richness values at the species level (mean

correlation 0.3, 95% credible intervals -0.55 - 0.86), suggesting that high diversity of one microbial

group does not necessarily reflect high diversity of the other. The bivariate model also revealed that

species identity explained 33.9% [22.2 – 44.2%] of variation in bacterial diversity, and 22.4% [9.8 –

35.5%] of variation in fungal diversity.

Phylogenetic analyses supported these general patterns (Fig. S2). For fungi, we detected

phylogenetic signal in patterns of both Inverse Simpson index (C_{mean} = 0.22, p = 0.021) and number of

observed amplicon sequence variants (ASVs) (C_{mean} = 0.26, p = 0.016). For bacteria, phylogenetic

signal was evident for number of ASVs (C_{mean} = 0.28, p = 0.016) but not inverse Simpson index (C_{mean}

= 0.114, p = 0.100).

FIGURE 1

Host phylogeny and diet as predictors of host bacterial and fungal alpha diversity. **(A)** Boxplots and

raw data (points) of inverse Simpson indices for bacterial (green) and fungal (orange) communities

across a range of host species. **(B)** Raw data (points) and model predictions (shaded area and lines)

of models examining the relationship between host diet and microbiome alpha diversity. In mammals,

an increase the in the amount of plant material in the diet (more negative PC1 values) drives

increases in richness. There was no corresponding relationship between diet and richness for fungi in

mammals, nor for bacteria and fungi in birds. Shaded areas represent 95% credible intervals.

Limited Evidence of Covariation Between Host Diet and Fungal Microbiome

Alpha Diversity: Models exploring the influence of diet on microbial richness yielded mixed

results. In mammals, only a relationship between *bacterial* richness and diet was evident (interaction

between microbe (fungi vs bacteria) and the primary axis of a PCA of dietary variation; Fig 1B). This

indicates that bacterial alpha diversity increases in tandem with the proportion of plant matter in the

diet. However, this relationship was absent in birds (Fig. 1B). Similarly, there was no relationship

between *fungal* richness and diet for birds or mammals (credible intervals for slopes all include zero).

Beta Diversity: Patterns of variation in microbial community *structure* broadly followed those

for alpha diversity above. While for mammals there was a significant correlation between host-

associated bacterial community composition and diet ($r = 0.334$, $p = 0.002$), and a near-significant

relationship between fungal community composition and diet ($r = 0.142$, $p = 0.067$), for birds there

was no significant relationship between dietary data and bacterial community composition ($r = 0.087$,

$p = 0.211$) or fungal community composition ($r = 0.026$, $p = 0.386$). Further, taxonomic differences in

microbiome composition based on differences in crude dietary patterns were not clear for either

bacteria or fungi when the microbiome composition was visualised at the family level (Figs. S3, S4).

That said, Alphaproteobacteria and Eurotiomycete fungi were notably absent from species that

primarily ate vegetation (i.e. grasses etc) and Neocallimastigomycete fungi were the predominant

fungal class associated with two out of four of these host species (Figs. S3, S4).

Strong Evidence of Correlated Phylosymbiosis in Both Microbial Groups

Our data revealed consistent variation in fungal and bacterial community structure across the host

phylogeny (Fig. 2A). PERMANOVA analyses on centred-log ratio (CLR) transformed ASV

abundances revealed significant phylogenetic effects of host class, order and species, as well as

effects of sample storage and library preparation protocol for both microbial groups (Table 2; Figs. S5

& S6). For both bacteria and fungi, host species identity explained more variation than host class or

order, and this pattern remained when re-running the models without sample preparation protocol

effects, though this inflated the estimate of R^2 for all taxonomic groupings (Table 2).

157   Consistent with our predictions, the similarity between the microbial communities of a given

158   pair of host species was proportional to the phylogenetic distance between them (e.g. ASV level:

159   fungal cor. = 0.26; p = 0.001; bacterial cor. = 0.37; p = 0.001; Fig. 2B). Correlations for both bacterial

160   and fungal communities became stronger when aggregating microbial taxonomy to broader

161   taxonomic levels (Fig. 2B). Notably, the bacterial correlation was stronger than the fungal equivalent

162   at most taxonomic levels (Fig. 2B), indicating stronger patterns of phylosymbiosis for bacteria.

163   We also detected a strong, significant correlation between fungal and bacterial community

structure of individual samples at the level of ASVs using Procrustes rotation (cor. = 0.29, p < 0.001;

Fig. 2C). Collapsing ASV taxonomy to genus, family, and order resulted in even stronger correlations

(cor. = 0.44, 0.48 & 0.43, respectively; all p < 0.001; Fig. 2C). These data indicate a coupling between

the structures of fungal and bacterial communities, whereby shifts in structure of one community

across the phylogeny also reflect consistent shifts in the other microbial group.

FIGURE 2

(A) Phylogenetic tree of host species, with branches coloured by class and node points coloured by

order. Barplots show proportional composition of fungal and bacterial phyla for each host species,

aligned to tree tips. **(B)** Correlation between microbial and host genetic distances (generated from the

phylogenetic tree in A) for both bacteria (green) and fungi (orange) across all host species. Microbial

taxonomy was either raw ASVs or grouped into higher taxonomic levels. Aggregation to higher

taxonomy tended to result in higher correlations for both microbial groups, and the correlation was

always stronger in bacteria. **(C)** Correlation between fungal and bacterial community structure derived

from Procrustes rotation on PCA ordinations of each microbial group. Microbial communities were

aggregated at various taxonomic groupings (order, family, genus), or as raw Amplicon Sequence

Variant (ASV) taxonomy. For both B and C, distributions of correlation values were generated using

resampling of 90% of available samples for that microbial group to generate 95% intervals (shaded

areas on graphs). Empty bars in panel 2A mean samples were not available for a particular species

and so would not have been included in the calculations in panel B or C.

TABLE 2

PERMANOVA results for (a) fungi and (b) bacteria of factors explaining variation in microbial

community structure. Terms were added in the order shown in the table to marginalise effects of

sample storage and preparation protocols before calculating % variance explained for taxonomic

groupings. Species ID was the dominant source of variation in the data for both taxonomic groups, but

there were also strong effects of sample storage and wet lab protocol, particularly for bacteria.

(a) FUNGI				Taxonomic Effects Only		
Predictor	df	R^2	p value	df	R^2	p value
Sample Type	7	0.05	0.001			
Tissue Storage	5	0.04	0.001			
Extraction Kit	7	0.07	0.001			
Class	2	0.02	0.001	6	0.05	0.001
Order	6	0.05	0.001	13	0.12	0.001
Species	18	0.09	0.001	26	0.14	0.001
Residuals	303	0.68		303	0.68	

(b) BACTERIA				Taxonomic Effects Only		
Predictor	df	R^2	p value	df	R^2	p value
Sample Type	6	0.06	0.001			
Tissue Storage	6	0.16	0.001			
Extraction Kit	7	0.12	0.001			

Class	2	0.02	0.001	6	0.09	0.001
Order	6	0.09	0.001	12	0.21	0.001
Species	18	0.12	0.001	27	0.27	0.001
Residuals	273	0.42		273	0.42	

Strength of Interactions Between Bacteria and Fungi May Vary Across Host Taxonomy

Analysis of correlations among fungal and bacterial abundances revealed differences in network

structure at both the host class (Fig. 3A) and host species level (Figs. S7; S8). In particular, fungi of

the phylum Ascomycota appeared frequently in the putative interaction networks of birds, mammals

and amphibians (Fig. 3A). There was also systematic variation in network structure among taxonomic

groups. Using the class-level network data in Fig. 3A, we estimated that Mammalia exhibited the

fewest components, fewest communities, and lowest modularity (Table 2), indicating lower overall

network subdivision relative to other animal classes. Mean betweenness of fungal nodes also varied

by host class; randomisations revealed that mean fungal betweenness was significantly lower than

expected by chance in Aves (2-tailed $p = 0.044$, Fig. 3B) but not Mammalia (2-tailed $p=0.6$, Fig 3B).

Models of species-level network data (Fig. S7, S8) revealed the frequency of positive co-occurrence

between pairs of microbes also varied by class; Mammalia exhibited the highest proportion of positive

edges (Fig. 3C), being significantly greater than those of birds (mean diff. 0.042 [0.017-0.067]) and

amphibians (mean diff. 0.05 [0.002-0.112]). Notably, insects had a markedly lower proportion of

positive edges compared to all other taxa (Fig. 3C). Class explained 93.2% [92.9-93.4%] of variation

in edge sign.

TABLE 2: Network statistics from class-specific microbial networks in Figure 3 in the main

manuscript. 'Modularity' and 'Groups' statistics are derived from the cluster_fast_greedy function

applied to *igraph* network objects. 'Components' data were extracted directly from the networks.

Modularity was positively correlated with both number of groups (cor = 0.76) and number of

components (cor = 0.86).

Class	Modularity	Groups	Components
Mammalia	0.658	7	1

Aves	0.719	23	14	215
Insecta	0.781	10	6	
Actinopterygii	0.806	16	11	216
Amphibia	0.923	35	35	
				217

FIGURE 3

(A) Putative microbial interaction networks between bacterial (circles) and fungal (squares) taxa,

coloured by microbial phylum. Networks were constructed using the R package *SpiecEasi* on CLR-

transformed abundance values to detect non-random co-occurrence between groups of microbes. **(B)**

Permutational testing revealed that mean fungal betweenness was significantly lower than expected

by chance in Aves, but not Mammalia, indicating heterogeneity in network structure. **(C)** Analysis of

network structural traits from species-specific networks comprising 39 species from five Classes.

There were significant differences in the proportion of positive edges (correlations between paired

microbial abundance values) among classes. Vertical dashed line indicates equal proportion of

positive and negative edges.

DISCUSSION

Our study represents the most wide-ranging evaluation of animal mycobiome composition, and its

covariation with the bacterial microbiome, undertaken to date. Our data provide novel evidence for

mycobiome phylosymbiosis in wild animals, indicative of close evolutionary coupling between hosts

and their resident fungal communities. Consistent with previous studies, we also find evidence of

phylosymbiosis in the bacterial microbiome (29), but crucially, we demonstrate strong and consistent

covariation between fungal and bacterial communities across host phylogeny, especially at higher

microbial taxonomic levels. These patterns are supported by complementary network analysis

illustrating frequent correlative links between fungal and bacterial taxa, whereby certain pairs of

microbes from different kingdoms are much more likely to co-occur in the microbiome than expected

by chance. Taken together, these data provide novel evidence of host recruitment for specific fungal

and bacterial communities, which in turn may reflect host selection for interactions between bacteria

and fungi critical for host physiology and health.

We found marked variation among host species in microbial community richness and

composition for both bacteria and fungi. Though our data suggest many species support a diverse

assemblage of host-associated fungi, we show that bacterial diversity tends to be higher on average

relative to fungal diversity, and that there is no signal of positive covariance between fungal and

bacterial richness within species, suggesting more ASV-rich bacterial microbiomes are not

consistently associated with more ASV-rich mycobiomes. These patterns could arise because of

competition for niche space within the gut, where high bacterial diversity may reflect stronger

competition that prevents proliferation of fungal diversity. Understanding patterns of niche competition

within and among microbial groups requires that we are able to define those niches by measuring

microbial gene function, and quantifying degree of overlap or redundancy in functional genomic

profiles across bacteria and fungi.

We detected strong phylosymbiosis for both fungi and bacteria across a broad host

phylogeny encompassing both vertebrate and invertebrate classes. This pattern was significantly

stronger in bacteria than for fungi. In both microbial kingdoms, the signal of phylosymbiosis

strengthened when aggregating microbial assignments to broader taxonomic levels, a phenomenon

that has previously been shown for bacterial communities (30). That this pattern also occurs in fungi

suggests either that host recruitment is weaker at finer-scale taxonomies, or our ability to detect that

signal is weaker at the relatively noisy taxonomic scale of ASVs. Stronger signals of phylosymbiosis

at family and order-level taxonomies may reflect the deep evolutionary relationships between hosts

and their bacterial and fungal communities, as well as the propensity for microbial communities to

allow closely related microbes to establish whilst repelling less related organisms (31). That is, higher-

order microbial taxonomy may better approximate functional guilds within the microbiome, such as the

ability to degrade cellulose (25, 30), which are otherwise obscured by taxonomic patterns of ASVs.

Resolving this requires the integration of functional genomic data from the fungal and bacterial

microbiota into the phylogeny.

In addition to microbe-specific patterns of phylosymbiosis, a key novel finding of our work is

discovery of strong covariation between fungal and bacterial community composition across the host

phylogeny. These patterns are consistent with host recruitment for particular suites of fungal and

bacterial taxa, which may represent bacteria-fungi metabolic interactions beneficial to the host.

Bacterial-fungal interactions have previously been demonstrated for a handful of animal species (8, 9,

17, 32, 33), but here we show these are widespread across multiple animal classes. Both bacteria

and fungi have considerable enzymatic properties that facilitate the liberation of nutrients for use by

other microbes, thus facilitating cross-kingdom colonisation (34–36) and promoting metabolic inter-

dependencies (37–39). We also identified numerous associations between bacteria and fungi for

many of our host species. The frequency and predicted direction of these relationships varied

considerably among host classes, with the mammalian network exhibiting i) a lower modularity,

indicating weaker clustering into fewer discrete units (both distinct components and interlinked

communities); and ii) a higher frequency of positive correlations between microbes compared to most

other classes, in particular birds and insects. Comparisons of networks are challenging when they

differ in size (i.e., number of nodes) and structure, and differences between classes in traits like

modularity will also be affected by species replication within each class. However, proportional traits

like interaction structure (proportion of positive interactions) are unlikely to be driven solely by sample

size, suggesting marked biological variation in strength of fungi-bacteria interactions across the host

phylogeny. These putative interaction networks provide novel candidates for further investigation in

controlled systems, where microbiome composition and therefore the interactions among microbes

can be manipulated to test the influence of such interactions on host physiology.

The drivers of phylosymbiosis remain unclear, even for bacterial communities; is a

phylogenetic signal indicative of host-microbiome coevolution, or simply a product of "ecological

filtering" of the microbiome in the host organism either via extrinsic (e.g. diet, habitat) or intrinsic

sources (e.g. gut pH, immune system function) (26, 29, 40)? Our results indicate host diet may play a

role in determining bacterial composition in mammals, but not fungal composition in either mammals

or birds. These results are broadly consistent with previous work, where the influence of diet on

bacterial microbiome was most evident in mammals (25). However, Li et al. (16) showed that the

composition and diversity of both fungal and bacterial communities of faecal samples differed

between phytophagous and insectivorous bats, and Heisel et al. (17) demonstrated changes in fungal

community composition in mice fed a high fat diet. Our study was not designed to test for the effects

of ecological variation in diet on fungal microbiome *within* a species, nor can we discount the

possibility that at finer taxonomic scales within classes, signals of the effect of *among* species

variation in diet on mycobiome may become stronger (e.g. (16)). It is also worth noting that the signals

produced from faecal and true gut samples may differ; evidence suggests faecal samples may

indicate diet is the predominant driver of "gut" microbiome composition when gastrointestinal samples

indicate host species is the predominant determinant (41). Moreover, faecal samples may only

represent a small proportion of the gastrointestinal microbiome (41–43). Our data also show that

sample type has a significant effect on both fungal and bacterial community composition (as well as

DNA extraction method and storage method; see (44–47) for other examples of this). As such, a more

thorough analysis of true gut communities is required to determine the extent to which mycobiome

phylosymbiosis and dietary signals occur across wild animals, and what other ecological and host-

associated factors influence mycobiome composition and function. We hypothesise that evolutionary

processes play a large role in shaping host-associated microbiomes, with selection for microbiome

function rather than taxonomic groupings per se.

Within animals, the roles of host-associated fungal communities are not well understood, yet

our data highlight that fungi are important components of microbiome structure that are often

overlooked. Our knowledge of the range of functions provided by the host mycobiome, and how these

alter or complement those provided by the bacterial microbiome, remains limited. We hypothesise that

host-associated fungi and bacteria produce mutually beneficial metabolites that facilitate the

colonisation, reproduction and function of cross-kingdom metabolic networks (28). Though we provide

evidence for consistent variation among host class in fungal community structure, and the role of fungi

within putative interaction networks, for many researchers the questions of key interest will focus on

what governs variation at the level of the individual. Clear gaps in our knowledge remain regarding the

relative contributions of host genomic (48–50) and environmental variation to host mycobiome

structure, function and stability. We argue that there is an urgent need to incorporate quantitative

estimates of microbial function into microbiome studies, which are crucial for understanding the forces

of selection shaping host-microbe interactions at both the individual and species level.

MATERIALS AND METHODS

Sample collection

DNA was extracted from tissue or faecal samples of 49 host species using a variety of DNA extraction

methods (Table S1) and normalised to ~10 ng/ul. Samples were largely collated from previous studies

and/or those available from numerous researchers and as such, DNA extraction and storage

techniques were not standardised across species. We sequenced a median of 10 samples per

species (range of 5 to 12; Table S1).

ITS1F-2 and 16S rRNA amplicon sequencing

Full details are provided in Supplementary Materials. Briefly, we amplified the ITS1F-2 rRNA gene to

identify fungal communities using single index reverse primers and a modified protocol of Smith &

Peay (51) and Nguyen et al. (52), as detailed in Griffiths et al. (13). To identify bacterial communities,

we amplified DNA for the 16S rRNA V4 region using dual indexed forward and reverse primers

according to Kozich et al. (53) and Griffiths et al. (49). The two libraries were sequenced separately

using paired-end reads (2 x 250bp) with v2 chemistry on an Illumina MiSeq.

We conducted amplicon sequence data processing in DADA2 v1.5 (54) in RStudio v1.2.1335

for R (55, 56) for both ITS rRNA and 16S rRNA amplicon data. After data processing, we obtained a

median of 1425 reads per sample (range of 153 to 424,527) from the ITS data, and a median of 3273

reads (range of 153 to 425,179) for the 16S rRNA data.

To compare alpha-diversity between species and microbial kingdoms, we rarefied libraries to

500 reads per sample, yielding 292 samples from 46 species and 307 samples from 47 species for

fungal and bacterial kingdoms respectively. Alpha-diversity measures remained relatively stable within

a host species whether data were rarefied to 500, 1000, or 2500 reads (Figs. 1, S1, S2; see

Supplementary Material for more details).

Host phylogeny

As many of our host species lack genomic resources from which to construct a genome-based

phylogeny, we built a dated phylogeny of host species using TimeTree (57). The phylogenetic tree

contained 42 species, of which 36 were directly represented in the TimeTree database. A further six

species had no direct match in TimeTree and so we used a congener as a substitute (*Amietia,*

Glossina, Portunus, Ircinia, Amblyomma, Cinachyrella). We calculated patristic distance among

species based on shared branch length in the phylogeny using the 'cophenetic' function in the *ape*

package (58) in R. We visualised and annotated the phylogeny using the R package *ggtree* (59). To

create a phylogeny for all samples, we grafted sample-level tips onto the species phylogeny with

negligible branch lengths following Youngblut et al. (25).

Fungal and bacterial community analysis

A fully reproducible workflow of all analyses is provided in supplementary material as an R

Markdown document. We used the R package *brms* (60, 61) to fit (generalized) linear mixed effects

models [(G)LMMs] to test for differences in alpha diversity and calculated r^2 of models using the

'bayes_R2' function. We assessed the importance of terms based on whether 95% credible intervals

of the parameter estimates of interest crossed zero. We used *ggplot* (62), *cowplot* (63) and *tidybayes*

(64) for raw data and plotting of posterior model estimates.

To support these analyses, we also used the R packages *phylobase* (65) and *phylosignal*

(66) to estimate the phylogenetic signal in patterns of alpha diversity for both bacteria and fungi, using

both Inverse Simpson Index and number of observed ASVs as outcome variables. We calculated

Abouheif's C_{mean} for each diversity-microbe combination and corrected p values for multiple testing

using Benjamini-Hochberg correction.

To identify taxonomic differences in microbiome and mycobiome composition between host

species, we used centred-log-ratio (CLR) transformation in the *microbiome* (67) package to normalise

microbial abundance data, which obviates the need to lose data through rarefying (68). To quantify

differences in beta-diversity among kingdoms and species whilst simultaneously accounting for

sample storage and library preparation differences among samples, we conducted a PERMANOVA

analysis on among-sample Euclidean distances of CLR-transformed abundances using the *adonis*

function in *vegan* (69) with 999 permutations. For both kingdoms, we specified effects in the following

order: sample type, tissue storage, extraction kit, class, order, species. This marginalises the effects

of sample metadata variables first, before partitioning the remaining variance into that accounted for

by host phylogeny. The results were similar when amplicon data were converted to relative

abundance or rarefied to 500 reads (data not presented).

To test the hypothesis that inter-individual differences in microbial community composition

were preserved between microbial kingdoms, we performed Procrustes rotation of the two PCA

ordinations for bacterial and fungal abundance matrices, respectively (n = 277 paired samples from

46 species). We also repeated this analysis with ASVs agglomerated into progressively higher

taxonomic rankings from genus to order (see (30)). To provide a formal test of differences in strength

of covariation at different taxonomic levels, we conducted a bootstrap resampling analysis where for

each kingdom at each iteration, we randomly sampled 90% of the data and recalculated the

correlation metric. We repeated this process 999 times to build a distribution of correlation values at

each taxonomic grouping. To examine the hypothesis that inter-individual distance in microbial

community composition varies in concert with interspecific phylogenetic distance, we performed a

Procrustes rotation on the paired matrix of microbial distance (Euclidean distance of CLR-transformed

abundances) and patristic distance from the phylogenetic tree.

To identify potential co-occurrence relationships between fungal and bacterial communities,

we conducted two analyses; 1) We used the R package *SpiecEasi* (70) to identify correlations

between unrarefied, CLR-transformed ASVs abundances at the host class level (with insects

grouped), and 2) we used co-occurrence analysis at the species level, by rarefying the bacterial and

fungal data sets to 500 reads each, and agglomerated taxonomy family level, resulting in 117

bacterial groups and 110 fungal groups. We then merged the *phyloseq* objects for bacterial and

fungal communities for each sample, with sufficient data retained to conduct the co-occurrence

analysis for 40 host species. Using these cross-kingdom data, we calculated the co-occurrence

between each pair of microbial genera by constructing a Spearman's correlation coefficient matrix in

the *bioDist* package (71, 72). We visualised those with rho > 0.50 (strong positive interactions) and

rho < -0.50 (strong negative interactions) for each host species separately using network plots

produced in *igraph* (73). We calculated modularity of the class-level microbial networks comprising

both positive and negative interactions using the modularity function after greedy clustering

implemented in the *igraph* package. We used binomial GLM to test the hypothesis that the proportion

of positive edges (correlations) varies by host class, and permutation approaches on betweenness

values of fungal nodes to test the hypothesis that fungi form critical components of microbial

networks.

To determine the effect of diet on bacterial and fungal community composition, we used only

samples from the bird and mammal species and agglomerated the data for each host species using

the merge_samples function in *phyloseq* (74). This gave us a representative microbiome for each

host species, which we rarefied to the lowest number of reads for each combination of kingdom and

host taxon (2,916 – 9,160 reads; bacterial read counts were low for lesser horseshoe bats and so this

species was removed from this analysis) and extracted Euclidean distance matrices for each. We

then correlated these with dietary data obtained from the EltonTraits database (75) using Mantel tests

with Kendall rank correlations in the *vegan* package (69). We agglomerated the microbial data to

class level and visualised the bacterial and fungal community compositions for mammals alongside

pie charts displaying EltonTrait dietary data for each species. We also used a primary axis of the

ordination of EltonTrait data to derive a 'dietary variation axis' used as a predictor for alpha diversity of

Birds and Mammals.

ACKNOWLEDGEMENTS

We would like to thank Miran Aprahamian, Chris Williams (Environment Agency) Patrick Abila

(National Livestock Resources Research Institute) and Dr Patrick Vudriko (Makerere University) for

providing samples, as well as BEI Resources, the US Forest Service and the University of Wisconsin-

Madison for providing mock communities. We are grateful to Devenish Nutrition for funding the red

deer research at Dowth Hall, Co. Meath, Ireland. The sampling of shrews was funded by a Heredity

Fieldwork Grant awarded by the Genetics Society. The collection of dove and pigeon faecal samples

was jointly funded by the Royal Society for the Protection of Birds and Natural England through the

Action for Birds in England (AfBiE) partnership. Fieldwork enabling collection of avian faecal samples

from Lincolnshire was funded by The Royal Society Research Grant RG170086 to JCD. Small

mammal sampling in the Chernobyl Exclusion Zone was supported by the TREE

(https://tree.ceh.ac.uk/) and RED FIRE (https://www.ceh.ac.uk/redfire) projects. TREE was funded by

the Natural Environment Research Council (NERC), Radioactive Waste Management Ltd. and the

Environment Agency as part of the RATE Programme; RED FIRE was a NERC Urgency Grant.

Northern muriqui research at Caparaó National Park, Brazil, was funded by CAPES (BEX 1298/15-1).

COMPETING INTERESTS

The authors have no competing interests to declare.

DATA ACCESSIBILITY STATEMENT

Sequence data are deposited in the NCBI SRA database under BioProject numbers PRJNA593927

and PRJNA593220. A fully-reproducible analysis workflow has been provided as supplementary

material at https://github.com/xavharrison/Mycobiome2020

REFERENCES

1.   E. A. Archie, J. Tung, Social behavior and the microbiome. *Curr. Opin. Behav. Sci.* **6**, 28–34
        (2015).

2.   M. McFall-Ngai, *et al.*, Animals in a bacterial world, a new imperative for the life sciences.

459   *Proc. Natl. Acad. Sci. U. S. A.* **110**, 3229–3236 (2013).

460   3.   S. Bahrndorff, S. Bahrndorff, T. Alemu, T. Alemneh, J. L. Nielsen, The Microbiome of Animals :

461   Implications for Conservation Biology. *Int. J. Genomics* **2016**, 5304028 (2016).

462   4.   R. J. Dillon, C. T. Vennard, A. Buckling, A. K. Charnley, Diversity of locust gut bacteria

463   protects against pathogen invasion. *Ecol. Lett.* **8**, 1291–1298 (2005).

464   5.   K. A. Bates, *et al.*, Amphibian chytridiomycosis outbreak dynamics are linked with host skin

465   bacterial community structure. *Nat. Commun.* **9**, 1–11 (2018).

466   6.   B. S. Walsh, C. Heys, Z. Lewis, Gut microbiota influences female choice and fecundity in the

467   nuptial gift-giving species, Drosophila subobscura (Diptera: Drosophilidae). *Eur. J. Entomol.*

468   **114**, 439–445 (2017).

469   7.   A. Apprill, Marine animal microbiomes: Toward understanding host-microbiome interactions in

470   a changing ocean. *Front. Mar. Sci.* **4**, 1–9 (2017).

471   8.   Q. H. Sam, M. W. Chang, L. Y. A. Chai, The fungal mycobiome and its interaction with gut

472   bacteria in the host. *Int. J. Mol. Sci.* **18**, 330 (2017).

473   9.   R. Enaud, *et al.*, The Mycobiome: A Neglected Component in the Microbiota-Gut-Brain Axis.

474   *Microorganisms* **6**, 22 (2018).

475   10.   P. Frey-Klett, *et al.*, Bacterial-Fungal Interactions: Hyphens between Agricultural, Clinical,

476   Environmental, and Food Microbiologists. *Microbiol. Mol. Biol. Rev.* **75**, 583–609 (2011).

477   11.   B. Jakuschkin, *et al.*, Deciphering the Pathobiome: Intra- and Interkingdom Interactions

478   Involving the Pathogen Erysiphe alphitoides. *Microb. Ecol.* **72**, 870–880 (2016).

479   12.   L. S. van Overbeek, K. Saikkonen, Impact of Bacterial-Fungal Interactions on the Colonization

480   of the Endosphere. *Trends Plant Sci.* **21**, 230–242 (2016).

481   13.   S. M. Griffiths, *et al.*, Complex associations between cross-kingdom microbial endophytes and

482   host genotype in ash dieback disease dynamics. *J. Ecol.*, 1–19 (2019).

483   14.   M. A. Ghannoum, *et al.*, Characterization of the oral fungal microbiome (mycobiome) in

484   healthy individuals. *PLoS Pathog.* **6** (2010).

485   15.   A. S. Liggenstoffer, N. H. Youssef, M. B. Couger, M. S. Elshahed, Phylogenetic diversity and

486   community structure of anaerobic gut fungi (phylum Neocallimastigomycota) in ruminant and

487   non-ruminant herbivores. *ISME J.* **4**, 1225–1235 (2010).

488   16.   J. Li, *et al.*, Fecal Bacteriome and Mycobiome in Bats with Diverse Diets in South China. *Curr.*

489   *Microbiol.* **75**, 1352–1361 (2018).

490   17.   T. Heisel, *et al.*, High-Fat Diet Changes Fungal Microbiomes and Interkingdom Relationships

491   in the Murine Gut. *mSphere* **2**, 1–14 (2017).

492   18.   L. Wegley, R. Edwards, B. Rodriguez-Brito, H. Liu, F. Rohwer, Metagenomic analysis of the

493   microbial community associated with the coral Porites astreoides. *Environ. Microbiol.* **9**, 2707–

494   2719 (2007).

495   19.   S. Yang, *et al.*, Metagenomic Analysis of Bacteria, Fungi, Bacteriophages, and Helminths in

496   the Gut of Giant Pandas. *Front. Microbiol.* **9**, 1–16 (2018).

497   20.   P. J. Kearns, *et al.*, Fight fungi with fungi: Antifungal properties of the amphibian mycobiome.

498   *Front. Microbiol.* **8**, 1–12 (2017).

499   21.   F. Yeung, *et al.*, Altered Immunity of Laboratory Mice in the Natural Environment Is Associated

500   with Fungal Colonization. *Cell Host Microbe*, 1–14 (2020).

501   22.   M. Lu, M. J. Wingfield, N. E. Gillette, S. R. Mori, J. H. Sun, Complex interactions among host

502   pines and fungi vectored by an invasive bark beetle. *New Phytol.* **187**, 859–866 (2010).

503   23.   E. R. Davenport, *et al.*, The human microbiome in evolution. *BMC Biol.* **15**, 1–12 (2017).

504   24.   S. C. L. Knowles, R. M. Eccles, L. Baltrūnaitė, Species identity dominates over environment in

505   shaping the microbiota of small mammals. *Ecol. Lett.* **22**, 826–837 (2019).

506   25.   N. D. Youngblut, *et al.*, Host diet and evolutionary history explain different aspects of gut

507   microbiome diversity among vertebrate clades. *Nat. Commun.* **10**, 1–15 (2019).

508   26.   K. R. Amato, *et al.*, Evolutionary trends in host physiology outweigh dietary niche in structuring

509   primate gut microbiomes. *ISME J.* **13**, 576–587 (2019).

510   27.   S. J. Song, *et al.*, Comparative Analyses of Vertebrate Gut Microbiomes Reveal Convergence

511   between Birds and Bats. *MBio* **11**, 1–14 (2020).

28.   K. A. Krautkramer, J. Fan, F. Bäckhed, Gut microbial metabolites as multi-kingdom
intermediates. *Nat. Rev. Microbiol.* (2020) https:/doi.org/https://doi.org/10.1038/s41579-020-
0438-4.

29.   A. W. Brooks, K. D. Kohl, R. M. Brucker, E. J. van Opstal, S. R. Bordenstein, Phylosymbiosis:
Relationships and Functional Effects of Microbial Communities across Host Evolutionary
History. *PLoS Biol.* **14**, 1–29 (2016).

30.   P. Rausch, *et al.*, Comparative analysis of amplicon and metagenomic sequencing methods
reveals key features in the evolution of animal metaorganisms. *Microbiome* **7**, 1–19 (2019).

31.   B. Stecher, *et al.*, Like will to like: Abundances of closely related species can predict
susceptibility to intestinal colonization by pathogenic and commensal bacteria. *PLoS Pathog.*
6, e1000711 (2010).

32.   Q. S. Mcfrederick, U. G. Mueller, R. R. James, Interactions between fungi and bacteria
influence microbial community structure in the Megachile rotundata larval gut Interactions
between fungi and bacteria influence microbial community structure in the Megachile
rotundata larval gut. *Proc. R. Soc. B Biol. Sci.* **281**, 20132653 (2014).

33.   J. D. Forbes, C. N. Bernstein, H. Tremlett, G. Van Domselaar, N. C. Knox, A fungal world:
Could the gut mycobiome be involved in neurological disease? *Front. Microbiol.* **10**, 1–13
(2019).

34.   S. Pareek, *et al.*, Comparison of Japanese and Indian intestinal microbiota shows diet-
dependent interaction between bacteria and fungi. *npj Biofilms Microbiomes* **5** (2019).

35.   K. G. Peay, P. G. Kennedy, T. D. Bruns, Fungal Community Ecology: A Hybrid Beast with a
Molecular Master. *Bioscience* **58**, 799–810 (2008).

36.   N. Fierer, Embracing the unknown: Disentangling the complexities of the soil microbiome. *Nat.*
Rev. Microbiol. **15**, 579–590 (2017).

37.   C. Hoffmann, *et al.*, Archaea and Fungi of the Human Gut Microbiome: Correlations with Diet
and Bacterial Residents. *PLoS One* **8**, e66019 (2013).

38.   W. Kim, S. B. Levy, K. R. Foster, Rapid radiation in bacteria leads to a division of labour. *Nat.*

Commun. **7**, 10508 (2016).

39.   S. Rakoff-Nahoum, K. R. Foster, L. E. Comstock, The evolution of cooperation within the gut
microbiota. *Nature* **533**, 255–259 (2016).

40.   F. Mazel, *et al.*, Is Host Filtering the Main Driver of Phylosymbiosis across the Tree of Life?
mSystems **3**, 1–15 (2018).

41.   M. R. Ingala, *et al.*, Comparing microbiome sampling methods in a wild mammal: Fecal and
intestinal samples record different signals of host ecology, evolution. *Front. Microbiol.* **9**, 1–13
(2018).

42.   E. Videvall, M. Strandh, A. Engelbrecht, S. Cloete, C. K. Cornwallis, Measuring the gut
microbiome in birds: Comparison of faecal and cloacal sampling. *Mol. Ecol. Resour.* **18**, 424–
434 (2018).

43.   G. Leite, *et al.*, Sa1911 – Analysis of the Small Intestinal Microbiome Reveals Marked
Differences from Stool Microbiome in a Large Scale Human Cohort: Redefining the "Gut
Microbiome." *Gastroenterology* **156**, S-449-S-450 (2019).

44.   S. Yuan, D. B. Cohen, J. Ravel, Z. Abdo, L. J. Forney, Evaluation of methods for the extraction
and purification of DNA from the human microbiome. *PLoS One* **7** (2012).

45.   L. Weber, E. DeForce, A. Apprill, Optimization of DNA extraction for advancing coral
microbiota investigations. *Microbiome* **5**, 1–14 (2017).

46.   X. Zhou, S. Nanayakkara, J. L. Gao, K. A. Nguyen, C. J. Adler, Storage media and not
extraction method has the biggest impact on recovery of bacteria from the oral microbiome.
Sci. Rep. **9**, 1–10 (2019).

47.   A. E. Asangba, *et al.*, Variations in the microbiome due to storage preservatives are not large
enough to obscure variations due to factors such as host population, host species, body site,
and captivity. *Am. J. Primatol.* **81**, 1–12 (2019).

48.   J. K. Goodrich, *et al.*, Human genetics shape the gut microbiome. *Cell* **159**, 789–799 (2014).

49.   S. M. Griffiths, *et al.*, Genetic variability and ontogeny predict microbiome structure in a
disease-challenged montane amphibian. *ISME J.* **12**, 2506–2517 (2018).

50.   S. M. Griffiths, *et al.*, Host genetics and geography influence microbiome composition in the
sponge Ircinia campana. *J. Anim. Ecol.* **88**, 1684–1695 (2019).

51.   D. P. Smith, K. G. Peay, Sequence depth, not PCR replication, improves ecological inference
from next generation DNA sequencing. *PLoS One* **9**, e90234 (2014).

52.   N. H. Nguyen, D. Smith, K. Peay, P. Kennedy, Parsing ecological signal from noise in next
generation amplicon sequencing (2014).

53.   J. J. Kozich, S. L. Westcott, N. T. Baxter, S. K. Highlander, P. D. Schloss, Development of a
dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on
the miseq illumina sequencing platform. *Appl. Environ. Microbiol.* **79**, 5112–5120 (2013).

54.   B. J. Callahan, *et al.*, DADA2: High-resolution sample inference from Illumina amplicon data.
Nat. Methods **13**, 581–583 (2016).

55.   RStudio Team, RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL
http://www.rstudio.c (2016).

56.   R Core Team, R: A language and environment for statistical computing. R Foundation for
Statistical Computing. Vienna, Austria. URL https://www.R-project.org/. (2017).

57.   S. Kumar, G. Stecher, M. Suleski, S. B. Hedges, TimeTree: A Resource for Timelines,
Timetrees, and Divergence Times. *Mol. Biol. Evol.* **34**, 1812–1819 (2017).

58.   E. Paradis, K. Schliep, Ape 5.0: An environment for modern phylogenetics and evolutionary
analyses in R. *Bioinformatics* **35**, 526–528 (2019).

59.   G. Yu, D. K. Smith, H. Zhu, Y. Guan, T. T. Y. Lam, Ggtree: an R Package for Visualization and
Annotation of Phylogenetic Trees With Their Covariates and Other Associated Data. *Methods*
Ecol. Evol. **8**, 28–36 (2017).

60.   P. C. Bürkner, Advanced Bayesian multilevel modeling with the R package brms. *R J.* **10**,
395–411 (2018).

61.   P. C. Bürkner, brms: An R package for Bayesian multilevel models using Stan. *J. Stat. Softw.*
80 (2017).

62.   H. Wickham, *ggplot2: Elegant Graphics for Data Analysis.* (Springer-Verlag New York, 2009).

63.   C. O. Wilke, cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2". R package

version 1.0.0. https://CRAN.R-project.org/package=cowplot (2019).

64.   M. Kay, tidybayes: Tidy Data and Geoms for Bayesian Models. R package version 2.0.3,

http://mjskay.github.io/tidybayes/ (2020).

65.   R Hackathon, phylobase: Base Package for Phylogenetic Structures and Comparative Data. R

package version 0.8.10. https://CRAN.R-project.org/package=phylobase (2020).

66.   F. Keck, F. Rimet, A. Bouchez, A. Franc, phylosignal: an R package to measure, test, and

explore the phylogenetic signal. *Ecol. Evol.* **6**, 2774–2780 (2016).

67.   L. Lahti, S. Shetty, Tools for microbiome analysis in R. Microbiome package version

1.1.10013. http://microbiome.github.com/microbiome. (2017).

68.   G. B. Gloor, J. M. Macklaim, V. Pawlowsky-Glahn, J. J. Egozcue, Microbiome datasets are

compositional: And this is not optional. *Front. Microbiol.* **8**, 1–6 (2017).

69.   J. Oksanen, *et al.*, vegan: Community Ecology Package. (2018).

70.   Z. D. Kurtz, *et al.*, Sparse and Compositionally Robust Inference of Microbial Ecological

Networks. *PLoS Comput. Biol.* **11**, 1–25 (2015).

71.   R. J. Williams, A. Howe, K. S. Hofmockel, Demonstrating microbial co-occurrence pattern

analyses within and between ecosystems. *Front. Microbiol.* **5**, 1–10 (2014).

72.   B. Ding, R. Gentleman, V. Carey, bioDist: Different distance measures. R package version

1.54.0. (2018).

73.   G. Csárdi, T. Nepusz, The igraph software package for complex network research.

InterJournal Complex Syst. **1695**, 1695 (2006).

74.   P. J. McMurdie, S. Holmes, Phyloseq: An R Package for Reproducible Interactive Analysis

and Graphics of Microbiome Census Data. *PLoS One* **8**, e61217 (2013).

75.   H. Wilman, *et al.*, EltonTraits 1.0 : Species-level foraging attributes of the world's birds and

mammals. *Ecology* **95**, 2027 (2014).

618