Host-associated fungal communities are determined by host phylogeny and exhibit widespread associations with the bacterial microbiome

Xavier A. Harrison[1] #, Allan D. McDevitt[2], Jenny C. Dunn[3], Sarah Griffiths[4], Chiara Benvenuto[2], Richard Birtles[2], Jean P. Boubli[2], Kevin Bown[2], Calum Bridson[4,5], Darren Brooks[2], Samuel S. Browett[2], Ruth F. Carden[6,7], Julian Chantrey[8], Friederike Clever[4,9], Ilaria Coscia[2], Katie L. Edwards[10], Natalie Ferry[2], Ian Goodhead[2], Andrew Highlands[2], Jane Hopper[11], Joseph Jackson[2], Robert Jehle[2], Mariane da Cruz Kaizer[2], Tony King[11,12], Jessica M. D. Lea[5], Jessica L. Lenka[2], Alexandra McCubbin[13], Jack McKenzie[2], Bárbara Lins Caldas de Moraes[14], Denise B. O'Meara[15], Poppy Pescod[2], Richard F. Preziosi[4], Jennifer K. Rowntree[4], Susanne Shultz[5], Matthew J. Silk[1], Jennifer E. Stockdale[13,16], William O. C. Symondson[13], Mariana Villalba de la Pena[5], Susan L. Walker[10], Michael D. Wood[2], Rachael E. Antwis[2] #

Corresponding authors:

Dr Rachael Antwis, University of Salford, r.e.antwis@salford.ac.uk

Dr Xavier Harrison, University of Exeter, x.harrison@exeter.ac.uk

1. School of Biosciences, University of Exeter, UK. 2. School of Science, Engineering and Environment, University of Salford, UK. 3. School of Life Sciences, Joseph Banks Laboratories, University of Lincoln, UK 4. Ecology and Environment Research Centre, Department of Natural Sciences, Manchester Metropolitan University, UK. 5. School of Earth and Environmental Sciences, University of Manchester, UK. 6. School of Archaeology, University College Dublin, Ireland. 7. Wildlife Ecological & Osteological Consultancy, Wicklow, Ireland. 8. Institute of Veterinary Science, University of Liverpool, UK. 9. Smithsonian Tropical Research Institute, Ancon, Republic of Panama. 10. North of England Zoological Society, Chester Zoo, Upton-by-Chester, UK. 11. The Aspinall Foundation, Port Lympne Reserve, Hythe, Kent, UK. 12. School of Anthropology and Conservation, University of Kent. 13. School of Biosciences, University of Cardiff, UK. 14. Department of Zoology, Federal University of Pernambuco, Recife, Brazil. 15. School of Science & Computing, Waterford Institute of Technology, Ireland. 16. School of Life Sciences, University of Nottingham, UK

ABSTRACT

Interactions between hosts and their resident microbial communities are a fundamental component of

fitness for both agents. Despite a recent proliferation of research on interactions between animals and

their associated bacterial communities, comparative evidence from fungal communities is lacking,

especially in natural populations. This disparity means knowledge of host-microbe interactions is

biased towards the bacterial microbiome. Using samples from 49 species from eight metazoan

classes, we demonstrate that the ecological distance between both fungal and bacterial components

of the microbiome shift in tandem with host phylogenetic distance. Though so-called phylosymbiosis

has been shown in bacterial communities, we extend previous knowledge by demonstrating that the

magnitude of shifts in fungal and bacterial community structure across host phylogeny are correlated.

These data are indicative of coordinated recruitment by hosts for specific suites of microbes, and

potentially selection for bacterial-fungal interactions across a broad taxonomic range of host species.

Using co-occurrence networks comprising both microbial groups, we illustrate that fungi form a critical

component of microbial interaction networks, and that the strength and frequency of such interactions

vary across host taxonomic groups. Collectively these data indicate fungal microbiomes may play a

key role in host fitness and suggest an urgent need to study multiple agents of the animal microbiome

to accurately determine the strength and ecological significance of host-microbe interactions.

INTRODUCTION

Multicellular organisms support diverse microbial communities that are critical for physiological

functioning, immunity, development, evolution, behaviour, and even conservation[1–3]. Variability in

host-associated microbiome composition may explain asymmetries among hosts in key traits

including susceptibility to disease[4,5], fecundity[6], and resilience to environmental change[7]. Although the

microbiota is a complex assemblage of bacteria, fungi, archaea, viruses and protozoa, the

overwhelming majority of research published to date has focused solely on exploring interactions

between the host and the bacterial component of its microbiome[8,9]. Despite fungal-bacterial

interactions being relatively well documented in soils and plants[10–13], relatively few studies have

examined the dynamics of non-bacterial components of the microbiome in animal hosts (but see [14–16]), especially in non-model organisms or wild systems. As such, our current understanding of the

strength of host-microbe interactions is likely skewed by a bacteria-centric view of the microbiome.

The roles of animal-associated mycobiomes are currently not well understood; however, a

small but growing body of research has identified the potential importance of resident fungal

microbiota, termed the 'mycobiome', for host animal health. This may include diverse functions such

as fat, carbon and nitrogen metabolism[17,18], degradation of cellulose and other carbohydrates[19],

pathogen resistance[20], initiating immune pathways and regulating inflammatory responses[9,21], and

even host dispersal[22]. Host phylogeny has repeatedly been shown to be an important predictor of

bacterial microbiome structure in multiple vertebrate clades, a phenomenon known as

'phylosymbiosis'[23–27]. However, evidence of phylosymbiosis from other microbial kingdoms or

domains is lacking and remains a major gap in our knowledge. Addressing this shortfall is vital as we

likely underestimate the strength and importance of coevolution between animal hosts and their

resident fungal communities. In addition, studying multiple microbial groups in concert will allow us to

identify positive and negative covariances across microbial kingdoms in the abundance of taxonomic

and/or functional groups that may reflect selection for interactions among microbes that are crucial

determinants of animal health.

Here we used ITS and 16S rRNA amplicon sequencing to characterise fungal and bacterial

communities of primarily gut and faecal samples from 49 host species across eight classes, including

both vertebrates and invertebrates (Table 1). We predicted that both fungal and bacterial microbiomes

demonstrated strong signals of phylosymbiosis across the broad host taxonomic range we test here.

Specifically, we predicted that patterns of phylosymbiosis within microbial kingdoms will also drive

significant positive covariance in patterns of microbial community structure between microbial

kingdoms within individual hosts, suggestive of evolutionary constraints that favour co-selection of

specific bacterial and fungal communities in tandem. Additionally, we used network analysis to identify

key bacteria-fungi interactions whilst quantifying variation in the frequency and strength of bacteria-

fungi interaction networks across host taxonomic groups. Finally, we investigate the prediction that

cross-kingdom phylosymbiosis may be partially driven by similarity in host dietary niche across the 32

bird and mammal species.

RESULTS

Alpha-diversity

Our data revealed consistent patterns in fungal and bacterial alpha-diversity across host taxonomic

groups. Bacterial community alpha-diversity was generally greater than, or similar to, fungal

community alpha-diversity at the host species level (Fig. 1A), although two species exhibited greater

fungal diversity than bacterial (great tit, tsetse fly; Fig. 1A). Alpha-diversity measures remained

relatively stable within a host species whether data were rarefied to 500, 1000, or 2500 reads (Figs. 1,

S1, S2; Supplementary Material). Comparisons between microbial richness values within individuals

(i.e. *relative* richness) using a binomial GLMM supported these patterns, indicating that bacterial

richness was higher on average in 80% of cases [95% credible interval (CI) 0.55 - 0.95]. When

conditioning on Class, samples from both Mammalia and Insecta were more likely to have higher

bacterial diversity than fungal diversity (credible intervals not crossing zero on the link scale; Fig. 1B),

though all classes had posterior mean values >0 (>50% on the probability scale). Indeed, Mammalia

were more likely to have higher bacterial relative to fungal diversity than Aves in our study organisms

(mean difference in probability 22.9% [1.6 - 45.7%). Variation among species in this model explained

19.5% [7.3 - 31.2%] of the variation in relative microbial richness. Using a bivariate model with both

fungal and bacterial diversity as response variables to examine patterns of *absolute* microbial

richness across host taxonomy, only Mammalia exhibited bacterial diversity that was consistently

higher than fungal diversity when controlling for variation among species (mean difference in index

5.16; [3.33 - 6.96]; Fig. 1C). This model differs from the binomial version because it accounts for the

magnitude of difference in microbial richness, rather than just relative richness. There was no

evidence of positive covariance between fungal and bacterial richness values at the species level

(mean correlation 0.3, 95% credible intervals -0.55 - 0.86), suggesting that high diversity of one

microbial group does not necessarily reflect high diversity of the other. The bivariate model also

revealed that species identity explained 33.9% [22.2 – 44.2%] of variation in bacterial diversity, and

22.4% [9.8 – 35.5%] of variation in fungal diversity. Phylogenetic analyses supported these general

patterns (Fig. S2). For fungi, we detected phylogenetic signal in patterns of both Inverse Simpson

index (C_{mean} = 0.22, p = 0.021) and number of observed amplicon sequence variants (ASVs) (C_{mean} =

0.26, p = 0.016). For bacteria, phylogenetic signal was evident for number of ASVs (C_{mean} = 0.28, p =

0.016) but not inverse Simpson index (C_{mean} = 0.114, p = 0.100).

Beta-diversity

The five most abundant classes of fungi across all host species were Dothideomycetes,

Eurotiomycetes, Lecanoromycetes, Pezizomycetes and Sordariomycetes, for which Dothideomycetes

and Eurotiomycetes showed the most variation between host species (Fig. S3). The five most

abundant classes of bacteria were Actinobacteria, Alphaproteobacteria, Bacilli, Bacteroidia, and

Gammaproteobacteria, which all varied considerably among host species (Fig. S3).

Both fungal and bacterial community composition at the phylum level varied considerably in

concert with host class and order (Fig. 2A). However, notable trends include dominance of the fungal

phylum Ascomycota in most species, with the exception of the two Perissodactyla (odd-toed ungulate)

species, where most fungal ASVs belonged to the Neocallimastigomycota. Likewise, the bacterial

phylum Firmicutes dominated most mammal species, but Proteobacteria were more common on

average in birds and fish (Fig. 2A). Analysis of factors affecting bacterial and fungal community

structure using PERMANOVA on centred-log ratio (CLR) transformed ASV abundances revealed

significant phylogenetic effects of host class, order and species, as well as effects of sample storage

and library preparation protocol (Table 2; Fig. S4). For both microbial kingdoms, host species identity

explained more variation than host class or order, and this pattern remained when re-running the

models without sample preparation protocol effects, though this inflated the estimate of R^2 for all

taxonomic groupings (Table 2).

Phylosymbiosis

Consistent with our predictions, the similarity between the microbial communities of a given pair of

host species was proportional to the phylogenetic distance between them (ASV level: fungal cor. =

0.26; p = 0.001; bacterial cor. = 0.37; p = 0.001; Fig. 2B). Correlations for both bacterial and fungal

communities became stronger when aggregating microbial taxonomy to family (Fig. 2B). Critically, the

bacterial correlation was stronger than the fungal equivalent at both ASV (mean diff. 0.077 [0.053 -

0.103]) and family (mean diff. 0.10 [0.082 - 0.13]) levels (Fig. 2B), indicating stronger patterns of

phylosymbiosis for bacteria.

We detected a strong, significant correlation between fungal and bacterial community

structure of individual samples at the level of ASVs using Procrustes rotation (cor. = 0.29, p < 0.001;

Fig. 2C). Collapsing ASV taxonomy to genus, family, and order resulted in even stronger correlations

(cor. = 0.44, 0.48 & 0.43, respectively; all p < 0.001; Fig. 2C). These data indicate a coupling between

the structures of fungal and bacterial communities, whereby shifts in structure of one community

across the phylogeny also reflect consistent shifts in the other microbial group.

Network analysis

Analysis of correlations among fungal and bacterial abundances revealed differences in network

structure at both the host class (Fig. 3) and host species level (Figs. S5; S6). In particular, fungi of the

phylum Ascomycota appeared frequently in the putative interaction networks of birds, mammals and

amphibians (Fig. 3). There was also systematic variation in network structure among taxonomic

groups. Using the class-level network data in Fig. 3, we detected that Mammalia exhibited the fewest

components, fewest communities, and lowest modularity (Fig. 4A, Table S1), indicating lower overall

network subdivision relative to other animal classes. Models of species-level network data (Fig. S5)

revealed the frequency of positive co-occurrence between pairs of microbes also varied by class;

Mammalia exhibited the highest proportion of positive edges (Fig. 4B), being significantly greater than

those of birds (mean diff. 0.042 [0.017-0.067]) and amphibians (mean diff. 0.05 [0.002-0.112]).

Notably, invertebrates had a markedly lower proportion of positive edges compared to all other taxa

(Fig. 4). Class explained 93.2% [92.9-93.4%] of variation in edge sign. There was also clear variation

at the species level; for some host species, there were considerably more positive interactions (e.g.

yellowhammers, pygmy shrews, greater white-toothed shrews, wood mouse, woodpigeon, yellow-

necked mouse; Fig. S6). In some species, there were slightly more negative interactions than positive

(e.g. blackcap, goldfinch; Fig. S5; S6).

Dietary analysis

While there was a significant correlation between host-associated bacterial community composition

and dietary data for mammals (r = 0.334, p = 0.002), and a near-significant relationship between

fungal community composition and diet (r = 0.142, p = 0.067), there was no significant relationship

between dietary data and bacterial community composition (r = 0.087, p = 0.211) or fungal community

composition (r = 0.026, p = 0.386) for birds. Further, taxonomic differences in microbiome composition

based on differences in crude dietary patterns were not clear for either bacteria or fungi when the

microbiome composition was visualised at the family level (Figs. S7, S8). That said,

Alphaproteobacteria and Eurotiomycete fungi were notably absent from species that primarily ate

vegetation (i.e. grasses etc) and Neocallimastigomycete fungi were the predominant fungal class

associated with two out of four of these host species (Figs. S7, S8).

DISCUSSION

Our study represents the most wide-ranging evaluation of animal mycobiome composition, and its

covariation with the bacterial microbiome, undertaken to date. Our data provide novel evidence for

mycobiome phylosymbiosis in wild animals, indicative of close evolutionary coupling between hosts

and their resident fungal communities. Consistent with previous studies, we also find evidence of

phylosymbiosis in the bacterial microbiome[28], but crucially, we demonstrate strong and consistent

covariation between fungal and bacterial communities across host phylogeny, especially at higher

microbial taxonomic levels. These patterns are supported by complementary network analysis

illustrating frequent correlative links between fungal and bacterial taxa, whereby certain pairs of

microbes from different kingdoms are much more likely to co-occur in the microbiome than expected

by chance. Taken together, these data provide novel evidence consistent with recruitment by animal

hosts for specific fungal and bacterial communities, which in turn may reflect selection for interactions

between bacteria and fungi critical for host physiology and health.

We found marked variation among host species in microbial community richness and

composition for both bacteria and fungi. Complementary analyses using mixed models and

phylogenetic models both detected a signal of host phylogeny in determining fungal and bacterial

microbiome diversity. Though our data suggest many species support a diverse assemblage of host-

associated fungi, critically we show that i) bacterial diversity tends to be higher on average relative to

fungal diversity; and ii) there is no signal of positive covariance between fungal and bacterial richness

within species, suggesting more ASV-rich bacterial microbiomes are not consistently associated with

more ASV-rich mycobiomes. These patterns could arise because of competition for niche space

within the gut, where high bacterial diversity may reflect stronger competition that prevents

proliferation of fungal diversity. Fungi are enzymatically very active, and their crucial roles in

degrading organic molecules and redistributing carbon and nitrogen in soil and plant ecosystems are

well documented[29,30]. Understanding patterns of niche competition within and among microbial groups

requires that we are able to define those niches by measuring microbial gene function, and

quantifying degree of overlap or redundancy in functional genomic profiles across bacteria and fungi.

We detected strong phylosymbiosis for both fungi and bacteria across a broad host

phylogeny encompassing both vertebrate and invertebrate classes. This pattern was significantly

stronger in bacteria than for fungi. In both microbial kingdoms, the signal of phylosymbiosis

strengthened when aggregating microbial taxonomic assignments to family level, a phenomenon that

has previously been shown for bacterial communities[31]. That this pattern also occurs in fungi suggests

either that host recruitment is weaker at finer-scale taxonomies, or our ability to detect that signal is

weaker at the relatively noisy taxonomic scale of ASVs. Stronger signals of phylosymbiosis at family-

level taxonomies may reflect the deep evolutionary relationships between hosts and their bacterial

and fungal communities, as well as the propensity for microbial communities to allow closely related

microbes to establish whilst repelling less related organisms[32]. That is, higher-order microbial

taxonomy may better approximate functional guilds within the microbiome, such as the ability to

degrade cellulose[25,31], which are otherwise obscured by taxonomic patterns of ASVs. Resolving this

requires the integration of functional genomic data from the fungal and bacterial microbiota into the

phylogeny.

In addition to microbe-specific patterns of phylosymbiosis, a key novel finding of our work is

that fungal and bacterial community composition correlate strongly across the host phylogeny. These

patterns are consistent with host recruitment for particular suites of fungal and bacterial taxa, which

may represent bacteria-fungi metabolic interactions beneficial to the host. Bacteria-fungi interactions

have previously been demonstrated for a handful of host species[8,9,17,33,34], but here we show these are

widespread across multiple animal classes. Both bacteria and fungi have considerable enzymatic

properties that facilitate the liberation of nutrients for use by other microbes, thus facilitating cross-

kingdom colonisation[35] and promoting metabolic inter-dependencies[36–38]. These findings are

supported by our network analyses, which identified numerous putative interactions between bacteria

and fungi for many of our host species. Critically, the frequency and predicted direction of these

relationships varied considerably among host classes, with the mammalian network exhibiting i) a

lower modularity, indicating weaker clustering into fewer discrete units (both distinct components and

interlinked communities); and ii) a higher frequency of positive correlations between microbes

compared to most other classes, in particular birds and insects. Comparisons of networks are

challenging when they differ in size (i.e. number of nodes) and structure, and differences between

classes in traits like modularity will also be affected by species replication within each class. However,

proportional traits like interaction structure (proportion of positive interactions) are unlikely to be driven

solely by sample size, suggesting marked biological variation in strength of fungi-bacteria interactions

across the host phylogeny. These putative interaction networks provide novel candidates for further

investigation in controlled systems, where microbiome composition and therefore the interactions

among microbes can be manipulated to test the influence of such interactions on host physiology.

The drivers of phylosymbiosis remain unclear, even for bacterial communities; is it a

phylogenetic signal indicative of host-microbiome coevolution, or simply a product of "ecological

filtering" of the microbiome in the host organism either via extrinsic (e.g. diet, habitat) or intrinsic

sources (e.g. gut pH, immune system function)[26,28,39]? Our results indicate host diet may play a role in

determining overall fungal community composition, although the relationship is weak and only evident

for bacteria in mammals. These results are broadly consistent with previous work, where the influence

of diet on bacterial microbiome was most evident in mammals[25]. However, Li et al.[15] showed that the

composition and diversity of both fungal and bacterial communities of faecal samples differed

between phytophagous and insectivorous bats, and Heisel et al.[17] demonstrated changes in fungal

community composition in mice fed a high fat diet. Our study was not designed to test for the effects

of ecological variation in diet on fungal microbiome *within* a species. Nor can we discount the

possibility that at finer taxonomic scales within classes, signals of the effect of *among* species

variation in diet on mycobiome become stronger (e.g. [15]). Although diet is thought to be a predominant

driver of bacterial microbiome composition in host organisms[25,40], there is also evidence that the

signals produced from faecal and true gut samples differ; that is, data generated from faecal samples

indicate diet is the predominant driver of "gut" microbiome composition, whereas data from

gastrointestinal samples indicate host species is the predominant determinant[41]. Moreover, faecal

samples may only represent a small proportion of the gastrointestinal microbiome[41–43]. Our data also

show that sample type has a significant effect on both fungal and bacterial community composition

(as well as DNA extraction method and storage method; see [44–47] for other examples of this). As such,

a more thorough analysis of true gut communities is required to determine the extent to which

mycobiome phylosymbiosis occurs across host taxa, and what other ecological and host-associated

factors influence mycobiome composition and function.

These data provide strong evidence that both fungal symbionts and fungi-bacteria

interactions are likely to be critical for host functioning and health. Within animals, the roles of host-

associated fungal communities are less well understood, yet these data highlight that fungi are

important components of microbiome structure that are often overlooked. Key priorities for future work

are to i) understand the range of functions provided by the host mycobiome, and how these alter or

complement those provided by the bacterial microbiome; and ii) characterise the functional

interactions between bacteria and fungi and how they influence key host metabolic processes and life

history.

METHODS

Sample collection

DNA was extracted from tissue or faecal samples of 49 host species using a variety of DNA extraction

methods (Table 1) and normalised to ~10 ng/ul. Samples were largely collated from previous studies

and/or those available from numerous researchers and as such, DNA extraction and storage

techniques were not standardised across species. We sequenced a median of 10 samples per

species (range of 5 to 12; Table 1).

ITS1F-2 and 16S rRNA amplicon sequencing and pre-processing

To identify fungal communities, we amplified DNA for the ITS1F-2 rRNA gene using single index

reverse primers and a modified protocol of Smith & Peay[48] and Nguyen et al.[49], as detailed in Griffiths

et al.[13] (see supplementary material for additional information). To identify bacterial communities, we

amplified DNA for the 16S rRNA V4 region using dual indexed forward and reverse primers according

to Kozich et al.[50] and Griffiths et al.[51] (see supplementary material for additional information). We

sequenced normalised libraries using paired-end reads (2 x 250bp) with Illumina v2 chemistry on the

MiSeq platform at the University of Salford. We ran the ITS rRNA library twice to increase sequencing

depth, and combined data within samples across these two runs in the data pre-processing stage.

We conducted all data processing and analysis in RStudio v1.2.1335 for R[52,53] (see

supplementary files for full code). We conducted amplicon sequence processing in DADA2 v1.5[54] for

both ITS rRNA and 16S rRNA amplicon data (see supplementary material for additional information).

After data processing, we obtained a median of 1425 reads per sample (range of 153 to 424,527) for

ITS rRNA libraries, and a median of 3273 reads per sample (range of 153 to 425,179) for 16S rRNA

libraries.

Host Phylogenetic Distances

As many of our host species lack genomic resources from which to construct a genome-based

phylogeny, we built a dated phylogeny of host species using TimeTree[55]. The phylogenetic tree

contained 42 species, of which 36 were directly represented in the TimeTree database. A further six

species had no direct match in TimeTree and so we used a congener as a substitute (*Amietia,*

Glossina, Portunus, Ircinia, Amblyomma, Cinachyrella). We calculated patristic distance among

species based on shared branch length in the phylogeny using the 'cophenetic' function in the *ape*

package[56] in R. We visualised and annotated the phylogeny using the R package *ggtree*[57]. To create

a phylogeny for all samples, we grafted sample-level tips onto the species phylogeny with negligible

branch lengths following Youngblut et al.[25].

Fungal and bacterial community analysis

We used the R package *brms*[58,59] to fit the following (generalized) linear mixed effects models

[(G)LMMs]. For higher order taxonomic predictors and random effects, we binned all invertebrate

classes into a single grouping to improve model performance, as otherwise invertebrate class and

species were co-linear. All vertebrate taxonomic groupings were equivalent to class (Mammalia, Aves

etc). To compare alpha-diversity between species and microbial kingdoms, we rarefied libraries to

500 reads per sample, yielding 292 samples from 46 species and 307 samples from 47 species for

fungal and bacterial kingdoms respectively. Alpha-diversity measures remained relatively stable within

a host species whether data were rarefied to 500, 1000, or 2500 reads, with similar patterns exhibited

between the two kingdoms (Figs. 1, S1, S2; see Supplementary Material for more details). To

visualise differences between microbial richness within species, we filtered the data to species with at

least two samples per microbial kingdom, giving a total of 41 species from six classes. For model

fitting, we filtered the data to only those samples with paired metrics of microbial richness for both

kingdoms (201 observations from 42 species). We fitted two models to these data. First, to quantify

relative differences in richness between bacteria and fungi within a sample, we used GLMMs in the

brms package, with i) Bernoulli errors and a logit link; ii) a binary response of '1' if bacterial richness

was higher than fungal richness, and '0' otherwise; and iii) 'Species' nested within 'Class' as random

intercepts. We did not include intermediate levels of taxonomy because replication at Order and

Family levels was low relative to Class. We did not use a phylogenetic mixed model as not all species

were represented in the TimeTree phylogeny. Second, to quantify absolute differences in microbial

richness, we fitted a bivariate response LMM with both fungal and bacterial richness values as a two-

column response with Class as a fixed effect, and Species as a random intercept. For all models, we

used uninformative Cauchy priors for the random effects and Gaussian priors for fixed effects

coefficients. We assessed model adequacy using visual inspection of chains to assess mixing and

stationarity properties, as well as posterior predictive checks using the 'pp_check' function in *brms*.

We calculated r^2 of models using the 'bayes_R2' function. We assessed the importance of terms

based on whether 95% credible intervals of the parameter estimates of interest crossed zero. We

used *ggplot*[60], *cowplot*[61] and *tidybayes*[62] for raw data and model estimate plotting. To support these

analyses, we also used the R packages *phylobase*[63] and *phylosignal*[64] to estimate the phylogenetic

signal in patterns of alpha diversity for both bacteria and fungi, using both Inverse Simpson Index and

number of observed ASVs as outcome variables. We calculated Abouheif's C_{mean} for each diversity-

microbe combination and corrected p values for multiple testing using Benjamini-Hochberg correction.

To identify taxonomic differences in microbiome and mycobiome composition between host

species, we used centred-log-ratio (CLR) transformation in the *microbiome*[65] package to normalise

microbial abundance data, which obviates the need to lose data through rarefying[66]. To visualise

differences in microbial community structure among samples, we i) plotted proportional abundance of

microbial groups at the phylum level, aligned to the host phylogenetic tree, ii) agglomerated the data

to class level and visualised the variation in CLR-transformed ratios for the five most abundant

microbial classes in each kingdom for each species using jitter plots, and iii) conducted principal

components analysis (PCA) using CLR-transformed abundance matrices for each kingdom. To

quantify differences in beta-diversity among kingdoms and species whilst simultaneously accounting

for sample storage and library preparation differences among samples, we conducted a

PERMANOVA analysis on among-sample Euclidean distances of CLR-transformed abundances

using the adonis function in *vegan*[67] with 999 permutations. For both kingdoms, we specified effects in

the following order: sample type, tissue storage, extraction kit, Class, Order, Species. This

marginalises the effects of sample metadata variables first, before partitioning the remaining variance

into that accounted for by host phylogeny. The results were similar when amplicon data were

converted to relative abundance or rarefied to 500 reads (data not presented).

To test the hypothesis that inter-individual differences in microbial community composition

were preserved between microbial kingdoms, we performed Procrustes rotation of the two PCA

ordinations for bacterial and fungal abundance matrices, respectively (n = 277 paired samples from

46 species). We also repeated this analysis with ASVs agglomerated into progressively higher

taxonomic rankings from genus to order (see [31]). To provide a formal test of differences in strength of

correlation at different taxonomic levels, we conducted a bootstrap resampling analysis where for

each kingdom at each iteration, we randomly sampled 90% of the data and recalculated the

correlation metric. We repeated this process 999 times to build a distribution of correlation values at

each taxonomic grouping.

To examine the hypothesis that inter-individual distance in microbial community composition

varies in concert with inter-host phylogenetic distance, we performed a Procrustes rotation on the

paired matrix of microbial distance (Euclidean distance of CLR-transformed abundances) and patristic

distance from the phylogenetic tree. We also repeated the analysis but binning the microbial data to

family level. As above, we conducted a bootstrap resampling procedure, selecting 90% of the data at

random and recalculating the correlation, for a total of 1000 permutations. This allowed us to test for

significant differences in strength of correlation *within* kingdoms across taxonomic grouping levels,

and *across* kingdoms within a particular taxonomic grouping.

To determine the effect of diet on bacterial and fungal community composition, we used only

samples from the bird and mammal species and agglomerated the data for each host species using

the merge_samples function in *phyloseq*[68]. This gave us an representative microbiome for each host

species, which we rarefied to the lowest number of reads for each combination of kingdom and host

taxon (2,916 – 9,160 reads; bacterial read counts were low for lesser horseshoe bats and so this

species was removed from this analysis) and extracted Euclidean distance matrices for each. We

then correlated these with dietary data obtained from the EltonTraits database[69] using Mantel tests

with Kendall rank correlations in the *vegan* package[67]. We agglomerated the microbial data to class

level and visualised the bacterial and fungal community compositions for mammals alongside pie

charts displaying EltonTrait dietary data for each species.

To identify potential relationships between fungal and bacterial communities, we conducted

two analyses; 1) We used the R package *SpiecEasi*[70] to identify correlations between unrarefied,

CLR-transformed ASVs abundances at the host class level (with Invertebrates grouped), and 2) we

used co-occurrence analysis at the species level, by rarefying the bacterial and fungal data sets to

500 reads each, and agglomerated these to family level, resulting in 117 bacterial groups and 110

fungal groups. We then merged the phyloseq objects for bacterial and fungal communities for each

sample, with sufficient data retained to conduct the co-occurrence analysis for 40 host species. Using

these cross-kingdom data, we calculated the co-occurrence between each pair of microbial genera by

constructing a Spearman's correlation coefficient matrix in the *bioDist* package[71,72]. We visualised

those with rho > 0.50 (strong positive interactions) and rho < -0.50 (strong negative interactions) for

each host species separately using network plots produced in *igraph*[73]. We calculated modularity of

the class-level microbial networks comprising both positive and negative interactions using the

modularity() function after greedy clustering implemented in the *igraph* package. We resampled 90%

of nodes in each network 1000 times to build distributions of modularity with which to quantify

differences among animal classes. We used binomial GLM to test the hypothesis that the proportion

of positive edges (correlations) varies by host class.

ACKNOWLEDGEMENTS

We would like to thank Miran Aprahamian and Chris Williams (Environment Agency) for providing

samples, as well as BEI Resources, the US Forest Service and the University of Wisconsin-Madison

for providing mock communities. We are grateful to Devenish Nutrition for funding the red deer

research at Dowth Hall, Co. Meath, Ireland. The sampling of shrews was funded by a Heredity

Fieldwork Grant awarded by the Genetics Society. The collection of dove and pigeon faecal samples

was jointly funded by the Royal Society for the Protection of Birds and Natural England through the

Action for Birds in England (AfBiE) partnership. Fieldwork enabling collection of avian faecal samples

from Lincolnshire was funded by The Royal Society Research Grant RG170086 to JCD. Small

mammal sampling in the Chernobyl Exclusion Zone was supported by the TREE

(https://tree.ceh.ac.uk/) and RED FIRE (https://www.ceh.ac.uk/redfire) projects. TREE was funded by

the Natural Environment Research Council (NERC), Radioactive Waste Management Ltd. and the

Environment Agency as part of the RATE Programme; RED FIRE was a NERC Urgency Grant.

Northern muriqui research at Caparaó National Park, Brazil, was funded by CAPES (BEX 1298/15-1).

DATA ACCESSIBILITY STATEMENT

Sequence data are deposited in the NCBI SRA database under BioProject numbers PRJNA593927

and PRJNA593220. Full analysis code has been provided as supplementary material.

REFERENCES

1. Archie, E. A. & Tung, J. Social behavior and the microbiome. *Curr. Opin. Behav. Sci.* **6**, 28–34 (2015).

2. McFall-Ngai, M. *et al.* Animals in a bacterial world, a new imperative for the life sciences. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 3229–3236 (2013).

3. Bahrndorff, S., Bahrndorff, S., Alemu, T., Alemneh, T. & Nielsen, J. L. The Microbiome of Animals : Implications for Conservation Biology. *Int. J. Genomics* **2016**, 5304028 (2016).

4. Dillon, R. J., Vennard, C. T., Buckling, A. & Charnley, A. K. Diversity of locust gut bacteria protects against pathogen invasion. *Ecol. Lett.* **8**, 1291–1298 (2005).

5. Bates, K. A. *et al.* Amphibian chytridiomycosis outbreak dynamics are linked with host skin bacterial community structure. *Nat. Commun.* **9**, 1–11 (2018).

6. Walsh, B. S., Heys, C. & Lewis, Z. Gut microbiota influences female choice and fecundity in the nuptial gift-giving species, Drosophila subobscura (Diptera: Drosophilidae). *Eur. J. Entomol.* **114**, 439–445 (2017).

7. Apprill, A. Marine animal microbiomes: Toward understanding host-microbiome interactions in a changing ocean. *Front. Mar. Sci.* **4**, 1–9 (2017).

8. Sam, Q. H., Chang, M. W. & Chai, L. Y. A. The fungal mycobiome and its interaction with gut bacteria in the host. *Int. J. Mol. Sci.* **18**, 330 (2017).

9. Enaud, R. *et al.* The Mycobiome: A Neglected Component in the Microbiota-Gut-Brain Axis. *Microorganisms* **6**, 22 (2018).

10. Frey-Klett, P. *et al.* Bacterial-Fungal Interactions: Hyphens between Agricultural, Clinical, Environmental, and Food Microbiologists. *Microbiol. Mol. Biol. Rev.* **75**, 583–609 (2011).

11. Jakuschkin, B. *et al.* Deciphering the Pathobiome: Intra- and Interkingdom Interactions Involving the Pathogen Erysiphe alphitoides. *Microb. Ecol.* **72**, 870–880 (2016).

12. van Overbeek, L. S. & Saikkonen, K. Impact of Bacterial-Fungal Interactions on the Colonization of the Endosphere. *Trends Plant Sci.* **21**, 230–242 (2016).

13.   Griffiths, S. M. *et al.* Complex associations between cross-kingdom microbial endophytes and
host genotype in ash dieback disease dynamics. *J. Ecol.* 1–19 (2019). doi:10.1111/1365-
2745.13302

14.   Ghannoum, M. A. *et al.* Characterization of the oral fungal microbiome (mycobiome) in healthy
individuals. *PLoS Pathog.* **6**, (2010).

15.   Li, J. *et al.* Fecal Bacteriome and Mycobiome in Bats with Diverse Diets in South China. *Curr.*
Microbiol. **75**, 1352–1361 (2018).

16.   Liggenstoffer, A. S., Youssef, N. H., Couger, M. B. & Elshahed, M. S. Phylogenetic diversity
and community structure of anaerobic gut fungi (phylum Neocallimastigomycota) in ruminant
and non-ruminant herbivores. *ISME J.* **4**, 1225–1235 (2010).

17.   Heisel, T. *et al.* High-Fat Diet Changes Fungal Microbiomes and Interkingdom Relationships in
the Murine Gut. *mSphere* **2**, 1–14 (2017).

18.   Wegley, L., Edwards, R., Rodriguez-Brito, B., Liu, H. & Rohwer, F. Metagenomic analysis of
the microbial community associated with the coral Porites astreoides. *Environ. Microbiol.* **9**,
2707–2719 (2007).

19.   Yang, S. *et al.* Metagenomic Analysis of Bacteria, Fungi, Bacteriophages, and Helminths in the
Gut of Giant Pandas. *Front. Microbiol.* **9**, 1–16 (2018).

20.   Kearns, P. J. *et al.* Fight fungi with fungi: Antifungal properties of the amphibian mycobiome.
Front. Microbiol. **8**, 1–12 (2017).

21.   Yeung, F. *et al.* Altered Immunity of Laboratory Mice in the Natural Environment Is Associated
with Fungal Colonization. *Cell Host Microbe* 1–14 (2020). doi:10.1016/j.chom.2020.02.015

22.   Lu, M., Wingfield, M. J., Gillette, N. E., Mori, S. R. & Sun, J. H. Complex interactions among
host pines and fungi vectored by an invasive bark beetle. *New Phytol.* **187**, 859–866 (2010).

23.   Davenport, E. R. *et al.* The human microbiome in evolution. *BMC Biol.* **15**, 1–12 (2017).

24.   Knowles, S. C. L., Eccles, R. M. & Baltrūnaitė, L. Species identity dominates over environment
in shaping the microbiota of small mammals. *Ecol. Lett.* **22**, 826–837 (2019).

25.   Youngblut, N. D. *et al.* Host diet and evolutionary history explain different aspects of gut

microbiome diversity among vertebrate clades. *Nat. Commun.* **10**, 1–15 (2019).

26.   Amato, K. R. *et al.* Evolutionary trends in host physiology outweigh dietary niche in structuring

primate gut microbiomes. *ISME J.* **13**, 576–587 (2019).

27.   Song, S. J. *et al.* Comparative Analyses of Vertebrate Gut Microbiomes Reveal Convergence

between Birds and Bats. *MBio* **11**, 1–14 (2020).

28.   Brooks, A. W., Kohl, K. D., Brucker, R. M., van Opstal, E. J. & Bordenstein, S. R.

Phylosymbiosis: Relationships and Functional Effects of Microbial Communities across Host

Evolutionary History. *PLoS Biol.* **14**, 1–29 (2016).

29.   Peay, K. G., Kennedy, P. G. & Bruns, T. D. Fungal Community Ecology: A Hybrid Beast with a

Molecular Master. *Bioscience* **58**, 799–810 (2008).

30.   Fierer, N. Embracing the unknown: Disentangling the complexities of the soil microbiome. *Nat.*

Rev. Microbiol. **15**, 579–590 (2017).

31.   Rausch, P. *et al.* Comparative analysis of amplicon and metagenomic sequencing methods

reveals key features in the evolution of animal metaorganisms. *Microbiome* **7**, 1–19 (2019).

32.   Stecher, B. *et al.* Like will to like: Abundances of closely related species can predict

susceptibility to intestinal colonization by pathogenic and commensal bacteria. *PLoS Pathog.*

6, e1000711 (2010).

33.   Mcfrederick, Q. S., Mueller, U. G. & James, R. R. Interactions between fungi and bacteria

influence microbial community structure in the Megachile rotundata larval gut Interactions

between fungi and bacteria influence microbial community structure in the Megachile

rotundata larval gut. *Proc. R. Soc. B Biol. Sci.* **281**, 20132653 (2014).

34.   Forbes, J. D., Bernstein, C. N., Tremlett, H., Van Domselaar, G. & Knox, N. C. A fungal world:

Could the gut mycobiome be involved in neurological disease? *Front. Microbiol.* **10**, 1–13

(2019).

35.   Pareek, S. *et al.* Comparison of Japanese and Indian intestinal microbiota shows diet-

dependent interaction between bacteria and fungi. *npj Biofilms Microbiomes* **5**, (2019).

36.   Hoffmann, C. *et al.* Archaea and Fungi of the Human Gut Microbiome: Correlations with Diet

500  and Bacterial Residents. *PLoS One* **8**, e66019 (2013).

501 37. Kim, W., Levy, S. B. & Foster, K. R. Rapid radiation in bacteria leads to a division of labour.

502  *Nat. Commun.* **7**, 10508 (2016).

503 38. Rakoff-Nahoum, S., Foster, K. R. & Comstock, L. E. The evolution of cooperation within the

504  gut microbiota. *Nature* **533**, 255–259 (2016).

505 39. Mazel, F. *et al.* Is Host Filtering the Main Driver of Phylosymbiosis across the Tree of Life?

506  *mSystems* **3**, 1–15 (2018).

507 40. Muegge, B. D. *et al.* Diet drives convergence in gut microbiome functions across mammalian

508  phylogeny and within humans. *Science (80-.).* **332**, 970–974 (2008).

509 41. Ingala, M. R. *et al.* Comparing microbiome sampling methods in a wild mammal: Fecal and

510  intestinal samples record different signals of host ecology, evolution. *Front. Microbiol.* **9**, 1–13

511  (2018).

512 42. Videvall, E., Strandh, M., Engelbrecht, A., Cloete, S. & Cornwallis, C. K. Measuring the gut

513  microbiome in birds: Comparison of faecal and cloacal sampling. *Mol. Ecol. Resour.* **18**, 424–

514  434 (2018).

515 43. Leite, G. *et al.* Sa1911 – Analysis of the Small Intestinal Microbiome Reveals Marked

516  Differences from Stool Microbiome in a Large Scale Human Cohort: Redefining the "Gut

517  Microbiome". *Gastroenterology* **156**, S-449-S-450 (2019).

518 44. Yuan, S., Cohen, D. B., Ravel, J., Abdo, Z. & Forney, L. J. Evaluation of methods for the

519  extraction and purification of DNA from the human microbiome. *PLoS One* **7**, (2012).

520 45. Weber, L., DeForce, E. & Apprill, A. Optimization of DNA extraction for advancing coral

521  microbiota investigations. *Microbiome* **5**, 1–14 (2017).

522 46. Zhou, X., Nanayakkara, S., Gao, J. L., Nguyen, K. A. & Adler, C. J. Storage media and not

523  extraction method has the biggest impact on recovery of bacteria from the oral microbiome.

524  *Sci. Rep.* **9**, 1–10 (2019).

525 47. Asangba, A. E. *et al.* Variations in the microbiome due to storage preservatives are not large

526  enough to obscure variations due to factors such as host population, host species, body site,

and captivity. *Am. J. Primatol.* **81**, 1–12 (2019).

48.   Smith, D. P. & Peay, K. G. Sequence depth, not PCR replication, improves ecological

inference from next generation DNA sequencing. *PLoS One* **9**, e90234 (2014).

49.   Nguyen, N. H., Smith, D., Peay, K. & Kennedy, P. Parsing ecological signal from noise in next

generation amplicon sequencing. (2014).

50.   Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K. & Schloss, P. D. Development of

a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data

on the miseq illumina sequencing platform. *Appl. Environ. Microbiol.* **79**, 5112–5120 (2013).

51.   Griffiths, S. M. *et al.* Genetic variability and ontogeny predict microbiome structure in a

disease-challenged montane amphibian. *ISME J.* 1 (2018). doi:10.1038/s41396-018-0167-0

52.   RStudio Team. RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL

http://www.rstudio.c (2016).

53.   R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing. Vienna, Austria. URL https://www.R-project.org/. (2017).

54.   Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon data.

Nat. Methods **13**, 581–583 (2016).

55.   Kumar, S., Stecher, G., Suleski, M. & Hedges, S. B. TimeTree: A Resource for Timelines,

Timetrees, and Divergence Times. *Mol. Biol. Evol.* **34**, 1812–1819 (2017).

56.   Paradis, E. & Schliep, K. Ape 5.0: An environment for modern phylogenetics and evolutionary

analyses in R. *Bioinformatics* **35**, 526–528 (2019).

57.   Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. Y. Ggtree: an R Package for Visualization

and Annotation of Phylogenetic Trees With Their Covariates and Other Associated Data.

Methods Ecol. Evol. **8**, 28–36 (2017).

58.   Bürkner, P. C. Advanced Bayesian multilevel modeling with the R package brms. *R J.* **10**,

395–411 (2018).

59.   Bürkner, P. C. brms: An R package for Bayesian multilevel models using Stan. *J. Stat. Softw.*

80, (2017).

60.   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis.* (Springer-Verlag New York, 2009).

61.   Wilke, C. O. cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'. R package

556   version 1.0.0. https://CRAN.R-project.org/package=cowplot. (2019).

557   62.   Kay, M. tidybayes: Tidy Data and Geoms for Bayesian Models. R package version 2.0.3,

558   http://mjskay.github.io/tidybayes/. (2020).

559   63.   R Hackathon. phylobase: Base Package for Phylogenetic Structures and Comparative Data. R

560   package version 0.8.10. https://CRAN.R-project.org/package=phylobase (2020).

561   64.   Keck, F., Rimet, F., Bouchez, A. & Franc, A. phylosignal: an R package to measure, test, and

562   explore the phylogenetic signal. *Ecol. Evol.* **6**, 2774–2780 (2016).

563   65.   Lahti, L. & Shetty, S. Tools for microbiome analysis in R. Microbiome package version

564   1.1.10013. http://microbiome.github.com/microbiome. (2017).

565   66.   Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V. & Egozcue, J. J. Microbiome datasets are

566   compositional: And this is not optional. *Front. Microbiol.* **8**, 1–6 (2017).

567   67.   Oksanen, J. *et al.* vegan: Community Ecology Package. (2018).

568   68.   McMurdie, P. J. & Holmes, S. Phyloseq: An R Package for Reproducible Interactive Analysis

569   and Graphics of Microbiome Census Data. *PLoS One* **8**, e61217 (2013).

570   69.   Wilman, H. *et al.* EltonTraits 1.0 : Species-level foraging attributes of the world's birds and

571   mammals. *Ecology* **95**, 2027 (2014).

572   70.   Kurtz, Z. D. *et al.* Sparse and Compositionally Robust Inference of Microbial Ecological

573   Networks. *PLoS Comput. Biol.* **11**, 1–25 (2015).

574   71.   Williams, R. J., Howe, A. & Hofmockel, K. S. Demonstrating microbial co-occurrence pattern

575   analyses within and between ecosystems. *Front. Microbiol.* **5**, 1–10 (2014).

576   72.   Ding, B., Gentleman, R. & Carey, V. bioDist: Different distance measures. R package version

577   1.54.0. (2018).

578   73.   Csárdi, G. & Nepusz, T. The igraph software package for complex network research.

579   *InterJournal Complex Syst.* **1695**, 1695 (2006).

FIGURE 1

(A) Boxplots and raw data (points) of inverse Simpson indices for bacterial (green) and fungal

(orange) communities across a range of host species. **(B)** Posterior estimates of a binomial mixed

effects model investigating the influence of host taxonomy on the probability of a host's bacterial

diversity being higher than its fungal diversity. Mammalia and Insecta had posterior estimates

consistent with a >50% probability of an average animal sample having higher bacterial diversity

(credible intervals do not cross zero on the link (logit) scale, equivalent to 50% on the probability

scale; vertical dashed line). **(C)** Posterior estimates of a bivariate mixed effects model examining

variation in average Inverse Simpson index of bacteria (green) and fungi (orange) across animal taxa.

Estimates are in logs. Points are posterior means, and error bars show 66% and 95% credible

intervals.

FIGURE 2

(A) Phylogenetic tree of host species, with branches coloured by class and node points coloured by

order. Barplots show proportional composition of fungal and bacterial phyla for each host species,

aligned to tree tips. **(B)** Correlation of fungal and bacterial community structures (inter-sample

distance) derived from Procrustes rotation on principal component ordinations of each microbial

group. Microbial communities were aggregated at various taxonomic groupings (order, family, genus),

or as raw Amplicon Sequence Variant (ASV) taxonomy **(C)** Correlation between matrices of inter-

sample distance of microbial communities and host genetic distances generated from the

phylogenetic tree in A for both bacteria (green) and fungi (orange). As for B, microbial taxonomy was

either raw ASVs or grouped into family level. Aggregation to family resulted in higher correlations for

both microbial groups, and the correlation was always stronger in bacteria. For both B and C,

distributions of correlation values were generated using resampling of 90% of available samples for

that microbial group. Empty bars mean samples were not available for a particular species and so

would not have been included in the calculations in panel B.

FIGURE 3

Putative microbial interaction networks between bacterial (circles) and fungal (squares) taxa, coloured

by microbial phylum. Networks were constructed using the R package *SpiecEasi* on CLR-transformed

abundance values to detect non-random co-occurrence between groups of microbes.

FIGURE 4

Analysis of network structural traits of five metazoan classes comprising 39 species. There were

significant differences in **(A)** network modularity; and **(B)** proportion of positive edges (correlations

between paired microbial abundance values) among classes. Vertical dashed line indicates equal

proportion of positive and negative edges.

622 **TABLE 1:** Details of host species and their origins, sex ratios, sample sizes and types, and storage and extraction methods for the study.

Class	Common name	Latin name	N	Sex ratio (M: F: J: unknown: N/A)	Captive or Wild	Origin	Sample Type	Collection Year	Tissue Storage	Extraction Kit
Demospongia	Vase sponge	*Ircinia campana*	10	0: 0: 0: 0: 10	Wild	Long Key, Florida, USA	Tissue (choanosome)	2014	95% ethanol	Qiagen Blood and Tissue kit with proteinase K
Demospongia	Golfball sponge	*Tethya aurantium*	10	0: 0: 0: 0: 10	Wild	Long Key, Florida, USA	Tissue (choanosome)	2014	95% ethanol	Qiagen Blood and Tissue kit with proteinase K
Arachnida	Hard tick	*Amblyomma rotundatum*	10	0: 0: 0: 10: 0	Wild	Montserrat, Caribbean	Whole organism	2014	70% ethanol	Alkaline digest and ethanol precipitation
Malacostraca	Blue swimming crab	*Portunus segnis*	5	0: 0: 0: 5: 0	Wild	Malta	Gut	2018	70% ethanol	Quigen QIAamp Fast DNA Stool Mini kit
Malacostraca	Brown shrimp	*Crangon crangon*	10	0: 0: 0: 10: 0	Wild	Liverpool, Lancashire, England	Gut	2018	Buffer AE and frozen at -20°C	Qiagen Blood and Tissue kit with proteinase K
Insecta	Cockroach	*Diploptera punctata*	11	7: 1: 0: 3: 0	Captive	Manchester Metropolitan University, Manchester, UK	Gut	2018	Liquid nitrogen and frozen at -80°C	Qiagen Blood and Tissue kit with proteinase K and lysozyme
Insecta	Honey bee	*Apis mellifera*	10	0: 10: 0: 0: 0	Wild	North West of England, UK	Gut	2016	100% ethanol and frozen at -20°C	Qiagen Blood and Tissue kit with proteinase K and lysozyme
Insecta	Tsetse fly	*Glossina fuscipes*	9	2: 7: 0: 0: 0	Wild	Patira East, Uganda	Whole organism	2019	70% ethanol	Qiagen Blood and Tissue kit with proteinase K
Insecta	African palm weevil larvae	*Rhynchophorus phoenicis*	6	0: 0: 0: 6: 0	Wild	Sapele Town, Delta State, Nigeria	Gut	2019	Frozen at -20°C	ZymoBIOMICS DNA mini kit
Actinopterygii	European eel	*Anguilla anguilla*	10	0: 0: 0: 10: 0	Wild	Cumbria, England	Gut	2009	Frozen at -20°C	Qiagen PowerSoil kit
Actinopterygii	Foureye butterflyfish	*Chaetodon capistratus*	10	0: 0: 0: 10: 0	Wild	Bocas del Toro, Bahia Almirante, Panama	Gut	2018	95% ethanol	Qiagen PowerSoil kit with proteinase K
Actinopterygii	Yellowhead wrasse	*Halichoeres garnoti*	10	0: 0: 0: 10: 0	Wild	Caye Caulker, Belize	Gut	2015	95% ethanol	Qiagen PowerSoil kit with proteinase K
Actinopterygii	Barred hamlet	*Hypoplectrus puella*	12	0: 0: 0: 12: 0	Wild	Bocas del Toro, Bahia Almirante, Panama	Gut	2018	95% ethanol	Qiagen PowerSoil kit with proteinase K
Amphibia	Common midwife toad	*Alytes obstetricans*	11	0: 0: 0: 11: 0	Captive	London Zoo, London, UK	Skin swab	2015	Frozen at -20°C	Qiagen DNEasy kit
Amphibia	Phofung river frog	*Amietia hymenopus*	10	0: 0: 10: 0: 0	Wild	Drakensberg National Park, South Africa	Tadpole mouthparts	2015	95% ethanol	Qiagen Blood and Tissue kit with proteinase K
Amphibia	Common toad	*Bufo bufo*	10	0: 0: 10: 0: 0	Wild	Norway	Whole organism	2009	70% ethanol	Phenol chlorphorm
Amphibia	Great-crested newt	*Triturus cristatus*	10	5: 5: 0: 0: 0	Wild	Lancashire, England	Toe clip	2015	70% ethanol	Phenol chlorphorm
Aves	Reed warbler	*Acrocephalus scirpaceus*	8	3: 3: 0: 2: 0	Wild	Lincolnshire, UK	Faeces	2018/19	Frozen at -20°C	Qiagen PowerSoil Pro kit
Aves	Light-bellied brent goose	*Branta bernicla hrota*	10	0: 0: 0: 10: 0	Wild	Iceland	Faeces	2017	Frozen at -20°C	Qiagen PowerSoil kit

Aves	Goldfinch	*Carduelis carduelis*	8	5: 1: 0: 2: 0	Wild	Lincolnshire, UK	Faeces	2018/19	Frozen at -20°C	Qiagen PowerSoil Pro kit
Aves	Stock dove	*Columba oenas*	10	0: 0: 0: 10: 0	Wild	East Anglia, UK	Faeces	2014	Frozen at -20°C	Qiagen QIAamp Fast DNA Stool Mini kit
Aves	Woodpigeon	*Columba palumbus*	5	0: 0: 0: 5: 0	Wild	East Anglia, UK	Faeces	2012	Frozen at -20°C	Qiagen QIAamp Fast DNA Stool Mini kit
Aves	Carrion crow	*Corvus corone*	12	0: 0: 0: 12: 0	Wild	Cumbria, UK	Gut	2019	Frozen at -20°C	Qiagen Microbiome kit
Aves	Blue tit	*Cyanistes caeruleus*	8	0: 0: 0: 8: 0	Wild	Lincolnshire, UK	Faeces	2018	Frozen at -20°C	Qiagen PowerSoil Pro kit
Aves	Yellowhammer	*Emberiza citrinella*	8	1: 1: 0: 6: 0	Wild	Lincolnshire, UK	Faeces	2018	Frozen at -20°C	Qiagen PowerSoil Pro kit
Aves	Reed bunting	*Emberiza schoeniclus*	8	4: 3: 0: 1: 0	Wild	Lincolnshire, UK	Faeces	2018	Frozen at -20°C	Qiagen PowerSoil Pro kit
Aves	Robin	*Erithacus rubecula*	8	1: 1: 0: 6: 0	Wild	Lincolnshire, UK	Faeces	2018	Frozen at -20°C	Qiagen PowerSoil Pro kit
Aves	Great tit	*Parus major*	8	3: 3: 0: 2: 0	Wild	Lincolnshire, UK	Faeces	2018/19	Frozen at -20°C	Qiagen PowerSoil Pro kit
Aves	Chiffchaff	*Phylloscopus collybita*	8	0: 1: 0: 7: 0	Wild	Lincolnshire, UK	Faeces	2018/19	Frozen at -20°C	Qiagen PowerSoil Pro kit
Aves	Collared dove	*Streptopelia decaocto*	8	0: 0: 0: 8: 0	Wild	East Anglia, UK	Faeces	2014	Frozen at -20°C	Qiagen QIAamp Fast DNA Stool Mini kit
Aves	Turtle dove	*Streptopelia turtur*	7	0: 0: 0: 7: 0	Wild	East Anglia, UK	Faeces	2014	Frozen at -20°C	Quigen QIAamp Fast DNA Stool Mini kit
Aves	Blackcap	*Sylvia atricapilla*	8	3: 2: 0: 3: 0	Wild	Lincolnshire, UK	Faeces	2018	Frozen at -20°C	Qiagen PowerSoil Pro kit
Aves	Song thrush	*Turdus philomelos*	8	0: 0: 0: 8: 0	Wild	Lincolnshire, UK	Faeces	2018	Frozen at -20°C	Qiagen PowerSoil Pro kit
Mammalia	Striped field mouse	*Apodemus agrarius*	10	5: 5: 0: 0: 0	Wild	Chernobyl Exclusion Zone, Ukraine	Faeces	2017	100% ethanol and frozen at -20°C	Invitrogen Microbiome kit
Mammalia	Yellow-necked mouse	*Apodemus flavicollis*	10	5: 5: 0: 0: 0	Wild	Chernobyl Exclusion Zone, Ukraine	Faeces	2017	100% ethanol and frozen at -20°C	Invitrogen Microbiome kit
Mammalia	Wood mouse	*Apodemus sylvaticus*	10	6: 4: 0: 0: 0	Wild	Chernobyl Exclusion Zone, Ukraine	Faeces	2017	100% ethanol and frozen at -20°C	Invitrogen Microbiome kit
Mammalia	Northern muriqui	*Brachyteles hypoxanthus*	10	0: 0: 0: 10: 0	Wild	Caparao National Park, Espirito Santo, Brazil	Faeces	2017/18	RNA Later and frozen at -20°C	Qiagen QIAamp Fast DNA Stool Mini kit
Mammalia	Roe deer	*Capreolus capreolus*	7	7: 0: 0: 0: 0	Wild	Cumbria, UK	Faeces	2019	Frozen at -20°C	Qiagen Microbiome kit
Mammalia	Red deer	*Cervus elaphus*	10	0: 0: 0: 10: 0	Wild	County Meath, Ireland	Faeces	2018	Frozen at -20°C	Qiagen QIAamp Fast DNA Stool Mini kit
Mammalia	Greater white-toothed shrew	*Crocidura russula*	10	5: 5: 0: 0: 0	Wild	Belle Ile, France	Gut	2018	100% ethanol and frozen at -20°C	Qiagen PowerSoil kit
Mammalia	Eastern black rhino	*Diceros bicornis michaeli*	10	0: 10: 0: 0: 0	Captive	Chester Zoo and Port Lympne Wild Animal Park, UK	Faeces	2011	Frozen at -20°C	Qiagen QIAamp Fast DNA Stool Mini kit
Mammalia	Wild pony	*Equus ferus caballus*	10	5: 5: 0: 0: 0	Wild	Snowdonia National Park, Wales	Faeces	2013	Frozen at -20°C	Qiagen QIAamp Fast DNA Stool Mini kit
Mammalia	Hedgehog	*Erinaceus europaeus*	12	0: 0: 0: 12: 0	Wild	Cumbria, UK	Faeces	2019	Frozen at -20°C	Qiagen Microbiome kit

Mammalia	Bank vole	*Myodes glareolus*	10	7: 3: 0: 0: 0	Wild	Chernobyl, Ukraine	Gut	2017	100% ethanol and frozen at -20°C	Invitrogen Microbiome kit
Mammalia	Lesser horseshoe bat	*Rhinolophus hipposideros*	10	3: 5: 2: 0: 0	Wild	County Kerry, Ireland	Faeces	2016	Frozen at -20°C	Zymo DNA Extraction kit
Mammalia	Capuchin monkey	*Sapajus libidinosus*	10	0: 0: 0: 10: 0	Wild	Serra Talhada, State of Pernambuco/Minas Gerais, Brazil	Faeces	2017	Frozen at -20°C	Qiagen QIAamp Fast DNA Stool Mini kit
Mammalia	Grey squirrel	*Sciurus carolinensis*	12	0: 0: 0: 12: 0	Wild	Cumbria, UK	Faeces	2019	Frozen at -20°C	Qiagen Microbiome kit
Mammalia	Red squirrel	*Sciurus vulgaris*	12	0: 0: 0: 12: 0	Wild	Cumbria, UK	Faeces	2019	Frozen at -20°C	Qiagen Microbiome kit
Mammalia	Pygmy shrew	*Sorex minutus*	10	5: 5: 0: 0: 0	Wild	Belle Ile, France	Gut	2018	100% ethanol and frozen at -20°C	Qiagen PowerSoil kit

624
625

TABLE 2

PERMANOVA results for (a) fungi and (b) bacteria of factors explaining variation in microbial

community structure. Terms were added in the order shown in the table to marginalise effects of

sample storage and preparation protocols before calculating % variance explained for taxonomic

groupings. Species ID was the dominant source of variation in the data for both taxonomic groups, but

there were also strong effects of sample storage and wet lab protocol, particularly for bacteria.

(a) FUNGI				Taxonomic Effects Only		
Predictor	df	R2	p value	df	R2	p value
Sample Type	7	0.05	0.001			
Tissue Storage	5	0.04	0.001			
Extraction Kit	7	0.07	0.001			
Class	2	0.02	0.001	6	0.05	0.001
Order	6	0.05	0.001	13	0.12	0.001
Species	18	0.09	0.001	26	0.14	0.001
Residuals	303	0.68		303	0.68	

(b) BACTERIA				Taxonomic Effects Only		
Predictor	df	R2	p value	df	R2	p value
Sample Type	6	0.06	0.001			
Tissue Storage	6	0.16	0.001			
Extraction Kit	7	0.12	0.001			
Class	2	0.02	0.001	6	0.09	0.001
Order	6	0.09	0.001	12	0.21	0.001
Species	18	0.12	0.001	27	0.27	0.001
Residuals	273	0.42		273	0.42	

635