Adenoviral chromatin organization primes for early gene activation

Uwe Schwartz[a], Tetsuro Komatsu[b,c], Claudia Huber[a], Floriane Lagadec[c,d], Conradin

Baumgartl[a], Elisabeth Silberhorn[a], Margit Nuetzel[f], Fabienne Rayne[c], Eugenia Basyuk[c],

Edouard Bertrand[e], Michael Rehli[f,g], Harald Wodrich[c#] and Gernot Laengst[a#]

[a]Biochemie Zentrum Regensburg, University of Regensburg, Regensburg, Germany.

[b]Laboratory of Epigenetics and Metabolism, Institute for Molecular and Cellular Regulation,

Gunma University, Gunma, Japan

[c]CNRS UMR 5234, Microbiologie Fondamentale et Pathogénicité, Université de Bordeaux,

Bordeaux, France.

[d]Department of Molecular Biology, Faculty of Medicine, Göttingen Center of Biosciences

(GZMB), Georg-August-University Göttingen, Humboldtallee 23, 37073 Göttingen,

Germany.

[e]CNRS UMR 5355, Institut de Généthique Moléculaire de Montpellier, Montpellier, France

[f]Department of Internal Medicine III, University Hospital Regensburg, 93053, Regensburg,

Germany.

[g]Leibniz Institute for Immunotherapy, Regensburg, Germany, c/o University Hospital

Regensburg, 93053 Regensburg, Germany

[#]Address correspondence: harald.wodrich@u-bordeaux.fr or gernot.laengst@ur.de

Running title: Adenovirus chromatin structure

SUMMARY

Within the virion, adenovirus DNA associates with the virus-encoded, protamine-like structural protein pVII. Whether this association is organized, and how genome packaging changes during infection and subsequent transcriptional activation is currently unknown. Here, we combined RNA-seq, MNase-seq, ChIP-seq and single genome imaging during early adenovirus infection to unveil the structure- and time-resolved dynamics of viral chromatin changes as well as their correlation with gene transcription. Our MNase mapping data indicates that the viral genome is arranged in precisely positioned nucleoprotein particles (Adenosomes), like nucleosomes. We identified 238 Adenosomes, being positioned by a DNA sequence code and protecting about 60 to 70bp of DNA. The incoming genome is more accessible at early gene loci that undergo additional chromatin de-condensation upon infection. Histone H3.3 containing nucleosomes specifically replace pVII at distinct genomic sites and at the transcription start sites of early genes. Acetylation of H3.3 is predominant at the transcription start sites, preceding transcriptional activation. Based on our results we propose a central role for the viral pVII nucleoprotein-architecture, which is required for the dynamic structural changes during early infection, including the regulation of nucleosome assembly prior to transcription initiation. Our study thus may aid the rational development of recombinant adenoviral vectors exhibiting sustained expression in gene therapy.

INTRODUCTION

Most DNA viruses translocate their genomes into the nucleus of infected target cells to initiate viral gene expression and genome replication. During cytoplasmic trafficking, genomes are highly compacted inside the viral capsid to protect them from nucleic acid sensing molecules. It is generally assumed that genomes are decondensed upon entering the nucleus and thereby activated for viral gene expression and replication. Replicated genomes are then re-condensed for packaging into progeny. Alternatively, nuclear viral genomes could undergo different parallel fates, i.e. one part serving as transcription unit, whereas others are replicated and/or packaged (Brison et al., 1979). While these remarkable and reversible genome functions are highly specialized and efficient, little is known about the structural organization of the genome, its dynamic changes or the regulatory cues initiating structural transitions during virus entry.

Adenoviruses (Ad) are prototypic DNA viruses containing a linear double stranded DNA genome of 30-40 kb that is incorporated into a ~ 90 nm non-enveloped icosahedral capsid

shell. Ad enter cells by receptor-mediated endocytosis. Stepwise entry cues liberate the internal capsid protein VI and permit the virus to escape the endosomal compartment (Greber et al., 1993; Wickham et al., 1993; Wiethoff et al., 2005). Cytoplasmic capsids are transported to the nuclear pore complex (NPC) where they dock, disassemble and release the viral genome for nuclear import (Cassany et al., 2015; Strunze et al., 2011). Within hours after cell entry, nuclear genomes initiate gene expression from the immediate early promoter. In a simplified view this promoter encodes the E1A gene, encoding a *trans*-activator driving early genes responsible for cell and immune modulation (E1B, E3, E4) and replication (E2). In contrast activation of late genes (L1-L5), encoding most structural proteins, requires prior genome replication (Akusjarvi, 2008; Flint, 1982).

Packaged Ad genomes are devoid of histones. Instead they associate with basic core proteins VII (pVII), V (pV) and polypeptide μ (Brown et al., 1975; Corden et al., 1976). Protein VII is the most tightly bound and most abundant protein with over 500 copies per virion and responsible for packaging the viral genome into a structure appearing as irregular "beads-on-a-string" by electron microscopy (Benevento et al., 2014). Protein VII exhibits sequence similarities to protamines and may use its positive charges to neutralise and compact the viral DNA (Anderson et al., 1989; Johnson et al., 2004; Keller et al., 2002). Cellular factors and/or post-transcriptional modification control the association of pVII with viral genomes in the producer cell prior or during genome packaging. A recent study showed that viral genomes can be packaged without pVII, but that its presence is a prerequisite for infection (Avgousti et al., 2016; Genoveso et al., 2020; Inturi et al., 2017; Mun and Punga, 2018; Ostapchuk et al., 2017; Samad et al., 2012). The genomic binding sites of pVII as well as the structural organisation it imposes on viral DNA, inside the virion, post infection and post nuclear genome delivery, remain elusive (Giberson et al., 2011; Mirza and Weber, 1982). Protein VII stays associated with the viral genome imported to the nucleus, while pV dissociates from the viral DNA following ubiquitylation (Puntener et al., 2011). The fate of the μ-peptide is not known.

Cellular TAF-Iβ/SET associates with pVII on nuclear viral genomes and is thought to stimulate transcription and prevent the activation of DNA damage responses (Avgousti et al., 2017; Giberson et al., 2011; Haruki et al., 2003; Komatsu et al., 2015). Protein VII was also shown to functionally interact with the viral *trans*-activator E1A, pinpointing to a direct role in transcriptional activation (Johnson et al., 2004). Some studies suggested that pVII remains attached throughout the early phase of infection while others argue for a gradual and at least partial removal from the incoming genome (Haruki et al., 2003; Johnson et al., 2004;

Komatsu et al., 2011, 2015; Xue et al., 2005). Conflicting reports postulated that pVII

turnover either requires transcription (Chen et al., 2007) or occurs independent of

transcription, at least on viral vector genomes (Ross et al., 2011). In contrast chromatin

immune precipitation (ChIP) analysis proposed that pVII remains associated with viral

chromatin for several hours (Chatterjee et al., 1986; Haruki et al., 2003; Johnson et al., 2004;

Komatsu et al., 2011; Xue et al., 2005). In addition, cellular histones may associate with

incoming genomes prior to replication (Giberson et al., 2018; Ross et al., 2011).

Functional chromatin extraction using high and low concentrations of micrococcal nuclease

(MNase) to hydrolyze cellular chromatin, combined with high-throughput sequencing, is a

way to analyse protein binding to DNA and to reveal dynamic changes in chromatin structure

(Mueller et al., 2017; Schwartz et al., 2018). In this study we combined time-resolved MNase,

ChIP and RNA sequencing as well as single molecule imaging to correlate chromatin

structure changes with transcription at very early time points of adenoviral infection. This

approach revealed the specific nucleoprotein architecture of the virus, which dynamically

changes during early infection. Protein VII positioning is driven by sequence and is

specifically organized at early genes, correlating with the early de-condensation of these

regions upon infection. Defined pVII nucleoprotein complexes are replaced by nucleosomes

containing H3.3 on early gene promoters and few additional genomic sites. pVII remodeling,

nucleosome assembly and the specific acetylation of the promoter nucleosomes are preceding

transcriptional activation. The time resolved single molecule imaging of transcribing genomes

support our findings, identifying the morphological correlate at the single genome level at

subcellular resolution. The defined positioning of pVII on the viral genome and its functional

involvement in dynamic chromatin rearrangement during infection, suggest a key role in

regulating the viral life cycle. We identify a DNA sequence driven adenovirus nucleoprotein

architecture, which may aid the design of improved Ad vectors for gene-therapy or

vaccination.

RESULTS

Time resolved analysis of the early adenovirus transcriptome by RNA-sequencing

To investigate the early spatio-temporal reorganization of viral chromatin and transcription

activation we infected H1299, a p53-deficient non-small cell lung carcinoma cell line, or

U2OS, a human osteosarcoma cell line, with a partially E3-deleted but replication competent

human adenovirus C5 (HAd-C5). The inoculum was replaced with fresh medium after 30

min, defining time point zero of the experimental setup (Fig. 1A). On time point 0, 0.5, 1, 2,

and 4 hours post infection (hpi) infected cells were subjected to immune fluorescence analysis

(IF), RNA-sequencing (RNA-seq), ChIP-seq and functional chromatin extraction combined

with high-throughput sequencing of protected DNA fragments (MNase-seq) (Fig. 1A). To

verify the progression of infection, cells were first analysed by immunofluorescence (Fig.

S1A) for the internal capsid protein VI (pVI), which can be transiently detected during

endosomal escape (Martinez et al., 2015), while nuclear import is marked by accessibility of

the genome bound pVII in the nucleus (Komatsu et al., 2015). Protein VI and pVII

quantification showed that endosomal escape occurred within the first 30 min (Fig. S1A and

S1B) followed by rapid nuclear delivery of viral genomes starting at 0.5 hpi, with most

genomes being imported 1 hpi (Fig. S1A and S1C). The data is confirming a synchronized

infection covering pre- and post-nuclear genome delivery phases.

To investigate viral transcription, we performed mRNA sequencing of two independent

infection time-courses yielding on average 26 million reads per sample (Table S1). The reads

were mapped to an extended version of the human genome containing the HAd-C5 reference

genome and gene annotation. The first mature Ad transcripts were detected at 2 hpi, and

strongly increased levels were observed 4 hpi (Fig. 1B and 1C; whole genome profile shown

as individual replicates in Fig. S2 and quantification in Table S1). Consistent with the spatial

organisation of the Ad transcription program (Crisostomo et al., 2019; Glenn and Ricciardi,

1988), transcription is initiated at both ends of the Ad genome covering the early genes (Fig.

1B). We detected E1A as the first viral transcript (Nevins et al., 1979), along with significant

levels of E3 at 2 hpi (Fig. 1B and 1C). At 4 hpi viral transcription is markedly enhanced with

high levels of transcripts originating from all early genes and significant signals arising from

the E2A region. Taken together we observed an efficient utilization of the host cell

transcription machinery within the first hours of infection, revealing a transcriptional

competent viral template.

***In situ* detection of E1A mRNA identifies transcribing Ad genomes.**

To directly correlate the progression of infection and the onset of transcription, the import of

genome associated pVII and the expression of the E1A gene was monitored simultaneously

by immunofluorescence (Fig. 2A). We used RNAscope technology for the direct detection of

accumulating individual viral transcripts at the single cell level (Pheasant et al., 2018). A

specific set of probes for the E1A mRNA was designed (ACDBio) and validated on cells

infected with either HAd-C5 virus or an E1-deleted vector (Ad5-GFP). At 4 hpi specific E1A

transcripts could be readily detected in virus- but not in E1-deleted GFP expressing viral

vector infected cells (Fig. 2B). Protein VII immunofluorescence analysis showed that most

genomes were imported before 1 hpi with no significant additional import after this time point

(Fig. 2C). While the nuclear import of genomes was saturated 1 hpi, RNA-seq and RNAscope

revealed that E1A transcripts were detected only at 2 hpi and accumulated at 4 hpi (Fig. 2D

and Fig. 1C). Ad transcription initiation and mRNA synthesis exhibited a significant lag time

after nuclear entry, suggesting a requirement for remodelling processes prior to the assembly

of the transcription machinery.

E1A transcript accumulation could be related to the continued expression from a stable pool

of transcriptionally active genomes or by an increasing number of genomes initiating

transcription over time. We quantified the colocalization of nuclear genomes with nuclear

E1A transcripts to discriminate between both possibilities. Colocalizations inside the nucleus

resulting from association by chance are expected to increase over time when active genome

numbers remain constant and E1A nuclear transcript numbers increase over time. However,

our results showed that E1A positive nuclear genomes rapidly plateau at ~20% despite the

strong accumulation of nuclear E1A transcripts (Fig. 2E). It is shown that only a relatively

small subpopulation of imported genomes contributes to early viral gene expression until 4

hpi, which was also observed in a recent study (Suomalainen et al., 2020). These observed

~20% transcribing viral genomes could also be the result of a larger proportion of genomes

that undergo transcriptional bursting. This interpretation would not contradict the observed

results.

***In vivo* detection of E1A mRNA identifies a subpopulation of transcribing Ad genomes.**

The RNAscope analysis confirmed RNA-seq results and indicated, that active viral genomes

undergo a lag phase in the nucleus, probably acquiring structural changes that allow to

activate early gene expression. To confirm this observation and further characterize

transcribing and non-transcribing genomes we developed an imaging system for living cells.

To mark E1A transcripts in living cells we inserted 24x MS2 binding sites into the 3'UTR of

the E1A gene of a replication competent virus. MS2 binding sites are short stem-loop

structures derived from the MS2 bacteriophage and can be detected by binding of multiple

copies of fluorophore tagged MS2 binding protein (i.e. MS2BP-NLS-GFP) forming a

fluorescent spot *in situ* (Bertrand et al., 1998). Production and amplification of the virus

resulted in normal yields with unaltered infectivity or protein composition compared to the

parental virus (data not shown). To detect Ad genomes in living cells we used our previously

established live cell imaging system based on fluorescent TAF-Iβ, which associates with the

pVII of incoming genomes, thereby generating a fluorescent spot *in situ* (Komatsu et al., 2015). A stable mCherry-TAF-Iβ expressing cell line was transduced with an expression vector for MS2BP-NLS-GFP to generate double fluorescent stable cell lines. Cells were left either uninfected, or infected with untagged replicative control virus, or infected with E1A-MS2 tagged replicative virus and at ~ 4 hpi analysed by live cell imaging at high temporal resolution (1 frame/sec) (Movies S1-S4). To display the results in a single image we superimposed individual movie frames (60 sec t-projections, Fig. 3A). In non-infected control cells both, mCherry-TAF-Iβ and MS2BP-NLS-GFP localized to the nucleoplasm, with some unspecific enrichment of the MS2 signal in nucleoli (Fig. 3A top row, taken from Movie S1). When cells were infected with non-MS2 containing control virus, genomes formed mCherry-TAF-Iβ fluorescent spots in the nucleoplasm with limited confined motion as previously described (Komatsu et al., 2015), while the MS2 signal remained unchanged compared to non-infected cells (Fig. 3A 2[nd] row taken from Movie S2). In contrast cells infected with virus expressing MS2-repeat-tagged E1A mRNA formed mCherry-TAF-Iβ spots indicating viral genomes but also rapidly moving GFP spots visible inside the nucleoplasm and in the cytoplasm (Movie S3/S4). GFP spots accumulated in most cells around 3-5 hpi in agreement with the RNAscope experiment. Upon merging the mCherry-TAF-Iβ channel (viral genomes) with the MS2BP-NLS-GFP channel (viral E1AmRNA) using t-projections, some genomes were clearly double positive suggesting transcribing genomes (Fig. 3A 3[rd] and 4[th] row, taken from Movie S3 and S4). Similar observations were made when we used a different MS2BP construct (MS2BP-NES-GFP) predominantly localizing to the cytosol or when we lowered the input MOI resulting in fewer (<10) genome copies per imaging field (Fig. S3). To quantify the number of E1A mRNA positive genomes we performed kymograph analysis using dual-channel t-projections of individual cells. This approach was possible because the genome position was stable over time allowing to draw a line plot through all nuclear genomes using the signal of mCherry-TAF-Iβ. When plotted as kymograph over time each genome is represented as a red line. Genomes positive for the MS2BP-NLS-GFP are represented as a green line and scored as transcribing genomes (Fig. 3B). Kymograph analysis of several cells showed that ~20% of detected genomes inside a given nucleus are positive for MS2BP-NLS-GFP (Fig. 3C), which was very similar to the RNAscope approach (Fig. 3D). Taken together the analysis of transcriptionally active genomes in living cells supports our findings that early in infection transcription occurs from a subpopulation of imported genomes that probably require to actively remodel the viral chromatin for transcription onset.

228   Longer observations of individual nuclei (>20min) confirmed the existence of transcribing
229   and non-transcribing genomes with no indication of transcriptional alternation between
230   genomes (data not shown).
231
Chromatin architecture of the invading Ad genome
Transcripts accumulated with a delay of 1 hour after nuclear entry, opening the question of
whether structural changes may be required for assembly of the transcriptional machinery. To
address the Ad pVII nucleoprotein organisation, we partially digested viral and cellular
genomic DNA with the endo-nuclease MNase (Fig. 4A). This enzyme preferentially
hydrolyses DNA in the linker region between nucleosomes and is restricted by stable protein-
DNA complexes (Noll et al., 1975). Incubation of cellular chromatin with increasing
concentration of MNase resulted in the appearance of a typical MNase ladder, reflecting the
monomer and multimers of the nucleosomal DNA fragments (Fig. 4A). MNase
concentrations were titrated such that one sample exhibited less than 20% mono-nucleosomal
DNA (low digestion) and the other sample was fully hydrolysed to the mono-nucleosomal
DNA (high digestion; Fig. S4A). The latter is the common condition used for MNase-seq
experiments in order to study nucleosome positions (Diermeier et al., 2014; Schones et al.,
2008; Valouev et al., 2011), whereas approaches using limited MNase concentrations have
been used to analyse chromatin accessibility (Hu et al., 2019; Mieczkowski et al., 2016;
Schwartz et al., 2018). High and low digestion conditions were used at each infection time
point to isolate the mono-nucleosomal DNA band from two replicates. The nucleosomal DNA
of 17 individual experiments was used for library preparation and paired-end high-throughput
sequencing, yielding in total 3.8 billion paired-end reads (Table S2).
DNA fragments were aligned to the human and the Ad genome, first analysing the read length
distributions on both genomes (Fig. 4B). For the initial viral chromatin analysis we chose the
samples at time point 0 hpi. At this stage of infection, the genome has not been released from
the capsid (Fig. S1A) and therefore the MNase digestion pattern displays the nucleoprotein
organization of the virus particles. When comparing the virus and host genomes digested at
the high MNase concentration, we observed a clear difference in the read-length distribution
on the two genomes (Fig. 4B). Whereas most DNA fragments exhibited the typical
nucleosomal DNA size on the human genome (147 bp), the protected DNA fragments on the
Ad genome were significantly shorter with a mean size of 60 to 70 bp. The MNase-seq
approach revealed a stable adenoviral packaging unit that most probably represents pVII
bound to DNA, which is the only abundant genome binding protein present prior to genome

release from the capsid. Previous studies showed that nuclease digestion of Ad nucleoprotein
resulted in discrete populations of protected DNA fragments, suggesting a chromatin-like
configuration with sizes varying between 180 bp and 200 bp, between 30 and 90 bp, or
between 30 and 60 bp depending on the conditions used (Corden et al., 1976; Daniell et al.,
1981; Johnson et al., 2004). Assuming that the larger DNA fragments represent partial
nuclease hydrolysis products, the 60 to 70 bp long Ad-DNA fragments mirror the previously
described pVII-DNA protein complexes. Detection of these complexes (referred to as pVII
peaks) allows us for the first time to analyse the occupancy and positioning of Ad pVII. We
identified defined pVII peaks over the Ad genome, showing specific pVII-DNA positions on
the viral DNA (Fig. 4D). The MNase profiles and peak calling were highly reproducible
between biological replicates at individual time points, suggesting that specific Ad genome
positions are occupied with pVII, equivalent to nucleosomes on the host genome (Fig. S4C-
E). The Ad exhibits a defined nucleoprotein architecture that differs along its functional
regions. Interestingly, a central region of the viral genome (Late3) and a region between the
E3 and E4 genes exhibited almost no peaks (Fig. 4D).
Regions lacking pVII peaks could result either from the absence of pVII protecting the DNA
from continuous MNase degradation or from tight compaction excluding DNA from MNase
digestion and therefore resulting in DNA fragments of higher molecular weight. To test both
possibilities we digested Ad chromatin directly within virus particles or protein-free viral
DNA (gDNA, to control for MNase cleavage preferences) with a very low amount (5U) of
MNase (Fig. S5A and S5B). While cleavage patterns were uniform in Ad gDNA, the highest
cleavage frequencies in virus particles corresponded to the regions lacking pVII signals after
infection (Late3 and between E3 and E4), suggesting the presence of "unprotected" domains
in the Ad genome, which are completely hydrolyzed under the MNase digestion conditions
used in infected cells.
287
pVII exhibits DNA sequence dependent positioning patterns
We next asked whether intrinsic DNA sequence patterns may establish pVII peaks. Such a
sequence dependent positioning code was shown for nucleosomes (Kaplan et al., 2009).
Sequence specific bending of DNA reduces the energy requirements for wrapping the DNA
around the histone octamer surface, revealed by characteristic di-nucleotide repeats in the
nucleosomal DNA sequence at 10 bp intervals (Widom, 2001). This pattern was clearly
observed in the human host genome (Fig. 4C – upper panel).

Interestingly, we also observed a specific di-nucleotide pattern when computing the di-
nucleotide frequency of the mid-point centered pVII peak fragments, suggesting that a
specific DNA code underlies pVII positioning on the Ad genome (Fig. 4C – lower panel).
However, to our surprise, a periodic pattern of 5.4 bp was obtained for WW- (where W is A
or T) and SS-dinucleotides (where S is G or C), much shorter than the nucleosomal DNA
repeat pattern, implying a distinct packaging mode. Such a repeat pattern indicates a spring-
like winding of the DNA around pVII. Even without knowledge of the exact DNA-pVII
topology, the sequence pattern and the specific binding sites of pVII on DNA clearly suggest
the organisation of the Ad genome into a specific nucleoprotein-structure that may be
essential for its function.
On average, we detected 238 pVII peaks covering the Ad genome, which was consistent
throughout all timepoints (Fig. 4E). Next, we quantified the mean pVII-DNA particle repeat
length (pVII-RL), the distances between individual pVII peaks, by measuring the center-
center distance of adjacent pVII peaks in each sample (Fig. 4F and S4B). The median pVII-
RL measures 97.3 bp, suggesting that we isolated defined pVII-DNA particles spaced by
accessible DNA linkers of 20 to 40 bp in length.
To address the presence of functionally distinct regions, we systematically subdivided the Ad
genome into 10 domains of similar size, named after the included gene and classified by the
timing of expression (color code: early to late = light to dark blue) (Fig. 4D). Plotting the
pVII-RL showed a clear correlation of the repeat length with expression timing (Fig. 4G).
Increased repeat lengths in the regions of the early genes may serve to increase genome
accessibility and to enable the binding of regulatory factors. Strengthening this finding, the
pVII occupancy and stability, reflected by the peak height, showed an anti-correlation with
early timing in gene expression (Fig. 4H). In general, less pVII peaks (Fig. 4I) with larger
repeat lengths and increased instability were found at early expressed domains (Fig S4D-E)
indicating an open chromatin conformation, which could be confirmed with limited MNase
digestions of virus particles (Fig. S5D).
Our high-resolution analysis of Ad genome packaging provides, for the first time, information
on pVII-DNA peak positioning and occupancy. The analysis revealed a defined viral
nucleoprotein architecture, with an open configuration at early expressed genes potentially
enabling immediate access of the host cell transcription machinery.
326
Dynamic changes of viral genome accessibility over time

One parameter determining viral genome accessibility and activity is the specific occupation
with pVII, another parameter is the higher order structure of the Ad-chromatin and its global
accessibility. Assessing the nucleoprotein structure with high MNase concentrations is
providing information about stable protein particles occupying the DNA (Schwartz et al.,
2018). In contrast, limited MNase concentrations, in analogy to the analysis of regulatory sites
by DNase I hypersensitivity mapping (Längst et al., 1997), can reveal accessible genomic
regions and regulatory elements (Schwartz et al., 2018).

Only a fraction of about 20% of the viral genomes in the nucleus are transcriptionally active
(Fig. 2E and 3C), explaining why we observed only little variation between the different time
points treated with high MNase concentrations (Fig. 5A). In contrast, at low MNase
concentration (Fig. 5B and Fig. 5C) clear changes over infection time are detected, signifying
gross structural changes of the viral chromatin during early infection. We hypothesize, that
the active viruses undergo de-compaction that can be detected by limited MNase hydrolysis.
When monitoring the pVII peak positioning at 0 hpi, the low MNase signal was well
correlated with the high MNase signal (R = 0.91), essentially showing the same pVII peak
positions in both conditions (Fig. S6A, B). However, when inspecting the viral chromatin
maps at later time points, we detected specific structural changes, with changes in peak
intensities and the appearance of novel peaks in the low MNase signal (Fig. 5C). To identify
structural changes over time we used a linear regression approach. The average signal
intensity at each pVII position was quantified at each time point and fitted by linear
regression. The slope of the regression line was used as a score for chromatin accessibility
changes (accessibility score; Fig. 5C). Genomic regions with a positive accessibility score are
opening up, being more MNase accessible, whereas negative values correlate with DNA
remaining condensed. A similar strategy was used in previous studies to measure the local
chromatin accessibility (Chereji et al., 2019; Mieczkowski et al., 2016; Mueller et al., 2017).

The increase of Ad chromatin accessibility is correlating with viral transcriptional activity,
with Ad chromatin remodeling and de-condensation at the early gene regions (Fig. 5D).
Interestingly, chromatin opening at E1A is not as pronounced as at other early genes. As this
is the first gene to be transcribed, we suggest that this region already exists in an open
chromatin configuration, correlating with the pre-existing low pVII peak density, and
therefore no additional opening is observed over time.

The Late3 domain also exhibited a positive accessibility score, suggesting Ad chromatin de-
condensation at this site (Fig. 5C). Additionally, in this domain we observe a discrete peak
appearing at low MNase concentrations at 4 hpi (Fig. 5C shaded region and peak marked by

asterik; Fig. S6C). The occurrence of this local nuclease hypersensitive site suggests the
binding of regulatory factors (Chereji et al., 2017; Kent et al., 2011; Mieczkowski et al.,
2016). Transcription factor motif analysis revealed potential binding sites for EN1, CREM,
C/EBP-alpha, and C/EBP-beta in the center of the peak (Fig. S6C). Interestingly, the
transcripts of EN1, CREM, and C/EBP-beta in the host cell were upregulated up to twofold
upon viral infection (Fig. S6D).

In summary, the time-resolved chromatin accessibility analysis shows, that in addition to the
predefined pVII binding sites on the genome, Ad chromatin compaction specifically changes
over time. Chromatin opening of the early gene regions starts at 0.5 hpi, prior to
transcriptional activation. We suggest that changes in early gene regions are a pre-requisite
for gene activation process.

374 Dynamic changes of Ad genome packaging

Previous studies showed that histones are deposited onto the Ad genome (Giberson et al.,
2018; Komatsu and Nagata, 2012; Komatsu et al., 2011). However, at which position and
whether these are functional nucleosomes is not known.

Analysing the MNase-seq data of the time points 1, 2 and 4 hpi revealed the appearance of
147 bp long, protected DNA fragments at 2 and 4 hpi, after nuclear import of the viruses and
prior to early gene activation (Fig. 6A). The nucleosome sized peaks increased in intensity,
within the background of the more abundant pVII protected DNA fragments, suggesting co-
occupation of the viral genome by pVII and nucleosomes. To quantify the fraction of
nucleosomes on the viral genome, we simulated the peak sizes by computationally adding
nucleosomal reads to the pVII fragment data of time point 1 hpi (Fig. S7A). The analysis
suggested that at 2 hpi 8% and at 4 hpi 14% of the viral genomes are occupied with
nucleosomes.

To verify this indirect prediction of nucleosomal occupancy on the Ad genome, we generated
U2OS cell lines, stably expressing FLAG-tagged histone H3.1 (U2OS31) and the histone
variant H3.3 (U2OS33) (Fig. S7B, C). Stable cell lines were infected with Ad and ChIP-seq
experiments were performed at 0, 1, 2, and 4 hpi, using either FLAG- or H3K27ac-antibody.

At 0 hpi we barely detected any ChIP-seq reads mapping to the Ad genome, showing the high
specificity of our ChIP-seq approach (Fig. 6B). The number of precipitated Ad DNA
fragments increased with infection time for histone H3.3 (FLAG IP: U2OS33), in contrast to
the canonical histone H3.1 DNA fragments (FLAG IP: U2O31) (Fig. 6B). The low number of
histone H3.1 DNA fragments clustered with the samples taken at 0 hpi on PCA and

hierarchical clustering analysis (Fig. 6C, Fig. S7D), and did not show any specific sites of

enrichment on the Ad genome (Fig. S7E, upper panel), suggesting no specific recruitment of

histone H3.1. However, we detected specific H3.3 binding sites on the Ad genome, clearly

showing that this histone variant is exclusively deposited onto incoming viral genomes (Fig.

6D). The H3.3 ChIP-Signal was most pronounced at early transcribed gene promoters,

including E1A, E3, and E4, but not E1B. Nevertheless, H3.3 also accumulated at additional

loci throughout the Ad genome, like the E2B and Late2 regions (Fig 6D, E).

Next, we performed H3K27ac ChIP-seq experiments, showing efficient histone acetylation,

without any detectable delay after histone deposition on the Ad genome. Interestingly, we

observed histone acetylation at specific sites on viral genomes, representing a subset of all

H3.3 binding sites, marking these as active regions (Fig. 6B, D). As the H3K27ac antibody

does not discriminate between H3.1 and H3.3, the results obtained in U2OS31 and U2OS33

are very similar (Fig. 6B, C and Fig. S7D). The U2OS33 cell line exhibited only slightly

higher H3K27ac levels due to additional expression and incorporation of histone H3.3.

Histone acetylation was identified at all early gene promoters, except E1B, with the strongest

signal at E1A, which was already detectable 1 hpi (Fig. 6D, F). Additionally, we observed

pronounced K27 acetylation at two central regions covering the E2B or Late2 gene.

We expected that sites marked by H3K27ac would promote chromatin de-condensation.

However, neither H3.3 deposition nor H3K27 acetylation did correlate with chromatin

accessibility changes ($R^2 < 0.1$, Fig. S8A, B). A closer inspection showed that acetylation at

the E1A and E4 genes were associated with open chromatin (Fig. S8C, F), whereas the

acetylated Late2 and E2B sites in the genome centre remain condensed (Fig. S8D, E).

In summary, we suggest that the histone variant H3.3 is preferentially deposited onto viral

genomes and a subset of binding sites get acetylated at K27. H3.3 acetylation is detected

preferentially at early gene promoters, being nucleosomal and acetylated prior to transcription

initiation.

Early genes gain a positioned +1 nucleosome

The sonicated DNA in the ChIP experiments is not short enough to reveal the exact sites of

nucleosome assembly. Therefore, we combined the ChIP-seq data with the MNase-seq

derived nucleosomal DNA fragments (size between 137-157 bp; time point 4 hpi), allowing

us to reveal the exact positions of the nucleosome core particle. A cut-off of 2 CPM was used

to define the genomic sites of preferential nucleosome assembly from U2OS2 FLAG IPs (Fig.

S8G, blue boxes) and the exact nucleosome positions were determined using MNase size

selected fragments (Fig. S8G, golden boxes). The nucleosome profile was distinct from the

pVII landscape and nucleosome calling was reproducible between replicates.

Nucleosome assembly occurs in the genomic regions associated with the early genes,

exhibiting a lower pVII peak occupancy and increased pVII peak repeat-length. An analysis

of the incoming nucleosome positions with respect to pVII peak occupancy revealed that

nucleosomes directly formed over the central positions of the pVII-nucleoprotein particles

(Fig. S8H) suggesting a specific mechanism to replace pVII.

A detailed analysis of the nucleosome assembly sites with respect to the early genes revealed

that nucleosomes occupy discrete positions. Remarkably nucleosome positions coincide with

the 5'-end of the synthesized E1A, E4, E3, and E2A RNA, correlating with the highly

positioned +1 nucleosome found in active human genes (Fig. 6G, H and Fig. S8I) (Schones et

al., 2008). Additional nucleosomes are detected in the coding region of the early genes. The

+1 nucleosome marks the transcription start sites of genes and flanks the regulatory sequences

of the promoters, residing in the upstream DNA linker region.

The time-resolved chromatin analysis shows, that the predefined pVII chromatin organisation

is remodeled, exchanging pVII for histone H3.3. Histone H3.3 deposition and acetylation

occur mainly at the +1 region of early genes and happens prior to transcription onset. Our

results imply the requirement of specific nucleosome assembly for transcription initiation at

early genes.

DISCUSSION

The adenovirus exhibits a defined nucleoprotein architecture

Early studies suggested that pVII binds adenoviral DNA at a 1:1 ratio like histones and

protects DNA from endonuclease cleavage (Lischwe and Sung, 1977; Sato and Hosokawa,

1984). However, no defined pVII-DNA particle has been identified on functional Ad genomes

yet, as the protected DNA had a variable size range of 30 to 250 bp depending on

experimental conditions and the degree of endonuclease digestion (Corden et al., 1976;

Daniell et al., 1981; Mirza and Weber, 1982). In addition, it is still not known whether pVII

binds to specific sites on the viral genome, creating a defined viral nucleoprotein architecture,

like the cellular nucleosomes. Here, our MNase-seq approach digesting Ad together with the

host chromatin at different time points post infection, revealed the 147 bp long DNA

fragments of the host nucleosomes and identified a shorter, protected DNA species with a

mean size of about 60 to 70 bp originating from the viral sequences. With pVII being the

major DNA binding protein of the adenovirus, and the protected fragment species being

present prior to genome release from the capsid, the observed DNA fragments are very likely

protected by their association with pVII. The viral MNase digestion pattern is not as regular

as the nucleosomal DNA fragments, suggesting heterogeneity in pVII-DNA complexes, as

observed in the early studies. Such a digestion pattern could also be the result of a discrete

pVII core complex (60 bp) that tightly associates with neighbouring pVII complexes, partially

protecting the interconnecting linker DNA from endonuclease cleavage. Such structures were

indeed visible in early electron microscopic studies (Mirza and Weber, 1982).

pVII organizes the viral DNA into a specific nucleoprotein structure

Annotation of the putative pVII protected DNA fragments to the viral genome revealed for

the first time a specific positioning of pVII peaks on the genome. We show that the Ad

genome is covered by an average of 238 defined pVII peaks located at defined positions that

are spaced by DNA linkers of about 20 to 40 bp. Precise pVII peak positions suggest a highly

organized and defined nucleoprotein architecture shared by all virus particles. In host cells,

DNA structure determines histone binding preferences, as shown by the DNA sequence

dependent nucleosome positioning code (Kaplan et al., 2009). DNA bending allows wrapping

of the DNA around a protein core, requiring site specific SS and WW di-nucleotide

enrichments. SS and WW repeats differ in the wideness of the DNA major/minor groove

thereby bending the DNA (Kaplan et al., 2009). If the sequence dependent DNA shape

follows the protein surface path it increases binding affinity and thereby determines binding

site preferences (Widom, 2001). Whereas the nucleosomal di-nucleotide repeats are spaced by

10.3 bp, pVII exhibits a much shorter repeat length of 5.4 bp. Nucleosomes and pVII-DNA

complexes are organized differently, and we suggest that the DNA associates in form of a

spiral around the pVII core (Fig. 4C).

Our results suggest that pVII folds Ad genomes into an organized higher order structure,

driven by specific binding sites pre-determined by the Ad DNA sequence. A recent report

showed that pVII is not required for genome packaging into virions, albeit the lack of pVII

resulted in non-infectious Ad (Ostapchuk et al., 2017). The specific organization of the viral

genome with pVII may be functionally linked with endosomal escape, in addition to potential

gene regulatory effects. However, as endosomal escape of the Ad lacking pVII is blocked and

genomes are not delivered to the nucleus, the effect of pVII absence on gene expression and

the specific exchange of positioned pVII peaks by nucleosomes cannot be concluded. Our

observation that pVII peaks are reproducible across biological replicates strongly indicates a

specific positioning code. Protein VII peak densities and the changes in pVII occupancy upon

nuclear delivery (see below) correlate with the onset of the transcriptional program and

cellular factors, such as TAF-Iβ. TAF-Iβ was shown to positively influence viral

transcription, being recruited to the incoming genomes via pVII (Haruki et al., 2003;

Matsumoto et al., 1995). One possible approach to validate our findings of an underlying

sequence code for pVII positioning may be the rational (sequence-)design of therapeutic

DNAs packed into adenoviral vectors to achieve high and sustained transgene expression. *In*

vitro assembly of pVII-DNA complexes showed that pVII binds to DNA regardless of the

DNA sequence (Avgousti et al., 2016; Komatsu et al., 2011). However, it was not addressed,

whether sequence motifs are preferred by pVII, like it was observed in nucleosome

assemblies. Specific assembly factors and ATP dependent remodeling enzymes may be

required to adjust the precise positioning of pVII on DNA, in addition to the DNA assembly

code. Like nucleosomes, pVII does not bind as isolated protein to DNA but is forming

homotypic complexes. The number of pVII proteins per Ad genome was estimated to be

about 500 copies (Benevento et al., 2014) and early biochemical studies suggested the

occurrence of pVII dimers (Chatterjee et al., 1985; Everitt et al., 1975). Taking into account

the median number of 238 pVII positions we identified on the Ad genome (plus 19 on average

for the 1871 bp deleted E3 region) our data would be compatible with two pVII molecules

forming dimers at 238 discrete sites. This number would also come close to the ~200 "beads-

on-string" estimated for purified viral cores (Vayda et al., 1983).

Viral chromatin dynamics drives early gene expression

Here we show that nuclear delivery of viral genomes takes about 30 min but requires an

additional hour before viral transcripts appear. In agreement with earlier observations, full

activation of immediate early gene expression is reached after 4 h (reviewed in (Pied and

Wodrich, 2019)). Correlating gene expression with our differential MNase digestion

conditions, we show that early genes exhibit a reduced density of MNase protected regions

that most probably represent pVII peaks, and increased pVII peak repeat lengths. Direct

detection of genomes and of the E1A mRNA showed that genomes imported to the nucleus

require about two hours to activate gene expression and to synthesize the mRNA (i.e. E1A

transcription). In this lag phase pVII is selectively replaced by nucleosomes at early genes,

preceding gene activation. ChIP-seq experiments revealed that the histone variant H3.3 is

incorporated at many specific genomic locations on the Ad genome, including the early gene

regions, shortly after nuclear import. A subset of these nucleosomal sites, corresponding

mainly to the early gene positions, are acetylated at H3K27 prior to transcriptional activation.

We identified that 80% of the imported Ad genomes are transcriptionally inactive, but the

remaining 20% do actively rearrange the chromatin structure to allow transcription factor and

RNA Polymerase binding. Once entering the nucleus, we showed that the chromatin structure

of active viral genomes decondensed and viral DNA becomes more accessible. A delay for

E1A gene expression and heterogeneity of transcriptional activity for imported genomes was

recently reported supporting our findings (Suomalainen et al., 2020). The limited activation of

incoming genomes and subsequent de-compaction was also observed during herpes simplex

virus type 1 (HSV-1) infections showing common features of nuclear replicating DNA viruses

(Hu et al., 2019; Sekine et al., 2017). However, in the case of HSV-1 gene activation, it was

suggested that the accessibility over the complete viral genome increased (Hu et al., 2019),

whereas active Ad genomes selectively de-condense the early gene regions and a specific

domain in the Late3 gene. The overlap of open viral chromatin with histone H3K27

acetylation and early transcription, suggests an organization that allows the binding of

regulatory factors for gene activation. In contrast, for the Late3 region no functionality in

gene expression control has been assigned yet. Our observation that increased DNA

accessibility overlaps with binding motifs for transcriptional regulators may indicate a role as

a transcriptional enhancer.

pVII is exchanged for nucleosomes at early genes

Genome de-compaction correlates with low pVII occupancy and coincided with a site-

specific exchange of pVII protein for histone octamers. The characteristic nucleosomal

fragment length of 147 bp, observed in MNase-seq, confirms the release of pVII from the

viral genome, as pVII-nucleosome complexes would result in longer fragments after MNase

digestion as previously shown (Avgousti et al., 2016). The ChIP-seq experiments clearly

show an overlap of the 147 bp long protected DNA fragments and the location of the histone

variant H3.3. Furthermore, the newly assembled nucleosome is located on a previous position

of a pVII-DNA complex. Once pVII is set free, it is able to sequester HMGB1 and modulate

the cellular chromatin structure, thereby downregulating the immune response of the host cell

(Avgousti et al., 2016). Protein VII molecules were exchanged for positioned nucleosomes at

the +1 site of actively transcribed genes (i.e. the E1A and E4 gene, Fig. 6G and 6H). The

calculated and experimentally validated number of nucleosomes and their positions on DNA,

as determined by the 147bp long DNA fragments and the ChIP-seq experiments, corresponds

to the number of ~ 20% transcribing genomes, exhibiting single nucleosomes positioned at

the +1 position of early genes. As nucleosome exchange is commencing at 0.5 hpi and is

completed at 2 hpi when transcription activation is detectable by mature transcript accumulation, we suggest that nucleosome positioning is a prerequisite for gene activation. At this point we can only speculate why only a subpopulation of genomes is transcriptionally active. Reasons could be that the few genomes undergoing nucleosome assembly and active transcription produce the replication enzymes, whereas the bulk of genomes enters replication without activation as an elegant way to avoid repeated chromatinization. Alternatively, cellular defense mechanisms and/or the cell cycle could influence transcription levels (Suomalainen et al., 2020).

Interestingly, pVII removal and nucleosome placement at early genes preceded transcription, suggesting that pVII removal is driving transcription and not vice versa, as previously reported (Chen et al., 2007). This observation is in agreement with recent pVII-ChIP experiments showing transcription and replication independent pVII removal in early infection (Giberson et al., 2018; Komatsu and Nagata, 2012; Komatsu et al., 2011).

Future work has to reveal which machinery drives the pVII nucleosome exchange and determines specificity. Generally, active promoters are free of nucleosomes (nucleosome depleted region, NDR) and the transcription start site is typically located upstream of a well-positioned '+1' nucleosome influencing recruitment of the transcriptional machinery and initiation of transcription (Jiang and Pugh, 2009; Kubik et al., 2015). In the case of the Ad genome, our data show that the E1A promoter region is already pVII free and exists in an accessible nucleoprotein conformation when entering the cell, potentially allowing fast assembly of the +1 nucleosome and enabling immediate gene activation. This organization might thus be defined in the producer cell during genome packaging, which occurs in a sequential packaging sequence driven manner (Hammarskjöld and Winberg, 1980; Hearing et al., 1987). In contrast, other early regions were actively de-compacted with nuclear entry of the virus, correlating with slightly delayed transcriptional onset. This effect may be driven by pVII interaction with transcription promoting factors like TAF-Iβ and E1A (Haruki et al., 2006; Johnson et al., 2004). Besides pVII is post-translationally poly(ADP)-ribosylated or acetylated in infected cells, which might further promote chromatin relaxation (Avgousti et al., 2016; Déry et al., 1986).

Taken together our analysis provides unprecedented insight into the adenoviral nucleoprotein structure and its dynamic reorganization during the onset of viral gene expression. We reveal genome sequence, pVII-DNA occupancy and nucleosome positioning as driving forces for transcriptional activation. Reproducing such organization in adenoviral vectors could result in efficient and sustained transgene expression.

EXPERIMENTAL PROCEDURES

RNA library preparation and sequencing

Total cellular RNA was isolated from infected H1299 cells at 0, 0.5, 1, 2, 4 hpi using TRI Reagent (Molecular Research Center, Inc.). Two time series sets of total RNA were prepared from two independent infections. Library preparation, comprising rRNA depletion, polyA enrichment, fragmentation to ~270 nucleotide length, reverse transcription to cDNA using random hexamer primers and adapter ligation, was accomplished using TruSeq RNA Sample Preparation Kit v2 (Illumina) according to the manufacturers protocol by EMBL GeneCore facility in Heidelberg. Libraries were sequenced on Illumina HiSeq2000 platform resulting in 18-75 Mio 50 bp single-end reads per sample.

Differential MNase-seq, library preparation and sequencing

H1299 cells infected with Ad were permeabilized with the detergent IGEPAL-CA630 and digested at 0, 0.5, 1, 2 and 4 hpi with 100U/600U of MNase for 4/5 min at 37°, denoted as 'low'/'high' MNase. Mnase treatment was performed as described in (Schwartz et al., 2018). After fragmented DNA has been separated on 1.3% agarose gels, mono-nucleosomal fractions were isolated. Paired-end sequencing on Illumina HiSeq2000 platform (EMBL GeneCore facility, Heidelberg), yielding about 220 million read pairs of 50 bp length per sample, was accomplished following library preparation using NEBNext DNA library prep Master Mix Kit (New England Bioloabs)

ChIP-seq, library preparation and sequencing

Chromatin immunoprecipitation (ChIP) was performed as described previously with slight modifications (Minderjahn et al., 2020). Chromatin of cell lines encoding Flag-tagged H3.1 or H3.3 (U2OS31 and U2OS33, respectively) was harvested at 0, 0.5, 1, 2 and 4 hpi in triplicates of independent Ad infections. Briefly, for anti-H3K27ac and anti-Flag ChIP-seq, cells were crosslinked with 1% formaldehyde for 10 min at room temperature and the reaction was quenched with glycine at a final concentration of 0.125 M. Chromatin was sheared using ultrasonication (Covaris S2). A total of 2.5 μg of antibody against H3K27ac (Abcam, ab4729), or FLAG (M2, Sigma Aldrich, F3165), was bound to 20 μl pre-washed Dynabeads Protein A (Thermo Fisher Scientific) in a total volume of 200 μl PBS containing 0.02% Tween-20 for 1h at room temperature rotating at 6 rpm. The buffer was removed on a magnet and antibody-coupled beads were resuspended in 120μl dilution buffer (20mM Tris pH7.4, 2mM EDTA, 100mM NaCl, 0.5% Triton x-100) containing proteinase inhibitors (cOmplete™, EDTA-free Protease Inhibitor Cocktail, 50x, Sigma). 80μl sonicated chromatin

was added to the antibody-coupled Dynabeads (sonicated chromatin of approx. 1.7×10^6 cells)

and incubated for 3h at room temperature rotating at 6 rpm. Beads were washed on a magnet

and chromatin was eluted. After crosslink reversal, RNase A and proteinase K treatment,

DNA was extracted with the Monarch PCR & DNA Cleanup kit (NEB). Sequencing libraries

were prepared with the NEBNext Ultra II DNA Library Prep Kit for Illumina (NEB)

according to manufacturer's instructions. The quality of dsDNA libraries was analyzed using

the High Sensitivity D1000 ScreenTape Kit (Agilent) and concentrations were assessed with

the Qubit dsDNA HS Kit (Thermo Fisher Scientific). Libraries were sequenced paired end on

a NextSeq550 (Illumina).

Viruses and virus production

Bacterial artificial chromosome (BAC) mutagenesis was used to generate recombinant

adenoviruses following the protocol described in (Ruzsics et al., 2014) and (Martinez et al.,

2015). Briefly, bp 1-3513 of the HAdV-C5 genome including the left inverted terminal repeat

(ITR) and the whole E1 region was replaced by an FRT (Flippase Recognition Target) site

and maintained as BAC (termed B12) to perform lambda red recombination essentially as

described. Plasmid pO6-Ad5-E1-FRT (encoding the left ITR and the complete E1 region

followed by an FRT recombination site) was used to clone 24 copies of the MS2 binding site

repeats into the 3'UTR of the E1A gene. Subsequently flp-recombination was used (Ruzsics

et al., 2014) to recombine plasmids pO6-Ad5-E1-FRT-MS2 and BAC B12 to reconstitute

BxAd5-E1-MS2 a BAC encoding the whole AdV genome including the E1A gene tagged

with 24 copies of the MS2 binding site. To generate the virus the BxAd5-E1-MS2 BAC was

digested by *PacI* restriction enzyme to release the viral genome and transfected into HEK293

cells to reconstitute the recombinant viruses. Two rounds of plaque purification were

performed followed by large scale production in HEK293 cells and purification through

double $CsCl_2$ gradients banding. Replicative E3-deleted HAd-C5 virus or E1/E3-deleted

HAd-C5 GFP expressing control virus were essentially produced in the same way. Purified

viruses were extensively dialyzed against PBS/10% glycerol and stored in aliquots at -80°C

and quantified as physical particles per μL following the method described in (Mittereder et

al., 1996).

Cell culture and Infection

U2OS (ATCC #HTB-96) and H1299 (ATCC #CRL-5803) and HEK293 (kindly provided by

G. Nemerow, TSRI, La Jolla USA) cells were maintained at 37°C and 5% CO_2 in Dulbecco's

modified Eagle's medium (DMEM)-Glutamax supplemented with 10% fetal calf serum (FCS)

and 1% of Penicillin/Streptavidin. U2OS cells stably expressing mCherry-TAF-Iβ (Komatsu
et al., 2016) were transduced with lentiviruses expressing MS2BP-NLS-GFP and double
positive cells were sorted by fluorescence activated cell sorting (FACS). The resulting cell
population was maintained without further selection. For control experiments using cytosolic
MS2BP (Fig. S6) stable mCherry-TAF-Iβ cells were transfected with an expression vector for
MS2BP-NES-GFP. For RNAscope and immunofluorescene (IF) analysis H1299 cells were
grown on 15mm glass coverslips. Subsequently cells were infected with 3000 physical
particles/cell of replicative E3-deleted HAd-C5 virus or E1/E3-deleted HAd-C5 GFP
expressing control virus, washed once with PBS and fixed at different time points with
paraformaldehyde 4%/PBS for 10min at room temperature. Fixed cells were washed 3 times
with PBS and slowly dehydrated in 50%/70%/100% ethanol for 5 min each and stored at -
20°C in 100% ethanol. For the RNAscope protocol cells were rehydrated for 2 min in
70%/50% ethanol and equilibrated for 10 min in 1x PBS.

Preparation of stable cell lines expression FLAG tagged H3.1 and H3.3

U2OS were seeded in 6-well plates (250 000 cells/well) and transfected with 2 µg of
pcDNA3-Histone H3.1-FLAG or pcDNA3-Histone H3.3-FLAG (kindly provided by K.
Nagata, Tsukuba University, Japan) using Lipofectamine 2000 (Invitrogen). Two days after
transfection, cells were selected with 3 mg/ml G418 for 13 days. Resistant cells were cloned
by limiting dilution and the uniform expression of the tagged histones was monitored by
immunofluorescence and western-blot on individual clones.

RNAscope

RNAscope experiment were performed using reagents and protocols from the
RNAscope® Multiplex Fluorescent Assay essentially as indicated by the supplier (ACD Bio).
Briefly, cells were incubated with Protease III (freshly diluted 1:30 with PBS) for 15 min at
room temperature (RT) and washed 3 times with PBS. Hybridisation with the E1A RNAscope
probe was performed for 2h at 40° in a humidified chamber. Hybridized cells were washed
twice for 2 min and hybridized with signal amplifier probes (final amplifier Amp 4 AltB-FL)
following the manufacturer's instructions. Hybridized cells were washed twice in PBS
followed by incubation in IF buffer (PBS/10% FCS/0.1% Saponin) for 15 min at room
temperature to saturate unspecific binding sites followed by IF analysis. Primary antibodies
(mouse anti-pVII) were diluted in IF buffer and incubated with cells for 1h at RT. Antibody
was removed by washing in PBS followed by incubation with secondary antibodies (mouse

Alexa 647) diluted in IF buffer for 1h at RT. Cells were washed with PBS, ddH$_2$O and

absolute ethanol, air-dried and mounted in DAKO mixed with DAPI (1:1000).

Live-cell imaging and image analysis

For live-cell imaging, U2OS cells stably expressing mCherry-TAF-Iβ and MS2BP-NLS-GFP

were seeded in ibidi μ-slide VI$^{0.4}$ (Ibidi). The next day, the medium in the ibidi μ-slide was

replaced by imaging medium (FluoBrite DMEM supplemented with Prolong antifade reagent,

both Life technologies) containing 3000 physical particles/cell of replicative BxAd5-E1-MS2

for 30min followed by washing with imaging medium to remove the inoculum. Infection and

subsequent cultivation were performed at 37°C. Cells were imaged using a Leica spinning-

disk microscopy system (x100 objective) equipped with an environmental chamber at a frame

rate of 1 frame per second using MetaMorph software. To generate kymographs, movies were

processed using the ImageJ KymographBuilder plugin

(https://imagej.net/KymographBuilder_:_Yet_Another_Kymograph_Fiji_plugin) on confocal

sections. Line width was adapted to cover all viral genomes throughout the length of the

movie and double positive genomes were scored as transcribing genomes. Mounted cells for

IF and RNAscope analysis were imaged using a SP8 confocal microscope (Leica) at the

Bordeaux Imaging Center (BIC) platform. Stacks (7x) were taken every 0,3μm using a

pinhole of 1 at a pixel size of 85 nm. Quantifications were done using an adapted semi-

automated macro in ImageJ (provided upon request). Briefly, channels were split and stacks

were Z-projected and the cell periphery was outlined manually. A threshold was applied to

every channel and single objects or colocalized objects exceeding a defined area of pixels

were counted. Statistical analysis was done using one-way ANOVA multicomparison test.

QUANTIFICATION AND STATISTICAL ANALYSIS

Virus annotation

723 The reference genome sequence and gene annotation of Human Adenovirus C serotype 5

724 (accession: AY339865.1) was downloaded from NCBI. The annotation was adapted to the

725 used Adenovirus strain by excising nucleotides 28,593-30,464 from the full version (modified

726 virus reference genome and gene annotation available at the github project page).

RNA-seq analysis

STAR (Dobin et al., 2013) was used to align the RNA-seq reads simultaneously to the human

and the adenoviral genome. Therefore, a STAR index was generated based on the hg19

reference, the modified adenoviral genome and gene annotation (see section *Virus annotation*

for details) and the GENCODE human transcript annotation version 25. Read mapping was
conducted using following parameters: --outSAMmultNmax 1 --outMultimapperOrder
Random --outFilterType BySJout --outFilterIntronMotifs RemoveNoncanonical --
outSAMstrandField intronMotif --outFilterMultimapNmax 20 --alignSJoverhangMin 8 --
alignSJDBoverhangMin 1 --alignIntronMin 20 --alignIntronMax 1000000
To quantify the RNA abundance for each gene the number of uniquely mapped reads per gene
was counted using the featureCounts function of the subread package (Liao et al., 2014). The
read counts were normalized to transcripts per million (TPMs).

MNase-seq analysis
A bowtie2 index comprised of the human reference genome hg19 and the sequence of the
used Adenovirus strain (see section *Virus annotation* for details) was built (code available at
(Schwartz and Längst, 2015)) and paired-end reads were mapped to the index using bowtie2-
aligner (Langmead and Salzberg, 2012) with following parameters: --very-sensitive-local --
no-discordant. Reads were filtered to mapping quality $\geq$ 30 and only reads still mapped in
proper pairs were kept using samtools (Li et al., 2009). Reads aligned to the Adenovirus
genome were extracted for further analysis. The fragment size distribution was plotted using
the CollectInsertSizeMetrics function from the Picard toolkit
(https://broadinstitute.github.io/picard; Broad Institute).

pVII positioning analysis
Fragments mapped to the Adenovirus reference with a maximum insert size of 139 bp were
selected and used for pVII positioning analysis. pVII positions were called using the
DANPOS2 toolkit (Chen et al., 2013). First, a pVII occupancy profile for each sample was
generated and normalized to 20,000 reads using the dpos function with following parameters:
-u 1e-5; -a 1; -z 5; -p 0.001, --extend 35 and --mifrsz 0. Next, the profiles were normalized by
a genome wide quantile normalization to the first replicate at 0 hpi of each MNase condition
(high/low) using the wiq script. Finally, pVII positions were obtained from the normalized
pVII occupancy profiles using the dpos function with following parameters: -jd 50; -q 10; -z
5; -a 1.

Rotational positioning analysis
49 to 51 bp sized fragments mapped to the Adenovirus reference of all high MNase samples
were selected and pooled together. Frequencies of A/T or G/C di-nucleotides around the
fragment center were calculated using the annotatePeaks.pl script from the HOMER suit
(Heinz et al., 2010). Noise was filtered using the Fast Fourier Transform build in the filterFFT

function of the nucleR Bioconductor package (Flores and Orozco, 2011). The cutoff was set
with the option pcKeepComp=0.2. Finally, to reflect the symmetry of the pVII binding the
reverse di-nucleotide frequency was added, and the total frequency was divided by 2. The
same procedure was conducted to derive the di-nucleotide periodicity of nucleosomal
fragments, except that 147 bp fragments of the first replicate at 0 hpi were selected.

Ad genome accessibility

DNA fragments released under low MNase conditions were used to identify structural
changes of Ad chromatin over time. Therefore, fragments mapped to the Ad genome were
filtered by a maximum length of 140 bp and replicates of each timepoint were pooled. Next,
all timepoints were pooled to call peaks using the DANPOS2 toolkit (Chen et al., 2013) with
following parameters: dpos -m 1 --extend 35 -c 34062 -u 0 -z 20 -jd 70 -q 15 -a 1 -e 1.
Fragments of each time were counted at the determined peaks using featureCounts (Liao et
al., 2014) with following parameters: --fracOverlap 0.7 -p -B –largestOverlap. Peaks
exhibiting in total less than 30 fragment counts were discarded. Fragment counts per peak
were normalized to CPMs (counts per million). The accessibility score was obtained for each
peak by fitting a linear regression on the logarithmical normalized count frequencies
computed against the logarithmical infection time in min. The slope of the linear fit was used
as a measure of dynamic changes of viral genome accessibility (accessibility score).

Transcription factor motif identification

The Ad DNA sequence from position 19,890 to 19,940 was extracted and uploaded to the
PROMO web server (Messeguer et al., 2002). Motif search was performed against known
human and mouse motifs stored in TRANSFAC data base using the SearchSites tool. The
maximum dissimilarity rate was set to 2 %. The obtained motifs were further filtered to have
at least a recognition sequence of 5 nucleotides and that the corresponding transcription factor
is expressed in the infected H1299 cells.

ChIP-seq analysis

Paired-end reads were mapped against viral and human reference genomes and subsequently
filtered as described in the MNase-seq analysis section. CPM-normalized coverage tracks
were generated using the bamCoverage script of the deepTools package (Ramírez et al.,
2016). PCA and correlation heatmap were generated using the multiBigwigSummary (50 bp
bin size), plotPCA and plotCorrelation deepTools functions.

Sites of nucleosome assembly

795
The R Bioconductor package nucleR (Flores and Orozco, 2011) was used to call sites of
nucleosome assembly onto the Ad genome. First, fragments specific to the Ad genome were
loaded into the R environment. Fragments between 137 to 157 bp from time points 1, 2 and 4
hpi of the high MNase condition were extracted for further analysis. The coverage was
normalized to reads per million Ad mapped reads and noise was removed using the built-in
Fast Fourier Transform (filterFFT with option pcKeepComp=0.007). Potential nucleosome
positions were called based on the normalized profile using the peakDetection function with
the option width=147. Peaks at genomic sites, which exhibit a mean H3.3 ChIP signal
(U2OS33) of at least 2 CPM at 4 hpi were used as nucleosome assembly positions.
805
806

ACCESSION NUMBERS

The Gene Expression Omnibus accession number for the MNase-seq, ChIP-seq and RNA-seq

data reported in this paper is GSE136550. The code used to analyze the sequencing data is

available at GitHub: https://github.com/uschwartz/AdVir

SUPPLEMENTAL INFORMATION

Supplemental Information includes 8 figures, 4 movies, 2 tables.

COMPETING INTEREST

The authors declare no competing interests

AUTHOR CONTRIBUTIONS

Conceptualization H.W. and G.L.; Methodology F.L., T.K., E.S., C.H., M.N., E.B., F.R.,

E.B., M.R., H.W. and G.L.; Formal Analysis F.L., U.S. and H.W.; Investigation T.K., C.H.,

F.L., C.B., E.S., M.R., H.W. and G.L., Data Curation U.S; Visualization F.L., U.S., C.B. and

H.W.; Writing – Original Draft U.S. G.L. and H.W.; Writing – Review & Editing E.B., T.K.,

U.S., M.R., G.L. and H.W., Resources E.B., H.W and G.L; Supervision H.W and G.L;

Funding Acquisition G.L. and H.W.

ACKNOWLEDGEMENTS

REFERENCES

Akusjarvi, G. (2008). Temporal regulation of adenovirus major late alternative RNA splicing.

Front. Biosci. *13*, 5006–5015.

Anderson, C.W., Young, M.E., and Flint, S.J. (1989). Characterization of the adenovirus 2

virion protein, mu. Virology *172*, 506–512.

Avgousti, D.C., Herrmann, C., Kulej, K., Pancholi, N.J., Sekulic, N., Petrescu, J., Molden,

R.C., Blumenthal, D., Paris, A.J., Reyes, E.D., et al. (2016). A core viral protein binds host

nucleosomes to sequester immune danger signals. Nature *535*, 173–177.

Avgousti, D.C., Della Fera, A.N., Otter, C.J., Herrmann, C., Pancholi, N.J., and Weitzman,

M.D. (2017). Adenovirus Core Protein VII Downregulates the DNA Damage Response on the

Host Genome. J. Virol. *91*.

Benevento, M., Di Palma, S., Snijder, J., Moyer, C.L., Reddy, V.S., Nemerow, G.R., and

Heck, A.J.R. (2014). Adenovirus composition, proteolysis, and disassembly studied by in-

depth qualitative and quantitative proteomics. J. Biol. Chem. *289*, 11421–11430.

Bertrand, E., Chartrand, P., Schaefer, M., Shenoy, S.M., Singer, R.H., and Long, R.M. (1998).

Localization of ASH1 mRNA particles in living yeast. Mol. Cell *2*, 437–445.

Brison, O., Kédinger, C., and Chambon, P. (1979). Adenovirus DNA template for late

transcription is not a replicative intermediate. J. Virol. *32*.

Brown, D.T., Westphal, M., Burlingham, B.T., Winterhoff, U., and Doerfler, W. (1975).

Structure and composition of the adenovirus type 2 core. J. Virol. *16*, 366–387.

Cassany, A., Ragues, J., Guan, T., Bégu, D., Wodrich, H., Kann, M., Nemerow, G.R., and

Gerace, L. (2015). Nuclear Import of Adenovirus DNA Involves Direct Interaction of Hexon

with an N-Terminal Domain of the Nucleoporin Nup214. J. Virol. *89*, 1719–1730.

Chatterjee, P.K., Vayda, M.E., and Flint, S.J. (1985). Interactions among the three adenovirus

core proteins. J. Virol. *55*, 379–386.

Chatterjee, P.K., Vayda, M.E., and Flint, S.J. (1986). Adenoviral protein VII packages

intracellular viral DNA throughout the early phase of infection. EMBO J. *5*, 1633–1644.

Chen, J., Morral, N., and Engel, D.A. (2007). Transcription releases protein VII from

adenovirus chromatin. Virology *369*, 411–422.

Chen, K., Xi, Y., Pan, X., Li, Z., Kaestner, K., Tyler, J., Dent, S., He, X., and Li, W. (2013).

DANPOS: Dynamic analysis of nucleosome position and occupancy by sequencing. Genome

Res. *23*, 341–351.

Chereji, R. V., Ocampo, J., and Clark, D.J. (2017). MNase-Sensitive Complexes in Yeast:

Nucleosomes and Non-histone Barriers. Mol. Cell *65*, 565-577.e3.

Chereji, R. V., Bryson, T.D., and Henikoff, S. (2019). Quantitative MNase-seq accurately

maps nucleosome occupancy levels. Genome Biol. *20*, 198.

Corden, J., Engelking, H.M., and Pearson, G.D. (1976). Chromatin-like organization of the

adenovirus chromosome. Proc. Natl. Acad. Sci. U. S. A. *73*, 401–404.

Crisostomo, L., Soriano, A.M., Mendez, M., Graves, D., and Pelka, P. (2019). Temporal

dynamics of adenovirus 5 gene expression in normal human cells. PLoS One *14*, e0211192.

Daniell, E., Groff, D.E., and Fedor, M.J. (1981). Adenovirus chromatin structure at different

stages of infection. Mol. Cell. Biol. *1*, 1094–1105.

Déry, C. V., de Murcia, G., Lamarre, D., Morin, N., Poirier, G.G., and Weber, J. (1986).

Possible role of ADP-ribosylation of adenovirus core proteins in virus infection. Virus Res. *4*,

313–329.

Diermeier, S., Kolovos, P., Heizinger, L., Schwartz, U., Georgomanolis, T., Zirkel, A.,

Wedemann, G., Grosveld, F., Knoch, T.A., Merkl, R., et al. (2014). TNFα signalling primes

chromatin for NF-κB binding and induces rapid and widespread nucleosome repositioning.

Genome Biol. *15*, 536.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,

M., and Gingeras, T.R. (2013). STAR: Ultrafast universal RNA-seq aligner. Bioinformatics

29, 15–21.

Everitt, E., Lutter, L., and Philipson, L. (1975). Structural proteins of adenoviruses. XII.

Location and neighbor relationship among proteins of adenovirion type 2 as revealed by

enzymatic Iodination, immunoprecipitation and chemical cross-linking. Virology *67*, 197–

208.

Flint, S.J. (1982). Expression of adenoviral genetic information in productively infected cells.

Biochim. Biophys. Acta *651*, 175–208.

Flores, O., and Orozco, M. (2011). nucleR: a package for non-parametric nucleosome

positioning. Bioinformatics *27*, 2149–2150.

Gao, J., Mese, K., Bunz, O., and Ehrhardt, A. (2019). State-of-the-art human adenovirus

vectorology for therapeutic approaches. FEBS Lett. *593*, 3609–3622.

Genoveso, M.J., Hisaoka, M., Komatsu, T., Wodrich, H., Nagata, K., and Okuwaki, M.

(2020). Formation of adenovirus DNA replication compartments and viral DNA accumulation

sites by host chromatin regulatory proteins including NPM1. FEBS J. *287*, 205–217.

Giberson, A.N., Davidson, A.R., and Parks, R.J. (2011). Chromatin structure of adenovirus

DNA throughout infection. Nucleic Acids Res. *40*, 2369–2376.

Giberson, A.N., Saha, B., Campbell, K., Christou, C., Poulin, K.L., and Parks, R.J. (2018).

Human adenoviral DNA association with nucleosomes containing histone variant H3.3 during

the early phase of infection is not dependent on viral transcription or replication. Biochem.

Cell Biol. *96*, 797–807.

Glenn, G.M., and Ricciardi, R.P. (1988). Detailed kinetics of adenovirus type-5 steady-state

transcripts during early infection. Virus Res. *9*, 73–91.

Greber, U.F., Willetts, M., Webster, P., and Helenius, A. (1993). Stepwise dismantling of

adenovirus 2 during entry into cells. Cell *75*, 477–486.

Hammarskjöld, M.L., and Winberg, G. (1980). Encapsidation of adenovirus 16 DNA is

directed by a small DNA sequence at the left end of the genome. Cell *20*, 787–795.

Haruki, H., Gyurcsik, B., Okuwaki, M., and Nagata, K. (2003). Ternary complex formation

between DNA-adenovirus core protein VII and TAF-Ibeta/SET, an acidic molecular

chaperone. FEBS Lett. *555*, 521–527.

Haruki, H., Okuwaki, M., Miyagishi, M., Taira, K., and Nagata, K. (2006). Involvement of

template-activating factor I/SET in transcription of adenovirus early genes as a positive-acting

factor. J. Virol. *80*, 794–801.

Hearing, P., Samulski, R.J., Wishart, W.L., and Shenk, T. (1987). Identification of a repeated

sequence element required for efficient encapsidation of the adenovirus type 5 chromosome.

J. Virol. *61*, 2555–2558.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,

Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription

factors prime cis-regulatory elements required for macrophage and B cell identities. Mol. Cell

38, 576–589.

Hu, M., Depledge, D.P., Flores Cortes, E., Breuer, J., and Schang, L.M. (2019). Chromatin

dynamics and the transcriptional competence of HSV-1 genomes during lytic infections.

PLOS Pathog. *15*, e1008076.

Inturi, R., Mun, K., Singethan, K., Schreiner, S., and Punga, T. (2017). Human Adenovirus

Infection Causes Cellular E3 Ubiquitin Ligase MKRN1 Degradation Involving the Viral Core

Protein pVII. J. Virol. *92*.

Jiang, C., and Pugh, B.F. (2009). Nucleosome positioning and gene regulation: advances

through genomics. Nat. Rev. Genet. *10*, 161–172.

Johnson, J.S., Osheim, Y.N., Xue, Y., Emanuel, M.R., Lewis, P.W., Bankovich, A., Beyer,

A.L., and Engel, D.A. (2004). Adenovirus Protein VII Condenses DNA, Represses

Transcription, and Associates with Transcriptional Activator E1A. J. Virol. *78*, 6459–6468.

Kaplan, N., Moore, I.K., Fondufe-Mittendorf, Y., Gossett, A.J., Tillo, D., Field, Y., LeProust,

E.M., Hughes, T.R., Lieb, J.D., Widom, J., et al. (2009). The DNA-encoded nucleosome

organization of a eukaryotic genome. Nature *458*, 362–366.

Keller, M., Tagawa, T., Preuss, M., and Miller, A.D. (2002). Biophysical characterization of

the DNA binding and condensing properties of adenoviral core peptide mu. Biochemistry *41*,

652–659.

Kent, N.A., Adams, S., Moorhouse, A., and Paszkiewicz, K. (2011). Chromatin particle

spectrum analysis: a method for comparative chromatin structure analysis using paired-end

mode next-generation DNA sequencing. Nucleic Acids Res. *39*.

Komatsu, T., and Nagata, K. (2012). Replication-Uncoupled Histone Deposition during

Adenovirus DNA Replication. J. Virol. *86*, 6701–6711.

Komatsu, T., Haruki, H., and Nagata, K. (2011). Cellular and viral chromatin proteins are

positive factors in the regulation of adenovirus gene expression. Nucleic Acids Res. *39*, 889–

901.

Komatsu, T., Dacheux, D., Kreppel, F., Nagata, K., and Wodrich, H. (2015). A method for

visualization of incoming adenovirus chromatin complexes in fixed and living cells. PLoS

One *10*, e0137102.

Komatsu, T., Nagata, K., and Wodrich, H. (2016). An Adenovirus DNA Replication Factor,

but Not Incoming Genome Complexes, Targets PML Nuclear Bodies. J. Virol. *90*, 1657–

1667.

Kubik, S., Bruzzone, M.J., Jacquet, P., Falcone, J.L., Rougemont, J., and Shore, D. (2015).

Nucleosome Stability Distinguishes Two Different Promoter Types at All Protein-Coding

Genes in Yeast. Mol. Cell *60*, 422–434.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat.

Methods *9*, 357–359.

Längst, G., Schätz, T., Langowski, J., and Grummt, I. (1997). Structural analysis of mouse

rDNA: Coincidence between nuclease hypersensitive sites, DNA curvature and regulatory

elements in the intergenic spacer. Nucleic Acids Res. *25*, 511–517.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

and Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics

25, 2078–2079.

Liao, Y., Smyth, G.K., and Shi, W. (2014). FeatureCounts: An efficient general purpose

program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–930.

Lischwe, M.A., and Sung, M.T. (1977). A histone-like protein from adenovirus chromatin

[37]. Nature *267*, 552–554.

Martinez, R., Schellenberger, P., Vasishtan, D., Aknin, C., Austin, S., Dacheux, D., Rayne, F.,

Siebert, A., Ruzsics, Z., Gruenewald, K., et al. (2015). The Amphipathic Helix of Adenovirus

Capsid Protein VI Contributes to Penton Release and Postentry Sorting. J. Virol. *89*, 2121–

2135.

Matsumoto, K., Okuwaki, M., Kawase, H., Handa, H., Hanaoka, F., and Nagata, K. (1995).

Stimulation of DNA transcription by the replication factor from the adenovirus genome in a

chromatin-like structure. J. Biol. Chem. *270*, 9645–9650.

Messeguer, X., Escudero, R., Farre, D., Nunez, O., Martinez, J., and Alba, M.M. (2002).

PROMO: detection of known transcription regulatory elements using species-tailored

searches. Bioinformatics *18*, 333–334.

Mieczkowski, J., Cook, A., Bowman, S.K., Mueller, B., Alver, B.H., Kundu, S., Deaton,

A.M., Urban, J.A., Larschan, E., Park, P.J., et al. (2016). MNase titration reveals differences

between nucleosome occupancy and chromatin accessibility. Nat. Commun. *7*, 11485.

Minderjahn, J., Schmidt, A., Fuchs, A., Schill, R., Raithel, J., Babina, M., Schmidl, C.,

Gebhard, C., Schmidhofer, S., Mendes, K., et al. (2020). Mechanisms governing the

pioneering and redistribution capabilities of the non-classical pioneer PU.1. Nat. Commun.

11, 402.

Mirza, M.A., and Weber, J. (1982). Structure of adenovirus chromatin. Biochim. Biophys.

Acta *696*, 76–86.

Mittereder, N., March, K.L., and Trapnell, B.C. (1996). Evaluation of the concentration and

bioactivity of adenovirus vectors for gene therapy. J. Virol. *70*, 7498–7509.

Mueller, B., Mieczkowski, J., Kundu, S., Wang, P., Sadreyev, R., Tolstorukov, M.Y., and

Kingston, R.E. (2017). Widespread changes in nucleosome accessibility without changes in

nucleosome occupancy during a rapid transcriptional induction. Genes Dev. *31*, 451–462.

Mun, K., and Punga, T. (2018). Cellular Zinc Finger Protein 622 Hinders Human Adenovirus

Lytic Growth and Limits Binding of the Viral pVII Protein to Virus DNA. J. Virol. *93*.

Nevins, J.R., Ginsberg, H.S., Blanchard, J.M., Wilson, M.C., and Darnell, J.E. (1979).

Regulation of the primary expression of the early adenovirus transcription units. J. Virol. *32*,

727–733.

Noll, M., Thomas, J.O., and Kornberg, R.D. (1975). Preparation of Native Chromatin and

Damage Caused by Shearing. Science (80-.). *187*, 1203–1206.

Ostapchuk, P., Suomalainen, M., Zheng, Y., Boucke, K., Greber, U.F., and Hearing, P.

(2017). The adenovirus major core protein VII is dispensable for virion assembly but is

essential for lytic infection. PLoS Pathog. *13*.

Pheasant, K., Möller-Levet, C.S., Jones, J., Depledge, D., Breuer, J., and Elliott, G. (2018).

Nuclear-cytoplasmic compartmentalization of the herpes simplex virus 1 infected cell

transcriptome is co-ordinated by the viral endoribonuclease vhs and cofactors to facilitate the

translation of late proteins. PLoS Pathog. *14*.

Pied, N., and Wodrich, H. (2019). Imaging the adenovirus infection cycle. FEBS Lett. *593*,

3419–3448.

Puntener, D., Engelke, M.F., Ruzsics, Z., Strunze, S., Wilhelm, C., and Greber, U.F. (2011).

Stepwise Loss of Fluorescent Core Protein V from Human Adenovirus during Entry into

Cells. J. Virol. *85*, 481–496.

Ramírez, F., Ryan, D.P., Bj¨, B., Grüning, B., Grüning, G., Bhardwaj, V., Kilpert, F., Richter,

A.S., Heyne, S., Undar, F.D.¨, et al. (2016). deepTools2: a next generation web server for

deep-sequencing data analysis. Web Serv. Issue Publ. Online *44*.

Ross, P.J., Kennedy, M. a, Christou, C., Risco Quiroz, M., Poulin, K.L., and Parks, R.J.

(2011). Assembly of helper-dependent adenovirus DNA into chromatin promotes efficient

gene expression. J. Virol. *85*, 3950–3958.

Ruzsics, Z., Lemnitzer, F., and Thirion, C. (2014). Engineering adenovirus genome by

bacterial artificial chromosome (BAC) technology. Methods Mol. Biol. *1089*, 143–158.

Samad, M.A., Komatsu, T., Okuwaki, M., and Nagata, K. (2012). B23/nucleophosmin is

involved in regulation of adenovirus chromatin structure at late infection stages, but not in

virus replication and transcription. J. Gen. Virol. *93*, 1328–1338.

Sato, K., and Hosokawa, K. (1984). Analysis of the interaction between DNA and major core

protein in adenovirus chromatin by circular dichroism and ultraviolet light induced cross-

linking. J. Biochem. *95*, 1031–1039.

Schones, D.E., Cui, K., Cuddapah, S., Roh, T.Y., Barski, A., Wang, Z., Wei, G., and Zhao, K.

(2008). Dynamic Regulation of Nucleosome Positioning in the Human Genome. Cell *132*,

887–898.

Schwartz, U., and Längst, G. (2015). Bioinformatic Analysis of ChIP-seq Data on the

Repetitive Ribosomal RNA Gene. Encycl. Cell Biol. *3*, 254–269.

Schwartz, U., Németh, A., Diermeier, S., Exler, J.H., Hansch, S., Maldonado, R., Heizinger,

L., Merkl, R., and Längst, G. (2018). Characterizing the nuclease accessibility of DNA in

human cells to map higher order structures of chromatin. Nucleic Acids Res. *47*, 1239–1254.

Sekine, E., Schmidt, N., Gaboriau, D., and O'Hare, P. (2017). Spatiotemporal dynamics of

HSV genome nuclear entry and compaction state transitions using bioorthogonal chemistry

and super-resolution microscopy. PLoS Pathog. *13*, e1006721.

Strunze, S., Engelke, M.F., Wang, I.-H., Puntener, D., Boucke, K., Schleich, S., Way, M.,

Schoenenberger, P., Burckhardt, C.J., and Greber, U.F. (2011). Kinesin-1-Mediated Capsid

Disassembly and Disruption of the Nuclear Pore Complex Promote Virus Infection. Cell Host

Microbe *10*, 210–223.

Suomalainen, M., Prasad, V., Kannan, A., and Greber, U.F. (2020). Cell-to-cell and genome-

to-genome variability of adenovirus transcription tuned by the cell cycle. J. Cell Sci. *134*.

Valouev, A., Johnson, S.M., Boyd, S.D., Smith, C.L., Fire, A.Z., and Sidow, A. (2011).

Determinants of nucleosome organization in primary human cells. Nature *474*, 516–520.

Vayda, M.E., Rogers, A.E., and Flint, S.J. (1983). The structure of nucleoprotein cores

released from adenovirions. Nucleic Acids Res. *11*, 441–460.

Wickham, T.J., Mathias, P., Cheresh, D.A., and Nemerow, G.R. (1993). Integrins alpha v beta

3 and alpha v beta 5 promote adenovirus internalization but not virus attachment. Cell *73*,

309–319.

Widom, J. (2001). Role of DNA sequence in nucleosome stability and dynamics. Q. Rev.

Biophys. *34*, 269–324.

Wiethoff, C.M., Wodrich, H., Gerace, L., and Nemerow, G.R. (2005). Adenovirus protein VI

mediates membrane disruption following capsid disassembly. J. Virol. *79*, 1992–2000.

Xue, Y., Johnson, J.S., Ornelles, D.A., Lieberman, J., and Engel, D.A. (2005). Adenovirus

Protein VII Functions throughout Early Phase and Interacts with Cellular Proteins SET and

pp32. J. Virol. *79*, 2474–2483.

1062

FIGURES AND LEGENDS

1064

1065

1066

Figure 1. Spatio-temporal analysis of adenovirus transcription using RNA-seq. **A)**
Experimental design. Virus was added (+Ad) and removed (Wash) after 30 min and cell were
processed as indicated (hpi = hours post infection) using immunofluorescence (IF), RNA
sequencing (RNA-seq), nucleosome sequencing (MNase-seq) and ChIP sequencing (ChIP-
seq). **B)** Profile illustrating Ad genome coverage by RNA-seq reads. Reads were normalized
to RPM (reads per million mapped reads) and the average of two replicates is shown.
Genomic location and underlying gene annotation of HAd-C5 (NCBI accession:
AY339865.1) are indicated at the bottom. The area deleted in the HAd-C5 strain is indicated
by the triangle. **C)** Transcript abundance estimation of early Ad genes. TPMs (Transcripts Per
Kilobase Million) for early expressed genes were calculated at each timepoint. The barplots
show the average of two replicates and each replicate is indicated by a black dot.

1078

1079

Figure 2. E1A transcription at the single cell level. **A)** Representative confocal images of
cells infected with HAd-C5 at different time points as indicated. Detection of E1A transcript
with specific RNAscope probes (left column, green signal), genomes using anti-pVII
antibodies (middle column, magenta signal) and merged with DAPI detecting the nucleus
(right column, grey signal). Scale bar is 10µm. **B)** Representative confocal images of cells
infected with HAd-C5 (Ad5, top row) or an E1-deleted HAd-C5 vector (Ad5-GFP, bottom
row). Scale bar is 10µm. **C)** Quantification of nuclear pVII foci over time as indicated on the
x-axis. The distribution of normalized pVII foci at each time point is shown as scatter plot
(Mean +/- SD, n = 30 cells per time point). **D)** Graph as in (C) showing the distribution of
nuclear E1A transcript foci per cell as scatter plot (Mean +/- SD, n = 30 cells per time point).
E) Scatter plot showing the distribution of nuclear pVII foci (C) coinciding with E1A
transcript foci (D) (Mean +/- SD, n = 30 nuclei per time point). The number of E1A positive
genomes was normalized by the total number of genomes per nucleus. Statistical analysis for
C-E via one-way ANOVA multi-comparison test.

1094

Figure 3. E1A transcription in living cells. **A)** Representative images showing t-projections of
60 frames (frame rate 1F/sec) imaged at ~4 hpi. Cells express mCherry-TAF-Iβ showing

viral genomes (first column, red signal in merge) and MS2BP-NLS-GFP showing MS2-repeat
tagged E1AmRNAs (second column, green signal in merge) and an overlay (third column,
merge) of both channels. The first row shows non-infected control cells, the second row cells
infected with untagged control virus and the third and fourth row examples of cells infected
with virus expressing E1AmRNA tagged with MS2-repeats. Black arrows point at non-
transcribing and white arrows at transcribing genomes. Details are shown as magnifications of
the boxed area. Scale bar is 10µm. **B)** Kymograph analysis of the bottom two cells shown in
A and as described in the material and methods section. Cell overview to the left, kymograph
separated into the individual channels and a superimposition of both channels to the right.
Genome position over time appears as red vertical line, E1AmRNA signal as green vertical
line. The boxed area in the overview corresponds to the boxed area in the kymograph. Green
arrows point at the E1AmRNA signal, orange arrows at transcribing (double positive) and red
arrow at non-transcribing (single positive) genomes. **C)** Quantitative kymograph analysis.
Scatter plot representation of the percentage of MS2-BP positive genomes per cell (n = 11
cells) vs. control cells infected with viruses expressing untagged E1AmRNA (n = 5 cells)
determined by kymograph analysis. **D)** Scatter plot representation comparing the percentage
of E1AmRNA positive genomes per cell determined with the MS2-BP method (left, n = 11
cells) vs. the RNAscope method (right, n = 22 cells). Unpaired t-test was used for statistical
analysis in C and D.
1116

1117

Figure 4. Characterization of Ad DNA packaging by pVII. **A)** Schematic view of MNase
digestion of infected cells. Agarose gel shows nucleosome ladders of a MNase titration.
Mono-, di- and tri-nucleosomes are indicated at the right. The mono-nucleosomal DNA
fraction was isolated after MNase digestions and subjected to high-throughput sequencing. **B)**
Fragment size distribution of paired-end reads either mapped to the human genome (red) or
the Ad genome (purple). **C)** Average WW (A/T)- and SS (C/G)-di-nucleotide frequency of
nucleosomal (upper panel) and pVII fragments (lower panel). Size-selected fragments were
centred at the midpoint and the di-nucleotide frequency was calculated. Every 10.3
(nucleosome) / 5.4 (pVII) bp a dashed line highlights the periodicity of the signal. **D)** Profile
illustrates Ad genome coverage by MNase-seq reads. pVII positions, which have been called
with the DANPOS2 package (Chen et al., 2013), are represented by grey boxes. Partitioning
of the genome into functional domains and the timing of their transcription is indicated by the
color code (lightblue = early, darkblue=late). Genomic location and underlying gene
annotation of HAd-C5 (NCBI accession: AY339865.1) are indicated at the bottom. **E)**
Boxplot showing the number of detected pVII peaks for each sample. **F)** Boxplot showing the
genome-wide pVII repeat length (pVII-RL) for each sample. **G)** and **H)** Boxplots showing the
distribution of pVII center-center distances / peak heights in each domain. Individual

1135 replicates are shown side by side. **I)** Barplot showing the number of pVII peaks per kb
1136 detected in the domains. Individual replicates are shown as black dots.

1137

Figure 5. Early decondensation of Ad chromatin. **A)** and **B)** Sample-wise Pearson
correlations of **A)** high MNase or **B)** low MNase profiles. **C)** Profile illustrates Ad genome
coverage by low MNase-seq fragments. The average of two replicates is shown, except at
timepoint 0 hpi where only one replicate was available. The accessibility score was calculated
as the slope of a linear regression fit of the average signal intensity across each timepoint. The
score was assessed for each peak. Yellow bars depict increased accessibility and blue bar
show peaks with reduced DNA accessibility. Domains of open and closed Ad chromatin are
shown below. A new accessibility peak arising during infection in the Late3 region is marked
by an asterisk. **D)** Barplot showing the average accessibility score in each domain.

1147

1148

1149

Figure 6. Nucleosome assembly onto the Ad genome. **A)** Fragment size distribution of
paired-end reads from high MNase experiments (see Fig. 4) either mapping to the human
genome (grey) or the Ad genome (shades of purple). Nucleosome sized fragments are
highlighted between two dashed lines at 137 bp and 157 bp. **B)** Quantification of ChIP-seq
fragments mapping to the Ad genome over time. Flag/H3K27ac immunoprecipitation (IP)
was conducted in U2OS cells expressing either Flag-tagged H3.1 (U2OS31, blue shades) or
H3.3 (U2OS33, red shades). The barplots show the average of 2-3 replicates (black dots) of
each time point and condition. The antibody used in the IP is indicated at the top. **C)** Principal
component analysis (PCA) of ChIP-seq samples. **D)** H3.3 and H3K27ac profiles along the Ad
genome. Normalized coverage of Flag (blue shades) / H3K27ac (green shades) ChIP-seq
experiments in U2OS33. The average signal of 3 replicates is shown and the 95% confidence
interval is indicated. CPM = counts per million mapped fragments **E)** Average coverage of
Flag-tagged H3.3 or **F)** H3K27ac in specific Ad domains. **G)** and **H)** Genome browser tracks
showing pVII occupancy at 0 hpi (upper panel), the H3.3 (blue shades) / H3K27ac (green

1165   shades) accumulation over time, predicted sites of nucleosome assembly (grey boxes) using

1166   nucleosome-sized fragments (137-157 bp) from 4 hpi MNase-seq samples and expression

1167   levels of early genes (middle panel).

1168

1169

1170